



Title	EULER CHARACTERISTICS ON A CLASS OF FINITELY GENERATED NILPOTENT GROUPS
Author(s)	Hamrouni, Hatem
Citation	Osaka Journal of Mathematics. 2013, 50(2), p. 339-346
Version Type	VoR
URL	<a href="https://doi.org/10.18910/25090">https://doi.org/10.18910/25090</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

# EULER CHARACTERISTICS ON A CLASS OF FINITELY GENERATED NILPOTENT GROUPS

HATEM HAMROUNI

(Received July 19, 2011)

## Abstract

A finitely generated torsion free nilpotent group is called an  $\mathcal{F}$ -group. To each  $\mathcal{F}$ -group  $\Gamma$  there is associated a connected, simply connected nilpotent Lie group  $G_\Gamma$ . Let  $\text{TUF}$  be the class of all  $\mathcal{F}$ -group  $\Gamma$  such that  $G_\Gamma$  is totally unimodular. A group in  $\text{TUF}$  is called  $\text{TUF}$ -group. In this paper, we are interested in finding non-zero Euler characteristic on the class  $\text{TUF}$  and therefore, on  $\text{TUFF}$ , the class of groups  $K$  having a subgroup  $\Gamma$  of finite index in  $\text{TUF}$ . An immediate consequence we obtain that any two isomorphic finite index subgroups of a  $\text{TUFF}$ -group have the same index. As applications, we give two results, the first is a generalization of Belegradek's result, in which we prove that every  $\text{TUFF}$ -group is co-hopfian. The second is a known result due to G.C. Smith, asserting that every  $\text{TUFF}$ -group is not compressible.

## 1. Introduction and main results

We follow [5, p.222] in defining an Euler characteristic on a class of groups as follows (see also [2, p.1]).

**DEFINITION 1.1** (Euler characteristic). Let  $\mathfrak{X}$  be a class of groups closed under taking subgroups of finite index. By an Euler characteristic on  $\mathfrak{X}$  it meant a function  $\chi : \mathfrak{X} \rightarrow \mathbb{R}$  satisfying

(Ec1) If  $K$  and  $H$  are in  $\mathfrak{X}$ , and  $K$  is isomorphic to  $H$ , then  $\chi(K) = \chi(H)$ .

(Ec2) If  $K$  is in  $\mathfrak{X}$ , and  $H$  is a subgroup of  $K$  of finite index, then  $\chi(H) = [K : H]\chi(K)$ , where  $[K : H]$  denotes the index of  $H$  in  $K$ .

In this paper, we are interested in finding non zero Euler characteristics defined on a class of finitely generated nilpotent groups.

Let  $G$  be a connected Lie group and  $\text{Aut}(G)$  its group of continuous automorphisms. Let  $\mu$  be a Haar measure on  $G$ . For every  $\alpha \in \text{Aut}(G)$  we have

$$\alpha_*^{-1}\mu = \Delta(\alpha)\mu,$$

where  $\alpha_*^{-1}\mu$  is the push forward of  $\mu$  under  $\alpha^{-1}$ , and  $\Delta: \text{Aut}(G) \rightarrow \mathbb{R}_+^*$  is a homomorphism of  $\text{Aut}(G)$  into the multiplicative group of the positive reals. If  $G$  is a connected, simply connected nilpotent Lie group, then

$$\Delta(\alpha) = |\det(\alpha)|.$$

**DEFINITION 1.2** ([10, p. 627]). A connected, simply connected nilpotent Lie group  $G$  is called totally unimodular if the image of  $\Delta$  is  $\{1\}$ .

Let  $\text{TULG}$  be the class of connected, simply connected totally unimodular nilpotent Lie groups.

A real Lie algebra is called *characteristically nilpotent* if all its derivations are nilpotent ([4, p. 157], [6, p. 623]). We note that a characteristically nilpotent Lie algebra is nilpotent. Let  $\text{CNLG}$  be the class of connected, simply connected nilpotent Lie groups  $G = \exp \mathfrak{g}$  such that  $\mathfrak{g}$  is a characteristically nilpotent Lie algebra.

**Proposition 1.3** ([10, (1.1)]). *We have*

$$\text{CNLG} \subset \text{TULG}.$$

A finitely generated torsion free nilpotent group is called an  $\mathcal{F}$ -group. Any  $\mathcal{F}$ -group  $\Gamma$  is isomorphic to a discrete uniform subgroup of a connected, simply connected nilpotent Lie group  $G_\Gamma$  whose Lie algebra  $\mathfrak{g}_\Gamma$  has rational structure constants ([8, Theorem 6]). Let  $\text{TUF}$  be the class of all  $\mathcal{F}$ -groups  $\Gamma$  such that  $G_\Gamma \in \text{TULG}$ . We call a group  $\Gamma$  a TUF-group if  $\Gamma \in \text{TUF}$ . For every integer  $n \geq 7$  there exists a  $n$ -dimensional characteristically nilpotent Lie algebra with rational structure ([14, Theorem 5]). By the Mal'cev rationality criterion (Theorem 2.1) we derive the following.

**Proposition 1.4.** *For every integer  $n \geq 7$  there exists a TUF-group with Hirsch length  $n$ .*

The main result of this paper is the following.

**Theorem 1.5.** *The class TUF admits Euler characteristics.*

In Section 3, we give an explicit Euler characteristic on  $\text{TUF}$ .

By [5, p. 222] (see also [15], [2]) every Euler characteristic  $\chi$  on  $\text{TUF}$  can be extended to  $\text{TUFF}$ , the class of groups  $K$  having a subgroup  $\Gamma$  of finite index in  $\text{TUF}$ , by setting

$$\chi(K) = \frac{1}{[K : \Gamma]} \chi(\Gamma).$$

As an immediate consequence we have the following.

**Proposition 1.6.** *Any two isomorphic finite index subgroups of a TUFF-group have the same index.*

DEFINITION 1.7 (Co-hopfian group). A group is called co-hopfian if it satisfies the following equivalent conditions:

- (1) It is not isomorphic to any proper subgroup.
- (2) Every injective endomorphism of the group is an automorphism.

As an easy consequence of Proposition 1.6, we obtain a generalization for I. Belegradek's result ([1, Corollary 2.4]).

**Proposition 1.8.** *Every TUFF-group is co-hopfian.*

We introduce the following definition due to G.C. Smith ([13, Definition 1]).

DEFINITION 1.9 (Compressible group). A group  $G$  is called compressible if any finite index subgroup of  $G$  contains a finite index subgroup isomorphic to  $G$ .

The following proposition which is due to G.C. Smith ([13, Proposition 4]) is an immediate consequence of Proposition 1.6.

**Proposition 1.10.** *Every TUFF-group is not compressible.*

## 2. Rational structures and discrete uniform subgroups

General references for the material in this section are [3] and [11] as well as the original paper of Mal'cev [8].

Let  $G$  be a connected and simply connected nilpotent Lie group with Lie algebra  $\mathfrak{g}$ . Then the exponential map  $\exp: \mathfrak{g} \rightarrow G$  is a diffeomorphism. Let  $\log: G \rightarrow \mathfrak{g}$  denote the inverse of  $\exp$ .

**2.1. Rational structures.** Let  $G$  be a nilpotent, connected and simply connected real Lie group and let  $\mathfrak{g}$  be its Lie algebra. We say that  $\mathfrak{g}$  (or  $G$ ) has a *rational structure* if there is a Lie algebra  $\mathfrak{g}(\mathbb{Q})$  over  $\mathbb{Q}$  such that  $\mathfrak{g} \cong \mathfrak{g}(\mathbb{Q}) \otimes \mathbb{R}$ . It is clear that  $\mathfrak{g}$  has a rational structure if and only if  $\mathfrak{g}$  has an  $\mathbb{R}$ -basis  $(X_1, \dots, X_n)$  with rational structure constants.

**2.2. Uniform subgroups.** A discrete subgroup  $\Gamma$  is called *uniform* in  $G$  if the quotient space  $G/\Gamma$  is compact. A proof of the next result can be found in Theorem 7 of [8] or in Theorem 2.12 of [11].

**Theorem 2.1** (The Malcev rationality criterion). *Let  $G$  be a simply connected nilpotent Lie group, and let  $\mathfrak{g}$  be its Lie algebra. Then  $G$  admits a uniform subgroup  $\Gamma$  if and only if  $\mathfrak{g}$  admits a basis  $(X_1, \dots, X_n)$  such that*

$$[X_i, X_j] = \sum_{k=1}^n c_{ijk} X_k, \quad (\forall 1 \leq i, j \leq n),$$

where the constants  $c_{ijk}$  are all rational.

**2.3. The Malcev rigidity theorem.** The following is a theorem of Mal'cev ([8, Theorem 5]); see also ([9, Theorem 4]).

**Theorem 2.2** (Malcev rigidity theorem). *Let  $G_1$  and  $G_2$  be connected simply connected nilpotent Lie groups and  $\Gamma_1, \Gamma_2$  discrete uniform subgroups of  $G_1$  and  $G_2$ . Any abstract group isomorphism  $\phi$  between  $\Gamma_1$  and  $\Gamma_2$  extends uniquely to an isomorphism  $M(\phi)$  of  $G_1$  on  $G_2$ ; that is, the following diagram*

$$(2.1) \quad \begin{array}{ccc} \Gamma_1 & \xrightarrow{\phi} & \Gamma_2 \\ i \downarrow & & \downarrow i \\ G_1 & \xrightarrow{M(\phi)} & G_2 \end{array}$$

is commutative, where  $i$  is the inclusion mapping. The isomorphism  $M(\phi)$  is called the Mal'cev extension of  $\phi$ .

### 3. An explicit Euler characteristic on TUF. Proof of Theorem 1.5

Let  $G$  be a connected Lie group,  $\mathcal{S}(G)$  be the space of discrete uniform (i.e., cocompact) subgroups of  $G$ . Let  $\mu$  be a right Haar measure of  $G$ . Let  $\Gamma \in \mathcal{S}(G)$ , the measure  $\mu$  induces a finite measure  $\bar{\mu}$  over the homogeneous space  $G/\Gamma$ . Let

$$V_G^\mu: \mathcal{S}(G) \rightarrow \mathbb{R}_+$$

defined for  $\Gamma \in \mathcal{S}(G)$  by

$$V_G^\mu(\Gamma) = \bar{\mu}(G/\Gamma).$$

REMARK 3.1. We recall that if  $F$  is a fundamental domain for  $G/\Gamma$  then  $\bar{\mu}(G/\Gamma) = \mu(F)$  ([7, p. 430]).

The notation  $H \leq_f K$  signifies that  $H$  is a finite index subgroup of the group  $K$ . A proof of the following proposition can be found in Lemma 3.2 of [7].

**Proposition 3.2.** *If  $H, K \in \mathcal{S}(G)$  and if  $H \leq_f K$  then we have*

$$(3.1) \quad V_G^\mu(H) = [K : H]V_G^\mu(K).$$

**Proposition 3.3.** *Let  $G$  in TULG and  $\mu$  a Haar measure on  $G$ . Let  $\Gamma_1, \Gamma_2$  be two isomorphic subgroups of  $\mathcal{S}(G)$ . Then we have*

$$(3.2) \quad V_G^\mu(\Gamma_1) = V_G^\mu(\Gamma_2).$$

Proof. Let  $\phi$  be an isomorphism of  $\Gamma_1$  onto  $\Gamma_2$ . Let  $F$  be a fundamental domain of  $G/\Gamma_1$  and compute

$$\begin{aligned} V_G^\mu(\Gamma_1) &= \mu(F) \\ &= \mu(\mathbf{M}(\phi)(F)) \\ &= V_G^\mu(\Gamma_2). \end{aligned}$$

□

We define an equivalence relation  $\simeq$  on TULG by

$$G_1 \simeq G_2 \iff G_1, G_2 \text{ are isomorphic.}$$

For  $G \in \text{TULG}$ , let  $[G]$  be the equivalence class containing  $G$ . Let  $T$  be a transversal for the equivalence relation  $\simeq$ .

Let  $H, K$  be two groups (resp. Lie groups), the set of all isomorphisms (resp. Lie groups isomorphisms) of  $H$  onto  $K$  is denoted by  $\mathcal{R}(H, K)$ .

**Lemma 3.4.** *Let  $G_0 \in T$  and  $G \in [G_0]$ . For every  $\phi, \psi \in \mathcal{R}(G_0, G)$  we have*

$$\phi_*\mu_0 = \psi_*\mu_0,$$

where  $\phi_*\mu_0$  (resp.  $\psi_*\mu_0$ ) is the push forward of  $\mu_0$  under  $\phi$  (resp.  $\psi$ ).

Proof. Let  $F$  be a measurable set and compute

$$\begin{aligned} \phi_*\mu_0(F) &= \mu_0(\phi^{-1}(F)) \\ &= \mu_0(\psi^{-1}\phi(\phi^{-1}(F))) \quad (\psi^{-1}\phi \in \text{Aut}(G_0)) \\ &= \mu_0(\psi^{-1}(F)) \\ &= \psi_*\mu_0(F). \end{aligned}$$

□

Let  $G_0 \in T$  and  $\mu_0$  a Haar measure on  $G_0$ . Let  $G \in [G_0]$  and  $\Gamma \in \mathcal{S}(G)$ . The function

$$\mathcal{R}(G_0, G) \rightarrow \mathbb{R}, \quad \phi \rightarrow V_G^{\phi_*\mu_0}(\Gamma)$$

is constant. In the sequel, we note

$$V[G_0, \mu_0, G](\Gamma) = V_G^{\phi_* \mu_0}(\Gamma) \quad (\forall \phi \in \mathcal{R}(G_0, G)).$$

Let  $G_1, G_2 \in [G_0]$ . For every  $\psi \in \mathcal{R}(G_0, G_2)$  and  $\phi \in \mathcal{R}(G_1, G_2)$ , we note

$$V[G_0, \mu_0, G_1] * \phi = V[G_0, \mu_0, G_2] \circ \phi^*,$$

$$\psi * V[G_0, \mu_0, G_1] = V[\psi(G_0), \psi_* \mu_0, G_1],$$

where  $\phi^* : \mathcal{S}(G_1) \rightarrow \mathcal{S}(G_2)$ ,  $\Gamma \mapsto \phi(\Gamma)$ .

**Proposition 3.5.** *With the same notation as above we have:*

$$(3.3a) \quad V[G_0, \mu_0, G_1] * \phi = V[G_0, \mu_0, G_1],$$

$$(3.3b) \quad \psi * V[G_0, \mu_0, G_1] = V[G_0, \mu_0, G_1].$$

Proof. Let  $\Gamma \in \mathcal{S}(G_1)$ . Let  $F$  be a fundamental domain for  $G_1/\Gamma$  and compute

$$\begin{aligned} V[G_0, \mu_0, G_1] * \phi(\Gamma) &= V[G_0, \mu_0, G_2](\phi(\Gamma)) \\ &= V_{G_2}^{\phi_* \mu_0}(\phi(\Gamma)) \quad (\phi \in \mathcal{R}(G_0, G_2)) \\ &= \phi_* \mu_0(\phi(F)) \\ &= (\phi^{-1} \phi)_* \mu_0(F) \\ &= V_{G_1}^{(\phi^{-1} \phi)_* \mu_0}(F) \\ &= V[G_0, \mu_0, G_1](\Gamma). \end{aligned}$$

Similarly, we prove (3.3b). □

We come now to the principal theorem of this paper, in which we give an explicit Euler characteristic on TUF.

**Theorem 3.6.** *The mapping*

$$\chi : TUF \rightarrow \mathbb{R}, \quad \Gamma \mapsto V[G_0, \mu_0, G_\Gamma](\Gamma),$$

where  $\{G_0\} = [G_\Gamma] \cap T$ , is an Euler characteristic on TUF, which does not depend of the choice of transversal.

Proof. Let  $\Gamma$  be a TUF-group and  $\Gamma_0 \leq_f \Gamma$ . Then  $\Gamma_0$  and  $\Gamma$  have the same Hirsch length, it follows that  $\mathfrak{g}_{\Gamma_0} = \mathfrak{g}_\Gamma$  and hence the class TUF is closed under taking subgroups of finite index. Let  $\Gamma_1$  and  $\Gamma_2$  be two isomorphic TUF-groups. By Theorem 2.2,

the Lie groups  $G_{\Gamma_1}$  and  $G_{\Gamma_2}$  are isomorphic and hence  $[G_{\Gamma_1}] = [G_{\Gamma_2}]$ . Let

$$\{G_0\} = [G_{\Gamma_1}] \cap T.$$

Let  $\phi \in \mathcal{R}(\Gamma_1, \Gamma_2)$  and compute

$$\begin{aligned} \chi(\Gamma_2) &= V[G_0, \mu_0, G_{\Gamma_2}](\Gamma_2) \\ &= V[G_0, \mu_0, G_{\Gamma_1}] * M(\phi)(\Gamma_1) \\ &= V[G_0, \mu_0, G_{\Gamma_1}](\Gamma_1) \quad (\text{by (3.3a)}) \\ &= \chi(\Gamma_1). \end{aligned}$$

This completes the proof of (Ec1). (Ec2) follows from (3.1). Finally, the formula (3.3b) implies that the mapping  $\chi$  is independent of the choice of the transversal  $T$ .  $\square$

---

### References

- [1] I. Belegradek: *On co-Hopfian nilpotent groups*, Bull. London Math. Soc. **35** (2003), 805–811.
- [2] I.M. Chiswell: *Euler characteristics of groups*, Math. Z. **147** (1976), 1–11.
- [3] L.J. Corwin and F.P. Greenleaf: *Representations of Nilpotent Lie Groups and Their Applications, Part I*, Cambridge Studies in Advanced Mathematics **18**, Cambridge Univ. Press, Cambridge, 1990.
- [4] J. Dixmier and W.G. Lister: *Derivations of nilpotent Lie algebras*, Proc. Amer. Math. Soc. **8** (1957), 155–158.
- [5] A. Karrass, A. Pietrowski and D. Solitar: *An improved subgroup theorem for HNN groups with some applications*, Canad. J. Math. **26** (1974), 214–224.
- [6] G. Leger and S. Tôgô: *Characteristically nilpotent Lie algebras*, Duke Math. J. **26** (1959), 623–628.
- [7] A.M. Macbeath and S. Świerczkowski: *Limits of lattices in a compactly generated group*, Canad. J. Math. **12** (1960), 427–437.
- [8] A.I. Malcev: *On a class of homogeneous spaces*, Amer. Math. Soc. Translation **1951** (1951).
- [9] Y. Matsushima: *On the discrete subgroups and homogeneous spaces of nilpotent Lie groups*, Nagoya Math. J. **2** (1951), 95–110.
- [10] L.P. Polek: *Small neighborhoods of the identity of a real nilpotent group*, Proc. Amer. Math. Soc. **42** (1973), 627–630.
- [11] M.S. Raghunathan: *Discrete Subgroups of Lie Groups*, Springer, New York, 1972.
- [12] A. Reznikov: *Volumes of discrete groups and topological complexity of homology spheres*, Math. Ann. **306** (1996), 547–554.
- [13] G.C. Smith: *Compressibility in nilpotent groups*, Bull. London Math. Soc. **17** (1985), 453–457.
- [14] S. Yamaguchi: *Derivations and affine structures of some nilpotent Lie algebras*, Mem. Fac. Sci. Kyushu Univ. Ser. A **34** (1980), 151–170.
- [15] C.T.C. Wall: *Rational Euler characteristics*, Proc. Cambridge Philos. Soc. **57** (1961), 182–184.

Department of Mathematics  
Faculty of Sciences at Sfax  
University of Sfax  
Route Soukra, B.P. 1171, 3000 Sfax  
Tunisia  
e-mail: [hatemhhamrouni@voila.fr](mailto:hatemhhamrouni@voila.fr)