

Title	Multi-level functional redundancy mechanisms of Saccharomyces cerevisiae in response to high extracellular calcium stress			
Author(s)	Laviña, Walter Alvarez			
Citation	大阪大学, 2013, 博士論文			
Version Type	VoR			
URL	https://doi.org/10.18910/26205			
rights				
Note				

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

**Doctoral Dissertation** 

### Multi-level functional redundancy mechanisms of *Saccharomyces cerevisiae* in response to high extracellular calcium stress

Walter Alvarez Laviña July 2013

Graduate School of Engineering, Osaka University

### Abstract

Upon the completion of the Yeast Genome Project in 1996, *Saccharomyces cerevisiae* became the first eukaryotic genome to be sequenced fully; defining 5,885 potential proteinencoding genes. More importantly, the complete sequence revealed the magnitude of research that was needed to be done as more than 50% of the sequenced genes were unknown and uncharacterized by previously established methods such as mutant hunting. Furthermore, the yeast genome showed a large amount of genes with apparent genetic redundancy. This particular problem necessitated the need to systematically characterize the biological function of the all genes in the yeast genome and to elucidate functional redundancy in more detail.

Sakumoto et al. (2002) have previously reported that the disruption of two protein phosphatase genes, *PTP2* and *MSG5*, caused calcium sensitivity indicating that functional redundancy exists between the two protein phosphatases in response to high extracellular calcium. Furthermore, Hermansyah et al. (2010) found that additional disruption of protein kinases *BCK1*, *MKK1*, *SLT2*, *MCK1*, *YAK*1 and *SSK2* in the *ptp2*\Delta*msg5*\Delta background conferred calcium tolerance. In this work, the suppression mechanisms conferred by the different protein kinase disruptions are described. First, it was found that the inactivation of calcineurin by the disruption of *CNB1* or treatment with FK506 can suppress the calcium sensitive phenotype of the *ptp2*\Delta*msg5*\Delta double disruptant. In the wake of a calciuminduced, calcineurin-driven signaling pathway activation, the calcium sensitivity of the *ptp2*\Delta*msg5*\Delta double disruptant can be suppressed by regulating the SLT2 pathway through the disruption of the major kinases in the SLT2 signal cascade that include *BCK1*, *MKK1* and *SLT2*. Also, this study showed that *PTP2* and *MSG5* are key regulatory phosphatases that prevent over-activation of the calcium-induced signaling cascade under the parallel control of the SLT2 and calcineurin pathways. Next, genetic analysis showed a HOG-pathway

independent function for Ssk2 in response to high extracellular calcium. Genome-wide transcriptional analysis found 19 induced genes in the calcium-sensitive  $ptp2\Delta msg5\Delta$  double disruptant that were subsequently repressed in the calcium-tolerant  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant ("rise and fall" patterned expression). Also, this study reports new suppressors,  $bcy1\Delta$  and  $msn2\Delta$ , which are related to the previously determined suppressors  $ssk2\Delta$  and  $yak1\Delta$  via the regulation of the "rise and fall" genes. This study provides evidence to the existence of several alternative pathways that mediate calcium signaling in the  $ptp2\Delta msg5\Delta$ double disruptant.

### **Table of Contents**

		Subject	Page
Abstract			2
Chapter 1		General introduction	7
1.1		The Yeast Genome Project and functional redundancy in yeasts	7
1.2		Cellular response to environmental pertubation and stress	8
1.3		Reversible phosphorylation dynamics (Protein kinases and phosphatases	8
1.4		Ptp2 and Msg5 protein phosphatases	12
1.5		Calcineurin and calcium homeostasis	13
1.6		SLT2 pathway	14
1.7		HOG pathway	14
1.8		cAMP-protein kinase A and Yak1	15
1.9		Environmental Stress Response (ESR) and transcription factors Msn2 and Msn4	16
1.10		Objective of the study	17
Chapter 2		Functionally redundant protein phosphatase genes <i>PTP2</i> and <i>MSG5</i> co-regulate the calcium signaling pathway in <i>Saccharomyces cerevisiae</i> upon exposure to high extracellular calcium concentration	19
2.1		Introduction	19
2.2		Materials and methods	22
2	2.2.1	Strains and media	22
2	2.2.2	Genetic manipulations	23
2	2.2.3	RNA isolation	23
2	2.2.4	Microarray analysis	23

		Subject	Page
	2.2.5	mRNA level determination using quantitative PCR	24
	2.2.6	Phenotypic analysis	25
	2.2.7	Fluorescence microscopy	25
2.3		Results	30
	2.3.1	An active calcineurin pathway is inhibitory to the growth of the calcium-exposed $ptp2\Delta msg5\Delta$ double disruptant	30
	2.3.2	Up-regulated cell wall genes in the $ptp2\Delta msg5\Delta$ double disruptant share common transcription factors Sok2 and Ste12	35
	2.3.3	SLT2 pathway is hyper-activated in the $ptp2\Delta msg5\Delta$ double disruptant upon exposure to high extracellular calcium	37
	2.3.4	Calcium sensitivity in the $ptp2\Delta msg5\Delta$ double disruptant results from a hyper-activated, calcium-induced signaling pathway caused by the improper activation of the SLT2 pathway and an active calcineurin.	38
2.4		Discussion	40
Chapter	3	Suppression mechanism of the calcium sensitivity in Saccharomyces cerevisiae $ptp2\Delta msg5\Delta$ double disruptant involves a novel HOG-independent function of Ssk2, transcription factor Msn2 and the PKA component Bcy1	47
3.1		Introduction	47
3.2		Materials and methods	49
	3.2.1	Strains, media and general methods	49
	3.2.2	Protein isolation and immunoblot analysis	49
	3.2.3	RNA isolation and microarray analysis	50
	3.2.4	Gene enrichment analysis using Genowiz	50
	3.2.5	Determination of mRNA level using quantitative PCR	51
3.3		Results	55

		Subject	Page
3	3.3.1	The PTP2 and MSG5-mediated calcium response involves a novel function for Ssk2, a protein kinase component of HOG pathway	55
3	3.3.2	19 induced genes in the calcium sensitive $ptp2\Delta msg5\Delta$ double disruptant are repressed in the calcium tolerant $ptp2\Delta msg5\Delta ssk2\Delta$ triple disruptant	56
3	3.3.3	"Rise and fall" genes share several transcription factors that may be involved in the suppression of the calcium sensitive phenotype of the $ptp2\Delta msg5\Delta$ double disruptant by <i>SSK2</i> disruption	61
3	3.3.4	Disruption of $msn2\Delta$ confers calcium tolerance to the $ptp2\Delta msg5\Delta$ double disruptant	62
3	3.3.5	Constitutive activation of PKA suppresses the calcium sensitive phenotype of the $ptp2\Delta msg5\Delta$ double disruptant	64
3.4		Discussion	67
Chapter 4		Conclusion and general discussion	71
References			74
Publications			85
Acknowledgeme	ents		86

### Chapter 1

### **General Introduction**

### 1.1 The Yeast Genome Project and functional redundancy in yeasts

In 1990, the Human Genome Project (HGP) was initiated by the US Department of Energy (DOE) and National Institutes of Health (NIH) with the primary goal of sequencing, identifying and mapping the entire 20,000 to 25,000 genes of the human genome (www.genome.gov). Several genome projects paralled from this international and collaborative research initiative using model organisms such as *Escherichia coli*, *Saccharomyces cerevisiae*, *Caenorhabditis elegans*, *Drosophila* sp. and *Arabidopsis thaliana* (www.nigms.nih.gov). In 1996, the genome of the yeast *S. cerevisiae* was completely sequenced with 12,068 kilobases defining 5885 potential protein-encoding genes, (including 140 ribosomal RNA genes, 40 genes specifying for small nuclear RNA molecules and 275 transfer RNA genes). (Goffeau et al., 1996). Aside from the fact that the *S. cerevisiae* was the first eukaryotic genome to be fully sequenced, yeast has been considered an ideal model organism to study biological phenomena due to its intrinsic advantages as an experimental system such as an ideal life cycle perfectly suited for genetic analysis and ease of genetic manipulation.

Although *S. cerevisiae* has been one of the best characterized organisms from the genetics and physiological point of view, results of the Yeast Genome Project revealed the magnitude of research that was still needed to be done as more than 50% of the sequenced yeast genes were unknown and uncharacterized by previously established methods such as mutant hunting (Piskur and Langkjaer, 2004). Such considerable amount of uncharacterized genes was accounted to genetic redundancy, which was believed to be responsible for cellular robustness (Goffeau et al., 1996; Li et al., 2010). This unexpected discovery necessitated the

need to systematically characterize the biological function of the all genes in the yeast genome and to elucidate functional redundancy in more detail.

### 1.2 Cellular response to environmental perturbation and stress

The ability to respond to changes in environmental conditions for the purpose of cell protection or damage repair is an intrinsic property of the cell, regardless whether it is prokaryotic or eukaryotic. To achieve this, cells require a complex network of sensory and signal transduction mechanisms that lead to adaptations in cell growth and proliferation, usually by modification in gene expression and metabolic activities (Hohmann and Mager, 2002). Specifically, stress refers to the state of environment that is not optimal and often detrimental to the survival of cells. When cells encounter drastic changes in the environments, survival largely depends on maintaining cellular homeostasis thus, cells developed various ways of keeping the cellular state in equilibrium. For example, in the advent of stress response, cell wall integrity and the calcineurin pathway is observed (Klis et al., 2006). Conversely, it is also essential for the cells to possess alternative "fail-safe" mechanisms in case of improper activation of these pathways that can be detrimental to the organism.

### **1.3 Reversible phosphorylation dynamics (Protein kinase and phosphatases)**

Cell signaling plays a very crucial role in the survival of organisms where a slight change in the cell dynamics can be detrimental. Hence, organisms evolved to have several parallel pathways to prevent unwanted activation or deactivation of signaling cascades. Furthermore, each signaling cascade possesses several mechanisms of regulation, ranging from transcriptional to post-translational. Reversible phosphorylation is one among the many regulatory mechanisms that cells employ to mediate cell signaling. Under this regulation,

proteins are phosphorylated with protein kinases (PKases) while protein phosphatases (PPases) are responsible for dephosphorylation. This process plays a key role in adaptation to environmental stress and cell proliferation (Martin et al., 2005).





In *S. cerevisiae*, there are 129 PKases and 40 PPases among approximately 6,600 genes that are involved in various aspects of signaling and other cellular aspects (www.yeastkinome.org; Sakumoto et al., 2002). Yeast protein phosphatases are divided into four groups namely: 1) serine threonine PPases, 2) protein tyrosine PPases, 3) dual specificity PPases and PTEN and myotubularin PPases. These groups of proteins have their corresponding homologs in humans and other higher eukaryotes although the number in each group is much larger.

Family/		
Subfamily	ORF/ Gene	Function
PPP family		
PP1 subfamily	YER133w/GLC7	Involved in glycogen metabolism, sporulation and
		mitosis
	YML016c/PPZ1	Involved in salt homeostasis
	YDR436w/PPZ2	Involved in salt homeostasis
	YPL179w/SAL6	Regulation of protein synthesis
PP2A subfamily	YDL047w/SIT4	Cell cycle
	YDL134c/PPH21	Cell cycle, cell wall maintenance
	YDL188c/PPH22	Highly homologous to Pph21p
	YDR075w/PPH3	Cell structure
	YNR032w/PPG1	Involved in glycogen accumulation
PP2B subfamily	YLR433c/CNA1	Some ion transport, cell polarity, mating response
	YML057w/CNA2	Some ion transport, mating response
Other	YGR123c/PPT1	Human PP5-related PPase
	YLL010c/PSR1	Response to stress
	YLR019w/PSR2	Functionally redundant Psr1p homolog
	YMR277w/FCP1	Carboxy-terminal domain (CTD) phosphatase
	YLR361c/DCR2	Interacts with Sic1p, an inhibitor of mitotic
		cyclin/cyclin-dependent kinase complexes.
	YHR004c/NEM1	Regulates nuclear growth
	YNL222w/SSU72	Phosphatase and transcription/RNA processing factor
PPM family		
PP2C subfamily	YDL006w/PTC1	Down-regulate the HOG pathway
	YER089c/PTC2	Cdc28p dephosphorylation, involved in HOG pathway
	YBL056w/PTC3	Cdc28p dephosphorylation, Involved in HOG pathway
	YBR125c/PTC4	Cytoplasmic type 2C PPase
	YOR090c/PTC5	Involved in regulation of pyruvate dehydrogenase activity
	YCR079w/PTC6	Mitochondria type 2C PPase with similarity to mammalian PPk1s
	YHR076w/PTC7	Mitochondrially localized type 2C PPase

### Table 1. Protein phosphatase (PPase) genes present in S. cerevisiae

PTP family				
PTP subfamily	YDL230w/PTP1	Protein tyrosine phosphatase		
-	YDR208w/PTP2	Down-regulate Hog1p and Fus3p MAPK, sporulation		
	YER075c/PTP3	Down-regulate Hog1p and Fus3p MAPK, sporulation		
	YPR073c/LTP1	Similar to PPase from human placenta		
	YNL032w/SIW14	Involved in nutritional control of the cell cycle		
	YMR036c/MIH1	S.pombe cdc25p homologue		
	YJR110w/YMR1	Phosphatidylinositol 3-phosphate phosphatase		
	YNL128w/TEP1	Similar to human tumor suppressor gene		
DSP subfamily	YFR028c/CDC14	Function at the late stage of the cell cycle		
-	YBR276c/PPS1	A role in the DNA synthesis phase of the cell cycle		
	YIR026c/YVH1	Yeast homologue of Vaccinia virus PTP, VH1		
	YNL053w/MSG5	Dephosphorylate Fus3p		
	YIL113w/SDP1	Negatively regulates Slt2p MAPK		
	YNL099c/OCA1	Protein tyrosine/serine/threonine phosphatase activity		
Other	YNR002c/ATO2	Weak similarity to PPase		

Elucidation of the protein functions of PPases and PKases contribute greatly to the understanding of various signaling pathways in the cell, paving the way to the clarification of the nature of diverse unknown cellular phenomena. Although knowledge about PKases and PPases has steadily grown over the past years, there are still a fair number of PPases that needs to be addressed. Between the two proteins, analysis of the functional redundancy of PPases seemed more practical thus it was undertaken. In our laboratory, a library of 30 PPase single disruptants and double disruptants in all 435 possible combinations was created to identify the role of every phosphatase gene in various cellular functions of S. cerevisiae. Systematic genetic screening led to the discovery of several interesting phenotypes such as temperature sensitivity, caffeine sensitivity and cation sensitivity. Through this work, we found that disruption of both PTP2 and MSG5 in S. cerevisiae causes calcium sensitivity, while single disruption of PTP2 or MSG5 does not (Sakumoto et al., 1999 & 2002). The synthetic phenotype displayed by the  $ptp2\Delta msg5\Delta$  double disruptant suggests a functional redundancy existing between PTP2 and MSG5 in response to high extracellular calcium stress. We were intrigued with the first report of the involvement of redundant function of established MAPK regulators, Ptp2 and Msg5, in the calcium-induced signaling pathway

thus, the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant was further studied.

### 1.4 Ptp2 and Msg5 protein phosphatases

PTP2 (YOR208W) encodes a 750-amino acid residue, 86 KDa protein tyrosine phosphatase that has 36-46% amino acid sequence similarity with at least 35 mammalian PPases that includes PTPRC (PTP receptor type T) and PTPRT (PTP receptor type T) (Guan et al., 1992; Wang et al., 2004). On the other hand, MSG5 (YNL053W) codes for a 489amino acid residue, 54 KDa dual-specificity protein phosphatase that has 42-72% amino acid sequence similarity with at least 27 mammalian PPases that includes MKP5, DUSP26 and DUSP3 (Guan et al., 1992; Rahmouni et al., 2006). Although Ptp2p and Msg5p are both included in the PTP family of PPases and contain an active-site signature motif HCX5R, Ptp2p dephosphorylates phosphotyrosine residues while Msg5 acts on both phosphoserine/phosphothreonine and phosphotyrosine residues (Guan et al., 1992; Doi et al., 1994; Sakumoto et al., 1999; 2002). Previous genetic and biochemical studies implicated protein tyrosine phosphatases (PTPases) with the negative regulation of MAP kinases (Zhan et al., 2000). Conventionally, Ptp2, a phosphotyrosine-specific PPase, is important in the inactivation of Hog1 MAPK during high osmolarity (Wurgler-Murphy et al., 1997) and heat stress (Winkler et al., 2002) while Msg5, a dual-specificity PPase, is involved in the regulation of the mating response pathway (Doi et al., 1994; Blackwell et al., 2007) and cell integrity pathway (Flandez et al., 2004). However, there has been evidence that PTP2 and MSG5 also play important roles in calcium signaling in yeasts (Sakumoto et al., 1999).



Figure 2. MAPK cascades in S. cerevisiae (Adapted from Martin et al., 2005)

### 1.5 Calcineurin and calcium homeostasis

Adaptation to changing environments require cells to activate signaling pathways that require messengers whose concentration varies with time (Clapham, 2007). Calcium ion, which is the most ubiquitous and versatile signaling molecule in the eukaryotic cell, fit this specific role perfectly thus making it arguably the most important signaling molecule in the cell. In *S. cerevisiae*, exposure to calcium activates the Ca<sup>2+/</sup> calmodulin-dependent protein phosphatase calcineurin that regulates the Cr21p transcription factor (Yoshimoto et al., 2002). Calcineurin is a heterodimeric protein comprised of a catalytic (A) and regulatory (B) sub-units that mediates the nuclear localization of the transcription factor Cr21. Furthermore, calcineurin is specifically inhibited by immunosuppressant drugs cyclosporin A and tacrolimus (FK506) thus making it a very valuable protein for clinical studies (Sugiura et al., 2002; Cyert 2003). Under normal conditions when cytosolic Ca<sup>2+</sup> levels are low, calcineurin is inactive but in response to stress, there is a rise in the cytosolic Ca<sup>2+</sup> levels that activates calcineurin, subsequently dephosphorylating several key proteins to promote survival (Cyert, 2003). One important function of calcineurin-dependent signaling is the

regulation of the transcription factor Crz1. Crz1p directs a calcium-induced, calcineurin dependent gene expression by binding to calcineurin-dependent response element (CDRE) (Stathopoulos et al., 1997). Calcineurin-induced activation leads to up-regulation of 163 genes including *GSC2*, *PMC1*, *PMR1* and *ENA1* (Yoshimoto et al., 2002). Calcineurin and Crz1-regulated genes belong to various functional classes such as integral membrane proteins, plasma membrane and cell wall components, lipid synthesis, protein degradation and vesicle trafficking (Cyert, 2003).

### 1.6 SLT2 pathway

The Slt2/Mpk1 mitogen-activated protein kinase (MAPK) cell integrity pathway is responsible for the monitoring and proper response to stresses that challenge the integrity of the cell wall as well as the maintenance of cell shape and integrity during vegetative growth and mating in *S. cerevisiae* (Jung and Levin, 1999; Hahn and Thiele, 2002). Slt2 pathway sensors Wsc1 and Mid2 interact with Rom2 to trigger Rho1 GTP loading which then activates Pkc1. Afterwards, there is a sequential activation of the MAPK cascade consisting of MAPK kinase kinase Bck1, MAPK kinase Mkk1/Mkk2 and MAP kinase Skt2 (Cid et al., 1995; Levin , 2005).

### 1.7 HOG pathway

Upon exposure to increased osmolarity, S. cerevisiae responds by activating the High Osmolarity Glycerol (HOG) MAPK pathway. The cascade is made up of three sequenitially activating kinases; the redundant MAPK kinase kinases, Ssk2 and Ssk22, MAPK kinase Pbs2 and the MAP kinase Hog1 (Saito and Tatebayashi, 2004). Three upstream, independent osmosensors regulate the HOG pathway namely the membrane-spanning proteins Sln1p and Sho1p and a third Msb2 branch (O'Rourke and Herskowitz, 2002). The Sln1 transmembrane

protein contains a cytoplasmic histidine kinase (HK) and receiver (Rec) domains and has strong similarities with the bacterial two-component signal transducers (Saito and Tatebayashi, 2004). The phosphate is transferred from Sln1 to Ypd1 via a phospho-relay mechanism until it reaches another Rec domain protein, Ssk1 (Posas et al., 1996). Upon exposure to hyper-osmotic conditions, Ssk1 is rapidly dephosphorylated by Ypd1 and binds the functionally redundant Ssk2/Ssk22 MAPKK kinases. Ssk2/Ssk22 in turn activates Pbs2 MAPKK. On the other hand, the SHO1 branch involves a transient formation of cell surface protein complexes consisting of at least Sho1p and Pbs2p (Reiser et al., 2000). Cdc24, Cdc42, Ste20, Ste50 and Ste11 are also reported to have involvement in this signaling cascade (Hohmann, 2002). The signals from each branch converge to Pbs2 which in turn activates Hog1. Activation of the key element Hog1 through the phosphorylation of the conserved Thr174 and Tyr176 residues results to a change in the transcriptional pattern in response to osmotic stress; which includes the expression of several genes such as *GPD1, CTT1*, and *HSP12* (Wurgler-Murphy et al., 1997).

#### 1.8 cAMP-protein kinase A Pathway and Yak1

In *S. cerevisiae*, cAMP-Protein Kinase A (PKA) pathway plays a major role in the regulating cell growth, metabolism and stress resistance (Tamaki, 2007). PKA is essential for cell cycle progression since the absence of PKA activity results in cell arrest at G1 and growth inhibition (Garrett et al., 1991). PKA is composed of a catalytic sub-unit encoded by three genes (*TPK1, TPK2* and *TPK3*) and a regulatory sub-unit encoded by *BCY1* (Thevelein and de Winde, 1999). Although the main function of the cAMP-PKA pathway is for nutrient sensing, it was also reported to control the expression of around 30 STRE (stress-responsive element) genes via the repression of the transcription factors Msn2 and Msn4 (Wilson and Roach, 2002).

*YAK1* encodes for a serine-threonine protein kinase and a known component of the glucose-sensing system initially identified as a suppressor of the lethal phenotype associated with loss of RAS function or loss of the catalytic unit of PKA (Pratt et al., 2007). Yak1 functions downstream of PKA and is a negative regulator of growth.

### 1.9 Environmental Stress Response (ESR) and transcription factors Msn2 and Msn4

Due to the constant fluctuations in environmental conditions, cells are prone to changes in their internal milieu that can disrupt normal cellular processes. Therefore, cells must maintain their internal system at a stable state to maintain viability. A mechanism employed by yeast to protect their internal system from drastic changes is the environmental stress response (ESR). ESR includes approximately 900 genes that are normally altered transcriptionally when exposed to a stressful environment (Martinez-Pastor et al., 1996; Gasch, 2002). Specifically, ESR response consists of around 300 induced genes related to stress response and 600 repressed genes related to protein synthesis (Berry and Gasch, 2008). Although the simultaneous expression of this group of genes is the main characteristic of ESR, regulation of the gene expression is gene-specific and condition-specific (Gasch, 2002).

The best characterized regulators of ESR gene expression are the transcription factors Msn2p and Msn4p. These transcription factors specifically bind to the STRE promoter sequence and induce the transcription of ESR genes in response to stress. Even though both factors were initially thought to be redundant, Msn2 and Msn4 appear to respond differently based on the genotype and condition with Msn2 being regarded as the stronger transcription factor (Hohmann, 2002). Furthermore, results by Berry and Gasch (2008) indicated that Msn2 and Msn4 play distinct roles in response to various types of stresses. However, both Msn2 and Msn4 are required by many genes for full induction (Berry and Gasch, 2008).

### 1.10 Objective of the study

In an effort to study the functional redundancy of protein phosphatases in yeast, Sakumoto and colleagues (2002) constructed a PPase library of single and double disruptants and systematically screened for interesting phenotypes. One such phenotype was the calcium sensitivity exhibited by the strain with the disruption of both PTP2 and MSG5 but not the single disruptants  $ptp2\Delta$  or  $msg5\Delta$  (Sakumoto et al., 1999; 2002). Furthermore, disruption of an additional protein kinase in the *ptp2\Deltamsg5\Delta* double disruptant revealed *bck1\Delta*, *mkk1\Delta* and *slt2* $\Delta$  of the Cell Wall Integrity pathway, *mck1* $\Delta$  which is involved in chromosome segregation and meiosis,  $yakl\Delta$  of the glucose sensing pathway and  $ssk2\Delta$  of the HOG pathway as suppressors of the calcium sensitive phenotype (Hermansyah et al., 2010). Interestingly, since the PKase suppressors belong to various signaling pathways, this indicates that the Ptp2 and Msg5-mediated calcium signaling pathway involves a complex interaction between various signal cascades. Thus, the purpose of this study is to elucidate the suppression mechanism/s of the calcium-sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant conferred by the different PKase suppressors and discover their interrelationships. In addition, we searched for other genes that may be related to the calcium-induced signaling mediated by PPases Ptp2 and Msg5.

In Chapter 1 of this dissertation, the importance of cell signaling and reversible phosphorylation on the cell's response to the environment was described. In addition, several important signaling cascades are outlined namely the calcineurin, SLT2, HOG and PKA pathways. In Chapter 2, the suppression of the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant via the inactivation of calcineurin either by the disruption of the calcineurin regulatory subunit, *CNB1* or treatment with a calcineurin inhibitor, FK506 was described. Also, evidence of a functional redundancy at the cascade level between SLT2 and calcineurin pathways in response to high extracellular calcium was shown. In this model,

the roles of Bck1, Mkk1 and Slt2 in the calcium-induced, Ptp2 and Msg5-mediated signaling was clarified. Chapter 3 described two new suppressors, *msn2* $\Delta$  and *bcy1* $\Delta$  that are genetically related to the previously identified suppressor *ssk2* $\Delta$  via the regulation of a specific gene cluster. Moreover, a novel, HOG-independent suppressor function of Ssk2 and 19 genes with "rise and fall" pattern of expression that is likely involved in the calcium phenotype of the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant were identified. Results suggest the interrelationship of a HOG-independent function of Ssk2, transcription factor Msn2, PKA-related protein Bcy1 and 19 "rise and fall" genes as responsible for the suppression mechanism of the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant by *ssk2* $\Delta$  disruption. The conclusion and general discussion was described in Chapter 4.

### Chapter 2

# Functionally redundant protein phosphatase genes *PTP2* and *MSG5* co-regulate the calcium signaling pathway in *Saccharomyces cerevisiae* upon exposure to high extracellular calcium concentration

### **2.1 Introduction**

Upon exposure to stress, *S. cerevisiae* employs a variety of signaling cascades such as the mitogen-activated protein kinase (MAPK) pathways that control patterns of gene expression and protein activity to cope with the deleterious changes in the environment (Gustin et al., 1998). The ability of the cells to correctly perceive and respond to the environment is critical to the organism's survival thus; inappropriate activation of cell signaling pathways often has dire consequences on the viability of the cell. Accordingly, regulation is an integral part of the signaling cascade architecture that ensures the maintenance of cellular homeostasis upon fluctuations in environmental conditions. Since over-activation of a particular signal can have damaging effects, cells possess a myriad of ways to combat improper activation of signaling pathways, one of which is regulation via protein phosphorylation and dephosphorylation catalyzed by protein kinases (PKases) and protein phosphatases (PPases) respectively.

In *S. cerevisiae*, high extracellular levels of calcium usually elicit the activation of calcineurin, a calcium/calmodulin-dependent serine/threonine PPase (Cyert, 2003). Calcineurin is known to regulate Ca<sup>2+</sup> pumps and exchangers responsible for calcium homeostasis in yeast to maintain the cytoplasmic concentration in the range of 100-300 nM (Cunningham and Fink, 1994; Davis, 1995). These include the vacuolar H<sup>+</sup> ATPases encoded by *VCX1* (5), vacuolar Ca<sup>2+</sup> ATPase Pmc1p (Cunningham and Fink, 1994) and high affinity Ca<sup>2+</sup>/Mn<sup>2+</sup> P-type ATPase Pmr1p (Rudolph et al., 1989). A rigid control of the

intracellular Ca<sup>2+</sup> concentration is an important feature of calcium signaling in which transient alteration of the cytosolic calcium concentration leads to the activation of several signal transduction pathways (Muller et al., 2001). In the regulation of these signaling pathways, calcineurin plays various physiological roles including recovery from  $\alpha$ -factor-induced growth arrest, salt and temperature tolerance, calcium and ion homeostasis, regulation of cell wall biogenesis and Mn<sup>2+</sup> tolerance (Stark, 1996; Miyakawa and Mizunuma, 2007).



Fig. 3. Calcineurin/Crz1 signaling pathway in *S. cerevisiae* (Adapted from Yoshimoto et al., 2002)

We have previously reported that the disruption of two PPases namely *PTP2* and *MSG5* has a detrimental effect on the growth of yeast in a calcium-rich medium. Furthermore, the fact that  $ptp2\Delta$  and  $msg5\Delta$  single disruptants were calcium tolerant implies that there is functional redundancy existing between the two PPases in response to high extracellular calcium (Sakumoto et al., 2002). The calcium-sensitive  $ptp2\Delta msg5\Delta$  double disruptant was further described to exhibit G1 delay, reduced *CLN2* expression and vacuole fragmentation (Hermansyah et al., 2009). In addition, we identified six PKases namely *BCK1*, *MKK1*, *SLT2*, *MCK1*, *YAK1* and *SSK2* whose additional disruption in the  $ptp2\Delta msg5\Delta$  double

disruptant background confers calcium tolerance (Hermansyah et al., 2010). Interestingly, these suppressors fall into two groups based on their capacity to restore cell cycle progression as shown by FACS analysis. Disruption of *BCK1*, *MKK1*, *SLT2* or *MCK1* in the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant was unable to alleviate the G1 transition defect while *SKK2* or *YAK1* disruption were able to initiate G1 to S transition, implying that there are at least two mechanisms governing the suppression of the calcium sensitive phenotype of the *ptp2* $\Delta$ *msg5* $\Delta$ double disruptant. Although Msg5 and Ptp2 are known to be regulators of the *SLT2* pathway (Martin et al., 2005), their specific involvement in the calcium-induced signaling related to growth is not yet fully understood.

In this chapter, we show the importance of the functionally redundant *PTP2* and *MSG5* in the growth of *S. cerevisiae* in high extracellular calcium conditions. Furthermore, we found another suppressor of the calcium sensitive phenotype of the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant, *cnb1* $\Delta$ , which for the first time implicates the calcineurin pathway in the calcium-induced signaling pathway regulated by the functionally redundant *PTP2* and *MSG5*. We also describe how growth is inhibited in the calcium-exposed *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant in relation to calcineurin and SLT2 pathways. Lastly, we explain the mechanism of suppression conferred by the previously discovered suppressors (*bck1* $\Delta$ , *mkk1* $\Delta$  and *slt2* $\Delta$ ) of the calcium sensitivity of the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant in relation to the redundant regulatory function exhibited by SLT2 and calcineurin pathways in cell processes related to growth.

### 2.2 Materials and methods

### 2.2.1 Strains and media

Yeast strains used in this study are listed in Table 2. FY833 or SH5209 (MATa ura3-52 his3- $\Delta 200 \ leu 2\Delta 1 \ lys 2\Delta 202 \ trp 1\Delta 63$ ) was used as the wild type strain. All experiments using yeast strains were conducted at 30°C using standard recipes for YPDA (Sigma-Aldrich Co., St. Louis, MO, USA), SC dropout and Sporulation medium (SPM). YPDA was supplemented with 0.6 M CaCl<sub>2</sub> for the calcium phenotype test while 0.3M CaCl<sub>2</sub> was used for total RNA isolation. For the calcineurin phenotype test, YPDA was added with 1 µg/ml FK506 (a calcineurin inhibitor). Escherichia coli strains used as sources of plasmids namely BYP1804 for p1804 (CgHIS3), BYP1805 for p1805 (CgTRP1), BYP1419 for p1807 (CgLEU2), SHB3276 for p3276 (ScURA3), BYP2446 for p2446 (GFP-CgTPRP1) were acquired from NBRP, YGRC, Japan and cultivated in LB medium containing 100 [g/ml ampicillin at 37°C as previously described (Sambrook et al., 1989). CRZ1 gene was tagged with GFP using the PCR-mediated gene tagging method described by Gerami-Nejad et al. (2001) with some modifications. Briefly, the CRZ1-GFP-CgTRP1 tagging cassette for transformation of the wild type and  $ptp2\Delta msg5\Delta$  double disruptant strains was constructed using primers CRZ1-GFP F: 5' -GTACGAA GAAGCCAGACAGGAGAAATCGGGAC AAGAGAGTTAA-3' and CRZ1-GFP R: 5'-AAAAAAAATTCCTATTCAAAGCTTAA upstream and downstream of the CRZ1 gene stop codon. The plasmid p2446 (containing the GFP-CgTRP1 fragment) from BYP2446 was used as template for PCR.

### 2.2.2 Genetic Manipulations

Construction of single, double and triple disruptants was carried out using the PCRmediated gene disruption method (Gietz and Schiestl, 2007) using *Candida glabrata HIS3, C. glabrata LEU2* or *S. cerevisiae* genes (*CgHIS3, CgLEU2* or *ScURA3*, respectively) or mating as previously described by Hermansyah. (2009; 2010). Confirmation PCR using specific primers was used to verify the correct disruption of genes. The primers for used in this study are listed in the Table 3.

### 2.2.3 RNA isolation

RNA was isolated using the hot phenol method (Spellman et al., 1998) with some modifications. Briefly, cells grown to mid-log phase at  $30^{\circ}$ C in YPD with or without 0.3M CaCl<sub>2</sub> were harvested by centrifugation. Total RNA was extracted by sequential treatment of TES/distilled water (DW)-saturated phenol, Trizol-LS (BRL), chloroform, and isopropanol before precipitation with Sodium Acetate (pH 5.2) – 99.9% EtOH (DEPC) mixture. The resulting RNA was dissolved in pre-chilled distilled water.

### 2.2.4 Microarray analysis

Total RNA was prepared using the hot phenol method (Spellman et al., 1998) with some modifications. First-strand cDNA targets were synthesized using Amino Allyl MessageAmp<sup>™</sup> II aRNA amplification kit (Ambion, Applied Biosystems) after which, they were labeled with Cy3-dUTP or Cy5-dUTP (Amersham Biosciences AB, Uppsala, Sweden). For DNA microarray, 3D-Gene<sup>™</sup> Yeast Oligo chip *S. cerevisiae* 6k containing 5,888 spotted genes was used (DNA Chip Research, Inc., Yokohama, Japan) and hybridization was done using the dual color method. Microarrays were scanned with ScanArray Lite (PerkinElmer Inc., Waltham, Massachusetts, USA). Differential expression analysis of the microarray

data was carried out using Genowiz<sup>TM</sup> 4.0 microarray data analysis software (Ocimum Biosolutions,Hyderabad, India). Replicated values for genes were merged and the median values of the expression ratios were considered for the dataset while empty spots were removed by filtering. Fold change analysis was done to detect highly expressed genes. Genes with 2 folds up/down-regulation were considered as differentially expressed at a pvalue < 0.05, Student's t-test. Functional classification of the genes was performed using gene ontology and pathway analysis (GOstat). The list of calcium-regulated calcineurindependent genes considered was taken from the genome-wide gene expression studies conducted by Yoshimoto and colleagues (Yoshimoto et al., 2002).

### 2.2.5 mRNA level determination using quantitative PCR

The differential expression data of selected genes were validated by quantitative RT-PCR. Using the total RNA isolated by the hot phenol extraction, first strand cDNA synthesis was carried out using the High Capacity cDNA Reverse Transcription kit (Applied Biosystems AB, Uppsala, Sweden). The synthesized cDNA was used as template for realtime PCR. Quantitative PCR was performed using 20 µl reaction mixtures on an Applied Biosystems 7300 Real Time PCR System with SYBR<sup>®</sup> Green PCR Master Mix (Applied Biosystems AB, Uppsala, Sweden). PCR was performed using the following amplification program: initial denaturation for 10 minutes at 95°C, followed by 40 cycles of denaturation for 15 seconds at 95°C and annealing for 1 minute at 60°C. *ACT1* was used as the reference gene for normalization of mRNA levels. The RT-PCR primers used in this study were designed using Primer Express version 2.0 (Applied Biosystems AB, Uppsala, Sweden) and are listed in Table 4.

### 2.2.6 Phenotypic analysis

Sensitivity of the strains to calcium and FK506 was assessed by spot plating on solid media. Cells suspended in water at 1 X  $10^6$  cells/ml were spotted onto YPDA and YPDA with 0.6 M CaCl2 and 1 µg/ml FK506. The plates were incubated at 30°C for 2 – 4 days.

### **2.2.7 Fluorescence microscopy**

Cells expressing GFP-tagged Crz1 proteins were cultured in YPD medium with or without 0.3 M CaCl<sub>2</sub> at 30°C until the mid-log phase after which the intracellular localization of the GFP-tagged protein was observed using a fluorescence microscope (BX61, Olympus) and photographed using a CCD camera (CCD-EX1, Universal Imaging Co., USA). DNA was stained for fluorescence using 4',6-diamidino-2-phenylindole (DAPI).

Strains	Alias	Genotype	Source/Description
SH6314	BY4742	MAT $\alpha$ his 3 $\Delta 1$ leu 2 $\Delta 0$ lys 2 $\Delta 0$ ura 3 $\Delta 0$	Invitrogen
SH5209	BY5209	MAT a ura3-52 his3- $\Delta 200$ leu2 $\Delta 1$ lys2 $\Delta 202$ trp1 $\Delta 63$	NBRP, YGRC*
SH5210	BY5210	MAT $\alpha$ ura3-52 his3- $\Delta$ 200 leu2 $\Delta$ 1 lys2 $\Delta$ 202 trp1 $\Delta$ 63	NBRP, YGRC*
SH5406	BY5210 <i>CRZ1-GFP</i>	MAT $\alpha$ ura3-52 his3- $\Delta$ 200 leu2 $\Delta$ 1 lys2 $\Delta$ 202 trp1 $\Delta$ 63 CRZ1-GFP::CgTRP1	Trp <sup>+</sup> transformant of SH5210 with <i>CRZ1-GFP</i> cassette
SH6790	$ptp2\Delta$	<i>MAT</i> a $\Delta ptp2::CgHIS3$ ura3-52 his3- $\Delta 20$ leu2 $\Delta 1$ lys2 $\Delta 202$ trp1 $\Delta 63$	0SH5209 disruptant
SH6791	$msg5\Delta$	MAT α Δmsg5::CgLEU2 ura3-52 his3- Δ200 leu2 Δ1 lys2 Δ202 trp1 Δ63	SH5210 disruptant
SH6792	$ptp2\Delta$ msg5 $\Delta$	MAT α Δptp2::CgHIS3 Δmsg5::CgLEU2 ura3-52 his3- Δ200 leu2 Δ1 lys2 Δ202 trp1 Δ63	(SH6790 x SH6791)- 1B
SH6793	$ptp2\Delta$ msg5 $\Delta$	$MAT$ a $\Delta ptp2::CgHIS3 \Delta msg5::CgLEU2$ ura3-52 his3- $\Delta 200$ leu2 $\Delta 1$ lys2 $\Delta 202$ trn1 $\Delta 63$	(SH6790 x SH6791)- 2B
SH8966	$ptp2\Delta$ msg5 $\Delta$	MAT a $\Delta ptp2::CgHIS3 \Delta msg5::CgLEU2$ ura3-52 his3- $\Delta 200$ leu2 $\Delta 1$ lys2 $\Delta 202$ trp1 $\Delta 63$	Leu <sup>+</sup> disruptant of SH6790 single disruptant using p1807 ( <i>Cal FU</i> 2)
SH8971	ptp2∆ msg5∆ CRZ1-GFP	MAT α Δptp2::CgHIS3 Δmsg5::CgLEU2 ura3-52 his3-Δ200 leu2 Δ1 lys2 Δ202 trn1 Δ63 CRZ1-GFP··CgTRP1	$Trp^+$ transformant of SH6792 with <i>CRZ1</i> - <i>GFP</i> cassette
SH8928	cna1 $\Delta$	MAT $\alpha \ \Delta cna1::KanMX4 \ his3 \ \Delta 1 \ leu2 \ \Delta 0 \ lys2 \ \Delta 0 \ ura3\Delta 0$	Invitrogen
SH8933	ptp2∆ msg5∆ cna1∆	MAT a $\Delta ptp2::CgHIS3 \Delta msg5::CgLEU2 \Delta cna1::KanMX4 ura3-52 (or ura3 \Delta 0)his3-\Delta 200 (or his3 \Delta 1) leu2\Delta 1 (leu2 \Delta 0 lys2\Delta 202 (lys2 \Delta 0) trp1\Delta 63$	4(SH6793 x SH8928)- 4C 9)
SH8929	$cna2\Delta$	MAT $\alpha$ $\Delta cna2::KanMX4$ his $3\Delta 1$ leu $2\Delta 0$ lvs $2\Lambda 0$ ura $3\Lambda 0$	Invitrogen
SH8934	ptp2∆ msg5∆ cna2∆	MAT a $\Delta ptp2::CgHIS3 \Delta msg5::CgLEU2 \Delta cna2::KanMX4 ura3-52 (or ura3 \Delta 0)his3-\Delta 200 (or his3 \Delta 1) leu2\Delta 1 (leu2 \Delta 0lvs2\Delta 202 (lvs2 \Delta 0) trp1\Delta 63$	4(SH6793 x SH8929)- 3C 9)
SH8930	$cnb1\Delta$	$MAT \propto \Delta cnb1::KanMX4 his3\Delta1 leu2\Delta0$ lys2 $\Delta0$ ura3 $\Delta0$	Invitrogen
SH8935	$ptp2\Delta$ $msg5\Delta$ $cnb1\Delta$	MAT a $\Delta ptp2::CgHIS3 \Delta msg5::CgLEU2 \Delta cnb1::KanMX4 ura3-52 (or ura3 \Delta 0)his3-\Delta 200 (or his3 \Delta 1) leu2\Delta 1 (leu2 \Delta 0lys2\Delta 202 (lys2 \Delta 0) trp1\Delta 63$	4(SH6793 x SH8930)- 13A 9)
SH8967	$crz1\Delta$	$MAT \propto \Delta crz1::KanMX4 his3\Delta1 leu2\Delta0$ lvs2A0 ura3A0	Invitrogen
SH8968	ptp2∆ msg5∆ crz1∆	MAT a $\Delta ptp2::CgHIS3 \Delta msg5::CgLEU2$ $\Delta crz1::KanMX4 ura3-52 (or ura3 \Delta 0)$ $his3-\Delta 200 (or his3 \Delta 1) leu2\Delta 1 (leu2 \Delta 0)$ $lys2\Delta 202 (lys2 \Delta 0) trp1\Delta 63$	(SH6793 x SH8967)- 4B ))

Table 2. S. cerevisiae strains used in the study

SH8931	$slt2\Delta$	MAT $\alpha \Delta slt2::KanMX4$ his $3\Delta 1$ leu $2\Delta 0$	Invitrogen
SH8972	$ptp2\Delta$ msg5 $\Delta$	MAT (Δptp2::CgHIS3 Δmsg5::CgLEU2	(SH6793 x SH8931)- 2B
	$slt2\Delta$	$his_3 \Delta 200 \text{ (or } his_3 \Delta 1)  leu 2\Delta 1 \text{ (leu 2 } \Delta 0)$	2D /)
SH8986	$ptp2\Delta$ msg5 $\Delta$	MAT α Δptp2::CgHIS3 Δmsg5::CgLEU2 Δste12::CgTRP1 urg3-52 his3- 4200 leu	$\operatorname{Trp}^+$ disruptant of 2 SH6792 double
	$stel 2\Delta$	$\Delta 1 \ lys2 \ \Delta 202 \ trp1 \ \Delta 63$	disruptant using p1805
SH8987	$ptp2\Delta$ msg5 $\Delta$	MAT α Δptp2::CgHIS3 Δmsg5::CgLEU2 Δsok2::ScURA3 ura3-52 his3- Δ200 leu2	Ura <sup>+</sup> disruptant of SH6792 double
	$sok2\Delta$	Δ1 lys2 Δ202 trp1 Δ63	disruptant using p3276 ( <i>ScURA3</i> )
SH8988	$ptp2\Delta$	MAT α Δptp2::CgHIS3 Δmsg5::CgLEU2	Ura <sup>+</sup> disruptant of
	msg5 $\Delta$	<i>∆ste12::CgTRP1 ∆sok2::ScURA3</i> ura3-5	2SH8987 triple
	$stel 2\Delta$	his3- Δ200 leu2 Δ1 lys2 Δ202 trp1 Δ63	disruptant using p3276
	$sok2\Delta$	2 1	(ScURA3)

\*NBRP/YGRC, National BioResource Project/Yeast Genetic Research Center, Japan (http://yeast.lab.nig.ac.jp/nig/index\_en.html)

Primer name	Primer sequence
Gd PTP2 F	5' ATAACGGCAATAGAATGGCTTCTTCCGCTATATCGGAA
	AACACAGG AAACAGCTATGACC 3'
Gd PTP2 R	5' GTAGCAATATACTTGAAATCAGGATTAATTTGCGTGAG
	CTGTTGTAAAACGACGGCCAGT 3'
Co PTP2 F	5' CTCAAGCTTGGAC ACTCGTTTAATTTAGCCA 3'
Co PTP2 R	5' CTCAAGCTTATTCGGTATTGG CACAAACTTT 3'
Gd MSG5 F	5' ACATCGATTTCAAGCCAAACTCACCGCGTTCCTTACAA
	AACACAGGAAACAGCTATGACC 3'
Gd MSG5 R	5' TCGTTGTCCACAGAAGCTTCCAGTGAATCTGCGGGTTG
	AGGTTGTAAAACGACGGCCAGT 3'
Co MSG5 F	5' CTCGGATCCGTAGTGATGGATGATGTGATTT 3'
Co MSG5 R	5' CTCGGATCCGTGCCCATGGTAATTTTTGACG 3'
Co CNA1 F	5' CAATAAATACAAACTGTTTG 3'
Co CNA1 R	5' ATTTGAATTTAGAACCGCTT 3'
Co CNA2 F	5' TCTTCTACGTATTTTTGTAT 3'
Co CNA2 R	5' TTCAAAATGGTGAAGTTAGA 3'
Co CNB1 F	5' AATTTGTTTCCCTCGACTTC 3'
Co CNB1 R	5' AGATATCAAAGCTATTAATA 3'
Co CRZ1 F	5' CGTCGAGGACCCTACCCCTG 3'
Co CRZ1 R	5' GAATTGAGCTACTTTTGTCT 3'
Gd STE12 F	5' ATAGCGGAACCGCTTTCTTTATTTGAATTGTCTTGTTC
	ACCAAGGCACAGGAAACAGCTATGACC 3'
Gd STE12 R	5' CCGCATTTTTAATTCTTGTATCATAAATTCAAAAATT
	ATATTATAGTTGTAAAACGACGGCCAGT 3'
Co STE12 F	5' ACTCTTCGCGGTCAGGTCTC 3'
Co STE12 R	5' GTCATGTTAAGAAACTCATT 3'
Gd SOK2 F	5' CAGTACAAAATCATCCTTATATAACCCTGGTAAGGT
	CCTTTTGTCCTTCGTACGCTGCAG 3'
Gd SOK2 R	5' GTTTTGATTAAAGTAACATAATTATCCAAGGAATT
	CATAGTTGTTGCCACTAGTGGATCT 3'
Co SOK2 F	5' GAAAAAAGTTCATCTTTAAC 3'
Co SOK2 R	5' AGTTTCTCGTTTAAAAAA TC 3'

Table 3. Primers used for disruption and confirmation PCR

Table 4. Primers used in RT-PCR

Primer name	Primer sequence
rt ENA1 F	5' TTGGATCCCTCGCTTTGGT 3'
rt ENA1 R	5' CAAACTCACGTTGCCCTCATT 3'
rt GSC2 F	5' CACAGACCTTCACCGCATCTT 3'
rt GSC2 R	5' GACTCTGCATATTTAGCAGCAAAAAC 3'
rt PMC1 F	5' GCCATAAGTGTTGCCCACAA 3'
rt PMC1 R	5' GAGTCTTTATCACGCAGCTGTTCA 3'
rt PMR1 F	5' ACCACCGCGGCCTGTA 3'
rt PMR1 R	5' TACCATCTTCGGCCATCTCTTT 3'

### 2.3 Results

## 2.3.1 An active calcineurin pathway is inhibitory to the growth of the calcium-exposed $ptp2 \Delta msg5 \Delta$ double disruptant

S. cerevisiae cells generally respond to high extracellular calcium exposure by eliciting the calcium/calmodulin-dependent signaling pathway to maintain cell viability (Cyert, 2003). To determine whether activation of calcineurin occurs in the  $ptp2\Delta msg5\Delta$ double disruptant upon exposure to calcium, we examined the expression of 120 genes that are known to be regulated by calcineurin using microarray (see Materials and methods). Based on Gene Ontology (GO) categories, 46 out of 102 calcineurin-dependent genes belong to "molecular function" while 23 and 33 out of 102 belong to "cell component" and "biological process", respectively.

Microarray results showed up-regulation of calcineurin-regulated genes in both the  $ptp2\Delta msg5\Delta$  double disruptant (89.1%) and wild-type (87.7%) strains upon exposure to high extracellular calcium. (Fig. 4, column 1 and column 3, respectively). Interestingly, the expression values corresponding to the up-regulated calcineurin genes in the calcium-exposed  $ptp2\Delta msg5\Delta$  double disruptant did not vary significantly to that of the calcium-treated wild-type strain although most exhibited slight decrease in expression (Fig.4, column 2).

To confirm these results, the transcription level of several representative genes known to be up-regulated in a calcineurin-dependent manner was validated using real-time RT-PCR (Cunningham and Fink, 1994). These include the β-1,3-glucan synthase gene *GSC2* (*FKS2*), *PMC1*, *PMR1* and *ENA1*(*PMR2*), in which the latter three encode for P-type ATPases required for ion homeostasis (Yoshimoto et al., 2002). Results of the RT-PCR analysis revealed a similar trend in the expression with the microarray data wherein exposure to calcium caused an increase in the expression level of *GSC2*, *PMC1* and *ENA1* in the

 $ptp2\Delta msg5\Delta$  double disruptant. Interestingly, expression of *PMR1* was not induced either in the wild type or  $ptp2\Delta msg5\Delta$  double disruptant upon exposure to calcium probably because *PMR1* is controlled by other calcium-dependent mechanisms. These results indicate that the calcineurin pathway is active in the  $ptp2\Delta msg5\Delta$  double disruptant and the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant is not due to a defective calcineurin pathway (Fig. 5).

To strengthen the above conclusion, we studied the localization of Crz1 in the wild type and the  $ptp2\Delta msg5\Delta$  double disruptant after calcium exposure. Since calcineurin activation is known to cause nuclear localization of Crz1 (Stathopoulos-Genontides et al., 1999), we tagged Crz1 with GFP in both wild type and  $ptp2\Delta msg5\Delta$  double disruptant. As expected, Crz1 was localized in the nucleus of the calcium-exposed wild type. Similarly, nuclear localization was observed in the  $ptp2\Delta msg5\Delta$  double disruptant exposed to calcium thereby indicating an active calcineurin pathway in the  $ptp2\Delta msg5\Delta$  double disruptant upon calcium exposure (Fig. 6).

We inhibited the calcineurin pathway either by FK506 (a calcineurin inhibitor) treatment or disruption of *cnb1* $\Delta$  (calcineurin regulatory subunit) to determine the effect of inactivation of the calcineurin pathway on the phenotype of the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant. As expected, disruption of either *cna1* $\Delta$  or *cna2* $\Delta$  in the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant did not cause calcium tolerance since *CNA1* and *CNA2* redundantly codes for the calcineurin catalytic unit (Fig. 7B). However, addition of FK506 in the medium (Fig. 7A) or disruption of *CNB1* (Fig. 7B) was able to suppress the calcium sensitive phenotype of the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant, indicating that deactivation of the calcineurin pathway in the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant can restore growth in a calcium-exposed environment. Taken together, we conclude that transcription of calcineurin genes is not impaired and calcineurin

YOR209C	YOLIS8C	YGLU53W
YOR288C	YMR096W	YDR319C -2.56
YNR064C	YDL128W	YLR046C
YPL056C	YHR138C	YNL074C
YPR167C	YDR039C_YDR040C	YML125C
YOL015W	YKLO01C	YAR031W
YMR095C	YJR078W	YGL156W
YOR035C	YMR107W	YDR129C
YJLO16W	YGR144W	YJL213W
YOR237W	YPL149W	YDR210W
YOR338W	YGR032W	YLR120C
YER144C	YGR110W	YHR071W
YBR182C	YGR131W	YER184C
YGR237C	YOL082W	YJR161C
YPR006C	YGR295C	YLR303W
YNL092W	YOR134W	YOR273C
YML123C	YGL167C	YDL012C
YOR019W	YLR350W	YLR149C
YKL213C	YDL173W	YNL210W
YAR028W	YLL057C	YKL218C
YBR036C	YLR054C	YNR010W
YBR284W	YCR009C	YBR296C
YNR063W	YML118W	YLR136C
YDR038C	YMR026C	YDL248W
YNL020C	YAR027W	YGR268C
YGL001C	YOR299W	YLR099C
YLR142W	YHR124W	YOR137C
YPL233W	YAL053W	YCR007C
YGR213C	YPR165W	YDR380W
YNL044W	YBR127C	YKR053C
YLR327C	YBR287W	YKL126W
YOL163W	YML119W	YMR040W
YMR020W	YJL094C	YKL157W
YKL159C	YPR198W	YLR257W
YGR212W	YNL208W	YMRU16C
YDL185W	YGL006W	YMR305C
YNL336W	YDL010W	YOL162W
YMR094W	YDL146W	YERUSUC
YMR238W	YMR062C	YORU36W
YBR203W	YDR482C	

Figure 4. Heatmap of the expression response of calcium regulated-calcineurin genes in the wild

1.91

6.38

type and  $ptp2\Delta msg5\Delta$  disruptant with or without calcium exposure. The column number indicates the set-up as follows: 1) comparison between Ca<sup>2+</sup>-exposed  $ptp2\Delta msg5\Delta$  disruptant with untreated  $ptp2\Delta msg5\Delta$  disruptant; 2) comparison between Ca<sup>2+</sup>-exposed  $ptp2\Delta msg5\Delta$  disruptant with  $Ca^{2+}$ -exposed wild type; 3) comparison between  $Ca^{2+}$ -exposed wild type with untreated wild type. All strains were cultivated in YPD media and calcium treatment was done by supplementing the YPD medium with 0.3 M CaCl<sub>2</sub>. Four independent DNA microarray data were analyzed using Genowiz (Ocimum Biosolutions). The data were normalized prior to statistical analysis using the LOWESS algorithm. Genes showing more than two-fold induction correspond to the rows and the columns represent the experimental set-ups. Red and green colors represent upregulation and down-regulation of expression respectively. The color saturation represents the magnitude of the expression ratio, as indicated by the scale at the right side of the figure.



■Wild type (- Ca<sup>2+</sup>) ■Wild type (+ Ca<sup>2+</sup>) ■*ptp2∆msg5∆* (- Ca<sup>2+</sup>) ■*ptp2∆msg5∆* (+ Ca<sup>2+</sup>)

Figure 5. Calcineurin activation occurs in the  $ptp2\Delta msg5\Delta$  double disruptant. Representative calcineurin genes in the  $ptp2\Delta msg5\Delta$  double disruptant are up-regulated upon calcium exposure. cDNA synthesized from total RNA isolated from the sample strains grown with (0.3 M CaCl<sub>2</sub>) or without calcium were used as templates for RT-PCR using SYBR Green and specific primers for *GSC2*, *PMC1*, *ENA1* and *PMR1*. *ACT1* was used as control. *GSC2*, *PMC1* and *ENA1* showed increased expression while *PMR1* expression remained the same in the  $ptp2\Delta msg5\Delta$  double disruptant upon calcium exposure. The results are means of three experiments from three independent preparations.





calcineurin activation has a negative effect on the growth of the  $ptp2\Delta msg5\Delta$  double disruptant upon exposure to calcium.

Interestingly, additional disruption of *CRZ1* in the  $ptp2\Delta msg5\Delta$  double disruptant background did not confer calcium tolerance. The fact that the  $crz1\Delta$  single disruptant is also calcium sensitive implies that Crz1 is essential to the cell's viability under high extracellular calcium stress conditions. Our result also indicate that *CRZ1* is located downstream of the calcium-induced, growth-related redundant pathways mediated by Ptp2 and Msg5 where the parallel pathways converge (Fig. 7B).



Figure 7A. Inhibition of calcineurin by FK506 treatment suppresses the calcium sensitivity of the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant. BY5209 (wild type), ptp2 $\Delta$ , *msg5* $\Delta$ , *slt2* $\Delta$ , *ptp2* $\Delta$ *msg5* $\Delta$  *and ptp2* $\Delta$ *msg5* $\Delta$ , *slt2* $\Delta$ , *were grown in YPD to mid-log phase. Ten-fold serial dilutions of the cell suspensions were prepared and spot plated on to YPDA, YPDA supplemented with 0.6 M CaCl<sub>2</sub>, YPDA added with FK506 (calcineurin inhibitor) and YPDA with both CaCl<sub>2</sub> and FK506. The plates were incubated at 30°C for 2-4 days.* 



Figure 7B. Inhibition of calcineurin by *CNB1* disruption suppresses the calcium sensitivity of the *ptp2*\Delta*msg5*\Delta double disruptant. The subunits of calcineurin were individually disrupted in the *ptp2*\Delta*msg5*\Delta background to produce triple disruptants. *CNA1* and *CNA2* are catalytic units while *CNB1* is the regulatory subunit of calcineurin. BY5209 (wild type), *cna1*\Delta, *cna2*\Delta, *cnb1*\Delta, *crz1*\Delta, *ptp2*\Delta*msg5*\Delta, *ptp2*\Delta*msg5*\Delta*cna1*\Delta, *ptp2*\Delta*msg5*\Delta*cna2*\Delta, *ptp2*\Delta*msg5*\Delta*cnb1*\Delta and *ptp2*\Delta*msg5*\Delta*crz1*\Delta were grown in YPD to mid-log phase. Ten-fold serial dilutions of the cell suspensions were prepared and spot plated on to YPDA with and without 0.6 M CaCl<sub>2</sub>. The plates were incubated at 30°C for 2-4 days.

### 2.3.2 Up-regulated cell wall genes in the *ptp2\Deltamsg5\Delta* double disruptant share common

### transcription factors Sok2 and Ste12

Activation of the SLT2 pathway leads to the induction of transcription factors like

Rlm1 and Swi4/Swi6 which in turn stimulates the expression of downstream or effector genes

that play roles in the maintenance of cell wall integrity (Watanabe and Matsumoto, 1995;

Levin, 2005; Jung and Levin, 1999). Based on this, we inferred that hyper-activation of the

SLT2 pathway essentially causes up-regulation of several downstream genes that eventually

leads to cell toxicity (Watanabe and Matsumoto, 1995; Hahn and Thiele, 2002). Since many
of the effector genes downstream of the SLT2 pathway are involved in cell wall construction and cell wall-related processes, we used the microarray data to search for cell wall-related genes that are differentially regulated in the calcium exposed-*ptp2* $\Delta$ *msg5* $\Delta$  double disruptant. Out of the 31 cell wall-related genes that are reported to be differentially expressed during cell cycle, we found five genes namely *CWP1*, *CHS1*, *PST1*, *SCW11* and *EXG1* that are upregulated while another gene, *CWP2*, was down-regulated in *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant exposed to calcium (Table 5) (Jung and Levin, 1999; Klis et al., 2006).

Table 5. Cell wall genes that are differentially regulated in the  $ptp2\Delta msg5\Delta$  double disruptant upon exposure to calcium

Gene name	Molecular function	Log2 expression
CWP1	Structural constituent of cell wall	1.1
CWP2	Structural constituent of cell wall	-1.2
CHS1	Chitin synthase activity	1.2
PST1	Molecular function unknown	1.6
SCW11	Glucan 1,3-beta-glucosidase activity	1.9
EXG1	Glucan 1,3-beta-glucosidase activity	1.1

To find a link between the cell wall genes that are differentially expressed, we looked for transcription factors that were common among the differentially regulated genes. YEASTRACT analysis identified two probable transcription factors shared by the altered genes as Sok2 and Ste12 (Teixeira et al., 2005). However, compared to the wild type, *SOK2* and *STE12* were not differentially expressed in the *ptp2*\Delta*msg5*\Delta double disruptant; indicating that the mode of regulation involved is post-transcriptional (unpublished data).

To further elucidate the roles of the transcription factors Sok2 and Ste12 in the calcium phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant, we disrupted SOK2 and STE12 independently and compositely in the  $ptp2\Delta msg5\Delta$  double disruptant background. Results showed that neither disruption of STE12 or SOK2 conferred calcium tolerance to the  $ptp2\Delta msg5\Delta$  double disruptant. However, disruption of both STE12 and SOK2 was able to

confer a slightly calcium tolerant phenotype as shown in the quadruple disruptant,  $ptp2\Delta msg5\Delta ste12\Delta sok2\Delta$  (Fig. 8). This indicates that both *STE12* and *SOK2* play roles in the calcium phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant and their synergistic relationship negatively affects growth in the presence of calcium.



Figure 8. Disruption of both transcription factors Ste12 and Sok2 confers partial calcium tolerance to  $ptp2\Delta msg5\Delta$  double disruptant. BY5209 (wild type),  $ptp2\Delta msg5\Delta$ ,  $ptp2\Delta msg5\Delta ste12\Delta$ ,  $ptp2\Delta msg5\Delta ste12\Delta$ ,  $ptp2\Delta msg5\Delta ste12\Delta$  and  $ptp2\Delta msg5\Delta ste12\Delta$  were grown in YPD to mid-log phase. Ten-fold serial dilutions of the cell suspensions were prepared and spot plated on to YPDA with and without 0.6 M CaCl<sub>2</sub>. The plates were incubated at 30°C for 2-4 days.

# 2.3.3 SLT2 pathway is hyper-activated in the *ptp2\Deltamsg5\Delta* double disruptant upon

## exposure to high extracellular calcium

MAPK signaling cascades are activated by the process of phosphorylation of cascades of protein kinases (Zhao et al., 1998). Furthermore, regulation of these pathways is via dephosphorylation of MAPK by protein phosphatases such as Ptc1, Ptc2, Ptc3, Ptp2, Ptp3 and Msg5 (Martin et al., 2005). Since Ptp2 and Msg5 are known to negatively regulate Slt2 phosphorylation, we confirmed if Slt2 was indeed hyper-phosphorylated in the *ptp2*\Delta*msg5*\Delta double disruptant. As expected, Western blot analysis revealed that Slt2 was hyperphosphorylated in the *ptp2*\Delta*msg5*\Delta double disruptant compared to the wild type (Fig. 9). Our result is consistent with the earlier findings that disruption of both *PTP2* and *MSG5* PPase genes results in a hyper-phosphorylated state of Slt2 (Hermansyah et al., 2009).



Figure 9. Slt2 is hyper-phosphorylated in the calcium-exposed  $ptp2\Delta msg5\Delta$  double disruptant. Representative Western blot showing the phosphorylation level of Slt2p. Soluble protein extracts were prepared from cells grown in YPD with (0.3 M CaCl<sub>2</sub>) or without calcium at mid-log phase. Total Slt2p and the phosphorylated form of Slt2p were detected using anti-Mpk1p (Slt2p) and anti-phospho-p44/42 MAPK (Thr<sup>202</sup>/Tyr<sup>204</sup>) antibodies, respectively. Anti-Act1p was used as control.

2.3.4 Calcium sensitivity in the *ptp2\musg5\u03e4* double disruptant results from a hyper-

activated, calcium-induced signaling pathway caused by the improper activation of the

## SLT2 pathway and an active calcineurin

We presumed that the simultaneous activation of both SLT2 and calcineurin pathways might be toxic to cells as genetic evidence indicates a functionally redundant relationship between SLT2 and calcineurin pathways in growth-related functions (Nakamura et al., 1996). Thus, we tried to inactivate either one of the redundant pathways (calcineurin or SLT2 pathways) to determine if the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant can be suppressed. Indeed,  $slt2\Delta$  disruption was able to confer calcium tolerance in the  $ptp2\Delta msg5\Delta$  double disruptant (Hermansyah et al., 2012). Moreover, when calcineurin was inactivated in the  $ptp2\Delta msg5\Delta$  double disruptant by either disrupting *CNB1* (calcineurin regulatory subunit gene) or by treatment with a calcineurin inhibitor, FK506, we found that inactivation of calcineurin by either means was also able to suppress the calcium sensitivity of the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant (Fig. 7). Thus, calcineurin seems to have an inhibitory role in the growth of the calcium-exposed *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant under a hyperphosphorylated SLT2 state (Fig. 7).

Inactivation of both calcineurin (FK506 treatment) and SLT2 pathways (disruption of  $slt2\Delta$  in  $ptp2\Delta msg5\Delta$ ) led to a calcium sensitive phenotype, indicating that the presence of at least one of the redundant pathways is essential for viability under high calcium conditions (Fig. 7A). Taken together, these findings indicate that a hyper-activated signaling state caused by a simultaneously over-activated SLT2 pathway and an active calcineurin pathway led to a calcium sensitive phenotype in the  $ptp2\Delta msg5\Delta$  double disruptant upon exposure to high extracellular calcium.

#### **2.4 Discussion**

We previously reported a PPase double disruptant,  $ptp2\Delta msg5\Delta$ , exhibiting sensitivity at very high concentrations of calcium (0.6 M). This calcium sensitive phenotype was suppressed by an additional disruption of either one of the six PKases namely *BCK1*, *MKK1*, *SLT2*, *MCK1*, *SSK2* and *YAK1* (Hermansyah et al., 2010). Although we have reported in our earlier study that the SLT2 pathway is over-activated in the  $ptp2\Delta msg5\Delta$  double disruptant, the detailed mechanism on how a hyper-activated SLT2 pathway causes calcium sensitivity remained unclear. In this work, we clarified the conditions that lead to the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant in terms of functional redundancy at the protein and signal cascade levels.

At the protein level, Ptp2p and Msg5p, which are protein tyrosine (PTP) and dual specificity (DSP) PPases respectively, are known to co-regulate Slt2p (Mpk1p) of the SLT2 (Cell Wall Integrity or Mpk1) pathway (Martin et al., 2005; , Levin, 2005). Based on genetic analysis, we established the redundant roles of these two PPases in the calcium-mediated signaling process since single disruption of either  $ptp2\Delta$  or  $msg5\Delta$  is calcium tolerant (Sakumoto et al., 2002). In general, the MAPK cascade is comprised of a sequential activation of protein kinases. Specifically, MEK kinase (MEKK or MAPKKK) activates the MAPK kinase (MEK or MAPKK) which in turn activates the MAP kinase by dual phosphorylation of its TXY motif (Zhao et al., 2007). This cascade event will eventually result in the activation of various transcription factors or induction of specific sets of genes based on the environmental stimuli. Thus, we presumed that hyper-activation of the SLT2 pathway arises when there is accumulation of hyper-phosphorylated Slt2p in the absence of the negative regulators, Ptp2p and Msg5p (Fig. 10). In this case, a hyper-phosphorylated state of Slt2p is indicative of a hyper-activated SLT2 pathway since the activation of yeast MAPK cascades is through phosphorylation (Zhao et al., 2007). This hyper-activated state of the

SLT2 pathway induces G1 delay by possibly regulating the expression of effector genes such as *CLN2* via several transcription factors as we previously reported (Hermansyah et al., 2009). In connection with this, we reported the down-regulation of *CLN2* in the *ptp2\Delta msg5\Delta* double disruptant upon exposure to high extracellular calcium (Hermansyah et al., 2010). The *CLN2* down-regulation can be explained based on the model for regulation of Swi4 by Slt2 (Mpk1) wherein Slt2 protein is activated by a calcium-induced cell wall stress and is likely to phosphorylate Swi6, causing its exclusion to the nucleus. A non-nuclear localized Swi6 therefore cannot form the SBF complex, resulting in reduction or abolishment of transcription of late G1 genes including *CLN1* and *CLN2* (Levin, 2005).

The fact that the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant exhibits growth arrest (G1 arrest) similar to that of calcineurin-deficient mutants suggested the possible involvement of calcineurin in this signaling pathway. Although essential for survival during stressed conditions, we propose that an active calcineurin coupled with a hyper-activated SLT2 pathway becomes toxic to yeast cells upon exposure to high extracellular calcium (Fig. 7A and 7B). To our knowledge, this is the first report of the existence of a functionally redundant relationship between the SLT2 and calcineurin pathways in response to high levels of calcium. In this case where the SLT2 pathway is hyper-activated due to the absence of the negative regulators Ptp2p and Msg5p, one way to alleviate the harmful effects of an over-active calcium-induced signaling pathway is to inactivate calcineurin by cnb1 disruption or treatment with a calcineurin inhibitor, FK506 (Fig. 7A and 7B). Therefore, the redundant pathways related to growth serve as safeguards for unwanted effects of an inactive signaling pathway. In this system, a signal cascade assumes the function of another cascade when it is inactivated or impaired. However, improper activation of both signaling cascades becomes harmful to the cells due to signal hyper-activation that eventually leads to cessation of growth. In this study, we have

demonstrated that such is the case for strains lacking signaling cascade regulators such as

PPases Ptp2 and Msg5 when exposed to high extracellular calcium (Fig. 8).



Figure 10. Over-activation of the calcium-induced signaling pathway in the  $ptp2\Delta msg5\Delta$  double disruptant causes calcium sensitivity. Calcineurin and SLT2 pathways share redundant functions in cell processes related to growth. Upon exposure to high extracellular calcium, calcineurin is the preferred route while SLT2 pathway is inactivated through dephosphorylation of Slt2p by the functionally redundant Ptp2p and Msg5p. In the absence of both Ptp2 and Msg5 PPases, the SLT2 pathway is hyper-activated and causes growth delay due to a hyper-activated state of the calcium-induced signaling cascade when coupled with an active calcineurin pathway. The two parallel and redundant pathways that are induced by calcium converge at Crz1.

Similar to a published report (Mazur et al., 1995), our result showed that expression of GSC2 (*FKS2*) is low under normal growth conditions but immediately increases upon exposure to calcium. In the  $ptp2\Delta msg5\Delta$  double disruptant exposed to calcium, there was also an increase in GSC2 expression (2-fold) although at a much lower increment compared to the wild type (6-fold) (data not shown). This can be an indication that the calcineurin pathway is operational but its activity is slightly impaired in the  $ptp2\Delta msg5\Delta$  double disruptant. There

was an earlier report of a dual control of *GSC2* expression by PKC and calcineurin pathways induced by heat stress. If calcineurin activity is impaired, it is expected that calcium-induced expression of *GSC2* is also reduced because calcium induction of *GSC2* is not cell integrity pathway-dependent (Zhao et al., 1998). Moreover, *GSC2* expression in response to extracellular calcium is strictly calcineurin-dependent. Our results imply that hyper-activation of the SLT2 pathway possibly has an inhibitory effect on calcineurin activity.

In the advent of sudden exposure to high extracellular calcium, induction of stress response plays an integral role in the cell's survival. It is therefore imperative that S. cerevisiae possess parallel, alternative mechanisms to control calcium signaling in cases where the major pathway is impaired (http://oicr.on.ca). However, stringent regulation of these redundant pathways is essential to avoid over-activation. Our results show that the SLT2 and calcineurin pathways co-mediate cellular processes related to growth upon exposure to high extracellular calcium. It is highly probable that upon exposure to calcium, the calcineurin pathway is the primary signaling response that the yeast employs while the SLT2 pathway, being secondary, is only activated in conditions where the calcineurin pathway is completely impaired. In the wild type, this is accomplished by inhibition of the SLT2 pathway via Slt2p dephosphorylation with the functionally redundant PPases, Ptp2 or Msg5 (Fig. 10). However, over-activation of the calcium-induced signaling pathway can occur if both pathways are active and this state promotes growth inhibition. Our results signify that in the event of SLT2 pathway hyper-activation in the presence of high extracellular calcium, calcineurin function can be abrogated to prevent growth inhibition. However, we found that complete inhibition of both pathways by the disruption of *SLT2* pathway genes and treatment of FK506 led to a calcium sensitive phenotype (Fig. 7A), indicating that the presence of at least one of the redundant pathways is essential for viability in high calcium conditions (Fig. 10).

Our results revealed that Crz1 is an essential part of the calcium-induced, growthrelated signaling pathway manifested in the  $ptp2\Delta msg5\Delta$  double disruptant. Genetic analysis suggested that Crz1 is located downstream of the redundant SLT2 and calcineurin pathway and is possibly the point of convergence between the parallel pathways. We also found that common calcineurin and Crz1-induced genes like *PMR1*, *GSC2*, *PMC1* and *ENA1* were not the transcriptional targets of the calcium-induced signaling pathway in the  $ptp2\Delta msg5\Delta$ double disruptant (Fig. 5), thus pointing out that the parallel pathways might induce the expression of other genes related to growth and stress response. This assumption is reinforced by the fact that representative calcineurin-regulated genes *PMC1* and *ENA1* were induced in the  $ptp2\Delta msg5\Delta slt2\Delta$  but not in  $ptp2\Delta msg5\Delta cnb1\Delta$  upon exposure to high extracellular calcium conditions (Fig. 11). The identity of the transcriptional targets of the parallel SLT2 and calcineurin pathways mediated by Ptp2 and Msg5 is still unknown.

Interestingly, Sdp1 or Yill13p has been implicated with SLT2 pathway regulation via dephosphorylation of Slt2, which is similar to the function of Ptp2 and Msg5 in the SLT2 pathway (Mattison et al., 1999; Martin et al., 2000; Hahn and Thiele, 2002; Collister et al., 2002). In the *ptp2Amsg5A* double disruptant, the presence of a functional Sdp1p could not compensate for the absence of both Ptp2 and Msg5, resulting in the over-activation of the calcium-induced signaling pathway and eventually calcium sensitivity. This fact rules out the possibility that Sdp1 is functionally redundant to Ptp2 and Msg5 in relation to the calcium-induced signaling pathway in the *ptp2Amsg5A* double disruptant. In accordance with our previous statement, our unpublished data showed that the *sdp1A* single disruptant (BY4742 background) was calcium tolerant (0.6 M CaCl<sub>2</sub>) which is an indication that Sdp1 activity is stress specific and has no relation to calcium stress.



Figure 11. Calcineurin-regulated genes *PMC1* and *ENA1* are induced in the  $ptp2\Delta msg5\Delta slt2\Delta$  but not in  $ptp2\Delta msg5\Delta cnb1\Delta$  triple disruptant. cDNA synthesized from total RNA isolated from the sample strains grown with (0.3 M CaCl<sub>2</sub>) or without calcium were used as templates for RT-PCR using SYBR Green and specific primers for *GSC2*, *PMC1*, *ENA1* and *PMR1*. *ACT1* was used as control. *PMC1* and *ENA1* showed increased expression in the wild type,  $ptp2\Delta msg5\Delta$  and  $ptp2\Delta msg5\Delta slt2\Delta$  but not in  $ptp2\Delta msg5\Delta cnb1\Delta$  upon calcium exposure. *PMR1* expression remained the same in all strains upon calcium exposure. *GSC2* showed increased expression in both  $ptp2\Delta msg5\Delta slt2\Delta$  and  $ptp2\Delta msg5\Delta cnb1\Delta$ , indicating a calcineurin-independent gene induction. The results are means of three experiments from three independent preparations.

This conclusion is supported by reports showing that *YIL113W* disruptant (*sdp1* $\Delta$ ) exhibited normal response to mating pheromone and has no effect on the phosphorylation of Hog1p MAPK, thus pointing out that Sdp1 is not involved in regulating the response to pheromone and osmotic stress in *S. cerevisiae* (Collister et al., 2002). Results of these studies support the claim that Yil113p phosphatase is a specific regulator of signaling through the Slt2/Mpk1 MAPK pathway and it is highly probable that calcium-induced signaling mediated by the functionally redundant Ptp2 and Msg5 is independent of the Sdp1 function.

This chapter discussed the mechanism of calcium sensitivity in the  $ptp2\Delta msg5\Delta$ double disruptant and the suppression mechanism conferred by either disrupting SLT2 pathway genes or inactivating calcineurin. The disparity in the mechanism in which calcium tolerance is conferred by  $ssk2\Delta$  or  $yak1\Delta$  disruption remains to be elucidated. Also, the extent and cause of calcineurin impairment is not presently known and is the subject of future study. In this study, we found proof of the functional redundancy of Ptp2 and Msg5 at the protein level as well as the SLT2 and calcineurin pathways at the cascade level in response to high extracellular calcium conditions. Furthermore, our results imply the important roles of PPases Ptp2 and Msg5 in preventing hyper-activation of SLT2 pathway upon exposure to high calcium concentrations. Lastly, strict modulation of the calcium-induced signaling pathway mediated by the functionally redundant SLT2 and calcineurin pathways is imperative for cell growth under high calcium environments.

#### Chapter 3

# Suppression mechanism of the calcium sensitivity in *Saccharomyces cerevisiae ptp2∆msg5*∆ double disruptant involves a novel HOG-independent function of Ssk2, transcription factor Msn2 and the PKA component Bcy1

#### **3.1 Introduction**

The growth and survival of eukaryotic cells under stress depend largely on their ability to quickly sense and adapt to the perturbations in the environment. Cells have complex networks of sensing and signaling cascades that are activated appropriately upon specific stress conditions; causing changes in gene expression program, metabolic profile and other cellular features that eventually lead to adaptations of cell growth and proliferation (Hohmann and Mager, 2002). In fact, cells often possess multiple pathways in response to a particular stress thus preventing unwanted consequences of improper activation or inactivation. Previously, we found the existence of functional redundancy in Ptp2 and Msg5 protein phosphatases in response to high extracellular calcium stress (Sakumoto et al., 2002). Suppressor analysis employing protein kinases revealed six suppressors whose additional disruption in the *ptp2\Deltamsg5\Delta* double disruptant conferred calcium tolerance namely *BCK1*, MKK1 and SLT2 (Cell Wall Integrity pathway), MCK1 (chromosome segregation and meiosis), YAK1 (PKA pathway) and SSK2 (HOG pathway) (Hermansyah et al., 2010). The suppressor roles of  $bckl\Delta$ ,  $mkkl\Delta$ , and  $slt2\Delta$  disruptions have been recently explained in Chapter 2 to relate to the co-regulation of calcium signaling in the *ptp2\Delta msg5\Delta* double disruptant by calcineurin and SLT2 pathways that acted as safeguards against sensitivity to high extracellular calcium concentrations (Laviña et al., 2013). On the contrary, FACS analysis implied that  $ssk2\Delta$  and  $vak1\Delta$  suppressors belong to a different regulatory pathway

since they were able to recover from G1 delay unlike  $bck1\Delta$ ,  $mkk1\Delta$ , and  $slt2\Delta$  suppressors (Hermansyah et al., 2010).

Ssk2 is the upstream MAP kinase kinase kinase of the HOG pathway that phosphorylates Pbs2 after its interaction with Ssk1 (Hohmann, 2002). Pbs2, in turn, phosphorylates and activates the MAP kinase Hog1 that results in the expression of genes involved in glycerol synthesis (Gustin et al, 1998). SSK2 has a close homolog, SSK22, which is redundant with Ssk2 in terms of Pbs2 phosphorylation (Maeda et al., 1995; Posas and Saito, 1998). In addition, Ssk2 has a specialized function in facilitating actin cytoskeleton reassembly after osmotic stress and at the end of the cell cycle (Yuzyuk et al., 2002). These specific functions of Ssk2 have been reported to be unrelated to other known components of the HOG pathway.

Yak1 is a serine-threonine kinase that acts as an antagonist for cell growth (Garrett et al., 1991). Yak1 can be directly phosphorylated and indirectly activated transcriptionally by PKA indicating that Yak1 is downstream of PKA (Pratt et al., 2007). The current model states that an active PKA represses the activity of transcription factors Msn2 and Msn4, thereby leading to the down-regulation of STRE genes that include *YAK1* (Kassis et al., 2000).

In this chapter, we describe a HOG-independent function of Ssk2 that mediates the calcium-sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant. Also, we report new suppressors,  $msn2\Delta$  and  $bcy1\Delta$ , which are possibly related to the suppression mechanism conferred by  $ssk2\Delta$  disruption.

#### 3.2 Materials and methods

#### 3.2.1 Strains, media and general methods

S. *cerevisiae* strains used in this study are summarized in Table 6. FY833 strain (SH5209) (*MATa ura3-52 his3-\Delta200 leu2\Delta1 lys2\Delta202 trp1\Delta63) was used as the wild type strain. Yeast strains were grown at 30°C in standard rich medium (YPD) (Sigma-Aldrich), SC medium and SPM medium as described previously (Amberg et al., 2005). YPDA was supplemented with 0.6 M CaCl<sub>2</sub> for the calcium phenotype test while 0.3 M CaCl<sub>2</sub> was used for mRNA and protein isolation. <i>Escherichia coli* strains used as sources of plasmids including BYP1804 for p1804 (*CgHIS3*), BYP1805 for p1805 (*CgTRP1*), BYP1419 for p1807 (*CgLEU2*), SHB3276 for p3276 (*ScURA3*) were acquired from NBRP, YGRC, Japan and cultivated in LB medium containing 100 (g/ml ampicillin at 37°C.

Single, double and triple disruptants were constructed using PCR-mediated gene disruption method or mating as previously described (Amberg et al., 2005; Gietz and Schiestl, 2007; Sherman and Hicks, 1991). Confirmation PCR using specific primers was done to verify the correct disruption of genes. The complete list of primers used in the construction and confirmation of the different kinds of disruptants is described on the Table 7.

For spot assay, yeast strains were grown on YPD broth until mid-logarithmic phase, after which the cell number was determined using a spectrophotometer (Smartspec<sup>TM</sup>300, Biorad). The cell suspension was diluted ten-fold using sterile water in a 96-well plate and spotted on appropriate media (YPDA, YPDA with 0.6M CaCl<sub>2</sub> or SC medium).

#### 3.2.2 Protein isolation and immunoblot analysis

Protein extracts for Western blot were prepared using the trichloroacetic acid (TCA) method from cells grown until mid-log phase ( $OD_{660} = 1.0$ ) at 30°C in YPD medium with or

without 0.3M CaCl<sub>2</sub> (An et al., 2006). The collected supernatant was used for western blotting. Protein extracts fractionated by SDS-PAGE using 10% polyacrylamide gels were transferred to PVDF Immobilon transfer membranes (Millipore). Total Hog1 was detected with anti-Hog1 (yC-15) antibody (Santa Cruz Biotechnology) while phosphorylated Hog1 was detected with anti-phospho-p38 MAP Kinase (Thr<sup>180</sup>/Tyr<sup>182</sup>) antibody (Cell Signaling Technology). Immunoblots were developed with horseradish peroxidase-conjugated antigoat secondary antibody for total Hog1 and horseradish peroxidase-conjugated anti-rabbit secondary antibody for phosphorylated Hog1using ECL, western lightning<sup>TM</sup> chemiluminescence (PerkinElmer).

#### 3.2.3 RNA isolation and microarray analysis

RNA was isolated using the hot phenol method with some modifications as described previously (Laviña et al., 2013). Preparation of the cDNA targets, hybridization and washing of the DNA microarray, fluorescence intensity measurement and gene expression analyses were all carried out following the manufacturer's protocol (Ambion Inc., TX, USA) (Laviña et al., 2013). The microarray-based global gene expression analysis of the *ptp2* $\otimes$ *msg5* $\otimes$ double disruptant was performed in triplicates.

#### 3.2.4 Gene enrichment analysis using Genowiz

Microarray data were analyzed for distinctive expression using Genowiz<sup>™</sup> 4.0 (Ocimum Biosolutions, India). Replicated gene values were combined and mean values of the expression ratios were considered for the dataset while empty spots were removed by filtering. The data were normalized prior to statistical analysis using the LOWESS algorithm (Yang et al., 2002). To stabilize the variation in the dataset, log transformation (log2) and mean centering were performed to bring down the data distribution of dataset near to zero.

Fold change analysis was done for detection of highly expressed genes in which genes with 2 folds up-regulation or down-regulation were considered as differentially expressed. ANOVA and Student's t-test were applied on the data to further establish the highly expressed genes with high degree of confidence, where *p*-value < 0.05 was considered significant in all statistical analyses. Filtering for the highly fluctuated genes was conducted using the FOREST algorithm (Leman et al., 2007). Two unsupervised methods of clustering namely K-means and Ward's agglomeration method were employed to group the highly fluctuated genes into different clusters based on the gene expression similarity. Functional classification of the genes was performed using Gene Ontology (GO) and pathway analysis using the web-based Gene Ontology (GO) enrichment analysis tool GOstat (http://gostat.wehi.edu.au). Prediction of protein-protein interactions was done using the web based STRING 9.0 resource (http://string-db.org).

#### 3.2.5 Determination of mRNA level using quantitative PCR

Total RNA was isolated from yeast cells grown in YPD broth with or without calcium until mid-logarithmic phase using hot phenol extraction. First strand cDNA synthesis was done using the High Capacity cDNA Reverse Transcription kit (Applied Biosystems) and the synthesized cDNA was used as the template for real-time RT-PCR. Quantitative PCR was performed using 20 µl reaction mixtures on an Applied Biosystems 7300 Real-Time PCR System with SYBR<sup>®</sup> Green PCR Master Mix (Applied Biosystems). PCR was performed using the following amplification program: initial denaturation for 10 min at 95°C, followed by 40 cycles of denaturation for 15 s at 95°C and annealing for 1 min at 60°C. *ACT1* was used as the reference gene for normalization of mRNA levels. The RT-PCR primers used in this study were designed using Primer Express version 2.0 (Applied Biosystems) and are described on the Table 8.

Strains	Alias	Genotype	Source
SH6314	BY4742	MATα his $3\Delta 1$ leu $2\Delta 0$ lys $2\Delta 0$ ura $3\Delta 0$	Invitrogen
SH5209	BY5209 or FY833	MATa ura3-52 his3∆200 leu2∆1 lys2∆202 trp1∆63	NBRP, YGRC*
SH5210	BY5210 or FY834	$\dot{M}$ ATa ura3-52 his3 $\Delta$ 200 leu2 $\Delta$ 1 lys2 $\Delta$ 202 trp1 $\Delta$ 63	NBRP, YGRC*
SH6790	$ptp2\Delta$	$\dot{M}ATa \ ptp2\Delta::CgHIS3 \ ura3-52 \ his3\Delta200$ leu2 $\Delta1 \ lys2\Delta202 \ trp1\Delta63$	SH5209 disruptant
SH6791	$msg5\Delta$	MATα msg5Δ::CgLEU2 ura3-52 his3Δ200 leu2Δ1 lys2Δ202 trp1Δ63	SH5210 disruptant
SH6792	$ptp2\Delta msg5\Delta$	MATα ptp2Δ ::CgHIS3 msg5Δ::CgLEU2 ura3-52 his3Δ 200 leu2Δ 1 lys2Δ 202 trp1Δ 63	Meiotic segregant 1B from SH6790 x SH6791
SH6793	$ptp2\Delta msg5\Delta$	MATa ptp $2\Delta$ ::CgHIS3 msg $5\Delta$ ::CgLEU2 ura3- 52 his $3\Delta 200$ leu $2\Delta 1$ lys $2\Delta 202$ trp $1\Delta 63$	Meiotic segregant 2B from SH6790 x SH6791
SH8991	ptp2∆msg5∆ ssk2∆	MATα ptp2Δ ::CgHIS3 msg5Δ ::CgLEU2 ssk2Δ ::CgTRP1 ura3-52 his3Δ 200 leu2Δ 1 lys2Δ 202 trp1Δ 63	Trp <sup>+</sup> disruptant of SH6792 double disruptant using p1805 ( <i>CgTRP1</i> )
SH8992	ptp2∆msg5∆ ssk22∆	MATα ptp2Δ ::CgHIS3 msg5Δ ::CgLEU2 ssk22Δ ::CgTRP1 ura3-52 his3Δ 200 leu2Δ 1 lys2Δ 202 trp1Δ 63	Trp <sup>+</sup> disruptant of SH6792 double disruptant using p1805 ( <i>CeTRP1</i> )
SH8993	ptp2∆msg5∆ pbs2∆	MATα ptp2Δ ::CgHIS3 msg5Δ ::CgLEU2 pbs2Δ ::CgTRP1 ura3-52 his3Δ 200 leu2Δ 1 lys2Δ 202 trp1Δ 63	Trp <sup>+</sup> disruptant of SH6792 double disruptant using $p_{1805}$ ( <i>CgTRP1</i> )
SH8994	ptp2∆msg5∆ hog1∆	MATα ptp2Δ::CgHIS3 msg5Δ ::CgLEU2 hog1Δ ::CgTRP1 ura3-52 his3Δ 200 leu2Δ 1 lys2Δ 202 trp1Δ 63	Trp <sup>+</sup> disruptant of SH6792 double disruptant using $p_{1805}$ ( <i>CgTRP1</i> )
SH8995	ptp2∆msg5∆ bcy1∆	MATα ptp2Δ ::CgHIS3 msg5Δ ::CgLEU2 bcy1Δ ::CgTRP1 ura3-52 his3Δ 200 leu2Δ 1 lys2Δ 202 trp1Δ 63	Trp <sup>+</sup> disruptant of SH6792 double disruptant using p1805 ( <i>CgTRP1</i> )
SH8996	ptp2∆msg5∆ msn2∆	MATα ptp2Δ ::CgHIS3 msg5Δ ::CgLEU2 msn2Δ ::loxP-ScURA3-loxP ura3-52 his3Δ 200 leu2Δ 1 lys2Δ 202 trp1Δ 63	$Ura^+$ disruptant of SH6792 double disruptant using $n_{3276}$ ( <i>ScUR4</i> 3)
SH8997	$msn4\Delta$	MATa msn4∆∷loxP-CgTRP1-loxP ura3-52 his3∆200 leu2∆1 lys2∆202 trp1∆63	Trp <sup>+</sup> disruptant of BY5209 using p3010 ( <i>CgTPR1</i> )
SH8998	ptp2∆msg5∆ msn4∆	MATα ptp2Δ ::CgHIS3 msg5Δ ::CgLEU2 msn4Δ ::loxP-CgTRP1-loxP ura3-52 his3Δ 200 leu2Δ 1 lys2Δ 202 trp1Δ 63	Meiotic segregant 3C from SH8996 x SH8997
SH8999	ptp2∆msg5∆ msn2∆msn4∆	MATa $ptp2\Delta$ ::CgHIS3 msg5 $\Delta$ ::CgLEU2 msn2 $\Delta$ ::loxP-ScURA3-loxP msn4 $\Delta$ ::loxP- CgTRP1-loxP ura3-52 his3 $\Delta$ 200 leu2 $\Delta$ 1 lys2 $\Delta$ 202 trp1 $\Delta$ 63	Meiotic segregant 4D from SH8996 x SH8997

Table 6. S. cerevisiae strains used in the study

KF-PTP2       5' ATAACGGCAATAGAATGGCTTCTTCCGCTATATCGGAAAACAC AGGAAACAGCTATGACC 3'         Kr-PTP2       5' GTAGCAATATACTTGAAATCAGGATTAATTTGCGTGAGCTGTT GTAAAACGACGGCCAGT 3'         Kfe-PTP2       5' CTCAAGCTTATTCGGTATTGGCACACCGCGTTCCTTACAAAAC ACGGAAACAGCTATTGGCACACCGCGTTCCATTACAAAAC ACAGGAAACAGCTATGACC 3'         Kr-MSG5       5' TCGTTGTCACAGAAGCTACAGCGGGAATCTGCGGGGTTGAG GTTGTAAAACGACGGCCAGT 3'         Kr-MSG5       5' TCCGGATCCTGAGTGATGGATAATGTGATTT 3'         Kre-MSG5       5' CTCGGATCCTGAGGATAGGATAATGTGATTT 3'         Kre-MSG5       5' CTCGGATCCTGAGGAAGCGCCATGGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCCCATGGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCCAG 3'         gd SSK2       5' AAAAGAAGAAAGCCAAATCTGC 3'         gd SSK2       5' ACTAAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGGTGAAA CCACAGGAAACAGCCAAATCTGC 3'         co SSK2P       5' TGTAAAAGGACGCAAATCTGC 3'         gd SSK2       5' AATATACGTGGGTATAAAAGGACGCTTCTG 3'         gd SSK2P       5' TATATATCGTAGTATATCATATTTTAAGACGGTGAGCACT GTTGTAAAACGAACGGCCAGT 3'         co SSK2P       5' CTTCCTGGGAAGTTGAGCC 3'         gd SSK2P       5' ATTATTATATGTAGGAACGCCACG 3'         gd PBS2       5' ATTATTAACTAGTGAGCCACC 3'         gd PBS2       5' ATTATTAACTAGTGACGCCAGT 3'         co SSK2P       5' TGTCTACTAGTGACGCGCAGT 3'         gd HOGIF       5' AACAAGAAGCAGCTATGACCG 3'         gd HOSIF       5' AACAAGAAGAGCTATGACGCCAGT 3' </th <th>Primer name</th> <th>Primer sequence</th>	Primer name	Primer sequence
AGGAAACAGCTATGACC 3' Kr-PTP2 5' GTAGCAATATACTTGAAATCAGGATTAATTTGCGTGAGCTGTT GTAAAACGACGGCCAGT 3' Kfe-PTP2 5' CTCAAGCTTGAGCACTCGTTTAATTTAGCCA 3' Kre-PTP2 5' CTCAAGCTTATCGGTATTGGCACAAACTTT 3' Kf-MSG5 5' ACATCGATTTCAAGCCAAACTCACCGCGGTTCCTTACAAAAC ACAGGAAACAGCTATGACC 3' Kr-MSG5 5' TCGGTGTCCCACGAGAGCTTCCAGTGAATCTGCGGGGTTGAG GTTGTAAAACGACGGCCAGT 3' Kfe-MSG5 5' CTCGGATCCGTAGTGATGGATGAATGTGATTT 3' Kre-MSG5 5' CTCGGATCCGTGGCCATGGTAAATTTTGACAGG 3' gd SSK2 5' AAAAGAAGAGAAGCCTTTGCGTAAATTTTGACAGGCAC AAATACTTCGTACGCTGCGAG 3' gd SSK2 5' AAAAGAAGAAGCCATTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3' co SSK2F 5' TAGAAAGAAGCCAAATCTGC 3' gd SSK22 5' AAAAGAAGAGGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3' co SSK22 5' ACATAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3' gd SSK22F 5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCCTATGACC 3' gd SSK22F 5' GAAATTGTTAGGATAACCC 3' co SSK22F 5' GAAATTGTAGAGAAGCCCAGT 3' co SSK22F 5' GAAATTGTTAGGAAAACCC 3' gd PBS2 5' ATTATATCATAGCAGGCCAGT 3' co SSK22F 5' AAAGGAGAAGCCTTGGCCCAGT 3' co SSK22F 5' AAAGGAGAACAGCTATGACC 3' gd PBS2 5' ATTATATCATGTAGCAGCC 3' gd PBS2 5' TATATATCAGGTGGACCTTGAGCGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3' gd PBS2 5' TATATTCACGTGCCCGTGTTGCTTTATTGGATATTACG GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' GCAATATATGACGTCGAGGACGTTAAATTCCCAAGG CACAGGAAACAGCTATGACC 3' gd HOGIF 5' AAAGGGAAAACAGGGAAACTACAACTATCGTATATAATA CACAGGAAACAGGCTATGACC 3' gd HOGIF 5' AAAGGGAAAACAGGGAAACTACAACTATCGTATATATAA CACAGGAAACAGGCTATGACC 3' gd HOGIF 5' AAAGGGAAAACAGGGCAATTAAACCAACGTATGACATACAACACGT GTTGTAAAACGACGGCCAGT 3' co HOGIF 5' GTCTGGCGGCGTTATTATAC 3' co HOGIF 5' GTCTGGCGGCGCTATTATATCA 3' gd BCY1IF 5' TACAACAAGGAAACAGGGAAACTACAACACAGTAAGAATAA ACCGCTCGTACGACGTGCGAG 3' gd BCY1F 5' AAAGGAAAGGAAATTCATGTGGGACTTAAAGAACAACAGGTAAGAATAA ACCGCTCGACCTAGTGGGACTT3'COCCTTTCCCCTTTT TACCGCCACTAGTGGGAACTTATGCAT 3' co BCY1F 5' ATAAGCTAAGGAACTTATTGACT 3' co BCY1F 5' ATAAGCTAAGGAAATTACACGTGGACTTAACGACGCTTCCCCTTTT TACCGCACTAGTGGAACTTATGACT 3' co BCY	Kf-PTP2	5' ATAACGGCAATAGAATGGCTTCTTCCGCTATATCGGAAAACAC
<ul> <li>Kr-PTP2</li> <li>S' GTAGCAATATACTTGAAATCAGGATTAATTTGCGTGAGCTGTT GTAAAACGACGGCCAGT 3'</li> <li>Kfe-PTP2</li> <li>S' CTCAAGCTTGGACACTCGTTTAATTTAGCCA 3'</li> <li>Kre-PTP2</li> <li>S' CTCAAGCTTGGACACTCGTTTAATTGGCACAAACTTT 3'</li> <li>Kf-MSG5</li> <li>S' ACATCGATTTCAAGCCAAACTCCACCGCGTTCCTTACAAAAC ACAGGAAACAGCTATGACC 3'</li> <li>Kr-MSG5</li> <li>S' CTCGGATCCGTAGTGATGGATAATGTGATTT 3'</li> <li>Kfe-MSG5</li> <li>S' CTCGGATCCGTAGTGATGGATAATGTGATTT 3'</li> <li>Kre-MSG5</li> <li>S' CTCGGATCCGTGCCCATGGTAAATGTGATTT 3'</li> <li>Kre-MSG5</li> <li>S' CTCGGATCCGTGCCCATGGTAAATGTGATTT 3'</li> <li>Kre-MSG5</li> <li>S' CTCGGATCCGTGCCCATGGTAAATGTGATATTTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'</li> <li>gd SSK2</li> <li>S' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'</li> <li>co SSK2F</li> <li>S' TGTAAAAGCGAAGCCAAATCTGC 3'</li> <li>co SSK22R</li> <li>S' TATATAGGTGGTGACTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'</li> <li>gd SSK22F</li> <li>GAAATTGTTAGGAAGCCAATCTGC 3'</li> <li>co SSK22R</li> <li>S' CTTCCTGGGAAGTTGAGCC 3'</li> <li>gd PBS2</li> <li>S' TATATTGTTAGGAAACCC 3'</li> <li>co SSK22R</li> <li>CTTCCTGGGAAGTTGAGCC 3'</li> <li>gd PBS2</li> <li>S' TATATTGATGAGCGCCAGT 3'</li> <li>co PBS2F</li> <li>S' GTCTACTAGTGAGCCGCAGT 3'</li> <li>co PBS2F</li> <li>S' ACACAATATATTGACGTCCA 3'</li> <li>gd HOGIF</li> <li>AAAGGAAAACAGCTATGACC 3'</li> <li>gd HOGIF</li> <li>AAAGGAAAACAGCTATGACCA 3'</li> <li>gd HOGIF</li> <li>AAAGGAAACAGCTATATTTTCTAAAAAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOGI R</li> <li>AGATAGAATGAGTGATAAAC3'</li> <li>gd BCY1IF</li> <li>TACAACAAGCAGGATATTTTTCAAAAGAACAACAGGTAAGAATAA ACCCCCACTAGTGGATTATAAC 3'</li> <li>gd BCY1IF</li> <li>TACAACAAGCAGATATTCATGTGGAACTTAAGAACAACAGGTAAGAATAA ACCCCCACTAGTGGAACTTATGAT 3'</li> <li>co BCY1F</li> <li>ATAAGGTGAAATTCATGTGGGAACTTAAGAACAACAGGAAATAA ACCCCCACTAG</li></ul>		AGGAAACAGCTATGACC 3'
GTAAAACGACGGCCAGT 3' Kfe-PTP2 5' CTCAAGCTTGGACACCGTTTAATTTAGCCA 3' Kre-PTP2 5' CTCAAGCTTATTCGGTATTGGCACAAACTTT 3' KF-MSG5 5' CTCGATTTCAAGCCAAACTCACCGCGGTTCCTTACAAAAC ACAGGAAACAGCTATGACC 3' Kr-MSG5 5' CTCGGATCCGTAGTGATGGATATGTGATTT 3' Kre-MSG5 5' CTCGGATCCGTGCCCATGGTAATGTGATTT 3' Kre-MSG5 5' CTCGGATCCGTGCCCATGGTAATGTGATTT 3' Kre-MSG5 5' CTCGGATCCGTGCCCATGGTAAATGTGATTT 3' Kre-MSG5 5' CTCGGATCCGTGCCCATGGTAAATTTTTGACAGGCAC AAATACTTCGTACGCTGCGC3 3' gd SSK2 5' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3' gd SSK2 5' TAGAAAGAAGAGCCAAATCTGC 3' co SSK2F 5' TAGAAAGAAGCCATATGCG 3' co SSK2F 5' TAGAAAGAAGCCATATGACG 3' gd SSK2 5' TGTTAAAACGGATGTCTCT 3' gd SSK22 5' AATATCGTAGGTGATATAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3' gd SSK22 5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3' co SSK22F 5' GAAATTGTAGGATGAGCC 3' gd PSS2 5' ATTATTATATATAAGCAGATCGGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3' gd PSS2 5' ATTATTATATATAAGCAGATCGGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3' gd PBS2 5' ATTATTATATATAAGCAGGCCAGT 3' co PBS2F 5' TGTCACTGGGCGCTGTTGCCTTTTATTGGATATTACG GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' TGTCACTAGTGACCACT GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' TGTCACTAGTGACCACT GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' TGTCACTAGTGACCACT GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' TGTCACTAGTGACCACT GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' TGTCACTAGTGACCACCACT GTTGTAAAACGACGGCCAGT 3' co HOGI F 5' AAAGGGAAAACAGGGAAAACTACAACTATAGTATATATA CACAGGAAACAGGCTATGACCA 3' gd HOGI F 5' AAAGGGAAAACAGGGAAAACTACAACATATAAAACACGT GTTGTAAAACGACGGCCAGT 3' co HOGI F 5' GACTAGAAGGCATTATATTACA 3' gd BCY1F 5' TACAACAAGCAGATATATTTCAAAAGACAACAGTAAGAATAA ACCGCTACGTACGACGCCGCGCTTATTATACA 3' gd BCY1F 5' ATAAGCGAAAGGAAATCATGTGTGGATTTAAGATCGCTTCCCCTTTT TACCCCACACTAGTGGAACTTATCACAGACGCCACACAGAAAGAA	Kr-PTP2	5' GTAGCAATATACTTGAAATCAGGATTAATTTGCGTGAGCTGTT
<ul> <li>Kte-PTP2</li> <li>S' CTCAAGCTTGGACACTCGTTTAATTTAGCCA 3'</li> <li>Kre-PTP2</li> <li>S' CTCAAGCTTATTCGGTATTGGCACAACTTT 3'</li> <li>Kf-MSG5</li> <li>S' ACATCGATTTCAAGCCAAGCTACCGCGTTCCTTACAAAAC ACAGGAAACAGCTATGACC 3'</li> <li>Kre-MSG5</li> <li>S' TCGTTGTCCACAGAAGCTTCCAGTGAATCTGCGGGGTTGAG GTTGTAAAACGACGGCCAGT 3'</li> <li>Kre-MSG5</li> <li>S' CTCGGATCCGTGCGAGTGGATGATATGTGATTT 3'</li> <li>Kre-MSG5</li> <li>S' CTCGGATCCGTGCGCATGGTAATTTTGACAG 3'</li> <li>gd SSK2</li> <li>S' AAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'</li> <li>gd SSK2</li> <li>S' AAAGAAGAGAAGCCATTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'</li> <li>co SSK2F</li> <li>S' TAGAAAGAAGCCAAATCTGC 3'</li> <li>co SSK2F</li> <li>S' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'</li> <li>gd SSK22F</li> <li>S' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'</li> <li>gd SSK22F</li> <li>S' ATTATGTATGTATGATATCATATTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'</li> <li>co SSK22F</li> <li>S' GAAATTGTTAGGAAACCC 3'</li> <li>gd PBS2</li> <li>S' ATTATTATATTAAGCAGAGTCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2</li> <li>S' ATTATTATATTAAGCAGGCCAGT 3'</li> <li>co PBS2F</li> <li>S' TGTCTACTAGTGGCGATTT 3'</li> <li>co PBS2F</li> <li>S' ACACAATATATTGACGTCCA 3'</li> <li>gd HOGIF</li> <li>S' GAACTAAGAAGAGCTATGACC 3'</li> <li>gd HOGIF</li> <li>S' GAACTAAGAAGCGATAGGCCAGT 3'</li> <li>co HOGIR</li> <li>S' GAACAAGCTATGACTGACAACATCAACTATCGTATATATA</li></ul>		GTAAAACGACGGCCAGT 3'
<ul> <li>Kre-PTP2 5' CTCAAGCTTATTCGGTATTGGCACAAACTTT 3'</li> <li>KF-MSG5 5' ACATCGATTTCAAGCCAACTCACCGCGTCCTTACAAAAC ACAGGAAACAGCTATGACC 3'</li> <li>Kr-MSG5 5' CTCGGATCCGTAGTGATGGATAGTGTATTTGAACG 3'</li> <li>gd SSK2 5' AAAAGAAGAAAGCCTTTGCGTAATTTTGAACG 3'</li> <li>gd SSK2 5' AAAAGAAGAAGACCCTTGCGTAATTTTGAACGGCAC AAATACTTCGTACGCTGCAG 3'</li> <li>gd SSK2 5' AAAAGAAGAAGACAGCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'</li> <li>gd SSK2 5' AAAAGAAGAGAAGCCATTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'</li> <li>go SSK2F 5' TAGAAAGAAGAGAGCATTTCG 3'</li> <li>co SSK2F 5' TAGAAAGAAGCAATCTGC 3'</li> <li>co SSK22F 5' ACTAAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'</li> <li>gd SSK228 5' TATATATCGTAGTGATGAACCC 3'</li> <li>gd SSK228 5' CTTCTCTGGGAAGTTGAGCC 3'</li> <li>gd SSK228 5' CTTCTCTGGGAAGTTGAGCC 3'</li> <li>gd PBS2 5' ATTATTATATTAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' ATTATTATATATAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' ACACAATATATTGAAGTCGCCAGT 3'</li> <li>co PBS2F 5' TGTCTACTAGTGAGGCCAGT 3'</li> <li>co PBS2R 5' ACACAATATATTGACGTCGACACTATCGTATATAACG GTTGTAAAACGACGGCCAGT 3'</li> <li>gd HOGIF 5' AAAGGAAAACAGCTATGACC 3'</li> <li>gd HOGIF 5' GAAGTAGAGAGGGCAGTTAACTATCGTATATAATA CACAGGAAACAAGCTATGACC 3'</li> <li>gd HOGIF 5' GAAGTAAGAATGAGTGGTTGAGGACATTAAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOGIF 5' GAAGTAAGAATGAGTGGTTAGGCTAAGACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'</li> <li>gd HOGIF 5' AAAGGAAAACAGCAATATTTTCAAAAGAACAACAGCTATGACC 3'</li> <li>gd HOGIF 5' GAAGTAAGAAGAGCATATTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGAGCGCAGT 3'</li> <li>co HOGIF 5' GTCTGCCGGGCGTTATTTTCAAAAGACAACAGGTAAGAATAA ACGCTTCGTACGGCGCAGT 3'</li> <li>gd BCYIR 5' GAAGAAGGAAATTCATGTGGATATAACGATCGCTTCCCCTTTT TACGCCACTAGTGGAGTTATTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGTGCGAATATTTCAAAAGACAACAGGTAAGAATAA ACGCTTCGTACGTGGAATATTCATGGGATTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGAATTATTCAAAAGACAACAGCTAAGAATAA ACGCTTCGTACGTAGAATTATTCATGGAATTAACGATGGTTAAGTGGTTCCCCTTTT TACGCCACTAGTGGAAT</li></ul>	Kfc-PTP2	5' CTCAAGCTTGGACACTCGTTTAATTTAGCCA 3'
Kf-MSG55' ACATCGATTTCAAGCCAAACTCACCGCGTTCCTTACAAAAC ACAGGAAACAGCTATGACC 3'Kr-MSG55' TCGTTGTCCACAGAAGCTTGCAGTGAATCTGCGGGGTTGAG GTTGTAAAACGACGGCCAGT 3'Kfc-MSG55' CTCGGATCCGTGGCCATGGTAATTTTGACG 3'gd SSK25' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'gd SSK25' AAAAGAAGAGAAGCCATTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'gd SSK25' TAGAAAGAAGAAGCCAAATCTGC 3'co SSK2F5' TGTTAAAAGCGATGTCTTCT 3'gd SSK225' ACTTAGGGTGGCTATAAAAGGAAGTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'gd SSK2285' TGTTAAAAGCGATGTCTTCT 3'gd SSK2285' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'co SSK2285' CTCTCTGGGAAGTTGAGCC 3'gd PBS25' ATATTATTATTAAGCAGATCGAGACGTTAATTCATATCATATTCTAAAG CACAGGAAACAGCTATGACC 3'gd PBS25' TATATTATCACGTGCCTGTTTGCTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'co PBS285' TGTCTACTAGTGAGCGATTTGACCACT GTTGTAAAACGACGGCCAGT 3'co PBS295' ACACAATATTTGACGTCCA 3'gd HOGIF5' GAAGTAAGAATGAGGGAAAACTACAACTATCGTATATAAAA CACAGGAAACAGCTATGACC 3'gd HOGIR5' GAAGTAAGAATGAGGGGACATTAAAACAACGT GTTGTAAAACGACGGCCAGT 3'co HOGIR5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCAGGT 3'gd HOGIR5' GAAGAAACGAGGTATGAACC 3'gd HOGIR5' GAGAAAGCAGCTATGATATATCA ACGCTTCGTACGTGGAGGTATATATACAGACGACAACAGTAAGAATAA ACGCTTCGTACGTGGATATATTTTCAAAAGACAACAGTAAAAAACACGT GTGGAAAGGAAATTCATGTGGAAGTACAACAGGTAAGAATAA ACGCTTCGTACGTGGATATATCATGGGACATTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'co BCYIF5' ATAAGCAGACACTTATTG	Krc-PTP2	5' CTCAAGCTTATTCGGTATTGGCACAAACTTT 3'
Kr-MSG55' TCGTTGTCCACAGAAGCTTCCAGTGAATCTGCGGGTTGAG GTTGTAAAACGACGACGGCCAGT 3'Kfe-MSG55' CTCGGATCCGTAGTGATGGATAATGTGATTT 3'Kre-MSG55' CTCGGATCCGTGCCCATGGTAATTTTGACG 3'gd SSK25' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'gd SSK25' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'gd SSK25' TAGAAAGAAGCAAACCTTGC 3'co SSK2R5' TGTTAAAAGCGATGTCTTC 3'gd SSK22F5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'gd SSK22R5' TATATATCGTAGTATATCATATTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'co SSK22R5' CTCTCTGGGAAGTTGAGCC 3'gd SSK22F5' GAAATTTGTTAGGAAAACCC 3'co SSK22R5' CTTCTCTGGGAAGTTGAGCC 3'gd PBS25' ATTATATATAACGACGCCGCAGT 3'co SSK22R5' TATATATCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGCCACT 3'gd PBS25' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'co PBS2F5' TGTCTACAGTGAGCGCAGT 3'co PBS2F5' ACACAATATATTGACGTCCA 3'gd HOGIF5' AAAGGGAAAACAGGGAAAACACACAACTACAACTATCGTATATAATA CACAGGAAACAGGGGAAACTACAACTACAACTACAACAACACGT GTTGTAAAACGACGGCCACGT 3'co HOGI R5' GAGTAAGAATGAGTGGTATAACG 3'gd BCY1IF5' GAGAAAGGAAATCATGTGGAATTAAAGAACAACAGCTATGGCAATTAAAGAACAACAGCTATGGAATTAAAGAACAACAGCTATGGCTGCAG 3'gd BCY1R5' GAGAAAGGAAATCATGTGGAATTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGAATCATGGATTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGAATCATGATGAT 3'co BCY1F5' ATTTTGTATGTCACATTGATTAGT CO BCY1Rco BCY1R5' TTTTTGTATGTCACTTGATTAACT CO BCY1R	Kf-MSG5	5' ACATCGATTTCAAGCCAAACTCACCGCGTTCCTTACAAAAC ACAGGAAACAGCTATGACC 3'
Kfc-MSG55' CTCGGATCCGTAGTGATGGATGGATAATGTGATTT 3'Krc-MSG55' CTCGGATCCGTGCCCATGGTAATTTTGACG 3'gd SSK25' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'gd SSK25' AAAAGAAGAGACAGCCATTGCGTGAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'co SSK2F5' TAGAAAGAAGCCAAATCTGC 3'co SSK2F5' TGTTAAAAGCGATGTCTTCT 3'gd SSK22F5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'gd SSK22F5' ACTTAGGGTGGCTATAAAAGGCAGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'gd SSK22R5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'co SSK22R5' CTTCTCTGGGAAGTTGAGCC 3'gd PBS25' ATTATTATATATAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'gd PBS25' TATATTCACGTGCCGTGTTGCTTTTATTGGATATTAACG GTTGTAAAACGACGGCAGT 3'co PBS2F5' TGCTACTAGTGAGCGGCAGT 3'co PBS2F5' ACACAATATATTGACGTCCA 3'gd HOG1F5' AAAGGGAAAACAGGCTATGACC 3'gd HOG1F5' AAAGGGAAAACAGGCTATGACC 3'gd HOG1F5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCAGT 3'co HOG1 R5' AGATAGAAGTGGTGATAAAC 3'gd BCY11F5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCGCGCAG 3'gd BCY11F5' GAGAAAGGAAATCATCATGTGGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGAATCATGATGAT 3'co BCY1F5' ATAGCTGAACTACTGATTGAT 3'co BCY1F5' ATAGCTGAACTACGCATCGATTGAT 3'co BCY1F5' ATAGCTGAACTACGATTGAT 3'co BCY1F5' ATAGCTGAACTACGACTTGATTGAT 3'co BCY1F5' ATAGCTGAACTACGACTTGATTAAC	Kr-MSG5	5' TCGTTGTCCACAGAAGCTTCCAGTGAATCTGCGGGTTGAG GTTGTAAAACGACGGCCAGT 3'
Kre-MSG55' CTCGGATCCGTGCCCATGGTAATTTTTGACG 3'gd SSK25' AAAAGAAGAGAAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'gd SSK25' AAAAGAAGAGAAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'co SSK2F5' TAGAAAGACGAAACTGTCTTC 3'gd SSK22F5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'gd SSK22R5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'co SSK22R5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'co SSK22R5' CTCTCTCTGGGAAGTTGAGCC 3'gd PBS25' ATTATTATATTAAGCAGAGCGGCCAGT 3'co SSK2285' CTTCTCTGGGAAGTTGAGCC 3'gd PBS25' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG CACAGGAAACAGCTATGACC 3'gd PBS25' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'co PBS2F5' TGTCTACTAGTGAGCGAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'gd HOGIF5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'gd HOGIF5' AGATAGAAGGCGAGTTAATTTTCCAAAGACACGT GTTGTAAAACGACGGCCAGT 3'co HOGIF5' GTCTGGCGGCGTTATTATAC 3'co HOGIF5' AGATAGAAGGCGAGTATATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'gd BCY11F5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'co BCY1F5' ATAAGCTGAACTATATGCAT 3'co BCY1F5' ATTTTGTATGTCATGTGATATA'	Kfc-MSG5	5' CTCGGATCCGTAGTGATGGATAATGTGATTT 3'
gd SSK2 5' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3' gd SSK2 5' AAAAGAAGAGAAAGCCTATGCGAG 3' co SSK2F 5' TAGAAAGAGCGAATCTGC 3' gd SSK22F 5' TGTAAAAGCGATGTCTTCT 3' gd SSK22F 5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGAACC 3' gd SSK22R 5' TATATATCGTAGTATATCATATTTTTAGACGGTGACACT GTTGTAAAACGACGGCCAGT 3' co SSK22F 5' GAAATTGTTAGGAAAACCC 3' gd PSS2 5' CTTCTCTGGGAAGTTGAGCC 3' gd PBS2 5' TATATATATATATATATATATATTGACGGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3' gd PBS2 5' TATATATCATGTGGGCCAGT 3' co PBS2F 5' TGTCTACTAGTGGCCAGTGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3' gd PBS2 5' TATATTCACGTGCCGTTTGCTTTTATTGGATATTAACG GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' TGTCTACTAGTGAGCGAATT 3' co PBS2R 5' ACACAATATATTGACGGCCAGT 3' gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3' gd HOG1F 5' GAAGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3' gd HOG1F 5' GAAGTAGAAGGGTAAACTACAACTATAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3' co HOG1F 5' GAAGTAGAATGAGTGGTTAGGGACATTAAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3' co HOG1F 5' GAATAGAATGGCTGATAAAC 3' gd BCY11F 5' TACAACAAGAGAGATTATTTCCAAAAGACAACAGCTATGAACAACAGCTACGAACAGCAACAGCTATGAACAACAGCTTTC TACGCCACTAGTGGGATTATCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCA 3' co BCY1R 5' ATAAGCTGACGACTTATAACT 3' co BCY1R 5' TATAGCTGAACTACTACTATGCAT 3'	Krc-MSG5	5' CTCGGATCCGTGCCCATGGTAATTTTTGACG 3'
gd SSK2 5' AAAAGAAGAAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3' co SSK2F 5' TAGAAAGAAGCCAAATCTGC 3' gd SSK22F 5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3' gd SSK22R 5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3' co SSK22R 5' CATCTCTGGGAAGTTGAGCC 3' gd PBS2 5' AATTATTATATAAGAGAGACGACGATCAAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3' gd PBS2 5' ATTATTATATAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3' gd PBS2 5' TATATTCACGTGCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' TGTCTACTAGTGAGCGGCAGT 3' co PBS2R 5' ACACAATATATTGACGTCCA 3' gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAACA CACAGGAAACAGCTATGACC 3' gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3' gd HOG1F 5' AGAATAGAATGAGTGGTTAGGGACATTAAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3' co HOG1 F 5' GTCTGGCGGCGTTATTATAC 3' co HOG1 F 5' AGATAGAAGTGCTGATAAAC 3' gd BCY1IF 5' TACAACAAGCAGAGTTAATTTTCCAAAAGAACAGCTATCGTATAAAAAAACAACGATCGTGTGAGACAACAAGAACAGCTATGACC 3' gd BCY1R 5' GAGAAAGGAAATCATCAGAGTTAAAGAACAGCATCAAAACAAGAACAAGAATAA ACGCTTCGTACGCTGCAG 3' gd BCY1R 5' GAGAAAGGAAATTCATGTGGAATTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3' co BCY1F 5' ATAAGCTGAACTTATTGCAT 3' co BCY1F 5' ATAAGCTGAACTTATTGCACT 3'	gd SSK2	5' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'
<ul> <li>co SSK2F 5' TAGAAAGAAGCCAAATCTGC 3'</li> <li>co SSK2R 5' TGTTAAAAGCGATGTCTTCT 3'</li> <li>gd SSK22F 5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'</li> <li>gd SSK22R 5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'</li> <li>co SSK22R 5' CTTCTCTGGGAAGTTGAGCC 3'</li> <li>gd PBS2 5' ATTATTATATTAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' ATTATTATATATAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'</li> <li>co PBS2F 5' TGTCTACTAGTGAGCGATTT 3'</li> <li>co PBS2R 5' ACACAATATATTGACGTCCA 3'</li> <li>gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'</li> <li>gd HOG1F 5' GACTAGAAGAGTGCTGATATAAC3'</li> <li>co HOG1 F 5' GTCTGGCGGCGTTATTATATC 3'</li> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAAACAAGCAGATTATTTTCAAAAAGACAACAGTAAGAATAA ACGCTTCGTACGAGCTATGACC 3'</li> <li>gd BCY11F 5' GAAGTAGAATGCTGATATAAC 3'</li> <li>gd BCY11F 5' GAAGTAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11R 5' GAGAAAGCAGAGATTATTATCATGTGGAATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGAACTATGAGCA 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATGCAT 3'</li> <li>co BCY1R 5' TTTTTGTATGTCACTTGATT 3'</li> </ul>	gd SSK2	5' AAAAGAAGAGAAGCCTTTGCGTAAACTATTTGACAGGCAC AAATACTTCGTACGCTGCAG 3'
<ul> <li>co SSK2R 5' TGTTAAAAGCGATGTCTTCT 3'</li> <li>gd SSK22F 5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'</li> <li>gd SSK22R 5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'</li> <li>co SSK22F 5' GAAATTTGTTAGGAAAACCC 3'</li> <li>co SSK22R 5' CTTCTCTGGGAAGTTGAGCC 3'</li> <li>gd PBS2 5' ATTATTATATATAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' ATTATTATATATAGCAGATCGAGACGTTAATTAACG GTTGTAAAACGACGGCCAGT 3'</li> <li>co PBS2F 5' TGTCTACTAGTGAGCGATTT 3'</li> <li>co PBS2R 5' ACACAATATATTGACGTCCA 3'</li> <li>gd HOG1F 5' AAAGGGAAAACAGGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'</li> <li>gd HOG1R 5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOG1F 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCGCGATC 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATCAGTGGATTAAAGATCACCTTTC TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGGTGAAGTGCTCATT3'</li> </ul>	co SSK2F	5' TAGAAAGAAGCCAAATCTGC 3'
gd SSK22F5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA CACAGGAAACAGCTATGACC 3'gd SSK22R5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'co SSK22F5' GAAATTTGTTAGGAAAACCC 3'co SSK22R5' CTTCTCTGGGAAGTTGAGCC 3'gd PBS25' ATTATTATATATAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'gd PBS25' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'co PBS2F5' TGTCTACTAGTGAGCGATTT 3'co PBS2R5' ACACAATATATTGACGTCCA 3'gd HOGIF5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'gd HOGIR5' GAAGTAAGAATGAGTGGTTAAGGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'co HOG15' GAAGTAAGAATGAGTGCTGATAAAC 3'gd BCY11F5' TACAACAAGCAGAGTGCTGATAAAC 3'gd BCY11R5' GAGAAAGGAAATCATGTGGAATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'co BCY1F5' ATAAGCTGAACTTATTGCAT 3'co BCY185' TTTTTGTATGTCACTTGATT 3' TTTTTGTATGTCAATTGCAT 3'	co SSK2R	5' TGTTAAAAGCGATGTCTTCT 3'
CACAGGAAACAGCTATGACC 3'gd SSK22R5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAAACGACGGCCAGT 3'co SSK22F5' GAAATTTGTTAGGAAAACCC 3'co SSK22R5' CTTCTCTGGGAAGTTGAGCC 3'gd PBS25' ATTATTATATTAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'gd PBS25' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'co PBS2F5' TGTCTACTAGTGAGCGATTT 3'co PBS2R5' ACACAATATATTGACGTCCA 3'gd HOG1F5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'gd HOG1R5' GAAGTAAGAATGAGTGGTTATGGACATTAAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'co HOG1 R5' AGATAGAAGTGCTGATAAAC 3'gd BCY11F5' TACAACAAGCAGGATATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCAGCAGT 3'co BCY1F5' ATAAGCTGAACTATTGCAT 3'co BCY1F5' ATAAGCTGAACTTATTGCAT 3'co BCY1R5' TTTTTGTATGTACCACTTGATT 3'	gd SSK22F	5' ACTTAGGGTGGCTATAAAAGGTAGTTCCTTGTAGGTGAAA
<ul> <li>gd SSK22R 5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT GTTGTAAAACGACGGCCAGT 3'</li> <li>co SSK22F 5' GAAATTTGTTAGGAAAACCC 3'</li> <li>co SSK22R 5' CTTCTCTGGGAAGTTGAGCC 3'</li> <li>gd PBS2 5' ATTATTATATTAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'</li> <li>co PBS2F 5' TGTCTACTAGTGAGCGATTT 3'</li> <li>co PBS2R 5' ACACAATATATTGACGTCCA 3'</li> <li>gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'</li> <li>gd HOG1R 5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOG1 F 5' AGATAGAATGAGTGGTTATTATAC 3'</li> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGAGTTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTTGCAT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTATCCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> </ul>	0	CACAGGAAACAGCTATGACC 3'
GTTGTAAAACGACGGCCAGT 3' co SSK22F 5' GAAATTTGTTAGGAAAACCC 3' gd PBS2 5' ATTATTATATTAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3' gd PBS2 5' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3' co PBS2F 5' TGTCTACTAGTGAGCGATTT 3' co PBS2R 5' ACACAATATATTGACGTCCA 3' gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3' gd HOG1R 5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3' co HOG1F 5' GTCTGGCGGCGTTATTATAC 3' co HOG1 F 5' AGATAGAAGTGCTGATAAAC 3' gd BCY11F 5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3' gd BCY11F 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3' co BCY1F 5' ATAAGCTGAACTTATTGCAT 3' co BCY1F 5' ATAAGCTGAACTTATTGCAT 3'	gd SSK22R	5' TATATATCGTAGTATATCATATTTTTAGACGTTGACCACT
<ul> <li>co SSK22F 5' GAAATITIGITAGGAAAACCC 3'</li> <li>co SSK22R 5' CTTCTCTGGGAAGTTGAGCC 3'</li> <li>gd PBS2 5' ATTATTATATATAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'</li> <li>co PBS2F 5' TGTCTACTAGTGAGCGATTT 3'</li> <li>co PBS2R 5' ACACAATATATTGACGTCCA 3'</li> <li>gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'</li> <li>gd HOG1R 5' GAAGTAAGAATGAGGGGCCAGT 3'</li> <li>co HOG1F 5' GTCTGGCGGCGTTATTATAC 3'</li> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGAGATTATTTTCCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATCATGTGGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTATGTCACTTGATT 3'</li> </ul>	GGWAAD	GTTGTAAAACGACGGCCAGT 3'
<ul> <li>co SSK22R 5' CTTCTCTGGGAAGTTGAGCC 3'</li> <li>gd PBS2 5' ATTATTATATATAAGCAGAATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'</li> <li>co PBS2F 5' TGTCTACTAGTGAGCGATTT 3'</li> <li>co PBS2R 5' ACACAATATATTGACGTCCA 3'</li> <li>gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'</li> <li>gd HOG1R 5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOG1F 5' GTCTGGCGGCGTTATTATAC 3'</li> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGAATTCATGTGGATTTAAGAACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTGCAT 3'</li> </ul>	co SSK22F	5' GAAATTTGTTAGGAAAACCC 3'
<ul> <li>gd PBS2 5' ATTATTATATTAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'</li> <li>gd PBS2 5' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'</li> <li>co PBS2F 5' TGTCTACTAGTGAGCGATTT 3'</li> <li>co PBS2R 5' ACACAATATATTGACGTCCA 3'</li> <li>gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGGCTATGACC 3'</li> <li>gd HOG1R 5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOG1F 5' GTCTGGCGGCGTTATTATAC 3'</li> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATCATGTGGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTGCAT 3'</li> </ul>	co SSK22R	5' CTTCTCTGGGAAGTTGAGCC 3'
gd PBS25' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'co PBS2F5' TGTCTACTAGTGAGCGATTT 3'co PBS2R5' ACACAATATATTGACGTCCA 3'gd HOG1F5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'gd HOG1R5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'co HOG1F5' GTCTGGCGGCGTTATTATAC 3'co HOG1R5' AGATAGAAGTGCTGATAAAC 3'gd BCY11F5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'gd BCY1R5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'co BCY1F5' ATAAGCTGAACTTATTGCAT 3'co BCY1R5' TTTTTGTATGTCACTTGATT 3'	gd PBS2	5' ATTATTATATTAAGCAGATCGAGACGTTAATTTCTCAAAG CACAGGAAACAGCTATGACC 3'
co PBS2F5' TGTCTACTAGTGAGCGATTT 3'co PBS2R5' ACACAATATATTGACGTCCA 3'gd HOG1F5' AAAGGGAAAACAGGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'gd HOG1R5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'co HOG1F5' GTCTGGCGGCGTTATTATAC 3'co HOG1 R5' AGATAGAAGTGCTGATAAAC 3'gd BCY11F5' TACAACAAGCAGAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'gd BCY1R5' GAGAAAGGAAATTCATGTGGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'co BCY1F5' ATAAGCTGAACTTATTGCAT 3'co BCY1R5' TTTTTGTATGTCACTTGATT 3'	gd PBS2	5' TATATTCACGTGCCTGTTTGCTTTTATTTGGATATTAACG GTTGTAAAACGACGGCCAGT 3'
<ul> <li>co PBS2R 5' ACACAATATATTGACGTCCA 3'</li> <li>gd HOG1F 5' AAAGGGAAAACAGGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'</li> <li>gd HOG1R 5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOG1F 5' GTCTGGCGGCGTTATTATAC 3'</li> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGAGTTATTTTCAAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTGCAT 3'</li> <li>co BCY18 5' TTTTTGTATGTCACTTGATT 3'</li> </ul>	co PBS2F	5' TGTCTACTAGTGAGCGATTT 3'
<ul> <li>gd HOG1F 5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'</li> <li>gd HOG1R 5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOG1F 5' GTCTGGCGGCGTTATTATAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGATGCTGATAAAC 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTATGTCACTTGATT 3'</li> </ul>	co PBS2R	5' ACACAATATATTGACGTCCA 3'
<ul> <li>gd HOG1R 5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'</li> <li>co HOG1F 5' GTCTGGCGGCGTTATTATAC 3'</li> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGATTATTTTCAAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTGCAT 3'</li> </ul>	gd HOG1F	5' AAAGGGAAAACAGGGAAAACTACAACTATCGTATATAATA CACAGGAAACAGCTATGACC 3'
<ul> <li>co HOG1F 5' GTCTGGCGGCGTTATTATAC 3'</li> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGATTATTTTCAAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTGCAT 3'</li> <li>co BCY1R 5' TTTTTGTATGTCACTTGATT 3'</li> </ul>	gd HOG1R	5' GAAGTAAGAATGAGTGGTTAGGGACATTAAAAAAACACGT GTTGTAAAACGACGGCCAGT 3'
<ul> <li>co HOG1 R 5' AGATAGAAGTGCTGATAAAC 3'</li> <li>gd BCY11F 5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTGCAT 3'</li> <li>co BCY1R 5' TTTTTGTATGTCACTTGATT 3'</li> </ul>	co HOG1F	5' GTCTGGCGGCGTTATTATAC 3'
<ul> <li>gd BCY11F 5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'</li> <li>gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'</li> <li>co BCY1F 5' ATAAGCTGAACTTATTGCAT 3'</li> <li>co BCY1R 5' TTTTTGTATGTCACTTGATT 3'</li> </ul>	co HOG1 R	5' AGATAGAAGTGCTGATAAAC 3'
gd BCY1R 5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3' co BCY1F 5' ATAAGCTGAACTTATTGCAT 3' co BCY1R 5' TTTTTGTATGTCACTTGATT 3'	gd BCY11F	5' TACAACAAGCAGATTATTTTCAAAAGACAACAGTAAGAATAA ACGCTTCGTACGCTGCAG 3'
co BCY1F 5' ATAAGCTGAACTTATTGCAT 3' co BCY1R 5' TTTTTGTATGTCACTTGATT 3'	gd BCY1R	5' GAGAAAGGAAATTCATGTGGATTTAAGATCGCTTCCCCTTTT TACGCCACTAGTGGATCT 3'
co BCY1R 5' TTTTTGTATGTCACTTGATT 3'	co BCY1F	5' ATAAGCTGAACTTATTGCAT 3'
	co BCY1R	5' TTTTTGTATGTCACTTGATT 3'

Table 7. Primers used for gene disruption and confirmation

gd MSN2F	5' TTTCTTTTTCAACTTTTATTGCTCATAGAAGAACTAGATCT
0	AAACTTCGTACGCTGCAG 3'
gd MSN2R	5' TTATCTTATGAAGAAAGATCTATCGAATTAAAAAAATGGGGT
0	CTAGCCACTAGTGGATCT 3'
co MSN2F	5' CGGGAAGATCACAACAGTAG 3'
co MSN2R	5' ACCCCTCTTGCTTTTGTACG 3'
gd MSN4F	5' ATCAGTTCGGCTTTTTTTTTTTTTTTTTTTTTTTTTTTT
-	ATACTTCGTACGCTGCAG 3'
gd MSN4R	5' TACCGTAGCTTGTCTTGCTTTTATTTGCTTTTGACCTTATTT
-	TTTGCCACTAGTGGATCT 3'
co MSN4F	5' AACCCGAGCTAGAACTAGGA 3'
co MSN4R	5' CATACCGTAGCTTGTCTTGC 3'

Table 8. RT-PCR primers used in the study

Primer name	Sequence
rt GRE1F	5' TGCTCAAAGTAACCGCTACCAA 3'
rt GRE1R	5' TCGTTTCCTGACCCAGACAGA 3'
rt HSP12F	5' CCGAAAAAGGCAAGGATAACG 3'
rt HSP12R	5' CGGCTCCCATGTAATCTCTAGCT 3'
rt HSP26F	5' TGGGTGAAGGCGGCTTAA 3'
rt HSP26R	5' AGAATCCTTTGCGGGTGTGT 3'
rt NCA3F	5' CCCTTTCTTCTGTCGCATTTTC 3'
rt NCA3R	5' TCATCTTTGTGATGATCTTCATGGT 3'
rt PDC6F	5' GATGGGCTGGTAATGCAAATG 3'
rt PDC6R	5' CCTTGATGCGTGCGTAACC 3'
rt SIP18F	5' GGAATGAAGATGGGCCATGA 3'
rt SIP18R	5' TCCAATCGTTCGCAATTCCT 3'
rt SPS100F	5' CGAATTTTACGAGCGCACAA 3'
rt SPS100R	5' TGGACTGGAGGATGATGAGGAT 3'
rt TKL2F	5' CGATTTGACACCTTCGAATCTG 3'
rt TKL2R	5' TGGGTAATGGGAGGTTGGAA 3'
rt ALD3	5' TGGGTGAAGGCGGCTTAA 3'
rt ALD3	5' TCCTTTGCGGGTGTGTTTTG 3'
rt NQM1F	5' AATGCTGTTGGTATGGCAATAGC 3'
rt NQM1R	5' GGGAAAGCCATCCTCGTTATAA 3'

#### **3.3 Results**

# 3.3.1 The *PTP2* and *MSG5*-mediated calcium response involves a novel function for Ssk2, a protein kinase component of HOG pathway

Previously, we presumed that HOG pathway is involved in the mechanism mediating the calcium sensitivity of the *ptp2\Deltamsg5\Delta* double disruptant since additional *ssk2* $\otimes$  disruption acts as a suppressor of the phenotype (Hermansyah et al., 2010). To clarify the nature of HOG pathway's involvement, we checked the phosphorylation of Hog1 in the wild type,  $ptp2\Delta msg5\Delta$  double disruptant and  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant. Western blot analysis using anti-phospho-Hog1p demonstrated that exposure to calcium caused an increase in the phosphorylated Hog1 level in the wild type and  $ptp2\Delta msg5\Delta$  double disruptant with the phosphorylated Hog1 level being higher in the calcium-exposed  $ptp2\Delta msg5\Delta$  double disruptant than in the wild-type strain. Interestingly, there was a decrease in the phosphorylated Hog1 in the *ptp2\Deltamsg5\Deltassk2\Delta* triple disruptant upon exposure to calcium, which indicated that Hog1 phosphorylation is negatively affected by an additional  $ssk2\otimes$ disruption in the *ptp2\Deltamsg5\Delta* background. (Fig.12A). Next, we disrupted the other protein kinase components of the HOG cascade in the  $ptp2\Delta msg5\Delta$ . In the case of the  $ssk2\Delta$ suppressor, we determined if other elements of the HOG pathway are also involved in the same suppression mechanism since SSK2 and PTP2 are components of the HOG pathway as MAPKKK and negative regulator, respectively (Wurgler-Murphy et al., 1997; Tatebayashi et al., 2003). Results showed that additional disruption of SSK22, PBS2 or HOG1 in the  $ptp2\Delta msg5\Delta$  background did not confer calcium tolerance to the double disruptant (Fig. 12B). Our result signifies that the HOG pathway is not directly involved in the suppression mechanism of the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant. Thus, we

hypothesize that *SSK2* has a novel function in the calcium stress pathway that is independent of its putative function in the HOG pathway.



B. Calcium phenotype of HOG gene disruptants

Figure 12.  $ssk2\Delta$  disruption suppresses the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant via a HOG pathway-independent manner. A) Western blot analysis of the wild type,  $ptp2\Delta msg5\Delta$  double disruptant and  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant using anti-Hog1 (yC-15) antibody (Santa Cruz Biotechnology) for total Hog1 and anti-phospho-p38 MAP Kinase (Thr<sup>180</sup>/Tyr<sup>182</sup>) antibody (Cell Signaling Technology) for the phosphorylated Hog1. B) Additional  $ssk2\Delta$  disruption confers calcium tolerance to the  $ptp2\Delta msg5\Delta$  double disruptants were constructed by an additional disruption of a HOG pathway kinase in a  $ptp2\Delta msg5\Delta$  background. Other protein kinase components of the HOG pathway did not confer calcium tolerance to the  $ptp2\Delta msg5\Delta$  double disruptant.

# 3.3.2 19 induced genes in the calcium sensitive $ptp2 \Delta msg5 \Delta$ double disruptant are

# repressed in the calcium tolerant $ptp2 \triangle msg5 \triangle ssk2 \triangle$ triple disruptant

Using microarray analysis, we also attempted to determine genes that might be involved in the suppression mechanism conferred by  $ssk2\Delta$  disruption to the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant by searching for genes with altered expression in the  $ptp2\Delta msg5\Delta$  and  $ptp2\Delta msg5ssk2\Delta$  background compared to the wild type. From the initial chip set used, 5,725 usable gene spots were chosen for variance stabilization across seven experimental grouped samples based on the effects of calcium exposure and gene disruption ( $ptp2\Delta$ ,  $msg5\Delta$  and  $ssk2\Delta$ ) described in Fig 13. From the list of 42 genes that showed significant fluctuation in gene expression (confidence limit = 0.05), 19 genes with distinct "rise and fall" pattern of expression were selected for further analysis (Fig. 13, Table 9). More specifically, "rise and fall" patterned expression refers to induction of genes in the calcium-sensitive  $ptp2\Delta msg5\Delta$  double disruptant strains in comparison to the calcium-tolerant wild type following exposure to high extracellular calcium (Fig. 13, columns 3, 6 and 7) and subsequent repression of the same set of genes in the calcium-tolerant  $ptp2\Delta msg5\Delta ssk2\Delta$ triple disruptant (Fig. 13, column 1). The genes were selected based on the hypothesis that the gene expression pattern directly correlates with the calcium phenotype of the strains in which increased expression of a set of genes is responsible for the calcium sensitive phenotype and that down-regulation of the same genes confers calcium tolerance.

To confirm the validity of the "rise and fall" patterned expression, we selected the top 10 genes with the highest fluctuation (Standard Deviation) values and determined their expression by real time RT-PCR (Table 9). Results showed the characteristic "rise and fall" pattern in all the genes tested thereby confirming the up-regulation and down-regulation of the genes in the *ptp2*\Delta*msg5*\Delta and *ptp2*\Delta*msg5*\Delta*ssk2*\Delta, respectively (Fig. 14). Taken together, we infer that the "rise and fall" pattern in gene expression is related to the calcium phenotype of the strains and their down-regulation may play a role in the suppression of the calcium sensitivity of the *ptp2*\Delta*msg5*\Delta double disruptant.



Figure 13. Heat map showing 19 highly fluctuated genes with significant "rise and fall" patterned expression (3-fold). A) Different experimental set-ups are shown in columns and are listed as follows: 1) calcium-exposed  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant and calcium-exposed  $ptp2\Delta msg5\Delta$  double disruptant; 2) untreated  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant and untreated  $ptp2\Delta msg5\Delta$  double disruptant; 3) calcium-exposed  $ptp2\Delta msg5\Delta$  double disruptant and untreated  $ptp2\Delta msg5\Delta$  double disruptant; 4) calcium-exposed  $msg5\Delta$  single disruptant and calcium-exposed wild type; 5) calcium-exposed  $ptp2\Delta$  single disruptant and calcium-exposed wild type; 6) calcium-exposed  $ptp2\Delta msg5\Delta$  double disruptant and calcium-exposed wild type; 6) calcium-exposed wild type and untreated wild type. The group was determined by k-clustering with the genes arranged based on standard deviation values (SD) with the highest value on the top.

The genes were then analyzed for predicted and known interactions using the STRING web resource (http://string-dg.org) to obtain hints on the overall perspective of the interacting genes or proteins (Szklarczyk et al., 2010). All genes with the exception of *PCK1* and *YAP7* showed linkage in the STRING analysis of the "rise and fall" genes implying the importance of their interaction in the calcium signaling pathway mediated by Ptp2 and Msg5 (Fig. 15). Also, the hierarchical relationships (Gene Ontology) of the "rise and fall" gene cluster were determined using GOstat (http://gostat.wehi.edu.au) in which some genes including *GRE1*, *HSP12*, *ALD3*, *HSP26* and *SIP18* were over-represented in the "response to stress" category (p = 0.000177).

Systematic name	Standard name	Function	SD
YJL116C	NCA3	Regulation of mitochondrial expression of subunits the Fo-F1 ATP synthases	3.55
YHR139C	SPS100	Spore wall maturation	3.50
YPL223C	GRE1	Hydrophilin for dessication-rehydration process	3.07
YBR117C	TKL2	Transketolase	3.01
YGR043C	NQM1	Transalsolase	3.00
YBR072W	HSP26	Small heat shock protein with chaperone activity	2.82
YFL014W	HSP12	Involved in maintaining membrane organization in stress conditions	2.80
YMR175W	SIP18	Essential to desiccation-rehydration process	2.80
YGR087C	PDC6	Involved in amino acid catabolism	2.78
YMR169C	ALD3	Involved in beta-alanine synthesis	2.72
YMR107W	SPG4	Protein required for survival at high temperature during stationary phase	2.64
YOL028C	YAP7	Transcription factor	2.44
YHR096C	HXT5	Hexose transporter	2.43
YKR097W	PCK1	Key enzyme in gluconeogenesis	2.35
YGL156W	AMS1	Vacuolar alpha mannosidase	2.32
YDL223C	HBT1	Polarized cell morphogenesis	2.30
YDR070C	FMP16	Unknown	2.30
YDL204W	RTN2	Stabilizes membrane curvature	2.19
YOL084W	PHM7	Unknown	2.17

Table 9. Highly fluctuated genes exhibiting "rise and fall" pattern in the  $ptp2\Delta msg5\Delta$  and  $ptp2\Delta msg5\Delta ssk2\Delta$  disruptants compared to the wild-type strain (confidence limit = 0.05)



Figure 14. Representative genes showing "rise and fall" patterned expression in the  $ptp2\Delta msg5\Delta$  double disruptant and  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant as compared to the wild type. cDNA synthesized from total RNA isolated from the sample strains grown with (0.3 M CaCl<sub>2</sub>) or without calcium were used as templates for RT-PCR using SYBR Green and specific primers for each gene. *ACT1* was used as an internal control. A "rise and fall" pattern designates the up-regulation of genes in the  $ptp2\Delta msg5\Delta$  double disruptant and subsequent down-regulation in the  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant. The results are means of three experiments from three independent preparations.



Figure 15. STRING analysis of the "rise and fall" genes. Gene set containing the "rise and fall" genes were subjected to protein-protein interaction prediction using STRING 9.0 (string-db.org). Blue lines indicate interaction between two proteins (nodes). Thicker lines indicate stronger interaction.

In correlating the calcium phenotype of the yeast strains with the expression pattern of genes, two trends in gene expression were considered: a) the "rise and fall" patterned expression exemplified by the aforementioned genes and b) the opposing, "fall and rise" expression pattern characterized by the repression and induction of gene expression in the  $ptp2\Delta msg5\Delta$  double disruptant and  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant, respectively. From our microarray analysis, we were not able to find highly fluctuated genes bearing the "fall and rise" trend in gene expression thus; only the "rise and fall" genes were selected for further analysis.

# 3.3.3 "Rise and fall" genes share several transcription factors that may be involved in the suppression of the calcium sensitive phenotype of the $ptp2 \Delta msg5 \Delta$ double disruptant by *SSK2* disruption

We inferred that the genes with "rise and fall" patterned expression belong to a common pathway that is responsible for regulating their expression and thus, the cellular process they control. Therefore, we set to find the transcription factors that may be involved in calcium sensitivity of the *ptp2\Deltamsg5\Delta* double disruptant and its suppression by *ssk2\Delta* disruption. The list of potential transcription factors related to each "rise and fall" gene was collected from the YEASTRACT (Yeast Search for Transcriptional Regulators and Consensus Tracking) website (http://www.yeastract.com) after which the transcription factors shared by the majority of the genes were determined. Overlaps among some but not all of genes were common for each transcription factor listed (Table 10). Among the "rise and fall" genes, Aft1 was shared by 94.7% (18/19), Rpn4 by 84.2% (16/19), Msn2 by 78.9% (15/19), Sok2 by 78.9% (28/42) and Ste12 by 78.9% (15/19) (cut-off was set at 75%). The presence of various overlaps indicated that some transcription factors might be shared among the different clusters thus; we inferred that Aft1, Rpn4, Msn2, Sok2 and Ste12 might play important roles

in the suppression mechanism of the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant by  $ssk2\Delta$  disruption. However, we do not discount the possibility of other transcription factors being involved in the calcium-induced pathway in our strain.

Transcription factor	%	ORF/Genes
Aft1	94.7	HSP26, TKL2, RTN2, HBT1, FMP16, HSP12, AMS1, NQM1, PDC6, HXT5, SPS100, NCA3, PCK1, SPG4, ALD3, SIP18, PHM7, GRE1
Rpn4	84.2	HSP26, TKL2, RTN2, HBT1, FMP16, HSP12, AMS1, NQM1, PDC6, HXT5, SPS100, NCA3, PCK1, SPG4, ALD3, PHM7,
Msn2	78.9	HSP26, TKL2, RTN2, HBT1, FMP16, HSP12, NQM1, HXT5, SPS100, NCA3, SPG4, ALD3, SIP18, PHM7, GRE1
Sok2	78.9	HSP26, HBT1, FMP16, HSP12, AMS1, NQM1, PDC6, HXT5, SPS100, NCA3, PCK1, SPG4, ALD3, SIP18, PHM7
Ste12	78.9	HSP26, TKL2, RTN2, HSP12, AMS1, NQM1, PDC6, HXT5, SPS100, NCA3, PCK1, SPG4, ALD3, SIP18, GRE1

Table 10. Transcription factors shared by the "rise and fall" genes

# 3.3.4 Disruption of $msn2\Delta$ confers calcium tolerance to the $ptp2\Delta msg5\Delta$ double disruptant

To determine which transcription factor is directly involved in the suppression mechanism conferred by *ssk2* $\Delta$  disruption, we disrupted the five candidate transcription factors (*AFT1, RPN4, MSN2, STE12, SOK2*) individually in the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant background. Results showed that only an additional *msn2* $\Delta$  disruption conferred calcium tolerance to the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant (Fig. 16). Interestingly, we found that *ptp2* $\Delta$ *msg5* $\Delta$ *msn4* $\Delta$  was calcium sensitive, indicating that Msn2 and Msn4 do not share a redundant function in relation to calcium signaling although Msn4 may play a lesser role. However, the quadruple disruptant *ptp2* $\Delta$ *msg5* $\Delta$ *msn2* $\Delta$ *msn4* $\Delta$  showed better growth in the presence of calcium compared to *ptp2* $\Delta$ *msg5* $\Delta$ *msn2* $\Delta$ , suggesting that Msn4 has an additive effect with Msn2 in the regulation of the Ptp2 and Msg5-mediated calcium signaling (Fig. 16).



Figure 16. Msn2 mediates the expression of the "rise and fall" genes involved in the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant. Additional  $msn2\Delta$  disruption confers calcium tolerance to the  $ptp2\Delta msg5\Delta$  double disruptant. Triple disruptants were constructed by an additional disruption of the candidate transcription factor genes (*AFT1, RPN4, MSN2, STE12, SOK2*) in the  $ptp2\Delta msg5\Delta$  background. Only  $msn2\Delta$  disruption was able to confer calcium tolerance to the  $ptp2\Delta msg5\Delta$  double disruptant. Yeast strains were grown on YPDA and YPDA + 0.6 M CaCl<sub>2</sub> media at 30°C for 2 - 4 days.

To confirm the direct relationship between Msn2 and the suppression mechanism exhibited in the  $ptp2\Delta msg5\Delta ssk2\Delta$  double disruptant, we examined the expression of genes

with patterned expression in the *ptp2\Delta msg5 \Delta msn2\Delta* triple disruptant. We found that the same

rise and fall pattern of genes was exhibited in the wild type,  $ptp2\Delta msg5\Delta$  and

 $ptp2\Delta msg5\Delta msn2\Delta$  disruptant, indicating a positive correlation between Msn2 and the

suppression mechanism conferred by  $ssk2\Delta$  disruption (Fig. 17).



Figure 17. Representative genes showing "rise and fall" patterned expression in the  $ptp2\Delta msg5\Delta$  double disruptant and  $ptp2\Delta msg5\Delta msn2\Delta$  triple disruptant as compared to the wild type. cDNA synthesized from total RNA isolated from the sample strains grown with (0.3 M CaCl<sub>2</sub>) or without calcium were used as templates for RT-PCR using SYBR Green and specific primers for each gene. *ACT1* was used as an internal control. A "rise and fall" pattern designates the up-regulation of genes in the  $ptp2\Delta msg5\Delta$  double disruptant and subsequent down-regulation in the  $ptp2\Delta msg5\Delta msn2\Delta$  triple disruptant. The results are means of three experiments from three independent preparations.

## 3.3.5 Constitutive activation of PKA suppresses the calcium sensitive phenotype of the

## $ptp2 \Delta msg5 \Delta$ double disruptant

We hypothesized that finding their common upstream regulator can consolidate the heterogeneity of the "rise and fall" genes involved in the Ptp2 and Msg5-mediated calcium signaling. Previously, we have identified two suppressors of the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant namely PKase disruptions  $mck1\Delta$  and  $yak1\Delta$ , which have functional relationships with the PKA pathway, a major regulator of cell growth (Hermansyah et al., 2010, Santangelo, 2006). Furthermore, published reports describe genetic interactions between PKA components and the candidate transcription factors Aft1, Rpn4, Msn2, Sok2 and Ste12 described above (Ward et al., 1995; Robertson et al., 2000; , Wang et al., 2004; Smith et al., 1998; Pan and Heitman, 1999). Since PKA promotes cell growth, we disrupted the PKA regulatory subunit *BCY1* in the  $ptp2\Delta msg5\Delta$  background to determine if

PKA activation can suppress the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant. Interestingly, genetic analysis showed that the  $ptp2\Delta msg5\Delta bcy1\Delta$  triple disruptant exhibited calcium tolerance thus indicating that Bcy1 contributes to the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant, more likely by inhibiting PKA (Fig. 18).



Figure 18. Constitutive PKA activation suppresses the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$  double disruptant via the regulation of the "rise and fall" genes. Additional  $bcy1\Delta$  disruption in the  $ptp2\Delta msg5\Delta$  double disruptant background confers calcium tolerance.  $ptp2\Delta msg5\Delta bcy1\Delta$  disruptant was constructed by disrupting *BCY1* in the  $ptp2\Delta msg5\Delta$  double disruptant background by direct gene disruption. Yeast strains were grown on YPDA and YPDA + 0.6 M CaCl<sub>2</sub> media at 30°C for 2 - 4 days where  $ptp2\Delta msg5\Delta bcy1\Delta$  disruptant displayed calcium tolerance.

To determine whether the suppression mechanism of  $bcy1\Delta$  disruption is directly related to Ssk2 and the "rise and fall" genes, we checked the patterned expression in the calcium tolerant  $ptp2\Delta msg5\Delta bcy1\Delta$ . Results showed that the expression of the representative genes were also down-regulated in the  $ptp2\Delta msg5\Delta bcy1\Delta$  disruptant thereby showing the same "rise and fall" pattern as in the  $ptp2\Delta msg5\Delta ssk2\Delta$  disruptant (Fig. 19). Based on this, we conclude that the suppression mechanism of the  $bcy1\Delta$  disruption is related to Ssk2 through its regulation of the "rise and fall" genes.



Figure 19. Representative genes showing "rise and fall" patterned expression in the  $ptp2\Delta msg5\Delta$  double disruptant and  $ptp2\Delta msg5\Delta bcy1\Delta$  triple disruptant as compared to the wild type. cDNA synthesized from total RNA isolated from the sample strains grown with (0.3 M CaCl<sub>2</sub>) or without calcium were used as templates for RT-PCR using SYBR Green and specific primers for each gene. *ACT1* was used as an internal control. A "rise and fall" pattern designates the up-regulation of genes in the  $ptp2\Delta msg5\Delta$  double disruptant and subsequent down-regulation in the  $ptp2\Delta msg5\Delta bcy1\Delta$  triple disruptant. The results are means of three experiments from three independent preparations.

#### **3.4 Discussion**

The non-suppressor nature of the disruptions of other HOG pathway protein kinases implies that the reduced Hog1 phosphorylation in the  $ptp2\Delta msg5\Delta ssk2\Delta$  triple disruptant is not a major factor in the suppression of the calcium sensitive phenotype of the  $ptp2\Delta msg5\Delta$ disruptant rather it is a result of an attenuated HOG signaling response due to the disruption of the HOG MAPKKK component *SSK2*. Furthermore, the observed increase in Hog1 phosphorylation in the  $ptp2\Delta msg5\Delta$  double disruptant was probably due to the absence of the HOG regulator Ptp2 and was not the major cause of the calcium sensitive phenotype.

Our analysis revealed that Aft1, a transcription factor involved in iron utilization and homeostasis as well as oxidative response, may have a role in the suppression mechanism by ssk2A disruption (Yamaguchi-Iwai et al., 1995; Shakoury-Elizeh et al., 2004; Castells-Roca et al., 2011). It is interesting to note the involvement of iron homeostasis in the calcium-induced signaling pathway of the *ptp2\Delta msg5\Delta* double disruptant since null mutation of *AFT1* is known to cause delayed cell cycle progression in G1 phase similar to the calcium sensitive phenotype of the *ptp2\Deltamsg5\Delta* double disruptant (White et al., 2009; Hermansyah et al., 2009). Rpn4 was previously reported to be involved in the expression of proteosomal genes and calcineurin-mediated degradation of Yap1 that occurs to ensure a G2 delay in response to calcium stress. In particular, Yap1 negatively regulates Swe1 and Cln2 indirectly via the down-regulation of RPN4 expression (Yokohama et al., 2006). Assuming that the upregulation of the "rise and fall" genes is due to the proteasome activity modulated by Yap1 and Rpn4, Ptp2 and Msg5 may both act as negative regulators of Yap1 while Ssk2 promotes Yap1 and Rpn4 activity. On the other hand, Sok2 is a nuclear protein in the cyclic AMP (cAMP)-dependent protein kinase A (PKA) signaling pathway that negatively regulates pseudohyphal differentiation while Ste12 is a transcription factor related to mating pheromone pathway (Ward et al., 1995; Pan and Heitman, 2000; Santangelo, 2006). We

recently reported the synergistic roles of both Sok2 and Ste12 in the up-regulation of the cell wall genes of the calcium-sensitive  $ptp2\Delta msg5\Delta$  double disruptant thus; their involvement in the suppression mechanism conferred by  $ssk2\Delta$  disruption is not surprising (Laviña et al., 2013). YEASTRACT analysis revealed that 78.9% (15/19) of the "rise and fall" genes contained one or more STRE binding motif (CCCCT or AGGGG), indicating the commonality of the general stress response transcription factors Msn2 and Msn4. Presumably, Msn2 and Msn4 play a role in mediating the stress response genes (*HSP26, HSP12, ALD3, SIP18* and *GRE1*) that exhibit the "rise and fall" patterned expression in the  $ptp2\Delta msg5\Delta$ double disruptant. In this case, Msn2 and Msn4 may promote induction of the stress response genes that eventually leads to a calcium sensitive phenotype.

It has been widely accepted that Msn2 is a stronger transcription factor than Msn4 although both share functional similarities (Hohmann, 2002). Our data support this idea wherein disruption of *MSN2* alone can confer calcium tolerance while *msn4* $\Delta$  cannot. However, Msn2 and Msn4 were found to exhibit an additive effect as exemplified by the better growth of the *ptp2* $\Delta$ *msg5* $\Delta$ *msn2* $\Delta$ *msn4* $\Delta$  quadruple disruptant in YPDA supplemented with calcium (Fig. 16).

One possibility is that the calcium sensitivity was a result of growth inhibition initiated by Yak1 since *yak1* $\Delta$  disruption had the most profound change on the phenotype from calcium sensitive to tolerant (Hermansyah et al., 2010). In this case, we assumed that constitutive PKA activation (due to *bcy1* $\Delta$  disruption) was able to suppress the cell cycle arrest by overcoming Yak1 activity. Thus, we hypothesized that PKA served as an upstream regulator of the calcium-induced signaling pathway exhibited in the *ptp2* $\Delta$ *msg5* $\Delta$  double disruptant. We have also observed a similar "rise and fall" pattern in the *ptp2* $\Delta$ *msg5* $\Delta$ *yak1* $\Delta$ triple disruptant, prompting us to conclude that the suppression mechanism conferred by *ssk2* $\Delta$  has a direct relationship with PKA componentsYak1 and Bcy1 (Fig. 20). In addition,

PKA is known to phosphorylate at least five sites on Msn2 to prevent its activity (Gorner et al., 2002). This may indicate that the suppression mechanism of the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant through constitutive PKA activation ( $bcy1\Delta$  disruption) may also occur via the inhibition of Msn2 (and Msn4) activity.



Figure 20. Representative genes showing "rise and fall" patterned expression in the  $ptp2\Delta msg5\Delta$  double disruptant and  $ptp2\Delta msg5\Delta yak1\Delta$  triple disruptant as compared to the wild type. cDNA synthesized from total RNA isolated from the sample strains grown with (0.3 M CaCl<sub>2</sub>) or without calcium were used as templates for RT-PCR using SYBR Green and specific primers for each gene. *ACT1* was used as an internal control. A "rise and fall" pattern designates the up-regulation of genes in the  $ptp2\Delta msg5\Delta$  double disruptant and subsequent down-regulation in the  $ptp2\Delta msg5\Delta yak1\Delta$  triple disruptant. The results are means of three experiments from three independent preparations.

In conclusion, the calcium sensitivity and the subsequent tolerance conferred by the protein kinase suppressors show the complexity of signaling cascades present in eukaryotic cells. Specifically, the yeast cells possess a number of redundant proteins and pathways to combat inadvertent activation or repression of pathway signals. As we have shown previously, such is the case for our  $ptp2\Delta msg5\Delta$  double disruptant strain in response to high intracellular calcium where we found the redundant function of the SLT2 and calcineurin pathways (Laviña et al., 2013). Furthermore, the absence of the functionally redundant *PTP2* 

and *MSG5* has proven to be detrimental to cells exposed to high calcium concentrations. Based on our results, *PTP2* and *MSG5* are integral in maintaining signal specificity thus their absence can lead to cross-talk events involving several pathways. In this study, we found another alternative pathway that mediates the calcium phenotype of the *ptp2*\Delta*msg5*\Delta double disruptant involving a novel function of Ssk2 that is related to PKA-related proteins Yak1 and Bcy1, transcription factor Msn2 as well as a group of calcium-induced, stress response-related genes.

#### Chapter 4

#### **Conclusion and general discussion**

*S. cerevisiae* possesses an inherent ability to cope with the deleterious changes in the environment by activating a variety of signaling cascades that control patterns of gene expression and protein activity. Since inappropriate activation of cell signaling pathways often has dire consequences on the viability, cells possess a myriad of ways to combat improper activation of signaling pathways, one of which is regulation via reversible phosphorylation mediated by protein kinases and phosphatases. In an effort to study the functional redundancy of protein phosphatases in yeast, Sakumoto and colleagues (2002) constructed a PPase library of single and double disruptants and systematically screened for interesting phenotypes; one of which is that the disruption of two protein phosphatase genes, *PTP2* and *MSG5*, causes calcium sensitivity indicating that functional redundancy exists between the two PPases in response to high extracellular calcium. Furthermore, we found that additional disruption of protein kinases *BCK1*, *MKK1*, *SLT2*, *MCK1*, *YAK*1 and *SSK2* in the *ptp2*\Delta*msg5*\Delta background confers calcium tolerance.

This work describes the suppression mechanism conferred by the protein kinase disruptions. First, we discuss the suppression mechanism of the SLT2 pathway suppressors  $bck1\Delta$ ,  $mkk1\Delta$  and  $slt2\Delta$  as having functional redundancy with the calcineurin pathway. In the event of a calcium-induced, calcineurin-driven signaling pathway activation, the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant can be suppressed by preventing over-activation of the calcium-induced signaling cascade under the parallel control of the SLT2 and calcineurin pathways via inactivating either one of the two (Fig. 10).

In our study we have discovered functional redundancy existing at two levels: at the protein level between Ptp2 and Msg5 and at the cascade level between SLT2 and calcineurin pathways. At the protein level, Ptp2 and Msg5 have redundant functions as negative
regulators of the SLT2 pathway. This indicates the importance of preventing the hyperactivation of the calcium-induced signaling pathway mediated by the parallel SLT2 and calcineurin pathways. In our model, calcineurin is the primary signaling cascade activated upon calcium exposure while SLT2 acts as an alternate, secondary pathway if calcineurin is impaired. Thus, in the normal calcium-exposed setting, SLT2 pathway should be repressed via negative regulation by Ptp2 or Msg5. At the cascade level, we show the functional redundancy between SLT2 and calcineurin pathways in response to high extracellular calcium (Fig. 10). Conversely, functional redundancy also serves as safeguard against the lethal effects of an inactive signaling pathway. It is believed that genetic redundancy contributes to the genetic robustness of yeast cells against external changes as shown in fitness data with significantly higher probability of functional compensation for a duplicate gene (Gu et al., 2003) However, this theory contradicts population genetics prediction that genetic redundancy is evolutionary unstable (Li et al., 2010). Our data more likely supports the theory of genetic buffering which results in the concealment of phenotypic consequences of mutations. In fact, redundancy is the best characterized form of buffering relationship, where genes can compensate for the loss of another by their ability to share and takeover the same function (van Wageningen et al., 2010).

In our model, we hypothesized that the calcium-induced signaling pathway mediated by the parallel SLT2 and calcineurin pathways bottlenecks at Crz1 since  $crz1\Delta$  in the  $ptp2\Delta msg5\Delta$  background resulted in a calcium sensitive phenotype (Fig. 10). It has been reported that Crz1 is the major, if not the only, effector of the calcineurin-regulated gene expression in yeast (Yoshimoto et al., 2002). However, we did not find correlation between normally activated Crz1 targets and the Ptp2 and Msg5-mediated calcium signaling thus, we deduced that in this particular signaling cascade, Crz1 targets a different group of downstream genes. Currently, the identity of the Crz1 targets is still undetermined.

72

We have also predicted another alternative pathway that can suppress the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant that involves the PKase suppressors  $ssk2\Delta$ and  $yak1\Delta$  based on FACS analysis. Genome-wide transcriptional analysis of the  $ptp2\Delta msg5\Delta$  and  $ptp2\Delta msg5\Delta ssk2\Delta$  disruptants revealed 19 genes with a "rise and fall" patterned expression that is related to the calcium phenotype. We discovered that these genes were mediated by the transcription factor Msn2 (and partly by Msn4) which served as the convergence point between Ssk2 and PKA since this particular group of suppressors ( $ssk2\Delta$ ,  $yak1\Delta$ ,  $bcy1\Delta$  and  $msn2\Delta$ ) all exhibit the "rise and fall" patterned expression of effector genes.

In detail, upon exposure to high extracellular calcium, Ssk2 is responsible for the induction of 19 "rise and fall" genes that contribute to the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant via Yak1 and the transcription factor Msn2. Suppression of the induction of the "rise and fall" genes can occur by inactivating any component of the linear cascade namely Ssk2, Yak1 or Msn2. In addition, inhibiting Msn2 activity by constitutive PKA activation ( $bcy1\Delta$  disruption) also results in the repression of the "rise and fall" genes. However, the positions of Ssk2 and Yak1 have been arbitrarily placed in the linear pathway and the exact position of the proteins in the Ptp2 and Msg5-mediated calcium signaling is the subject of future experiments (Fig. 21).

Using genetic and transcriptional analyses, the roles of the previously isolated PKase suppressors  $bck1\Delta$ ,  $mkk1\Delta$ ,  $slt2\Delta$ ,  $ssk2\Delta$  and  $yak1\Delta$  in the calcium-induced, Ptp2 and Msg5mediated signaling in *S. cerevisiae* were clarified. In addition to the previously identified PKase suppressors, additional suppressors namely  $cnb1\Delta$ ,  $bcy1\Delta$  and  $msn2\Delta$  were identified in this study. It also provides evidence to the existence of several alternative pathways that mediate calcium signaling in the  $ptp2\Delta msg5\Delta$  double disruptant.

73



Figure 21. Suppression of the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant results from the repression of the Msn2-mediated "rise and fall" genes. High extracellular calcium triggers the induction of 19 "rise and fall" genes responsible for the calcium sensitivity of the  $ptp2\Delta msg5\Delta$  double disruptant via Ssk2, Yak1 and the transcription factor Msn2. Repression of the "rise and fall" genes can be achieved by disruption of any component of the linear pathway ( $ssk2\Delta$ ,  $yak1\Delta$ ,  $msn2\Delta$ ) or constitutive PKA activation ( $bcy1\Delta$ ).

In our model, Msn2 was placed downstream of Ssk2 based on their roles in the HOG pathway. However, a direct genetic relationship between Ssk2 and Msn2 was not established in our study. Furthermore, there are no published reports of a direct interaction between Ssk2 and Msn2 (BioGRID), indicating the possibility of an unknown Ssk2 substrate that mediates Msn2 regulation.

This study shows evidence of multi-level, functionally redundant mechanisms in *S. cerevisiae* responsible for buffering the lethal effects of genetic and environmental changes. Here, we have outlined a case where the detrimental effects of gene deletion were circumvented by the activation or deactivation of alternative pathways involving SLT2, calcineurin, PKA and a novel function of Ssk2.

## References

- An, X., Zhang, Z., Yang, K. and Huang, M. 2006. Cotransport of the heterodimer small subunit of the *Saccharomyces cerevisiae* ribonucleotide reductase between the nucleus and the cytoplasm, Genetics, 173, 63-73.
- Amberg, D. C., Burke, D. J. and Strathern, J. N. 2005. Methods in Yeast Genetics, 2005 ed., Cold Spring Harbor Press, New York, p. 199-202.
- Berry, D. B. and Gasch, A. P. 2008. Stress-activated genomic expression changes serve a preparative role for impending stress in yeast, Mol. Biol. Cell, 19, 4580-4587.
- Blackwell E., Kim, H. J. and Stone, D. E. 2007. The pheromone-induced nuclear accumulation of the Fus3 MAPK in yeast depends on its phosphorylation state and on Dig1 and Dig2. BMC Cell Biol. 8 (44), 1-15.
- Castells-Roca, L., Muhlenhoff, U., Lill, R., Herrero, E. and Belli, G. 2011. The oxidative stress response in yeast cells involves changes in the stability of Aft1 regulon mRNAs, Mol. Microbiol., 81 (1), 232-248.
- Clapham, D. E. 2007. Calcium signaling. Cell 131 (6), 1047-1058.
- Collister, M., Didmon, M. P., MacIsaac, F., Stark, M. J., MacDonald, N.Q. and Keyse, S.M.
   2002. YIL113W encodes a functional dual-specificity protein phosphatase which specifically interacts with and inactivates the Slt2/Mpk1p MAP kinase in *S. cerevisiae*, FEBS Letters, 527, 186-192.
- Cunningham, K. W. and Fink, G. R. 1994. Ca<sup>2+</sup> transport in *Saccharomyces cerevisiae*, J. Exp. Biol., 196, 157-166.
- Cyert, M. S. 2003. Calcineurin signaling in *Saccharomyces cerevisiae*: how yeast go crazy in response to stress. Biochem. Biophys Res. Comm., 311, 1143-1150.

- Cyert, M. S. 2002. Genome-wide analysis of gene expression regulated by the calcineurin/Crz1p signaling pathway in *Saccharomyces cerevisiae*, J. Biol. Chem., 277, 31079 - 31088.
- Cyert, M. S., Kunisawa, R., Kaim, D., and Thorner, J. 1991. Yeast has homologs (*CNA1* and *CNA2* gene products) of mammalian calcineurin, a calmodulin-regulated phosphoprotein phosphatase, PNAS, 88, 7376-7380.
- Davis, T. N. 1995. Calcium in *Saccharomyces cerevisiae*, Adv. Second Messenger Phosphoprotein Res., 30, 339-358.
- Doi, K., Gartner, A., Ammerer, G., Errede, B., Shinkawa, H., Sugimoto, K. and Matsumoto,
  K. 1994. *MSG5*, a novel protein phosphatase promotes adaptation to pheromone response in *S. cerevisiae*, EMBO J., 13, 61-70.
- Flandez, M., Cosano, I. C., Nombela, C., Martin, H. and Molina, M. 2004. Reciprocal regulation between Slt2 MAPK and isoforms of Msg5 dual-specificity protein phosphatase modulates the yeast cell integrity pathway, J. Biol. Chem., 279, 11027-11034.
- Garrett, S., Menold, M. M. and Broach, J. R. 1991. The *Saccharomyces cerevisiae YAK1* gene encodes a protein kinase that is induced by arrest early in the cell cycle, Mol. Cell. Biol., 11 (8), 4045-4052.
- Garrett-Engele, P., Moilanen, B. and Cyert, M. S. 1995. Calcineurin, the Ca<sup>2+/</sup>calmodulindependent protein phosphatase, is essential in yeast mutants with cell integrity defects and in mutants that lack a functional vacuolar H(+)-ATPase, Mol. Cell. Biol., 15, 4103-4114.
- Gasch, A. P. 2002. The environmental stress response: a common yeast response to diverse environmental stresses. In Yeast stress responses. S. Hohmann and P. Mager (ed.)

Volume 1, Topics in current genetics (series editor S. Hohmann) Springer-Verlag Heidelberg p. 11-70.

- Gerami-Nejad, M., Berman, J. and Gale, C. A. 2001. Cassettes for PCR-mediated construction of green, yellow and cyan fluorescent protein fusions in *Candida albicans*, Yeast, 18 (9), 859-864.
- Gietz, R. D. and Schiestl, R. H. 2007. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method, Nat. Protocols, 2, 31-34.
- Goffeau, A., Barrell, B. G., Bussey, H., Davis, R. W., Dujon, B., Feldmann, H., Galibert, F., Hoseisel, J. D., Jacq, C., Johnston, M., Louis, E. J., Mewes, H. W., Murakami, Y., Phillippsen, P., Tettelin, H., and Oliver, S. G. 1996. Life with 6000 genes, Science, 274, 546-567.
- Gorner, W., Durchschlag, E., Wolf, J., Brown, E. L., Ammerer, G., Ruis, H., and Schuller, C.
  2002. Acute glucose starvation activates the nuclear localization signal of a stress-specific yeast transcription factor, EMBO J., 21, 135-144.
- Gu, Z., Steinmetz, L. M., Gu, X., Scharfe, C., Davis, R. W. and Li, W-H. 2003. Role of duplicate genes in genetic robustness against null mutations, Nature, 421, 63-66.
- Guan, K.L., Deschenes, R.J. and Dixon, J.E. 1992. Isolation and characterization of a second protein tyrosine phosphatase gene, *PTP2*, from *Saccharomyces cerevisiae*, J. Biol. Chem., 267, 10024-10030.
- Gustin, M. C., Albertyn, J., Alexander, M. and Davenport, K. 1998. MAP kinase pathways in the yeast *Saccharomyces cerevisiae*, Microbiol. Mol. Biol. Rev., 62, 1264-1300.
- Hahn, J. S. and Thiele, D. J. 2002. Regulation of the *Saccharomyces cerevisiae* Slt2 kinase pathway by the stress-inducible Sdp1 dual specificity phosphatase, J. Biol. Chem., 277, 21278-21284.

Hermansyah, Laviña, W. A., Sugiyama, M., Kaneko, Y. and Harashima, S. 2010.
Identification of protein kinase disruptions as suppressors of the calcium sensitivity of *S. cerevisiae Δptp2Δmsg5* protein phosphatase double disruptant, Arch. Microbiol., 192, 157-165.

- Hermansyah, Sugiyama, M., Kaneko, Y. and Harashima, S. 2009. Yeast protein phosphatase Ptp2p and Msg5p are involved in G1-S transition, *CLN2* transcription, and vacuole morphogenesis, Arch. Microbiol., 191, 721-733.
- Hohmann, S. 2002. Osmotic stress signaling and osmoadaptation in yeasts, Microbiol. Mol. Biol. Rev., 66 (2), 300-372.
- Hohmann, S. and Mager, W. H. 2002. Introduction, p. 1-8. In: Hohmann, S. and P. Mager, P. (ed.), Yeast Stress Responses: Topics in Current Genetics. Vol. 1. Springer-Verlag, Heidelberg.
- Jung, U. S. and Levin, D. E. 1999. Genome-wide analysis of gene expression regulated by the yeast cell wall integrity signalling pathway, Mol. Microbiol., 34, 1049–1057.
- Kassis, S., Melhuish, T., Annan, R. S., Chen, S. L., Lee, J. C., Livi, G. P., and Creasy, C. L.
  2000. *Saccharomyces cerevisiae* Yak1p protein kinase autophosphorylates on tyrosine residues and phosphorylates myelin basic protein on a C-terminal serine residue.
  Biochem J., 348, 263-272.
- Kingsbury, T. J. and Cunningham, K. W. 2000. A conserved family of calcineurin regulators. Genes Dev., 14, 1595-1604.
- Klis, F. M., Boorsma, A. and De Groot, P. W. J. 2006. Cell construction in *Saccharomyces cerevisiae*, Yeast, 23, 185-202.
- Kuno, T., Tanaka, H., Mukai, H., Chang, C. D., Hiraga, K., Miyakawa, T. and Tanaka, C.
  1991. cDNA cloning of a calcineurin B homolog in *Saccharomyces cerevisiae*,
  Biochem. Biophys. Res. Commun, 180, 1159-1163.

- Laviña, W. A., Hermansyah, Sugiyama, M., Kaneko, Y. and Harashima, S. 2013. Functional redundancy of protein phosphatase genes *PTP2* and *MSG5* in the presence of high extracellular calcium is essential for stress response and growth in *Saccharomyces cerevisiae*, J. Biosci. Bioeng., 115 (2), 138-146.
- Leman, S. C., Uyenoyama, M. K., Lavine, M. and Chen, Y. 2007. The evolutionary forest algorithm, Bioinformatics, 23 (15), 1962-1968.
- Levin, D. E. 2005. Cell wall integrity signaling in *Saccharomyces cerevisiae*, Microbiol. Mol. Biol. Rev., 69, 262-291.
- Li, J., Yuan, Z. and Zhang, Z. 2010. The cellular robustness by genetic redundancy in budding yeast, PloS Genet., e1001187.doi:10.1371/journal.pgen.1001187.
- Maeda, T., Takekawa, M. and Saito, H. 1995. Activation of yeast PBS2 MAPKK by MAPKKKs or by binding of an SH3-containing osmosensor, Science, 269, 554-558.
- Mattison, C.P., Spencer, S.S., Kresge, K.A., Lee, J. and Ota, I.M. 1999. Differential regulation of the cell wall integrity mitogen-activated protein kinase pathway in budding yeast by the protein tyrosine phosphatases Ptp2 and Ptp3, Mol. Cell Biol., 19, 7651-7660.
- Martin, H., Flandez, M., Nombela, C. and Molina, M. 2005. Protein phosphatases in MAPK signalling: we keep learning from yeast, Mol. Microbiol., 58, 6-16.
- Martin, H., Rodriguez-Pachon, J.M., Ruiz, C., Nombela, C. and Molina, M. 2000. Regulatory mechanisms for modulation of signaling through the cell integrity Slt2-mediated pathway in *Saccharomyces cerevisiae*, J. Biol. Chem., 275, 1511-1519.
- Martinez-Pastor, M., Marchler, G., Schuller, C., Marchler-Bauer, A., Ruis, H. and Estruch, F. 1996. The *Saccharomyces cerevisiae* zinc finger proteins Msn2p and Msn4p are required for transcriptional induction through the stress response element (STRE), EMBO J., 15, 2227-2235.

- Mazur, P., Morin, N., Baginsky, W., el-Sherbeini, M., Clemas, J. A., Nielsen, J. B. and Foor,
  F. 1995. Differential expression and function of two homologous subunits of yeast
  1,3-β-D-glucan synthase, Mol. Cell Biol., 15, 5671-5681.
- Miyakawa, T. and Mizunuma, M. 2007. Physiological roles of calcineurin in *Saccharomyces cerevisiae* with special emphasis on its roles in G2/M cell-cycle regulation, Biosci.
   Biotecnol. Biochem., 71, 633-645.
- Mizunuma, M., Hirata, D., Miyahara, K., Tsuchiya, E. and Miyakawa, T. 1998. Role of calcineurin and Mpk1 in regulating the onset of mitosis in budding yeast, Nature, 392, 303-306.
- Muller, E. M., Locke, E. G. and Cunningham, K. W. 2001. Differential regulation of two Ca<sup>2+</sup> influx systems by pheromone signaling in *Saccharomyces cerevisiae*, Genetics, 159, 1527-1538.
- Nakamura, T., Ohmoto, T., Hirata, D., Tsuchiya. E. and Miyakawa, T. 1996. Genetic evidence for the functional redundancy of the calcineurin and Mpkl-mediated pathways in the regulation of cellular events important for growth in *Saccharomyces cerevisiae*. Mol. Gen. Genet., 251, 211-219.
- O'Rourke, S. M. and Herskowitz, I. 2002. A third osmosensing branch in *Saccharomyces cerevisiae* requires the Msb2 protein and functions in parallel with the Sho1 branch. Mol. Cell. Biol., 22 (13), 4739-4749.
- Pan, X. and Heitman, J. 2000. Sok2 regulates yeast pseudohyphal differentiation via a transcription factor cascade that regulates cell-cell adhesion, Mol. Cell Biol., 20 (22), 8364-8372.
- Pan, X. and Heitman, J. 1999. Cyclic AMP-dependent protein kinase regulates pseudohyphal differentiation in *Saccharomyces cerevisiae*, Mol. Cell Biol., 19 (7), 4874-4887.

- Piskur, J., and Langkjaer, R. B. 2004. Yeast genome sequencing: the power of comparative genomics, Mol. Microbiol., 53 (2), 381-389.
- Posas, F. and Saito, H. 1998. Activation of the yeast *SSK2* MAP kinase kinase kinase by the *SSK1* two-component response regulator, EMBO J., 17 (5), 1385-1394.
- Pratt, . L., Drehman, B. J., Miller, M. E. and Johnston, S. D. 2007. Mutual interdependence of MSII (CAC3) and YAK1 in Saccharomyces cerevisiae, J. Mol. Biol., 368, 30-43.
- Rahmouni S., Cerignoli, F., Alonso A., Tsutji, T., Henkens, R., Zhu, C., Louis-dit-Sully, C.,
  Moutschen, M., Jiang, W. and Mustelin, T. 2006. Loss of the VHR dual-specific
  phosphatase cause cell-cycle arrest and senescence, Nat. Cell Biol., 8 (5), 524-531.
- Reiser, V., Salah, S. M. and Ammerer, G. 2000. Polarized localization of yeast Pbs2 depends on osmostress, the membrane protein Sho1 and Cdc42, Nat. Cell Biol, 2, 620-627.
- Robertson, L. S., Causton, H. C., Young, R. A. and Fink, G. R. 2000. The yeast A kinases differentially regulate iron uptake and respiratory function, PNAS, 97 (11), 5984-5988.
- Rudolph, H. K., Antebi, A., Fink, G. R., Buckley, C. M., Dorman, T. E, Le Vitre, J.,
  Davidow, L. S., Mao, J. I. and Moir, D. T. 1989. The yeast secretory pathway is
  perturbed by mutations in *Pmr1*, a member of the Ca<sup>2+</sup> ATPase family, Cell, 58, 133-145.
- Saito, H. and Tatebayashi, K. 2004. Regulation of the osmoregulatory HOG MAPK cascade in yeast, J. Biochem., 136, 267-272.
- Sakumoto, N., Matsuoka, I., Mukai, Y., Ogawa, N., Kaneko, Y. and Harashima, S. 2002. A series of double disruptants for protein phosphatase genes in *Saccharomyces cerevisiae* and their phenotypic analysis, Yeast, 19, 587-599.
- Sakumoto, N., Mukai, Y., Uchida, K., Kouchi, T., Kuwajima, J., Nakagawa, Y., Sugioka, S., Yamamoto, E., Furuyama, T., Mizubuchi, H., Ohsugi, N., Sakuno, T., Kikuchi, K.,

Matsuoka, I., Ogawa, N., Kaneko, Y. and Harashima, S. 1999. A series of protein phosphatase gene disruptants in *Saccharomyces cerevisiae*. Yeast, 15, 1669-1679.

- Sambrook, J., Fritsch, E. F. and Maniatis, T. 1989. Molecular cloning: A laboratory manual, 2<sup>nd</sup> ed., Cold Spring Harbor, New York.
- Santangelo, G. M. 2006. Glucose signaling in *Saccharomyces cerevisiae*, Microbiol. Mol. Biol. Rev., 70 (1), 253-282.
- Shakoury-Elizeh, M., Tiederman, J., Rashford, J., Ferea, T., Demeter, J., Garcia, E., Rolfes,
  R., Brown, P. O., Botstein, D., and Philpott, C. C. 2004. Transcriptional remodeling in response to iron deprivation in *Saccharomyces cerevisiae*, Mol. Biol. Cell, 15 (3), 1233-1243.
- Sherman, F., and Hicks, J. 1991. Micromanipulation and dissection of asci, Methods Enzymol., 194, 21-37.
- Smith, A., Ward, M. P. and Garrett, S. 1998. Yeast PKA represses Msn2p/Msn4p-dependent gene expression to regulate growth, stress response and glycogen accumulation, EMBO J., 17 (13), 3556-3564.
- Spellman, P. T., Sherlock, G., Zhang, M. Q., Iyer, V. R., Anders, K., Eisen, M. B., Brown, P. O., Botstein, D. and Futcher, B. 1998. Comprehensive identification of cell-cycle regulated genes of the yeast *Saccharomyces cerevisiae* by microarray hybridization, Mol. Biol. Cell, 9, 3273-3297.
- Stark, M. J. R. 1996. Yeast protein serine/threonine phosphatases: multiple roles and diverse regulation, Yeast, 12, 1647-1675.
- Stathopoulos-Genontides, A., Guo, J. J. and Cyert, M. S. 1999. Yeast calcineurin regulates nuclear localization of the Crz1p transcription factor through dephosphorylation, Genes Dev., 13, 798-803.

- Stathooulos, A. M. and Cyert, M.S. 1997. Calcineurin acts through the *CRZ1/TCN1*-encoded transcription factor to regulate gene expression in yeast. Genes Dev., 11, 3432-3444.
- Sugiura, R., Sio, S. O., Shuntoh, H. and Kuno, T. 2002. Calcineurin phosphatase in signal transduction: lessons from fission yeast, Genes to cells, 7, 619-627.
- Szklarczyk, D., Franceschini, A., Kuhn, M., Simonivic, M., Roth, A., Minguez, P., Doerks,
  T., Stark, M., Muller, J., Bork, P., Jensen, L. J. and von Mering, C. 2010. The
  STRING database in 2011: functional interaction networks of proteins, globally
  integrated and scored, Nucleic Acids Res., 39, 561-568.
- Tamaki, H. 2007. Glucose-stimulated cAMP-protein kinase A pathway in yeast *Saccharomyces cerevisiae*, J. Biosc. Bioeng., 104 (4), 245 250.
- Tatebayashi, K., Takekawa, M. and Saito, H. 2003. A docking site determining specificity of Pbs2 MAPKK for Ssk2/Ssk22 MAPKKKs in the yeast HOG pathway, EMBO J., 22, 3624-3634.
- Teixeira, M. C., Monteiro, P., Jain, P., Tenreiro, S., Fernandes, A. R., Mira, N. P., Alenquer, M., Freitas, A. T., Oliveira, A. L., and Sa-Correia, I. 2005. The YEASTRACT database: a tool for the analysis of transcription regulatory associations in *Saccharomyces cerevisiae*. Nucleic Acids Res., 34, D446-D451.
- Thevelein J.M., and de Winde, J.H. 1999. Novel sensing mechanisms and targets for the cAMP-protein kinase A pathway in the yeast *Saccharomyces cerevisiae*. Mol. Microbiol. 33, 904-918.
- van Wageningen, S., Kemmeren, P., Lijnzaad, P., Margaritis, T., Benschop, J. J., de Castro, I.
  J., van Leenen, D., Groot Koerkamp, M. J. A., Ko, C. W., Miles, A. J., Brabers, N.,
  Brok, M. O., Lenstra, T. L., Fiedler, D., Fokkens, L., Aldecoa, R., Apweiler, E.,
  Taliadouros, V., Sameith, K., van de Pasch, L. A. L., van Hooff, S. R., Bakker, L. V.,

Krogan, N. J., Snel, B. and Holstege, F. C. P. 2010. Functional overlap and regulatory links shape genetic interactions between signaling pathways, Cell, 143 (6), 991-1004.

- Wang, Y., Pierce, M., Schneper, L., Guldal, C. G., Zhang, X., Tavazoie, S. and Broach, J. R. 2004. Ras and Gpa2 mediate one branch of a redundant glucose signaling pathway in yeast, PLoS Biol., 2 (5), 610-622.
- Ward, M. P., Gimeno, C. J., Fink, G. R. and Garrett, S. 1995. SOK2 may regulate cyclic AMP-dependent protein kinase-stimulated growth and pseudohyphal development by repressing transcription, Mol. Cell Biol., 15 (12), 6854-6863.
- Watanabe, Y., Irie, K. and Matsumoto, K. 1995. Yeast *RLM1* encodes a serum response factor-like protein that may function downstream of the Mpk1 (Slt2) mitogen-activated protein kinase pathway, Mol. Cell Biol., 15, 5740-5749.
- White, M. A., Riles, L. and Cohen, B. A. 2009. A systematic screen for transcriptional regulators of the yeast cell cycle, Genetics, 181, 435-446.
- Wurgler-Murphy, S. M., Maeda, T., Witten, E. A. and Saito, H. 1997. Regulation of the Saccharomyces cerevisiae HOG1 mitogen-activated protein kinase by the PTP1 and PTP3 protein tyrosine phosphatases, Mol. Cell Biol., 17 (3), 1289-1297.
- Yamaguchi-Iwai, Y., Dancis, A., Klausner, R. D. 1995. AFT1: a mediator of iron regulated transcriptional control in *Saccharomyces cerevisiae*, EMBO J., 14 (6), 1231-1239.
- Yang, Y. H., Dudoit, S., Luu, P., Lin, D. M., Peng, V., Ngai, J. and Speed, T. P. 2002.
  Normalization for cDNA microarray data: a robust composite method addressing single and multiple slide systematic variation, Nucleic Acids Res. 30 (4), 1-10.
- Yokohama, H., Mizunuma, M., Okamoto, M., Yamamoto, J., Hirata, D., and Miyakawa, T.
  2006. Involvement of calcineurin-dependent degradation of Yap1p in Ca2+ -induced
  G2 cell-cycle regulation in *Saccharomyces cerevisiae*, EMBO reports, 7 (5), 519-524.

- Yoshimoto, H., Saltsman, K., Gasch, A. P., Li, H. X., Ogawa, N., Botstein, D., Brown, P. O. and Cyert, M. S. 2002. Genome-wide analysis of gene expression regulated by the calcineurin/Crz1p signaling pathway in *Saccharomyces cerevisiae*, J. Biol. Chem., 277, 31079-31088.
- Yuzyuk, T., Foehr, M. and Amberg, D.C. 2002. The MEK kinase Ssk2p promotes actin cytoskeleton recovery after osmotic stress. Mol. Biol. Cell, 13 (8), 2869-2880.
- Zhan, X. L., Hong, Y., Zhu, T., Mitchell, A. P., Deschenes, R. J., and Guan K. L. 2000. Essential functions of protein tyrosine phosphatases Ptp2 and Ptp3 and Rim11 tyrosine phosphorylation in *Saccharomyces cerevisiae* meiosis and sporulation, Mol. Biol. Cell 11 (2), 663-676.
- Zhao, C., Jung, U. S., Garrett-Engele, P., Roe, T., Cyert, M. S. and Levin, D. E. 1998.
  Temperature-induced expression of yeast *FKS2* is under the dual control of protein kinase C and calcineurin, Mol. Cell Biol., 18, 1013-1022.
- Zhao, X., Mehrabi, R. and Xu, J-R. 2007. Mitogen-activated protein kinase pathways and fungal pathogenesis, Eukaryot. Cell, 6, 1701-1714.

## **Publications**

Laviña, W. A., Shahsavarani, H., Saidi, A., Sugiyama, M., Kaneko, Y., and Harashima, S.: Suppression mechanism of the calcium sensitivity in *Saccharomyces cerevisiae ptp2\Delta msg5\Delta* double disruptant involves a novel HOG-independent function of Ssk2, transcription factor Msn2 and the PKA component Bcy1, *J. Biosci. Bioeng.* (In press).

Laviña, W. A., Hermansyah, Sugiyama, M., Kaneko, Y., and Harashima, S.: Functionally redundant protein phosphatase genes *PTP2* and *MSG5* co-regulate the calcium signaling pathway in *Saccharomyces cerevisiae* upon exposure to high extracellular calcium concentration, *J. Biosci. Bioeng.*, **115** (2), 138-146 (2013).

Hermansyah, Laviña, W. A., Sugiyama, M., Kaneko, Y., and Harashima, S.: Identification of protein kinase disruptions as suppressors of the calcium sensitivity of *S. cerevisiae*  $\Delta ptp2$  $\Delta msg5$  protein phosphatase double disruptant, *Arch. Microbiol.*, **192**, 157-165 (2010).

## Acknowledgements

I would like to express my sincerest gratitude to Professor Satoshi Harashima for giving me the opportunity to learn about yeast genetics. I will be forever indebted for his kindness and generosity. I would like to thank Professor Yoshinobu Kaneko for his neverending support during the duration of this study. I also thank Associate Professor Minetaka Sugiyama for all his help during the conduct of the experiment. His advice and constructive comments were indeed very helpful. I also thank Assistant Professor Yu Sasano for the fruitful discussions.

I would like to thank the members of my advisory committee, Professor Kazuhito Fujiyama and Professor Hajime Watanabe for accepting the advisory role and helping with the completion of my study. Their comments, insights and suggestions were very helpful in constructing a more coherent story of my study.

I would like to thank the past and present members of Harashima Laboratory for the kindness during my stay in Osaka University.

This work was partially supported by a Grant-in Aid for Scientific Research on Priority Areas, 16013227, 2004 to 2006, and by a Grant-in-Aid for Scientific Research B, 19380193, 2007 to 2009, to S. H. from the Ministry of Education, Science, Sports and Culture of Japan.

We also thank Dr. Abbas Saidi and Hosein Shahsavarani for their help with the analysis of the microarray data using Genowiz.

My parents, siblings, wife and daughter, to whom I dedicate this work. And most importantly, to God.

87