

Title	A FAKE HOMFLY POLYNOMIAL OF A KNOT
Author(s)	Miyazawa, Yasuyuki
Citation	Osaka Journal of Mathematics. 2013, 50(4), p. 1073-1096
Version Type	VoR
URL	<a href="https://doi.org/10.18910/26827">https://doi.org/10.18910/26827</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## A FAKE HOMFLY POLYNOMIAL OF A KNOT

YASUYUKI MIYAZAWA

(Received July 12, 2011, revised April 10, 2012)

### Abstract

We introduce a fake HOMFLY polynomial of a knot and show existence of such polynomials of a given knot.

### 1. Introduction

In this paper, we deal with the HOMFLY polynomial [2, 10, 13] of a knot.

The HOMFLY polynomial  $P(L; v, z) \in \mathbb{Z}[v^{\pm 1}, z^{\pm 1}]$  of an oriented link  $L$  is an invariant of the isotopy type of  $L$ , which is defined by the following formulas:

- (1)  $P(U; v, z) = 1$ ;
- (2)  $v^{-1}P(L_+; v, z) - vP(L_-; v, z) = zP(L_0; v, z)$ ,

where  $U$  is the trivial knot and  $L_+$ ,  $L_-$  and  $L_0$  are three links that are identical except near one point where they are as in Fig. 1.

The reduced polynomial  $P(L; 1, z)$  of  $L$  is called the Conway polynomial [1] of  $L$  and denoted by  $\nabla(L; z)$ . The Jones polynomial  $V(L; t)$  [7] of  $L$  is defined as the reduced polynomial  $P(L; t, t^{1/2} - t^{-1/2})$ .

By [10], the HOMFLY polynomial of an oriented knot  $K$  is of the form

$$P(K; v, z) = \sum_{j \geq 0} P_{2j}(K; v) z^{2j},$$

where each Laurent polynomial  $P_{2j}(K; v) \in \mathbb{Z}[v^{\pm 2}]$  is called the  $2j$ -th coefficient polynomial of  $P(K; v, z)$  in  $z$  or the  $2j$ -th HOMFLY coefficient polynomial. The  $2j$ -th coefficient polynomial of a knot is said to be *trivial* if it coincides with that of the trivial knot.

In [11], the author shows that there are infinitely many 2-bridge knots with trivial 0-th HOMFLY coefficient polynomial. The purpose of the paper is to explore a little further into HOMFLY coefficient polynomials of a knot.

Let  $K$  and  $K'$  be oriented knots. The HOMFLY polynomial  $P(K'; v, z)$  of  $K'$  is said to be a *fake HOMFLY polynomial of  $K$  with identical order  $2m$*  if  $P(K'; v, z) \equiv$

---

2010 Mathematics Subject Classification. Primary 57M25; Secondary 57M27.

The author was partially supported by Grant-in-Aid for Scientific Research (C) (No. 23540092), Japan Society for the Promotion of Science.




Fig. 1. A skein triple.




Fig. 2. Pass-moves.

$P(K; v, z) \pmod{z^{2m+2}}$ , that is  $P_{2j}(K'; v, z) = P_{2j}(K; v, z)$ ,  $0 \leq j \leq m$ , in terms of HOMFLY coefficient polynomials of  $K$  and  $K'$ .

REMARK 1.1. If the HOMFLY polynomial of  $K'$  is a fake HOMFLY polynomial of  $K$  with identical order  $2m$ , then it is also a fake HOMFLY polynomial of  $K$  with identical order  $2j$ ,  $0 \leq j < m$ .

In order to state our main theorems we give some definitions.

The Gordian distance from  $K$  to  $K'$  is defined to be the minimum number of crossing changes needed to transform  $K$  into  $K'$ . We denote it by  $d_G(K, K')$ .

A pass-move [8] is a local move on a diagram of an oriented knot as in Fig. 2. Two knots are pass-equivalent if one can be obtained from the other by a combination of Reidemeister moves and pass-moves. If  $K$  and  $K'$  are pass-equivalent, then the pass distance from  $K$  to  $K'$  is defined to be the minimum number of pass-moves needed to change  $K$  into  $K'$ . We denote it by  $d_{\text{pass}}(K, K')$ .

Let  $f$  be an isotopy type invariant of an oriented link, which takes values in an abelian group. Then,  $f$  can be uniquely extended to a singular link invariant by the Vassiliev skein relation:

$$f(L_\times) = f(L_+) - f(L_-),$$

where  $L_\times$  is a singular link with a double point  $x$ , and  $L_+$  and  $L_-$  are ones obtained from  $L_\times$  by replacing  $x$  with a positive and a negative crossing, respectively. We call  $f$  a finite type invariant if there exists an integer  $q$  such that  $f(L) = 0$  for any singular link  $L$  with more than  $q$  double points. The smallest of such integers is the order of  $f$ . We denote a finite type invariant of  $L$  with order  $q$  by  $f_q(L)$ .

The signature of  $K$ ,  $\sigma(K)$ , is a cobordism invariant of a knot; see [12], Section 8F in [14].

**Theorem 1.2** ([9]). *Let  $m, l$  and  $q$  be integers with  $m \geq 0$  and  $l, q > 0$ . For an oriented knot  $K$ , there exist infinitely many knots  $\{K_n; n \in \mathbb{N}\}$  with the following properties:*

- (1) *The HOMFLY polynomial of  $K_n$  is a fake HOMFLY polynomial of  $K$  with identical order  $2m$ ;*
- (2)  $P(K_n; v, z) \equiv P(K; v, z) \pmod{(v^{2l} - 1)}$ ;
- (3)  $\nabla(K_n; z) = \nabla(K; z)$ ;
- (4)  $K_n$  *is a band sum of  $K$  and the trivial knot, and so  $K_n$  is cobordant to  $K$  and  $\sigma(K_n) = \sigma(K)$ ;*
- (5)  $d_G(K_n, K) = 1$ ;
- (6)  $d_{\text{pass}}(K_n, K) = 1$ ;
- (7)  $f_j(K_n) = f_j(K)$ ,  $0 \leq j \leq q$ .

REMARK 1.3. Kawauchi informed the author that Theorem 1.2 could be accomplished by using imitation theory, that is, appropriate almost identical link imitations of a given knot had the properties in the theorem. However, in this paper, the proof of the theorem is given by an elementary method of construction which shows explicitly diagrams of knots.

**Corollary 1.4.** *For a non-negative integer  $m$ , there exist infinitely many knots  $\{K_n; n \in \mathbb{N}\}$  whose  $2j$ -th HOMFLY coefficient polynomials,  $0 \leq j \leq m$ , are trivial.*

Since an almost identical link imitation of a knot has the same Alexander polynomial as the knot, imitation theory does not work on the following theorem.

**Theorem 1.5.** *Let  $m$  be a non-negative integer and  $q$  a positive integer. For an oriented knot  $K$ , there exist infinitely many knots  $\{K_n; n \in \mathbb{N}\}$  with the following properties:*

- (1) *The HOMFLY polynomial of  $K_n$  is a fake HOMFLY polynomial of  $K$  with identical order  $2m$ ;*
- (2)  $\nabla(K_n; z) \neq \nabla(K; z)$ , *and so  $K_n$  is not an imitation of  $K$ ;*
- (3)  $d_G(K_n, K) = 1$ ;
- (4)  $d_{\text{pass}}(K_n, K) = 1$ ;
- (5)  $\sigma(K_n) = \sigma(K)$ ;
- (6)  $f_j(K_n) = f_j(K)$ ,  $0 \leq j \leq q$ .

To prove theorems, we make use of polynomials derived from a tangle which comes from decomposition of a knot. We introduce them in Section 2. The proofs of theorems are given in Sections 4 and 6 after preliminaries in Sections 3 and 5.

**2. The normal coordinates of a tangle**

A tangle  $T$  is a pair  $(B^3, t)$  of a 3-ball  $B^3$  and a proper 1-submanifold  $t$  with  $\partial t \neq \emptyset$ .  $T$  is said to be a 2-string tangle if  $T$  consists of two arcs and some circle components. Each of  $\partial t$  is called an endpoint of  $T$ .  $T$  is called *properly oriented* if each arc of  $T$  is oriented as in Fig. 3.

The numerator (resp. denominator) of  $T$  denoted by  $N(T)$  (resp.  $D(T)$ ) is a link obtained from  $T$  by connecting four endpoints of  $T$  by two arcs outside  $T$  as in the left (resp. right) figure of Fig. 4.

A tangle  $T$  is said to be of *type*  $N_{\mu(N(T))}$  (resp. *type*  $D_{\mu(D(T))}$ ) or an  $N_{\mu(N(T))}$ -tangle (resp. a  $D_{\mu(D(T))}$ -tangle) if  $T$  is a properly oriented 2-string tangle and  $\mu(N(T)) < \mu(D(T))$  (resp.  $\mu(N(T)) > \mu(D(T))$ ), where  $\mu(L)$  denotes the number of components of a link  $L$ .

We denote by  $E_{2n}$ ,  $n \in \mathbb{Z}$ , and  $E_\infty$  tangles of type  $D_1$  and of type  $N_1$  as in Fig. 5, respectively.  $E_{2n}$  has  $2|n|$  positive (resp. negative) crossings if  $n > 0$  (resp.  $n < 0$ ) and  $E_0$  means horizontal parallel strings without crossings.

Let  $L(T)$ ,  $L(E_0)$  and  $L(E_\infty)$  be three links identical outside a ball and inside are a properly oriented 2-string tangle  $T$ , the  $D_1$ -tangle  $E_0$  and the  $N_1$ -tangle  $E_\infty$ , respectively.

**Lemma 2.1.** *Let  $T$  be a tangle of type  $D_{\mu(D(T))}$ . Then, there exists a unique pair  $(e_0(T; v, z), e_\infty(T; v, z))$  of polynomials for  $T$  so that*

$$\begin{aligned} P(L(T); v, z) &= (vz)^{1-\mu(D(T))} \{e_0(T; v, z)P(L(E_0); v, z) + vze_\infty(T; v, z)P(L(E_\infty); v, z)\}, \end{aligned}$$

where  $e_0(T; v, z), e_\infty(T; v, z) \in \mathbb{Z}[v^{\pm 2}, z^2]$ .

*Proof.* Linear skein theory gives a unique pair  $(e_0(T; v, z), e_\infty(T; v, z))$  of polynomials for  $T$  so that

$$\begin{aligned} P(L(T); v, z) &= (vz)^{1-\mu(D(T))} \{e_0(T; v, z)P(L(E_0); v, z) + vze_\infty(T; v, z)P(L(E_\infty); v, z)\}, \end{aligned}$$

where  $e_0(T; v, z), e_\infty(T; v, z) \in \mathbb{Z}[v^{\pm 1}, z^{\pm 1}]$ . We only have to show that  $e_0(T; v, z)$  and  $e_\infty(T; v, z)$  are elements of  $\mathbb{Z}[v^{\pm 2}, z^2]$ . Considering the HOMFLY polynomials of the numerator and the denominator of  $T$ , we have

$$P(D(T); v, z) = (vz)^{1-\mu(D(T))}h(D(T); v, z)$$

and

$$P(N(T); v, z) = (vz)^{1-\mu(D(T))}(vz)^{-1}h(N(T); v, z),$$




Fig. 3. A properly oriented 2-string tangle.



Fig. 4. The numerator and the denominator.



Fig. 5. Trivial tangles.

where  $h(D(T); v, z), h(N(T); v, z) \in \mathbb{Z}[v^{\pm 2}, z^2]$ , because  $\mu(N(T)) = \mu(D(T)) + 1$ . Thus,

$$h(D(T); v, z) = e_0(T; v, z)P(D(E_0); v, z) + vze_{\infty}(T; v, z)P(D(E_{\infty}); v, z)$$

and

$$h(N(T); v, z) = vze_0(T; v, z)P(N(E_0); v, z) + v^2z^2e_{\infty}(T; v, z)P(N(E_{\infty}); v, z).$$

Since  $D(E_0)$  and  $N(E_{\infty})$  are trivial knots and  $N(E_0)$  and  $D(E_{\infty})$  are 2-component trivial links, we obtain

$$h(D(T); v, z) = e_0(T; v, z) + (1 - v^2)e_{\infty}(T; v, z)$$

and

$$h(N(T); v, z) = (1 - v^2)e_0(T; v, z) + v^2z^2e_{\infty}(T; v, z).$$

From these equalities, we have

$$\{(1 - v^2)^2 - v^2z^2\}e_{\infty}(T; v, z) = (1 - v^2)h(D(T); v, z) - h(N(T); v, z).$$

It follows that  $e_\infty(T; v, z) \in \mathbb{Z}[v^{\pm 2}, z^2]$ , which leads to  $e_0(T; v, z) \in \mathbb{Z}[v^{\pm 2}, z^2]$ .  $\square$

**Lemma 2.2.** *Let  $T$  be a tangle of type  $N_{\mu(N(T))}$ . Then, there exists a unique pair  $(e_0(T; v, z), e_\infty(T; v, z))$  of polynomials for  $T$  so that*

$$\begin{aligned} &P(L(T); v, z) \\ &= (vz)^{1-\mu(N(T))}\{vze_0(T; v, z)P(L(E_0); v, z) + e_\infty(T; v, z)P(L(E_\infty); v, z)\}, \end{aligned}$$

where  $e_0(T; v, z), e_\infty(T; v, z) \in \mathbb{Z}[v^{\pm 2}, z^2]$ .

*Proof.* The proof of the lemma is similar to that of Lemma 2.1.  $\square$

The polynomials  $e_0(T; v, z)$  and  $e_\infty(T; v, z)$  which appear in Lemmas 2.1 or 2.2 are essentially determined by the tangle  $T$  only. So, a pair  $(e_0(T; v, z), e_\infty(T; v, z))$  of the polynomials is called the *normal coordinates* of  $T$ .

Let  $M_D$  and  $M_N$  be  $2 \times 2$  matrices whose entries are in  $\mathbb{Z}[v^{\pm 1}, z^{\pm 1}]$  defined by  $\begin{pmatrix} (v^{-1} - v)z^{-1} & vz \\ 1 & 1 - v^2 \end{pmatrix}$  and  $\begin{pmatrix} 1 - v^2 & 1 \\ vz & (v^{-1} - v)z^{-1} \end{pmatrix}$ , respectively.

Let  $3_1$  [14] be the trefoil knot. We put  $\lambda(v, z) = -(1 - v^2)^2 + v^2z^2 = P(3_1!; v, z) - 1$ , where  $K!$  denotes the mirror image of a knot  $K$ .

REMARK 2.3.  $\det M_D = \det M_N = -v^{-1}z^{-1}\lambda(v, z) \neq 0$ .

The following two lemmas are corollaries of Lemmas 2.1 and 2.2.

**Lemma 2.4.** *Let  $T$  be a tangle of type  $D_{\mu(D(T))}$ . Then, the normal coordinates  $(e_0(T; v, z), e_\infty(T; v, z))$  of  $T$  are expressed as follows:*

$$\begin{aligned} \begin{pmatrix} e_0(T; v, z) \\ e_\infty(T; v, z) \end{pmatrix} &= (vz)^{\mu(D(T))-1} M_D^{-1} \begin{pmatrix} P(N(T); v, z) \\ P(D(T); v, z) \end{pmatrix} \\ &= \frac{(vz)^{\mu(D(T))-1}}{\lambda(v, z)} \begin{pmatrix} -v(1 - v^2)zP(N(T); v, z) + v^2z^2P(D(T); v, z) \\ vzP(N(T); v, z) - (1 - v^2)P(D(T); v, z) \end{pmatrix}. \end{aligned}$$

*Proof.* Let  $v = (vz)^{\mu(D(T))-1}$ . By Lemma 2.1 and equalities  $P(N(E_0); v, z) = P(D(E_\infty); v, z) = (v^{-1} - v)z^{-1}$  and  $P(N(E_\infty); v, z) = P(D(E_0); v, z) = 1$ , we have

$$\begin{aligned} v \begin{pmatrix} P(N(T); v, z) \\ P(D(T); v, z) \end{pmatrix} &= \begin{pmatrix} P(N(E_0); v, z) & vzP(N(E_\infty); v, z) \\ P(D(E_0); v, z) & vzP(D(E_\infty); v, z) \end{pmatrix} \begin{pmatrix} e_0(T; v, z) \\ e_\infty(T; v, z) \end{pmatrix} \\ &= M_D \begin{pmatrix} e_0(T; v, z) \\ e_\infty(T; v, z) \end{pmatrix}. \end{aligned}$$

Thus, we obtain the first expression of the normal coordinates of  $T$ . Since  $M_D^{-1} = (-vz/\lambda(v, z)) \begin{pmatrix} 1 - v^2 & -vz \\ -1 & (v^{-1} - v)z^{-1} \end{pmatrix}$ , we have the second equality.  $\square$




Fig. 6. Addition of tangles.

**Lemma 2.5.** *Let  $T$  be a tangle of type  $N_{\mu(N(T))}$ . Then, the normal coordinates  $(e_0(T; v, z), e_\infty(T; v, z))$  of  $T$  are expressed as follows:*

$$\begin{aligned} \begin{pmatrix} e_0(T; v, z) \\ e_\infty(T; v, z) \end{pmatrix} &= (vz)^{\mu(N(T))-1} M_N^{-1} \begin{pmatrix} P(N(T); v, z) \\ P(D(T); v, z) \end{pmatrix} \\ &= \frac{(vz)^{\mu(N(T))-1}}{\lambda(v, z)} \begin{pmatrix} -(1 - v^2)zP(N(T); v, z) + vzP(D(T); v, z) \\ v^2z^2P(N(T); v, z) - v(1 - v^2)P(D(T); v, z) \end{pmatrix}. \end{aligned}$$

Proof. The proof of the lemma is similar to that of Lemma 2.4. □

Let  $T$  and  $S$  be properly oriented 2-string tangles. We define addition of tangles  $T$  and  $S$  by connecting endpoints of  $T$  and  $S$  as in Fig. 6 and denote it by  $T + S$ .

If  $T$  is a  $D_1$ -tangle and  $S$  is an  $N_1$ -tangle, then  $T + S$  is an  $N_1$ -tangle.

**Lemma 2.6.** *Let  $T$  be a  $D_1$ -tangle and  $S$  an  $N_1$ -tangle. Let  $(e_0(T; v, z), e_\infty(T; v, z))$  and  $(e_0(S; v, z), e_\infty(S; v, z))$  be the normal coordinates of  $T$  and  $S$ , respectively. Then, the normal coordinates  $(e_0(T + S; v, z), e_\infty(T + S; v, z))$  of the tangle  $T + S$  are expressed as follows:*

$$e_0(T + S; v, z) = e_0(T; v, z)e_0(S; v, z)$$

and

$$\begin{aligned} e_\infty(T + S; v, z) &= e_0(T; v, z)e_\infty(S; v, z) + v^2z^2e_\infty(T; v, z)e_0(S; v, z) \\ &\quad + (1 - v^2)e_\infty(T; v, z)e_\infty(S; v, z). \end{aligned}$$

Proof. Note that  $L(E_0 + S) = L(S)$  and  $L(E_\infty + S)$  is the connected sum of  $L(E_\infty)$  and  $D(S)$ . By using Lemma 2.1, we have

$$\begin{aligned} P(L(T + S); v, z) &= e_0(T; v, z)P(L(E_0 + S); v, z) \\ &\quad + vze_\infty(T; v, z)P(L(E_\infty + S); v, z) \\ &= e_0(T; v, z)P(L(S); v, z) \\ &\quad + vze_\infty(T; v, z)P(D(S); v, z)P(L(E_\infty); v, z). \end{aligned}$$



Since

$$\begin{aligned} P(D(S); v, z) &= vze_0(S; v, z)P(D(E_0); v, z) + e_\infty(S; v, z)P(D(E_\infty); v, z) \\ &= vze_0(S; v, z) + (v^{-1} - v)z^{-1}e_\infty(S; v, z), \end{aligned}$$

we obtain

$$\begin{aligned} &P(L(T + S); v, z) \\ &= e_0(T; v, z)\{vze_0(S; v, z)P(L(E_0); v, z) + e_\infty(S; v, z)P(L(E_\infty); v, z)\} \\ &\quad + vze_\infty(T; v, z)\{vze_0(S; v, z) + (v^{-1} - v)z^{-1}e_\infty(S; v, z)\}P(L(E_\infty); v, z) \\ &= vze_0(T; v, z)e_0(S; v, z)P(L(E_0); v, z) \\ &\quad + \{e_0(T; v, z)e_\infty(S; v, z) + v^2z^2e_\infty(T; v, z)e_0(S; v, z) \\ &\quad + (1 - v^2)e_\infty(T; v, z)e_\infty(S; v, z)\}P(L(E_\infty); v, z). \end{aligned}$$

This completes the proof.  $\square$

**Lemma 2.7.** *Under the same assumption as Lemma 2.6,*

$$\begin{aligned} P(N(T + S); v, z) &= (1 - v^2)e_0(T; v, z)e_0(S; v, z) + e_0(T; v, z)e_\infty(S; v, z) \\ &\quad + v^2z^2e_\infty(T; v, z)e_0(S; v, z) + (1 - v^2)e_\infty(T; v, z)e_\infty(S; v, z). \end{aligned}$$

*Proof.* Since  $P(N(E_0); v, z) = (v^{-1} - v)z^{-1}$  and  $P(N(E_\infty); v, z) = 1$ , Lemma 2.6 shows the claim.  $\square$

**Proposition 2.8.** *Under the same assumption as Lemma 2.6,*

$$P(N(T + S); v, z) = P(D(T); v, z)P(N(S); v, z) + \lambda(v, z)e_\infty(T; v, z)e_0(S; v, z).$$

*Proof.* Since  $P(D(T); v, z) = e_0(T; v, z) + (1 - v^2)e_\infty(T; v, z)$  and  $P(N(S); v, z) = (1 - v^2)e_0(S; v, z) + e_\infty(S; v, z)$ , by Lemma 2.7,

$$\begin{aligned} &P(D(T); v, z)P(N(S); v, z) \\ &= (1 - v^2)e_0(T; v, z)e_0(S; v, z) + e_0(T; v, z)e_\infty(S; v, z) \\ &\quad + (1 - v^2)^2e_\infty(T; v, z)e_0(S; v, z) + (1 - v^2)e_\infty(T; v, z)e_\infty(S; v, z) \\ &= P(N(T + S); v, z) + \{(1 - v^2)^2 - v^2z^2\}e_\infty(T; v, z)e_0(S; v, z), \end{aligned}$$

completing the proof.  $\square$

Combining Proposition 2.8 with Lemmas 2.4 and 2.5, we easily obtain the following.

**Corollary 2.9.** *Let  $T$  be a  $D_1$ -tangle and  $S$  an  $N_1$ -tangle. Then,*

$$\begin{aligned} &\lambda(v, z)P(N(T + S); v, z) \\ &= -v(1 - v^2)z\{P(N(T); v, z)P(N(S); v, z) + P(D(T); v, z)P(D(S); v, z)\} \\ &\quad + v^2z^2\{P(N(T); v, z)P(D(S); v, z) + P(D(T); v, z)P(N(S); v, z)\}. \end{aligned}$$

**Lemma 2.10.** *Let  $T$  and  $S$  be tangles of type  $D_1$ . Let  $(e_0(T; v, z), e_\infty(T; v, z))$  and  $(e_0(S; v, z), e_\infty(S; v, z))$  be normal coordinates of  $T$  and  $S$ , respectively. Then, the normal coordinates  $(e_0(T + S; v, z), e_\infty(T + S; v, z))$  of the  $D_1$ -tangle  $T + S$  are expressed as follows:*

$$e_0(T + S; v, z) = e_0(T; v, z)e_0(S; v, z)$$

and

$$\begin{aligned} e_\infty(T + S; v, z) &= e_0(T; v, z)e_\infty(S; v, z) \\ &\quad + e_\infty(T; v, z)\{e_0(S; v, z) + (1 - v^2)e_\infty(S; v, z)\}. \end{aligned}$$

*Proof.* Since  $L(E_0 + S) = L(S)$ , and  $L(E_\infty + S)$  is the connected sum of  $L(E_\infty)$  and  $D(S)$ , by Lemma 2.1, we have

$$\begin{aligned} P(L(T + S); v, z) &= e_0(T; v, z)P(L(E_0 + S); v, z) \\ &\quad + vze_\infty(T; v, z)P(L(E_\infty + S); v, z) \\ &= e_0(T; v, z)P(L(S); v, z) \\ &\quad + vze_\infty(T; v, z)P(D(S); v, z)P(L(E_\infty); v, z). \end{aligned}$$

Since  $P(L(S); v, z) = e_0(S; v, z)P(L(E_0); v, z) + vze_\infty(S; v, z)P(L(E_\infty); v, z)$ , the last expression is  $e_0(T; v, z)e_0(S; v, z)P(L(E_0); v, z) + vze_\infty(T; v, z)e_0(S; v, z) + e_\infty(T; v, z)P(D(S); v, z)P(L(E_\infty); v, z)$ . Thus, we obtain

$$e_0(T + S; v, z) = e_0(T; v, z)e_0(S; v, z)$$

and

$$e_\infty(T + S; v, z) = e_0(T; v, z)e_\infty(S; v, z) + e_\infty(T; v, z)P(D(S); v, z).$$

Since

$$\begin{aligned} P(D(S); v, z) &= e_0(S; v, z)P(D(E_0); v, z) + vze_\infty(S; v, z)P(L(E_\infty); v, z) \\ &= e_0(S; v, z) + (1 - v^2)e_\infty(S; v, z), \end{aligned}$$

we have the result. □

### 3. Preliminaries for the proof of Theorem 1.2

Let  $X$  and  $Y$  be properly oriented 2-string tangles and  $R(X, Y)$  the properly oriented 2-string tangle illustrated in Fig. 7.




Fig. 7. Tangle  $R(X, Y)$ .

Note that  $N(R(X, Y)) = D(Y + X)$ . It is easy to see the following.

**Lemma 3.1.** *Let  $U$  and  $U_2$  be the trivial knot and the 2-component trivial link respectively. Then,  $R(E_0, E_0) = E_\infty$ ,  $R(E_0, E_\infty) = E_\infty \sqcup U$ ,  $R(E_\infty, E_\infty) = E_\infty \sqcup U_2$ .*

**Lemma 3.2.** *Let  $X$  and  $Y$  be tangles of type  $D_1$ . Let  $(e_0(X; v, z), e_\infty(X; v, z))$ ,  $(e_0(Y; v, z), e_\infty(Y; v, z))$  and  $(e_0(R(E_\infty, E_0); v, z), e_\infty(R(E_\infty, E_0); v, z))$  be the normal coordinates of  $X$ ,  $Y$  and  $R(E_\infty, E_0)$ , respectively. Then, the normal coordinates  $(e_0(R(X, Y); v, z), e_\infty(R(X, Y); v, z))$  of the tangle  $R(X, Y)$  are expressed as follows:*

$$e_0(R(X, Y); v, z) = e_\infty(X; v, z)e_0(Y; v, z)e_0(R(E_\infty, E_0); v, z)$$

and

$$\begin{aligned} e_\infty(R(X, Y); v, z) &= e_0(X; v, z)e_0(Y; v, z) + (1 - v^2)e_0(X; v, z)e_\infty(Y; v, z) \\ &\quad + (1 - v^2)^2e_\infty(X; v, z)e_\infty(Y; v, z) \\ &\quad + e_\infty(X; v, z)e_0(Y; v, z)e_\infty(R(E_\infty, E_0); v, z). \end{aligned}$$

Proof. By using normal coordinates of  $X$  and  $Y$ , we have

$$\begin{aligned} P(L(R(X, Y)); v, z) &= e_0(X; v, z)P(L(R(E_0, Y)); v, z) \\ &\quad + vze_\infty(X; v, z)P(L(R(E_\infty, Y)); v, z) \\ &= e_0(X; v, z)\{e_0(Y; v, z)P(L(R(E_0, E_0)); v, z) \\ &\quad + vze_\infty(Y; v, z)P(L(R(E_0, E_\infty)); v, z)\} \\ &\quad + vze_\infty(X; v, z)\{e_0(Y; v, z)P(L(E_\infty, E_0)); v, z) \\ &\quad + vze_\infty(Y; v, z)P(L(R(E_\infty, E_\infty)); v, z)\} \end{aligned}$$

$$\begin{aligned}
 &= e_0(X; v, z)e_0(Y; v, z)P(L(R(E_0, E_0)); v, z) \\
 &\quad + vze_0(X; v, z)e_\infty(Y; v, z)P(L(R(E_0, E_\infty)); v, z) \\
 &\quad + vze_\infty(X; v, z)e_0(Y; v, z)P(L(R(E_\infty, E_0)); v, z) \\
 &\quad + v^2z^2e_\infty(X; v, z)e_\infty(Y; v, z)P(L(R(E_\infty, E_\infty)); v, z).
 \end{aligned}$$

By Lemma 3.1, we obtain

$$\begin{aligned}
 P(L(R(E_0, E_0)); v, z) &= P(L(E_\infty); v, z), \\
 P(L(R(E_0, E_\infty)); v, z) &= (v^{-1} - v)z^{-1}P(L(R(E_\infty)); v, z)
 \end{aligned}$$

and

$$P(L(R(E_\infty, E_\infty)); v, z) = \{(v^{-1} - v)z^{-1}\}^2P(L(R(E_\infty)); v, z).$$

Since  $R(E_\infty, E_0)$  is an  $N_2$ -tangle, we see that

$$\begin{aligned}
 P(L(R(E_\infty, E_0)); v, z) &= e_0(R(E_\infty, E_0); v, z)P(L(E_0); v, z) \\
 &\quad + (vz)^{-1}e_\infty(R(E_\infty, E_0); v, z)P(L(E_\infty); v, z).
 \end{aligned}$$

From these equalities, we find that  $P(L(R(X, Y)); v, z)$  is equal to

$$\begin{aligned}
 &vze_\infty(X; v, z)e_0(Y; v, z)e_0(R(E_\infty, E_0); v, z)P(L(E_0); v, z) \\
 &\quad + \{e_0(X; v, z)e_0(Y; v, z) \\
 &\quad \quad + (1 - v^2)e_0(X; v, z)e_\infty(Y; v, z) \\
 &\quad \quad + (1 - v^2)^2e_\infty(X; v, z)e_\infty(Y; v, z) \\
 &\quad \quad + e_\infty(X; v, z)e_0(Y; v, z)e_\infty(R(E_\infty, E_0); v, z)\}P(L(E_\infty); v, z).
 \end{aligned}$$

Since  $R(X, Y)$  is an  $N_1$ -tangle, we obtain the claim. □

The proof of the following lemma is straightforward.

**Lemma 3.3.**

$$P(N(R(E_\infty, E_0)); v, z) = (v^{-1} - v)z^{-1}$$

and

$$\begin{aligned}
 P(D(R(E_\infty, E_0)); v, z) &= v^{-2}(v^{-1} - v)^2\{v^2z^{-2} + 2(v^{-1} - v)^2 \\
 &\quad \quad \quad + (5v^{-2} + 12 + 5v^2)z^2 \\
 &\quad \quad \quad + (4v^{-2} - 13 + 4v^2)z^4 \\
 &\quad \quad \quad + (v^{-2} - 6 + v^2)z^6 - z^8\}.
 \end{aligned}$$

**Lemma 3.4.**

$$e_0(R(E_\infty, E_0); v, z) = -v^{-2}(v^{-1} - v)^2 z^2(1 + z^2)^2(2 + z^2).$$

Proof. Note that  $R(E_\infty, E_0)$  is an  $N_2$ -tangle. Put  $T = R(E_\infty, E_0)$ . By Lemma 2.5, we have

$$e_0(T; v, z) = \frac{vz}{\lambda(v, z)} \{- (1 - v^2)P(N(T); v, z) + vzP(D(T); v, z)\}.$$

Lemma 3.3 ensures the claim. □

For a properly oriented 2-string tangle  $T$ , we denote by  $T^\perp$  the properly oriented 2-string tangle obtained from  $T$  by reversing orientations of strings after rotating through angle  $\pi/2$  in the anticlockwise direction about an axis perpendicular to the projective plane. We call it the *rotation* of  $T$ .

**Lemma 3.5.** *Let  $(e_0(T; v, z), e_\infty(T; v, z))$  and  $(e_0(T^\perp; v, z), e_\infty(T^\perp; v, z))$  be the normal coordinates of an  $N_1$ -tangle  $T$  and its rotation  $T^\perp$ . Then,*

$$(e_0(T^\perp; v, z), e_\infty(T^\perp; v, z)) = (e_\infty(T; v, z), e_0(T; v, z)).$$

Proof. Let  $\Psi$  be a resolution tree for  $L(T)$  obtained by switching or smoothing crossings of  $T$ . We may assume that each end node  $L(S_j)$ ,  $1 \leq j \leq n$ , of  $\Psi$  is disjoint union of  $L(E_0)$  or  $L(E_\infty)$  with some circle components. Then, we obtain a resolution tree for  $L(T^\perp)$  from  $\Psi$  by replacing each end node  $L(S_j)$  of  $\Psi$  with  $L(S_j^\perp)$ . From the two resolution trees, we see that the HOMFLY polynomials of  $L(T)$  and  $L(T^\perp)$  can be written as

$$P(L(T); v, z) = \sum_{j=1}^n h(L(S_j); v, z) P(L(S_j); v, z)$$

and

$$P(L(T^\perp); v, z) = \sum_{j=1}^n h(L(S_j); v, z) P(L(S_j^\perp); v, z),$$

where  $h(L(S_j); v, z) \in \mathbb{Z}[v^{\pm 1}, z^{\pm 1}]$ . Thus, if

$$P(L(T); v, z) = g_0(L(T); v, z)P(L(E_0); v, z) + g_\infty(L(T); v, z)P(L(E_\infty); v, z),$$

where  $g_0(L(T); v, z), g_\infty(L(T); v, z) \in \mathbb{Z}[v^{\pm 1}, z^{\pm 1}]$ , then we obtain

$$P(L(T^\perp); v, z) = g_\infty(L(T); v, z)P(L(E_0); v, z) + g_0(L(T); v, z)P(L(E_\infty); v, z).$$

Since the tangle  $T^\perp$  is of type  $D_1$ , we have the claim. □

Let  $X_0$  be an  $N_1$ -tangle and  $\{Y_n; n \in \mathbb{N}\}$  a set of  $D_1$ -tangles. We define a sequence  $\{W_n; n \in \mathbb{N}\}$  of  $N_1$ -tangles by the following recursive formulas:

- (1)  $W_1 = R(X_0^\perp, Y_1)$ ;
- (2)  $W_n = R(W_{n-1}^\perp, Y_n), n \geq 2$ .

**Proposition 3.6.** *For any positive integer  $n$ ,*

$$e_0(W_n; v, z) = e_0(X_0; v, z) \left( \prod_{j=1}^n e_0(Y_j; v, z) \right) e_0(R(E_\infty, E_0); v, z)^n.$$

*Proof.* The proof is by an induction on the number  $n$ . If  $n = 1$ , then Lemmas 3.2 and 3.5 show that

$$\begin{aligned} e_0(W_1; v, z) &= e_0(R(X_0^\perp, Y_1); v, z) \\ &= e_\infty(X_0^\perp; v, z) e_0(Y_1; v, z) e_0(R(E_\infty, E_0); v, z) \\ &= e_0(X_0; v, z) e_0(Y_1; v, z) e_0(R(E_\infty, E_0); v, z). \end{aligned}$$

Thus, the claim is true. Suppose that  $n > 1$ . By Lemmas 3.2 and 3.5 and the inductive hypothesis, we have

$$\begin{aligned} e_0(W_n; v, z) &= e_0(R(W_{n-1}^\perp, Y_n); v, z) \\ &= e_\infty(W_{n-1}^\perp; v, z) e_0(Y_n; v, z) e_0(R(E_\infty, E_0); v, z) \\ &= e_0(W_{n-1}; v, z) e_0(Y_n; v, z) e_0(R(E_\infty, E_0); v, z) \\ &= e_0(X_0; v, z) \left( \prod_{j=1}^n e_0(Y_j; v, z) \right) e_0(R(E_\infty, E_0); v, z)^n. \end{aligned}$$

This completes the proof. □

The normal coordinates of the tangle  $E_{2n}, n \in \mathbb{Z}$ , are the following.

**Lemma 3.7.**  $(e_0(E_{2n}; v, z), e_\infty(E_{2n}; v, z)) = (v^{2n}, (v^{2n} - 1)/(v^2 - 1))$ .

*Proof.* Let  $\varepsilon_n = n/|n|$ . Since  $P(L(E_{2(n-\varepsilon)}); v, z) = e_0(E_{2(n-\varepsilon)}; v, z)P(L(E_0); v, z) + vze_\infty(E_{2(n-\varepsilon)}; v, z)P(L(E_\infty); v, z)$ , we have

$$\begin{aligned} P(L(E_{2n}); v, z) &= v^{2\varepsilon_n} P(L(E_{2(n-\varepsilon)}); v, z) + \varepsilon_n v^{\varepsilon_n} z P(L(E_\infty); v, z) \\ &= v^{2\varepsilon_n} e_0(E_{2(n-\varepsilon)}; v, z) P(L(E_0); v, z) \\ &\quad + v z \{ v^{2\varepsilon_n} e_\infty(E_{2(n-\varepsilon)}; v, z) + \varepsilon_n v^{1-\varepsilon_n} \} P(L(E_\infty); v, z). \end{aligned}$$




Fig. 8. Tangle \$H\_q\$.

Thus, we obtain

$$\begin{pmatrix} e_0(E_{2n}; v, z) \\ e_\infty(E_{2n}; v, z) \end{pmatrix} = v^{2n} \begin{pmatrix} e_0(E_0; v, z) \\ e_\infty(E_0; v, z) \end{pmatrix} + \begin{pmatrix} 0 \\ \frac{v^{2n} - 1}{v^2 - 1} \end{pmatrix}.$$

Since \$e\_0(E\_0; v, z) = 1\$ and \$e\_\infty(E\_0; v, z) = 0\$, we have the desired polynomials. □

Lemmas 3.5 and 3.7 give the normal coordinates of \$E\_{2n}^\perp\$, \$n \in \mathbb{Z}\$.

**Corollary 3.8.** \$(e\_0(E\_{2n}^\perp; v, z), e\_\infty(E\_{2n}^\perp; v, z)) = ((v^{2n} - 1)/(v^2 - 1), v^{2n})\$.

Let \$n\$ and \$l\$ be positive integers. We denote an \$n\$-tuple \$(r\_1, r\_2, \dots, r\_n)\$ of integers \$r\_1, r\_2, \dots, r\_n\$ by \$\langle r \rangle\_n\$. We denote by \$W\_{(n,l,\langle r \rangle\_n)}\$ the \$N\_1\$-tangle obtained from \$W\_n\$ by putting \$X\_0 = E\_{2l}^\perp\$ and \$Y\_j = E\_{2r\_j}\$, \$1 \le j \le n\$.

By Proposition 3.6, Lemmas 3.4 and 3.7, and Corollary 3.8, we have the following.

**Lemma 3.9.**

$$\begin{aligned} e_0(W_{(n,l,\langle r \rangle_n)}; v, z) \\ = (-1)^{n+1} v^{2(\sum_{j=1}^n r_j - 2n)} (v^{2l} - 1) (1 - v^2)^{2n-1} z^{2n} (1 + z^2)^{2n} (2 + z^2)^n. \end{aligned}$$

Let \$T\$ be a properly oriented 2-string tangle and \$c\$ a crossing on \$T\$. We denote by \$S\_c T\$ the tangle obtained from \$T\$ by switching the crossing \$c\$. We also denote by \$Z\_c T\$ the tangle obtained from \$T\$ by smoothing the crossing \$c\$.

Let \$H\_q\$, \$q \ge 3\$, be the \$N\_1\$-tangle depicted in Fig. 8.

Note that \$N(H\_q)\$ is the trivial knot. We also see that \$N(S\_{c\_1} S\_{c\_2} H\_q)\$ and \$N(S\_{c\_3} H\_q)\$ are trivial knots because \$S\_{c\_1} S\_{c\_2} H\_q = E\_\infty\$ and \$S\_{c\_3} H\_q = E\_\infty\$.

Let  $K$  be an oriented knot and  $T_K$  a  $D_1$ -tangle with  $D(T_K) = K$ . For a positive integer  $q$ ,  $q \geq 2$ , we denote the  $N_1$ -tangle  $R(H_{q+1}^\perp, T_K)$  by  $\mathcal{Q}_{(K,q)}$ . Note that  $N(\mathcal{Q}_{(K,q)}) = D(T) = K$ . We also denote a knot  $N(\mathcal{Q}_{(K,q)}^\perp + W_{(n,l,(r)_n)})$  by  $K[n, l, \langle r \rangle_n, q]$ , where  $W_{(n,l,(r)_n)}$  denotes the  $N_1$ -tangle introduced after Corollary 3.8.

**Proposition 3.10.**

$$P(K[n, l, \langle r \rangle_n, q]; v, z) - P(K; v, z) = \lambda(v, z)e_0(R(E_\infty, E_0); v, z)e_0(H_{q+1}; v, z)e_0(T_K; v, z)e_0(W_{(n,l,(r)_n)}; v, z).$$

Proof. By Proposition 2.8, we have

$$\begin{aligned} P(K[n, l, \langle r \rangle_n, q]; v, z) &= P(D(\mathcal{Q}_{(K,q)}^\perp); v, z)P(N(W_{(n,l,(r)_n)}); v, z) \\ &\quad + \lambda(v, z)e_\infty(\mathcal{Q}_{(K,q)}^\perp; v, z)e_0(W_{(n,l,(r)_n)}; v, z) \\ &= P(N(\mathcal{Q}_{(K,q)}); v, z)P(N(W_{(n,l,(r)_n)}); v, z) \\ &\quad + \lambda(v, z)e_0(\mathcal{Q}_{(K,q)}; v, z)e_0(W_{(n,l,(r)_n)}; v, z). \end{aligned}$$

Since  $P(N(\mathcal{Q}_{(K,q)}); v, z) = P(K; v, z)$ ,  $P(N(W_{(n,l,(r)_n)}); v, z) = P(N(E_{\frac{1}{2n}}^\perp); v, z) = P(U; v, z) = 1$ , and  $e_0(\mathcal{Q}_{(K,q)}; v, z) = e_0(H_{q+1}; v, z)e_0(T_K; v, z)e_0(R(E_\infty, E_0); v, z)$  from Proposition 3.6, we obtain the result.  $\square$

**Lemma 3.11.**  $e_0(T_K; v, z) \neq 0$ .

Proof. By Lemma 2.4 and  $D(T_K) = K$ , we obtain

$$e_0(T_K; v, z) = \frac{1}{\lambda(v, z)}\{-v(1 - v^2)zP(N(T_K); v, z) + v^2z^2P(K; v, z)\}.$$

It follows that  $e_0(T_K; 1, z) = \nabla(K; z)$ . Since  $K$  is a knot, we have  $\nabla(K; z) \neq 0$ , and thus,  $e_0(T_K; v, z) \neq 0$ .  $\square$

**Lemma 3.12.** For  $q \geq 3$ ,  $\nabla(D(H_q); z) = (-1)^q z^{2q-1}$ .

Proof. The proof is by an induction on the number  $q$ . If  $q = 3$ , then we see that  $\nabla(D(H_3); z) = -z^5$  by a direct calculation. Suppose that  $q > 3$ . Let  $c_3$  and  $c_4$  be crossings in  $H_q$  as in Fig. 8. From recursive formulas for skein triples  $(D(S_{c_3}H_q) = U_2, D(H_q), D(Z_{c_3}H_q))$  and  $(D(S_{c_4}Z_{c_3}H_q), D(Z_{c_3}H_q), D(Z_{c_4}Z_{c_3}H_q) = U_2)$ , we have  $\nabla(D(H_q); z) = -z\nabla(D(S_{c_4}Z_{c_3}H_q); z)$  because  $\nabla(U_2; z) = 0$ . Since  $D(S_{c_4}Z_{c_3}H_q)$  is the connected sum of  $D(H_q)$  and a diagram of the positive Hopf link, we obtain  $\nabla(D(H_q); z) = -z^2\nabla(D(H_{q-1}))$ . By the inductive hypothesis, we have the claim.  $\square$



**Lemma 3.13.** For  $q \geq 3$ ,  $e_0(H_q; v, z) \neq 0$ .

Proof. By Lemma 2.5, we have

$$e_0(H_q; v, z) = \frac{1}{\lambda(v, z)} \{-(1 - v^2)P(N(H_q); v, z) + vzP(D(H_q); v, z)\},$$

and thus,  $e_0(H_q; 1, z) = z^{-1}P(D(H_q); 1, z) = z^{-1}\nabla(D(H_q); z)$ . Lemma 3.12 completes the proof.  $\square$

**4. Proof of Theorem 1.2**

For a 2-variable polynomial  $h \in \mathbb{Z}[v^{\pm 1}, z^{\pm 1}]$ , we denote the minimal degree of  $h$  in  $z$  by  $\min \deg_z h$ .

Let  $K$  be an oriented knot. Let  $m, l$  and  $q$  be integers with  $m \geq 0, l > 0$  and  $q > 1$ . Let  $\langle r \rangle_m \in \mathbb{Z}^m$ .

**Proposition 4.1.**  $K[m, l, \langle r \rangle_m, q]$  is distinct from  $K$  and the HOMFLY polynomial of  $K[m, l, \langle r \rangle_m, q]$  is a fake HOMFLY polynomial of  $K$  with identical order  $2m$ .

Proof. Recall that  $\lambda(v, z) \neq 0$ . Proposition 3.10 and Lemmas 3.4, 3.9, 3.11 and 3.13 show that  $P(K[m, l, \langle r \rangle_m, q]; v, z) \neq P(K; v, z)$ . It follows that  $K[m, l, \langle r \rangle_m, q]$  is distinct from  $K$ . From the definitions of  $\lambda(v, z)$  and the normal coordinates of a tangle, we have

$$\min \deg_z \lambda(v, z) = 0, \quad \min \deg_z e_0(H_q; v, z) \geq 0 \quad \text{and} \quad \min \deg_z e_0(T_K; v, z) \geq 0.$$

By Lemmas 3.4 and 3.9, we obtain

$$\min \deg_z e_0(R(E_\infty, E_0); v, z) = 2 \quad \text{and} \quad \min \deg_z e_0(W_{(m, l, \langle r \rangle_m)}; v, z) \geq 2m.$$

From Proposition 3.10, it follows that

$$\min \deg_z (P(K[m, l, \langle r \rangle_m, q]; v, z) - P(K; v, z)) \geq 2m + 2,$$

completing the proof.  $\square$

**Lemma 4.2.**  $P(K[m, l, \langle r \rangle_m, q]; v, z) \equiv P(K; v, z) \pmod{(v^{2l} - 1)}$ .

Proof. Since  $e_0(W_{(m, l, \langle r \rangle_m)}; v, z) \equiv 0 \pmod{(v^{2l} - 1)}$  from Lemma 3.9, the claim is given by Proposition 3.10.  $\square$

**Lemma 4.3.**  $\nabla(K[m, l, \langle r \rangle_m, q]; z) = \nabla(K; z)$ .

Proof. By Lemma 4.2, we have

$$\nabla(K[m, l, \langle r \rangle_m, q]; z) = P(K[m, l, \langle r \rangle_m, q]; 1, z) = P(K; 1, z) = \nabla(K; z). \quad \square$$

**Lemma 4.4.**  $d_G(K[m, l, \langle r \rangle_m, q], K) = 1$ .

Proof. Since the tangle  $H_{q+1}$  can be changed into  $E_\infty$  by switching the crossing  $c_3$  in  $H_{q+1}$  indicated in Fig. 8, the tangle  $R(H_{q+1}^\perp, T_K)$  becomes the tangle  $R(E_0, T_K)$  by a single crossing change. Then, the knot  $K[m, l, \langle r \rangle_m, q]$  which is equal to  $N(Q_{(K,q)}^\perp + W_{(m,l,\langle r \rangle_m)}) = N(R(H_{q+1}^\perp, T_K)^\perp + W_{(m,l,\langle r \rangle_m)})$  becomes a knot  $N(E_0, T_K)^\perp + W_{(m,l,\langle r \rangle_m)} = D(T_K) \# N(W_{(m,l,\langle r \rangle_m)}) = K \# N(W_{(m,l,\langle r \rangle_m)})$ . Since  $N(W_{(m,l,\langle r \rangle_m)}) = N(E_{2l}^\perp) = D(E_{2l}) = U$ , we have the result.  $\square$

**Lemma 4.5.**  $K[m, l, \langle r \rangle_m, q]$  is a band sum of  $K$  and the trivial knot.

Proof. Since  $N(W_{(m,l,\langle r \rangle_m)}) = U$  and  $N(H_{q+1}) = U$ ,  $K[m, l, \langle r \rangle_m, q]$  can be changed into  $K \sqcup U$  by a hyperbolic transformation along the band  $B$  in the tangle  $Q_{(K,q)}$  as in Fig. 7. This completes the proof.  $\square$

**Lemma 4.6.**  $K[m, l, \langle r \rangle_m, q]$  can be changed into  $K$  by a single pass-move.

Proof. Let  $a_1, a_2, a_3$  and  $a_4$  be crossings in the tangle  $Q_{(K,q)}$  depicted in Fig. 7. If the four crossings are switched simultaneously, then  $K[m, l, \langle r \rangle_m, q]$  is changed into the connected sum  $N(W_{(m,l,\langle r \rangle_m)}) \# D(T_K) \# N(H_{q+1})$ , which is equivalent to  $K$  because  $N(W_{(m,l,\langle r \rangle_m)})$  and  $N(H_{q+1})$  are trivial knots. Since such an operation is a pass-move, the claim is true.  $\square$

**Lemma 4.7.** Let  $f$  be a finite type invariant with order less than  $q + 1$ . Then,  $f(K[m, l, \langle r \rangle_m, q]) = f(K)$ .

Proof. Let  $c_1$  and  $c_2$  be crossings in  $H_{q+1}$  as shown in Fig. 8. Since  $S_{c_1} S_{c_2} H_{q+1} = E_\infty$ , we see that  $S_{c_1} S_{c_2} K[m, l, \langle r \rangle_m, q] = K$ . Since switching the crossings  $c_1$  and  $c_2$  can be realized by applying a  $C_{q+1}$ -move [5] in  $H_{q+1}$ ,  $K$  and  $K[m, l, \langle r \rangle_m, q]$  are  $C_{q+1}$ -equivalent. Thus, by [4, 6] we have the result.  $\square$

We are now ready to give the proof of Theorem 1.2.

Proof of Theorem 1.2. For distinct integers  $s$  and  $t$ , let  $\langle r \rangle_m = (s, 0, \dots, 0)$  and  $\langle r' \rangle_m = (t, 0, \dots, 0)$ , where  $\langle r \rangle_m, \langle r' \rangle_m \in \mathbb{Z}^m$ . Then, by Lemma 3.9, we obtain

$$\begin{aligned} & e_0(W_{(m,l,\langle r \rangle_m)}; v, z) - e_0(W_{(m,l,\langle r' \rangle_m)}; v, z) \\ &= (-1)^{m+1} v^{-4m} (v^{2s} - v^{2t})(v^{2l} - 1)(1 - v^2)^{2m-1} z^{2m} (1 + z^2)^{2m} (2 + z^2)^m \neq 0. \end{aligned}$$

Thus, by Proposition 3.10 and Lemmas 3.4, 3.11 and 3.13, we have

$$\begin{aligned} & \frac{P(K[m, l, \langle r \rangle_m, q]; v, z) - P(K[m, l, \langle r' \rangle_m, q]; v, z)}{\lambda(v, z)e_0(R(E_\infty, E_0); v, z)e_0(H_{q+1}; v, z)e_0(T_K; v, z)} \\ &= e_0(W_{(m,l,\langle r \rangle_m)}; v, z) - e_0(W_{(m,l,\langle r' \rangle_m)}; v, z) \neq 0. \end{aligned}$$

Let  $K_n = K[m, l, \langle r \rangle_m, q]$ , where  $n \in \mathbb{N}$  and  $\langle r \rangle_m = (n, 0, \dots, 0) \in \mathbb{Z}^m$ . Then,  $K_i$  and  $K_j$ ,  $i \neq j$ , are distinct because of the above argument. Proposition 4.1 and Lemmas 4.2–4.7 show that the knots  $\{K_n; n \in \mathbb{N}\}$  are desired ones. □

For  $\langle r \rangle_m \in \mathbb{Z}^m$ , we denote  $\sum_{j=1}^m r_j$  by  $\|r\|_m$ . Then, we have the following.

**Lemma 4.8.**  $P(K[m, l, \langle r \rangle_m, q]; v, z) = P(K[m, l, \langle r' \rangle_m, q]; v, z)$  if  $\|r\|_m = \|r'\|_m$ .

Proof. By Lemma 3.9 and the assumption of the lemma, we find that

$$e_0(W_{(m,l,\langle r \rangle_m)}; v, z) - e_0(W_{(m,l,\langle r' \rangle_m)}; v, z) = 0.$$

From Proposition 3.10, we obtain the result. □

**Conjecture 4.9.**  $K[m, l, \langle r \rangle_m, q]$  and  $K[m, l, \langle r' \rangle_m, q]$  are distinct even though  $\|r\|_m = \|r'\|_m$ .

REMARK 4.10. Almost identical link imitations which give a solution of Theorem 1.2 have the same HOMFLY polynomial.

**Proposition 4.11.** For an oriented knot  $K$ , there exist infinitely many knots with the same reduced HOMFLY polynomial at  $z = \sqrt{-2}$  as  $K$ . In particular, there exist infinitely many knots  $\{K_n; n \in \mathbb{N}\}$  with  $P(K_n; v, \sqrt{-2}) = 1$ .

Proof. From Lemma 3.9 and Proposition 3.10, it follows that knots in Theorem 1.2 have the same reduced HOMFLY polynomial at  $z = \sqrt{-2}$  as  $K$ . □

**Proposition 4.12.** Let  $K$  be an oriented knot. Let  $p$  be a positive integer and  $\xi$  a primitive  $2p$ -th root of unity. Then, there exist infinitely many knots  $\{K_n; n \in \mathbb{N}\}$  with  $V(K_n; \xi) = V(K; \xi)$ .

Proof. We show that knots in Theorem 1.2 have the same value of the Jones polynomial at  $t = \xi$  as  $K$ . From the second property in Theorem 1.2, there exists

a polynomial  $g(v, z) \in \mathbb{Z}[v^{\pm 1}, z^{\pm 1}]$  so that  $P(K_n; v, z) - P(K; v, z) = (v^{2p} - 1)g(v, z)$ . Thus, we have

$$\begin{aligned} V(K_n; \xi) - V(K; \xi) &= P(K_n; \xi, \xi^{1/2} - \xi^{-1/2}) - P(K; \xi, \xi^{1/2} - \xi^{-1/2}) \\ &= (\xi^{2p} - 1)g(\xi, \xi^{1/2} - \xi^{-1/2}) \\ &= 0, \end{aligned}$$

completing the proof. □

**5. Preliminaries for the proof of Theorem 1.5**

Let  $X$  be a properly oriented 2-string tangle and  $R(X)$  the properly oriented 2-string tangle illustrated in Fig. 9.

Note that  $N(R(X)) = D(X)$ , and that  $R(X)$  is an  $N_1$ -tangle if  $X$  is a  $D_1$ -tangle. The following lemma is easily obtained.

**Lemma 5.1.**  $R(E_0) = E_\infty$ .

It is easy to see that  $N(R(E_\infty))$  is the trivial 2-component link. Since it is shown that  $D(R(E_\infty))$  is the link  $9^3_{18}$  [14] with appropriate orientations, we have the following.

**Lemma 5.2.**  $P(N(R(E_\infty)); v, z) = (v^{-1} - v)z^{-1}$  and  $P(D(R(E_\infty)); v, z) = (v^{-2} - 2 + v^2)z^{-2} + (-v^{-4} + 3v^{-2} - 3 + v^2) + (-v^{-4} + 3v^{-2} - 2)z^2 + v^{-2}z^4$ .

Let  $\alpha(v, z) = v^{-2}z^2(1 - v^2 + z^2)$ . Since  $R(E_\infty)$  is a tangle of type  $N_2$ , Lemma 2.5 gives the following.

**Lemma 5.3.**  $(e_0(R(E_\infty); v, z), e_\infty(R(E_\infty); v, z)) = (\alpha(v, z), (1 - v^2)(1 - \alpha(v, z)))$ .

**Lemma 5.4.** *Let  $X$  be a  $D_1$ -tangle and  $(e_0(X; v, z), e_\infty(X; v, z))$  the normal coordinates of  $X$ . Then, the normal coordinates  $(e_0(R(X); v, z), e_\infty(R(X); v, z))$  of the tangle  $R(X)$  are expressed as follows:*

$$e_0(R(X); v, z) = e_\infty(X; v, z)e_0(R(E_\infty); v, z)$$

and

$$e_\infty(R(X); v, z) = e_0(X; v, z) + e_\infty(X; v, z)e_\infty(R(E_\infty); v, z).$$

*Proof.* Since  $X$  is a tangle of type  $D_1$ , by using the normal coordinates of  $X$ , we have

$$P(L(R(X); v, z) = e_0(X; v, z)P(L(R(E_0)); v, z) + vze_\infty(X; v, z)P(L(R(E_\infty)); v, z).$$




Fig. 9. Tangle  $R(X)$ .

Lemma 5.1 shows that  $P(L(R(E_0))); v, z = P(L(E_\infty); v, z)$ . Since  $R(E_\infty)$  is an  $N_2$ -tangle, Lemma 2.2 gives

$$P(L(R(E_\infty))); v, z = e_0(R(E_\infty); v, z)P(L(E_0); v, z) + v^{-1}z^{-1}e_\infty(R(E_\infty); v, z)P(L(E_\infty); v, z).$$

Thus, we obtain

$$P(L(R(X))); v, z = vze_\infty(X; v, z)e_0(R(E_\infty); v, z)P(L(E_0); v, z) + \{e_0(X; v, z) + e_\infty(X; v, z)e_\infty(R(E_\infty); v, z)\}P(L(E_\infty); v, z).$$

Since  $R(X)$  is an  $N_1$ -tangle, we have the desired formulas. □

Let  $q, q \geq 2$ , be an integer. We define a sequence  $\{W_{(n,q)}; n \in \mathbb{N}\}$  of  $N_1$ -tangles by the following recursive formulas:

- (1)  $W_{(1,q)} = R(H_{q+1}^\perp)$ ;
- (2)  $W_{(n,q)} = R(W_{(n-1,q)}^\perp), n \geq 2$ .

REMARK 5.5.  $N(W_{(n,q)}) = D(H_{q+1}^\perp) = N(H_{q+1}) = U$ .

**Proposition 5.6.** For any positive integer  $n$ ,

$$e_0(W_{(n,q)}; v, z) = e_0(H_{q+1}; v, z)e_0(R(E_\infty); v, z)^n.$$

Proof. The proof is by an induction on the number  $n$ . If  $n = 1$ , then Lemmas 3.5 and 5.4 show that

$$e_0(W_{(1,q)}; v, z) = e_\infty(H_{q+1}^\perp; v, z)e_0(R(E_\infty); v, z) = e_0(H_{q+1}; v, z)e_0(R(E_\infty); v, z).$$

Thus, the claim is true. Suppose that  $n > 1$ . By the inductive hypothesis and Lemmas 3.5 and 5.4, we have

$$\begin{aligned} e_0(W_{(n,q)}; v, z) &= e_\infty(W_{(n-1,q)}^\perp; v, z)e_0(R(E_\infty); v, z) \\ &= e_0(W_{(n-1,q)}; v, z)e_0(R(E_\infty); v, z) \\ &= e_0(H_{q+1}; v, z)e_0(R(E_\infty); v, z)^n. \end{aligned}$$

This completes the proof. □

Let  $K$  be an oriented knot and  $T_K$  a  $D_1$ -tangle with  $D(T_K) = K$ .

**Lemma 5.7.** *The normal coordinates of the tangle  $T_K + E_{2l}$ ,  $l \in \mathbb{Z}$ , are*

$$\begin{aligned} &(e_0(T_K + E_{2l}; v, z), e_\infty(T_K + E_{2l}; v, z)) \\ &= \left( v^{2l}e_0(T_K; v, z), \frac{v^{2l} - 1}{v^2 - 1}e_0(T_K; v, z) + e_\infty(T_K; v, z) \right). \end{aligned}$$

Proof. Since Lemma 3.7 gives  $e_0(E_{2l}; v, z) + (1 - v^2)e_\infty(E_{2l}; v, z) = 1$ , the lemma immediately comes from Lemma 2.10. □

We denote a knot  $N((T_K + E_{2l}) + W_{(n,q)})$  by  $K[n, q, l]$ .

**Proposition 5.8.**

$$\begin{aligned} &P(K[n, q, l]; v, z) - P(K; v, z) \\ &= \lambda(v, z)e_0(H_{q+1}; v, z)e_0(R(E_\infty); v, z)^n \left( \frac{v^{2l} - 1}{v^2 - 1}e_0(T_K; v, z) + e_\infty(T_K; v, z) \right). \end{aligned}$$

Proof. By Proposition 2.8, we have

$$\begin{aligned} P(K[n, q, l]; v, z) &= P(D(T_K + E_{2l}); v, z)P(N(W_{(n,q)}); v, z) \\ &\quad + \lambda(v, z)e_\infty(T_K + E_{2l}; v, z)e_0(W_{(n,q)}; v, z). \end{aligned}$$

Since  $D(T_K + E_{2l}) = D(T_K) = K$  and  $N(W_{(n,q)}) = U$ , the first term on the right-hand side of the above equality is equal to  $P(K; v, z)$ . Proposition 5.6 and Lemma 5.7 lead to the claim. □

**Corollary 5.9.**

$$\nabla(K[n, q, l]; z) = \nabla(K; z) + (-1)^{q+1}z^{4n+2q+2}(l\nabla(K; z) + e_\infty(T_K; 1, z)).$$

Proof. Note that

$$e_0(T_K; 1, z) = P(K; 1, z) = \nabla(K; z)$$

and

$$e_0(H_{q+1}; 1, z) = z^{-1}P(D(H_{q+1}); 1, z) = z^{-1}\nabla(D(H_{q+1}); z).$$

By Lemma 5.3, we have the result. □

From the definition of the normal coordinates of a tangle,  $e_\infty(T_K; 1, 2\sqrt{-1})$  is an integer.

**Lemma 5.10.**

$$\frac{\nabla(K[n, q, l]; 2\sqrt{-1})}{\nabla(K; 2\sqrt{-1})} > 1$$

if  $l > |e_\infty(T_K; 1, 2\sqrt{-1})/\nabla(K; 2\sqrt{-1})|$ .

Proof. Note that  $n \geq 1$  and  $q \geq 2$ . By Corollary 5.9, we obtain

$$\frac{\nabla(K[n, q, l]; 2\sqrt{-1})}{\nabla(K; 2\sqrt{-1})} = 1 + 4^{2n+q+1} \left( l + \frac{e_\infty(T_K; 1, 2\sqrt{-1})}{\nabla(K; 2\sqrt{-1})} \right).$$

Since  $|e_\infty(T_K; 1, 2\sqrt{-1})/\nabla(K; 2\sqrt{-1})| + e_\infty(T_K; 1, 2\sqrt{-1})/\nabla(K; 2\sqrt{-1}) \geq 0$ , we have the desired inequality. □

**6. Proof of Theorem 1.5**

Let  $K$  be an oriented knot and  $T_K$  a  $D_1$ -tangle with  $D(T_K) = K$ . Let  $m, l$  and  $q$  be integers with  $m \geq 0$ ,  $l > l_0 = |e_\infty(T_K; 1, 2\sqrt{-1})/\nabla(K; 2\sqrt{-1})|$  and  $q > 1$ .

The following is an immediate consequence of Lemma 5.10.

**Lemma 6.1.**  $\nabla(K[m + 1, q, l]; z) \neq \nabla(K; z)$ , that is,  $K[m + 1, q, l]$  is distinct from  $K$ .

**Proposition 6.2.** *The HOMFLY polynomial of  $K[m + 1, q, l]$  is a fake HOMFLY polynomial of  $K$  with identical order  $2m$ .*

Proof. Note that Lemma 6.1 shows  $P(K[m + 1, q, l]; v, z) \neq P(K; v, z)$ . From the definitions of  $\lambda(v, z)$  and the normal coordinates of a tangle, we obtain four inequalities:  $\min \deg_z \lambda(v, z) = 0$ ,  $\min \deg_z e_0(H_{q+1}; v, z) \geq 0$ ,  $\min \deg_z e_0(T_K; v, z) \geq 0$  and  $\min \deg_z e_\infty(T_K; v, z) \geq 0$ . By Lemma 5.3, we have  $\min \deg_z e_0(R(E_\infty); v, z) = 2$ . From Proposition 5.8, it follows that

$$\min \deg_z (P(K[m + 1, q, l]; v, z) - P(K; v, z)) \geq 2m + 2,$$

completing the proof. □

**Lemma 6.3.**  $d_G(K[m + 1, q, l], K) = 1$ .

Proof. Since the tangle  $H_{q+1}$  can be changed into  $E_\infty$  by switching the crossing  $c_3$  in  $H_{q+1}$  indicated in Fig. 8, the tangle  $R(H_{q+1}^\perp)$  becomes  $R(E_0) = E_\infty$  by a single crossing change. Hence, the tangle  $W_{(m+1,q)}$  is changed into  $E_\infty$  by a single crossing change, and thus,  $K[m + 1, q, l]$  becomes  $N(T_K + E_{2l} + E_\infty) = D(T_K + E_{2l}) = D(K) = K$ . □

The value  $\nabla(K; 2\sqrt{-1})$ , whose absolute value is equal to the determinant of  $K$ , is an integer. By [3], it is known that the signature  $\sigma(K)$  of  $K$  has the following properties:

- (1)  $\nabla(K; 2\sqrt{-1})/|\nabla(K; 2\sqrt{-1})| = \sqrt{-1}^{\sigma(K)}$ ;
- (2)  $|\sigma(K) - \sigma(K')| \leq 2$ ,

where  $K'$  is a knot obtained from  $K$  by switching a crossing of  $K$ .

**Lemma 6.4.**  $\sigma(K[m + 1, q, l]) = \sigma(K)$ .

Proof. Since two integers  $\nabla(K[m + 1, q, l]; 2\sqrt{-1})$  and  $\nabla(K; 2\sqrt{-1})$  have the same signature by Lemma 5.10, we have  $\sigma(K[m + 1, q, l]) \equiv \sigma(K) \pmod{4}$ . Since  $d_G(K[m + 1, q, l], K) = 1$  by Lemma 6.3, we obtain  $|\sigma(K[m + 1, q, l]) - \sigma(K)| \leq 2$ . These two relations give the claim. □

**Lemma 6.5.**  $K[m + 1, q, l]$  can be changed into  $K$  by a single pass-move.

Proof. We consider the 3-string tangle in the disk  $F$  bounded by dotted segments depicted in the tangle  $H_{q+1}$  as in Fig. 8. If we apply a  $\Gamma$ -move [8] to the 3-string tangle, then we find that the tangle  $H_{q+1}$  can be changed into the tangle  $E_\infty$ . Thus,  $K[m + 1, q, l]$  can be changed into  $K$  by a single  $\Gamma$ -move as the proof of Lemma 6.3. Since a  $\Gamma$ -move can be accomplished by a combination of Reidemeister moves and a pass-move, we complete the proof. □

**Lemma 6.6.** Let  $f$  be a finite type invariant with order less than  $q + 1$ . Then,  $f(K[m + 1, q, l]) = f(K)$ .

Proof. The proof is similar to that of Lemma 4.7. □

Proof of Theorem 1.5. For two integers  $l$  and  $l'$  with  $l' > l > l_0$ , by Corollary 5.9, we obtain

$$\begin{aligned} \nabla(K[m + 1, q, l']; z) - \nabla(K[m + 1, q, l]; z) &= (-1)^{q+1} z^{4(m+1)+2q+2} \nabla(K; z)(l' - l) \\ &\neq 0. \end{aligned}$$



Hence,  $K[m + 1, q, l']$  and  $K[m + 1, q, l]$  are distinct. Proposition 6.2 and Lemmas 6.1 and 6.3–6.6 show that knots  $\{K[m + 1, q, l]; l_0 < l \in \mathbb{N}\}$  are desired ones.  $\square$

**Proposition 6.7.** *For an oriented knot  $K$ , there exist infinitely many knots with the same reduced HOMFLY polynomial at  $z = \sqrt{v^2 - 1}$  as  $K$ . In particular, there exist infinitely many knots  $\{K_n; n \in \mathbb{N}\}$  with  $P(K_n; v, \sqrt{v^2 - 1}) = 1$ .*

Proof. From Lemma 5.3 and Proposition 5.8, it follows that the knots  $K_n$  in Theorem 1.5 have the same reduced HOMFLY polynomial at  $z = \sqrt{v^2 - 1}$  as  $K$ .  $\square$

---

### References

- [1] J.H. Conway: *An enumeration of knots and links*, in Computational Problems in Abstract Algebra, Pergamon Press, New York, 1969, 329–358.
- [2] P. Freyd, D. Yetter, J. Hoste, W.B.R. Lickorish, K.C. Millett and A. Ocneanu: *A new polynomial invariant of knots and links*, Bull. Amer. Math. Soc. (N.S.) **12** (1985), 239–246.
- [3] C.A. Giller: *A family of links and the Conway calculus*, Trans. Amer. Math. Soc. **270** (1982), 75–109.
- [4] M.N. Goussarov: *Knotted graphs and a geometrical technique of  $n$ -equivalence*, POMI Sankt Petersburg preprint, circa 1995, (Russian).
- [5] K. Habiro: Master Thesis, University of Tokyo, 1994.
- [6] K. Habiro: *Claspers and finite type invariants of links*, Geom. Topol. **4** (2000), 1–83.
- [7] V.F.R. Jones: *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math. (2) **126** (1987), 335–388.
- [8] L.H. Kauffman: *On Knots*, Annals of Mathematics Studies **115**, Princeton Univ. Press, Princeton, NJ, 1987.
- [9] A. Kawachi: *Almost identical link imitations and the skein polynomial*; in Knots 90 (Osaka, 1990), de Gruyter, Berlin, 1992, 465–476.
- [10] W.B.R. Lickorish and K.C. Millett: *A polynomial invariant of oriented links*, Topology **26** (1987), 107–141.
- [11] Y. Miyazawa: *Knots with a trivial coefficient polynomial*, Kyungpook Math. J. **49** (2009), 801–809.
- [12] K. Murasugi: *On a certain numerical invariant of link types*, Trans. Amer. Math. Soc. **117** (1965), 387–422.
- [13] J.H. Przytycki and P. Traczyk: *Invariants of links of Conway type*, Kobe J. Math. **4** (1987), 115–139.
- [14] D. Rolfsen: *Knots and Links*, Publish or Perish, Berkeley, CA, 1976.

Department of Mathematical Sciences  
 Yamaguchi University  
 Yamaguchi 753-8512  
 Japan  
 e-mail: miyazawa@yamaguchi-u.ac.jp