

| Title | Strategy for enhancement of lactic-acid resistance of Saccharomyces cerevisiae to produce lactic acid without neutralization |
|--------------|--|
| Author(s) | Suzuki, Toshihiro |
| Citation | 大阪大学, 2013, 博士論文 |
| Version Type | VoR |
| URL | https://doi.org/10.18910/26844 |
| rights | |
| Note | |

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Doctoral Dissertation

Strategy for enhancement of lactic-acid resistance of Saccharomyces cerevisiae to produce lactic acid without neutralization

Toshihiro Suzuki

February 2013

Department of Biotechnology, Graduate School of Engineering

Osaka University

Doctoral Dissertation

Strategy for enhancement of lactic-acid resistance of Saccharomyces cerevisiae to produce lactic acid without neutralization

Toshihiro Suzuki

February 2013

Department of Biotechnology, Graduate School of Engineering

Osaka University

Table of contents

Subject

٠

| P | age |
|--|-----|
| Chapter 1 General Introduction | 1 |
| 1.1 Lactic acid production | 1 |
| 1.2 Poly-lactic acid and a novel method of lactic-acid production | 2 |
| 1.3 Effect of organic acid on cell mechanisms | 5 |
| 1.4 Response to weak-organic acid | 6 |
| 1.5 Objective of the study | 8 |
| Chapter 2 Disruption of multiple genes whose deletion causes lactic-acid | |
| resistance improves lactic-acid resistance and productivity in | |
| S. cerevisiae | 10 |
| 2.1 Introduction | 10 |
| 2.2 Materials and Methods | 12 |
| 2.2.1 Strains and media | 12 |
| 2.2.2 Construction of plasmid vectors | 14 |
| 2.2.3 Construction of single and multiple disruptants | 15 |
| 2.2.4 Screening for L-lactic-acid-resistant disruptants | 16 |
| 2.2.5 Spot dilution growth assay | 17 |
| 2.2.6 Measurement of lactic acid production | 17 |
| 2.3 Results | 18 |
| 2.3.1 Genome-wide screening for S. cerevisiae L-lactic-acid-resistant | |
| disruptants | 18 |
| 2.3.2 Multiple disruptants show superior lactic-acid resistance | 23 |
| 2.3.3 A lactic-acid-resistant strain that produces lactic acid more | |
| efficiently than wild type | 27 |
| 2.4 Discussion | 31 |
| 2.5 Summary | 33 |
| Chapter 3 Lactic-acid stress causes vacuolar fragmentation and impairs | |
| intracellular amino-acid homeostasis in S. cerevisiae | 35 |
| 3.1 Introduction | 35 |

-

| 3.2 | Ma | terials and Methods | 36 |
|-------------------------|--------------------|--|----|
| 3.2 | 2.1 | Strains, plasmids, and media | 36 |
| 3.2 | 2.2 | Screening for L-lactic acid hypersensitive mutants | 37 |
| 3.2 | 2.3 | Spot dilution growth assays | 37 |
| 3.2 | 2.4 | Active staining of the vacuole with quinacrine | 38 |
| 3.2 | 2.5 | Labeling the vacuolar membrane with FM4-64 | 38 |
| 3.2 | 2.6 | Quantification of intracellular amino acids | 39 |
| 3.2 | 2.7 | Cell viability | 40 |
| 3.3 | Res | sults | 40 |
| 3.3 | 3.1 | A genome-wide screening for lactic-acid-hypersensitive alleles | 40 |
| 3.3 | 3.2 | Almost half of the lactic-acid-sensitive disruptants show vacuolar | |
| | | acidification defect. | 48 |
| 3.3 | 3.3 | Vacuole is subjected to fragmentation by lactic-acid stress | 51 |
| 3. | 3.4 | Amount of intracellular amino acid is important for the growth | |
| | | under lactic-acid stress | 54 |
| 3.3 | 3.5 | Amino acid supplementation recovers adaptation deficiency to | |
| | | lactic acid | 57 |
| 3.4 | Dis | cussion | 59 |
| 3.5 | Sui | nmary | 62 |
| Chapte | er 4 | General discussion and conclusion | 64 |
| Refere | nces | | 71 |
| List of Publications 88 | | | 88 |
| Ackno | Acknowledgements 8 | | 89 |

.

Chapter 1

General Introduction

1.1 Lactic acid production

Lactic acid (2-hydroxypropionic acid, CH₃CHOHCOOH) is widely used in foods (acidulants, preservatives, flavors, pH regulators, etc.), cosmetics (moisturizers, skin-lightening agents, etc.), pharmaceutical agents (mineral preparations, dialysis solution, etc.), and industrial chemical products (descaling agents, chiral intermediates, etc.). Lactic acid is also one of the major metabolic intermediates in most organisms from prokaryotes to human (Vickroy, 1985; Datta et al., 1995). Lactic acid was first isolated from sour milk by Scheele, C. W. in 1780 and was commercially produced in 1881 by Avery, C. E. in Littleton, Massachusetts, USA (Vickroy, 1985). Lactic acid can be produced by chemical synthesis or carbohydrate fermentation. Chemical synthesis is mainly by hydrolysis of lactonitrile using a strong acid; this method produces a racemic mixture of D- and L-lactic acid. Production of lactic acid by fermentation has several advantages compared to chemical synthesis, such as low cost of substrates, low production temperature, low energy consumption, and high product specificity, *i.e.*, it can produce the desired stereoisomer, either pure L-(+)- or D-(-)-lactic acid, in an optically pure form (Pandey et al., 2001).

1

The commercial production of lactic acid uses homolactic bacteria, such as Lactobacillus delbrueckii, L. bulgaricus, and L. leichmanii (Datta et al., 1995). Homolactic bacteria almost exclusively produce a single fermentation product, *i.e.*, lactic acid, whereas heterolactic bacteria produce a mix of products including ethanol, diacetyl, formate, acetoin, acetic acid, and carbon dioxide along with lactic acid. Homolactic- and heterolactic bacteria utilize pyruvic acid, which is the end product of Embden-Meyerhof pathway, and convert it to lactic acid. L- or D-lactate dehydrogenase (LDH) plays a key role in this conversion. Lactic-acid yields of approximately 90% (w/w) from a glucose equivalent of carbohydrate are obtained. The pH of the culture medium drops during fermentation, affecting growth of the bacteria and their productivity of L-lactic acid. To counteract this, calcium carbonate is added to the culture medium to neutralize the pH during fermentation by producing calcium lactate. The culture solution containing calcium lactate is filtered to remove cells with carbon treatment, evaporated, and , acidified with sulfuric acid to convert the calcium lactate salt into lactic acid and a precipitate of calcium sulfate, which is removed by filtration. The filtrate is further purified over carbon columns and ion exchange, and subsequently evaporated to produce technical- and food-grade lactic acid. The technical-grade lactic acid is esterified and the ester is recovered by distillation, hydrolyzed and evaporated, producing a highly pure product.

1.2 Poly-lactic acid and a novel method of lactic-acid production

The application of lactic acid to its polymer is recently desired. Poly-lactic acid is a semicrystalline polymer has high tensile strength, low elongation with high modulus, a melting point of 175-178°C, and slow degradation time. These physical properties are suitable for medical applications (orthopedic, vascular, dental, and intestinal application and sutures, etc.) (Wee et al., 2006). Furthermore. the application of this polymer to environmental-friendly alternative products is recently desired because shift of petroleum-based society to carbon-neutral society is needed to solve serious problems of global warming and escalating petroleum costs. Poly-lactic acid is being developed as a renewable alternative to conventional petroleum-based plastics (Arntzen and Dale, 1999; Chotani et al., 2000; Ohara *et al.*, 2001). Poly-lactic acid has been receiving increasing attention as a carbon neutral material that can contribute to reducing atmospheric CO₂ emissions. Although this type of plastic should be applicable globally to products such as films and chassis, among others, the relatively high production costs of producing poly-lactic-acid plastics in comparison to petroleum-based plastics such as polyethylene prevented their widespread application.

Lactic-acid bacteria can produce lactic acid in high yields of more than 0.9 g lactic acid per gram of glucose consumed (Wee *et al.*, 2006), based on a maximum theoretical yield of 1.0 g lactic acid per gram of glucose. However, the optical purity of the lactic acid produced by some lactic-acid bacteria is not 100% because some lactic-acid bacteria have both the L-*LDH* and D-*LDH* genes (Hofvendahl and Hahn-Hägerdal, 2000). The low optical purity is not suitable for the regulation of the physical properties of poly-lactic acid (Tsuji, 2002). It was also reported that

3

stereocomplex-type poly lactic acid has a high melting temperature point compared with L- or D-homopolymers (Ikada et al., 1987). However, because poly-DL-lactic acid is not crystalline and soften at a low glass-transition point (54°C) (Tsuji, 2002), it is important that poly-L-lactic acid and poly-D-lactic acid of high optical purity are blended during the synthesis of the stereocomplex polymer. In addition. because of the difficulties in cultivating lactic-acid bacteria at high density and their highly auxotrophic growth requirements (Hofvendahl and Hahn-Hagerdal, 2000), some other organisms are now being used for lactic-acid production, for example, Escherichia coli (Dien et al., 2001), Rhizopus oryzae (Bai et al., 2008), and Bacillus coagulans (Sakai and Ezaki, 2006). It has been recognized that Saccharomyces cerevisiae has advantages for lactic-acid production and new methods for the large-scale production of L-lactic acid with engineered yeast have been developed (Dequin and Barre, 1994; Porro et al., 1995). S. cerevisiae is more tolerant to low pH than lactic-acid bacteria (Skory, 2003) and can grow on inexpensive synthetic However, wild-type S. cerevisiae hardly produce lactic acid owing to the • media. conversion of pyruvate to ethanol by pyruvate decarboxylase (PDC) and alcohol dehydrogenase; however, genetically engineered yeast with an introduced exogenous L-LDH gene can produce lactic acid through the glycolytic pathway (Dequin and Barre, 1994; Porro et al., 1995; Ishida et al., 2005). Genetically engineered yeast can produce very high optical purity of L-lactic acid (at least 99.9%) (Saitoh et al., 2005). These advantages offer the potential to reduce cost because the engineered yeast can produce lactic acid without requiring the process of desalination of lactate. Although, as mentioned above, S. cerevisiae is more

tolerant to low pH than lactic-acid bacteria, the decrease in yield of L-lactic acid due to the decrease in culture pH resulting from the accumulation of lactic acid still occurs (Ishida *et al.*, 2005).

1.3 Effect of organic acid on cell mechanisms

In *S. cerevisiae*, the monocarboxylate permease Jen1 mediates the transport of lactic, pyruvic, acetic, and propionic acid (Casal *et al.*, 1999) but is repressed in the presence of glucose (Chambers *et al.*, 2003). It has been suggested that undissociated lactic-acid molecule ($pK_a = 3.86$) freely diffuse through the plasma membrane (Cássio *et al.*, 1987). Under acidic conditions, undissociated organic acids can diffuse through the plasma membrane and acidify the cytoplasm following its dissociation inside the cell (Mollapour *et al.*, 2008). The dissociated anions accumulate in the cytoplasm, their negative charge prevents them from easily crossing the plasma membrane, and such a high anion pool can potentially generate abnormally high turgor pressure (Hatzixanthis *et al.*, 2003). Cells can counteract this intracellular acidification by extruding protons through the plasma membrane ATPase, but this is performed at the expense of ATP and leads to a reduction in growth yield (Viegas *et al.*, 1998).

Organic-acid stress is not only because of the toxic effect of a high hydrogen ion concentration, but also depends on the specific chemical nature of the organic acid to which the organism is exposed (Bayrock and Ingledew, 2004). For example, acetic acid is far less inhibitory than the more lipophilic sorbic acid, even though these two carboxylate compounds have a $pK_a = 4.76$ (Piper *et al.*, 1998; Bracey *et al.*, 1998). In addition, benzoic acid inhibits nitrogen starvation-induced macroautophagy, whereas sorbic acid does not (Hazan *et al.*, 2004). Because of the effects of low extracellular pH on the cell wall, inhibition of growth by weak-organic acids has been proposed to be due to a number of actions, including membrane disruption, inhibition of essential metabolic reactions, stress on intracellular pH homeostasis, and the accumulation of toxic anions (Holyoak *et al.*, 1999).

1.4 Response to weak-organic acid

The adaptation and resistance of *S. cerevisiae* to weak-organic acids involves several mechanisms. (i) The plasma membrane H⁺-ATPase, Pam1, pumps protons out of the cell in an ATP-dependent manner (Eraso and Gancedo, 1987; Viegas and Sà-Correia, 1991). Under acidic conditions, the vacuolar ATPase also plays a role in cytosolic pH homeostasis (Kane, 2006). (ii) The transcription factors Msn2/Msn4 are involved in the general stress response (Schüller *et al.*, 2004). Most genes up-regulated by Msn2 and Msn4 in response to weak acid stress encode proteins of the environmental stress response such as molecular chaperones (*e.g.*, Hsp26, Sse2), enzymes involved in carbohydrate metabolism (*e.g.*, Hxk1, Gpd1), and antioxidant defense system (*e.g.*, Ctt1, Gpx1) (Schuller *et al.*, 2004; Simões *et al.*, 2006). (iii) War1 is required for weak acid-induced transcriptional activation of *PDR12*, which encodes a plasma membrane ABC efflux pump and is presumably involved in the efflux of benzoic and sorbic acid (Piper *et al.*, 1998; Piper *et al.*,

2001; Schüller et al., 2004). Pdr12 catalyzes the active efflux of weak organic-acid anions from the cytosol and mediates broad resistance to water-soluble weak acids, including monocarboxylic acids of aliphatic chain length from C1-7 (Holvoak et al., 1999). (iv) Extracellular undissociated acetic acid enters glucose-repressed cells primarily by facilitated diffusion through the plasma membrane aquaglyceroporin Fps1 (Mollapour and Piper, 2007). Hog1 is transiently activated by acetic-acid stress (Mollapour and Piper, 2006) and activated Hog1 directly phosphorylates Fps1 (Mollapour and Piper, 2007). This phosphorylation is the signal for Fps1 to become ubiquitinated and endocytosed into the vacuole (Mollapour and Piper, 2007). This removal of Fps1 from the plasma membrane seems to be essential for downregulating the acetic-acid influx to the cell (Mollapour and Piper, 2007). (v) Haa1, a transcription factor, is required for adaptation and resistance to low pH, particularly to the acids that are more hydrophilic, such as acetic and propionic acids (Fernandes et al., 2005). A number of genes are activated directly or indirectly by Haa1 in response to acetic acid (Mira et al., 2010). Expression of SAP30 and HRK1 gene is the most significant protective effect (Mira et al., 2010). SAP30 encodes a subunit of the Rpd3L histone deacetylase complex, whereas HRK1 encodes a protein kinase dedicated to the posttranslational regulation of plasma membrane transporters (Goossens et al., Other Haal-regulated genes that encode the drug:H⁺ antiporters Tpo2 and 2000). Tpo3 which propose to mediate acetate export (Fernandes et al., 2005) and the cell wall-related protein Ygp1 which may contribute to the acid-induced remodeling of the cell-wall structure (Fernandes et al., 2005) also confer resistance to acetic acid.

(vi) Rim101, which is involved in response to alkaline pH and cell-wall assembly is also required for resistance to benzoic, propionic, and acetic acid (Mira *et al.*, 2009). For example, Rim101 was found to regulate a small subset of the genes transcriptionally activated in response to propionic acid, including *KNH1* and *CWP1* which are involved in cell-wall integrity, and the uncharacterized gene *YIL029c* (Mira *et al.*, 2009).

1.5 Objective of the study

The fermentation of lactic acid under low pH leads to minimization of the risk of contamination (van Maris *et al.*, 2006) and elimination of the necessity for neutralization process. To increase the productivity of L-lactic acid without requiring pH control, the resistance of *S. cerevisiae* to L-lactic acid must be enhanced. To uncover the detailed mechanisms underlying the adaptation of *S. cerevisiae* to lactic-acid stress, I perform a functional genomic analysis. This study deals with the response to lactic acid among weak organic acids and the finding in this study may be applicable to the response of other weak organic acids.

In chapter 1, I describe the background information for this study: lactic-acid production and its effects on the cell. In chapter 2, I systematically screen for the yeast haploid collection for single disruptants that are resistant to 6% lactic acid, construct the lactic acid resistant strains, and then evaluate the lactic-acid productivity of these resistant strains in the absence of neutralization. Because there is currently no effective method for enhancing the lactic-acid resistance of *S. cerevisiae*, I adapt a way of the combining gene disruptions which cause lactic-acid

resistance, expecting improvement of the lactic-acid resistance. In chapter 3, I systematically screen the single disruptants that are hypersensitive to lactic acid stress and attempt to examine the intracellular mechanisms which are important for adaptation to lactic acid stress. In chapter 4, I summarize the results obtained in this study and discuss the validity of strain constructed in this study for lactic-acid production and the approach to further enhancement of lactic-acid resistance.

Chapter 2

Disruption of multiple genes whose deletion causes lactic-acid resistance improves lactic-acid resistance and productivity in *S. cerevisiae*

2.1 Introduction

While wild-type S. cerevisiae produces very little lactic acid, genetically engineered yeast harboring an exogenous L-lactate dehydrogenase gene (L-LDH) can produce lactic acid through the glycolytic pathway (Dequin and Barre, 1994; Porro et al., 1995; Ishida et al., 2005). A genetically engineered strain that has L-LDH also produces ethanol (Porro et al., 1995; Adachi et al., 1998). Because pyruvic acid is consumed to produce ethanol during fermentation, the production of ethanol will affect any improvement in lactic-acid yield. In S. cerevisiae, the , pyruvate decarboxylase genes PDC1, PDC5, and PDC6 are responsible for conversion of pyruvic acid into ethanol. Pdc1 is the main PDC that functions during glucose fermentation, whereas Pdc5 is expressed only in the absence of PDC1 (Seeboth et al., 1990) or under thiamine limitation (Muller et al., 1999). Pdc6 is not expressed during glucose fermentation and relates to the activity of pyruvate decarboxylase in an ethanol-based medium (Hohmann^a, 1991). It has been reported that disruption of PDC1 by replacement with L-LDH results in the efficient production of L-lactic acid (Ishida et al., 2005). To increase the productivity of L-lactic acid without relying on pH control, the L-lactic acid

resistance of *S. cerevisiae* must be enhanced. However, findings about the mechanisms of adaptation to lactic acid are limited. For example, it was reported that the concentration of intracellular ATP was decreased by lactic acid (Thomsson and Larsson, 2006), and *FIT2*, *ARN1* and *ARN2*, involved in metal metabolism regulated by Aft1 which is a transcription factor that responds to intracellular iron, were induced by lactic-acid stress (Kawahata *et al.*, 2006). In addition, the transcription factor Ace2 and Swi5 involved in cell cycle progression from M to G1 and target genes of transcription factor Haa1, *TPO2*, *TPO3*, *YGP1*, and *YRO2*, were highly induced by lactic acid while Haa1 was not strongly induced by lactic acid and disruptants of these target genes had no effect of growth rates in the presence of lactic acid (Abbott *et al.*, 2008).

To identify the cellular mechanisms which prevent the exhibition of lactic-acid resistance toward construction of superior lactic-acid resistant yeast, I performed functional genomic analysis. I found that 94 gene disruptants showed resistance to 6% lactic acid and that lactic-acid resistance was further enhanced by combining the disruption of several genes. I have evaluated the effect of those multiple disruptions on the production of lactic acid and found that, when those disruptions were incorporated in a strain with an exogenous bovine L-LDH gene, quadruple disruptant ($\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$) showed improved productivity of lactic acid as compared with the wild-type strain. The findings in this chapter strongly suggest that a combination of disruptions that show lactic-acid resistance is an effective approach to improve lactic-acid resistance and that this approach contributes to improvements in lactic-acid productivity without neutralization.

2.2 Materials and Methods

2.2.1 Strains and media

The nonessential genes deletion collection (Openbiosystems, Huntsville, AL, USA), which contains 4828 yeast strains with all nonessential ORFs disrupted by the *kanMX4* cassette (http://www-sequence.stanford.edu/group/yeast_deletion_project/deletions3.html), was used to screen for disruptants that showed superior lactic-acid resistance. Other strains used in this chapter are listed in Table 1. The parental strain of the deletion collection, BY4742 (*MATa leu2A0 his3A1 ura3A0 lys2A0*) (Brachmann *et al.*, 1998), was used as a wild-type strain. The standard complete medium YPDA consisted of 1% (w/v) yeast extract (Sigma-Aldrich, St. Louis, MO, USA), 2% (w/v) peptone (Sigma-Aldrich, St. Louis, MO, USA), 2% (w/v) adenine (Wako Pure Chemical Industries, Osaka). L-Lactic acid (NACALAI TESQUE, Kyoto) was added to YPDA after autoclaving. Agar (2%) was added to media to make culture plates.

| Yeast strains | Genotype | Origin |
|------------------|--|----------------------------|
| BY4742 | $MAT\alpha$ his3 $\Delta 1$ leu2 $\Delta 0$ lys2 $\Delta 0$ ura3 $\Delta 0$ | Euroscarf |
| BY4741 | MATa his $3\Delta 1$ leu $2\Delta 0$ met $15\Delta 0$ ura $3\Delta 0$ | Euroscarf |
| SH6704 | MATα $\Delta dse2$::kanMX his $3\Delta1$ leu $2\Delta0$ ura $3\Delta0$ | OpenBio |
| SH6705 | MATa $\Delta scw11$::loxP-CgLEU2-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ | systems This chapter |

Table 1. Yeast strains used in this chapter.

| SH6706 | MATα $\Delta eaf3$::loxP-CgHIS3-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ lys2 $\Delta 0$ met15 $\Delta 0$ | This |
|--------|--|---------|
| | | chapter |
| SH6707 | MATa $\Delta sed1::loxP$ -zeocin-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ lys2 $\Delta 0$ met15 $\Delta 0$ | This |
| | | chapter |
| SH6708 | MATa $\Delta dse2::kanMX \Delta scw11::loxP-CgLEU2-loxP his3\Delta1 leu2\Delta0 ura3\Delta0$ | This |
| | $met15\Delta0$ | chapter |
| SH6709 | MAT $\alpha \Delta dse2::kanMX \Delta eaf3::loxP-CgHIS3-loxP his3\Delta1 leu2\Delta0 ura3\Delta0$ | This |
| | $met15\Delta0$ | chapter |
| SH6710 | MATa $\Delta dse2::kanMX \ \Delta sed1::loxP-zeocin-loxP \ his3\Delta1 \ leu2\Delta0 \ ura3\Delta0$ | This |
| | $lvs2\Delta0$ | chapter |
| SH6711 | MATa $\Delta scw11::loxP-CgLEU2-loxP$ $\Delta eaf3::loxP-CgHIS3-loxP$ his $3\Delta 1$ | This |
| | $leu 2\Delta 0 ura 3\Delta 0 lvs 2\Delta 0 met 15\Delta 0$ | chanter |
| | | Th:- |
| SH6712 | MATa $\Delta scw11::loxP-CgLEU2-loxP \Delta sed1::loxP-zeocin-loxP miss \Delta 1$ | I nis |
| | $leu2\Delta 0$ ura $3\Delta 0$ | chapter |
| SH6713 | MATa $\Delta eaf3::loxP-CgHIS3-loxP \Delta sed1::loxP-zeocin-loxP his3\Delta1 leu2\Delta0$ | Inis |
| | $ura3\Delta 0$ | chapter |
| SH6714 | MATo. $\Delta dse2::kanMX \Delta scw11::loxP-CgLEU2-loxP \Delta eaf5::loxP-CgHIS3-loxP h1s3 \Delta 1$ | This |
| | $leu2\Delta 0 \ ura3\Delta 0 \ met15\Delta 0$ | chapter |
| SH6715 | MATo. $\Delta dse2::kanMX \Delta scw11::loxP-CgLEU2-loxP \Delta sed1::loxP-zeocin-loxP his 3 \Delta 1$ | This |
| | $leu2\Delta 0$ | chapter |
| SH6716 | $MAT \alpha \ \Delta dse2::kanMX \ \Delta eaf3::loxP-CgHIS3-loxP \ \Delta sed1::loxP-zeocin-loxP$ | This |
| | his $3\Delta 1$ leu $2\Delta 0$ ura $3\Delta 0$ lys $2\Delta 0$ | chapter |
| SH6717 | $MAT \alpha \ \Delta scw11::loxP-CgLEU2-loxP \ \Delta eaf3::loxP-CgHIS3-loxP \ \Delta sed1::loxP$ | This |
| | -zeocin-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ | chapter |
| SH6718 | $MAT \alpha \ \Delta dse2::kanMX \ \Delta scw11::loxP-CgLEU2-loxP \ \Delta eaf3::loxP-CgHIS3-loxP$ | This |
| • | $\Delta sed1::loxP$ -zeocin-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ lys2 $\Delta 0$ | chapter |
| SH6764 | MATα his $3\Delta 1$ leu $2\Delta 0$ lys $2\Delta 0$ ura $3\Delta 0$ $\Delta pdc1$::LDH-HPH | This |
| | | chapter |
| SH6765 | MATα $\Delta dse2$::kanMX his3 $\Delta1$ leu2 $\Delta0$ ura3 $\Delta0$ $\Delta pdc1$::LDH-HPH | This |
| | | chapter |
| SH6766 | $MAT \alpha \ \Delta scw11::loxP-CgLEU2-loxP \ his3\Delta1 \ leu2\Delta0 \ ura3\Delta0$ | This |
| | Δpdc1::LDH-HPH | chapter |
| SH6767 | MATa $\Delta eaf3::loxP$ -CgHIS3-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ lys2 $\Delta 0$ met15 $\Delta 0$ | This |
| | Δpdc1::LDH-HPH | chapter |
| SH6768 | MATa $\Delta sed1::loxP$ -zeocin-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ lys2 $\Delta 0$ met15 $\Delta 0$ | This |
| | Δpdc1::LDH-HPH | chapter |
| SH6769 | MATα $\Delta dse2$::kanMX $\Delta scw11$::loxP-CgLEU2-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ | This |
| | met $15\Delta0 \ \Delta pdc1::LDH-HPH$ | chapter |
| SH6770 | MATa $\Delta dse2::kanMX \ \Delta eaf3::loxP-CeHIS3-loxP \ his3\Delta1 \ leu2\Delta0 \ ura3\Delta0$ | This |
| | met $15\Delta0 \Delta pdc1::LDH-HPH$ | chapter |
| SH6771 | MATa $\Delta dse2$::kanMX $\Delta sed1$::loxP-zeocin-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ | This |
| | lys2Δ0 Δpdc1::LDH-HPH | chapter |
| | × 1 | - |

.

| SH6772 | $MAT \alpha \ \Delta scw11::loxP-CgLEU2-loxP \ \ \Delta eaf3::loxP-CgHIS3-loxP \ his3\Delta 1$ | This |
|--------|--|---------|
| | $leu2\Delta 0 \ ura3\Delta 0 \ lys2\Delta 0 \ met15\Delta 0 \ \Delta pdc1::LDH-HPH$ | chapter |
| SH6773 | $MATa \Delta scw11::loxP-CgLEU2-loxP \Delta sed1::loxP-zeocin-loxP his3\Delta1$ | This |
| | $leu2\Delta 0 ura3\Delta 0 \Delta pdc1::LDH-HPH$ | chapter |
| SH6774 | MATa $\Delta eaf3::loxP-CgHIS3-loxP \Delta sed1::loxP-zeocin-loxP his3\Delta1 leu2\Delta0$ | This |
| | $ura3\Delta0 \Delta pdc1::LDH-HPH$ | chapter |
| SH6775 | MATα Δdse2::kanMX Δscw11::loxP-CgLEU2-loxP Δeaf3::loxP-CgHIS3-loxP | This |
| | his $3\Delta 1 \ leu 2\Delta 0 \ ura 3\Delta 0 \ met 15\Delta 0 \ \Delta pdc 1::LDH-HPH$ | chapter |
| SH6776 | $MAT \alpha \ \Delta dse2::kanMX \ \Delta scw11::loxP-CgLEU2-loxP \ \Delta sed1::loxP-zeocin-loxP$ | This |
| | his $3\Delta 1$ leu $2\Delta 0$ ura $3\Delta 0$ lys $2\Delta 0$ $\Delta pdc1::LDH-HPH$ | chapter |
| SH6777 | MATα Δdse2::kanMX Δeaf3::loxP-CgHIS3-loxP Δsed1::loxP-zeocin-loxP | This |
| | his $3\Delta 1 \ leu 2\Delta 0 \ ura 3\Delta 0 \ lys 2\Delta 0 \ \Delta pdc 1::LDH-HPH$ | chapter |
| SH6778 | $MATa \Delta scw11::loxP-CgLEU2-loxP \Delta eaf3::loxP-CgHIS3-loxP$ | This |
| | $\Delta sed1::loxP$ -zeocin-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ $\Delta pdc1::LDH$ -HPH | chapter |
| SH6779 | MATα Δdse2::kanMX Δscw11::loxP-CgLEU2-loxP Δeaf3::loxP-CgHIS3-loxP | This |
| | $\Delta sed1::loxP$ -zeocin-loxP his3 $\Delta 1$ leu2 $\Delta 0$ ura3 $\Delta 0$ lys2 $\Delta 0$ $\Delta pdc1::LDH$ -HPH | chapter |

2.2.2 Construction of plasmid vectors

Integration vector pBHPH-PDC1P-LDHKCB (Fig. 1), which was based on pBTRP-PDC1P-LDHKCB plasmid (Ishida *et al.*, 2005), consist of the *PDC1* promoter, L-LDH from bovine (GeneBank No. D90141), *TDH3* terminator, hygromycin B phosphotransferase (*HPH*, GenBank No. K01193) (Gritz and Davies, 1983), and *PDC1-3'* fragment. *HPH* fragment was isolated by PCR using the genomic DNA of the *Escherichia coli* K12 strain as a template and fused between the *S. cerevisiae TDH3* promoter and *CYC1* terminator. This fragment was replaced by *TRP1* fragment of pBTRP-PDC1P-LDHKCB plasmid. For construction of plasmid having *loxP*-zeocin-*loxP* cassette, zeocin resistant gene was amplified from pREMI-Z (van Dijk *et al.*, 2001) using Zeocin F (Table 2), containing a *Bgl*II restriction site (underlined) and Zeocin R (Table 2), containing a *Xho*I restriction site (underlined). The amplified fragment treated with *Bgl*II and *Xho*I, was ligated to pUG6 vector to yield plasmid pUG6-zeocin^r.



Figure 1. Structure of the plasmid pBHPH-PDC1P-LDHKCB and breeding of transgenic *S. cerevisiae.* The constructed DNA fragment, which was obtained by digesting the plasmid pBHPH-PDC1P-LDHKCB with *Sac* I and *Kpn* I, was integrated into the *PDC1* ORF region of the *S. cerevisiae* disruptant strains by homologous recombination.

2.2.3 Construction of single and multiple disruptants

For construction of single disruptants, *DSE2*, *SCW11*, *EAF3*, or *SED1* ORF was replaced with *kanMX4* (amplified from the genomic DNA of SH6704 using the primers DSE2 F and DSE2 R, Table 2), *loxP-CgLEU2-loxP* cassette [amplified from p3008 (Sugiyama *et al.*, 2005) using the primers SCW11 F and SCW11 R, Table 2], *loxP-CgHIS3-loxP* cassette [amplified from p3009 (Sugiyama *et al.*, 2005) using the primers EAF3 F and EAF3 R, Table 2], and *loxP-zeocin-loxP* cassette (amplified from pUG6-zeocin^r using the primers SED1 F and SED1 R, Table 2), respectively. Multiple disruptants were constructed by the following procedure: (i) *DSE2* and *SCW11* of BY4742 were disrupted with homologous recombination using *kanMX4*

and loxP-CgLEU2-loxP cassette, respectively. (ii) *EAF3* and *SED1* of BY4741 (*MATa his3* Δ 1 *leu2* Δ 0 *met15* Δ 0 *ura3* Δ 0) (Brachmann *et al.*, 1998) were disrupted with homologous recombination using *loxP-CgHIS3-loxP* cassette and *loxP-zeocin-loxP* cassette, respectively. (iii) These double disruptants were crossed, and multiple gene disruptants were constructed by tetrad analysis as showing G418 resistant, leucine protorophic, histidine protorophic, and/or zeocin resistant phenotype. The strains for production of lactic acid were constructed by transforming a DNA fragment, which was obtained by digesting the pBHPH-PDC1P-LDHKCB plasmid with *SacI* and *KpnI*, into each disruptant strain (Fig. 1).

| Oligo- nucleotide | Sequence |
|----------------------|--|
| Zeocin F | 5'-CTC <u>AGATCT</u> CCCACACACCATAGCTTCAA-3' |
| Zeocin R | 5'-CTC <u>CTCGAG</u> CCAGCTTGCAAATTAAAGCC-3' |
| DSE2 F | 5'-CAGAGTAGAAATAAAGCCACTCCTTTAACAAATTACAAAGAAATGCTTCGTACGCTGCAG-3' |
| DSE2 R | 5'-GAAAACAAATAGAAAAGAATGCACCACCATTATGCCTAAGCGCTAGCCACTAGTGGATCT-3' |
| SCW11 F | 5'-TTACGCTACACTCATTGATATAATATCTAATAGAAAACCATCATGCTTCGTACGCTGCAG-3' |
| SCW11 R | 5'-CTGATCTTTTATATGCATGTTTCTCTATTCGATTTAGAAAACTTAGCCACTAGTGGATCT-3' |
| EAF3 F | 5'-GTGAGGCCTCGTCACTGGATTTACCCTATTGAAGAACGTATAATGCTTCGTACGCTGCAG-3' |
| EAF3 R | 5'-AACTAAATACTAGAAATAATCCCAAGCTAGAATATAAACGTCTCAGCCACTAGTGGATCT-3' |
| SED1 F | 5'-ACTACAAAGACAAGCAAAATAAAATACGTTCGCTCTATTAAGATGCTTCGTACGCTGCAG-3' |
| SED1 R | 5'-GAAAGAAAGCATTAAGAAGGCGGATGTGTCAAACACCACCGTTTAGCCACTAGTGGATCT-3' |

Table 2. Sequence of the deoxy-oligonucleotides used in this chapter.

Underline represents restriction site.

2.2.4 Screening for L-lactic-acid-resistant disruptants

To screen for L-lactic-acid-resistant strains, the nonessential genes deletion collection was stamped onto YPDA and 6% L-lactic acid plates with a 96-prong

replicator. Growth of the disruptants was scored visually by comparison with that of the parent strain (BY4742), which was stamped similarly onto YPDA and 6% L-lactic acid plates. Before screening, single-gene disruptants were tested for growth on YPDA medium containing 300 mg/l of G418 disulfate (Wako Pure Chemical Industries, Osaka).

2.2.5 Spot dilution growth assay

Yeast cells were grown in YPDA at 30°C to mid-log phase ($OD_{660} = 1.0$). Cells were harvested by centrifugation, washed once with water, and resuspended in water. The cell density was normalized to 1×10^6 cells/5 µl. A 10-fold serial dilution of this culture was made, and 5 µl of each dilution was spotted onto YPDA medium or lactic-acid medium. Cells were incubated at 30°C and scored visually by comparison with that of BY4742.

2.2.6 Measurement of lactic acid production

Fermentation experiments were performed according to the method of Ishida *et al.* (2005). The inoculum of each strain was prepared by cultivating overnight in YPDA 5 ml at 30°C. Overnight culture ($OD_{660} = 1.2$) 2 ml was transferred to 30 ml YPD10 medium (1% Bacto yeast extract, 2% Bacto peptone, 10% D-glucose). Each culture was incubated by static culture at 30°C for 72 h without neutralization. The glucose, lactic acid, and ethanol concentrations were measured with a biosensor BF-5 instrument (Oji Keisoku Kiki, Hyogo). The *post hoc* Tukey's test was used for multiple comparisons between strains with statistical significance set at a *p*

value lower than 0.05 (Toothaker, 1993). Data analysis was carried out with IBM SPSS Statistics version 21 (IBM SPSS Inc., Chicago, USA).

2.3 Results

2.3.1 Genome-wide screening for *S. cerevisiae* L-lactic-acid-resistant disruptants

The purpose of this chapter was to construct a superior lactic-acid-resistant strain by combining gene disruptions that lead to lactic-acid resistance, and to investigate the possibility of enhancing lactic-acid productivity without Although Kawahata et al. (2006) reported 46 genes whose neutralization. disruption led to resistance to 5.1% lactic acid, in this chapter I intended to screen for strains showing stronger resistance under more severe conditions. Α concentration of at least 5% lactic acid in culture media (pH 2.7) seems to exert toxic effects on yeast (Ishida et al., 2005). To obtain more highly lactic-acid-resistant S. cerevisiae strains, I first systematically screened for viable yeast haploid single disruptants (4828 strains) that were resistant to 6% L-lactic acid stress (pH 2.6), a condition in which the wild-type strain (BY4742) hardly grew (Fig. 2). Cell suspensions of freshly grown cultures of all of the disruption mutants were transferred to YPDA plates containing 0% and 6% L-lactic acid by using a 96-prong replicator and incubated for 1 day (0%) or 6 days (6%), and growth was assessed visually. The screen was carried out in triplicate. Disruption alleles of candidate disruptants were structurally confirmed by PCR

analysis (data not shown).

As a result of screening, I identified 94 genes (1.9% of the total) that, when disrupted, led to a phenotype resistant to 6% L-lactic acid (Table 3). These genes were categorized by gene function according to the Munich Information Center for Protein Sequences (MIPS, http://mips.gsf.de/) (Mewes *et al.*, 2002) and *Saccharomyces* Genome Database (SGD, http://www.yeastgenome.org/) (Dwight *et al.*, 2002). Although 25 of the 94 genes had previously been reported to increase resistance to 5.1% lactic acid in a synthetic glucose minimal medium (Kawahata *et al.*, 2006), my analysis newly identified 69 genes whose gene disruption increased resistance to 6% lactic acid in YPDA glucose-rich medium. Although my screening condition was more severe than that used in Kawahata's work (2006), not all genes that have previously been reported (Kawahata *et al.*, 2006) were identified in my screening. One reason for this may be that I used a glucose-rich medium (YPDA) containing lactic acid as a screening medium, whereas Kawahata *et al.* (2006) used a synthetic glucose minimal medium containing lactic acid.

| ORF | Description | | phosphatase |
|--|---------------------------------|-----------------|-----------------------------------|
| Cellular transport, transport facilities and | | <u>PTC7</u> | 2C protein phosphatase |
| transport routes | | <u>RUP1</u> | Protein involved in regulation |
| i) vacuolar/lysosomal transport | | | of Rsp5 |
| ATG22 | Breakdown of autophagic | <u>SHR5</u> | Subunit of a almitoyltransferase |
| | vesicles inside the vacuole | <u>YPL150w</u> | Putative Ser/Thr protein kinase |
| SYN8 | Endosomal SNARE protein | <u>YPL236c</u> | Protein kinase localised to |
| <u>VPS62</u> | Vacuolar protein sorting | | vacuolar membrane |
| <u>VPS70</u> | Protein involved in vacuolar | ii) protein/per | otide degradation |
| | sorting | <u>CDH1</u> | Substrate-specific activator of |
| <u>VPS74</u> | Protein involved in vacuolar | | APC-dependent proteolysis |
| | sorting | <u>MMS2</u> | Ubiquitin-conjugating enzyme |
| VTC3 | Subunit of the vacuolar | <u>RKR1</u> | Nuclear RING domain E3 |
| | transporter chaperone complex | | ubiquitin ligase |
| ii) vesicular transport (Golgi network, etc.) | | <u>TRE1</u> | Transferrin receptor-like protein |
| <u>ALY1</u> | α arrestin that controls | UBP2 | Ubiquitin-specific proteinase |
| | intracellular sorting of Gap1 | Transcription | a |
| <u>ERP2</u> | P24 protein involved in | <u>SR09</u> | La motif-containing proteins |
| | membrane trafficking | | that modulate mRNA translation |
| ERV25 | Component of the COPII-coated | i) RNA synthe | esis |
| | vesicles | ACE2 | Metallothionein expression |
| MVB12 | ESCRT-I subunit | | activator |
| iii) transport f | acilities | <u>ASH1</u> | Negative regulator of HO |
| ATP4 | F1F0-ATPase complex | | transcription |
| <u>THI72</u> | Transporter of thiamine or | <u>CBF1</u> | Centromere binding factor 1 |
| | related compound | EAF3 | Esa1p-associated factor |
| iv) nuclear tra | nsport | EAF5 | Esalp-associated factor |
| <u>YHL010c</u> | Protein of unknown function | EAF6 | Esa1p-associated factor |
| Protein fate (folding, modification, estination) | | <u>SFL1</u> | Transcriptional repressor and |
| i) protein mod | lification | | activator |
| <u>FRK1</u> | Putative protein kinase | YAF9 | Subunit of the NuA4 and the |
| <u>MNN2</u> | Type II membrane protein | | SWR1 complex |
| MSG5 Dual-specifity protein | | ii) RNA proce | essing |

٠

Table 3. Genes whose loss results in resistance to 6% L-lactic acid.

20

. .

| <u>CWC15</u> | Protein involved in mRNA | | to glucanases |
|--|-----------------------------------|--------------------------------|---------------------------------------|
| | splicing | <u>SP07</u> | Meiotic protein |
| ISY1 Pre-mRNA splicing factor | | ii) DNA processing | |
| Metabolism | | <u>MEI4</u> | Meiosis-specific protein |
| i) lipid, fatty a | acid and isoprenoid metabolism | iii) DNA repa | ir |
| <u>ACB1</u> | Acyl-coenzyme-A-binding | UBC13 | E2 ubiquitin-conjugating enzyme |
| | protein | Cell type diff | erentiation |
| <u>FENI</u> | Fatty acid elongase | i) fungal/micro | oorganismic cell type differentiation |
| <u>LDH1</u> | Serine hydrolase | <u>AXL2</u> | Protein required for axial |
| <u>PSD2</u> | Lipid, fatty acid and isoprenoid | | pattern of budding |
| | metabolism | <u>BUD4</u> | Budding protein |
| ii) C-compour | nd and carbohydrate metabolism | <u>DIT1</u> | Spore wall maturation protein |
| <u>GND1</u> | 6-phosphogluconate | <u>RBD2</u> | Protein of unknown function |
| | dehydrogenase | Biogenesis of | cellular components |
| <u>TCO89</u> | TOR complex 1 | i) cell wall | |
| iii) nucleotide/nucleoside/nucleobase metabolism | | LRG1 | GTPase-activating protein of the |
| <u>URA5</u> | Orotate phosphoribosyltransferase | | Rho/Rac family |
| <u>YND1</u> | Nucleoside diphosphatase | SED1 | Abundant cell surface glycoprotein |
| iv) metabolism of vitamins, cofactors, and | | ii) eukaryotic plasma membrane | |
| prosthetic gro | ups | <u>SEY1</u> | Synthetic enhancement with |
| · <u>SIS2</u> | Negative regulatory subunit of | | YOP1 |
| | Ppz1 and a subunit of PPCDC | Cellular com | munication/Signal transduction |
| v) amino acid | metabolism | mechanism | |
| <u>TKL1</u> | Transketolase 1 | <u>VAC17</u> | Vacuole-specific receptor of |
| Cell cycle and | I DNA processing | | Myo2 |
| i) cell cycle | | i) cellular sigr | aling |
| <u>BUD3</u> | Budding protein | <u>GIS4</u> | CAAX box containing protein |
| CTS1 | Endochitinase | Cell fate | |
| <u>СҮКЗ</u> | SH3-domain protein | i) cell growth | |
| DSE2 | Daughter cell-specific secreted | <u>CKA2</u> | α catalytic subunit of casein |
| | protein with similarity to | | kinase II |
| | glucanases | ii) cell aging | |
| EGT2 | Cell-cycle regulation protein | SUN4 | Protein involved in the aging |
| SCW11 | Cell wall protein with similarity | | process |

| i) stress respon | se |
|------------------|-------------------------------------|
| SIP5 | Protein facilitates the interaction |
| | between Glc7 and Snf1 |
| Interaction wi | th the environment |
| i) homeostasis | of cations |
| <u>FRE6</u> | Putative ferric reductase |
| Unclassified p | roteins |
| <u>CEX1</u> | Protein of unknown function |
| <u>FMP49</u> | Questionable protein |
| IRC8 | Protein of unknown function |
| <u>IRC9</u> | Questionable protein |
| <u>MRNI</u> | RNA-binding protein |
| <u>NBA1</u> | Protein of unknown function |
| <u>SLP1</u> | Protein of unknown function |
| SVF1 | Protein with a potential role in |
| | cell survival pathways |
| TOS6 | Protein of unknown function |
| <u>YBR062c</u> | Protein of unknown function |
| <u>YBR187w</u> | Protein of unknown function |
| • <u>YCL062w</u> | Questionable protein |
| <u>YDR357c</u> | Protein of unknown function |
| <u>YHR078w</u> | Conserved hypothetical protein |
| <u>YHL017w</u> | Protein of unknown function |
| <u>YIL141w</u> | Questionable protein |
| <u>YKL063c</u> | Protein of unknown function |
| <u>YKR073c</u> | Questionable protein |
| <u>YLR346c</u> | Protein of unknown function |
| YMR294w-A | Questionable protein |
| <u>YNR005c</u> | Questionable protein |
| <u>YOR139c</u> | Questionable protein |

.

Cell rescue, defense and virulence

| <u>YOR289w</u> | Protein of unknown function |
|----------------|-----------------------------|
| YPL191c | Protein of unknown function |
| <u>ZRG8</u> | Protein of unknown function |

Underline represents newly identified gene whose disruption results in resistance to lactic acid. Other genes were also identified by Kawahata *et al.* (2006).

ź.

2.3.2 Multiple disruptants show superior lactic-acid resistance

My analysis identified disruptants of DSE2, SCW11, EAF3, and SED1, which belonged to the group showing strong resistance to 6% lactic acid as compared with the parental strain BY4742 (Fig. 2). Kawahata et al. (2006) reported that $\Delta scw11$ and $\Delta eaf3$ disruptants showed 0.5% acetic-acid resistance while $\Delta dse2$, $\Delta scw11$, $\Delta eaf3$, and $\Delta sed1$ disruptants did not show 0.28% hydrochloric-acid resistance. The lactic-acid-resistant phenotype of these disruptants was subsequently confirmed to be due to a single disruption, as indicated by tetrad analysis of 6-10 asci, which lactic-acid-resistance/2 lactic-acid-sensitive segregation, showed 2 and cosegregation with the disruption marker (kanMX) (data not shown). In the following experiments, I primarily used these 4 disruptants as lactic-acid resistance had been confirmed to be due to a single disruption.

I expected that multiple disruptants harboring various combinations of the $\Delta dse2$, $\Delta scw11$, $\Delta eaf3$, and $\Delta sed1$ disruptions might show more superior lactic-acid resistance. Therefore, I constructed single, double, triple, and quadruple disruptants of these genes by tetrad analysis as described in Materials and methods (Table 1), and examined the lactic-acid resistance of these multiple disruptants (Fig. 2). There was no significant difference between the growth curve of BY4742 and that of 15 disruptants (SH6704–SH6718) in YPDA without lactic acid (data not shown). As shown in Fig. 2, all of the multiple disruptants showed resistance to 7% lactic acid (pH 2.5), and almost all disruptants except for $\Delta dse2\Delta scw11$ (SH6708) and $\Delta dse2\Delta scd1$ (SH6710) displayed enhanced lactic-acid resistance as compared with the single-gene disruptants.



Figure 2. Assay of growth resistance of single and multiple disruptants to lactic acid. A ten-fold dilution of each yeast strain was spotted onto YPDA medium, 6% lactic-acid medium (pH 2.6), and 7% lactic-acid medium (pH 2.5) as described in Materials and methods. Cells were incubated at 30°C for 24 h (YPDA medium), 5 days (6% lactic-acid medium), and 10 days (7% lactic-acid medium). The strains assayed were $\Delta dse2$ (SH6704), $\Delta scw11$ (SH6705), $\Delta eaf3$ (SH6706), $\Delta sed1$ (SH6707), $\Delta dse2\Delta scw11$ (SH6708), $\Delta dse2\Delta eaf3$ (SH6709), $\Delta dse2\Delta sed1$ (SH6710), $\Delta scw11\Delta eaf3$ (SH6711), $\Delta scw11\Delta sed1$ (SH6712), $\Delta eaf3\Delta sed1$ (SH6713), $\Delta dse2\Delta scw11\Delta eaf3$ (SH6714), $\Delta dse2\Delta scw11\Delta sed1$ (SH6715), $\Delta dse2\Delta eaf3\Delta sed1$ (SH6716), $\Delta scw11\Delta eaf3\Delta sed1$ (SH6717), and $\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$ (SH6718).

The multiple disruptant showing the highest lactic-acid resistance seemed to be the $\Delta dse2\Delta scw11\Delta sed1$ disruptant (SH6715) judging from Fig. 2 since it showed the highest number of colonies in each serial dilution spots on 7% lactic-acid medium. However, its colony size was very small when compared with those of other multiple disruptants. This may come from a complex effect of highest survival

rate but slower growth rate that the $\Delta dse2\Delta scw11\Delta sed1$ disruptant displays against lactic-acid stress. Since the difference of level of resistance among multiple disruptants appeared to be obscure on lactic-acid plate and weak organic acid such as sorbic acid causes the extended lag phase (Lambert et al., 1999), I also examined the growth behavior of the multiple disruptants in liquid culture under 6% lactic-acid condition to see what extent of the lactic-acid resistance of the multiple disruptants was enhanced as compared with the single-gene disruptants (Fig. 3). Six strains, $\Delta eaf3$ (SH6706), $\Delta dse2\Delta eaf3$ (SH6709), $\Delta dse2\Delta sed1$ (SH6710), $\Delta scw11\Delta eaf3$ (SH6711), $\Delta dse2\Delta scw11\Delta eaf3$ (SH6714), and $\Delta dse2\Delta eaf3\Delta sed1$ (SH6716), showed almost the same pattern of growth as other strains and are not shown in Fig. 3. That is, the growth curve of $\Delta eaf3$ (SH6706) was almost the same as that of $\Delta scw11$ (SH6705). Similarly, $\Delta dse2\Delta eaf3$ (SH6709) and $\Delta scw11\Delta ead3$ (SH6711) were the same as $\Delta dse2\Delta scw11$ (SH6708); $\Delta dse2\Delta sed1$ (SH6710) and $\Delta dse2\Delta scw11\Delta eaf3$ (SH6714) were the same as $\Delta scw11\Delta sed1$ (SH6712); and $\Delta dse2\Delta eaf3\Delta sed1$ (SH6716) was the same as $\Delta scw11\Delta eaf3\Delta sed1$ (SH6717). Since pH of culture media of strains including quadruple disruptant was not changed from the initial pH value (pH 2.6) during 120 h cultivation, I believe that the resistance was not acquired by raising the pH of culture media. As shown in Fig. 3, all of the single-gene disruptants showed lower cell density than the multiple disruptants after 120 h incubation and the specific growth rate (μ) of $\Delta sed1$ (SH6707), $\Delta scw11\Delta sed1$ (SH6715), (SH6712), $\Delta dse2\Delta scw11\Delta sed1$ $\Delta scw11\Delta eaf3\Delta sed1$ (SH6717), $\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$ (SH6718) was almost same [$\mu = 0.085$ h⁻¹ (SH6704). 0.074 h⁻¹ (SH6706), 0.10 h⁻¹ (SH6707), 0.072 h⁻¹ (SH6708), 0.10 h⁻¹ (SH6712),

25

0.082 h⁻¹ (SH6713), 0.10 h⁻¹ (SH6715), 0.10 h⁻¹ (SH6717), 0.11 h⁻¹ (SH6718)]. The quadruple disruptant ($\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$; SH6718) displayed both the shortest lag phase and the highest cell density than any other disruptants after 120 h of incubation. Based upon these results, I concluded that the highest resistant-strain to 6% lactic acid was $\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$ quadruple disruptant (SH6718). I also examined the resistance of all the 15 disruptants (SH6704-SH6718) against hydrochloric acid (0.4%). Hydrochloric-acid resistance of all of 15 disruptants was not stronger than that of wild-type strain (BY4742) (data not shown). This result suggests that all the 15 disruptants show the resistance to the effect of lactic-acid anion but not proton. I show that a strategy of constructing strains with multiple disruptions of genes whose disruption leads to lactic-acid resistance is an effective way to enhance resistance to lactic acid.



Figure 3. Growth curve of single and multiple disruptants under 6% lactic acid without neutralization. Each strain was pre-cultured on YPDA at 30 °C and transferred to 6% lactic-acid medium ($OD_{660} = 0.1$); the OD_{660} of the culture was then monitored periodically for 120 h. Symbols: closed diamond ($\Delta dse2$, SH6704), closed triangle ($\Delta scw11$, SH6705), asterisk ($\Delta sed1$, SH6707), closed square ($\Delta dse2\Delta scw11$, SH6708), open triangle ($\Delta scw11\Delta sed1$, SH6712), open diamond ($\Delta eaf3\Delta sed1$, SH6713), closed circle ($\Delta dse2\Delta scw11\Delta sed1$, SH6715), open square ($\Delta scw11\Delta eaf3\Delta sed1$, SH6717), open circle ($\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$, SH6718). Broken line indicates the growth curve of the wild-type strain (BY4742). Error bars show the standard deviation of three independent experiments.

2.3.3 A lactic-acid-resistant strain that produces lactic acid more efficiently than wild type

I expected that lactic-acid production in the lactic-acid resistant strains would be improved as compared with that in the wild-type strain under non-neutralizing conditions. *S. cerevisiae* converts pyruvic acid to ethanol and Pdc1 is the main pyruvate decarboxylase. Therefore, to decrease the production of ethanol and to generate efficient production of lactic acid, the bovine L-lactate dehydrogenase gene (L-*LDH*) was introduced into the *PDC1* locus of the wild-type strain (BY4742), as well as single and multiple disruptants, by homologous recombination with a DNA fragment obtained by digesting the plasmid pBHPH-PDC1P-LDHKCB (Fig. 1). The *LDH*-harboring BY4742 (SH6764) strain produced lactic acid and, as expected, the ethanol productivity of this strain decreased as compared with that of BY4742 (data not shown).

Because the *LDH*-harboring strains acquired the ability to intrinsically produce lactic acid, I examined to what extent single and multiple disruptants harboring the *L-LDH* gene displayed lactic-acid resistance (Fig. 4). As shown in Fig. 4, all single and multiple disruptants with bovine *LDH* showed resistance to 6% lactic-acid, whereas the wild-type strain harboring *LDH* could not. Interestingly, the *LDH*-harboring $\Delta dse2\Delta scw11$ (SH6769), $\Delta dse2\Delta eaf3$ (SH6770), $\Delta scw11\Delta eaf3\Delta sed1$ (SH6778), and $\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$ (SH6779) strains displayed high resistance to 7% lactic acid, although for almost all strains the degree of resistance to 6% and 7% lactic acid was different from that shown in Fig.

. 2.



Figure 4. Assay of growth resistance of *LDH*-harboring single and multiple disruptants to lactic acid. A ten-fold dilution of each yeast strain was spotted onto YPDA medium, 6% lactic-acid medium (pH 2.6), and 7% lactic-acid medium (pH 2.5) as described in Materials and methods. Cells were incubated at 30°C for 24 h (YPDA medium) and 6 days (6% and 7% ·lactic-acid medium). The strains assayed were BY4742-*LDH* (SH6764), $\Delta dse2$ -*LDH* (SH6765), $\Delta scw11$ -*LDH* (SH6766), $\Delta eaf3$ -*LDH* (SH6767), $\Delta sed1$ -*LDH* (SH6768), $\Delta dse2\Delta scw11$ -*LDH* (SH6769), $\Delta dse2\Delta eaf3$ -*LDH* (SH6770), $\Delta dse2\Delta sed1$ -*LDH* (SH6771), $\Delta scw11\Delta eaf3$ -*LDH* (SH6772), $\Delta scw11\Delta sed1$ -*LDH* (SH6773), $\Delta eaf3\Delta sed1$ -*LDH* (SH6776), $\Delta dse2\Delta scw11\Delta eaf3$ -*LDH* (SH6777), $\Delta scw11\Delta eaf3\Delta sed1$ -*LDH* (SH6778), and $\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$ -*LDH* (SH6777).

I then examined the productivity of lactic acid in the disruptants harboring *LDH* without neutralization after 72 h cultivation, at which time all glucose in medium was consumed in almost all disruptants (Fig. 5). The multiple comparisons of productivity between strains were carried out by Tukey's test

(Toothaker, 1993). As shown in Fig. 5, the *LDH*-harboring quadruple disruptant $(\Delta dse2\Delta scw11\Delta eaf3\Delta sed1;$ SH6779) which showed the strongest lactic-acid resistance produced significantly high concentration of lactic acid (48 g/litter, 72h) as compared with wild-type strain (SH6764) (38 g/litter, 72 h) (p = 0.024), while that of other 14 disruptants did not exhibit significant increase as compared with SH6764.



Figure 5. Lactic-acid productivity of LDH-harboring single and multiple disruptants. The amount of lactic acid was measured as described in Materials and methods. Error bars show the standard deviation of three independent experiments. Asterisk represents a significant difference (p < 0.05) from LDH-harboring wild-type strain (SH6764) (Tukey's multiple comparisons). The strains assayed were BY4742-LDH (SH6764), ∆dse2-LDH (SH6765), $\Delta scw11$ -LDH (SH6766), $\Delta eaf3$ -LDH (SH6767), $\Delta sed1$ -LDH (SH6768), $\Delta dse2\Delta scw11$ -LDH (SH6769), $\Delta dse2\Delta eaf3$ -LDH (SH6770), $\Delta dse2\Delta sed1$ -LDH (SH6771), $\Delta scw11\Delta eaf3-LDH$ (SH6772), $\Delta scw11\Delta sed1-LDH$ (SH6773), $\Delta eaf3\Delta sed1-LDH$ (SH6774), $\Delta dse2\Delta scw11\Delta eaf3$ -LDH (SH6775), $\Delta dse2\Delta scw11\Delta sed1$ -LDH (SH6776), $\Delta dse_{2}\Delta eaf_{3}\Delta sed_{1}$ -LDH (SH6777), $\Delta scw_{11}\Delta eaf_{3}\Delta sed_{1}$ -LDH (SH6778), and $\Delta dse2\Delta scw11\Delta eaf3\Delta sed1-LDH$ (SH6779).

2.4 Discussion

The newly identified genes belong to a variety of functional categories. For example, the deletion of genes related to ubiquitination (RUP1, MMS2, RKR1, and UBC13) was found for the first time to lead to lactic-acid resistance. Ubiquitination is involved in resistance to various types of stress, such as UV, canavanine, high temperature (39°C), LiCl, sorbitol, ethanol, and H₂O₂ (Hofmann and Pickart, 1999; Seufert and Jentsch, 1990; Hiraishi et al., 2006). Ubiquitin/proteasome-dependent proteolysis consumes ATP in order to degrade abnormal proteins (Hochstrasser, 1995). The role of ATPase, which consumes much ATP during growth (van der Rest et al., 1995), is also important for regulating intracellular pH under acidic conditions. Therefore, it may be advantageous for the survival of yeast cells in a high concentration of lactic acid if ATP is utilized to remove protons from the cytosol via ATPase rather than to degrade proteins by the ubiquitin-proteasome pathway, although an accumulation of misfolded proteins in the endoplasmic reticulum causes endoplasmic reticulum stress (Schröder and Kaufman, 2005).

 $\Delta dse2$, $\Delta scw11$, $\Delta eaf3$, and $\Delta sed1$ disruptants showed strong resistance to 6% lactic acid. According to the SGD database, Dse2 and Scw11 both function as glucanases that are involved in the separation of daughter cells from mother cells (Cappellaro *et al.*, 1998; Colman-Lerner *et al.*, 2001; Ufano *et al.*, 2004). Because the continuous synthesis of glucan may lead to wall thickening (Dominguez *et al.*, 1978), the $\Delta scw11$ and $\Delta dse2$ disruptants may have a thick cell wall and therefore show increased resistance to lactic acid and/or other substances. Eaf3 is a
component of the NuA4 acetyltransferase complex. In the $\Delta eaf3$ disruptant, histone acetylation is decreased at promoter regions and increased at coding regions, and the transcriptional level of a few yeast genes increase (Reid *et al.*, 2004). *FET3*, which is negatively regulated by Eaf3, is involved in high-affinity iron uptake, and it has been suggested that iron metabolism may be important for growth under acidic conditions (Kawahata *et al.*, 2006). Therefore, enhancement of lactic-acid resistance in the $\Delta eaf3$ strain may be caused by increased expression of Eaf3-downregulated genes such as *FET3*. Sed1 is a major stress-induced GPI-cell wall glycoprotein that associates with translating ribosomes and is involved in mitochondrial genome maintenance (van der Vaart *et al.*, 1996; Shimoi *et al.*, 1998). Because the $\Delta sed1$ disruptant shows somewhat increased resistance to calcofluor white and Congo red (Caro *et al.*, 1998), this disruption may lead to an increase in the maintenance of cell-wall integrity and thus to enhancement of lactic-acid resistance.

Lactic-acid resistance of the $\Delta dse2\Delta eaf3$ double disruptant was higher than that of the $\Delta dse2\Delta scw11\Delta eaf3$ and $\Delta dse2\Delta eaf3\Delta sed1$ triple disruptants in experiment shown in Fig. 4. However, I did not see clear difference in the level of the resistance among these three disruptants in Fig. 2. Therefore, I think that the difference in resistance between the $\Delta dse2\Delta eaf3$ double disruptant and the $\Delta dse2\Delta scw11\Delta eaf3$ and $\Delta dse2\Delta eaf3\Delta sed1$ triple disruptants shown in Fig. 4 seems to come from a consequence of the expression of the *LDH* that was integrated into the genome of those strains shown in Fig. 4. Although the mechanisms are unknown, one possible interpretation would be that this difference in resistance in Fig. 4 was caused by the alteration of pyruvate metabolism. Since at least some amount of pyruvate is fluxed to the synthesis of lactic acid in *LDH*-containing disruptant, TCA cycle may be slow down and consequently ATP synthesis may decrease compared with disruptant without *LDH* gene. The amount of ATP required for growth under lactic acid may differ with disruption genotype. Another possibility would be that the detrimental effects of intracellularly produced lactic acid in addition to extracellularly added lactic acid to cells may differ with disruption genotype.

In conclusion, I showed that lactic-acid resistance can be enhanced by combining multiple gene disruptions, each of which causes lactic-acid resistance, and also showed that the quadruple disruptant displaying highest lactic-acid resistance led to significant improvement in lactic-acid production without neutralization as compared with wild-type strain.

2.5 Summary

To create strains that have high productivity of lactic acid without neutralization, a genome-wide screening for strains showing hyper-resistance to 6% L-lactic acid (pH 2.6) was performed using the gene deletion collection of *S. cerevisiae*. I identified 94 genes whose disruption led to resistance to 6% lactic acid in rich medium. I also found that multiple combinations of $\Delta dse2$, $\Delta scw11$, $\Delta eaf3$, and $\Delta sed1$ disruption led to enhanced resistance to 7% lactic acid (pH 2.5). In particular, the quadruple disruptant $\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$ grew well in 6% lactic acid with a short lag phase. I then generated strains harboring these multiple disruptions along with an exogenous lactate dehydrogenase gene (*LDH*) gene, and found that some of the multiple disruptants that showed stronger lactic-acid resistance displayed high productivity of lactic acid. In particular, the *LDH*-harboring quadruple disruptant, which showed the strongest lactic-acid resistant strains, showed an increase in lactic-acid productivity of approximately 27% as compared with the *LDH*-harboring wild-type strain (BY4742) (48 g/litter, 72 h). These observations suggest that disruption of multiple genes each of whose deletion leads to lactic-acid resistance is an effective way to enhance resistance to lactic acid, leading to high lactic-acid productivity without neutralization.

Chapter 3

Lactic-acid stress causes vacuolar fragmentation and impairs intracellular amino-acid homeostasis in *S. cerevisiae*

3.1 Introduction

In *S. cerevisiae*, it is well known that adaptation to lipophilic weak organic acids such as sorbic and benzoic acids involves the extraction of organic acid anions by the Pdr12p plasma membrane ATP-binding cassette transporter whose expression is regulated by the War1p zinc-finger transcription factor (Mollapour *et al.*, 2008; Holyoak *et al.*, 1999). On the other hand, adaptation to acetic acid that is more hydrophilic than sorbic or benzoic acid involves the attenuation of influx of acetic acid into the cell through degradation of acetic acid uptake channel Fps1 on the plasma membrane (Mollapour *et al.*, 2008; Mollapour and Piper, 2007). Although in chapter 2 I described that enhancement of lactic-acid resistance contributes to improvements in lactic-acid productivity without neutralization, the detailed mechanism of adaptation response to lactic-acid stress in *S. cerevisiae* remains unclear, and factors and mechanisms involved in this response are not fully elucidated.

To elucidate the cellular mechanisms which are injured by lactic-acid, I performed functional genomic analysis. I identified 107 genes that when disrupted reduced the resistance of yeast cells to 4% lactic acid. Of these 107 genes, 34 (>30% in my analysis) had not been mentioned in previous studies and were newly

identified to be involved in mechanisms for adaptation response to lactic acid. Intracellular transport including vacuolar transport and amino acid metabolism were suggested to be involved in adaptation to lactic-acid stress. The analysis of this chapter revealed that vacuole of wild-type strain fragments under lactic- and hydrochloric-acid conditions and the amount of intracellular amino acids significantly reduced by addition of lactic acid. These data suggest for the first time that lactic-acid stress impairs the vacuole integrity and intracellular amino acids are related to the growth under lactic-acid stress.

3.2 Materials and Methods

3.2.1 Strains, plasmids, and media

The nonessential genes deletion collection (OpenBiosystems, Huntsville, AL, USA), which contains 4828 yeast strains with all nonessential ORFs disrupted by the *kanMX4* cassette (http://www-sequence.stanford.edu/group/yeast_deletion_ project/deletions3.html), was used for screening in this chapter. A parental strain of the deletion collection, BY4742, (*MAT* α *leu2* Δ 0 *his3* Δ 1 *ura3* Δ 0 *lys2* Δ 0) (Brachmann *et al.*, 1998) was used as a wild-type strain in this chapter. The standard complete medium, YPDA, consisted of 1% (weight per volume, w/v) yeast extract, 2% Bacto-peptone, 2% glucose, and 0.04% adenine. L-Lactic acid (4%) or hydrochloric acid (0.13% and 0.3%) was added to YPDA after autoclaving. The pH of both the 4% lactic-acid and the 0.13% hydrochloric-acid media was 2.8. Agar (2%) was added to media to make culture plates.

3.2.2 Screening for L-lactic acid hypersensitive mutants

To screen for L-lactic-acid-hypersensitive strains, the nonessential genes deletion collection was replica stamped onto YPDA and 4% L-lactic acid plates with a 96-prong replicator. Growth of the disruptants was scored visually by comparison with that of the parent strain (BY4742), which was stamped similarly on YPDA and 4% L-lactic acid plates. Before screening, single-gene disruptants were tested for their growth on YPDA medium containing 300 mg/l G418 disulfate (Wako Pure Chemical Industries, Osaka). The L-lactic-acid-sensitive phenotype of all of the disruptants identified in the screen was subsequently confirmed to be due to a single disruption, as indicated by tetrad analysis of 6-10 asci, which showed 2 lactic-acid-resistance/2 lactic-acid-sensitive segregation and cosegregation with the disruption marker (*kanMX4*). After screening, the sensitivity of the disruptants to 1% (pH 3.7), 2% (pH 3.0), 3% (pH 2.9), and 4% (pH 2.8) lactic acid and 0.3% hydrochloric acid (pH 2.3) was tested and scored.

3.2.3 Spot dilution growth assays

Yeast cells were grown in YPDA at 30 °C to mid-log phase ($OD_{660} = 1.0$). Cells were harvested by centrifugation, washed once with water, and resuspended in water. The cell density was normalized to 1×10^6 cells/5 µl. A 10-fold serial dilution of this culture was made, and 5 µl of each dilution was spotted onto YPDA medium, 4% lactic-acid medium, or 4% lactic-acid medium supplemented with amino acids (10 mM Ile, 50 mM Tyr, 50 mM Thr, 50 mM Gly, or 50 mM Leu). Each amino acid-supplemented medium was adjusted to pH 2.8 with hydrochloric acid. Cells were incubated at 30 °C and scored visually by comparison with that of parent strain (BY4742).

3.2.4 Active staining of the vacuole with quinacrine

In vivo quinacrine staining of the vacuole was performed by a slightly modified method from Phelan *et al.* (2006). Yeast cells were grown in YPDA at 30 °C to mid-log phase ($OD_{660} = 1.0$), and 1 ml of the sample was collected. The cell pellet was resuspended in 1 ml HEPES-glucose buffer [100 mM HEPES/KOH (pH 7.5), 3% (w/v) glucose] containing 200 μ M quinacrine (Sigma-Aldrich, St. Louis, MO, USA), and incubated for 5 min in the dark at room temperature. Cells were then washed twice with HEPES-glucose buffer. After centrifugation, the cell pellet was resuspended in HEPES-glucose buffer and photographed by fluorescence microscopy (Olympus BX61; Olympus, Tokyo) within 1 h after quinacrine staining. To quantify quinacrine staining, a culture grown in YPDA was divided into two, and the quinacrine fluorescence in 1 ml of cell suspension processed as described above was measured directly using a spectrofluorometer (Hitachi F-2500; Hitachi, Tokyo), while the remainder of the cells was collected and measured for optical density. Relative fluorescence was then determined as total fluorescence per number of cells.

3.2.5 Labeling the vacuolar membrane with FM4-64

FM4-64 [*N*-(3-triethylammoniumpropyl)-4-(ρ-diethylaminophenyl-hexatrienyl)

pyridinium dibromide; Invitrogen, Carlsbad, CA, USA] labeling was performed by a slightly modified method described by Conibear *et al.* (2002). Yeast cells were grown in YPDA at 30 °C to mid-log phase (OD_{660} = 0.8), and 1 ml of the sample was collected. The cell pellet was resuspended in 100 µl YPDA. FM4-64 was added to final concentration of 40 µM [from a 16 mM stock in dimethyl sulfoxide (DMSO)], and cells were incubated for 15 min in the dark at room temperature. Cells were then washed once with YPDA. After centrifugation, the cell pellet was resuspended in YPDA and incubated for 30 min in the dark at 30 °C. Finally, cells were collected, resuspended in YPDA, and photographed by fluorescence microscopy (Olympus BX61). In the lactic-acid or hydrochloric-acid condition, cells were similarly incubated for 30 min in the dark at 30 °C, transferred to 4% lactic acid or 0.13% hydrochloric acid medium, and then photographed over time.

3.2.6 Quantification of intracellular amino acids

In the non-stressed condition, yeast cells were grown to mid-log phase (OD_{660} = 1.0) in YPDA and then collected. In the lactic-acid condition, yeast cells were grown to mid-log phase in YPDA, collected and then incubated for 2 h or 24 h in YPDA medium supplemented with 4% lactic acid. Collected cells were suspended in 0.5 N perchloric acid and disrupted with glass beads. The homogenate was kept on ice for 10 min, and then the supernatant was collected by centrifuging at 13,200 rpm for 15 min. The sample was adjusted to pH 2.2 with lithium hydroxide and was kept on ice for 10 min. Next, to remove the precipitate, the sample was centrifuged at 13,200 rpm for 15 min. Free-form amino acids were quantified by

an amino acid analyzer (Hitachi L-8500; Hitachi, Tokyo). To normalize the content of amino acids, total protein was extracted by the TCA method and determined with a Quant-iTTM Protein Assay Kit (Invitrogen, Carlsbad, CA, USA).

3.2.7 Cell viability

Cell viability was determined by exclusion of the vital dye trypan blue, based on previously published protocols (Xu *et al.*, 1999). Mid-log phase cultures were centrifuged at 3300 rpm for 5 min at room temperature, re-suspended in YPDA with 4% L-lactic acid, and incubated with aeration at 30 °C. Samples of each culture were taken at 0, 24 and 48 hours of induction. Samples were once washed with phosphate buffered saline (PBS), mixed with equal volumes of 0.4% trypan blue (Sigma-Aldrich, St. Louis, MO, USA) in PBS, incubated at room temperature for 15 min and washed two times with PBS. The numbers of dead (blue) and live (no color) cells were determined by counting using microscope.

3.3 Results

3.3.1 A genome-wide screening for lactic-acid-hypersensitive alleles

To gain more insight into the adaptation response to lactic-acid stress, I systematically screened the *S. cerevisiae* haploid collection for single disruptants that were hypersensitive to L-lactic acid stress. I used YPDA medium with 4% L-lactic acid (pH 2.8) as the screening medium because a concentration of at least 5% L-lactic acid in culture media (pH 2.7) seems to exert strong toxic effects on

yeast (Ishida *et al.*, 2005). Cell suspensions of freshly grown cultures of all of the deletion mutants (4828 strains) were transferred to YPDA plates containing 0% and 4% L-lactic acid media by using a 96-prong replicator and incubated for 1 day (0%) or 2 days (4%), and growth was assessed visually. In these conditions, the 4% lactic acid medium caused growth retardation (of about 16 h) of the wild-type strain as compared to the 0% medium. The screen was carried out in triplicate, and all lactic-acid-sensitive disruptants identified in the screening were subsequently confirmed by tetrad analysis (data not shown). Disruption alleles were also structurally confirmed by PCR (data not shown).

As a result of screening, 107 genes (2.2% of the total genes) were identified to lead to a hypersensitive phenotype to L-lactic acid when deleted (Table 4). Of the 107 genes, 34 (> 30%) had not been mentioned in a previous study (Kawahata et al., 2006). For example, I found for the first time that the depletion of Uba4 that is involved in protein urmylation (Goehring et al., 2003) caused lactic-acid hypersensitivity, suggesting that protein urmylation appears to be associated with mechanisms underlying adaptation to lactic-acid stress. A N-terminal acetyltransferase, Nat3 (Caesar et al., 2006), has also never been reported to be involved in lactic acid adaptation mechanisms. I also found that disruptions of 6 genes (RNY1, YEL059w, GUP1, RIC1, YKR041w, and MRF1) reported by Kawahata et al. (2006) did not cosegregate with the 4% L-lactic-acid-sensitive phenotype as judged by the disruption marker in tetrad analysis. Therefore, those disruptants In addition, I evaluated degree of the were excluded for further analysis. sensitivity of the 107 disruptants to 1% (pH 3.7), 2% (pH 3.0), 3% (pH 2.9), and 4%

41

(pH 2.8) lactic acid (Table 4). I examined the sensitivity of 107 disruptants to 0.5% acetic acid and 0.3% hydrochloric acid. Nine disruptants showed hydrochloric-acid sensitivity and 21 disruptants showed acetic acid sensitivity. Seven disruptants ($\Delta clc1$, $\Delta vps15$, $\Delta vps34$, $\Delta ppa1$, $\Delta hsp150$, $\Delta ypr022c$, $\Delta ecm3$) did not show the sensitivity to both acetic acid and hydrochloric acid (data not shown), indicating that these 7 genes play a role for the response to lactic-acid anion. *PMA1* that should be important for adaptation to acidic condition (Viegas *et al.*, 1998) is an essential gene for yeast growth, thereby was not employed in my screening. In my screening, disruption of *WAR1*, *PDR12*, *AFT1*, or *HAA1*, which has been reported to be involved in weak-acid stress responses (Mollapour *et al.*, 2008; Kawahata *et al.*, 2006; Abbott *et al.*, 2008; Gregory *et al.*, 2008; Hatzixanthis *et al.*, 2003) did not cause sensitivity to at least 4% lactic acid stress.

| ORF | Description | Lactic acid |
|---------------|---|-------------|
| <u>ADA2</u> | General transcriptional adaptor or co-activator | |
| <u>ALF1</u> | Protein implicated in folding of alpha tubulin | |
| <u>ANP1</u> | Protein required for protein glycosylation in the golgi | - |
| <u>ARG82</u> | Arginine metabolism transcription factor | - |
| ARO1 | Arom pentafunctional enzyme | |
| ARO2 | Chorismate synthase | |
| <u>ARO7</u> | Chorismate mutase | |
| BAP2 | Broad-specificity amino-acid permease | |
| BCKI | Ser/Thr protein kinase of the MEKK family | |
| BRP1 | Questionable protein | |
| BSTI | Negative regulator of COPII vesicle formation | |
| <u>BUD16</u> | Protein involved in bud-site selection | - |
| <u>BUD19</u> | Questionable protein | |
| <u>BUD20</u> | Protein involved in bud-site selection | |
| BUD23 | Protein involved in bud-site selection | |
| BUD32 | Protein involved in bud-site selection, putative O-sialoglyco | |
| | -endopeptidase | |
| · <u>CAX4</u> | Protein required for full levels of dolichol-linked oligosaccharides in the | - |
| | endoplasmic reticulum | |
| CCH1 | Calcium channel (α subunit) of the plasma membrane | - |
| CHC1 | Clathrin heavy chain | |
| CLC1 | Clathrin light chain | |
| CNBI | Calcineurin B, regulatory subunit | - |
| <u>COG1</u> | Conserved oligomeric golgi complex | - |
| CSG2 | Calcium dependent regulatory protein | |
| DEF1 | Coordinates repair and RNA pol II proteolysis in response to DNA | |
| | damage | |
| ЕСМ3 | Protein involved in cell wall biogenesis and architecture | |
| <u>EFG1</u> | Essential protein required for maturation of 18S rRNA | |
| ERG2 | C-8 sterol isomerase | - |
| ERG3 | C-5 sterol desaturase | |
| ERG4 | Sterol C-24 reductase | |

Table 4. Genes whose loss results in sensitivity to 4% L-lactic acid.

| ERG6 | S-adenosyl-methionine delta-24-sterol-c-methyltransferase | | | | | | | |
|--------------|---|---|--|--|--|--|--|--|
| ERG24 | C-14 sterol reductase | | | | | | | |
| FYV8 | Protein required for survival upon exposure to K1 killer toxin - | | | | | | | |
| <u>GAL11</u> | DNA-directed RNA polymerase II holoenzyme and Kornberg's | | | | | | | |
| | mediator subcomplex subunit | | | | | | | |
| GASI | Glycophospholipid-anchored surface glycoprotein | - | | | | | | |
| GLYI | L-threonine aldolase | | | | | | | |
| HFI1 | Transcriptional coactivator | | | | | | | |
| HOG1 | Ser/Thr protein kinase of MAP kinase family | - | | | | | | |
| HSP150 | Member of the Pir1/Hsp150/Pir3 family | - | | | | | | |
| ILV1 | Anabolic serine and threonine dehydratase precursor | | | | | | | |
| <u>KEMI</u> | Multifunctional nuclease | | | | | | | |
| LEM3 | Membrane protein of the plasma membrane and ER | | | | | | | |
| <u>MDJ1</u> | Heat shock protein-chaperone | | | | | | | |
| MID1 | Mechanosensitive Ca ²⁺ -permeable channel of the plasma membrane | - | | | | | | |
| MON2 | Peripheral membrane protein with a role in endocytosis and vacuole | | | | | | | |
| | integrity | | | | | | | |
| <u>NAT3</u> | N-acetyltransferases | - | | | | | | |
| NHX1 | Na^+/H^+ exchanger of the prevacuolar compartment | | | | | | | |
| OCH1 | α-1,6-mannosyltransferase· | | | | | | | |
| PBS2 | Tyrosine protein kinase of the MAP kinase kinase family | | | | | | | |
| <u>PDX3</u> | Pyridoxamine-phosphate oxidase | | | | | | | |
| PEP3 | Vacuolar membrane protein | | | | | | | |
| PEP5 | Vacuolar biogenesis protein | | | | | | | |
| PEP7 | Vacuolar segregation protein | | | | | | | |
| <u>PHO85</u> | Cyclin-dependent protein kinase | - | | | | | | |
| POP2 | Protein required for glucose derepression | | | | | | | |
| PPA1 | H^+ -ATPase V_0 domain | | | | | | | |
| RAD6 | E2 ubiquitin-conjugating enzyme | | | | | | | |
| RCYI | Protein involved in recycling of the SNARE Snc1p | | | | | | | |
| RGD1 | GTPase activating protein (GAP) (putative) | - | | | | | | |
| RGP1 | Reduced growth phenotype protein | | | | | | | |
| RPB9 | DNA-directed RNA polymerase II | | | | | | | |
| <u>RPL1B</u> | 60S large subunit ribosomal protein | | | | | | | |

•

| <u>RRG1</u> | Protein of unknown function localized to mitochondria, required for | |
|--------------|---|---|
| | vacuolar acidification | |
| <u>RTF1</u> | Regulates DNA binding properties of TBP | |
| SAC1 | Recessive suppressor of secretory defect | |
| <u>SAP30</u> | Subunit of the histone deacetylase B complex | - |
| <u>SFP1</u> | Zinc finger protein | |
| SNF5 | Component of SWI/SNF transcription activator complex | |
| SNF12 | Component of SWI/SNF global transcription activator complex | |
| SHE4 | Protein required for mother cell-specific gene expression | |
| <u>SHP1</u> | Potential regulatory subunit for Glc7 | |
| SLA2 | Cytoskeleton assembly control protein | |
| SLT2 | Ser/Thr protein kinase of MAP kinase family | |
| <u>SPT20</u> | Member of the TBP class of SPT proteins that alter transcription site | |
| | selection | |
| SRV2 | Adenylate cyclase-associated protein | |
| SWI3 | Transcription regulatory protein | |
| <u>TFP3</u> | H^+ -ATPase V_0 domain | - |
| THRI | Homoserine kinase | |
| THR4 | Threonine synthase (o-p-homoserine p-lyase) | - |
| TLG2 | Member of the syntaxin family of t-SNAREs | - |
| TRK1 | Potassium transporter I | |
| TYRI | Prephenate dehydrogenase | |
| <u>UBA4</u> | Protein that activates Urm1 before its conjugation to proteins | - |
| VAM10 | Protein required for normal tethering of vacuoles prior to fusion | - |
| <u>VMA1</u> | Encodes 3 region protein which is self-spliced into Tfp1 and PI-SceI | |
| VMA4 | H ⁺ -ATPase V ₁ domain | - |
| VMA5 | H ⁺ -ATPase V ₁ domain | - |
| VMA6 | H^+ -ATPase V_0 domain | - |
| VMA7 | H ⁺ -ATPase V ₁ domain | |
| <u>VMA8</u> | H^+ -ATPase V_1 domain | - |
| <u>VMA10</u> | H^+ -ATPase V_1 domain | |
| <u>VMA22</u> | Vacuolar ATPase assembly protein | |
| VPS1 | Member of the dynamin family of GTPases | |
| VPS3 | Vacuolar sorting protein | |
| VPS5 | Protein involved in Golgi retention and vacuolar sorting | |
| | | |

.

| VPS15 | Ser/Thr protein kinase | | | | | | | |
|---|---|---|--|--|--|--|--|--|
| VPS16 | Vacuolar sorting protein | | | | | | | |
| VPS20 | Vacuolar protein sorting | | | | | | | |
| VPS34 | Phosphatidylinositol 3-kinase | | | | | | | |
| VPS35 | Protein-sorting protein | | | | | | | |
| VPS45 | Vacuolar protein sorting-associated protein | | | | | | | |
| VPS51 | Subunit of VP51-54 complex, required for protein sorting at the | | | | | | | |
| | yeast late Golgi | | | | | | | |
| <u>VPS52</u> | Subunit of VP51-54 complex, required for protein sorting at the | | | | | | | |
| | yeast late Golgi | | | | | | | |
| <u>VPS54</u> | Subunit of VP51-54 complex, required for protein sorting at the | | | | | | | |
| | yeast late Golgi | | | | | | | |
| <u>VRP1</u> | Verprolin | | | | | | | |
| YCL007c | Questionable protein | | | | | | | |
| <u>YPR022c</u> Protein of unknown function localized to cytoplasm and nucleus | | | | | | | | |
| YPT6 | GTP-binding protein of the Rab family | - | | | | | | |

Underline represents newly identified genes in this chapter whose disruption causes lactic acid hypersensitivity. Others were also identified in Kawahata *et al.* (2006). Scores for lactic acid sensitivity were assigned as follows: "- - - -", "- - -", and "-" represent hypersensitivity to >1%, to >2%, to >3%, and to >4% lactic acid when compared with wild type, respectively.

I initially classified the 107 genes into functional categories using Munich Information Centre for Protein Sequence (MIPS) database with FunSpec program (Robinson *et al.*, 2002). These classifications are useful for understanding which biological functions are important for adaptation responses to lactic acid. Table 5 lists the representative functional categories of genes whose deletions caused lactic acid sensitivity (p < 0.01). A variety of cellular functions were suggested to be involved in the adaptation response to lactic acid. The p value is a good statistical indicator of overrepresentation of genes in a given functional category (Robinson *et al.*, 2002). The category associated with vacuolar transport containing 23 genes (VPS15, PEP7, NHX1, VPS3, VMA8, VPS45, CHC1, VMA7, CLC1, VMA10, VPS35, VMA5, VPS1, VPS51, PEP3, VPS34, VMA6, VPS20, PEP5, MON2, TLG2, VMA4, and VPS16) exhibited lowest p values (Table 5), suggesting that many cellular functions, especially functions associated with vacuolar transport play important roles in adaptation response under lactic acid stress condition.

| to 4% L-lactic acid stress. | | | |
|---|---|--------------------------|-----------------------|
| MIPS functional category ^a | No. of genes identified in this chapter | No. of total genes | p-value ^b |
| Vacuolar transport [20.09.13] | 23 | 153 | $< 1 \times 10^{-14}$ |
| Cation transport (H ⁺ , Na ⁺ , K ⁺ , Ca ²⁺ , NH ₄ ⁺ , etc.) [20.01.01.01] | 10 | 68 | 8.3×10^{-8} |
| Vesicular transport (Golgi network, etc.) [20.09.07] | 10 | 72 | 1.4×10^{-7} |
| Homeostasis of protons [34.01.01.03] | 8 | 47 | 5.5×10^{-7} |
| Protein targeting, sorting and translocation [14.04] | 15 | 281 | 3.0×10^{-5} |
| Vacuole [42.25] | 6 | 44 | 5.7×10^{-5} |
| Intra Golgi transport [20.09.07.05] | 5 | 33 | 1.5×10^{-4} |
| Transport ATPases [20.03.22] | 6 | 53 | 1.7×10^{-4} |
| Osmosensing and response [34.11.03.13] | 5 | 35 | 2.0×10^{-4} |
| Metabolism of tryptophan [01.01.09.06] | 3 | 8 | 2.0×10^{-4} |
| Tetracyclic and pentacyclic triterpenes metabolism [01.06.06.11] | 5 | 36 | 2.3×10^{-4} |
| Electron transport [20.01.15] | 7 | 83 | 3.0×10^{-4} |
| Budding, cell polarity and filament formation [43.01.03.05] | 14 | 312 | 3.5×10^{-4} |
| Cellular import [20.09.18] | 7 | 90 | 5.0×10^{-4} |
| Biosynthesis of threonine [01.01.06.04.01] | 2 | 3 | 7.3×10^{-4} |
| MAPKKK cascade [30.01.05.01.03] | 4 | 27 | 7.7×10^{-4} |
| Metabolism of phenylalanine [01.01.09.04] | 3 | 13 | 9.7×10^{-4} |
| Metabolism of tyrosine [01.01.09.05] | 3 | 14 | 1.2×10^{-3} |
| Phosphate metabolism [01.04] | 15 | 401 | 1.4×10^{-3} |
| Degradation of threonine [01.01.06.04.02] | 2 | 4 | 1.4×10^{-3} |
| Exocytosis [20.09.16.09.03] | 4 | 33 | 1.7×10^{-3} |

Table 5. Representative functional categories of genes whose disruption causes sensitivity to 4% L-lactic acid stress.

| Modification by phosphorylation, dephosphorylation, | 9 | 186 | 2.5×10^{-3} | | | | | | |
|---|----|-----|----------------------|--|--|--|--|--|--|
| autophosphorylation [14.07.03] | | | | | | | | | |
| Transcriptional control [11.02.03.04] | 15 | 426 | 2.6×10^{-3} | | | | | | |
| Pheromone response, mating-type determination, sex-specific | 9 | 189 | 2.8×10^{-3} | | | | | | |
| proteins [34.11.03.07] | | | | | | | | | |
| Homeostasis of cations [34.01.01] | 3 | 19 | 3.1×10^{-3} | | | | | | |
| Protein binding [16.01] | 14 | 391 | 3.1×10^{-3} | | | | | | |
| Polyphosphoinositol mediated signal transduction [30.01.09.11] | 3 | 20 | 3.6×10^{-3} | | | | | | |
| Regulation of C-compound and carbohydrate metabolism [01.05.25] | 7 | 126 | 3.6×10^{-3} | | | | | | |
| Actin cytoskeleton [42.04.03] | 6 | 96 | 3.9×10^{-3} | | | | | | |
| Modification by acetylation, deacetylation [14.07.04] | 5 | 69 | 4.5×10^{-3} | | | | | | |
| Cell wall [42.01] | 9 | 213 | 6.2×10^{-3} | | | | | | |
| Metabolism of serine [01.01.09.02] | 2 | 8 | 6.5×10^{-3} | | | | | | |

^a Number in parenthesis is MIPS functional category number.

^b p-values represent the probability that the intersection of a given list with any given category occurs by chance (Robinson *et al.*, 2002).

3.3.2 Almost half of the lactic-acid-sensitive disruptants show vacuolar acidification defect.

Many disruptants for genes involved in the vacuolar transport such as VPS genes are known to show defects of vacuole (Rothman and stevens, 1986; Rothman *et al.*, 1989; Seeley *et al.*, 2002). The yeast vacuole, an acidic compartment responsible for the degradation of cellular components, is essential for many cellular processes including endocytosis, pH homeostasis, metabolite storage, and sequestration of toxic substances such as excess metals (Corson *et al.*, 1999; Klionsky *et al.*, 1990; Kane, 2006). Vacuolar acidification is important for stress-related functions. Strains with a defect in acidification of the vacuole show a growth defect in medium of inadequate pH (Nelson and Nelson, 1990). Vacuolar acidification activity is also essential for efficient cytosolic pH homeostasis that plays important roles in expressing proper activity of many cellular proteins

(Martínez-Muñoz and Kane, 2008). Furthermore, vacuolar acidification defect causes acutely sensitivity to a wide variety of heavy metals (Corson et al., 1999) and hypersensitivity to many different stresses including ethanol (Teixeira et al., 2009), acids (Kawahata et al., 2006), and air-drying (Shima et al., 2008). When the 107 genes were classified into yeast biological process categories using the FunSpec program (Robinson et al., 2002), the category associated with vacuolar acidification exhibited lowest p value (1.2×10^{-10}) (data not shown). In addition. 18 of the 23 lactic-acid-sensitive disruptants for genes classified in vacuolar transport category were found to exhibit abnormal acidification of vacuole even in the absence of lactic acid by staining with quinacrine which is an indicator of the acidic interior of a vacuole (Conibear and Stevens, 2002) (Table 6). Because these data suggested importance of maintenance of vacuolar functions especially vacuolar acidification for adaptation for lactic-acid stress, I next examined the presence of vacuolar acidification defects in all of the lactic-acid-sensitive disruptants with quinacrine.

| Acidification | Morphology | Genes |
|---------------|------------|--|
| Normal | Fragmented | VAM10, CCH1, SLA2, MON2, VPS35, VPS5, BST1, SAP30, CHC1 |
| Decreased | Normal | ECM3, BUD19, BUD23, RGD1, MID1, VMA8, VPS1, COG1, CSG2, PDX3, CAX4, CNB1, SWI3, SFP1, VPS20, NHX1, VPS3, VPS34, SRV2 |
| Decreased | Fragmented | OCH1, YPT6, ALF1, BUD32, VPS54, VRP1 |
| Lost | Normal | PPA1, VPS45, PEP5, SHP1, KEM1, VPS15, VMA22, PEP3, VMA5, VMA7, VMA6, VMA1, VPS16, PEP7, VPS52, VMA10, VMA4, RRG1 |
| Lost | Fragmented | TFP3, VPS51, ANP1, MDJ1, RPL1B |

Table 6. Vacuolar acidification and morphology of L-lactic acid sensitive disruptants under lactic-acid free condition.

Interestingly, I found that 48 of the lactic-acid-sensitive disruptants (45% of the total) exhibited abnormal acidification of vacuole, even in the absence of lactic-acid stress (Table 6). In this analysis, 20 genes (ECM3, BUD19, BUD23, RGD1, MID1, COG1, CSG2, PDX3, CAX4, CNB1, SWI3, SRV2, SFP1, OCH1, ALF1, BUD32, KEM1, ANP1, MDJ1, and RPL1B; Table 6) were newly found to be involved in vacuole acidification by confirming that these abnormal vacuolar phenotypes were indeed caused by disruption of the corresponding genes by tetrad analysis (data not shown). For example, the vacuole of the $\Delta kem1$ and $\Delta mdj1$ disruptants did not show positive quinacrine staining, and the quinacrine fluorescence intensity of the $\Delta cnb1$ and $\Delta och1$ disruptants was significantly reduced as compared with the parental strain, indicating that these disruptants failed to acidify the vacuole properly (Fig. 6A). The decrease in fluorescence intensity of quinacrine in these disruptants was also confirmed by using a spectrofluorometer (Fig. 6B). I also found that the vacuole stained with FM4-64, which was used for visualization of the vacuolar membrane (Conibear and Stevens, 2002), of the $\Delta och1$ and $\Delta m d i l$ disruptants was fragmented (Fig. 6A). Because vacuolar acidification is important for adaptation to many stress conditions (Phelan et al., 2006; Corson et al., 1999; Teixeira et al., 2009), it may be suggested that these disruptants displaying vacuolar acidification defect even under lactic acid free condition showed sensitivity to lactic acid due to its vacuolar acidification defect. Moreover. disruptants showing defects in vacuolar acidification may be suggested to have some defects in intracellular pH homeostasis which is important to maintain the activity of intracellular proteins because proper acidification of vacuole was

essential for stable localization of Pma1, an essential proton pump, at the plasma membrane (Martínez-Muñoz and Kane, 2008; Perzov *et al.*, 2000; Hirata and Takatsuki, 2001).



Figure 6. Vacuolar acidification and morphology of lactic-acid-sensitive disruptants. (A) Quinacrine and FM4-64 staining. BY4742 (WT) and lactic-acid-sensitive disruptants ($\Delta cnb1$, $\Delta mdj1$, $\Delta kem1$, and $\Delta och1$) were stained as described in Materials and methods, and examined by differential interference contrast (DIC) and fluorescence microscopy. A decrease in fluorescence intensity indicates defective vacuolar acidification. (B) Quantification of quinacrine fluorescence intensity. Fluorescence intensity values were calculated as described in Materials and methods and were expressed relative to those of the BY4742 (WT) strain (defined as 100%); error bars show the standard deviation of three independent experiments.

3.3.3 Vacuole is subjected to fragmentation by lactic-acid stress

I next examined vacuolar integrity from the view point of vacuolar morphology during adaptation period to lactic-acid stress in the wild-type strain

since fragmentation of the vacuole also causes increased sensitivities against low and high pH (Corson *et al.*, 1999) and ethanol and heat shock (Matsuura and Takagi, 2005). When I observed the vacuolar morphology of the wild-type strain with FM4-64 under the 4% lactic-acid condition in a time-dependent manner, I found that the vacuole became markedly smaller and heavily fragmented as the exposure time to 4% lactic acid increased (Fig. 7A). In YPDA medium without lactic acid, the wild-type strain did not show vacuole fragmentation (Fig. 7B). Although I was not able to observe vacuolar morphology with FM4-64 staining by unknown reason when observation was extended to 24 h where wild-type strain restarted to grow (Fig. 8; see below), I was able to stain vacuole of a portion of 24 h cells with quinacrine and found that at least 53% of cells had vacuole as a single round structure like normal morphology (data not shown), indicating that vacuolar acidification and fragmentation were becoming normal after 24 h incubation in lactic-acid medium. These results indicate that vacuole of yeast cells is subjected to fragmentation by exposure to lactic acid but the fragmented vacuole is gradually being restored to normal state through adaptation to lactic acid.

Osmotic stress (0.4 M NaCl or 0.5 M sorbitol) is also reported to induce vacuolar fragmentation (Bonangelino *et al.*, 2002). This response may serve to regulate the surface area to volume ratio of the vacuole by changing the number and size of the vacuole lobes under conditions where the vacuole loses volume due to loss of water (Bonangelino *et al.*, 2002). Thus, it is possible that the vacuolar fragmentation caused by 4% lactic acid (0.44 M) might be due to osmotic stress. Therefore, I tested whether another acid stress that has microbial growth inhibitory

effects at a lower concentration than lactic acid also causes vacuole fragmentation. The results revealed that 0.13% (35 mM) hydrochloric acid (pH 2.8) stress that causes similar growth retardation to 4% lactic acid stress (data not shown) also causes vacuolar fragmentation (Fig. 7C). Although I cannot fully rule out the possibility that some of the effects of 4% lactic acid are due to osmostress, these results highly suggest that vacuolar fragmentation seems to be a common mechanism underlying adaptation response to acid stress.



Figure 7. Time-dependent change in vacuolar morphology caused by acid stress. BY4742 cells were stained with FM4-64 as described in Materials and methods, transferred to 4% lactic acid (A), fresh YPDA (B), and 0.13% hydrochloric acid (C) medium (time = 0 min), and then examined by differential interference contrast (upper panel) and fluorescence microscopy (lower panel) for 120 min.

3.3.4 Amount of intracellular amino acid is important for the growth under lactic-acid stress

To facilitate my understanding of mechanisms involved in adaptation to lactic-acid stress in S. cerevisiae, I next examined the involvement of intracellular amino acid homeostasis in the adaptation because the category associated with tryptophan, threonine, phenylalanine, tyrosine, and serine metabolisms also exhibited extremely low p values ($p = 2.0 \times 10^{-4}$, 7.3×10^{-4} , 9.7×10^{-4} , 1.2×10^{-3} , 6.5×10^{-3} for Metabolism of tryptophan, Biosynthesis of threonine, Metabolism of phenylalanine, Metabolism of tyrosine, and Metabolism of serine, respectively, according to MIPS functional category) (Table 5) and genes involved in amino-acid biosynthesis and uptake formed the largest group (18%, 9 disruptants, $\Delta glyl$, $\Delta ilvl$, $\Delta aro1$, $\Delta aro2$, $\Delta aro7$, $\Delta thr1$, $\Delta thr4$, $\Delta tyr1$, and $\Delta bap2$) in the remaining 50 lactic-acid-sensitive disruptants which did not exhibit abnormal acidification and/or morphology in Table 4. Gly1 is involved in glycine biosynthesis; thus, $\Delta gly1$ shows a glycine auxotrophic phenotype (McNeil et al., 1994). Similarly, Ilv1 catalyzes the first step in isoleucine biosynthesis; thus, $\Delta i l v l$ displays isoleucine and valine auxotrophy (Holmberg and Petersen, 1988). Aro1, Aro2, Aro7, and Tyr1 proteins are essential for the biosynthesis of aromatic amino acids such as phenylalanine, tryptophan, and tyrosine (Jones and Fink, 1982; Mannhaupt et al., 1989). Thr1 and Thr4 proteins are required for threonine biosynthesis (Ramos and Calderon, 1994). BAP2 encodes a plasma membrane permease for a branched-chain amino acid (leucine, isoleucine, and valine).

To see the relationship between intracellular amount of amino-acid and growth retardation under lactic-acid stress condition, growth of wild-type strain under lactic-acid condition was examined (Fig. 8). After addition of 4% lactic-acid stress, the growth of wild-type cells seemed to be hampered within 2 h, whereas the cells began to grow slowly after 12 h incubation and to grow more robustly after 24 h incubation (Fig. 8). I also examined the cell viability of wild-type strain after 2 h and 24 h incubation. Results showed that the cell viability decreased to 23.8% ± 4.2 (2 h incubation), whereas it recovered to 44.7% ± 4.1 (24 h incubation) compared with 0 h incubation. This suggests that the cell growth increased as the cell viability recovered.

I next examined intracellular amino-acid homeostasis under lactic-acid stress condition. Cells of wild-type strain were collected after 0 h, 2 h, and 24 h of incubation in 4% L-lactic acid medium and subjected to measurement of the content of free-form amino acids. The results revealed that the amount of intracellular amino acids markedly decreased after 2 h of incubation with lactic acid (Table 7), a time point at which the growth of wild-type strain seemed to be almost hampered (Fig. 8). For example, the intracellular amount of lysine, which showed the greatest decrease, was reduced to 1% as compared with 0 h, and that of phenylalanine, which showed the least decrease, was reduced to 13.2% as compared with 0 h. I also checked the intracellular amount of amino acids after 24 h of incubation and found that the total amount of amino acids was restored about 3.7-fold as compared with 2 h (Table 7). These results indicate that the cell viability (23.8% \pm 3.1 for 2 h incubation and 44.7% \pm 8.1 for 24 h incubation compared with 0 h incubation) showed a link with the intracellular amino acid concentration during the adaptation period to lactic-acid stress. The disruptant of

55

BAP2, which encodes a high-affinity branched-chain amino acid permease and causes lactic-acid sensitivity when deleted, displayed a much smaller amount of intracellular amino acids including not only branched-chain amino acid but also other amino acids after 0 h, 2 h, and 24 h of incubation than did the wild-type strain (Table 7). Importantly, the growth of the $\Delta bap2$ disruptant was still suppressed even in the 24 h of incubation under lactic-acid stress condition (Fig. 8) and the cell viability of the $\Delta bap2$ disruptant was also much less than that of the wild-type strain (0.4% ± 0.03 for 2 h incubation and 1.8% ±0.11 for 24 h incubation compared with 0 h incubation). These results indicate that intracellular amino acids reduce by addition of lactic acid, suggesting that the amount of intracellular amino acid is important for the growth under lactic-acid stress.



Figure 8. Effect of lactic-acid stress on cell growth. BY4742 (WT; open circles) and $\Delta bap2$ (closed square) strains were grown in YPDA at 30 °C to mid-log phase (OD₆₆₀ = 1.0). Cells were harvested by centrifugation and resuspended in 4% lactic-acid medium, and then the OD₆₆₀ of the culture was monitored periodically for 24 h.

| Amino acid | BY4742 | | | Δ <i>bap</i> 2 | | | |
|--------------------|------------------|----------------------|-----------------------|------------------|----------------------|-----------------------|--|
| | – Lactic acid | + Lactic acid 2 h | + Lactic acid 24 h | – Lactic acid | + Lactic acid 2 h | + Lactic acid 24 h | |
| Asp | 865.3 | 15.2 | 93.3 | 450.3 | 4.1 | 4.8 | |
| Thr | 591.3 | 11.0 | 48.9 | 263.5 | 3.2 | 4.3 | |
| Ser | 542.9 | 15.4 | 89.2 | 280.5 | 4.9 | 6.2 | |
| $AspNH_2$ | 184.5 | 8.1 | 20.2 | 77.9 | 2.6 | 3.5 | |
| Glu | 4250.4 | 82.6 | 451.5 | 2744.1 | 14.4 | 14.9 | |
| GluNH ₂ | 921.2 | 22.9 | 110.9 | 96.0 | N.D. | N.D. | |
| Gly | 861.5 | 15.3 | 56.0 | 521.8 | 5.4 | 7.0 | |
| Ala | 964.7 | 32.0 | 65.4 | 523.3 | 12.0 | 13.0 | |
| Val | 271.3 | 13.8 | 24.1 | 115.1 | 5.0 | 6.5 | |
| Met | 56.9 | 4.4 | 8.3 | 39.2 | 1.8 | 2.1 | |
| Ile | 143.6 | 10.6 | 11.2 | 70.6 | 4.5 | 5.8 | |
| Leu | 266.9 | 25.2 | 29.1 | 125.5 | 10.2 | 11.7 | |
| Tyr | 88.9 | 9.5 | 4.5 | 51.8 | 3.9 | 3.6 | |
| Phe | 151.1 | 20.0 | 15.8 | 80.7 | 8.3 | 9.0 | |
| Orn | 248.7 | 5.8 | 24.7 | 160.7 | 0.9 | 1.8 | |
| Lys | 4610.2 | 68.9 | 405.6 | 2528.0 | 12.5 | 11.4 | |
| His | 536.0 | 8.2 | 42.5 | 278.6 | 2.4 | 2.6 | |
| Arg | 3512.6 | 121.1 | 313.3 | 2031.0 | 36.0 | 32.9 | |
| · Pro | 242.3 | 5.4 | 25.3 | 90.3 | 1.3 | 2.1 | |
| Total | 19310.2 | 495.4 | 1839.9 | 10528.6 | 133.4 | 143.2 | |

Table 7. Amount of intracellular amino acids (nmol/mg protein).

N.D., not detected

3.3.5 Amino acid supplementation recovers adaptation deficiency to lactic acid To further investigate an idea that the amount of intracellular amino acid is important for the growth under lactic-acid stress, I examined whether addition of an amino acids to the media could rescue the growth defect of 9 disruptants ($\Delta gly1$, $\Delta ilv1$, $\Delta aro1$, $\Delta aro2$, $\Delta aro7$, $\Delta thr1$, $\Delta thr4$, $\Delta tyr1$, and $\Delta bap2$) under the lactic-acid stress condition. A serial dilution-spot assay revealed that amino-acid supplementation alleviated the lactic-acid sensitivity of those disruptants with a

defect in amino-acid biosynthesis and transport (Fig. 9). For example, the

lactic-acid sensitivity of the $\Delta aro1$, $\Delta aro2$, $\Delta aro7$, and $\Delta tyr1$ disruptants, which cannot synthesize Tyr, was alleviated by adding 50 mM Tyr to the 4% lactic-acid medium. Similarly, the sensitivity of disruptants requiring Ile ($\Delta i l v l$), Thr ($\Delta t h r l$ and $\Delta thr4$) or Gly ($\Delta gly1$, $\Delta thr1$, and $\Delta thr4$) was alleviated by adding 10 mM Ile, 50 mM Thr, or 50 mM Gly, respectively. The sensitivity of the $\Delta bap2$ disruptant was also alleviated by adding 50 mM Leu to the 4% lactic-acid medium. On the other hand, the addition of 10 or 50 mM Val, 10 or 50 mM Trp, or 5, 10, or 50 mM Phe did not relieve the sensitivity of disruptants requiring Val ($\Delta bap2$), Trp ($\Delta aro1$, $\Delta aro2$, and $\Delta aro7$) or Phe ($\Delta aro1$, $\Delta aro2$, $\Delta aro7$, and $\Delta tyr1$), respectively (data not Since ARO1, ARO2, ARO7, and TYR1, or BAP2 and ILV1 are involved in shown). biosynthesis of Tyr, Trp, and Phe, or transport of Leu, Ile, and Val, respectively, the growth of their disruptants was examined on 4% lactic-acid media containing Tyr (5 mM or 50 mM), Trp (5 mM or 10 mM), and Phe (1 mM or 3 mM), or Leu (50 mM or 25 mM), Ile (10 mM or 5 mM), and Val (10 mM or 5 mM), simultaneously. However, the growth of the $\Delta aro1$, $\Delta aro2$, $\Delta aro7$, $\Delta tyr1$, $\Delta bap2$, and $\Delta ilv1$ disruptants on those media was not enhanced further when compared with those on the media containing single amino acid, *i.e.*, Tyr (ARO1, ARO2, and ARO7) or Leu (BAP2 and ILV1) (data not shown). These results suggest that specific amino acids but not any amino acids are important for the growth under lactic-acid stress. Taken together, the amount of intracellular amino acids, especially for Thr, Gly, Ile, Tyr, and Leu, may be important for mitigation of the growth defect by lactic-acid stress.



YPDA + 4% lactic acid

Figure 9. Growth assay of lactic-acid-sensitive disruptants on lactic-acid medium supplemented with amino acids. A ten-fold serial dilution of each yeast strain was spotted onto YPDA medium, 4% lactic-acid medium (pH 2.8), and 4% lactic-acid medium supplemented with the indicated amino acids (50 mM Tyr, 10 mM Ile, 50 mM Thr, 50 mM Gly, or 50 mM Leu) (pH 2.8) as described in Materials and methods. Cells were incubated at 30 °C for 2 days.

3.4 Discussion

In this chapter, my analysis newly identified 34 genes (>30% in my analysis) (Table 4) that were not mentioned previously but are involved in lactic-acid resistance. For example, the depletion of Uba4 was found for the first time to cause lactic-acid hyper-sensitivity. Since Uba4 was suggested to regulate the function of the antioxidant protein Ahp1 through its urmylation in response to oxidative stress (Goehring *et al.*, 2003), it may be possible to speculate that Uba4 is also involved in the urmylation of proteins that are important for adaptation to lactic-acid stress. A N-terminal acetyltransferase, Nat3 (Caesar *et al.*, 2006), was also found to be involved in lactic acid adaptation mechanisms. Since Nat3-mediated N-terminal acetylation of proteins is involved in many physiological processes such as regulation of actin cytoskeleton or cell cycle (Caesar *et al.*, 2006), N-terminal acetylation of specific targets is also suggested to play important roles in

adaptation responses of lactic acid.

In this analysis, I found that remarkable reduction of intracellular amino acids occurred under lactic acid condition (Table 7). This may be caused by inhibition of the uptake of amino acids because amino-acid uptake across the plasma membrane is catalyzed by transporters whose activity is dependent on the proton electrochemical gradient (Seaston et al., 1973), and acid stress likely reduces this gradient. Consistent with this idea, it has been reported that sorbic acid inhibits the uptake of aromatic amino acids (Bauer et al., 2003). Disturbance of vacuolar integrity, which is suggested to be important for recycling of membrane proteins (Bugnicourt et al., 2004), may also impair the correct localization of amino-acid transporters, thereby leading to amino-acid starvation. Moreover, vacuolar fragmentation caused by lactic acid is likely to affect the intracellular amount of amino acids because the vacuole is involved in storage of amino acids, and a fragmented vacuole loses storage capacity and activity of vacuolar enzymes (Klionsky et al., 1990; Jacquemin-Faure et al., 1994; Kitamoto et al., 1988). In addition, since yeast cells can utilize peptides that are contained in much amount in YPD broth as a source of amino acids by transporting and hydrolyzing them (Perry et al., 1994; Hauser et al., 2001), my result (Table 7) suggests that decreased transport and hydrolysis activities toward peptides may also occur under lactic-acid condition. Since amino-acid starvation caused by lactic-acid stress may in turn cause a general reduction in protein synthesis, which is important for producing stress-response proteins, maintaining amino acid homeostasis plays an important role in adaptation responses against lactic-acid stress. As observed in Fig. 9, the

lactic-acid sensitivities of disruptants for genes involved in amino-acid metabolism were alleviated but partially even when excess amount of amino acids were added to the media. Since the proton electrochemical gradient is important for amino-acid uptake activity across the plasma membrane (Seaston *et al.*, 1973) and acid stress may reduce this gradient, it may be possible to expect that amino acids were not transported sufficiently into the cells, thereby I think that the disruptants did not show complete recovery.

Lactic acid sensitive disruptants having vacuolar acidification defects possibly had insufficient sequestration abilities of proton at the vacuole, thereby showing the Moreover, since proper acidification of vacuole was essential for sensitivity. stable localization of Pma1 at the plasma membrane (Martínez-Muñoz et al., 2008; Perzov et al., 2000; Hirata and Takatsuki, 2001), lactic acid sensitive disruptants showing defects in vacuolar acidification may also likely have some defects in intracellular pH homeostasis which is important to maintain the activity of intracellular proteins. It is also known that fragmentation of vacuoles causes shortage of storage capacity of amino acids (Kitamoto *et al.*, 1988). Attenuation of the uptake activity of amino acids is also imagined since amino-acid uptake across the plasma membrane is catalyzed by transporters whose activity is dependent on the proton electrochemical gradient (Seaston et al., 1973), and lactic-acid stress may reduce this gradient across the plasma membrane. Therefore, severe depletion of amino acids should be caused subsequently as observed in this chapter. In the stressed condition, maintenance of intracellular amino acid homeostasis is critical because cells have to produce a variety of stress response

proteins using intracellular amino acids. Starvation of amino acids also causes a marked accumulation of oxygen radicals (Eisler et al., 2004), thereby producing an additional stress. Moreover, as reported in bacteria, for example Escherichia coli (Foster, 2004), amino acids play important roles to maintain intracellular pH by enhancing consumption of protons through decarboxylation of glutamate or arginine under acidic condition. In addition to decreased ATP level resulting from activation of Pma1 under acidic condition (Viegas et al., 1998), these may prevent the proliferation of yeast. However, yeast cell is supposed to acquire abilities to generate ATP by progressing the phase of stress adaptation, for example, the sequestration of the proton and cell-wall reorganization that may attenuate the influx of lactic acid into the cell (Kawahata et al., 2006) and by possibly increasing pH of the medium through consumption of proton. ATP may be important for active efflux of the anion from cell by unknown transporters and for import of peptides or amino acids into cell. These effects would further facilitate the adaptation responses, thereby supporting the restart of yeast growth under lactic-acid condition.

3.5 Summary

To gain more insight into adaptation response to lactic-acid stress in yeast, a genome-wide screening for genes whose disruption cause hypersensitivity to 4% L-lactic acid (pH 2.8) was performed using the gene deletion collection of *S. cerevisiae*. I identified 107 genes that contributed significantly to the ability of yeast cells to adapt lactic-acid stress. More than 30% of the genes identified in

this screening were newly identified to be involved in mechanisms for adaptation response to lactic acid. For example, I found for the first time that protein urmylation by Uba4 and N-terminal acetylation by Nat3 were involved in lactic acid adaptation mechanisms. Functional categorization of the genes followed by microscopic analysis revealed that a variety of cellular functions were involved in adaptation response to lactic acid and function associated with vacuolar transport played important roles in adaptation response to lactic acid. I also found that vacuole fragmented immediately upon exposure to lactic- and hydrochloric-acid stress. In addition, my analysis revealed that the amount of intracellular amino acids significantly reduced by addition of lactic acid, suggesting that the amount of intracellular amino acids may be important for mitigation of the growth defect by lactic-acid stress.

Chapter 4

General discussion and conclusion

In this study, I first screened single gene disruptants which showed resistance to 6% lactic acid (Chapter 2) or hypersensitivity to 4% lactic-acid stress (Chapter 3) although Kawahata et al. (2006) reported the lactic-acid sensitive or resistant disruptants screened under different concentration of lactic acid (3.1% and 5.1%, respectively) from those (4% and 6%) used in this study. Lactic acid (5%) exerts toxic effect to yeast cells (Ishida et al., 2005) and wild-type strain showed transient growth retardation in the presence of 4% lactic acid (Chapter 3). Kawahata et al. (2006) identified 46 genes whose disruption led to resistance to 5.1% lactic acid and 128 genes whose disruption led to sensitivity to 3.1% lactic acid. This study aimed to identify novel genes involved in resistance and sensitivity against lactic acid by employing more severe conditions than previous studies. This screen identified 69 additional disruptants that were resistance to lactic acid and 34 additional disruptants sensitive to lactic acid. Because many genes involved in severe lactic-acid stress were newly identified, it is considered that understanding the function of these genes will help to elucidate the detailed response mechanisms to lactic acid in yeast.

In chapter 2, the lactic-acid productivity of the *LDH*-harboring quadruple disruptant ($\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$; SH6779), which showed the strongest lactic-acid resistance, significantly increased (approximately 27%, 48 g/litter, 72h)

64

compared with wild-type strain (SH6764; 38 g/litter, 72 h) (Fig. 5). Table 8 compares lactic-acid productivity with those reported in previous studies which were performed in batch cultures, with or without neutralization. As shown in Table 8, although *LDH*-harboring quadruple disruptant ($\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$; SH6779) did not show the highest productivity and yield among all cases, SH6779 had the highest productivity and yield of any published strain grown without neutralization (Dien *et al.*, 2001 and Gao *et al.* 2011), and despite the non-neutralization condition, SH6779 productivity was comparable to studies employing neutralization. Thus, I propose that SH6779 has great potential and its lactic-acid productivity could be additionally enhanced with other refinements.

| Organism | LDH (copy number) | Neutralization | Initial glucose | Lactic acid (g/l) | Producti vity (g/l/h) | Yield (g/g) | Disruption of EtOH pathway | Reference |
|---------------------------|--------------------------------------|-------------------|--------------------|-------------------------|-----------------------------|----------------|----------------------------------|--------------------|
| L. casei | Native (1) | CaCO ₃ | 9% (9 g) | 82.0 | 1.71 | 0.91 | | Hujanen (1996) |
| R. oryzae | Native (1) | NaOH | 15% (15 g) | 76.2 | 2.21 | 0.51 | — | Bai (2003) |
| E. coli | S. bovis (low copy) | KOH NaOH | 10% (35 g) | 64.0 | 0.40 | 0.64 | _ | Dien (2001) |
| E. coli | S. bovis (low copy) | — | 10% (35 g) | 11.4 | 0.07 | 0.11 | _ | Dien (2001) |
| S. cerevisiae | R. oryzae (over expression) | NaOH | 9% (2.7 g) | 38.0 | 1.27 | 0.42 | <u></u> | Skoy (2003) |
| S. cerevisiae | L. plantarum (over expression) | КОН | 20% (200 g) | 58.0 | 0.64 | 0.29 | ∆pdc1 | Colombie (2003) |
| S. cerevisiae | Bovin (2) | CaCO ₃ | 10% (4 g) | 55.6 | 0.76 | 0.56 | ∆pdc1 | Ishida (2005) |
| S. cerevisiae | Bovin (2) | CaCO ₃ | 10% (4 g) | 82.3 | 0.38 | 0.82 | Δpdc1 Δpdc5 | Ishida (2006) |
| S. cerevisiae | Bovin (4) | NaOH | 10% (4 g) | 74.2 | 1.55 | 0.74 | ∆pdc1 ∆adh1 | Tokuhiro (2009) |
| S. cerevisiae | Bovin (2) | | 10% (10 g) | 36.0 | 0.50 | 0.36 | ∆pdc1 | Gao (2011) |
| S. cerevisiae (SH6779) | Bovin (1) | — | 10% (3 g) | 48.0 | 0.67 | 0.48 | ∆pdc1 | This study |

Table 8. Comparison of lactic-acid production

All single and multiple disruptants (SH6765–SH6779) retained the *PDC5* gene. Because Pdc1 and Pdc5 are key enzymes for ethanol production (Hohmann and Cederberg, 1990) and *PDC5* transcription is enhanced about five-fold in $\Delta pdc1$ disruptants (Hohmann^b, 1991), Pdc5 activity may allow ethanol production in the *LDH*-harboring disruptants. Consistent with this, ethanol was detectable in all single and multiple disruptants (SH6765–SH6779; data not shown). However, because the $\Delta pdc1\Delta pdc5$ double disruptant failed to grow using glucose as a sole carbon source (Hohmann^b, 1991), additional disruption of PDC5 may not be suitable for construction of an industrial strain. It has been also reported that the lactic-acid productivity of a double $\Delta pdc1 \Delta adh1$ disruptant is higher than that of a single $\Delta pdc1$ disruptant, although growth rate of the $\Delta pdc1\Delta adh1$ disruptant on glucose slightly decreased as compared with the $\Delta pdcl$ disruptant (Tokuhiro *et al.*, To minimize the negative effect on growth, other approaches may be 2009). suitable to increase the productivity of lactic acid. For example, lactic-acid production by the S. cerevisiae isogenic diploid strain is higher than the haploid In addition, increasing the LDH gene copy number on strain (Skory, 2003). genome to six enhances LDH activity 2.8-fold versus two gene copies (Saitoh et al., However, as noted in Fig. 4, the resistance of some disruptants, such as 2005). $\Delta dse2\Delta scw11\Delta eaf3$ and $\Delta dse2\Delta eaf3\Delta sed1$, differed from the disruptants in Fig. 2, possibly related to expression of LDH. Therefore, although increasing the LDH copy number is an effective way to enhance the lactic-acid productivity, it should be kept in mind that this may also decrease the degree of lactic-acid resistance in some On the basis of these findings, it appears appropriate to construct disruptants. industrial strains by either disruption of ADH1, a construction of isogenic diploid strain, and/or increasing the LDH copy number of LDH gene. In addition, because undissociated lactic acid freely diffuses through the cell membrane, lactic acid production at a pH lower than 3.86, which is the pK_a of lactic acid, may facilitate higher lactic-acid concentrations in the medium owing to increased efflux. Therefore, I propose that the enhancement of lactic-acid resistance is an important and viable method for breeding yeast strains for lactic-acid production.
Fragmentation of the vacuole occurred rapidly under lactic acid conditions (Fig. Undissociated lactic acid diffuses through the plasma membrane and 7). dissociates in the cytoplasm because of higher intracellular pH, which generates protons that acidify the cytoplasm and also increases concentration of lactate anion, which in turn increases turgor pressure. Upon exposure to lactic-acid stress, the growth of yeast cells was inhibited and the cell viability decreased (Fig. 8). Intracellular amino acid homeostasis was also disturbed severely after addition of lactic acid (Table 7). However, after incubation in lactic acid-containing medium for 24 h, the yeast cells resumed growth and survival rate was recovered Intracellular amino acid content was also restored in significantly (Fig. 8). comparison to the level after 2 h exposure to lactic-acid (Table 7). During this period, the vacuole began to fragment (Fig. 7) but seemed to be restored to normal after 24 h incubation in lactic-acid medium (data not shown). Although the mechanisms for the induction of the vacuolar fragmentation under lactic-acid conditions is not clear, vacuolar fragmentation may be an adaptive response to lactic-acid stress: the resulting increase in the surface area/volume ratio may maximize sequestration of proton and/or the anion by the vacuolar proton-ATPase and/or anion transporters which are not yet identified on the vacuolar membrane. This kind of response has been reported in the calcium and sodium-stress responses to increase the uptake efficiency of ions by vacuolar transporters (Kellermayer et al., In the present study, vacuolar morphology in the SH6718 lactic acid 2003). resistant strain ($\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$) under 4% lactic acid was examined, considering that the vacuole of lactic acid resistant strain was not fragmented.

68

Although the vacuole of the $\Delta dse2\Delta scw11\Delta eaf3\Delta sed1$ disruptant was fragmented after 2 h incubation in 4% lactic acid, the ratio of the fragmented cells was slightly lower than in wild-type strain (unpublished data).

The amount of intracellular amino acids decreased after 2 h of incubation with lactic acid (Table 7) and addition of excess amino acids to lactic-acid media could rescue the growth defect of disruptants (Fig. 9). However, it was not clear whether the decrease in intracellular amino acids played a role in lactic-acid sensitivity. It is possible that lactic-acid sensitivity was caused by some other factor and as a result, the amount of intracellular amino acids decreased. However, when the amino-acid requirements were complemented by introducing each of three plasmids bearing have LEU2, HIS3, or LYS2 as a selection marker into BY4742 (MAT α $leu2\Delta 0$ his $3\Delta 1$ ura $3\Delta 0$ lys $2\Delta 0$), the transformants showed higher lactic-acid resistance in 6% lactic-acid medium as compared with BY4742 and the total amount of intracellular amino acids of these strains was higher than that of BY4742 after 2 h incubation with 4% lactic acid (unpublished data). It is also known that acetic-acid sensitivity of S. cerevisiae was alleviated by adding of Tyr to acetic-acid medium and that acetic-acid resistance is enhanced by overexpression of TAT2, which encodes a tyrosine permease (Shima et al., unpublished data). These data suggest the possibility that the amount of intracellular amino acids is one factors affecting the resistance of cells to acid stress and the improvement of cellular mechanisms to increase amino acid availability, such as the storage capacity of vacuole and uptake of amino acids may contribute to increased resistance to acid-stress.

69

This study is beneficial not only for breeding of yeast strains with high lactic acid production without neutralization but also for a full-scale investigation of the response mechanisms to organic and inorganic acid stress in *S. cerevisiae*. The goal of this study was to establish a strategy to improve lactic-acid resistance in yeast and construct a superior lactic acid resistant yeast strain producing lactic acid efficiently without neutralization. The results show successful construction of yeast strains with improved lactic-acid resistance and suggest that lactic-acid stress causes vacuolar fragmentation and impairs intracellular amino-acid homeostasis.

References

- Abbott, D. A., Suir, E., van Maris, A. J., and Pronk, J. T.: Physiological and transcriptional responses to high concentrations of lactic acid in anaerobic chemostat cultures of *Saccharomyces cerevisiae*, Appl. Environ. Microbiol., 74, 5759–5768 (2008).
- Adachi, E., Torigoe, M., Sugiyama, M., Nikawa, J., and Shimizu, K.: Modification of metabolic pathways of *Saccharomyces cerevisiae* by the expression of lactate dehydrogenase and deletion of pyruvate decarboxylase genes for the lactic acid fermentation at low pH value, J. Ferment. Bioeng., **86**, 284–289 (1998).
- Arntzen, C. E., and Dale, B. E.: Biobased industrial products, priorities for research and commercialization, National Academy Press, Washington, D.C. (1999).
- Bai, D. M., Jia, M. Z., Zhao, X. M., Ban, R., Shen, F., Li, X. G., and Xu, S. M.: L (+)-lactic acid production by pellet-form *Rhizopus oryzae* R1021 in a stirred tank fermentor, Chem. Eng. Sci., 58, 785–791 (2003).
- Bai, D. M., Li, S. Z., Liu, Z. L., and Cui, Z. F.: Enhanced L-(+)-lactic acid production by an adapted strain of *Rhizopus oryzae* using corncob hydrolysate. Appl. Biochem. Biotechnol., 144, 79–85 (2008).
- Bauer, B. E., Rossington, D., Mollapour, M., Mamnun, Y., Kuchler, K., and Piper, P. W.: Weak organic acid stress inhibits aromatic amino acid uptake by yeast, causing a strong influence of amino acid auxotrophies on the phenotypes

of membrane transporter mutants, Eur. J. Biochem., 270, 3189-3195 (2003).

- Bayrock, D. P. and Ingledew, W. M.: Inhibition of yeast by lactic acid bacteria in continuous culture: nutrient depletion and/or acid toxicity?, J. Ind. Microbiol. Biotechnol., 31, 362–368 (2004).
- Benninga, J. E.: A history of lactic acid making, Kluyver Academic Publishers, Dordrecht, The Netherlands (1990).
- Bonangelino, C. J., Nau, J. J., Duex, J. E., Brinkman, M., Wurmser, A. E., Gray, J. D., Emr, S. D., and Weisman, L. S.: Osmotic stress-induced increase of phosphatidylinositol 3,5-bisphosphate requires Vac14p, an activator of the lipid kinase Fab1p, J. Cell Biol., 156, 1015–1028 (2002).
- Bracey, D., Holyoak, C. D., and Coote, P. J.: Comparison of the inhibitory effect of sorbic acid and amphotericin B on *Saccharomyces cerevisiae*: is growth inhibition dependent on reduced intracellular pH? J. Appl. Microbiol., **85**, 1056–1066 (1998).
- Brachmann, C. B., Davies, A., Cost, G. J., Caputo, E., Li, J., Hieter, P. and Boeke, J. D.: Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications, Yeast, 14, 115–132 (1998).
- Bugnicourt, A., Froissard, M., Sereti, K., Ulrich, H. D., Haguenauer-Tsapis, R., and Galan, J. M.: Antagonistic roles of ESCRT and Vps class C/HOPS complexes in the recycling of yeast membrane proteins, Mol. Biol. Cell, 15, 4203–214 (2004).

Caesar, R., Warringer, J., and Blomberg, A.: Physiological importance and

identification of novel targets for the N-terminal acetyltransferase NatB, Eukaryot. Cell, 5, 368–378 (2006).

- Cappellaro, C., Mrsa, V., and Tanner, W.: New potential cell wall glucanases of Saccharomyces cerevisiae and their involvement in mating, J. Bacteriol., 180, 5030–5037 (1998).
- Caro, L. H., Smits, G. J., van Egmond, P., Chapman, J. W., and Klis, F. M.: Transcription of multiple cell wall protein-encoding genes in *Saccharomyces cerevisiae* is differentially regulated during the cell cycle, FEMS Microbiol. Lett., 161, 345–349 (1998).
- Casal, M., Paiva, S., Andrade, R. P., and Leão, C.: The lactate-proton symport of Saccharomyces cerevisiae is encoded by JEN1, J. Bacteriol., 181, 2620–2623 (1999).
- Cássio, F., Leão, C., and van Uden, N.: Transport of lactate and other short chain monocarboxylates in the yeast *Saccharomyces cerevisiae*, Appl. Environ. Microbiol., 53, 509–513 (1987).
- Chambers, P., Issaka, A., and Palecek, S. P.: Saccharomyces cerevisiae JEN1 promoter activity is inversely related to concentration of repression sugar, Appl. Environ. Microbiol., **70**, 8–17 (2004).
- Chotani, G., Dodge, T., Hsu, A., Kumar, M., LaDuca, R., Trimbur, D., Weyler,
 W., and Sanford, K.: The commercial production of chemicals using pathway engineering, Biochim. Biophys. Acta, 1543, 434–455 (2000).
- Colman-Lerner, A., Chin, T. E., and Brent, R.: Yeast Cbk1 and Mob2 activate daughter-specific genetic programs to induce asymmetric cell fates, Cell, 107,

739-750 (2001).

- Colombié, S., Dequin, S., and Sablayrolles, J. M.: Control of lactate production by *Saccharomyces cerevisiae* expressing a bacterial *LDH* gene, Enzyme Microb. Technol., 33, 38–46 (2003).
- Conibear, E. and Stevens, T. H.: Studying yeast vacuoles. In Christine Guthrie and Gerald R. Fink (eds.) Guide to Yeast Genetics and Molecular and Cell Biology, Pt C, Methods in Enzymology, 351, 408–432 (2002).
- Corson, L. B., Folmer, J., Strain, J. J., Culotta, V. C., and Cleveland, D. W.: Oxidative stress and iron are implicated in fragmenting vacuoles of *Saccharomyces cerevisiae* lacking Cu,Zn-superoxide dismutase, J. Biol. Chem., 274, 27590–27596 (1999).
- Datta, R., Tsai, S. P., Bonsignore, P., Moon, S. H., and Frank, J. R.: Technological and economic potential of poly (lactic acid) and lactic acid derivatives, FEMS Microbiol. Rev., 16, 221–231 (1995).
- Dequin, S., and Barre, P.: Mixed lactic acid-alcoholic fermentation by Saccharomyces cerevisiae expressing the Lactobacillus casei L(+)-LDH, Bio/Technology., 12, 173–177 (1994).
- Dien, B. S., Nichols, N. N., Bothast, R. J.: Recombinant Escherichia coli engineered for production of L-lactic acid from hexose and pentose sugars, J. Ind. Microbiol. Biotechnol., 27, 259–264 (2001).
- Dominguez, A., Villanueva, J. R., and Sentandreu, R.: Inositol deficiency in Saccharomyces cerevisiae NCYC 86, Antonie Van Leeuwenhoek, 44, 25–34 (1978).

- Dwight, S. S., Harris, M. A., Dolinski, K., Ball, C. A., Binkley, G., Christie, K.
 R., Fisk, D. G., Issel-Tarver, L., Schroeder, M., Sherlock, G., and other 4 suthors:. Saccharomyces genome database (SGD) provides secondary gene annotation using the gene ontology (GO), Nucleic Acids Res., 30, 69–72 (2002).
- Eisler, H., Fröhlich, K. U., and Heidenreich, E.: Starvation for an essential amino acid induces apoptosis and oxidative stress in yeast, Exp. Cell Res., 300, 345–353 (2004).
- Eraso, P., and Gancedo, J. M.: Activation of yeast plasma membrane ATPase by acid pH during growth, FEBS Lett., 224, 187–192 (1987).
- Fernandes, A. R., Mira, N. P., Vargas, R. C., Canelhas, I., and Sá-Correia, I.: Saccharomyces cerevisiae adaptation to weak acids involves the transcription factor Haa1p and Haa1p-regulated genes, Biochem. Biophys. Res. Commun., 337, 95–103 (2005).
- Foster, P. L.: Adaptive mutation in *Escherichia coli*, J. Bacteriol., 186, 4846–4852 (2004).
- Gao, M. T., Shimamura, T., Ishida, N., and Takahashi, H.: pH-Uncontrolled lactic acid fermentation with activated carbon as an adsorbent, Enzyme Microb. Technol., 48, 526–530 (2011).
- Goehring, A. S., Rivers, D., and Sprague, G. F.: Attachment of the ubiquitin-related protein Urm1p to the antioxidant protein Ahp1p, Eukaryot. Cell, 2, 930–936 (2003).
- Goossens, A., de la Fuente, N., Forment, J., Serrano, R., and Portillo, F.: Regulation of yeast H⁺-ATPase by protein kinases belonging to a family

dedicated to activation of plasma membrane transporters, Mol. Cell Biol., 20, 7654–7661 (2000).

- Gregory, S. L., Ebrahimi, S., Milverton, J., Jones, W. M., Bejsovec, A., and Saint, R.: Cell division requires a direct link between microtubule-bound RacGAP and Anillin in the contractile ring, Curr. Biol., 18, 25–29 (2008).
- Gritz, L. and Davies, J.: Plasmid-encoded hygromycin B resistance: the sequence of hygromycin B phosphotransferase gene and its expression in *Escherichia coli* and *Saccharomyces cerevisiae*, Gene, 25, 179–188 (1983).
- Hatzixanthis, K., Mollapour, M., Seymour, I., Bauer, E. B., Krapf, G., Schüller,
 C., Kuchler, K., and Piper, P. W.: Moderately lipophilic carboxylate compounds are the selective inducers of the *Saccharomyces cerevisiae* Pdr12p ATP-binding cassette transporter, Yeast, 20, 575–585 (2003).
- Hauser, M., Narita, V., Donhardt, A. M., Naider, F., and Becker, J. M.: Multiplicity and regulation of genes encoding peptide transporters in Saccharomyces cerevisiae, Mol. Membrane Biol., 18, 105–112 (2001).
- Hazan, R., Levine, A., and Abeliovich, H.: Benzoic acid, a weak organic acid food preservative, exerts specific effects on intracellular membrane trafficking pathways in *Saccharomyces cerevisiae*, Appl. Environ. Microbiol., **70**, 4449–4457 (2004).
- Hiraishi, H., Mochizuki, M., and Takagi, H.: Enhancement of stress tolerance in Saccharomyces cerevisiae by overexpression of ubiquitin ligase Rsp5 and ubiquitin-conjugating enzymes, Biosci. Biotechnol. Biochem., 70, 2762–2765 (2006).

- Hirata, R. and Takatsuki, A.: Role of organelle acidification in intracellular protein transport, RIKEN Review, 41, 90–91 (2001).
- Hochstrasser, M.: Ubiquitin, proteasomes, and the regulation of intracellular protein degradation, Curr. Opin. Cell Biol., 7, 215–223 (1995).
- Hofmann, R. M., and Pickart, C. M.: Noncanonical MMS2-encoded ubiquitin-conjugating enzyme functions in assembly of novel polyubiquitin chains for DNA repair, Cell, 96, 645–653 (1999).
- Hofvendahl, K., and Hahn-Hägerdal, B.: Factors affecting the fermentative lactic acid production from renewable resources, Enzyme Microb. Technol., 26, 87–107 (2000).
- Hohmann, S.^a: Characterization of PDC6, a third structural gene for pyruvate decarboxylase in Saccharomyces cerevisiae, J. Bacteriol., 173, 7963–7969 (1991).
- Hohmann, S.^b: PDC6, a weakly expressed pyruvate decarboxylase gene from yeast, is activated when fused spontaneously under the control of the *PDC1* promoter, Curr. Genet., **20**, 373–378 (1991).
- Hohmann, S., and Cederberg, H.: Autoregulation may control the expression of yeast pyruvate decarboxylase structural genes *PDC1* and *PDC5*, Eur. J. Biochem., 188, 615–621 (1990).
- Holmberg, S. and Petersen, J. G.: Regulation of isoleucine-valine biosynthesis in Saccharomyces cerevisiae, Curr. Genet., 13, 207–217 (1988).
- Holyoak, C. D., Bracey, D., Piper, P. W., Kuchler, K., and Coote, P. J.: The Saccharomyces cerevisiae weak-acid-inducible ABC transporter Pdr12 transports

fluorescein and preservative anions from the cytosol by an energy-dependent mechanism, J. Bacteriol., **181**, 4644–4652 (1999).

- Hujanen, M., and Linko, Y. Y.: Effect of temperature and various nitrogen sources on L (+)-lactic acid production by *Lactobacillus casei*, Appl. Microbiol Biotech., 45, 307–313 (1996).
- Ikada, Y., Jamshidi, k., Tsuji, H., and Hyon, S. H.: Stereocomplex formation between enantiomeric poly (lactides), Macromolecules, 20, 904–906 (1987).
- Ishida, N., Saitoh, S., Tokuhiro, K., Nagamori, E., Matsuyama, T., Kitamoto, K., and Takahashi, H.: Efficient production of L-lactic acid by metabolically engineered *Saccharomyces cerevisiae* with a genome-integrated L-lactate dehydrogenase gene, Appl. Environ. Microbiol., **71**, 1964–1970 (2005).
- Ishida, N., Satitoh, S, Onishi, T., Tokuhiro, K., Nagamori, E., Kitamoto, K., and
 Takahashi, H.: The effect of pyruvate decarboxylase gene knockout in
 Saccharomyces cerevisiae on L-lactic acid production, Enzyme Microb. Technol.
 33, 38–46 (2006).
- Jacquemin-Faure, I., Thomas, D., Laporte, J., Cibert, C., and Surdin-Kerjan,
 Y.: The vacuolar compartment is required for sulphur amino acid homeostasis in Saccharomyces cerevisiae, Mol. Gen. Genet., 244, 519–529 (1994).
- Jones, E. W. and Fink, G. R.: Regulation of amino acid and nucleotide biosynthesis in yeast. In: Strathern, J. N., Jones, E. W., Broach, J. R., Editors, 1982. The Molecular Biology of the Yeast *Saccharomyces*: Metabolism and Gene Expression, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, pp 181–299 (1982).

- Kane, P. M.: The where, when, and how of organelle acidification by the yeast vacuolar H⁺-ATPase, Microbiol. Mol. Biol. Rev., **70**, 177–191 (2006).
- Kawahata, M., Masaki, K., Fujii, T., and Iefuji, H.: Yeast genes involved in response to lactic acid and acetic acid: acidic conditions caused by the organic acids in *Saccharomyces cerevisiae* cultures induce expression of intracellular metal metabolism genes regulated by Aft1p, FEMS Yeast Res., 6, 924–936 (2006).
- Kellermayer, R., Aiello, D. P., Miseta, A., and Bedwell, D. M.: Extracellular Ca^{2+} sensing contributes to excess Ca^{2+} accumulation and vacuolar fragmentation in a *pmr1* mutant of *S. cerevisiae*, J. Cell Sci., **116**, 1637–1646 (2003).
- Kitamoto, K., Yoshizawa, K., Ohsumi, Y., and Anraku, Y.: Mutants of Saccharomyces cerevisiae with defective vacuolar function, J. Bacteriol., 170, 2687–2691 (1988).
- Klionsky, D. J., Herman, P. K., and Emr, S. D.: The fungal vacuole: composition, function, and biogenesis, Microbiol. Rev., 54, 266–292 (1990).
- Lambert, R. J., and Stratford, M.: Weak-acid preservatives: modeling microbial inhibition and response, J. Appl. Microbiol., 86, 157–164 (1999).
- Mannhaupt, G., Stucka, R., Pilz, U., Schwarzlose, C., and Feldmann, H.: Characterization of the prephenate dehydrogenase-encoding gene, *TYR1*, from *Saccharomyces cerevisiae*, Gene, **85**, 303–311 (1989).
- Marshall, V. M.: Lactic acid bacteria: starters for flavor, FEMS Microbiol. Rev., 46, 327–336 (1987).
- Martínez-Muñoz, G. A. and Kane, P.: Vacuolar and plasma membrane proton

pumps collaborate to achieve cytosolic pH homeostasis in yeast, J. Biol. Chem., **283**, 20309–20319 (2008).

- Matsuura, K. and Takagi, H.: Vacuolar functions are involved in stress-protective effect of intracellular proline in *Saccharomyces cerevisiae*, J. Biosci. Bioeng., 100, 538–544 (2005).
- McNeil, J. B., McIntosh, E. V., Taylor, B. V., Zhang, F. R., Tang, S., and Bognar,
 A. L.: Cloning and molecular characterization of three genes, including two genes encoding serine hydroxyltransferases, whose inactivation is required to render yeast auxotrophic for glycine, J. Biol. Chem., 269, 9155–9165 (1994).
- Mewes, H. W., Frishman, D., Guldener, U., Mannhaupt, G., Mayer, K., Mokrejs, M., Morgenstern, B., Munsterkotter, M., Rudd, S., and Weil, B.: MIPS: a database for genomes and protein sequences, Nucleic Acids Res., 30, 31–34 (2002).
- Mira, N. P., Becker, J., and Sá-Correia, I.: Genomic expression program involving the Haa1p-regulon in Saccharomyces cerevisiae response to acetic acid. OMICS, 14, 578–601 (2010).
- Mira, N. P., Lourenco, A. B., Fernandes, A. R., Becker, J. D., and Sá-Correia,
 I.: The *RIM101* pathway has a role in *Saccharomyces cerevisiae* adaptive response and resistance to propionic acid and other weak acids, FEMS Yeast Res.,
 9, 202–216 (2009).
- Mollapour, M., and Piper, P. W.: Hog1p MAP kinase determines acetic acid resistance in *Saccharomyces cerevisiae*, FEMS Yeast Res., 6, 1274–1280 (2006).
 Mollapour, M. and Piper, P. W.: Hog1 Mitogen-activated protein kinase

phosphorylation targets the yeast Fps1 aquaglyceroporin for endocytosis, thereby rendering cells resistant to acetic acid, Mol. Cell Biol., **27**: 6446–6556 (2007).

- Mollapour, M., Shepherd, A., and Piper, P. W.: Novel stress responses facilitate Saccharomyces cerevisiae growth in the presence of the monocarboxylate preservatives, Yeast, 25, 169–177 (2008).
- Muller, E. H., Richards, E. J., Norbeck. J., Byrne, K. L., Karlsson, K. A., Pretorius, G. H., Meacock, P. A., Blomberg, A., and Hohmann, S.: Thiamine repression and pyruvate decarboxylase autoregulation independently control the expression of the *Saccharomyces cerevisiae PDC5* gene, FEBS Lett., 449, 245–250 (1999).
- Nelson, H. and Nelson, N.: Disruption of genes encoding subunits of yeast vacuolar H⁺-ATPase causes conditional lethality, Proc. Natl. Acad. Sci. USA, 87, 3503–3507 (1990).
- Ohara, H., Okuyama, H, Sawa, S., Fujii, Y., and Hiyama, K.: Development of industrial production of high molecular weight poly-L-lactate from renewable resources, Nippon Kagaku Kaishi., 6, 323–331 (2001).
- Pandey, A., Soccol, C. R., Rodriguez-Leon, J. A., and Nigam, P.: Solid state fermentation in biotechnology: fundamentals and applications, Asiatech. Publishers, New Delhi (2001).
- Perry, J. R., Basrai, M. A., Steiner, H. Y., Naider, F., and Becker, J. M.: Isolation and characterization of a *Saccharomyces cerevisiae* peptide transport gene, Mol. Cell Biol., 14, 104–115 (1994).
- Perzov, N., Nelson, H., and Nelson, N.: Altered distribution of the yeast plasma

membrane H^+ -ATPase as a feature of vacuolar H^+ -ATPase null mutants, J. Biol. Chem., 275, 40088–40095 (2000).

- Phelan, J. P., Millson, S. H., Parker, P. J., Piper, P. W., and Cooke, F. T.: Fab1p and AP-1 are required for trafficking of endogenously ubiquitylated cargoes to the vacuole lumen in *S. cerevisiae*, J. Cell Sci., **119**, 4225–4234 (2006).
- Piper, P., Mahé, Y., Thompson, S., Pandjaitan, R., Holyoak, C., Egner, R., Mühlbauer, M., Coote, P., and Kuchler, K.: The pdr12 ABC transporter is required for the development of weak organic acid resistance in yeast, EMBO J., 17, 4257–4265 (1998).
- Piper, P. W.: Yeast superoxide dismutase mutants reveal a pro-oxidant action of weak organic acid food preservatives, Free Radic. Biol. Med., 27, 1219–1227 (1999).
- Porro, D., Barmbilla, L., Ranzi, B. M., Martegani, E., and Alberghina, L.: Development of metabolically engineered *Saccharomyces cerevisiae* cells for the production of lactic acid, Biotechnol. Prog., 11, 294–298 (1995).
- Ramos, C. and Calderon, I. L.: Biochemical evidence that the Saccharomyces cerevisiae THR4 gene encodes threonine synthetase, FEBS Lett., 351, 357–359 (1994).
- Reid, J. L., Moqtaderi, Z., and Struhl, K.: Eaf3 regulates the global pattern of histone acetylation in *Saccharomyces cerevisiae*, Mol. Cell Biol., 24, 757–764 (2004).
- Robinson, M. D., Grigull, J., Mohammad, N., and Hughes, T. R.: FunSpec: a web-based cluster interpreter for yeast, BMC Bioinformatics, 3, 35 (2002).

- Rothman, J. H., Howald, I., and Stevens, T. H.: Characterization of genes required for protein sorting and vacuolar function in the yeast *Saccharomyces cerevisiae*, EMBO J., 8, 2057–2065 (1989).
- Rothman, J. H. and Stevens, T. H.: Protein sorting in yeast: mutants defective in vacuole biogenesis mislocalize vacuolar proteins into the late secretory pathway, Cell, 26, 1041–1051 (1986).
- Sakai, K, and Ezaki, Y.: Open L-lactic acid fermentation of food refuse using thermophilic *Bacillus coagulans* and fluorescence *in situ* hybridization analysis of microfloea, J. Biosci. Bioeng., 101, 457–463 (2006).
- Saitoh, S., Ishida, N., Onishi, T., Tokuhiro, K., Nagamori, E., Kitamoto, K., and Takahashi, H.: Genetically engineered wine yeast produces a high concentration of L-lactic acid of extremely high optical purity, Appl. Environ. Microbiol, 71, 2789–2792 (2005).
- Schröder, M. and Kaufman, R. J.: ER stress and the unfolded protein response, Mutat. Res., 569, 29–63 (2005).
- Schuller, C., Mamnun, Y. M., Mollapour, M., Krapf, G., Schuster, M., Bauer, B.
 E., Piper P. W., and Kuchler, K.: Global phenotypic analysis and transcriptional profiling defines the weak acid stress response regulon in *Saccharomyces cerevisiae*, Mol. Biol. Cell, 15, 706–720 (2004).
- Seaston, A., Inkson, C., and Eddy, A. A.: The absorption of protons with specific amino acids and carbohydrates by yeast, Biochem. J., 134, 1031–1043 (1973).
- Seeboth, P. G., Bohnsack. K., and Hollenberg. C. P.: pdc1 mutants of *Saccharomyces cerevisiae* give evidence for an additional structural PDC gene:

cloning of *PDC5*, a gene homologous to *PDC1*, J. Bacteriol., **172**, 678–685 (1990).

- Seeley, E. S., Kato, M., Margolis, N., Wickner, W., and Eitzen, G.: Genomic analysis of homotypic vacuole fusion, Mol. Biol. Cell, 13, 782–794 (2002).
- Seufert, W. and Jentsch, S.: Ubiquitin-conjugating enzymes UBC4 and UBC5 mediate selective degradation of short-lived and abnormal proteins, EMBO J., 9, 543-550 (1990).
- Shima, J., Ando, A., and Takagi, H.: Possible roles of vacuolar H⁺-ATPase and mitochondrial function in tolerance to air-drying stress revealed by genome-wide screening of *Saccharomyces cerevisiae* deletion strains, Yeast, 25, 179–190 (2008).
- Shimoi, H., Kitagaki, H., Ohmori, H., Iimura, Y., and Ito, K.: Sed1p is a major cell wall protein of *Saccharomyces cerevisiae* in the stationary phase and is involved in lytic enzyme resistance, J. Bacteriol., **180**, 3381–3387 (1998).
- Simões, T., Mira, N. P., Fernandes, A. R., and Sá-Correia, I.: The SP11 gene, encoding a glycosylphosphatidylinositol (GPI)-anchored cell wall protein, plays a prominent role in the development of yeast resistance to lipophilic weak acids food preservatives, Appl. Environ. Microbiol., **72**, 7168–7175 (2006).
- Skory, C. D.: Lactic acid production by Saccharomyces cerevisiae expressing a Rhizopus oryzae lactate dehydrogenase gene, J. Ind. Microbiol. Biotechnol., 30, 22-27 (2003).
- Sugiyama, M., Ikushima, S., Nakazawa, T., Kaneko, Y., and Harashima, S.: PCR-mediated repeated chromosome splitting in *Saccharomyces cerevisiae*,

Biotechniques, 38, 909-914 (2005).

- Teixeira, M. C., Raposo, L. R., Mira, N. P., Lourenço, A. B., and Sá-Correia, I.: Genome-wide identification of *Saccharomyces cerevisiae* genes required for maximal tolerance to ethanol, Appl. Environ. Microbiol., 75, 5761–5772 (2009).
- Thomsson, E., and Larsson, C.: The effect of lactic acid on anaerobic carbon or nitrogen limited chemostat cultures of *Saccharomyces cerevisiae*, Appl. Microbiol. Biotechnol., 71, 533–542 (2006).
- Tokuhiro, K., Ishida, N., Nagamori, E., Saitoh, S., Onishi, T., Kondo, A., Takahashi, H.: Double mutation of the *PDC1* and *ADH1* genes improves lactate production in the yeast *Saccharomyces cerevisiae* expressing the bovine lactate dehydrogenase gene, Appl. Microbiol. Biotechnol., 82, 883–890 (2009).
- **Toothaker, L. E.:** Multiple Comparison Procedures. Sage University Paper Series on Quantitative Applications in the Social Sciences, Sage, Newbury Park, CA, (1993).
- Tsuji, F.: Autocatalytic hydrolysis of amorphous-made polylactides: effects of L-lactide content, tacticity, and enantiomeric polymer blending, Polymer, 43, 1789–1796 (2002).
- Ufano, S., Pablo, M. E., Calzada, A., del Rey, F., and Vázquez de Aldana, C. R.: The Swm1p subunit of the APC/Cyclosome is required for activation of the daughter-specific gene expression program mediated by Ace2p during growth at high temperature in *Saccharomyces cerevisiae*, J. Cell Sci., 117, 545–557 (2004).
- van der Rest, M., Kamminga, A. H., Nakano, A., Anraku, Y., Poolman, B., Konings, W. N.: The plasma membrane of *Saccharomyces cerevisiae*: Structure,

function and biogenesis, Microbiol. Rev., 59, 304-322 (1995).

- van der Vaart, J. M., van Schagen, F. S., Mooren, A. T, Chapman, J. W., Klis, F. M., and Verrips, C. T.: The retention mechanism of cell wall proteins in *Saccharomyces cerevisiae*. Wall-bound Cwp2p is beta-1,6-glucosylated, Biochim. Biophys. Acta., 1291, 206–214 (1996).
- van Dijk, R., Faber, K. N., Hammond, A. T., Glick, B. S., Veenhuis, M., and Kiel, J. A.: Tagging Hansenula polymorpha genes by random integration of linear DNA fragments (RALF), Mol. Genet. Genomics., 266, 646–656 (2001).
- van Maris, A. J., Abbott, D. A., Bellissimi, E., van den Brink, J., Kuyper, M., Luttik, M. A., Wisselink, H. W., Scheffers, W. A., van Dijken, J. P., and Pronk, J. T.: Alcoholic fermentation of carbon sources in biomass hydrolysates by *Saccharomyces cerevisiae*: current status, Antonie Van Leeuwenhoek, 90, 391–418 (2006).
- Vivkroy, T. B.: Comprehensive biotechnology, Vol. 3, ed. M. Moo-Young, Pergamon Press, Oxford, pp. 761–776 (1985).
- Viegas, C. A., Almeida, P. F., Cavaco, M., and Sá-Correia, I.: The H⁺-ATPase in the plasma membrane of *Saccharomyces cerevisiae* is activated during growth latency in octanoic acid-supplemented medium accompanying the decrease in intracellular pH and cell viability, Appl. Environ. Microbiol., 64, 779–783 (1998).
- Viegas, C. A., and Sà-Correia, I.: Activation of plasma membrane ATPase of Saccharomyces cerevisiae by octanoic acid, J. Gen. Microbiol., 137, 645–651 (1991).

- Wee, Y. J., Kim, J. N., Ryu, H. W.: Biotechnological production of lactic acid and its recent applications, Food Technol. Biotechnol., 44, 163–172 (2006).
- Xu, Q., Jürgensmeier, J. M., and Reed, J. C.: Methods of assaying Bcl-2 and Bax family Proteins in yeast, Methods, 17, 292–304 (1999).
- Zhang, D. X., Cheryan, M.: Direct fermentation of starch to lactic acid by Lactobacillus amylovorus, Biotechnol. Lett., 10, 733-738 (1991).

List of Publications

- Suzuki, T., Sakamoto, T., Sugiyama, M., Ishida, N., Kambe, H., Obata, S., Kaneko, Y., and Harashima, S.: Disruption of multiple genes whose deletion causes lactic-acid resistance improves lactic-acid resistance and productivity in *Saccharomyces cerevisiae*. J. Biosci. Bioeng. (In press).
- Suzuki, T., Sugiyama, M., Wakazono, K., Kaneko, Y., and Harashima, S.: Lactic-acid stress causes vacuolar fragmentation and impairs intracellular amino-acid homeostasis in *Saccharomyces cerevisiae*. J. Biosci. Bioeng., 113, 421–430 (2012).

Acknowledgements

I wish to express my deepest and sincere gratitude to Professor Dr. Satoshi Harashima for his kindness in giving me the great opportunity to study the yeast genetics and molecular biology in his laboratory, and for worthy guidance throughout this study. I would like to thank to Professor Dr. Yoshinobu Kaneko and Assistant Professor Dr. Yu Sasano for his helpful comments and support during this study. I am also grateful to Associate Professor Dr. Minetaka Sugiyama. His intellectual and close guidance were of inestimable value for this study. My appreciation is also expressed to Professor Dr. Toshiya Muranaka and Professor Dr. Kazuhito Fujiyama for their valuable comments and suggestions to improve the thesis.

My gratitude is also extended to all Harashima lab members. Special thanks are given to Mr. Takatoshi Sakamoto for construction of multiple disruptants, Mr. Kenta Wakazono for screening of lactic acid sensitive disruptants, and Mrs. Minori Numamoto for developing my idea and daily life.

I owe my thanks to my father Syogo Suzuki and my mother Motoko Suzuki for their encouragement, understanding, and support.

