

Title	Isospin symmetry of T_z = $\pm 3/2 \rightarrow \pm 1/2$ Gamow-Teller transitions in A=41 nucllei
Author(s)	Fujita, Y.; Shimbara, Y.; Adachi, T. et al.
Citation	Physical Review C - Nuclear Physics. 2004, 70, p. 54311
Version Type	VoR
URL	https://hdl.handle.net/11094/27150
rights	Y. Fujita, Y. Shimbara, T. Adachi, G.P.A. Berg, B.A. Brown, H. Fujita, K. Hatanaka, J. Kamiya, K. Nakanishi, Y. Sakemi, S. Sasaki, Y. Shimizu, Y. Tameshige, M. Uchida, T. Wakasa, M. Yosoi. Physical Review C - Nuclear Physics, 70, 54311, 2004 of publication. "Copyright (2004) by the American Physical Society.
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Isospin symmetry of $T_z = \pm 3/2 \rightarrow \pm 1/2$ Gamow-Teller transitions in A = 41 nuclei

Y. Fujita,^{1,*} Y. Shimbara,^{1,†} T. Adachi,¹ G. P. A. Berg,^{2,3} B. A. Brown,⁴ H. Fujita,^{1,‡} K. Hatanaka,² J. Kamiya,^{2,§} K. Nakanishi,² Y. Sakemi,² S. Sasaki,⁵ Y. Shimizu,² Y. Tameshige,² M. Uchida,⁶ T. Wakasa,^{2,∥} and M. Yosoi⁶

¹Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

²Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047, Japan

³KVI, Zernikelaan 25, 9747 AA Groningen, The Netherlands

⁴NSCL and Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA

⁵Shizuoka Institute of Technology, Fukuroi, Shizuoka 437-8555, Japan

⁶Department of Physics, Kyoto University, Kyoto 606-8502, Japan

(Received 31 July 2004; published 12 November 2004)

Under the assumption that isospin T is a good quantum number, isobaric analog states and various analogous transitions are expected in isobars with mass number A. The strengths of $T_z = \pm 3/2 \rightarrow \pm 1/2$ analogous Gamow-Teller (GT) transitions and analogous M1 transitions within the A=41 isobar quartet are compared in detail. The $T_{z}=+3/2 \rightarrow +1/2$ GT transitions from the $J^{\pi}=3/2^{+}$ ground state of ⁴¹K leading to excited J^{π} $=1/2^+$, $3/2^+$, and $5/2^+$ states in ⁴¹Ca were measured using the (³He, t) charge-exchange reaction. With a high energy resolution of 35 keV, many fragmented states were observed, and the GT strength distribution was determined up to 10 MeV excitation energy (E_x) . The main part of the strength was concentrated in the E_x =4-6 MeV region. A shell-model calculation could reproduce the concentration, but not so well details of the strength distribution. The obtained distribution was further compared with two results of 41 Ti β decay studying the analogous $T_z = -3/2 \rightarrow -1/2$ GT strengths. They reported contradicting distributions. One-to-one correspondences of analogous transitions and analog states were assigned up to $E_x=6$ MeV in the comparison with one of these ⁴¹Ti β -decay results. Combining the spectroscopic information of the analog states in ⁴¹Ca and ⁴¹Sc, the most probable J^{π} values were deduced for each pair of analog states. It was found that $5/2^+$ states carry the main part of the observed GT strength, while much less GT strength was carried by $1/2^+$ and $3/2^+$ states. The gross features of the GT strength distributions for each J were similar for the isospin analogous $T_z = \pm 3/2 \rightarrow \pm 1/2$ transitions, but the details were somewhat different. From the difference of the distributions, isospin-asymmetry matrix elements of ≈ 8 keV were deduced. The Coulomb displacement energy, which is sensitive to the configuration of states, showed a sudden increase of about 50 keV at the excitation energy of 3.8 MeV. The strengths of several M1 transitions to the IAS in 41 Ca were compared with the strengths of analogous GT transitions. It was found that ratios of the M1 and GT transition strengths were similar, suggesting that the contributions of the $\ell \tau$ term in M1 transitions are small.

DOI: 10.1103/PhysRevC.70.054311

PACS number(s): 25.55.Kr, 21.10.Hw, 21.60.Cs, 27.40.+z

I. INTRODUCTION

Isospin is a good quantum number under the assumption that the nuclear interaction is charge symmetric and that the effect of the electromagnetic interaction is relatively small. As such, an analogous structure is expected for the same mass A nuclei with different T_z (isobars), where $T_z[=(N$ -Z/2] is the z component of the isospin T (see, e.g., Refs. [1,2]). The corresponding states in different T_z nuclei are called isobaric analog states (or simply analog states). Transitions connecting various combinations of analog states are analogous. Such "analogous transitions" should have corresponding energies and strengths. Properties and isospin symmetry of analogous transitions in isobars have been discussed extensively in Refs. [1,3].

Gamow-Teller (GT) transitions, caused purely by the $\sigma\tau$ -type operator, are well suited for the study of analog states and properties of analogous transitions, because they can be studied in both β decay and hadron charge-exchange (CE) reactions. The GT transitions are characterized by ΔL =0, ΔS =1, and ΔT_z =±1, where L and S are the orbital and spin quantum numbers. An important physical quantity for understanding the nuclear structure is the reduced GT transition strength B(GT). The B(GT) values are usually obtained from studies of GT β decay. In addition, CE reactions, such as (p,n) or $({}^{3}\text{He},t)$, performed at intermediate energies (>100 MeV/nucleon) can be used as a means to map the GT strengths over a wide range of excitation energy (E_r) [4.5]. For this purpose, one relies upon the approximate proportionality between the reaction cross sections measured at the scattering angle $\Theta = 0^{\circ}$ and their B(GT) values [6].

The simplest analogous GT transitions are expected for the odd-mass mirror nuclei with $T_z = \pm 1/2$. By comparing the GT transitions from the ground states of $T_z = -1/2$ and

^{*}Electronic address: fujita@rcnp.osaka-u.ac.jp

[†]Present address: Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047, Japan.

⁴Present address: iThemba LABS, P.O. Box 722, Somerset West 7129, South Africa.

[§]Present address: JAERI, Tokai, Ibaraki 319-1195, Japan.

Present address: Department of Physics, Kyushu University, Higashi, Fukuoka 812-8581, Japan.

+1/2 nuclei to excited states of $T_z = +1/2$ and -1/2 nuclei studied in β decays and (p,n)-type CE reactions, respectively, the symmetry of analogous transitions and analog states, and thus the isospin symmetry of isobars, has been discussed in the low-lying region of various sd-shell nuclei [6–9]. Similarly, the symmetric nature of $T_z = \pm 1$ to $T_z = 0$ GT transitions has been examined for a few systems of light *sd*-shell nuclei, such as the A = 26 nuclei system (²⁶Mg, ²⁶Al, and 26 Si) [10,11] or the A=38 system (38 Ar, 38 K, and 38 Ca) [12] by comparing (p,n) or $({}^{3}\text{He},t)$ CE reactions and β -decay studies. A good correspondence of B(GT) values with a difference of less than 5% was observed for the four strong $T_z = \pm 1 \rightarrow T_z = 0$ GT transitions in the A=26 system [11], and thus a good isospin symmetry was found. However, such a comparison was possible only for the low-lying states up to $E_x \approx 3$ MeV due to the low $Q_{\rm EC}$ values of the relevant β decays [13]. An extension of the study to *fp*-shell nuclei with higher $Q_{\rm EC}$ values was discussed [14].

The symmetry of systems with larger isospin T is more challenging. The $T_z = -T$ nuclei, unlike stable $T_z = T$ nuclei, are usually far away from the β -stability line, and it is expected that the "isospin-asymmetric features," if any, are observed better. The isospin structure and analogous transitions of the so-called "T=3/2 system," after subtracting the Coulomb displacement energy, are schematically shown in Fig. 1. The isobaric analog states of the T=3/2 ground states of the $T_z = \pm 3/2$ nuclei are called the IAS in the $T_z = \pm 1/2$ nuclei. There are several stable $T_z = 3/2$ nuclei in the sd- and *fp*-shell regions. The GT transitions from the ground state (g.s.) of a $T_z = +3/2$ nucleus to $\Delta J^{\pi} = 1^+$ states (GT states) in a $T_z = +1/2$ nucleus can be studied via CE reactions, such as (p,n) or $({}^{3}\text{He},t)$. The measurements of the corresponding $T_z = -3/2$ to $T_z = -1/2$ transitions in β decays have become more accurate in recent years with the technical development of producing a larger number of unstable nuclei with greater purity. It should be noted that these β decays have relatively high $Q_{\rm EC}$ values of more than 10 MeV. Therefore, it is expected that not only the transitions to the low-lying states, but also the main part of the GT transition strengths can be studied in these β -decay measurements.

Among the candidates for the $T_z = \pm 3/2 \rightarrow \pm 1/2$ comparison, we find that analogous transitions in the A = 41 system i.e., ⁴¹K ($T_z = +3/2$) to ⁴¹Ca ($T_z = +1/2$) and ⁴¹Ti ($T_z = -3/2$) to ⁴¹Sc ($T_z = -1/2$)—are well suited for an accurate study of the analogous GT transitions and also the isospin-symmetry structure of isobars. The former can be studied by a CE reaction on a stable ⁴¹K target and the latter in the β decay of ⁴¹Ti. Owing to the high $Q_{\rm EC}$ value of 12.93 MeV in the ⁴¹Ti β decay [15], the *B*(GT) values have been reported for ⁴¹Sc up to about 8 MeV by several groups [16–19].

In this paper, we report the GT strengths of the T_z =+3/2→+1/2 GT transitions up to E_x =10.5 MeV deduced from a measurement of the ⁴¹K(³He, t)⁴¹Ca reaction at 0°. These *B*(GT) values are compared with those of the isospinsymmetric T_z =-3/2→-1/2 GT transitions from ⁴¹Ti β decay. In order to make a reliable comparison, high energy resolution is important in the CE reaction. Good angular resolution is also important in order to obtain a well-defined 0° spectrum. In our ⁴¹K(³He, t) measurement, a high energy



FIG. 1. (Color online) Schematic view of the isopin analog states and analogous transitions in the A=41, $T_z=\pm 3/2$ and $\pm 1/2$ isobar system. The Coulomb displacement energies are removed so that the isospin symmetry of the states and transitions become clearer. The type of the reaction or decay is shown along the arrow indicating the transition.

resolution of 35 keV and a good angular resolution were realized even at an intermediate incident energy of 420 MeV (140 MeV/nucleon) by using a magnetic spectrometer in combination with beam matching techniques [20].

II. EXPERIMENT

The GT states are predominantly excited in CE reactions at intermediate energies (≥100 MeV/nucleon) and at forward scattering angles including 0°. This is because of the L=0 nature of GT states and the dominance of the $\sigma\tau$ part of the effective nuclear interaction [21]. The experiment was performed at the Research Center for Nuclear Physics (RCNP), Osaka, by using a 140 MeV/nucleon ³He beam from the K=400 RCNP Ring Cyclotron and the QDD-type Grand Raiden spectrometer [22] placed at 0°. The beam was stopped in a Faraday cup inside the first dipole magnet. In a $({}^{3}\text{He}, t)$ experiment using a magnetic spectrometer, the measurement at 0° is relatively easy, because the magnetic rigidities of the ${}^{3}\text{He}^{2+}$ beam and the singly charged outgoing tritons differ by about a factor of 2. However, for the analysis of intermediate-energy tritons, a large bending power is required for the spectrometer. This requirement makes the Grand Raiden spectrometer ideally suited for the present measurement. The large difference of atomic energy losses of ³He and tritons in a target foil can cause a large energy spread of the outgoing tritons, which deteriorates the spectral resolution. Therefore, a thin self-supporting foil is desirable as a target.

In order to produce a thin ⁴¹K self-supporting target, a thin foil of ⁴¹K₂CO₃ supported by polyvinylalcohol (PVA) [23] was used. The enrichment of the ⁴¹K isotope was 99.2%. The total thickness of the target was approximately 1 mg/cm². Therefore, this target is a mixture of ⁴¹K, carbon isotopes ¹²C and ¹³C with natural abundance 98.9% and 1.1%, respectively, and oxygen isotopes ¹⁶O and ¹⁸O with



FIG. 2. (Color online) The ${}^{41}K({}^{3}He,t){}^{41}Ca$ spectrum at 0° with an angular range up to 0.5°. A high resolution of 35 keV has been achieved. The major GT states are indicated by their excitation energies. For other details, see text.

natural abundance 99.8% and 0.2%, respectively. The final nuclei after the (³He, *t*) charge-exchange reactions are ⁴¹Ca, ¹²N, ¹³N, ¹⁶F, and ¹⁸F, respectively, and their reaction Q values are -0.42, -17.36, -2.24, -15.44, and -1.67 MeV. Owing to the large difference of Q values, states in ⁴¹Ca can be observed without being affected by the strongly excited states in ¹²N and ¹⁶F. Since the Q values of ¹³C and ¹⁸O nuclei are comparable with that of ⁴¹Ca, the ground and excited states of ¹³N and ¹⁸F are expected in the ⁴¹Ca spectrum. The separation and identification of these states and the states of ⁴¹Ca were possible owing to the high energy resolution of this experiment, as will be described below.

The outgoing tritons were momentum analyzed without using any acceptance-defining aperture of the spectrometer. A spectrum covering the angular range of $\approx \pm 20$ mrad horizontally (x) and $\approx \pm 40$ mrad vertically (y) was obtained. The tritons were detected at the focal plane with a multiwire driftchamber system allowing track reconstruction for each ray [24]. The acceptance of the spectrometer was subdivided in the software analysis by using this track information.

An energy resolution far better than the energy spread of the beam was realized by applying the dispersion matching technique [25]. Using the new high-resolution "WS course" [26] for the beam transport and the "faint beam method" for the diagnosis of matching conditions [27,28], an unprecedented resolution of 35 keV [full width at half maximum (FWHM)] was achieved. This improved resolution allowed us to resolve states of ⁴¹Ca up to $E_r = 10$ MeV, as shown in Fig. 2. As can be seen, all strongly excited states are concentrated in the energy region between 4 and 6 MeV. The g.s. of ¹⁸F and the g.s. and the 3.502 MeV state of ¹³N were identified by comparing this spectrum with the spectrum of a ^{nat}MgCO₃+PVA target measured under the same condition as the ${}^{41}K_2CO_3$ +PVA target. They are indicated in Fig. 2. The higher excited states of ¹³N were also identified in the spectrum, but the contributions from these states were very small in the region $E_x < 10.5$ MeV. The possible contributions from the states of ³⁹Ca originating from the ³⁹K nuclei with 0.8% isotopic abundance in the target were examined by comparing with the spectrum from the ${}^{39}K_2CO_3+PVA$ target. The strongest g.s. of ³⁹Ca is expected at ≈ 6.1 MeV in the spectrum shown in Fig. 2, but no noticeable contribution was observed.

In order to accurately determine the scattering angle Θ near 0°, angle measurements in both the *x* direction (θ) and *y* direction (ϕ) are equally important, where Θ is defined by $\Theta = \sqrt{\theta^2 + \phi^2}$. Good θ and ϕ resolutions were achieved by applying the *angular dispersion matching* technique [25] and the "overfocus mode" in the spectrometer [29], respectively. The angle calibration was performed by using a multihole aperture. It is estimated that the angle resolution was better than 8 mrad [29]. The "0° spectrum" in Fig. 2 shows events for scattering angles $\Theta \leq 0.5^\circ$. As will be shown, all prominent states are of L=0 nature.

Accurate E_x values with errors of about 1 keV are given in Ref. [30] for the $\Delta J^{\pi} = 1^+$ GT states of ⁴¹Ca up to 6 MeV (see column 1 of Table I). However, we derived the E_x values independently. The ${}^{26}Mg({}^{3}He,t){}^{26}A1$ and ${}^{24}Mg({}^{3}He,t){}^{24}A1$ reactions have negative Q values of -4.02 MeV and -13.90 MeV, respectively. Accurate E_x values with errors of less than 1 keV are known for the 1⁺ states of ²⁶Al up to 7.9 MeV and for a few states in 24 Al [7]. The 16 O(3 He,t) 16 F reaction has a large negative Q value of -15.44 MeV, and sharp 1⁺ states are known up to 4.65 MeV [31]. On the other hand, the ${}^{13}C({}^{3}He,t){}^{13}N$ and ${}^{18}O({}^{3}He,t){}^{18}F$ reactions have small negative Q values of -2.24 MeV and -1.67 MeV, respectively. States from all of these reactions were observed in the spectrum using a ^{nat}MgCO₃+PVA target [23] measured under the same condition as with the ${}^{41}K_2CO_3 + PVA$ target. The E_x values of the states in the ${}^{41}K({}^{3}He, t)$ spectrum were determined with the help of kinematic calculations. Owing to the large variety of Q values of the states used in the calibration, all E_x values of ⁴¹Ca states shown in column 3 of Tables I and II could be determined by interpolation. The present values agree with the literature values for most states up to $E_x=6$ MeV within 3 keV. The estimated errors are 5 keV for the states up to 8 MeV and 7 keV for higher excited states, unless indicated otherwise.

The yields of individual peaks were derived by applying a peak fitting program using the peak shape of the well separated ³⁹Ca g.s. observed in the ³⁹K(³He, *t*) measurement performed under the same condition as the ⁴¹K measurement. No obvious broadening of the width of ⁴¹Ca states was observed even for states above the proton separation energy S_p of 6.3 MeV.

 $(^{3}\text{He}, t)^{b}$

L

0

0

0

≥1

≥1

≥1

0

≥1

(0)

0

≥1

0

0

0

(≥1)

≥1

(0)

≥1

≥1

0

≥1

≥1

0

0

0

≥1

≥1

(0)

(≥1)

≥1

(0)

≥1

0

≥1

(0)

≥1

≥1

(0)

(0)

 ≥ 1

≥1

0

≥1

0

B (GT)

0.018(2)

0.024(3)

0.026(3)

0.006(2)

0.006(2)

0.018(3)

0.010(4) 0.002(1)

0.008(2)

0.009(2)

0.016(3)

0.011(3)

0.009(2)

0.016(3)

0.006(2)

0.019(4)

0.014(3)

0.004(2)

0.009(3)

0.013(3)

0.023(4)

0.014(3)

TABLE I. States observed in the 41 K(3 He, t) 41 Ca reaction below 7 MeV excitation energy and the corresponding GT transition strengths B(GT). For details of the derivation and the errors of B(GT) values, see text.

TABLE II. Continuation of Table I for the excitation energies between 7 and 10.5 MeV.

 E_x (MeV)

7.225

7.296

7.332

7.37(2)

7.552

7.586

7.639

7.72(2)

7.792

7.854

7.901

7.986

8.046

8.144

8.272

8.347

8.406

8.468

8.515

8.587

8.653

8.702

8.861

8.926

9.013

9.081

9.183

9.230

9.324

9.40(2)

9.590

9.616

9.669

9.771

9.862

9.928

10.03(2)

10.113

10.161

10.194

10.238

10.286

10.339

10.421

 $2J^{\pi}$

 $1^+, 3^+, 5^+$

 $(5-11)^+$

$\frac{B(\text{GT}) \text{ values, see text.}}{\text{Evaluated values}^{\text{a}} \qquad (^{3}\text{He.}t)^{\text{b}}}$					Evaluated values ^a		
					E_x (MeV)	$2J^{2}$	
E_x (MeV)	$2J^{\pi};2T$	E_x (MeV)	L	B(GT)		_	
2.010	3+	2.012	0	0.031(3)	7.296	5-	
2.606	5+	2.607	0	0.020(3)			
2.671	1^{+}	2.676	(0)	0.003(1)			
2.884	7^{+}	2.884	≥1				
3.050	3+	3.050	0	0.004(1)			
3.400	1^{+}	3.400	0	0.067(6)			
3.527	3+	3.526	0	0.034(4)			
3.740	$(3,5)^+$	3.737	0	0.030(3)			
3.846	1^{+}	3.845	0	0.012(2)			
4.097	5+	4.093	0	0.446(33)	7 990(20)	1+ 3+	
4.184	(3,5)	4.182	0	0.034(4)	1.550(20)	1,5	
4.328	$(3 - 13)^{(+)}$	4.330	≥1				
4.415	3+	4.419	0	0.033(4)			
4.550		4.550	0	0.014(2)			
4.728	$(3)^{+}$	4 707	0	0.000(17)	8 402(2)	(5 - 1)	
4.733	$(5)^+$	4.727	0	0.220(17)	0.402(2)	(3-1)	
4.779	$(3)^+$	4.777	0	0.033(5)			
4.816	5+	4.815	0	0.149(12)			
		4.966	0	0.021(3)			
5.095	3+	5.097	0	0.011(2)			
5.283	5+	(5.28)		0.059(17)			
5.411	5+	5.406	(0)	0.045(10)			
5.482	$(3)^{+}$	5.480	0	0.079(7)			
5.649	(5)-	5.652	0	0.067(6)			
5.719	(5)-	5.717	0	0.190(15)			
5.760	$(5)^{+}$	5.756	0	0.051(6)			
5.817	$3^+; 3^c$	5.814	0	$0.148(25)^{d}$			
5.892	1-	5.890	(0)	0.021(4)			
5.976	$(3,5)^+$	5.969	0	0.192(15)			
		6.019	(0)	0.018(4)			
6.325(5)	$(5)^{+}$	6.326	0	0.031(3)			
6.462(10)	$(5-11)^+$	6.464	0	0.030(3)			
		6.544	≥1				
		6.596	0	0.033(3)			
6.647(2)	5-	6.653	≥1				
6.738(7)	$(5-11)^+$	6.744	≥1				
6.807(2)	$(3)^{+}$	6 873	>1				
6.822	1+ 5	0.823	≈1				
6.901(2)	5+	6.904	0	0.007(1)			
6.966(15)	3+,5+	∫6.959	≥1	0.013(2)			
	· · ·	6.984	0	~ /			

^aFrom Ref. [30].

^bPresent work.

^cThe IAS.

^dFor the details of the derivation, see text.

^aFrom Ref. [30].

^bPresent work.

III. DATA ANALYSIS

A. Identification of L=0 Gamow-Teller states

In order to identify L=0 GT states, yields were derived for all states in the spectra with angle cuts $\Theta = 0^{\circ} - 0.5^{\circ}$ (see Fig. 2), $0.5^{\circ}-1.0^{\circ}$, and $1.0^{\circ}-1.5^{\circ}$. The angular distributions of GT states peak at 0° . The relative decrease of the yields of each state in these three cuts was compared with that of the 4.093 MeV state, the most strongly excited state known to have a pure GT nature. It was found that for most of the well-observed states below $E_x=6$ MeV, the ratios of yields in the $1.0^{\circ}-1.5^{\circ}$ and $0^{\circ}-0.5^{\circ}$ angle cuts were similar to the ratio of the 4.093 MeV state within 20%, suggesting that they are all L=0 states. On the other hand, a few very weakly excited states in this region and many weakly excited states above $E_x=6$ MeV had ratios larger than 1.2 (typically 1.3– 1.5), suggesting that these states have $L \ge 1$. They are indicated by the sign " ≥ 1 " in column 4 of Tables I and II.

B. B(GT) evaluation from (³He, t) data

It is known that in CE reactions at 0° the cross sections for GT transitions are approximately proportional to *B*(GT) values [6,32,33]:

$$\frac{d\sigma_{\rm CE}}{d\Omega}(0^{\circ}) \simeq KN_{\sigma\tau} |J_{\sigma\tau}(0)|^2 B(\rm GT)$$
(1)

$$=\hat{\sigma}_{\rm GT}B({\rm GT}),\qquad(2)$$

where $J_{\sigma\tau}(0)$ is the volume integral of the effective interaction $V_{\sigma\tau}$ at momentum transfer q=0, K is the kinematic factor, $N_{\sigma\tau}$ is a distortion factor, and $\hat{\sigma}_{GT}$ is a unit cross section for the GT transition. For (³He, *t*) reactions, the approximate proportionality between the 0° cross sections and the GT transition strengths B(GT) was shown for the B(GT) values ≥ 0.04 from the studies of analogous GT transitions in A=27, T=1/2 mirror nuclei ²⁷Al and ²⁷Si [9] and A=26 nuclei ²⁶Mg, ²⁶Al, and ²⁶Si [11]. Here we use a unit system that gives a value of B(GT)=3 for the β decay of the free neutron.

The product $KN_{\sigma\tau}$ and therefore the unit cross section $\hat{\sigma}_{\rm GT}$ in Eq. (2) changes gradually as a function of mass number A as well as excitation energy [6]. In order to estimate the latter effect, a distorted-wave Born approximation (DWBA) calculation was performed by using the code DW81 [34], assuming a simple $d_{5/2} \rightarrow d_{3/2}$ transition for the excited GT states. The optical potential parameters used were those determined for ²⁸Si at an incident ³He energy of 150 MeV/nucleon [35]. For the outgoing triton channel, by following the arguments given in Ref. [36], the well depths were multiplied by a factor of 0.85 without changing the geometrical parameters of the optical potential. For the effective projectile-target interaction of the composite ³He particle, the form derived by Schaeffer [37] by applying a folding procedure was used. As the interaction strength $V_{\alpha\tau}$ and the range parameter R at 140 MeV/nucleon, we used the values of 2.1 MeV and 1.415 fm, respectively [38]. The calculated 0° cross section decreases by about 5% as E_x increases up to 10 MeV. This result was used to correct the peak intensity of each state.

In order to obtain B(GT) values from Eq. (2), we have to determine the unit cross section $\hat{\sigma}_{\mathrm{GT}}$ or the "unit GT intensity" for the transitions to the states observed in the 41 K(3 He, t) 41 Ca spectrum at 0°, shown in Fig. 2. However, there is no transition whose B(GT) value can be directly determined from a β -decay measurement. Therefore, we assume that the total sum of B(GT) values in the analogous mirror transitions—i.e., $T_z = \pm 3/2 \rightarrow \pm 1/2$ GT transitions observed in the ${}^{41}K({}^{3}He, t)$ reaction and in the ${}^{41}Ti \beta$ decay are the same (see Fig. 1). From Fig. 2, we see that most of the prominent levels are concentrated in the $E_r = 4-6$ MeV region. As discussed, most of them have L=0. A similar concentration of GT strength was reported in the corresponding $E_r = 4 - 6.2$ MeV region of ⁴¹Sc by Honkanen *et al.* [18] in one of the β -decay studies of ⁴¹Ti (for details, see Sec. V A). In addition, the sum of the B(GT) values of $\Sigma B(GT)$ =1.77(17) in this region of 41 Sc, except the GT strength in the IAS, was in good agreement within errors with the sum $\Sigma B(GT) = 1.60(15)$ reported in another β -decay study by Liu et al. [19]. Therefore, the unit GT intensity was obtained by assuming that the total intensity summed over all GT states in this region, after making a correction for excitation energy, corresponds to the B(GT) value of 1.68(12), which is the average of the $\Sigma B(GT)$ values of the two β -decay studies.

The B(GT) values for all excited states, except for the IAS, were calculated by using the same unit GT intensity from their peak intensities after making the excitation energy correction. The obtained B(GT) values are listed in column 5 of Tables I and II, and they are also displayed in Fig. 3(a). The errors quoted for these B(GT) values come from the statistical uncertainties of the experimental data and the ambiguities of B(GT) values in the β -decay measurement. On the other hand, the error associated with the possible isospin asymmetry of the strength in the $E_x=4-6$ MeV region, where the normalization of the B(GT) strength was performed, is not included. Since the proportionality given by Eq. (2) is experimentally established only for transitions with $B(GT) \ge 0.04$ [9], the uncertainty may be larger for weaker transitions.

The transition from the $J^{\pi}=3/2^+$ g.s. of ⁴¹K to the IAS can contain both Fermi (caused by the τ operator) and GT components. In order to extract the GT component in the IAS, we assume that the Fermi transition strength concentrates to this IAS and the strength is B(F)=N-Z=3. In addition, we used the fact that the ratio of GT and Fermi unit cross sections, denoted by R^2 in Ref. [6], is only weakly dependent on mass number A and can be deduced to be 8.7(7) for the A=41 nuclei by an interpolation between the known R^2 values of 6.6(4) and 10.5(5) for A = 26 [11] and A = 58 [14,39] nuclei, respectively. In this way, we estimated a B(GT) value of 0.148(25) in the transition to the IAS as the most probable value. In Ref. [18], an isospin impurity of $(10\pm8)\%$ is suggested for the IAS. It is calculated that this amount of isospin impurity would increase the B(GT) value in the IAS by 0.034 ± 0.028 .



FIG. 3. (Color online) Experimental and shell-model B(GT) distributions. (a) B(GT) distribution from the present ${}^{41}K({}^{3}He, t){}^{41}Ca$ measurement. (b) B(GT) distribution from a shell-model (SM) calculation for all allowed *J*. B(GT) distribution from a SM calculation (c) for J=1/2, (d) for J=3/2, and (e) for J=5/2. Note the change in size of the panels.

IV. GAMOW-TELLER STATES AND B(GT) DISTRIBUTION

A. Experimental results

Since the g.s. of ⁴¹K has $J^{\pi}=3/2^+$, the L=0, $\Delta J^{\pi}=1^+$ GT transitions can reach either $J^{\pi}=1/2^+$, $3/2^+$, or $5/2^+$ states. For most states assigned to be L=0 in the present analysis, corresponding states with either $J^{\pi}=1/2^+$, $3/2^+$, or $5/2^+$ can be found up to 7 MeV excitation energies in Ref. [30], as listed in Table I. Exceptions are the 4.550, 4.966, 5.652, 5.717, and 5.890 MeV states. For them either no J^{π} values or negative parity are assigned. No J^{π} values were given for the

4.550 MeV state, but possible J^{π} values of $(1/2, 3/2, 5/2^+)$ are given for the probable analog state at 4.644 MeV in ⁴¹Sc (see Table III and the discussion of Sec. VA2). The 4.966 MeV state is not known [30], but again its probable analog state with J^{π} values of $1/2^+$ is found at 5.023 MeV. In Ref. [30] it is suggested that the 5.652 and 5.717 MeV states have negative parity. However, they are rather strongly excited in the 0° spectrum, and L=0 is assigned in our analysis. In addition, possible analog states for these states with more or less corresponding strengths are observed at 5.774 and 5.840 MeV, respectively, in the ⁴¹Ti β -decay measurement studying the isospin mirror GT transitions (see Table III). Therefore, we suggest that they are GT states with positive parity. We also suggest a positive parity for the 5.890 MeV state, although it is weaker. The J^{π} values of $5/2^{-}$ are assigned to the 7.296 MeV state in Ref. [30]. Our analysis, however, shows L=0 for the transition to this state, and a positive parity is therefore suggested.

As seen from Fig. 3(a), the main part of the GT strength is concentrated at $E_x=4-6$ MeV, but it is very fragmented. Clustering of states is observed around 4.1, 4.75, 5.7, and 5.95 MeV.

B. B(GT) distribution from the shell model

The shell-model calculations for the A=41 system were carried out in the *sd-fp* model space. Relative to a closed *sd* shell for ⁴⁰Ca, particles are in the *fp* shell and holes are in the *sd* shell. The model space was truncated to one-particle (1p) for the low-lying T=1/2 negative parity states and to twoparticle one-hole (2*p*-1*h*) for the excited T=1/2 and 3/2positive parity states. We used the SDPFMW Hamiltonian [40] that was deduced for this model space and truncation. The computer code OXBASH [41] was used in the calculation. In Fig. 3, the experimentally extracted *B*(GT) values and the shell-model values are compared for excitation energies up to 10 MeV.

The results of shell-model calculation include the average renormalization factor (quenching factor) of 0.6 [13] for the B(GT) values. By comparing Figs. 3(a) and 3(b), we see that the calculated and experimental B(GT) distributions are generally in agreement. The observed concentration of the GT strength at $E_x=4-6$ MeV and the fragmented and weakly excited strengths above 6 MeV are well reproduced by the calculation. The calculated total B(GT) strength up to 6.5 MeV, where the main part of the strength is concentrated, is 2.43, which is in agreement with the corresponding experimental value of 2.12 within 15%. However, the observed fragmentation of the strength in the 4–6 MeV region was not reproduced. The calculated GT strength is concentrated in fewer states.

The calculation predicts that all T=3/2 GT states, except for the IAS, are above $E_x=7.5$ MeV. In addition, all of them have very small B(GT) values. Therefore, it is suggested that most of the GT strength is in the transitions to the T=1/2states. In a naive picture, the structure of ⁴¹K is described by the ⁴⁰Ca core plus two neutrons in the $f_{7/2}$ shell and one proton hole in the $d_{3/2}$ shell. The much weaker T=3/2strength compared to the T=1/2 strength can be understood

TABLE III. Comparison of $T_z+3/2 \rightarrow +1/2$ and $T_z-3/2 \rightarrow -1/2$ analogous GT transitions observed in the 41 K(3 He,t) 41 Ca reaction and the β decay of 41 Ti to 41 Sc, respectively. The comparison is for the transitions to the states below 6.2 MeV, where the main part of the strength is concentrated.

41 K(3 He, t) 41 Ca ^a			⁴¹ Ti µ	Deduced values			
E_x (MeV)	$2J^{\pi};2T$	B(GT)	E_x (MeV)	$2J^{\pi};2T$	B(GT)	$2J^{\pi d}$	ΔE_x (keV)
2.012	3+	0.031(3)	2.096	3+	0.036(7)	3+	84
2.607	5+	0.020(3)	2.667	5^{+}	0.067(13)	5^{+}	60
2.676	1^{+}	0.003(1)	2.719	1^{+}	0.010(3)	1^{+}	43
3.050	3+	0.004(1)				3+	
3.400	1^{+}	0.067(6)	3.411	1^{+}	0.061(7)	1^{+}	11
3.526	3+	0.034(4)	3.563	$(1,3,5)^+$	0.031(6)	3+	37
3.737	$(3,5)^+$	0.030(3)	3.781	(5^{+})	0.029(6)	5^{+}	44
3.845	1^{+}	0.012(2)	3.951(14)	1^{+}	0.009(3)	1^{+}	106
4.093	5+	0.446(33)	4.245(4)	5^{+}	0.360(30)	5+	152
4.182	(3,5)	0.034(4)	4.328(3)	5^{+}	0.016(4)	5+	146
4.419	3+	0.033(4)	4.502(5)	3+	0.014(4)	3+	83
4.550		0.014(2)	4.644(5)	1-	0.015(4)		94
4.727	$(3)^+, (5)^+$	0.220(17)	4.869(4)	5^{+}	0.015(20)	5^{+}	142
4.777	$(3)^{+}$	0.033(5)	4.777(5)	3+	0.056(9)	3+	0
4.815	5^{+}	0.149(12)	4.928(5)	$(1,3,5)^+$	0.250(30)	5^{+}	113
4.966		0.021(3)	5.023(5)	1^{+}	0.031(8)	1^+	57
5.097	3+	0.011(2)	5.084(5)	3+	0.031(8)	3+	-13
5.283	5+	0.059(17)	5.375(4)	5+	0.160(20)	5^{+}	92
5.406	5+	0.045(10)	5.493(5)	1^{+}	0.017(7)		87
5.480	$(3)^{+}$	0.079(7)	5.576(4)	3+, 5+	0.081(19)	3+	96
5.652	$(5)^{-}$	0.067(6)	5.774(4)	$(1,3,5)^+$	0.069(20)	5^{+}	122
5.717	$(5)^{-}$	0.190(15)	5.840(5)	3+,5+	0.320(40)	5^{+}	123
5.756	$(5)^{+}$	0.051(6)	5.886(12)	$(1,3,5)^+$	0.069(19)	5^{+}	130
5.814	3 ⁺ ;3 ^e	0.148(25)	5.939(4)	3 ⁺ ;3 ^e	0.110 ^c	3+	125
5.890	1-	0.021(4)	6.038(25)	$(1,3,5)^+$	0.051(7)	1^+	148
5.969	$(3,5)^+$	0.192(15)	6.083(20)	$(1,3,5)^+$	0.060(8)		114
6.019		0.018(4)	6.133(20)	3+,5+	0.053(7)		114

^aPresent work.

^bFrom Ref. [18].

^cFrom the shell-model calculation; see Ref. [18].

^dSuggested J^{π} values from the comparison.

^eThe IAS.

qualitatively with this picture in combination with the discussion given in Ref. [42] on the possible isospins for singleparticle and single-hole configurations.

The shell-model calculation predicts an additional concentration of T=1/2 strength around $E_x=10$ MeV, which is not observed in the experiment. This predicted GT strength is probably spread over many 5p-3h states that are not in the model space, and thus the GT strength in this high energy region around 10 MeV may be blended into the continuum. A further comparison of strengths for each J value will be made in Sec. V A 3. As will be discussed, the distributions for three different J states are not so well reproduced.

V. COMPARISON OF ANALOGOUS TRANSITIONS

From Fig. 1, we see that the GT transitions studied in the β decay of the T_z =-3/2 nucleus ⁴¹Ti to the T_z =-1/2 nucleus ⁴¹Sc is analogous to those studied in the ⁴¹K(³He,*t*) measurement. In addition, the *M*1 γ transitions from the IAS in ⁴¹Ca to lower excited states are also analogous. Our inter-

est is whether or not similarities of strengths are found for these different analogous transitions.

A. Isospin-symmetric GT transitions

1. GT transition strengths from β -decay measurements

The GT strength distribution with $T_{z} = +3/2 \rightarrow +1/2$ nature studied in the 41 K(3 He, t) reaction can be compared most directly with the result from the isospin-symmetric T_z $=-3/2 \rightarrow -1/2$ transitions studied in the ⁴¹Ti β decay. Owing to the high $Q_{\rm EC}$ value of this decay, the $B({\rm GT})$ values can be studied for states in ⁴¹Sc with relatively high excitation energies. In Figs. 4(b) and 4(c), two B(GT) distributions from the most recent ⁴¹Ti β -decay measurements from Honkanen et al. [18] and from Liu et al. [19], respectively, are displayed. Because of the low S_p value of 1.09 MeV [30] in ⁴¹Sc, both distributions were derived from measurements of delayed protons. It was assumed that possible γ decays to the g.s. of ⁴¹Sc were not important, although such γ decays may have impact in some cases. In reality, the influence seems to be small, because similar B(GT) values are observed in the $T_z = \pm 3/2 \rightarrow \pm 1/2$ isospin mirror transitions to the lowest $J^{\pi} = 3/2^+$ states (see Table III).

Both of these ⁴¹Ti β -decay results show that the main part of the strength is found at excitation energies above 4 MeV. Similar clustering strengths can be seen at 4.25, 4.9, and 5.8 MeV, although the strengths are distributed somewhat differently in each cluster. The differences may be explained by the difference of the resolutions in measuring the energies of delayed protons. They are 30 keV in Ref. [18] [Fig. 4(b)] and 70–100 keV in Ref. [19] [Fig. 4(c)]. The total sums of *B*(GT) values in the 4–6.2 MeV region, excluding the GT strength in the IAS, were 1.77(17) and 1.60(15) in Refs. [18,19], respectively, in agreement within the experimental errors. On the other hand, at higher excitations above 6.2 MeV, we see that these two β -decay results differ substantially.

2. Corresponding GT states in 41 Ca and 41 Sc

The GT strength distribution from our ⁴¹K(³He, *t*) study is shown in Fig. 4(a) for the energy region where the β -decay results are available. It is clear that this distribution is quite similar to the one of Honkanen *et al.* [18] shown in Fig. 4(b) with respect to the gross structure. Furthermore, since the experimental resolutions in this β -decay measurement (30 keV) and the (³He, *t*) reaction (35 keV) are both good and similar, we see a one-to-one correspondence of observed states and the GT transition strengths to them for the energy region below 6.2 MeV, where the main part of the GT strength is concentrated. By combining further the knowledge of J^{π} values of states in ⁴¹Ca and ⁴¹Sc evaluated in Ref. [30], correspondences of states are suggested in Table III, and details are discussed below.

In these two experiments detecting the isospin mirror transitions, we notice that the number of observed states with appreciable B(GT) strength is identical in ⁴¹Ca and ⁴¹Sc, except for the very weak transition to the 3.050 MeV state observed in the (³He, *t*) reaction. The residual interaction de-



FIG. 4. (Color online) Experimental B(GT) distributions from $T_z = \pm 3/2 \rightarrow \pm 1/2$ isospin mirror transitions. (a) B(GT) distribution from the present ⁴¹K(³He,t)⁴¹Ca reaction. (b) B(GT) distribution from the ⁴¹Ti \rightarrow ⁴¹Sc β decay reported in Ref. [18]. (c) B(GT) distribution from the ⁴¹Ti \rightarrow ⁴¹Sc β decay reported in Ref. [19].

pending on T_z —i.e., isospin-asymmetric interaction—can make the GT strength distribution somewhat different. However, since J is a good quantum number, the modification of the distribution is expected only among the same J states. Rather accurate excitation energies and J^{π} values have been evaluated for many of these states [30]. Therefore, the corresponding states can be assigned on the basis of excitation energies, J^{π} values, and the transition strengths from the (³He, t) and β -decay measurements. A clear disagreement of J values is seen only for the corresponding states at 5.406 MeV in ⁴¹Ca and 5.493 MeV in ⁴¹Sc. They are given J^{π} values of 5/2⁺ and 1/2⁺, respectively, in Ref. [30].

Since the number of observed states are practically the same, one may think that the corresponding states can be assigned by the ordering of the excitation energies, if we believe in the isospin symmetry. However, there are several clusters of states, and it is difficult to identify the corresponding states from only the sequence of excitation energies and the consistency of the assigned J^{π} values that sometimes have ambiguities. Clusters are at 4.75, 5.7, and 5.95 MeV in ⁴¹Ca and also in ⁴¹Sc at the corresponding energies [see Figs. 4(a) and 4(b). Clustering states are indicated by parentheses

between columns 3 and 4 in Table III. For these states, the corresponding states should be assigned taking the similarities of transition strengths into account in addition to the consistency of the assigned J^{π} values. As a result, it is suggested that the 4.727 and 4.777 MeV states of the first cluster in ⁴¹Ca correspond to the 4.869 and 4.777 MeV states in ⁴¹Sc, respectively; i.e., the order of the excitation energy of the corresponding states is reversed. Such a reversed order was noticed only for these pairs (see Table III).

For the analog states, the correlation of B(GT) values between analogous $T_z = \pm 3/2 \rightarrow \pm 1/2$ transitions was examined. As we see in Fig. 5, points are distributed more or less along the 45° line, but they are rather scattered, especially for pairs of states with smaller B(GT) values. As a result, we conclude that the gross symmetry of these isospin mirror transitions is rather well preserved, but the fine structures are somewhat different.

3. Strength distributions of J=1/2, 3/2, and 5/2 states

The J^{π} values of the states in ⁴¹Ca and ⁴¹Sc have been evaluated independently for each nucleus [30]. Since the correspondence between analogous GT transitions became apparent in the present analysis, the knowledge of the J^{π} values for a pair of analog GT states can now be combined. The most probable J^{π} values of each pair of states were deduced up to $E_x = 6$ MeV, and they are given in column 7 of Table III. We see that the number of $J^{\pi} = 5/2^+$ states is the largest, while the number of $1/2^+$ states is smaller, suggesting a large contribution of the $d_{5/2}$ hole. The B(GT) distributions of transitions to J=1/2, 3/2, and 5/2 states are shown in Figs. 6 and 7 for nuclei ⁴¹Ca and ⁴¹Sc, respectively. In both figures, the main part of the GT strength is found in the $5/2^+$ states. Among the $3/2^+$ states, the largest GT strength was seen in the T=3/2 IAS at 5.814 MeV. Therefore, in comparison among T=1/2 states, the dominance of the GT strength in $5/2^+$ states is even more prominent.

When these B(GT) distributions with three different J values are compared to the corresponding shell-model results shown in Figs. 3(c)–3(e), we see notable disagreements. In



FIG. 5. A correlation of B(GT) values between analogous $T_z \pm 3/2 \rightarrow \pm 1/2$ transitions. The $-3/2 \rightarrow -1/2$ $B(\text{GT})_\beta$ values are from ⁴¹Ti β decay [18] and the $+3/2 \rightarrow +1/2$ $B(\text{GT})_{\text{het}}$ values are from the present ⁴¹K(³He, t) measurement. Analog states with ideal correlation would follow the dashed line.



FIG. 6. (Color online) The $T_z = +3/2 \rightarrow +1/2 B(\text{GT})$ distributions deduced from the ⁴¹K(³He, *t*) measurement for the ⁴¹Ca states with (a) $J^{\pi} = 1/2^+$, (b) $J^{\pi} = 3/2^+$, and (c) $J^{\pi} = 5/2^+$.

the shell-model calculation, the J=1/2 and 3/2 strengths are stronger than in the experiment, while the J=5/2 strength is weaker. In addition, the fragmentation of the strength, especially that of the J=5/2 strength in the $E_x=5-6$ MeV region is not well reproduced. The larger number of levels, and thus the higher level density, above about 5 MeV is probably due to the states of 4p-3h levels that are not included in the model space truncation. These configurations cannot be reached directly from the 2p-1h, T=3/2 g.s. However, above about 5 MeV, they could mix with the 2p-1h doorway states and cause a spreading of the GT strength.



FIG. 7. (Color online) The $T_z = -3/2 \rightarrow -1/2$ B(GT) distributions deduced from the ⁴¹Ti β -decay measurement for the ⁴¹Sc states with (a) $J^{\pi} = 1/2^+$, (b) $J^{\pi} = 3/2^+$, and (c) $J^{\pi} = 5/2^+$.

4. Estimation of the isospin-asymmetry matrix element

By comparing Figs. 6 and 7, we again notice that the gross features of these two B(GT) distributions for the isospin mirror transitions are similar, but the details are somewhat different. One of the interesting features is that the strengths of two J=5/2 states at about 4.8 MeV and 4.9 MeV in ⁴¹Ca and ⁴¹Sc, respectively, are almost reversed. It is natural to think that these reversed strengths in the T_z $=\pm 3/2 \rightarrow \pm 1/2$ analog transitions are caused by the action of an interaction depending on T_z —i.e., an isospinasymmetric interaction. For simplicity, let us think only of the mixture between these J=5/2 doublet states. Then we can deduce that (A) the isospin-asymmetry matrix elements acting in these isospin mirror states have similar strengths but opposite signs and (B) without the isospin-asymmetry matrix elements the transition strengths to these doublet states are almost the same. We estimate the approximate values of isospin-asymmetry matrix elements on the basis of these assumptions.

The nuclear Hamiltonian is written as $\mathcal{H}=\mathcal{H}_0+\mathcal{H}_{IA}$, where \mathcal{H}_0 and \mathcal{H}_{IA} are the isospin-symmetry (conserving) term and isospin-asymmetry term, respectively. Two eigenstates of \mathcal{H}_0 are denoted by Φ_a and Φ_b . Since the Hamiltonian \mathcal{H}_0 is isospin symmetric, these wave functions are the same in both $T_z=1/2$ (⁴¹Ca) and $T_z=-1/2$ (⁴¹Sc) nuclei. The excitation energies E_a and E_b of observed states are the eigenvalues of the total Hamiltonian \mathcal{H} . Let us think of wave functions Ψ_a and Ψ_b that satisfy

$$(\mathcal{H}_0 + \mathcal{H}_{\mathrm{IA}})\Psi_a = E_a \Psi_a \tag{3}$$

and

$$(\mathcal{H}_0 + \mathcal{H}_{\mathrm{IA}})\Psi_b = E_b\Psi_b,\tag{4}$$

respectively. The form of these equations is formally the same for $T_z = \pm 1/2$ nuclei, but note here that the matrix elements in the isospin-asymmetry term \mathcal{H}_{IM} are different in the $T_z = \pm 1/2$ nuclei [assumption (A)]. Therefore, the wave functions Ψ_a and Ψ_b are different in $T_z = \pm 1/2$ nuclei.

We can formally write the states Ψ_a and Ψ_b in terms of linear combinations of the two states Φ_a and Φ_b as

$$\Psi_a = \alpha \Phi_a + \beta \Phi_b \tag{5}$$

and

$$\Psi_b = \beta \Phi_a - \alpha \Phi_b, \tag{6}$$

where $\alpha^2 + \beta^2 = 1$. Note again here that the coefficients α and β are different in the $T_z = \pm 1/2$ nuclei. Using these relationships, the matrix element of the off-diagonal matrix \mathcal{H}_{IA} —i.e., the isospin-asymmetry term—can be written as

$$\langle \Phi_a | \mathcal{H}_{\text{IA}} | \Phi_b \rangle = \alpha \beta (E_a - E_b). \tag{7}$$

Let us think of the GT transitions caused by the operator $\mathcal{O}=\sigma\tau$ starting from the g.s. Φ_0 of $T_z=\pm 3/2$ nuclei. For simplicity, we assume that the effect of isospin asymmetry in the g.s. is small. The GT transition strength B(GT) is proportional to the squared value of the transition matrix element of $\sigma\tau$ type, as will be discussed in Sec. V B. Therefore, the ratios of the B(GT) values R^0 and R^1 for the transitions to the

two excited states before and after the mixing, respectively, can be expressed as

$$R^{0} = \frac{B^{0}(\mathrm{GT})_{b}}{B^{0}(\mathrm{GT})_{a}} = \frac{|\langle \Phi_{b} | \mathcal{O} | \Phi_{0} \rangle|^{2}}{|\langle \Phi_{a} | \mathcal{O} | \Phi_{0} \rangle|^{2}}$$
(8)

and

$$R^{1} = \frac{B^{1}(\mathrm{GT})_{b}}{B^{1}(\mathrm{GT})_{a}} = \frac{|\langle \Psi_{b} | \mathcal{O} | \Phi_{0} \rangle|^{2}}{|\langle \Psi_{a} | \mathcal{O} | \Phi_{0} \rangle|^{2}},\tag{9}$$

where $B^0(\text{GT})$ and $B^1(\text{GT})$ are the B(GT) values before the mixing of states and the observed B(GT) values after the mixing of states, respectively. By putting Eqs. (5) and (6) into Eq. (9) and using R^0 , the ratio R^1 can be written as

$$R^{1} = \frac{|\langle \beta \Phi_{a} - \alpha \Phi_{b} | \mathcal{O} | \Phi_{0} \rangle|^{2}}{|\langle \alpha \Phi_{a} + \beta \Phi_{b} | \mathcal{O} | \Phi_{0} \rangle|^{2}}$$
(10)

$$\simeq \frac{|\beta - \alpha \sqrt{R^0}|^2}{|\alpha + \beta \sqrt{R^0}|^2}.$$
(11)

The transformation from Eq. (10) to Eq. (11) is not exact when the associated phases are different in the matrix elements $\langle \Phi_a | \mathcal{O} | \Phi_0 \rangle$ and $\langle \Phi_b | \mathcal{O} | \Phi_0 \rangle$.

There is no way to study the ratio R^0 experimentally. However, according to assumption (B), we can put $R^0 \approx 1$; i.e., the doublet states have almost equal B(GT) values without the influence of \mathcal{H}_{IA} . Using the B(GT) values listed in Table III, the values of R^1 are obtained for the doublet states observed at 4.8 MeV in the $({}^{3}\text{He}, t)$ study and also for the doublet states observed at 4.9 MeV in the β -decay study. A set of α and β is obtained for each of these R^1 values using Eq. (11) and the relationship $\alpha^2 + \beta^2 = 1$. The isospinasymmetry matrix element is calculated by putting a set of α and β and the difference of the excitation energies into Eq. (7). As a result, values of -8.3 keV and 7.5 keV are obtained for the $T_z = +3/2 \rightarrow +1/2$ and $T_z = -3/2 \rightarrow -1/2$ transitions, respectively. The signs of them are opposite and the absolute values are nearly the same, which is consistent with assumption (A). It is interesting that relatively small isospinasymmetry interactions of $\approx 8 \text{ keV}$ can make the reversed transition strengths observed for the isospin mirror transitions to the doublet states with $\Delta E \approx 70$ keV.

Isospin mixing was studied at a similar mass of A=37 for a pair of $J^{\pi}=3/2^+$ states. One of them was the T=3/2 IAS in ³⁷K at $E_x=5.051$ MeV and the other was a T=1/2 state lying 31 keV below. The relative proton widths of the two levels measured in a ³⁶Ar(\vec{p}, p_0) resonance reaction implied an isospin-mixing matrix element of 4.8 keV [43]. In addition the analysis of the β^+ -decay study from the g.s. of ³⁷Ca to these two states suggested a value of 5.9 keV [43,44]. It is interesting that similar values are deduced for the isospinasymmetry matrix element obtained here and the isospinmixing matrix element.

5. Difference of excitation energies in ⁴¹Ca and ⁴¹Sc

The Coulomb displacement energy (CDE) is dependent on the configuration of each individual state [45,46]. In order ISOSPIN SYMMETRY OF $T_z = \pm 3/2 \rightarrow \pm 1/2$ GAMOW-TELLER...

to study the CDE as a function of excitation energy, the difference of excitation energies,

$$\Delta E_x = E_x (^{41} \mathrm{Sc}) - E_x (^{41} \mathrm{Ca}), \qquad (12)$$

was calculated for each pair of states. The ΔE_x values are given in the last column of Table III. They are also plotted in Fig. 8 for different J^{π} states.

The ΔE_x values of the $J^{\pi}=3/2^+$ states vary largely from -13 to 125 keV [Fig. 8(b)], and the distribution is scattered as a function of E_x . The ΔE_x distributions of the $1/2^+$ and $5/2^+$ states are less scattered. It is interesting that the minimum at $E_x=5.2$ MeV in the $3/2^+$ distribution [Fig. 8(b)] is common with the local minimum in the distribution of the $5/2^+$ states [Fig. 8(c)].

We see a clear discrete increase of ΔE_x values at around $E_x=3.8$ MeV. In $1/2^+$ and $5/2^+$ distributions, a typical value of about 50 keV in the low-lying region changes suddenly up to more than 100 keV above 4 MeV excitation energies, where the main part of the strength appears. For the transitions in the low-lying region, the main configurations of $d_{3/2} \rightarrow d_{3/2}$ and $f_{7/2} \rightarrow f_{7/2}$ are expected in a naive single-particle model. On the other hand, at the higher excitation energies the main configurations of $d_{5/2} \rightarrow d_{3/2}$ and $f_{7/2} \rightarrow f_{5/2}$ are expected. The sudden increase of ΔE_x values, corresponding to the sudden change of the CDE, suggests a change of the main configurations of the wave functions at around $E_x=3.8$ MeV.

B. M1 γ transitions in ⁴¹Ca

The *M*1 γ transitions from the IAS at E_x =5.817 MeV to low-lying states in ⁴¹Ca are analogous to the corresponding GT transitions observed in the ⁴¹K(³He,*t*) reaction (see Fig. 1).

In order to compare M1 transition strengths and B(GT) strengths of analogous GT transitions, we have to examine the similarities and differences between these transitions. The GT operator has only an IV spin ($\sigma\tau$) term. The GT transition strength B(GT) reduced in isospin [47] is given by

$$B(\text{GT}) = \frac{1}{2J_i + 1} \frac{1}{2} \frac{C_{\text{GT}}^2}{2T_f + 1} [M_{\text{GT}}(\sigma\tau)]^2, \qquad (13)$$

where C_{GT} is the isospin Clebsch-Gordan (CG) coefficient $(T_iT_{zi}1\pm 1 | T_fT_{zf})$ with $T_{zf}=T_{zi}\pm 1$. The matrix element $M_{\text{GT}}(\sigma\tau)$ denotes the GT transition matrix element of $\sigma\tau$ type.

In addition to the IV spin $(\sigma \tau)$ term, the *M*1 operator has an IV orbital $(\ell \tau)$ term and an IS term. Since the *M*1 transitions of interest here are between the T=3/2 IAS and the T=1/2 GT states, only the IV terms can contribute. In which case, the *M*1 transition strength B(M1) reduced in isospin [47] is given by

$$B(M1) = \frac{1}{2J_i + 1} \frac{3}{4\pi} \mu_N^2 \frac{C_{M1}^2}{2T_f + 1} \left[g_\ell^{\text{IV}} M_{M1}(\ell \tau) + g_s^{\text{IV}} \frac{1}{2} M_{M1}(\sigma \tau) \right]^2, \qquad (14)$$

where C_{M1} expresses the isospin CG coefficient



FIG. 8. The difference of excitation energies ΔE_x for the corresponding states in ⁴¹Sc and ⁴¹Ca. The ΔE_x values are plotted for pairs of states with (a) J=1/2, (b) 3/2, and (c) 5/2. A clear sudden increase of ΔE_x values is observed at $E_x=3.8$ MeV for the pairs of J=1/2 and 5/2 states. The dotted lines are drawn to guide the eye.

 $(T_i T_{zi} 1 \ 0 | T_f T_{zf})$ with $T_{zf} = T_{zi}$. The matrix elements $M_{M1}(\sigma\tau)$ and $M_{M1}(\ell\tau)$ denote the $\sigma\tau$ -type and $\ell\tau$ -type components of the M1 transition matrix element, respectively. The IV combinations of spin and orbital g factors are expressed by g_s^{IV} $= \frac{1}{2}(g_s^{\pi} - g_s^{\nu})$ and $g_\ell^{IV} = \frac{1}{2}(g_\ell^{\pi} - g_\ell^{\nu})$, respectively. Using the bare spin and orbital g factors of protons and neutrons—i.e., g_s^{π} = 5.586, $g_s^{\nu} = -3.826$ and $g_l^{\pi} = 1$, $g_l^{\nu} = 0$, we get $g_s^{IV} = 4.706$ and $g_\ell^{IV} = 0.5$. Due to the large value of the coefficient g_s^{IV} , it is expected that the $\sigma\tau$ term is usually larger than the $\ell\tau$ term [48,49]. Therefore, corresponding strengths are expected for the analogous GT and M1 transitions (for details, see, e.g., Ref. [47]).

The relative intensities I_{γ} and the energies E_{γ} of the γ transitions from the IAS to the states at E_x =4.728, 4.097, 3.740, 3.400, and 2.606 MeV are given in Ref. [30] and are listed in column 4 of Table IV. Using these values, the transition strengths proportional to the *M*1 transition strengths B(M1) can be deduced under the assumption that the *E*2 and *M*1 mixing ratios δ are small for these transitions. The proportionality is given (see, e.g., Ref. [50]) by

$$B(M1) \propto \frac{1}{E_{\gamma}^3} I_{\gamma}.$$
 (15)

The relative strengths of the B(M1) values for these five M1 transitions are listed in column 5 of Table IV, and the relative strengths for the corresponding GT transitions are given in column 7, where the strongest M1 and GT strengths to the 4.097 MeV are normalized to unity. As seen, the relative intensities of the analogous M1 and GT transitions are in

TABLE IV. Comparison of analogous M1 and GT transitions in A=41 isobars. The γ transitions in ⁴¹Ca from the 5.817 MeV IAS to lower-lying states are analogous to the GT transitions to these states from the g.s. of ⁴¹K. Relative transition strengths are compared for these transitions assuming that these γ transitions are of pure M1 nature.

States in ⁴¹ Ca		γ transitions in ⁴¹ Ca			GT transitions to ⁴¹ Ca	
$E_x (\text{MeV})^{\text{a}}$	$2J^{\pi b}$	E_{γ} (MeV)	Intensity ^c	B(M1) ratio	B(GT)	B(GT) ratio
2.606	5+	3.211	23(4)	0.04(1)	0.020(3)	0.04(1)
3.400	1^{+}	2.417	30(4)	0.11(1)	0.067(6)	0.15(1)
3.740	5+	2.077	15(9)	0.09(5)	0.030(3)	0.07(1)
4.097	5^{+}	1.720	100(8)	1.00(8)	0.446(33)	1.00(7)
4.728	5+	1.089	21(4)	0.83(16)	0.220(17)	0.49(4)

^aFrom Ref. [30].

^bFrom the present analysis.

^cRelative intensity from Ref. [30].

good agreement, except for the transitions to the 4.728 MeV state. From the similarity of corresponding GT and *M*1 transitions strengths, it is suggested that the contributions of the $\ell \tau$ term in the transitions are relatively small. It was found that such contributions of $\ell \tau$ terms are very large for the deformed nuclei in the A=23-25 mass region, where large enhancements or suppressions of *M*1 strengths compared to the analogous GT transition strengths were observed [8,51,52].

VI. SUMMARY

Isospin analogous GT and M1 transitions were compared for A=41 nuclei. Especially, $T_z=\pm 3/2 \rightarrow \pm 1/2$ mirror GT transitions were compared and analyzed from various points of view.

The $T_7 = +3/2 \rightarrow +1/2$ GT transitions were measured by using the 41 K(3 He, t) 41 Ca reaction at 0° and at an intermediate incident energy of 140 MeV/nucleon. Thanks to the high energy resolution of 35 keV, states up to $E_r = 10$ MeV in ⁴¹Ca were clearly separated. This energy resolution of better than $\Delta E/E = 10^{-4}$ and also a good angle resolution of the scattering angle around 0° were achieved by the implementation of complete beam matching techniques between the magnetic spectrometer and the beamline. Highly fragmented discrete states were observed and the excitation energies of these states could be determined with an accuracy of less than 7 keV up to the excitation energy of 10 MeV. The B(GT) values in the (³He, t) reaction were calibrated by using B(GT) values derived in the GT β decay from ⁴¹Ti to ⁴¹Sc by assuming mirror symmetry of the $T_z = \pm 3/2$ $\rightarrow \pm 1/2$ transitions. A DWBA calculation was used to correct for the excitation energy dependence of the cross section. It was found that the main part of the GT strength is concentrated in the energy region between 3 and 6 MeV. The fragmentation of the GT strength was not so well reproduced by a shell-model calculation, suggesting more complicated structures of the A = 41 nuclei that extend over *sd*- and *f*-shell regions.

The mirror symmetric $T_z = -3/2 \rightarrow -1/2$ GT transitions can be studied in the β decay of ⁴¹Ti to ⁴¹Sc. Two independent β -decay measurements have been reported and the obtained B(GT) distributions were significantly different in the energy region above $E_x = 6$ MeV. If isospin is a good quantum number, the transition strengths of the $T_z = \pm 3/2$ $\rightarrow \pm 1/2$ GT mirror transitions should not be much different. It was found that the general feature of the B(GT) distribution deduced by Honkanen *et al.* [18] was similar to that observed in the present ${}^{41}K({}^{3}He,t)$ measurement. In addition, the good energy resolution achieved by Honkanen *et al.* in the delayed proton measurement after the β decay was comparable with our resolution.

Through a detailed comparison of the $T_z = \pm 3/2 \rightarrow \pm 1/2$ GT transitions, one-to-one correspondences of analog states in ⁴¹Ca and ⁴¹Sc could be identified up to the excitation energy of 6 MeV, where the main transition strength was observed. In the identification, it was assumed that the strengths of the analogous transitions should be similar. The GT states in ⁴¹Ca and ⁴¹Sc can have a J^{π} value of either $1/2^+$, $3/2^+$, or $5/2^+$, but the J^{π} assignment of individual states in ⁴¹Ca and ⁴¹Sc was not always clear. As a result of the identification of analog states, the most probable J^{π} values were assigned for each pair of analog states by combining the independent J^{π} information of the pair of corresponding states.

On the basis of the J^{π} assignment, the B(GT) strength distribution could be separately obtained for the three J^{π} values. It was found that the main part of the GT strength was carried by the $5/2^+$ states. The gross features of the GT strength distributions for the three J states were similar for the isospin analogous $T_z \pm 3/2 \rightarrow \pm 1/2$ transitions, but the details were somewhat different. From the difference of the distributions, isospin-asymmetry matrix elements of $\approx 8 \text{ keV}$ were deduced. The difference of excitation energies was studied for analog states as a function of excitation energy. An energy increase of about 50 keV was observed at E_x = 3.8 MeV, suggesting a change of configuration in the wave function of the GT states below and above this energy.

The strengths of *M*1 transitions in ⁴¹Ca from five excited states to the IAS were compared with the analogous GT transition strengths derived from the 41 K(3 He,t) 41 Ca study. It was found that ratios of the *M*1 and GT transition strengths, except for one transition, were similar. This suggests that the

contribution from the $\ell \tau$ term, which is inherent to an M1 transition and has no corresponding term in a GT transition, is small in these M1 transitions.

ACKNOWLEDGMENTS

The 41 K(3 He, t) 41 Ca experiment was performed at RCNP, Osaka University, under the Experimental Program E195. The authors are grateful to the accelerator group of RCNP, especially to Professor T. Saito and Dr. S. Ninomiya, for

- Isospin in Nuclear Physics, edited by D.H. Wilkinson (North-Holland, Amsterdam, 1969).
- [2] A. Bohr and B. Mottelson, *Nuclear Structure* (Benjamin, New York, 1969), Vol. 1, and references therein.
- [3] A. Bohr and B. Mottelson, *Nuclear Structure* (Benjamin, New York, 1975), Vol. 2, Chap. 6, and references therein.
- [4] F. Osterfeld, Rev. Mod. Phys. 64, 491 (1992), and references therein.
- [5] J. Rapaport and E. Sugarbaker, Annu. Rev. Nucl. Part. Sci. 44, 109 (1994).
- [6] T.N. Taddeucci, C.A. Goulding, T.A. Carey, R.C. Byrd, C.D. Goodman, C. Gaarde, J. Larsen, D. Horen, J. Rapaport, and E. Sugarbaker, Nucl. Phys. A469, 125 (1987), and references therein.
- [7] P.M. Endt, Nucl. Phys. A521, 1 (1990); A633, 1 (1998), and references therein.
- [8] Y. Fujita et al., Phys. Rev. C 66, 044313 (2002).
- [9] Y. Fujita et al., Phys. Rev. C 59, 90 (1999).
- [10] R. Madey, B.S. Flanders, B.D. Anderson, A.R. Baldwin, C. Lebo, J.W. Watson, S.M. Austin, A. Galonsky, B.H. Wildenthal, and C.C. Foster, Phys. Rev. C 35, 2011 (1987); 36, 1647 (1987).
- [11] Y. Fujita et al., Phys. Rev. C 67, 064312 (2003).
- [12] B.D. Anderson et al., Phys. Rev. C 54, 602 (1996).
- [13] B.A. Brown and B.H. Wildenthal, At. Data Nucl. Data Tables 33, 347 (1985).
- [14] Y. Fujita et al., Eur. Phys. J. A 13, 411 (2002).
- [15] G. Audi and A.H. Wapstra, Nucl. Phys. A565, 1 (1993); A595, 409 (1995).
- [16] R.G. Sextro, R.A. Gough, and J. Cerny, Phys. Rev. C 8, 258 (1973).
- [17] Z.Y. Zhou, E.C. Schloemer, M.D. Cable, M. Ahmed, J.E. Reiff, and J. Cerny, Phys. Rev. C 31, 1941 (1985).
- [18] A. Honkanen, P. Dendooven, M. Huhta, G. Lhersonneau, P.O. Lipas, M. Oinonen, J.-M. Parmonen, H. Penttilä, K. Peräjärvi, T. Siiskonen, and J. Äystö, Nucl. Phys. A621, 689 (1997).
- [19] W. Liu et al., Phys. Rev. C 58, 2677 (1998).
- [20] Y. Fujita et al., Nucl. Phys. A687, 311c (2001).
- [21] W.G. Love and M.A. Franey, Phys. Rev. C 24, 1073 (1981).
- [22] M. Fujiwara *et al.*, Nucl. Instrum. Methods Phys. Res. A **422**, 484 (1999).
- [23] Y. Shimbara *et al.*, Nucl. Instrum. Methods Phys. Res. A **522**, 205 (2004).
- [24] T. Noro *et al.*, RCNP (Osaka University), Annual Report, 1991, p.177.

their effort in providing a high-quality and stable ³He beam indispensable for the realization of matching conditions required for achieving good energy and angular resolution. Y.F. thanks Professor J. Äystö (Jyväskylä, Finland) for valuable discussions and Dr. F.D. Smit (iThemba LABS, South Africa) for his comments in preparing the manuscript. This work was supported in part by Monbukagakusho, Japan, under Grant No. 15540274. B.A.B. acknowledges support from NSF Grant No. PHY-0244453.

- [25] Y. Fujita, K. Hatanaka, G.P. A. Berg, K. Hosono, N. Matsuoka, S. Morinobu, T. Noro, M. Sato, K. Tamura, and H. Ueno, Nucl. Instrum. Methods Phys. Res. B 126, 274 (1997), and references therein.
- [26] T. Wakasa *et al.*, Nucl. Instrum. Methods Phys. Res. A **482**, 79 (2002).
- [27] H. Fujita *et al.*, Nucl. Instrum. Methods Phys. Res. A 484, 17 (2002).
- [28] Y. Fujita et al., J. Mass Spectrom. Soc. Jpn. 48, 306 (2000).
- [29] H. Fujita *et al.*, Nucl. Instrum. Methods Phys. Res. A 469, 55 (2001).
- [30] J.A. Cameron and B. Singh, Nucl. Data Sheets **94**, 429 (2001), and references therein.
- [31] D.R. Tilley, H.R. Weller, and C.M. Cheves, Nucl. Phys. A564, 1 (1993).
- [32] C.D. Goodman, C.A. Goulding, M.B. Greenfield, J. Rapaport, D.E. Bainum, C.C. Foster, W.G. Love, and F. Petrovich, Phys. Rev. Lett. 44, 1755 (1980).
- [33] W.G. Love, K. Nakayama, and M.A. Franey, Phys. Rev. Lett. 59, 1401 (1987).
- [34] DW81, a DWBA computer code by J.R. Comfort (1981) and updated version (1986), an extended version of DWBA70 by R. Schaeffer and J. Raynal (1970).
- [35] T. Yamagata, H. Utsunomiya, M. Tanaka, S. Nakayama, N. Koori, A. Tamii, Y. Fujita, K. Katori, M. Inoue, M. Fujiwara, and H. Ogata, Nucl. Phys. A589, 425 (1995).
- [36] S.Y. van der Werf, S. Brandenburg, P. Grasdijk, W.A. Sterrenburg, M.N. Harakeh, M.B. Greenfield, B.A. Brown, and M. Fujiwara, Nucl. Phys. A496, 305 (1989).
- [37] R. Schaeffer, Nucl. Phys. A164, 145 (1971).
- [38] R.G.T. Zegers *et al.*, Phys. Rev. Lett. **90**, 202501 (2003); S.Y. van der Werf and R.G.T. Zegers (private communication).
- [39] H. Fujita, Ph.D. thesis, Osaka University, 2002 (unpublished);H. Fujita *et al.*, (unpublished).
- [40] E.K. Warburton, J.A. Becker, and B.A. Brown, Phys. Rev. C 41, 1147 (1990).
- [41] B.A. Brown, A. Etchegoyen, and W.D.M. Rae, computer code OXBASH, MSU Cyclotron Laboratory Report No. 524, 1986.
- [42] A. Bohr and B. Mottelson, *Nuclear Structure* (Benjamin, New York, 1969), Vol. 1, illustrative example to Sec. 3-1.
- [43] N.I. Kaloskamis, A. García, S.E. Darden, E. Miller, W. Haeberli, P.A. Quin, B.P. Schwartz, E. Yacoub, and E.G. Adelberger, Phys. Rev. C 55, 640 (1997).
- [44] A. García, E.G. Adelberger, P.V. Magnus, H.E. Swanson, D.P. Wells, F.E. Wietfeldt, O. Tengblad, and ISOLDE Collabora-

tion, Phys. Rev. C 51, R439 (1995).

- [45] J.A. Nolen and J.P. Schiffer, Annu. Rev. Nucl. Sci. 19, 471 (1969).
- [46] N. Auerbach, Phys. Rep. 98, 273 (1983).
- [47] Y. Fujita, B.A. Brown, H. Ejiri, K. Katori, S. Mizutori, and H. Ueno, Phys. Rev. C 62, 044314 (2000), and references therein.
- [48] E.K. Warburton and J. Weneser, in *Isospin in Nuclear Physics*, edited by D.H. Wilkinson (North-Holland, Amsterdam, 1969), Chap. 5, and references therein.
- [49] S.S. Hanna, in *Isospin in Nuclear Physics*, edited by D.H. Wilkinson (North-Holland, Amsterdam, 1969), Chap. 12, and references therein.
- [50] H. Morinaga and T. Yamazaki, *In-Beam Gamma-Ray Spectroscopy* (North-Holland, Amsterdam, 1976), and references therein.
- [51] Y. Shimbara et al., Eur. Phys. J. A 19, 25 (2004).
- [52] Y. Fujita et al., Phys. Rev. Lett. 92, 062502 (2004).