

Title	Biochemical studies on the transcription antitermination function of the N gene product of coliphage $\lambda$
Author(s)	Ishii, Shunsuke
Citation	
Issue Date	
Text Version	ETD
URL	<a href="http://hdl.handle.net/11094/27746">http://hdl.handle.net/11094/27746</a>
DOI	
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Biochemical studies on the transcription antitermination function..

of the N gene product of coliphage  $\lambda$ .

Shunsuke Ishii

## SUMMARY

An in vitro assay has been developed for the activity of the N gene product (pN) of bacteriophage  $\lambda$ , based on the N-dependent stimulation of trp mRNA synthesis from the DNA templates of  $\lambda$ trp transducing phages. Using this assay system, we show (a) that the stimulation of trp mRNA synthesis by pN requires the nutL sequence on the  $\lambda$  chromosome and (b) that pN can participate in vitro in the formation of RNA polymerase complexes that are competent for the transcription antitermination observed in our system.

## INTRODUCTION

Efficient transcription of the delayed-early genes in phage  $\lambda$  is dependent upon the antitermination activity of the phage gene N product (pN). In the absence of pN function, transcription from the  $p_L$  and  $p_R$  phage promoters is terminated at sites,  $t_{L1}$  and  $t_{R1}$ , through the action of the host-specified rho factor (Roberts, 1969).

The mechanism of the antitermination activity exerted by pN is not clearly understood. Some evidence indicates that pN at some stage interacts with RNA polymerase to effect antitermination (Georgopoulos, 1971; Ghysen and Pironio, 1972; Epp and Pearson, 1976; Baumann and Friedman, 1976; Sternberg, 1976). Salstrom and Szybalski (1978) have suggested that this association may precede the initiation of transcription, or possibly occur during transcription, at the nut site on the  $\lambda$  chromosome. From analyses of genetic hybrids formed in crosses between related bacteriophages with diverse N-like gene functions (Salstrom and Szybalski, 1978; Salstrom *et al.*, 1979), and from the genetic and sequence analyses of nutL<sup>-</sup> phage mutants (Salstrom and Szybalski, 1978; Rosenberg *et al.*, 1977) defective in the recognition of pN for antitermination, the nutL site is inferred to just to the left of the  $p_L$  promoter.

Here we demonstrate *in vitro* that (a) pN stimulates trp mRNA synthesis on  $\lambda$ trp DNA by overcoming transcription termination barriers within the translocated trp operon segment, and (b) like *in vivo*, pN activity in this assay depends on the integrity of the nutL site of  $\lambda$ .

## MATERIALS AND METHODS

a) Strains. All bacteria, bacteriophages and plasmids employed in this study are listed in Table 1. The rif<sup>r</sup> mutant of E.coli A19trpAE1 was spontaneously isolated and mapped at the rpoB locus on the E.coli chromosome by Plkc transduction.

b) In vitro transcription system and RNA assay. The reaction mixture (0.42 ml) contained 40 mM Tris-HCl (pH 7.8), 6 mM  $\beta$ -mercaptoethanol, 50 mM KCl, 10 mM MgCl<sub>2</sub>, 0.2 mM each of ATP, GTP, CTP and [<sup>3</sup>H]-UTP (0.60 mCi/ $\mu$ mole), 1 mM K<sub>2</sub>HPO<sub>4</sub>, 5  $\mu$ g of phage DNA and various amounts of nucleic acid-free S-100 protein. In the experiments employing purified E.coli RNA polymerase, the reaction mixture (0.2 ml) contained 40 mM Tris-HCl (pH 7.8), 6 mM  $\beta$ -mercaptoethanol, 50 mM KCl, 10 mM MgCl<sub>2</sub>, 0.2 mM each of ATP, GTP, CTP and [<sup>3</sup>H]-UTP (1.2 mCi/ $\mu$ mole), 5  $\mu$ g of phage DNA and 3  $\mu$ g of E.coli RNA polymerase. After incubation, reaction mixture was treated with 5  $\mu$ g of pancreatic DNaseI for 5 min at 30° C and followed by the addition of 0.1 ml of 5 % SDS. The mixture was then quickly chilled to 0 °C. The RNA was extracted with phenol, precipitated with ethanol at -20° C and dissolved in 3 X SSC (1 X SSC: 0.15 M NaCl-0.015 M sodium citrate) containing 1 mM EDTA. The trp mRNA was assayed as described previously (Ishii and Imamoto 1976). The N mRNA was determined by DNA-RNA hybridization with l-strand DNA of  $\lambda$ bio256 and  $\lambda$ bio3h-1 phages.

c) Preparation of nucleic acid-free S-100 extracts. Cells of A19trpAE1, A19trpAE1rif<sup>r</sup> and A19trpAE1(pYS5) were grown at 30° C in L-broth to 4 x 10<sup>8</sup> cells/ml. The cultures were quickly heated to 42° C for 12 min and then rapidly cooled to 37° C for 10 min further incubation. In the case

of phage infection, cultures of  $\lambda 19\text{trpAEl}$  were grown to  $4 \times 10^8$  cells/ml in L-broth (30° C) and infected at a multiplicity of 5. After a 15 min (stationary) adsorption period at 30°C, the cultures were incubated aerobically for 10 min at 30°C. The cultures were mixed with sufficient crushed ice to lower the temperature below 4°C within a few seconds. The cells were harvested by centrifugation, washed with buffer<sub>I<sub>0.06</sub></sub> [10 mM Tris-HCl(pH 7.8), 60 mM KCl, 10 mM MgCl<sub>2</sub>, 6 mM β-mercaptoethanol] and then suspended in the same buffer. Cells were lysed by sonication and the lysate was centrifuged for 3 hrs at 37,000 rpm in the SW50.1 rotor of the Beckman model L ultracentrifuge. The clear supernatant was removed to yield the S-100. The extract was adjusted to 0.4 M KCl and loaded onto a DEAE-cellulose column previously equilibrated with buffer<sub>I<sub>0.4</sub></sub> [10 mM Tris-HCl(pH 7.8), 0.4 M KCl, 10 mM MgCl<sub>2</sub>, 6 mM β-mercaptoethanol]. The column was washed with the same buffer, and fractions containing the yellow color were pooled to yield the nucleic acid-free S-100. After addition of solid ammonium sulfate (56.1 g/100 ml of nucleic acid-free S-100), the resulting precipitate was resolved in buffer<sub>I<sub>0.06</sub></sub> and dialyzed against buffer<sub>I<sub>0.06</sub></sub>. After dialysis, the extract can be stored for up to 6 months at -70°C before pN activity is assayed.

d) Preparation of phage DNA. Phage DNA's were prepared as described previously (Ishii and Imamoto, 1978) and dissolved in a 10 mM Tris-HCl buffer (pH 7.8) after dialysis against the solution, and stored at 4°C. Isolated  $\lambda c190$ ,  $\lambda \text{bio}256$  and  $\lambda \text{bio}3h-1$  phage DNA strands were prepared by the poly (U,G)/CsCl technique (Bøvre et al., 1971).

e) The other materials and methods used in the present experiments were as described elsewhere (Ishii and Imamoto, 1976).

## RESULTS AND DISCUSSION

In vitro transcription system responding to pN function. Quantitative estimates of pN activity have been accomplished by using DNA-directed transcription-translation systems to synthesize particular proteins. Two such systems have been reported: in one, pN permits transcription originating at the  $p_R$  promoter to extend into gene  $R$ , resulting in the synthesis of endolysin (Greenblatt, 1972) ; and in the other, pN allows anthranilate synthetase synthesis as a result of readthrough from  $p_L$  into trp on a  $\lambda$ trp DNA template (Dottin and Pearson, 1973). We have developed a transcriptional assay for pN function in vitro based on the pN-dependent stimulation of trp mRNA synthesis from a  $\lambda$ trp DNA template, a reaction catalyzed by a nucleic acid-free extract of soluble protein (S-100) from E. coli. In the  $\lambda$ trp phages, the distal part of the trp operon (trp D,C,B,A) is fused to the N operon such that transcription of the trp operon segment depends on the  $p_L$  promoter (see Fig.1) and occurs in the absence of translational activities.

Transcription initiating at the  $p_L$  promoter of  $\lambda$ trp48 in the absence of N function is frequently terminated at at least two sites inside the trp operon segment (Nakamura et al., 1978a; Nakamura et al., 1978b). The first such site is located in the trpD gene and the second site is in the trpC gene (see Fig.1). However, termination of transcription at these sites in vivo appears to be leaky, since some expression of the distal trp genes is observed. In an in vitro transcription system employing purified E. coli RNA polymerase, this background or "read-through" transcription of trp genes C,B and A can be almost entirely eliminated (a) by deletion of the low level  $p_{2trp}$  promoter and (b) by the addition of a relatively high concentration (35  $\mu$ g/ml) of purified rho factor. In this system, the overall rate of trp D mRNA synthesis was reduced by about 30 % of the level observed in the absence

of rho factor (data not shown). Most of the trpD mRNA molecules produced in the presence of rho factor were short, sedimenting as a symmetrical peak of about 16 S (Fig.2a). Sedimentation of the RNA synthesized in the absence of rho factor, however, revealed the presence of an additional species of trpD mRNA of about 23 S (Fig.2b). This indicates the presence of at least one site where termination of transcription is facilitated by rho function at or near the middle of trp C-D region of  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub>.

When the nucleic acid-free S-100 extract of E.coli A19trpAEl was used instead of purified RNA polymerase, the overall synthesis of trp mRNA directed by  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> DNA was severely depressed (Table 1a). In these conditions the synthesis of N mRNA from the promoter-proximal portion of the N-trp operon proceeded at a virtually normal rate. The observed low-level of trp mRNA synthesis is therefore presumed to result from premature termination of p<sub>L</sub>-initiated transcription at the specific site near to the trp D-E boundary, where rho factor (endogenously supplied) could function more efficiently in the S-100 extract than in the purified system. The data presented in Table 1a reveals a remarkable stimulation (about 40-fold) of trp mRNA synthesis when the S-100 extract was used derived from the thermo-induced E.coli A19trpAEl carrying pYS5, a colE1-type plasmid constructed in vitro which carries the EcoRI-generated cIII-N-cI-O segment of phage  $\lambda$ cI857 (ts) croam6 DNA. The two-fold higher level of trpCBA mRNA relative to trp D mRNA reflects coordinate synthesis of trp mRNA from the whole region of the translocated trp operon segment in  $\lambda$ trp48 (cf. Nakamura et al., 1978a). This is consistent with antitermination exerted at otherwise termination sites within the trp operon segment.

Several lines of evidence indicate that it is pN itself that is stimulating



overall synthesis of trp mRNA in the in vitro system. First, the stimulatory effect of trp mRNA synthesis on  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> DNA was also observed with the nucleic acid-free S-100 extract from  $\lambda$  or  $\lambda$ cI857tof12 -infected A19trpAEl (Table 1). In these cases, however, production of mRNA from the promoter-distal trp genes was less than that seen with the extract from the strain carrying pYS5. Second, stimulation of trp mRNA synthesis was barely detectable in the S-100 extracts of A19trpAEl cells infected with  $\lambda$ Nam7am53cI857tof12 or with  $\lambda$ imm21 (Table 1). Third, the S-100 extract of  $\lambda$ Nts8-infected A19 trpAEl showed little or no stimulation of trp mRNA synthesis at the non-permissive temperature (40° C), but at 30° C stimulated trpCBA mRNA as well as  $\lambda$  or  $\lambda$ tof12 (see Table 1c). Therefore, we conclude that the observed stimulation of trp mRNA synthesis is due to the activity of pN rather than some other  $\lambda$ -coded protein whose synthesis is dependent on pN.

Comparing the relative factors of stimulation of trpD and trpCBA mRNA synthesis by the various pN-containing S-100 extracts on the  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> DNA template, we further conclude that the pYS5 plasmid carrying strain, designed to over-produce pN after temperature induction of this strain, indeed yields about 25 times more pN activity than in the thermoinduced  $\lambda$ cI857croam6 lysogen (unpublished data). Thus, using the pN-containing S-100 extract from pYS5-carrying cells, we compared the effectiveness of the  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> DNA template relative to several other  $\lambda$ trp DNA templates (see Fig. 1). The results presented in Table 2 reveal that the factor of the stimulation caused by the pN-containing extract was greater on templates with fewer transcriptional termination signals. The decreasing order of stimulation was :  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub>,  $\lambda$ trp48,  $\lambda$ trp60-3 (which retains a strong t<sub>N</sub>-site (Nakamura et al., 1978b), and  $\lambda$ trp34 (which bares two t<sub>L</sub> sites and at least

one  $t_N^-$  site) (see Fig.1). We note that in the experiment of Table 2, the individual reactions were carried out in S-100 extracts of  $\lambda 19trpAE1$  to which a smaller aliquote of the pN-containing S-100 extract was added. The amount of trp mRNA synthesized on  $\lambda trp48\Delta p_{2trp}$  DNA in the pN-free S-100 extract was nearly proportional to the amount of pN-containing extract added (see Fig. 4a). Further addition of the pN-containing extract (more than 1 mg/ml) caused a decrease in the level of trp mRNA synthesis. [ Under these conditions, the initiation frequency at  $p_L$  in  $\lambda trp$  DNA was not markedly affected by the addition of increasing amount of the pN-containing extract to the pN-free extract, indicating an excess of RNA polymerase in the latter extract (cf. N mRNA levels in Table 1a and unpublished data).]

Interaction of pN with RNA polymerase. Using the foregoing transcription assay system, we (a) demonstrated the functional association of pN and RNA polymerase in vitro, an interaction which could be catalyzed by the nutL sequences of  $\lambda$ , and (b) ascertained in vitro the effect of a nutL<sup>-</sup> mutation, a defect in the leftward site of N utilization. In the experiments described below, increasing amounts of the pN-containing S-100 extract of thermoinduced  $\lambda 19trpAE1(rif^+)$  carrying pYS5 were added to a constant amount of the pN-free S-100 extract of  $\lambda 19trpAE1(rif^r)$ , in the presence or absence of rifampicin, and the stimulation of trpmRNA synthesis from the  $\lambda trp48\Delta p_{2trp}$  DNA template was ascertained. First, in a control experiment, RNA polymerase from  $\lambda 19trpAE1$  bearing the rif<sup>r</sup> mutation was found to be quite resistant to rifampicin over the entire range of the drug concentrations tested (Fig. 3a); this is in contrast to the enzyme from the isogenic rif<sup>+</sup> strain and its pYS5-carrying derivative, which were found to be sensitive to very low levels of rifampicin (Fig.3, b and c). [ In the experiments shown in Fig.3, a relatively low

concentration of the extracts (one fifth amount of the protein employed for standard assay) was used to alleviate premature termination of transcription.] Evidence that pN and RNA polymerase interact in vitro was obtained in the experiments represented in Fig.4, a and b. Synthesis of trp mRNA from  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> DNA in an extract of A19trpAE1 rif<sup>r</sup> was progressively stimulated by increasing amounts of the pN-containing extract from the rif<sup>+</sup> strain carrying pYS5. The amounts of trp mRNA synthesized in the presence of rifampicin at all concentrations of pN tested (up to a maximum stimulation at a ratio of  $\frac{1}{4}$  of rif<sup>r</sup> to rif<sup>+</sup> RNA polymerase) was about 70 % that produced in the absence of rifampicin (compare Fig.4, a and b). These results suggest that newly formed pN-rif<sup>r</sup> RNA polymerase complexes participate efficiently in the transcription of the N-trp operon of  $\lambda$ trp DNA. We favor the possibility that pN forms its initial association with RNA polymerase (holoenzyme) in a post-initiation event, perhaps at the nutL site, to stimulate trpCBA mRNA synthesis.

When the DNA from  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> phage carrying the nutL44 mutation was used as template, the stimulation of trp mRNA synthesis by the addition of the pN-containing extract was barely detectable, both in the presence or absence of rifampicin (Fig.4, d and c). As a control to assess the overall capacity of the assay system to overcome transcription termination barriers on the template DNA, we examined the pN-dependence of transcription directed by the p<sub>R</sub> promoter and normally controlled in vivo by the nutR site. The transcripts arising from the p<sub>R</sub> promoter can be specifically assayed by using the r-strand of  $\lambda$ cI90 DNA as a probe in DNA-RNA hybridization. Under the conditions employing  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> nutL44 DNA as a template, the addition of the pN-containing extract stimulated transcription originating at the p<sub>R</sub> promoter. Stimulation was observed both in the presence and absence of

rifampicin, in contrast to the failure to stimulate transcription initiating at the  $p_L$  promoter (Fig.4, d and c). The stimulation of  $p_R$ -directed transcription was roughly linear, depending on the amount of pN-containing extract added, and was as efficient as that seen with  $\lambda_{trp48\Delta p_{2trp}}$  DNA (cf. Fig.4, a and b).

These results support a previous conclusion by Salstrom and Szybalski (1978) that the  $nutL^-$  mutation specifically prevents the action of pN on transcription from the  $p_L$  promoter. Furthermore, since the cell-free system used lacks ribosomes and other essential components for translation,  $N$ -mediated antitermination does not require any contemporaneous translation of the mRNA. The *in vitro* transcription assay system is anticipated to be useful to determine the details of the interaction of pN and RNA polymerase.

#### ACKNOWLEDGMENTS

We are deeply indebted to Dr. W. Szybalski who was very generous in critical review of the manuscript prior to its submission and providing manuscripts relating to their most recent work on the role and mode of action of pN. We are also grateful to Dr. D. Schlessinger for critical reading of the manuscript.

Legends to Figures and Tables

Table 1. Bacteria, bacteriophages and plasmids employed in this study.

Figure 1. Simplified molecular maps of  $\lambda$ trp phages.

The genetic maps of transducing phages are based on the data reported Fian dt et al., 1974; previously (Nishimune, 1973; Franklin, 1974; Jackson and Yanofsky, 1972). The map of phage  $\lambda$ trp34 is based on the data of heteroduplex mapping obtained by Fian dt and Szybalski (personal communication).  $t_N^-$  indicates pN-sensitive transcription termination sites within the trp operon, observed in the absence of pN function in  $\lambda$ trp. Interpretation of ( $t_N^-$ ) is in the text. The dotted lines indicate deleted portions of the genome.

Table 2. *In vitro* synthesis of  $p_L$ -directed mRNA from  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> DNA in the presence or absence of functional pN<sup>λ</sup>.

The reaction mixture containing 5  $\mu$ g of  $\lambda$ trp48 $\Delta$ p<sub>2trp</sub> phage DNA and 500  $\mu$ g of nucleic acid-free S-100 protein was incubated for 40 min at the temperature indicated. The other conditions are as described in Materials and Methods.

Table 3. pN-dependent stimulation of trp mRNA synthesis on various  $\lambda$ trp DNA templates.

150  $\mu$ g of nucleic acid-free S-100 protein from A19trpAE1(pYS5) or A19trpAE1 were added to the reaction mixtures containing 670  $\mu$ g of nucleic acid-free S-100 protein from A19trpAE1 and 5  $\mu$ g of template DNA from various phages. The levels of trpD mRNA and trpCBA mRNA synthesized in the reaction mixture containing 670  $\mu$ g of S-100 protein from A19trpAE1 was 143 cpm and 89 cpm, respectively. The other conditions are as described in Materials and Methods.

Figure 2. Sedimentation profiles of *trpD* mRNA synthesized by purified RNA polymerase in the presence or absence of *rho* factor.

The reaction mixture containing 5  $\mu\text{g}$  of  $\lambda_{\text{trp48}\Delta\text{p}_{2\text{trp}}}$  phage DNA and 3  $\mu\text{g}$  of *E. coli* RNA polymerase was incubated for 30 min at 37° C in the presence (a) or absence (b) of 7  $\mu\text{g}$  of *rho* factor. The other conditions are as described in Materials and Methods.

Figure 3. Effect of rifampicin on *trp* mRNA synthesis catalyzed by S-100 extracts of  $\text{Al9trpAElrif}^{\text{r}}$ ,  $\text{Al9trpAElrif}^{\text{+}}$  and  $\text{Al9trpAElrif}^{\text{+}}(\text{pYS5})$ .

The reaction mixture contained 130  $\mu\text{g}$  of nucleic acid-free S-100 protein from  $\text{Al9trpAElrif}^{\text{r}}$  (a),  $\text{Al9trpAElrif}^{\text{+}}$  (b) or  $\text{Al9trpAElrif}^{\text{+}}(\text{pYS5})$  (c). Rifampicin was present at the concentrations indicated.  $\text{O--O}$ , *trpD* mRNA;  $\bullet\text{--}\bullet$ , *trpCBA* mRNA.

Figure 4. pN-dependent synthesis of *trp* mRNA and  $\lambda$  mRNA from the  $\lambda_{\text{trp}}$  DNA template carrying the  $\text{nutL}^{\text{+}}$  or  $\text{nutL44}$  alleles.

The conditions employed were the same as described in Table 3 except that the reaction mixture contained 670  $\mu\text{g}$  of nucleic acid-free S-100 protein from  $\text{Al9trpAElrif}^{\text{r}}$  instead of that from  $\text{Al9trpAElrif}^{\text{+}}$ . (a)  $\lambda_{\text{trp48}\Delta\text{p}_{2\text{trp}}\text{nutL}^{\text{+}}}$  DNA template, in the absence of rifampicin; (b)  $\lambda_{\text{trp48}\Delta\text{p}_{2\text{trp}}\text{nutL}^{\text{+}}}$  DNA template, in the presence of rifampicin (100  $\mu\text{g}/\text{ml}$ ); (c)  $\lambda_{\text{trp48}\Delta\text{p}_{2\text{trp}}\text{nutL44}}$  DNA template, in the absence of rifampicin; (d)  $\lambda_{\text{trp48}\Delta\text{p}_{2\text{trp}}\text{nutL44}}$  DNA template, in the presence of rifampicin (100  $\mu\text{g}/\text{ml}$ ).  $\bullet\text{--}\bullet$ , *trpCBA* mRNA;  $\text{O--O}$ ,  $\text{p}_{\text{R}}$ -promoted  $\lambda$  mRNA. The other conditions are as described in Materials and Methods.

## REFERENCE

- Baumann, M.F. and Friedman, D.I., Cooperative effects of bacterial mutations affecting  $\lambda$  N gene expression II Isolation and characterization of mutations in the rif region, *Virology*, 73(1976) 128-138
- Bøvre, K., Lozeron, H.A. and Szybalski, W., Techniques of RNA-DNA hybridization in solution for the study of viral transcription, in K. Maramorosch and H. Koprowski (Eds.), *Methods in virology*, Vol. 5, Academic Press, New York, 1971, pp 271-292
- Deeb, S.S., Okamoto, K. and Hall, B., Isolation and characterization of non-defective transducing elements of bacteriophage  $\phi 80$ , *Virology*, 31(1967) 289-295
- Dottin, R.P. and Pearson, M.L., Regulation by N gene protein of phage lambda of anthranilate synthetase synthesis in vitro, *Proc. Natl. Acad. Sci., U.S.A.*, 70 (1973) 1078-1082
- Epp, C. and Pearson, M.L., Association of bacteriophage lambda N gene protein with E. coli RNA polymerase, in R. Losick and M. Chamberlin (Eds.), *RNA polymerase*, Cold Spring Harbor Laboratory, New York, 1976, pp 667-691
- Fianndt, M., Szybalski, W. and Imamoto, F., Physical mapping of trp endpoint in the N-t<sub>L</sub> segments of phage  $\lambda$  trpE-A, *Virology*, 61 (1974) 312-314
- Franklin, N.C., Altered reading of genetic signals fused to the N operon of bacteriophage  $\lambda$ : Genetic evidence for modification of polymerase by the protein product of the N gene, *J. Mol. Biol.*, 89 (1974) 33-48
- Georgopoulos, C.P., Bacterial mutants in which the gene N function of bacteriophage lambda is blocked have an altered RNA polymerase, *Proc. Natl. Acad. Sci. U.S.A.*, 68 (1971) 2977-2981
- Gesteland, R.F., Isolation and characterization of ribonuclease I mutants of Escherichia coli, *J. Mol. Biol.*, 16 (1966) 67-84
- Ghysen, A. and Pironio, M., Relationship between the N function of bacteriophage  $\lambda$  and host RNA polymerase, *J. Mol. Biol.*, 65 (1972) 259-272

- Greenblatt, J., Positive control of endolysin synthesis in vitro by the gene N protein of phage  $\lambda$ , Proc.Natl.Acad.Sci. U.S.A., 69 (1972) 3606-3610
- Ishii, S. and Imamoto, F., In vitro transcription of the tryptophan operon in isolated bacterial nucleoids, Mol.Gen.Genet., 148 (1976) 295-305
- Ishii, S. and Imamoto, F., Release of nascent trp mRNA from the operon DNA in chloramphenicol-treated cells of Escherichia coli, Mol.Gen.Genet., 161 (1978) 31-37
- Jackson, E.N. and Yanofsky, C., Internal promoter of the tryptophan operon of Escherichia coli is located in a structure gene, J.Mol.Biol., 69 (1972) 307-341
- Nakamura, H. Kano, Y. and Imamoto, F., Restoration of polarity by N-deficiency in lambda phage containing a translocated trp operon segment, Mol.Gen.Genet., 159 (1978, a) 13-20
- Nakamura, H., Yamamoto, T. and Imamoto, F., p<sub>L</sub>-promoted transcription of the promoter-proximal N-trp region is insensistive to chloramphenicol in the absence of N function, Mol.Gen.Genet., 159 (1978, b) 21-26
- Nishimune, Y., Prophage excision: Specific gene amplification of excision-impaired lysogen after induction, Virology, 53 (1973) 236-246
- Oppenheim, A. and Oppenheim, A.B., Suppresible mutations in the cro gene of bacteriophage  $\lambda$ , Virology, 75 (1976) 469-476
- Roberts, J.W., Termination factor for RNA synthesis, Nature (london), 224 (1969) 1168-1174
- Rosenberg, M., Court, D., Brady, C., Salstrom, J.S. and Szybalski, W., The nucleotide sequence for the N-recognition sites of coliphage lambda, Abst. Meet. Bacteriophage and Single-stranded DNA phage, Cold Spring Harbor Laboratory, New York, 1977, pp 36-37
- Salstrom, J.S. and Szybalski, W., Coliphage  $\lambda$  nutL<sup>-</sup> : A unique class of mutants



- defective in the site of gene N product utilization for antitermination of leftward transcription, *J.Mol.Biol.*, 124 (1978) 195-221
- Salstrom, J.S., Fianndt, M. and Szybalski, W., The site controlling the specificity of N action is outside the promoter-operator: A triple hybrid phage  $\lambda$ N21imm434nin5, *Gene*, 5 (1979) in press
- Sato, K. and Matsushiro, A., The tryptophan operon regulated by phage immunity, *J.Mol.Biol.*, 14 (1965) 608-610
- Szybalski, E. and Szybalski, W., Physical mapping of the att-N region of coliphage lambda: Apparent oversaturation of coding capacity in the gam-ral segment, *Biochimie*, 56 (1974) 1497-1503
- Sternberg, N., A class of rif<sup>r</sup> RNA polymerase mutations that interferes with the expression of coliphage  $\lambda$  late genes, *Virology*, 73 (1976) 139-154
- Takeda, Y., Ogata, K. and Matsubara, K., Genetic analysis of the tof gene in the bacteriophage  $\lambda$  genome, *Virology*, 65 (1975) 385-391

Table 1.

Strain	Relevant characteristics	Source or reference
Bacteria E.coli K12		
<u>λ</u> 9 <trpae1< td=""> <td>RNaseI<sup>-</sup><u>Δ</u>trpAE1 rif<sup>+</sup></td> <td>Gesteland(1969)</td> </trpae1<>	RNaseI <sup>-</sup> <u>Δ</u> trpAE1 rif <sup>+</sup>	Gesteland(1969)
<u>λ</u> 9 <trpaerif<sup>r</trpaerif<sup>	RNaseI <sup>-</sup> <u>Δ</u> trpAE1 rif <sup>r</sup>	This study
Bacteriophages		
<u>λ</u> trp48 <u>Δ</u> p <sub>2</sub> trp	<u>trp</u> <u>Δ</u> DC181 <u>Nam</u> 7am53fed <sup>-</sup> <u>cI</u> At2 <u>h</u> <sup>λ</sup>	Nakamura <u>et al.</u> (1978a)
<u>λ</u> trp48 <u>Δ</u> p <sub>2</sub> trp <u>nut</u> L44	<u>trp</u> <u>Δ</u> DC181 <u>Nam</u> 7am53fed <sup>-</sup> <u>cI</u> At2 <u>h</u> <sup>λ</sup> <u>nut</u> L44	Salstrom and Szybalski(1978)
<u>λ</u> trp48	<u>trp</u> (D,C,B,A) <sup>+</sup>	Franklin(1974)
<u>λ</u> trp60-3	<u>trp</u> (E,D,C,B,A) <sup>+</sup>	Nishimune(1973)
<u>λ</u> trp34	<u>trp</u> (C,B,A) <sup>+</sup>	This study
<u>λ</u> croam6 <u>cI</u> 857	<u>N</u> <sup>+</sup>	Oppenheim and Oppenheim(1976)
<u>λ</u> cI857 <u>tof</u> 12	<u>N</u> <sup>+</sup>	Takeda <u>et al.</u> (1975)
<u>λ</u> <u>Nam</u> 7am53 <u>cI</u> 857 <u>tof</u> 12	<u>NN</u> <sup>-</sup>	Takeda <u>et al.</u> (1975)
<u>λ</u> <u>Nts</u> 8	<u>Nts</u>	Tomizawa
<u>λ</u>	<u>N</u> <sup>λ</sup>	
<u>λ</u> <u>imm</u> 21	<u>N</u> <sup>21</sup>	
<u>λ</u> bio256 <u>cI</u> 857	<u>Δ</u> (att-cIII:57.3-72.55% <u>λ</u> )	Szybalski and Szybalski(1974)
<u>λ</u> bio3h- <u>lnin</u> 5	<u>Δ</u> (att-N:57.3-73.5% <u>λ</u> )	Szybalski and Szybalski(1974)
<u>λ</u> cI90		
<u>φ</u> 80 <trped< td=""> <td></td> <td>Sato and Matsushiro(1965)</td> </trped<>		Sato and Matsushiro(1965)
<u>φ</u> 80 <trpcba< td=""> <td></td> <td>Deeb <u>et al.</u>(1967)</td> </trpcba<>		Deeb <u>et al.</u> (1967)
Plasmids		
pYS1	Ap <sup>R*</sup> derivative of colE1	This study
pYS5	<u>λ</u> cIII-N-cI-O segment of <u>λ</u> c1857 <u>croam</u> 6 cloned in pYS1	This study

\* Ap<sup>R</sup>: ampicilin resistant

Table 2.

	Source of pN	Temperature of assay	Presence of functional pN <sup>λ</sup>	mRNA synthesis (cpm)		
				<u>trpCBA</u>	<u>trpD</u>	<u>N</u>
(a)	pYS5	30°C	+	2,342	1,260	1,040
	none	30°C	-	61	150	840
(b)	<u>λcI857tof12</u>	30°C	+	686	482	
	<u>λNam7am53cI857tof12</u>	30°C	-	41	76	
(c)	<u>λNts8</u>	30°C	+	467	318	
	<u>λNts8</u>	40°C	-	30	100	
	λ	30°C	+	494	305	
	λ	40°C	+	450	288	
(d)	λ	30°C	+	452	297	
	<u>λimm21</u>	30°C	-	101	323	

Table 3.

	DNA template	Presence of pN	mRNA synthesis (cpm)	
			<u>trpCBA</u>	<u>trpD</u>
(a)	$\lambda$ <u>trp60-3</u>	+	782	984
		-	78	85
(b)	$\lambda$ <u>trp48</u>	+	991	1,217
		-	107	117
(c)	$\lambda$ <u>trp48</u> $\Delta$ p <sub>2trp</sub>	+	1,104	1,461
		-	48	76
(d)	$\lambda$ <u>trp34</u>	+	289	318
		-	38	71







