



Title	EXISTENCE, NONEXISTENCE AND MULTIPLICITY OF POSITIVE SOLUTIONS FOR PARAMETRIC NONLINEAR ELLIPTIC EQUATIONS
Author(s)	Iannizzotto, Antonio; S. Papageorgiou, Nicolaos
Citation	Osaka Journal of Mathematics. 2014, 51(1), p. 179–202
Version Type	VoR
URL	<a href="https://doi.org/10.18910/29189">https://doi.org/10.18910/29189</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

# EXISTENCE, NONEXISTENCE AND MULTIPLICITY OF POSITIVE SOLUTIONS FOR PARAMETRIC NONLINEAR ELLIPTIC EQUATIONS

ANTONIO IANNIZZOTTO and NIKOLAOS S. PAPAGEORGIOU

(Received January 24, 2012, revised July 10, 2012)

## Abstract

We consider a parametric nonlinear elliptic equation driven by the Dirichlet  $p$ -Laplacian. We study the existence, nonexistence and multiplicity of positive solutions as the parameter  $\lambda$  varies in  $\mathbb{R}_0^+$  and the potential exhibits a  $p$ -superlinear growth, without satisfying the usual in such cases Ambrosetti–Rabinowitz condition. We prove a bifurcation-type result when the reaction has  $(p-1)$ -sublinear terms near zero (problem with concave and convex nonlinearities). We show that a similar bifurcation-type result is also true, if near zero the right hand side is  $(p-1)$ -linear.

## 1. Introduction

Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with a  $C^2$  boundary  $\partial\Omega$  and  $p > 1$  be a real number. In this paper we study the following nonlinear parametric Dirichlet problem:

$$(P_\lambda) \quad \begin{cases} -\Delta_p u = f(z, u, \lambda) & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

The aim of this study is to establish the existence, nonexistence and multiplicity of positive smooth solutions of  $(P_\lambda)$  as the parameter  $\lambda$  varies over  $]0, +\infty[$  and when the reaction term  $f(z, x, \lambda)$  exhibits a  $(p-1)$ -superlinear growth as  $x$  goes to  $+\infty$ . However, we do not employ the usual in such cases Ambrosetti–Rabinowitz condition (AR-condition for short). Instead, we use a weaker condition which permits a much slower growth for  $x \mapsto f(z, x, \lambda)$  near  $+\infty$ . Our setting incorporates, as a very special case, equations involving the combined effects of concave and convex nonlinearities. Such problems were studied by Ambrosetti, Brezis and Cerami [2] (semilinear equations, i.e.  $p = 2$ ) and by Garcia Azorero, Manfredi and Peral Alonso [7] and Guo and Zhang [12] (nonlinear equations, i.e.  $p \neq 2$ ; in Guo and Zhang [12] it is assumed that  $p \geq 2$ ). In all the aforementioned works, the reaction term has the form

$$f(x, \lambda) = \lambda|x|^{q-2}x + |x|^{r-2}x, \quad \text{for all } x \in \mathbb{R}, \lambda > 0, \text{ with } 1 < q < p < r < p^*$$

(recall that  $p^* = Np/(N - p)$  if  $p < N$  and  $p^* = \infty$  if  $p \geq N$ ).

Recently, Hu and Papageorgiou [14] extended these results by considering reactions of the form

$$f(z, x, \lambda) = \lambda|x|^{q-2}x + f_0(z, x), \quad \text{for all } x \in \mathbb{R}, \lambda > 0, \text{ with } 1 < q < p,$$

$f_0: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  being a Carathéodory function (i.e.,  $z \mapsto f_0(z, x)$  is measurable for all  $x \in \mathbb{R}$  and  $x \mapsto f_0(z, x)$  is continuous for a.a.  $z \in \Omega$ ) with subcritical growth in  $x$  and which satisfies the *AR*-condition.

We should mention that there are alternative ways to generalize the *AR*-condition and incorporate more general “superlinear” reactions. For more information in this direction, we refer to the works of Li and Yang [17] and Miyagaki and Souto [19].

Other parametric equations driven by the  $p$ -Laplacian were also considered by Brock, Iturriaga and Ubilla [4], Guo [11], Hu and Papageorgiou [13] and Takeuchi [22]. However, their hypotheses preclude  $(p - 1)$ -superlinear terms.

We will prove the following bifurcation-type result: there exists  $\lambda^* > 0$  s.t. for all  $0 < \lambda < \lambda^*$  problem  $(P_\lambda)$  admits at least two positive smooth solutions; for  $\lambda = \lambda^*$  there is at least one positive smooth solution; and for  $\lambda > \lambda^*$  there is no positive solution. This holds for both problems with  $(p - 1)$ -sublinear reaction near zero (see Theorem 10 below) and problems with  $(p - 1)$ -linear reaction near zero (see Theorem 13 below). Our approach is variational, based on the critical point theory coupled with suitable truncation techniques.

## 2. Mathematical background

In this section we recall some basic notions and analytical tools which we will use in the sequel. So, let  $X$  be a Banach space and  $X^*$  its topological dual. By  $\langle \cdot, \cdot \rangle$  we denote the duality brackets for the pair  $(X^*, X)$ . Let  $\varphi \in C^1(X)$  be a functional. A point  $x_0 \in X$  is called a *critical point* of  $\varphi$  if  $\varphi'(x_0) = 0$ . A number  $c \in \mathbb{R}$  is a *critical value* of  $\varphi$  if there exists a critical point  $x_0 \in X$  of  $\varphi$ , s.t.  $\varphi(x_0) = c$ .

We say that  $\varphi \in C^1(X)$  satisfies the *Cerami condition at level*  $c \in \mathbb{R}$  (the  $C_c$ -condition, for short), if the following holds: *every sequence*  $(x_n) \subset X$ , *s.t.*

$$\varphi(x_n) \rightarrow c \quad \text{and} \quad (1 + \|x_n\|)\varphi'(x_n) \rightarrow 0 \quad \text{in } X^* \quad \text{as } n \rightarrow \infty,$$

*admits a strongly convergent subsequence*. If this is true at every level  $c \in \mathbb{R}$ , then we say that  $\varphi$  satisfies the *Cerami condition* (*C*-condition, for short).

Using this compactness-type condition, we can have the following minimax characterization of certain critical values of a  $C^1$  functional. The result is known as the *mountain pass theorem*.

**Theorem 1.** *If  $X$  is a Banach space,  $\varphi \in C^1(X)$ ,  $x_0, x_1 \in X$ ,  $0 < \rho < \|x_1 - x_0\|$ ,*

$$\max\{\varphi(x_0), \varphi(x_1)\} \leq \inf_{\|x-x_0\|=\rho} \varphi(x) = \eta_\rho,$$

*and  $\varphi$  satisfies the  $C_c$ -condition, where*

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} \varphi(\gamma(t)) \quad \text{and} \quad \Gamma = \{\gamma \in C([0, 1], X) : \gamma(i) = x_i, i = 0, 1\},$$

*then  $c \geq \eta_\rho$  and  $c$  is a critical value of  $\varphi$ . Moreover, if  $c = \eta_\rho$ , then there exists a critical point  $x \in X$  s.t.  $\varphi(x) = c$  and  $\|x - x_0\| = \rho$ .*

In the study of problem  $(P_\lambda)$ , we will use the Sobolev space  $W = W_0^{1,p}(\Omega)$ , endowed with the norm  $\|u\| = \|Du\|_p$ , whose dual is the space  $W^* = W^{-1,p'}(\Omega)$  ( $1/p + 1/p' = 1$ ). We will also use the space

$$C_0^1(\bar{\Omega}) = \{u \in C^1(\bar{\Omega}) : u(z) = 0 \text{ for all } z \in \partial\Omega\}.$$

This is an ordered Banach space with positive cone

$$C_+ = \{u \in C_0^1(\bar{\Omega}) : u(z) \geq 0 \text{ for all } z \in \bar{\Omega}\}.$$

This cone has a nonempty interior, given by

$$\text{int}(C_+) = \left\{ u \in C_+ : u(z) > 0 \text{ for all } z \in \Omega, \frac{\partial u}{\partial n}(z) < 0 \text{ for all } z \in \partial\Omega \right\}.$$

Here  $n(z)$  denotes the outward unit normal to  $\partial\Omega$  at a point  $z$ .

Concerning ordered Banach spaces, in the sequel we will use the following simple fact about them.

**Lemma 2.** *If  $X$  is an ordered Banach space with positive (order) cone  $C$  and  $x_0 \in \text{int}(C)$ , then for every  $y \in X$  we can find  $t > 0$  s.t.  $tx_0 - y \in \text{int}(C)$ .*

A nonlinear map  $A: X \rightarrow X^*$  is of type  $(S)_+$  if, for every sequence  $(x_n) \subset X$  s.t.

$$x_n \rightharpoonup x \quad \text{in} \quad X \quad \text{and} \quad \limsup_n \langle A(x_n), x_n - x \rangle \leq 0,$$

we have  $x_n \rightarrow x$  in  $X$ .

Let  $A: W \rightarrow W^*$  be defined by

$$(1) \quad \langle A(u), v \rangle = \int_{\Omega} |Du|^{p-2} Du \cdot Dv \, dz \quad \text{for all } u, v \in W_0^{1,p}(\Omega).$$

We have the following result (see, for example, Papageorgiou and Kyritsi [20]).

**Proposition 3.** *The map  $A: W \rightarrow W^*$  defined by (1) is continuous, strictly monotone (hence maximal monotone too) and of type  $(S)_+$ .*

Next, let us recall some basic facts about the spectrum of the negative Dirichlet  $p$ -Laplacian. Let  $m \in L^\infty(\Omega)_+$ ,  $m \neq 0$  and consider the following nonlinear weighted eigenvalue problem:

$$(2) \quad \begin{cases} -\Delta_p u = \hat{\lambda} m(z)|u|^{p-2}u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

By an *eigenvalue* of (2) we mean a number  $\hat{\lambda}(m) \in \mathbb{R}$  s.t. problem (2) has a non-trivial solution  $u \in W$ . Nonlinear regularity theory (see, for example, Papageorgiou and Kyritsi [20], pp. 311–312) implies that  $u \in C_0^1(\bar{\Omega})$ . We know that (2) has a smallest eigenvalue  $\hat{\lambda}_1(m) > 0$ , which is simple and isolated. Moreover, the following variational characterization is available:

$$(3) \quad \hat{\lambda}_1(m) = \min_{u \in W \setminus \{0\}} \frac{\|Du\|_p^p}{\int_{\Omega} m(z)|u|^p dz}.$$

The minimum in (3) is attained on the one-dimensional eigenspace of  $\hat{\lambda}_1(m)$ . Note that, if  $m, m' \in L^\infty(\Omega)_+ \setminus \{0\}$ ,  $m \neq m'$  and  $m \leq m'$ , then because of (3) we see that  $\hat{\lambda}_1(m) > \hat{\lambda}_1(m')$ . If  $m = 1$ , we simply write  $\hat{\lambda}_1$  for  $\hat{\lambda}_1(1)$ . Let  $\hat{u}_1 \in C_0^1(\bar{\Omega})$  be the  $L^p$ -normalized eigenfunction corresponding to  $\hat{\lambda}_1$ . It is clear from (3) that  $\hat{u}_1$  does not change sign, and so we may assume  $\hat{u}_1 \in C_+$ . In fact the nonlinear maximum principle of Vázquez [23] implies that  $\hat{u}_1 \in \text{int}(C_+)$ . Every eigenfunction  $u$  corresponding to an eigenvalue  $\hat{\lambda} \neq \hat{\lambda}_1$  is necessarily *nodal* (i.e., sign changing).

Finally, in what follows we denote by  $|\cdot|_N$  the Lebesgue measure on  $\mathbb{R}^N$ . For all  $x \in \mathbb{R}$ , we set

$$x^\pm = \max\{\pm x, 0\}.$$

### 3. Problems with concave and convex nonlinearities

In this section, we consider problems with reactions which are concave (i.e.  $(p-1)$ -sublinear) near zero and convex (i.e.  $(p-1)$ -superlinear) near  $+\infty$ . More precisely, the hypotheses on  $f(z, x, \lambda)$  are the following (by  $p^*$  we denote the Sobolev critical exponent, defined as in Introduction):

**H**  $f: \Omega \times \mathbb{R} \times \mathbb{R}_0^+ \rightarrow \mathbb{R}$  is a Carathéodory function s.t.  $f(z, 0, \lambda) = 0$  for a.a.  $z \in \Omega$  and all  $\lambda > 0$ . We set

$$F(z, x, \lambda) = \int_0^x f(z, s, \lambda) ds \quad \text{for a.a. } z \in \Omega \quad \text{and all } x \in \mathbb{R}, \lambda > 0$$

and assume:

(i)  $f(z, x, \lambda) \leq a(z, \lambda) + c|x|^{r-1}$  for a.a.  $z \in \Omega$  and all  $x \in \mathbb{R}$ ,  $\lambda > 0$ , with  $p < r < p^*$  and  $a(\cdot, \lambda) \in L^\infty(\Omega)_+$  s.t. the function  $\lambda \mapsto \|a(\cdot, \lambda)\|_\infty$  is bounded on bounded sets and goes to 0 as  $\lambda \rightarrow 0^+$ ,  $c > 0$ ;

(ii) for all  $\lambda > 0$

$$\lim_{x \rightarrow +\infty} \frac{F(z, x, \lambda)}{x^p} = +\infty \quad \text{uniformly for a.a. } z \in \Omega,$$

and there exist  $\tau \in ](r-p)\max\{1, N/p\}, p^*[$  and, for all bounded  $I \subset \mathbb{R}_0^+$ , a real number  $\beta_0 > 0$  s.t.

$$(4) \quad \liminf_{x \rightarrow +\infty} \frac{f(z, x, \lambda)x - pF(z, x, \lambda)}{x^\tau} \geq \beta_0 \quad \text{for all } \lambda \in I;$$

(iii) there exist  $\delta_0 > 0$ ,  $\mu \in ]1, p[$  and  $\eta_0 > 0$  s.t.

$$f(z, x, \lambda) \geq \eta_0 x^{\mu-1} \quad \text{for a.a. } z \in \Omega \quad \text{and all } x \in [0, \delta_0], \lambda > 0;$$

(iv) for a.a.  $z \in \Omega$  and all  $x \geq 0$  the function  $\lambda \mapsto f(z, x, \lambda)$  is increasing, for all  $\lambda > \lambda' > 0$ ,  $s > 0$  there exists  $\mu_s > 0$  s.t.

$$f(z, x, \lambda) - f(z, x, \lambda') \geq \mu_s \quad \text{for a.a. } z \in \Omega \quad \text{and all } x \geq s$$

and for all compact  $K \subset \mathbb{R}_0^+$

$$\lim_{\lambda \rightarrow +\infty} f(z, x, \lambda) = +\infty \quad \text{uniformly for a.a. } z \in \Omega \quad \text{and all } x \in K;$$

(v) for all  $\xi > 0$  and every bounded interval  $I \subset \mathbb{R}_0^+$ , we can find  $\sigma_\xi^I > 0$  s.t. the function  $x \mapsto f(z, x, \lambda) + \sigma_\xi^I x^{p-1}$  is nondecreasing on  $[0, \xi]$  for a.a.  $z \in \Omega$  and all  $\lambda \in I$ .

**REMARK 4.** Since we are interested in positive solutions and hypotheses **H** (ii)–(v) concern only the positive semiaxis  $\mathbb{R}^+$ , by truncating things if necessary, we may (and will) assume that  $f(z, x, \lambda) = 0$  for a.a.  $z \in \Omega$  and all  $x \leq 0$ ,  $\lambda > 0$ . Hypothesis **H** (i) imposes a growth condition only from above, since from below the other hypotheses imply that for every  $\lambda > 0$  we can find  $\xi^* > 0$  s.t.  $f(z, x, \lambda) \geq -\xi^*$  for a.a.  $z \in \Omega$ , all  $x \geq 0$ . Indeed, from **H** (ii) we see that for  $x > 0$  large, say for  $x \geq M > 0$ , we have  $f(z, x, \lambda) \geq 0$  for a.a.  $z \in \Omega$ . Similarly, hypothesis **H** (iii) implies that  $f(z, x, \lambda) \geq 0$  for a.a.  $z \in \Omega$ , all  $x \in [0, \delta_0]$ . Finally, for  $x \in [\delta_0, M]$  we use **H** (v) and obtain the required bound from below. Hypothesis **H** (ii) classifies problem  $(P_\lambda)$  as *p-superlinear*, since it implies that near  $\infty$  the potential function  $x \mapsto F(z, x, \lambda)$  grows faster than  $x^p$ . Evidently, this is the case if  $x \mapsto f(z, x, \lambda)$  is  $(p-1)$ -superlinear near  $+\infty$ , i.e.

$$\lim_{x \rightarrow +\infty} \frac{f(z, x, \lambda)}{x^{p-1}} = +\infty \quad \text{uniformly for a.a. } z \in \Omega \quad \text{and all } \lambda > 0.$$

In the literature, such problems are usually studied using the *AR-condition*. We recall that  $f$  satisfies the (unilateral) *AR-condition* uniformly in  $\lambda > 0$ , if there exist  $M > 0$ ,  $\tau > p$  s.t.

$$(5) \quad 0 < \tau F(z, x, \lambda) \leq f(z, x, \lambda)x \quad \text{for a.a. } z \in \Omega \quad \text{and all } x \geq M, \lambda > 0.$$

Integrating (5), we obtain the weaker condition

$$(6) \quad c_1 x^\tau \leq F(z, x, \lambda) \quad \text{for a.a. } z \in \Omega \quad \text{and all } x \geq M, \lambda > 0 \quad (c_1 > 0).$$

Clearly (6) implies the much weaker condition

$$(7) \quad \lim_{x \rightarrow +\infty} \frac{F(z, x, \lambda)}{x^p} = +\infty \quad \text{uniformly for a.a. } z \in \Omega \quad \text{and all } \lambda > 0.$$

Here, instead of the *AR-condition* (5), we employ the more general conditions (7) and (4). Similar assumptions can be found in Costa and Magalhães [5] and Fei [6]. Other ways to relax the *AR-condition* in the study of  $p$ -superlinear problems can be found in the papers of Jeanjean [15], Miyagaki and Souto [19] and Schechter and Zou [21]. Finally, note that hypothesis **H** (iii) implies that  $x \mapsto F(z, x, \lambda)$  is  $p$ -sublinear near zero. Therefore hypotheses **H** correspond to problems with *concave and convex nonlinearities*.

**EXAMPLE 5.** The following functions  $f_i: \mathbb{R}^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}$  ( $i = 1, 2, 3$ ) satisfy hypotheses **H**:

$$\begin{aligned} f_1(x, \lambda) &= \lambda x^{q-1} + x^{r-1} \quad (1 < q < p < r < p^*), \\ f_2(x, \lambda) &= \lambda x^{q-1} + x^{p-1} \left( \ln(1+x) + \frac{1}{p} \frac{x}{1+x} \right) \quad (1 < q < p), \\ f_3(x, \lambda) &= \begin{cases} \lambda x^{q-1} & \text{if } 0 \leq x \leq 1, \\ p \lambda x^{p-1} \left( \ln(x) + \frac{1}{p} \right) & \text{if } x > 1, \end{cases} \quad (1 < q < p). \end{aligned}$$

Of course, we set  $f_i(x, \lambda) = 0$  for all  $x \leq 0$ ,  $\lambda > 0$  and for  $i = 1, 2, 3$ . Note that  $f_1(x, \lambda)$  is the reaction term used by Ambrosetti, Brezis and Cerami [2] (for  $p = 2$ ), by Garcia Azorero, Manfredi and Peral Alonso [7] (for  $p > 1$ ) and by Guo and Zhang [12] (for  $p \geq 2$ ). Functions  $f_2(x, \lambda)$  and  $f_3(x, \lambda)$  do not satisfy the *AR-condition*. So, our work generalizes significantly those in [7] and [12].

For all  $\lambda > 0$  and  $u \in W$ , we denote

$$(8) \quad N_f^\lambda(u)(z) = f(z, u(z), \lambda) \quad \text{for a.a. } z \in \Omega.$$

By a (*weak*) *solution* of  $(P_\lambda)$  we mean a function  $u \in W$  s.t.

$$A(u) = N_f^\lambda(u) \quad \text{in } W^*,$$

that is,

$$\int_{\Omega} |Du|^{p-2} Du \cdot Dv \, dz = \int_{\Omega} f(z, u, \lambda) v \, dz \quad \text{for all } v \in W.$$

We say that  $u$  is *positive* if  $u(z) > 0$  for a.a.  $z \in \Omega$ . Set

$$\mathcal{P} = \{\lambda \in \mathbb{R}_0^+ : (P_{\lambda}) \text{ has a positive solution}\}.$$

The following Propositions illustrate the properties of the set  $\mathcal{P}$ .

**Proposition 6.** *If hypotheses **H** hold, then  $\mathcal{P} \neq \emptyset$  and for all  $\lambda \in \mathcal{P}$ ,  $\mu \in ]0, \lambda[$  we have  $\mu \in \mathcal{P}$ .*

Proof. Let  $e \in W \setminus \{0\}$ ,  $e \geq 0$  be the unique solution of the following auxiliary Dirichlet problem:

$$(9) \quad \begin{cases} -\Delta_p e = 1 & \text{in } \Omega, \\ e = 0 & \text{on } \partial\Omega. \end{cases}$$

Nonlinear regularity theory (see [20]) and the nonlinear maximum principle (see Vázquez [23]) imply that  $e \in \text{int}(C_+)$ .

**Claim.** *There exists  $\tilde{\lambda} > 0$  s.t., for all  $\lambda \in ]0, \tilde{\lambda}[$ , we can find  $\tilde{\xi} > 0$  s.t.*

$$(10) \quad \|a(\cdot, \lambda)\|_{\infty} + c(\tilde{\xi} \|e\|_{\infty})^{r-1} < \tilde{\xi}^{p-1} \quad (c > 0 \text{ as in } \mathbf{H} \text{ (i)}).$$

We argue by contradiction. So, suppose we can find a sequence  $(\lambda_n) \subset \mathbb{R}_0^+$  s.t.  $\lambda_n \rightarrow 0$  and

$$\tilde{\xi}^{p-1} \leq \|a(\cdot, \lambda_n)\|_{\infty} + c(\tilde{\xi} \|e\|_{\infty})^{r-1} \quad \text{for all } n \in \mathbb{N}, \tilde{\xi} > 0.$$

Passing to the limit as  $n \rightarrow \infty$  and using hypothesis **H** (i), we obtain

$$1 \leq c\tilde{\xi}^{r-p} \|e\|_{\infty}^{r-1} \quad \text{for all } \tilde{\xi} > 0.$$

Since  $r > p$ , letting  $\tilde{\xi} \rightarrow 0^+$  we reach a contradiction. This proves the claim.

Now, we fix  $\lambda \in ]0, \tilde{\lambda}[$ . Set  $\tilde{u} = \tilde{\xi} e \in \text{int}(C_+)$ . We have

$$A(\tilde{u}) = \tilde{\xi}^{p-1} \quad (\text{see (9)}),$$

which implies

$$(11) \quad A(\tilde{u}) \geq N_f^{\lambda}(\tilde{u}) \quad \text{in } W^* \quad (\text{see (10) and } \mathbf{H} \text{ (i)}),$$

therefore  $\tilde{u}$  is an upper solution for problem  $(P_\lambda)$ . We consider the following truncation of  $f(z, x, \lambda)$ :

$$(12) \quad \tilde{f}(z, x, \lambda) = \begin{cases} f(z, x, \lambda) & \text{if } x < \tilde{u}(z), \\ f(z, \tilde{u}(z), \lambda) & \text{if } x \geq \tilde{u}(z), \end{cases} \quad \text{for a.a. } z \in \Omega \text{ and all } x \in \mathbb{R}, \lambda \in ]0, \tilde{\lambda}[.$$

Evidently,  $(z, x) \mapsto \tilde{f}(z, x, \lambda)$  is a Carathéodory function. We set

$$\tilde{F}(z, x, \lambda) = \int_0^x \tilde{f}(z, s, \lambda) ds$$

and consider the functional  $\tilde{\varphi}_\lambda: W \rightarrow \mathbb{R}$  defined by

$$\tilde{\varphi}_\lambda(u) = \frac{1}{p} \|Du\|_p^p - \int_{\Omega} \tilde{F}(z, u, \lambda) dz \quad \text{for all } u \in W.$$

It is clear from (12) that  $\tilde{\varphi}_\lambda \in C^1(W)$  is coercive. Also, exploiting the compact embedding of  $W$  into  $L^r(\Omega)$  (by the Sobolev embedding theorem), we can easily check that  $\tilde{\varphi}_\lambda$  is sequentially weakly l.s.c. Thus, by the Weierstrass theorem, we can find  $u_0 \in W$  s.t.

$$(13) \quad \tilde{\varphi}_\lambda(u_0) = \inf_{u \in W} \tilde{\varphi}_\lambda(u) = \tilde{m}_\lambda.$$

Let  $\delta_0 > 0$  be as postulated in hypothesis **H** (iii) and let  $t \in ]0, 1[$  be s.t.

$$0 \leq t\hat{u}_1(z) \leq \min\{\tilde{u}(z), \delta_0\} \quad \text{for all } z \in \overline{\Omega}$$

(recall that  $\tilde{u}, \hat{u}_1 \in \text{int}(C_+)$  and use Lemma 2). Then, by virtue of hypothesis **H** (iii), we have

$$(14) \quad F(z, t\hat{u}_1(z), \lambda) \geq \frac{\eta_0}{\mu} (t\hat{u}_1(z))^\mu \quad \text{for a.a. } z \in \Omega.$$

So, we get

$$\begin{aligned} \tilde{\varphi}_\lambda(t\hat{u}_1) &= \frac{t^p}{p} \|D\hat{u}_1\|_p^p - \int_{\Omega} F(z, t\hat{u}_1, \lambda) dz \quad (\text{see (12) and (14)}) \\ &\leq t^\mu \left[ \frac{t^{p-\mu}}{p} \hat{\lambda}_1 - \frac{\eta_0}{\mu} \|\hat{u}_1\|_\mu^\mu \right] \quad (\text{see (3), (14) and recall } \|\hat{u}_1\|_p = 1). \end{aligned}$$

Since  $\mu < p$  (see **H** (iii)), choosing  $t \in ]0, 1[$  even smaller if necessary, from the inequality above we infer that

$$\tilde{\varphi}_\lambda(t\hat{u}_1) < 0,$$

which in turn implies

$$\tilde{m}_\lambda < 0 = \tilde{\varphi}_\lambda(0).$$

So, by (13)  $u_0 \neq 0$ .

From (13) we deduce that  $u_0$  is a critical point of  $\tilde{\varphi}_\lambda$ , that is,

$$(15) \quad A(u_0) = N_{\tilde{f}}^\lambda(u_0) \quad \text{in } W^* \quad (N_{\tilde{f}}^\lambda \text{ defined as in (8), with } \tilde{f} \text{ instead of } f).$$

On (15) we act with  $u_0^- \in W$  and we obtain

$$\|Du_0^-\|_p = 0 \quad (\text{see (12)}),$$

i.e.  $u_0 \geq 0$  a.e. in  $\Omega$ .

Also, on (15) we act with  $(u_0 - \tilde{u})^+ \in W$ . Then,

$$\begin{aligned} \langle A(u_0), (u_0 - \tilde{u})^+ \rangle &= \int_{\Omega} \tilde{f}(z, u_0, \lambda)(u_0 - \tilde{u})^+ dz \\ &= \int_{\Omega} f(z, \tilde{u}, \lambda)(u_0 - \tilde{u})^+ dz \quad (\text{see (12)}) \\ &\leq \langle A(\tilde{u}), (u_0 - \tilde{u})^+ \rangle \quad (\text{see (11)}), \end{aligned}$$

that is,

$$\begin{aligned} \langle A(u_0) - A(\tilde{u}), (u_0 - \tilde{u})^+ \rangle &= \int_{\{u_0 > \tilde{u}\}} (|Du_0|^{p-2}Du_0 - |D\tilde{u}|^{p-2}D\tilde{u}) \cdot (Du_0 - D\tilde{u}) dz \\ &\leq 0. \end{aligned}$$

So we have

$$|\{u_0 > \tilde{u}\}|_N = 0,$$

i.e.  $u_0 \leq \tilde{u}$ . So (15) becomes

$$A(u_0) = N_{\tilde{f}}^\lambda(u_0) \quad \text{in } W^*.$$

We have proved that  $u_0 \in W \setminus \{0\}$ ,  $0 \leq u_0 \leq \tilde{u}$  and  $u_0$  solves problem  $(P_\lambda)$ . As before, nonlinear regularity theory (see [20]) assures that  $u_0 \in C_+ \setminus \{0\}$ . Set  $\xi = \|u_0\|_\infty$ ,  $I = ]0, \tilde{\lambda}[$  and find  $\tilde{\sigma} = \sigma_\xi^I$  as in hypothesis **H** (v). We have

$$-\Delta_p u_0(z) + \tilde{\sigma} u_0(z)^{p-1} = f(z, u_0(z), \lambda) + \tilde{\sigma} u_0(z)^{p-1} \geq 0 \quad \text{for a.a. } z \in \Omega,$$

so

$$\Delta_p u_0(z) \leq \tilde{\sigma} u_0(z)^{p-1} \quad \text{for a.a. } z \in \Omega,$$

hence  $u_0 \in \text{int}(C_+)$  (see [23]). Thus,  $u_0$  is a smooth positive solution of  $(P_\lambda)$ , in particular  $\lambda \in \mathcal{P}$ . Therefore  $]0, \tilde{\lambda}[ \subseteq \mathcal{P}$ , in particular  $\mathcal{P} \neq \emptyset$ .

Next, let  $\lambda \in \mathcal{P}$  and  $0 < \mu < \lambda$ . We can find a positive solution  $u_\lambda \in \text{int}(C_+)$  for problem  $(P_\lambda)$ . By hypothesis **H** (iv) we have

$$(16) \quad A(u_\lambda) = N_f^\lambda(u_\lambda) \geq N_f^\mu(u_\lambda) \quad \text{in } W^*,$$

therefore  $u_\lambda$  is an upper solution for problem  $(P_\mu)$ . We truncate  $x \mapsto f(z, x, \lambda)$  at  $u_\lambda(z)$  and we argue as above. Via the direct method (using this time (16) instead of (11)), we produce a positive solution  $u_\mu \in \text{int}(C_+)$  for problem  $(P_\mu)$ , s.t.  $0 \leq u_\mu \leq u_\lambda$  in  $\overline{\Omega}$ . Therefore,  $\mu \in \mathcal{P}$ .  $\square$

Denote

$$\lambda^* = \sup \mathcal{P}.$$

**Proposition 7.** *If hypotheses **H** hold, then  $\lambda^* < +\infty$ .*

Proof. Hypotheses **H** (ii), (iii) and (iv) imply that we can find  $\bar{\lambda} > 0$  large s.t.

$$(17) \quad f(z, x, \bar{\lambda}) \geq \hat{\lambda}_1 x^{p-1} \quad \text{for a.a. } z \in \Omega, \quad \text{all } x \geq 0.$$

To see (17) note that by choosing  $\delta_0 > 0$  even smaller if necessary, from **H** (iii) we have

$$f(z, x, \lambda) \geq \hat{\lambda}_1 x^{p-1} \quad \text{for a.a. } z \in \Omega, \quad \text{all } x \in [0, \delta_0].$$

Also, from hypothesis **H** (ii) we see that we can find  $M > 0$  large enough s.t.

$$f(z, x, \lambda) \geq \hat{\lambda}_1 x^{p-1} \quad \text{for a.a. } z \in \Omega, \quad \text{all } x \geq M.$$

Finally, invoking **H** (v), we infer that for all  $\lambda > 0$  big, we have

$$f(z, x, \lambda) \geq \hat{\lambda}_1 M^{p-1} \geq \hat{\lambda}_1 x^{p-1} \quad \text{for a.a. } z \in \Omega, \quad \text{all } x \in [\delta_0, M].$$

From these estimates we have (17) for  $\lambda > 0$  big.

We will prove that  $\lambda^* \leq \bar{\lambda}$ , arguing by contradiction. So, let  $\lambda > \bar{\lambda}$  and suppose that problem  $(P_\lambda)$  has a nontrivial positive solution  $u_\lambda \in W$ . As before, we obtain  $u_\lambda \in \text{int}(C_+)$ . By virtue of Lemma 2, we can find  $t > 0$  s.t.

$$t \hat{u}_1(z) \leq u_\lambda(z) \quad \text{for all } z \in \overline{\Omega}.$$

Let  $t > 0$  be the largest such positive real number. Let  $\xi = \|u_\lambda\|_\infty$ ,  $I = [\bar{\lambda}, \lambda]$  and

choose  $\bar{\sigma} = \sigma_\xi^I$  as in hypothesis **H** (v). We have

$$\begin{aligned}
& -\Delta_p u_\lambda + \bar{\sigma} u_\lambda^{p-1} \\
&= f(z, u_\lambda, \lambda) + \bar{\sigma} u_\lambda^{p-1} \\
&= f(z, u_\lambda, \bar{\lambda}) + \bar{\sigma} u_\lambda^{p-1} + \theta^*(z) \quad (\text{we set } \theta^*(z) = f(z, u_\lambda, \lambda) - f(z, u_\lambda, \bar{\lambda})) \\
&\geq \hat{\lambda}_1 u_\lambda^{p-1} + \bar{\sigma} u_\lambda^{p-1} + \theta^*(z) \quad (\text{see (17)}) \\
&\geq \hat{\lambda}_1 (t\hat{u}_1)^{p-1} + \bar{\sigma} (t\hat{u}_1)^{p-1} + \theta^*(z) \quad (\text{recall } t\hat{u}_1 \leq u_\lambda) \\
&= -\Delta_p (t\hat{u}_1) + \bar{\sigma} (t\hat{u}_1)^{p-1} + \theta^*(z).
\end{aligned}$$

Since  $u_\lambda \in \text{int}(C_+)$ , using hypothesis **H** (iv), we see that for every compact  $K \subset \Omega$  we can find  $\mu_K > 0$  s.t.

$$\theta^*(z) \geq \mu_K \quad \text{for a.a. } z \in K.$$

Then, from Proposition 2.6 of Arcaya and Ruiz [3], we infer that  $u_\lambda - t\hat{u}_1 \in \text{int}(C_+)$ , which contradicts the maximality of  $t > 0$ .

This proves that for  $\lambda > \bar{\lambda}$  problem  $(P_\lambda)$  has no nontrivial positive solution in  $W$  and so  $\lambda^* \leq \bar{\lambda}$ , in particular  $\lambda^* < +\infty$ .  $\square$

**Proposition 8.** *If hypotheses **H** hold, then  $\lambda^* \in \mathcal{P}$  and so  $\mathcal{P} = ]0, \lambda^*]$ .*

Proof. Let  $(\lambda_n) \subset ]0, \lambda^*] \subseteq \mathcal{P}$  be an increasing sequence s.t.  $\lambda_n \rightarrow \lambda^*$ . To each  $\lambda_n$  there corresponds a positive smooth solution  $u_n = u_{\lambda_n} \in \text{int}(C_+)$  for problem  $(P_{\lambda_n})$ . For all  $m > n \geq 1$  we have

$$(18) \quad A(u_m) = N_f^{\lambda_m}(u_m) \geq N_f^{\lambda_n}(u_m) \quad \text{in } W^* \quad (\text{see hypothesis } \mathbf{H} \text{ (iv)}).$$

Truncating  $x \mapsto f(z, x, \lambda_n)$  at  $u_m(z)$  and reasoning as in the proof of Proposition 6, using the direct method and (18) we obtain a smooth positive solution for  $(P_{\lambda_n})$  with values in  $[0, u_m(z)]$ , with negative energy. So, without any loss of generality, we may assume that

$$(19) \quad \varphi_{\lambda_n}(u_n) < 0 \quad \text{for all } n \in \mathbb{N},$$

with

$$\varphi_\lambda(u) = \frac{1}{p} \|Du\|_p^p - \int_\Omega F(z, u, \lambda) dz \quad \text{for all } \lambda > 0, u \in W.$$

Also, we have

$$(20) \quad A(u_n) = N_f^{\lambda_n}(u_n) \quad \text{for all } n \in \mathbb{N}.$$

From (19) we have

$$(21) \quad \|Du_n\|_p^p - \int_{\Omega} pF(z, u_n, \lambda_n) dz < 0 \quad \text{for all } n \in \mathbb{N}.$$

Acting on (20) with  $u_n \in W$ , we obtain

$$(22) \quad \|Du_n\|_p^p - \int_{\Omega} f(z, u_n, \lambda_n) u_n dz = 0 \quad \text{for all } n \in \mathbb{N}.$$

Subtracting (22) from (21), we get

$$(23) \quad \int_{\Omega} [f(z, u_n, \lambda_n) u_n - pF(z, u_n, \lambda_n)] dz < 0 \quad \text{for all } n \in \mathbb{N}.$$

Hypotheses **H** (i), (ii) imply that we can find  $\beta_1 \in ]0, \beta_0[$  and  $c_2 > 0$  s.t.

$$(24) \quad \beta_1 x^\tau - c_2 \leq f(z, x, \lambda) x - pF(z, x, \lambda) \quad \text{for a.a. } z \in \Omega \text{ and all } x \geq 0, \lambda \in ]0, \lambda^*].$$

Using (24) in (23), we see that

$$(25) \quad (u_n) \text{ is bounded in } L^\tau(\Omega).$$

**Claim.** *There exists  $u^* \in W$  s.t., up to a subsequence,*

$$(26) \quad u_n \rightharpoonup u^* \text{ in } W \quad \text{and} \quad u_n \rightarrow u^* \text{ in } L^r(\Omega) \quad \text{as } n \rightarrow \infty.$$

First, suppose that  $N \neq p$ . From hypothesis **H** (ii) it is clear that we can always assume  $\tau \leq r < p^*$ . So, we can find  $t \in [0, 1[$  s.t.

$$\frac{1}{r} = \frac{1-t}{\tau} + \frac{t}{p^*} \quad (\text{recall that } p^* = +\infty \text{ if } N < p).$$

From the interpolation inequality (see, for example, Gasiński and Papageorgiou [8], p. 905) we have

$$\|u_n\|_r \leq \|u_n\|_\tau^{1-t} \|u_n\|_{p^*}^t \quad \text{for all } n \in \mathbb{N},$$

which (together with (25) and the Sobolev embedding theorem) implies

$$(27) \quad \|u_n\|_r^r \leq c_3 \|Du_n\|_p^{tr} \quad \text{for all } n \in \mathbb{N} \quad (c_3 > 0).$$

From hypothesis **H** (i) we have

$$(28) \quad f(z, u_n(z), \lambda_n) u_n(z) \leq c_4 (1 + |u_n(z)|^r) \quad \text{for a.a. } z \in \Omega \text{ and all } n \in \mathbb{N} \quad (c_4 > 0).$$

From (20), we have for all  $n \in \mathbb{N}$  and some  $c_5, c_6 > 0$

$$\begin{aligned} \|Du_n\|_p^p &= \int_{\Omega} f(z, u_n, \lambda_n) u_n \, dz \\ &\leq c_5(1 + \|u_n\|_r^r) \quad (\text{see (28)}) \\ &\leq c_6(1 + \|Du_n\|_p^{tr}) \quad (\text{see (27)}). \end{aligned}$$

The restriction on  $\tau$  in hypothesis **H** (ii) implies that  $tr < p$ . So, from the inequality above we infer that  $(u_n)$  is bounded in  $W$  and we can find  $u^* \in W$  satisfying (26).

If  $N = p$ , then by the Sobolev theorem  $W$  is (compactly) embedded in  $L^\eta(\Omega)$  for all  $\eta \in [1, +\infty[$  (see, for example, Gasiński and Papageorgiou [8], p.222) while now  $p^* = +\infty$ . So, in the above argument, we replace  $p^*$  by some  $\eta > r$  large enough s.t.

$$tr = \frac{\eta(r - \tau)}{\eta - \tau} < p \quad (\text{see } \mathbf{H} \text{ (ii)}).$$

Then, again we deduce that  $(u_n)$  is bounded in  $W$  and (26) holds. So, the Claim is proved.

On (20) we act with  $u_n - u^* \in W$  and we pass to the limit as  $n \rightarrow \infty$ . We obtain

$$\lim_n \langle A(u_n), u_n - u^* \rangle = 0 \quad (\text{see (26)}),$$

which implies

$$(29) \quad u_n \rightarrow u^* \quad \text{in } W \quad (\text{see Proposition 3}).$$

Therefore, if on (20) we pass to the limit as  $n \rightarrow \infty$  and use (29), then

$$A(u^*) = N_f^{\lambda^*}(u^*),$$

i.e.  $u^* \in C_+$  (by nonlinear regularity theory) and it solves  $(P_{\lambda^*})$ .

We need to show that  $u^* \neq 0$ . We argue by contradiction. So, suppose  $u^* = 0$  and consider the following auxiliary Dirichlet problem:

$$(30) \quad \begin{cases} -\Delta_p w = \eta_0(w^+)^{\mu-1} & \text{in } \Omega, \\ w = 0 & \text{on } \partial\Omega \end{cases}$$

(see **H** (iii)). Since  $\mu < p$ , the energy functional for (30), defined by

$$\psi(w) = \frac{1}{p} \|Dw\|_p^p - \frac{\eta_0}{\mu} \|w^+\|_{\mu}^{\mu} \quad \text{for all } w \in W,$$

is coercive and of course it is sequentially weakly l.s.c. Hence, by the Weierstrass theorem, we can find a minimizer  $w \in W$  of  $\psi$ . Note that, since  $\mu < p$ , we have

$$\psi(w) = \inf_{u \in W} \psi(u) < 0 = \psi(0),$$

so  $w \in W \setminus \{0\}$ . Then

$$A(w) = \eta_0(w^+)^{\mu-1} \quad \text{in } W^*,$$

which implies  $w \in \text{int}(C_+)$  and it solves (30).

From Ladyzhenskaya and Uraltseva [16] (p. 286, see also [20], p. 311) we can find  $\hat{M} > 0$  s.t.  $\|u\|_\infty \leq \hat{M}$  for all  $n \geq 1$ . Then we can apply Theorem 1 of Lieberman [18] (see also [20], p. 312) and find  $\alpha \in ]0, 1[$  and  $c_7 > 0$  s.t.

$$u_n \in C_0^{1,\alpha}(\bar{\Omega}) \quad \text{and} \quad \|u_n\|_{C_0^{1,\alpha}(\bar{\Omega})} \leq c_7 \quad \text{for all } n \in \mathbb{N}.$$

Recalling that  $C_0^{1,\alpha}(\bar{\Omega})$  is compactly embedded in  $C^1(\bar{\Omega})$ , we may assume that  $u_n \rightarrow u^* = 0$  in  $C_0^1(\bar{\Omega})$  as  $n \rightarrow \infty$ , so there exists  $n_0 \in \mathbb{N}$  s.t.

$$(31) \quad 0 \leq u_n(z) \leq \delta_0 \quad \text{for all } z \in \bar{\Omega} \quad \text{and all } n \geq n_0.$$

Fix  $n \geq n_0$  and choose  $t_n > 0$  s.t.

$$t_n w(z) \leq u_n(z) \quad \text{for all } z \in \bar{\Omega} \quad (\text{recall } u_n \in \text{int}(C_+) \text{ and use Lemma 2}).$$

Let  $t_n$  be the biggest such number and suppose that  $t_n \in ]0, 1[$ . Set  $\xi = \|u_n\|_\infty$ ,  $I = ]0, \lambda^*[$  and let  $\sigma_n = \sigma_\xi^I$  be as in hypothesis **H** (v). Then

$$\begin{aligned} & -\Delta_p(t_n w) + \sigma_n(t_n w)^{p-1} \\ &= t_n^{p-1} \eta_0 w^{\mu-1} + \sigma_n(t_n w)^{p-1} \quad (\text{see (30)}) \\ &< \eta_0(t_n w)^{\mu-1} + \sigma_n(t_n w)^{p-1} \quad (\text{recall that } t_n \in ]0, 1[ \text{ and } \mu < p) \\ &\leq \eta_0 u_n^{\mu-1} + \sigma_n u_n^{p-1} \quad (\text{since } t_n w \leq u_n) \\ &\leq f(z, u_n, \lambda_n) + \sigma_n u_n^{p-1} \quad (\text{since } n \geq n_0, \text{ see (31) and hypothesis H (iii)}) \\ &= -\Delta_p u_n + \sigma_n u_n^{p-1}. \end{aligned}$$

Note that if we set

$$h_1(z) = t_n^{p-1} \eta_0 w^{\mu-1} + \sigma_n(t_n w)^{p-1}, \quad h_2(z) = \eta_0 u_n^{\mu-1} + \sigma_n u_n^{p-1},$$

then  $h_1, h_2 \in C(\bar{\Omega})$  and

$$h_1(z) < h_2(z) \quad \text{for all } z \in \Omega.$$

Moreover, we have

$$h_2(z) \leq f(z, u_n, \lambda_n) + \sigma_n u_n^{p-1} \quad \text{a.e. in } \Omega.$$

Therefore, we can apply Proposition 2.6 of Arcoya and Ruiz [3] (see also Guedda and Veron [10]) and we have

$$u_n - t_n w \in \text{int}(C_+),$$

which contradicts the maximality of  $t_n$ . Therefore  $t_n \geq 1$  and so we have  $w \leq u_n$  for all  $n \geq n_0$ , hence  $w \leq 0$ , a contradiction. Thus,  $u^* \neq 0$ .

As before, by using hypothesis **H** (v) and the nonlinear maximum principle of Vázquez [23], we have  $u^* \in \text{int}(C_+)$ . So,  $\lambda^* \in \mathcal{P}$ , i.e.,  $\mathcal{P} = ]0, \lambda^*]$ .  $\square$

**Proposition 9.** *If hypotheses **H** hold, then for all  $\lambda \in ]0, \lambda^*]$  problem  $(P_\lambda)$  has at least two positive smooth solutions  $u_0, \hat{u} \in \text{int}(C_+)$  s.t.  $u_0 \leq \hat{u}$  in  $\bar{\Omega}$  and  $u_0 \neq \hat{u}$ .*

Proof. From Proposition 8, we know that  $\lambda^* \in \mathcal{P}$ , i.e., there is a solution  $u^* \in \text{int}(C_+)$  for problem  $(P_{\lambda^*})$ . We have

$$(32) \quad A(u^*) = N_f^{\lambda^*}(u^*) \geq N_f^\lambda(u^*) \quad \text{in } W^* \quad (\text{see } \mathbf{H} \text{ (iv)}),$$

so  $u^*$  is an upper solution of  $(P_\lambda)$  when  $\lambda \in ]0, \lambda^*]$ . In what follows  $\lambda \in ]0, \lambda^*]$ . We truncate  $x \mapsto f(z, x, \lambda)$  at  $u^*(z)$  and, using the direct method and (32), as in the proof of Proposition 6, we obtain a solution  $u_0 \in \text{int}(C_+)$  for problem  $(P_\lambda)$ , s.t.  $0 \leq u_0(z) \leq u^*(z)$  for all  $z \in \bar{\Omega}$ . For  $\xi = \|u^*\|_\infty$  and  $I = ]0, \lambda^*]$ , let  $\hat{\sigma} = \sigma_\xi^I$  be as postulated by hypothesis **H** (v). We have

$$\begin{aligned} & -\Delta_p u_0 + \hat{\sigma} u_0^{p-1} \\ &= f(z, u_0, \lambda) + \hat{\sigma} u_0^{p-1} \\ &= f(z, u_0, \lambda^*) + \hat{\sigma} u_0^{p-1} + \hat{\theta}(z) \quad (\text{we set } \hat{\theta}(z) = f(z, u_0, \lambda) - f(z, u_0, \lambda^*)) \\ &\leq f(z, u^*, \lambda^*) + \hat{\sigma}(u^*)^{p-1} + \hat{\theta}(z) \quad (\text{see } \mathbf{H} \text{ (v)} \text{ and recall } u_0 \leq u^*) \\ &= -\Delta_p u^* + \hat{\sigma}(u^*)^{p-1} + \hat{\theta}(z). \end{aligned}$$

By virtue of hypothesis **H** (iv), for every compact  $K \subset \Omega$ , we have

$$\text{esssup}_K \hat{\theta} < 0.$$

Invoking Proposition 2.6 of Arcoya and Ruiz [3], we have

$$(33) \quad u^* - u_0 \in \text{int}(C_+).$$

We consider the following truncation of  $x \mapsto f(z, x, \lambda)$ :

(34)

$$g(z, x, \lambda) = \begin{cases} f(z, u_0(z), \lambda) & \text{if } x \leq u_0(z), \\ f(z, x, \lambda) & \text{if } x > u_0(z), \end{cases} \quad \text{for a.a. } z \in \Omega \text{ and all } x \in \mathbb{R}, \lambda \in ]0, \lambda^*].$$

This is a Carathéodory function. We set

$$G(z, x, \lambda) = \int_0^x g(z, s, \lambda) ds \quad \text{for a.a. } z \in \Omega \quad \text{and all } x \in \mathbb{R}, \lambda \in ]0, \lambda^*[$$

and consider the  $C^1$  functional  $\psi_\lambda: W \rightarrow \mathbb{R}$  defined by

$$\psi_\lambda(u) = \frac{1}{p} \|Du\|_p^p - \int_\Omega G(z, u, \lambda) dz \quad \text{for all } u \in W.$$

**Claim 1.**  $\psi_\lambda$  satisfies the  $C$ -condition.

Let  $(u_n) \in W$  be a sequence s.t.

$$(35) \quad |\psi_\lambda(u_n)| \leq c_8 \quad \text{for all } n \in \mathbb{N} \quad (c_8 > 0)$$

and

$$(36) \quad \lim_n (1 + \|u_n\|) \psi'_\lambda(u_n) = 0 \quad \text{in } W^*.$$

From (35) we have

$$(37) \quad \|Du_n\|_p^p - \int_\Omega pG(z, u_n, \lambda) dz \leq pc_8 \quad \text{for all } n \in \mathbb{N}.$$

From (36) we have

$$(38) \quad \left| A(u_n, v) - \int_\Omega g(z, u_n, \lambda) v dz \right| \leq \varepsilon_n \frac{\|v\|}{1 + \|u_n\|} \quad \text{for all } v \in W, n \in \mathbb{N} \quad (\varepsilon_n \rightarrow 0^+ \text{ as } n \rightarrow \infty).$$

In (38) we choose  $v = -u_n^- \in W$ . Then,

$$\begin{aligned} \|Du_n^-\|_p^p &\leq \varepsilon_n + \int_\Omega f(z, u_n^-, \lambda) (-u_n^-) dz \quad (\text{see (34)}) \\ &\leq c_9(1 + \|Du_n^-\|_p) \quad \text{for some } c_9 > 0 \quad (\text{see H (i)}), \end{aligned}$$

which implies that  $(u_n^-)$  is bounded in  $W$ .

Next, in (38) we choose  $v = u_n^+ \in W$ . Then,

$$(39) \quad -\|Du_n^+\|_p^p + \int_\Omega g(z, u_n^+, \lambda) u_n^+ dz \leq \varepsilon_n \quad \text{for all } n \in \mathbb{N}.$$

We add (37) and (39) and use (34) and the boundedness of  $(u_n^-)$  to obtain, for all  $n \in \mathbb{N}$ ,

$$(40) \quad \int_\Omega [f(z, u_n^+, \lambda) u_n^+ - pF(z, u_n^+, \lambda)] dz \leq c_{10} \quad (c_{10} > 0).$$

From (40), using hypothesis **H** (ii) and the interpolation inequality, as in the proof of Proposition 8, we show that  $(u_n^+)$  is bounded in  $W$  as well. Thus,  $(u_n)$  is bounded in  $W$ . So, we may assume that there exists  $u \in W$  s.t.

$$u_n \rightharpoonup u \quad \text{in } W \quad \text{and} \quad u_n \rightarrow u \quad \text{in } L^r(\Omega) \quad \text{as } n \rightarrow \infty,$$

from which, using as before Proposition 3, we show that  $u_n \rightarrow u$  in  $W$  (as in the proof of Proposition 8), hence  $\psi_\lambda$  satisfies the  $C$ -condition. This proves Claim 1.

**Claim 2.**  $u_0$  is a local minimizer of  $\psi_\lambda$ .

We can always assume that  $u_0$  is the only nontrivial positive solution of problem  $(P_\lambda)$  in the order interval

$$\mathcal{I} = \{u \in W : 0 \leq u(z) \leq u^*(z) \text{ for a.a. } z \in \Omega\},$$

or otherwise we already have a second nontrivial smooth solution and we are done (see also [9]).

We introduce the following truncation of  $x \mapsto g(z, x, \lambda)$ :

$$(41) \quad \hat{g}(z, x, \lambda) = \begin{cases} f(z, u_0(z), \lambda) & \text{if } x \leq u_0(z), \\ f(z, x, \lambda) & \text{if } u_0(z) < x < u^*(z), \\ f(z, u^*(z), \lambda) & \text{if } x \geq u^*(z), \end{cases}$$

for a.a.  $z \in \Omega$  and all  $x \in \mathbb{R}$ ,  $\lambda \in \mathbb{R}_0^+$ . This is a Carathéodory function. As usual, we set

$$\hat{G}(z, x, \lambda) = \int_0^x \hat{g}(z, s, \lambda) ds \quad \text{for a.a. } z \in \Omega \quad \text{and all } x \in \mathbb{R}, \lambda \in \mathbb{R}_0^+$$

and consider the functional  $\hat{\psi}_\lambda \in C^1(W)$  given by

$$\hat{\psi}_\lambda(u) = \frac{1}{p} \|Du\|_p^p - \int_{\Omega} \hat{G}(z, u, \lambda) dz \quad \text{for all } u \in W.$$

Evidently  $\hat{\psi}_\lambda$  is coercive (see (41)) and is as well sequentially weakly l.s.c. So, we can find  $\hat{u}_0 \in W$  s.t.

$$\hat{\psi}_\lambda(\hat{u}_0) = \inf_W \hat{\psi}_\lambda,$$

in particular  $\hat{u}_0$  is a critical point of  $\hat{\psi}_\lambda$ , i.e.

$$(42) \quad A(\hat{u}_0) = N_g^\lambda(\hat{u}_0) \quad \text{in } W^* \quad (N_g^\lambda \text{ defined as in (8)}).$$

On (42) we act with  $(u_0 - \hat{u}_0)^+ \in W$ . Then

$$\begin{aligned} \langle A(\hat{u}_0), (u_0 - \hat{u}_0)^+ \rangle &= \int_{\Omega} \hat{g}(z, \hat{u}_0, \lambda)(u_0 - \hat{u}_0)^+ dz \\ &= \int_{\Omega} f(z, u_0, \lambda)(u_0 - \hat{u}_0)^+ dz \quad (\text{since } u_0 \leq u^*, \text{ see (41)}) \\ &= \langle A(u_0), (u_0 - \hat{u}_0)^+ \rangle, \end{aligned}$$

which implies

$$\begin{aligned} \langle A(u_0) - A(\hat{u}_0), (u_0 - \hat{u}_0)^+ \rangle &= \int_{\{u_0 > \hat{u}_0\}} (|Du_0|^{p-2} Du_0 - |D\hat{u}_0|^{p-2} D\hat{u}_0) \cdot (Du_0 - D\hat{u}_0) dz \\ &= 0. \end{aligned}$$

So

$$|\{u_0 > \hat{u}_0\}|_N = 0,$$

i.e.  $u_0 \leq \hat{u}_0$ . Also, acting on (42) with  $(\hat{u}_0 - u^*)^+ \in W$ , we have

$$\begin{aligned} \langle A(\hat{u}_0), (\hat{u}_0 - u^*)^+ \rangle &= \int_{\Omega} \hat{g}(z, \hat{u}_0, \lambda)(\hat{u}_0 - u^*)^+ dz \\ &= \int_{\Omega} f(z, u^*, \lambda)(\hat{u}_0 - u^*)^+ dz \quad (\text{see (41) and recall } u_0 \leq u^*) \\ &\leq \langle A(u^*), (\hat{u}_0 - u^*)^+ \rangle \quad (\text{see (32)}), \end{aligned}$$

i.e.

$$\begin{aligned} \langle A(\hat{u}_0) - A(u^*), (\hat{u}_0 - u^*)^+ \rangle &= \int_{\{\hat{u}_0 > u^*\}} (|D\hat{u}_0|^{p-2} D\hat{u}_0 - |Du^*|^{p-2} Du^*) \cdot (D\hat{u}_0 - Du^*) dz \\ &\leq 0. \end{aligned}$$

So

$$|\{\hat{u}_0 > u^*\}|_N = 0,$$

i.e.  $\hat{u}_0 \leq u^*$ . Hence, (42) becomes

$$A(\hat{u}_0) = N_f^\lambda(\hat{u}_0) \quad \text{in } W^* \quad (\text{see (41) and (34)})$$

and  $\hat{u}_0 \in \text{int}(C_+) \cap \mathcal{I}$  is a solution of problem  $(P_\lambda)$ . This implies

$$\hat{u}_0 = u_0 \quad (\text{recall that } u_0 \text{ is the only nontrivial solution of } (P_\lambda) \text{ in } \mathcal{I}).$$

Note that

$$\hat{\psi}_\lambda(u) = \psi_\lambda(u) \quad \text{for all } u \in \mathcal{I}.$$

Recall, also, that  $u^* - u_0 \in \text{int}(C_+)$  (see (33)) and  $u_0 \in \text{int}(C_+)$ . Therefore,  $\mathcal{I}$  is a neighborhood of  $u_0$  in the topology of  $C_0^1(\bar{\Omega})$ , and so  $u_0$  is a local  $C_0^1(\bar{\Omega})$ -minimizer of  $\psi_\lambda$ . By virtue of Theorem 1.2 of Garcia Azorero, Manfredi and Peral Alonso [7], it is also a local  $W$ -minimizer of  $\psi_\lambda$ . This proves Claim 2.

We may assume that  $u_0$  is an isolated critical point of  $\psi_\lambda$  (otherwise we have a whole sequence of distinct positive smooth solutions converging to  $u_0$ ). Therefore we can find  $\rho \in ]0, 1[$  small enough s.t.

$$(43) \quad \psi_\lambda(u_0) < \inf_{\|u-u_0\|=\rho} \psi_\lambda(u) = \eta_\rho$$

(see Aizicovici, Papageorgiou and Staicu [1], proof of Proposition 29).

Clearly hypothesis **H** (ii) implies that

$$(44) \quad \lim_{t \rightarrow +\infty} \psi_\lambda(t\hat{u}_1) = -\infty.$$

Then, (43), (44) and Claim 1 permit the use of Theorem 1 (the mountain pass theorem). So, we obtain  $\hat{u} \in W$  s.t.

$$(45) \quad \psi_\lambda(u_0) < \eta_\rho \leq \psi_\lambda(\hat{u}) \quad (\text{see (43)})$$

and

$$(46) \quad \psi'_\lambda(\hat{u}) = 0.$$

From (45) we have  $\hat{u} \neq u_0$ . From (46), we have

$$(47) \quad A(\hat{u}) = N_g^\lambda(\hat{u}) \quad \text{in } W^*.$$

Acting on (47) with  $(u_0 - \hat{u})^+ \in W$ , as before we show that  $u_0 \leq \hat{u}$ . Hence (47) becomes

$$A(\hat{u}) = N_f^\lambda(\hat{u}) \quad \text{in } W^* \quad (\text{see (34)}),$$

so  $\hat{u} \in \text{int}(C_+)$  (nonlinear regularity) is a solution of  $(P_\lambda)$ .  $\square$

Summarizing the situation, we have the following bifurcation-type result for problem  $(P_\lambda)$ .

**Theorem 10.** *If hypotheses **H** hold, then there exists  $\lambda^* \in \mathbb{R}_0^+$  s.t.*

- (a) *for every  $\lambda \in ]0, \lambda^*[$  problem  $(P_\lambda)$  has at least two positive smooth solutions  $u_0, \hat{u} \in \text{int}(C_+)$  s.t.  $u_0 \leq \hat{u}$  in  $\bar{\Omega}$  and  $u \neq \hat{u}$ ;*
- (b) *for  $\lambda = \lambda^*$  problem  $(P_\lambda)$  has at least one positive smooth solution  $u^* \in \text{int}(C_+)$ ;*
- (c) *for every  $\lambda > \lambda^*$  problem  $(P_\lambda)$  has no positive solution.*

REMARK 11. If  $p = 2$  and  $0 < \lambda < \lambda^*$ , then the two positive solutions  $u_0, \hat{u} \in \text{int}(C_+)$  satisfy

$$\hat{u} - u_0 \in \text{int}(C_+).$$

Indeed, if  $\xi = \|\hat{u}\|_\infty$  and  $I = ]0, \lambda^*]$ , then we find  $\hat{\sigma} = \sigma_\xi^I$  as in hypothesis **H** (v) and we have

$$\begin{aligned} -\Delta(\hat{u} - u_0) + \hat{\sigma}(\hat{u} - u_0) &= f(z, \hat{u}, \lambda) + \hat{\sigma}\hat{u} - f(z, u_0, \lambda) - \hat{\sigma}u_0 \\ &\geq 0 \quad (\text{see } \mathbf{H} \text{ (v)}), \end{aligned}$$

i.e.

$$\Delta(\hat{u} - u_0) \leq \hat{\sigma}(\hat{u} - u_0) \quad \text{a.e. in } \Omega,$$

which implies

$$\hat{u} - u_0 \in \text{int}(C_+) \quad (\text{see Vázquez [23]}).$$

Finally, note that, if  $f(z, \cdot, \lambda) \in C^1(\mathbb{R})$ , then by the mean value theorem **H** (v) is automatically true.

#### 4. Problems with $(p - 1)$ -linear nonlinearities near zero

In the previous section, we examined problems in which the reaction was concave near the origin (see hypothesis **H** (iii)). Here, we consider equations in which  $x \mapsto f(z, x, \lambda)$  exhibits  $(p - 1)$ -linear growth near zero. We show that in this case we can still have a bifurcation-type theorem similar to Theorem 10.

The new hypotheses on the nonlinearity  $f(z, x, \lambda)$  are the following.

**H'**  $f: \Omega \times \mathbb{R} \times \mathbb{R}_0^+ \rightarrow \mathbb{R}$  is a Carathéodory function s.t.  $f(z, 0, \lambda) = 0$  for a.a.  $z \in \Omega$  and all  $\lambda > 0$ . We set

$$F(z, x, \lambda) = \int_0^x f(z, s, \lambda) ds \quad \text{for a.a. } z \in \Omega \quad \text{and all } x \in \mathbb{R}, \lambda > 0.$$

Let hypotheses **H'** (i), (ii), (iv), (v) be as **H** (i), (ii), (iv), (v) and

(iii) for all bounded  $I \subset \mathbb{R}_0^+$  there exist  $\eta_0 \in L^\infty(\Omega)$ ,  $\eta_0(z) \geq \hat{\lambda}_1$  for a.a.  $z \in \Omega$ ,  $\eta_0 \neq \hat{\lambda}_1$ , and  $\eta_0 > 0$  s.t.

$$\begin{aligned} \eta_0(z) &\leq \liminf_{x \rightarrow 0^+} \frac{f(z, x, \lambda)}{x^{p-1}} \leq \limsup_{x \rightarrow 0^+} \frac{f(z, x, \lambda)}{x^{p-1}} \\ &\leq \eta_1 \quad \text{uniformly for a.a. } z \in \Omega \quad \text{and all } \lambda \in I. \end{aligned}$$

EXAMPLE 12. Let  $\eta > \hat{\lambda}_1$ ,  $1 < q < p < r < p^*$ . The following function satisfies hypotheses **H'**:

$$f(x, \lambda) = \begin{cases} \lambda x^{r-1} + \eta x^{p-1} & \text{if } 0 \leq x \leq 1, \\ \lambda x^{q-1} + p\eta x^{p-1} \left( \ln(x) + \frac{1}{p} \right) & \text{if } x > 1, \end{cases} \quad \text{for a.a. } z \in \Omega \text{ and all } \lambda \in \mathbb{R}_0^+.$$

Again this  $(p-1)$ -superlinear function (at  $\infty$ ) does not satisfy the *AR*-condition.

A careful inspection of the proofs in Section 3 reveals that they remain essentially unchanged. The only two parts which need to be modified are the following:

(A) in the proof of Proposition 6, the part where we show that the minimizer  $u_0$  is nontrivial;

(B) in the proof of Proposition 8, the part where we show that  $u^* \neq 0$ .

First we deal with (A). By virtue of hypothesis **H'** (iii), given  $\varepsilon > 0$ , we can find  $\delta > 0$  s.t.

$$(48) \quad F(z, x, \lambda) \geq \frac{1}{p}(\eta_0(z) - \varepsilon)x^p \quad \text{for a.a. } z \in \Omega, \text{ all } x \in [0, \delta] \text{ and all } \lambda \in ]0, \tilde{\lambda}[.$$

Let  $t \in ]0, 1[$  be s.t.

$$(49) \quad 0 \leq t\hat{u}_1(z) \leq \min\{\delta, \tilde{u}(z)\} \quad \text{for all } z \in \overline{\Omega} \quad (\text{see Lemma 2}).$$

Then,

$$\begin{aligned} \tilde{\varphi}_\lambda(t\hat{u}_1) &= \frac{t^p}{p} \|D\hat{u}_1\|_p^p - \int_{\Omega} F(z, t\hat{u}_1, \lambda) dz \quad (\text{see (12) and (49)}) \\ &\leq \frac{t^p}{p} \int_{\Omega} (\hat{\lambda}_1 - \eta_0(z))\hat{u}_1(z)^p dz + \frac{t^p}{p}\varepsilon \quad (\text{see (48), (49) and recall } \|\hat{u}_1\|_p = 1). \end{aligned}$$

Since

$$\int_{\Omega} (\hat{\lambda}_1 - \eta_0(z))\hat{u}_1(z)^p dz < 0,$$

by choosing  $\varepsilon > 0$  small enough we see that

$$\tilde{\varphi}_\lambda(u_0) \leq \tilde{\varphi}_\lambda(t\hat{u}_1) < 0 \quad (\text{see (13)}),$$

i.e.  $u_0 \neq 0$ .

Next we deal with (B). Again we argue indirectly. So, suppose that  $u^* = 0$ . Then,  $u_n \rightarrow 0$  in  $W$  (see (29) and in fact, using Theorem 1 of Lieberman [18], we show that  $u_n \rightarrow 0$  in  $C_0^1(\overline{\Omega})$  as  $n \rightarrow \infty$  (see the proof of Proposition 8). Therefore we can find  $n_0 \in \mathbb{N}$  s.t.

$$0 \leq u_n(z) \leq 1 \quad \text{for all } n \geq n_0 \quad \text{and } z \in \overline{\Omega}.$$

Hypotheses **H'** (i), (iii) imply that

$$|f(z, x, \lambda)| \leq c_{11}|xt^{p-1}| \quad \text{for a.a. } z \in \Omega \text{ and all } x \in [0, 1], \lambda \in ]0, \lambda^*[ \quad (c_{11} > 0),$$

which implies

$$|f(z, u_n(z), \lambda_n)| \leq c_{11}|u_n(z)|^{p-1} \quad \text{for a.a. } z \in \Omega \text{ and all } n \geq n_0.$$

So, the sequence

$$\left( \frac{N_f^{\lambda_n}(u_n)}{\|u_n\|^{p-1}} \right)$$

is bounded in  $L^{p'}(\Omega)$ . Hence, passing if necessary to a subsequence, we may assume that

$$(50) \quad \frac{N_f^{\lambda_n}(u_n)}{\|u_n\|^{p-1}} \rightharpoonup h \quad \text{in} \quad L^{p'}(\Omega) \quad \text{as} \quad n \rightarrow \infty.$$

Set  $y_n = u_n/\|u_n\|$  for all  $n \in \mathbb{N}$ . Then  $|y_n| = 1$  for all  $n \in \mathbb{N}$  and so we may assume that

$$(51) \quad y_n \rightharpoonup y \quad \text{in} \quad W \quad \text{and} \quad y_n \rightarrow y \quad \text{in} \quad L^p(\Omega) \quad \text{as} \quad n \rightarrow \infty.$$

Recall that

$$(52) \quad A(y_n) = \frac{N_f^{\lambda_n}(u_n)}{\|u_n\|^{p-1}} \quad \text{for all} \quad n \in \mathbb{N} \quad (\text{see (20)}).$$

Acting on (52) with  $y_n - y \in W$ , passing to the limit as  $n \rightarrow \infty$  and using (50) and (51), we obtain

$$\lim_n \langle A(y_n), y_n - y \rangle = 0,$$

hence  $y_n \rightarrow y$  (see Proposition 3). In particular, we have

$$(53) \quad \|y\| = 1 \quad \text{and} \quad y(z) \geq 0 \quad \text{for a.a.} \quad z \in \Omega.$$

Moreover, using hypothesis **H'** (iii) and reasoning as in the proof of Theorem 2.8 of [14] (see also [1], proof of Proposition 31), we show that there exists  $m \in L^\infty(\Omega)$  s.t.

$$(54) \quad h(z) = m(z)y(z)^{p-1} \quad \text{and} \quad \eta_0(z) \leq m(z) \leq \eta_1 \quad \text{for a.a.} \quad z \in \Omega.$$

So, if in (52) we pass to the limit as  $n \rightarrow \infty$  and use (53) and (54), then

$$A(y) = m(z)y^{p-1},$$

i.e.,  $y \in W$  solves the Dirichlet problem

$$(55) \quad \begin{cases} -\Delta_p y = m(z)y^{p-1} & \text{in} \quad \Omega, \\ y = 0 & \text{on} \quad \partial\Omega. \end{cases}$$

But, note that

$$\hat{\lambda}_1(m) < \hat{\lambda}_1(\hat{\lambda}_1) = 1 \quad (\text{see (3) and (54)}).$$

So, from (55) it follows that  $y$  must be nodal, contradicting (53). This proves that  $u^* \neq 0$ .

So, we can state the following bifurcation-type theorem.

**Theorem 13.** *If hypotheses  $H'$  hold, then there exists  $\lambda^* \in \mathbb{R}_0^+$  s.t.*

- (a) *for every  $\lambda \in ]0, \lambda^*[$  problem  $(P_\lambda)$  has at least two positive smooth solutions  $u_0, \hat{u} \in \text{int}(C_+)$  s.t.  $u_0 \leq \hat{u}$  in  $\overline{\Omega}$  and  $u_0 \neq \hat{u}$ ;*
- (b) *for  $\lambda = \lambda^*$  problem  $(P_\lambda)$  has at least one positive smooth solution  $u^* \in \text{int}(C_+)$ ;*
- (c) *for every  $\lambda > \lambda^*$  problem  $(P_\lambda)$  has no positive solution.*

ACKNOWLEDGMENT. The authors wish to thank a knowledgeable Referee for her/his corrections and remarks.

---

### References

- [1] S. Aizicovici, N.S. Papageorgiou and V. Staicu: *Degree theory for operators of monotone type and nonlinear elliptic equations with inequality constraints*, Mem. Amer. Math. Soc. **196** (2008).
- [2] A. Ambrosetti, H. Brezis and G. Cerami: *Combined effects of concave and convex nonlinearities in some elliptic problems*, J. Funct. Anal. **122** (1994), 519–543.
- [3] D. Arcoya and D. Ruiz: *The Ambrosetti–Prodi problem for the  $p$ -Laplacian operator*, Comm. Partial Differential Equations **31** (2006), 849–865.
- [4] F. Brock, L. Iturriaga and P. Ubilla: *A multiplicity result for the  $p$ -Laplacian involving a parameter*, Ann. Henri Poincaré **9** (2008), 1371–1386.
- [5] D.G. Costa and C.A. Magalhães: *Existence results for perturbations of the  $p$ -Laplacian*, Nonlinear Anal. **24** (1995), 409–418.
- [6] G. Fei: *On periodic solutions of superquadratic Hamiltonian systems*, Electron. J. Differential Equations **2002**.
- [7] J.P. García Azorero, I. Peral Alonso and J.J. Manfredi: *Sobolev versus Hölder local minimizers and global multiplicity for some quasilinear elliptic equations*, Commun. Contemp. Math. **2** (2000), 385–404.
- [8] L. Gasiński and N.S. Papageorgiou: *Nonlinear Analysis*, Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [9] L. Gasiński and N.S. Papageorgiou: *Nodal and multiple constant sign solutions for resonant  $p$ -Laplacian equations with a nonsmooth potential*, Nonlinear Anal. **71** (2009), 5747–5772.
- [10] M. Guedda and L. Véron: *Quasilinear elliptic equations involving critical Sobolev exponents*, Nonlinear Anal. **13** (1989), 879–902.
- [11] Z.M. Guo: *Some existence and multiplicity results for a class of quasilinear elliptic eigenvalue problems*, Nonlinear Anal. **18** (1992), 957–971.
- [12] Z. Guo and Z. Zhang:  *$W^{1,p}$  versus  $C^1$  local minimizers and multiplicity results for quasilinear elliptic equations*, J. Math. Anal. Appl. **286** (2003), 32–50.
- [13] S. Hu and N.S. Papageorgiou: *Multiple positive solutions for nonlinear eigenvalue problems with the  $p$ -Laplacian*, Nonlinear Anal. **69** (2008), 4286–4300.
- [14] S. Hu and N.S. Papageorgiou: *Multiplicity of solutions for parametric  $p$ -Laplacian equations with nonlinearity concave near the origin*, Tohoku Math. J. (2) **62** (2010), 137–162.
- [15] L. Jeanjean: *On the existence of bounded Palais–Smale sequences and application to a Landesman–Lazer-type problem set on  $\mathbf{R}^N$* , Proc. Roy. Soc. Edinburgh Sect. A **129** (1999), 787–809.

- [16] O.A. Ladyzhenskaya and N.N. Ural'tseva: *Linear and Quasilinear Elliptic Equations*, Academic Press, New York, 1968.
- [17] G. Li and C. Yang: *The existence of a nontrivial solution to a nonlinear elliptic boundary value problem of  $p$ -Laplacian type without the Ambrosetti–Rabinowitz condition*, *Nonlinear Anal.* **72** (2010), 4602–4613.
- [18] G.M. Lieberman: *Boundary regularity for solutions of degenerate elliptic equations*, *Nonlinear Anal.* **12** (1988), 1203–1219.
- [19] O.H. Miyagaki and M.A.S. Souto: *Superlinear problems without Ambrosetti and Rabinowitz growth condition*, *J. Differential Equations* **245** (2008), 3628–3638.
- [20] N.S. Papageorgiou and S.Th. Kyritsi-Yiallourou: *Handbook of Applied Analysis*, Springer, New York, 2009.
- [21] M. Schechter and W. Zou: *Superlinear problems*, *Pacific J. Math.* **214** (2004), 145–160.
- [22] S. Takeuchi: *Multiplicity result for a degenerate elliptic equation with logistic reaction*, *J. Differential Equations* **173** (2001), 138–144.
- [23] J.L. Vázquez: *A strong maximum principle for some quasilinear elliptic equations*, *Appl. Math. Optim.* **12** (1984), 191–202.

Antonio Iannizzotto  
 Dipartimento di Informatica  
 Università degli Studi di Verona  
 Cá Vignal II, Strada Le Grazie 15  
 37134 Verona  
 Italy  
 e-mail: antonio.iannizzotto@univr.it

Nikolaos S. Papageorgiou  
 Department of Mathematics  
 National Technical University  
 Zografou Campus, 15780 Athens  
 Greece  
 e-mail: npapg@math.ntua.gr