

Title	SIGMA FUNCTIONS FOR TELESCOPIC CURVES
Author(s)	Ayano, Takanori
Citation	Osaka Journal of Mathematics. 51(2) P.459-P.480
Issue Date	2014-04
Text Version	publisher
URL	https://doi.org/10.18910/29199
DOI	10.18910/29199
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/repo/ouka/all/>

SIGMA FUNCTIONS FOR TELESCOPIC CURVES

TAKANORI AYANO

(Received January 6, 2012, revised September 19, 2012)

Abstract

In this paper we consider a symplectic basis of the first cohomology group and the sigma functions for algebraic curves expressed by a canonical form using a finite sequence (a_1, \dots, a_t) of positive integers whose greatest common divisor is equal to one (Miura [13]). The idea is to express a non-singular algebraic curve by affine equations of t variables whose orders at infinity are (a_1, \dots, a_t) . We construct a symplectic basis of the first cohomology group and the sigma functions for telescopic curves, i.e., the curves such that the number of defining equations is exactly $t - 1$ in the Miura canonical form. The largest class of curves for which such construction has been obtained thus far is (n, s) -curves ([4] [15]), which are telescopic because they are expressed in the Miura canonical form with $t = 2$, $a_1 = n$, and $a_2 = s$, and the number of defining equations is one.

1. Introduction

Recently the theory of Abelian functions is attracting increasing interest in mathematical physics and applied mathematics. In particular the sigma functions for algebraic curves have been studied actively. In this paper we construct sigma functions explicitly for a class of algebraic curves for which such construction has not been obtained thus far.

Let C be a compact Riemann surface of genus g and $H^1(C, \mathbb{C})$ the first cohomology group, which is defined by the linear space of second kind differentials modulo meromorphic exact forms. We say a meromorphic differential on C to be second kind if it is locally exact.

We consider a basis of $H^1(C, \mathbb{C})$ consisting of $\dim_{\mathbb{C}} H^1(C, \mathbb{C}) = 2g$ elements (cf. [11], pp.29–31, Theorems 8.1 and 8.2). In particular, in order to construct sigma functions explicitly, we wish to construct a basis (symplectic basis) $\{du_i, dr_i\}_{i=1}^g$ of $H^1(C, \mathbb{C})$ such that

1. du_i is holomorphic on C for each i , and
 2. $du_i \circ du_j = dr_i \circ dr_j = 0$ and $du_i \circ dr_j = \delta_{ij}$ for each i, j ,
- where the operator \circ is the intersection form on $H^1(C, \mathbb{C})$ defined by

$$\eta \circ \eta' = \sum_p \operatorname{Res} \left(\int^p \eta \right) \eta'(p)$$

for second kind differentials η, η' (the summation is over all the singular points of η and η' , and Res means taking a residue at a point).

In order to express defining equations of C , we use a canonical form for expressing non-singular algebraic curves introduced by Miura [13]. Given a finite sequence (a_1, \dots, a_t) of positive integers whose greatest common divisor is equal to one, Miura [13] introduced a non-singular algebraic curve determined by the sequence (a_1, \dots, a_t) . The idea is to express a non-singular algebraic curve by affine equations of t variables whose orders at infinity are (a_1, \dots, a_t) . Any non-singular algebraic curve is birationally equivalent to a curve expressed in the Miura canonical form (cf. [13]).

Klein [9] [10] extended the elliptic sigma functions to the case of hyperelliptic curves of genus g , which are expressed in the Miura canonical form with $t = 2$, $a_1 = 2$, and $a_2 = 2g + 1$. Bukhshtaber et al. [4] and Nakayashiki [15] extended Klein's sigma functions to the case of more general plane algebraic curves called (n, s) -curves, which are expressed in the Miura canonical form with $t = 2$, $a_1 = n$, and $a_2 = s$. In this paper we give an explicit construction of sigma functions for telescopic curves, i.e., the curves such that the number of defining equations is exactly $t - 1$ in the Miura canonical form. The telescopic curves contain the (n, s) -curves as special cases. Recently Matsutani [12] constructed sigma functions for $(3, 4, 5)$ -curves, which are not telescopic.

The plan of this paper is as follows. In Section 2 we recall the definition of the Miura canonical form. In Section 3 we construct the holomorphic 1-forms $\{du_i\}_{i=1}^g$ for the telescopic curves. In Section 4 we construct the second kind differentials $\{dr_i\}_{i=1}^g$ for the telescopic curves and show that the set $\{du_i, dr_i\}_{i=1}^g$ is a symplectic basis of the first cohomology group. In Section 5 we construct sigma functions for the telescopic curves.

Throughout this paper, $\mathbb{N}, \mathbb{N}_+, \mathbb{Z}$, and \mathbb{C} denote the set of non-negative integers, positive integers, integers, and complex numbers, respectively.

2. Miura canonical form

Miura [13] introduced a canonical form of defining equations for any non-singular algebraic curve. Here we recall the definition of the Miura canonical form.

Let $t \geq 2$, a_1, \dots, a_t positive integers such that $\text{GCD}\{a_1, \dots, a_t\} = 1$, $A_t = (a_1, \dots, a_t) \in \mathbb{N}_+^t$, and $\langle A_t \rangle = a_1\mathbb{N} + \dots + a_t\mathbb{N}$, assuming that the order of a_1, \dots, a_t is fixed. For the map $\Psi: \mathbb{N}^t \rightarrow \langle A_t \rangle$ defined by $\Psi((m_1, \dots, m_t)) = \sum_{i=1}^t a_i m_i$, we define the order $<$ in \mathbb{N}^t so that $M < M'$ for $M = (m_1, \dots, m_t)$ and $M' = (m'_1, \dots, m'_t)$ if

1. $\Psi(M) < \Psi(M')$ or
 2. $\Psi(M) = \Psi(M')$ and $m_1 = m'_1, \dots, m_{i-1} = m'_{i-1}, m_i > m'_i$ for some i ($1 \leq i \leq t$).
- Let $M(a)$ be the minimum element with respect to the order $<$ in \mathbb{N}^t satisfying $\Psi(M) = a \in \langle A_t \rangle$. We define $B(A_t) \subseteq \mathbb{N}^t$ and $V(A_t) \subseteq \mathbb{N}^t \setminus B(A_t)$ by

$$B(A_t) = \{M(a) \mid a \in \langle A_t \rangle\}$$

and

$$V(A_t) = \{L \in \mathbb{N}^t \setminus B(A_t) \mid L = M + N, M \in \mathbb{N}^t \setminus B(A_t), N \in \mathbb{N}^t \Rightarrow N = (0, \dots, 0)\},$$

respectively.

Hereafter $\mathbb{C}[X] := \mathbb{C}[X_1, \dots, X_t]$ denotes the polynomial ring over \mathbb{C} of t -variables X_1, \dots, X_t . For $A \subset \mathbb{C}[X]$, $\text{Span}\{A\}$ and (A) denote the linear space over \mathbb{C} generated by A and the ideal in $\mathbb{C}[X]$ generated by A , respectively. Also X^M , $M = (m_1, \dots, m_t)$, denotes $X^M = X_1^{m_1} \cdots X_t^{m_t}$ for simplicity.

For $M \in V(A_t)$ we define the polynomial $F_M(X) \in \mathbb{C}[X]$ by

$$(1) \quad F_M(X) = X^M - X^L - \sum_{\{N \in B(A_t) \mid \Psi(N) < \Psi(M)\}} \lambda_N X^N, \quad \lambda_N \in \mathbb{C},$$

where L is the element of $B(A_t)$ satisfying $\Psi(L) = \Psi(M)$. We assume that the set of polynomials $\{F_M \mid M \in V(A_t)\}$ satisfies the following condition:

$$(2) \quad \text{Span}\{X^N \mid N \in B(A_t)\} \cap (\{F_M \mid M \in V(A_t)\}) = \{0\}.$$

Let $I = (\{F_M \mid M \in V(A_t)\})$, $R = \mathbb{C}[X]/I$, x_i the image of X_i for the projection $\mathbb{C}[X] \rightarrow R$, and K the total quotient ring of R . Then we have the following three propositions. Because there exists no paper where proofs are written in English, we give complete proofs in Appendix.

- Proposition 2.1** (Miura [13]). (i) *The set $\{x^N \mid N \in B(A_t)\}$ is a basis of R over \mathbb{C} , where $x = (x_1, \dots, x_t)$.*
 (ii) *The ring R is an integral domain, therefore K is the quotient field of R .*
 (iii) *The field K is an algebraic function field of one variable over \mathbb{C} .*
 (iv) *There exists a discrete valuation v_∞ of K such that $(x_i)_\infty = a_i v_\infty$ for any i , where $(x_i)_\infty$ denotes the pole divisor of x_i (cf. [19] p. 19).*

Let $C^{\text{aff}} = \{(z_1, \dots, z_t) \in \mathbb{C}^t \mid f(z_1, \dots, z_t) = 0, \forall f \in I\}$. From Proposition 2.1 (ii) (iii), C^{aff} is an affine algebraic curve in \mathbb{C}^t . Hereafter we assume that C^{aff} is non-singular. For $k \in \mathbb{N}$ we define $L(kv_\infty) = \{f \in K \mid (f) + kv_\infty \geq 0\} \cup \{0\}$, where (f) denotes the divisor of f , i.e., $(f) = \sum_v v(f) \cdot v$.

- Proposition 2.2** (Miura [13]). (i) $R = \bigcup_{k=0}^\infty L(kv_\infty)$.
 (ii) *The map ϕ*

$$C^{\text{aff}} \rightarrow \{\text{discrete valuation of } K\} \setminus \{v_\infty\},$$

$$p \rightarrow v_p$$

is bijective, where v_p is the discrete valuation corresponding to $p \in C^{\text{aff}}$ (cf. [17], p. 21, 22).

Let C be the compact Riemann surface corresponding to C^{aff} . From Proposition 2.2 (ii), C is obtained from C^{aff} by adding one point, say ∞ , where the discrete valuation corresponding to ∞ is v_∞ . It is known that any non-singular algebraic curve is birationally equivalent to such C for some A_t (cf. [13]). Hereafter we represent each curve C by the sequence $A_t = (a_1, \dots, a_t)$ and call (a_1, \dots, a_t) -curve for short.

The sequence $A_t = (a_1, \dots, a_t)$ is called telescopic if for any i ($2 \leq i \leq t$)

$$\frac{a_i}{d_i} \in \frac{a_1}{d_{i-1}}\mathbb{N} + \dots + \frac{a_{i-1}}{d_{i-1}}\mathbb{N}, \quad d_i := \text{GCD}\{a_1, \dots, a_i\}.$$

Note that $A_2 = (a_1, a_2)$ is always telescopic.

Proposition 2.3 (Miura [13]). *If A_t is telescopic, then the condition (2) is satisfied and we have the following properties.*

- (i) $B(A_t) = \{(m_1, \dots, m_t) \in \mathbb{N}^t \mid 0 \leq m_i \leq d_{i-1}/d_i - 1, 2 \leq i \leq t\}$.
- (ii) $V(A_t) = \{(d_{i-1}/d_i)\mathbf{e}_i \mid 2 \leq i \leq t\}$, where \mathbf{e}_i is the i -th unit vector in \mathbb{Z}^t .
- (iii) The genus g of C is

$$(3) \quad g = \frac{1}{2} \left\{ (1 - a_1) + \sum_{i=2}^t \left(\frac{d_{i-1}}{d_i} - 1 \right) a_i \right\}.$$

Note that $\#V(A_t)$ is the number of defining equations, where $\#$ denotes the number of elements. From Lemma C.1 (iv) in Appendix, we obtain $\#V(A_t) \geq t - 1$. If A_t is telescopic, then from Proposition 2.3 (ii) we obtain $\#V(A_t) = t - 1$. On the other hand Suzuki [18] proved that if $\#V(A_t) = t - 1$, then A_t is telescopic.

From Proposition 2.3, the defining equations of a telescopic (a_1, \dots, a_t) -curve are given as follows: for $2 \leq i \leq t$,

$$F_i(X_1, \dots, X_t) = X_i^{d_{i-1}/d_i} - \prod_{j=1}^i X_j^{m_{ij}} - \sum \lambda_{j_1, \dots, j_i}^{(i)} X_1^{j_1} \cdots X_i^{j_i},$$

where $(m_{i1}, \dots, m_{it}) \in B(A_t)$ such that $\sum_{j=1}^i a_j m_{ij} = a_i d_{i-1}/d_i$, $\lambda_{j_1, \dots, j_i}^{(i)} \in \mathbb{C}$, and the sum is over all $(j_1, \dots, j_i) \in B(A_t)$ such that $\sum_{k=1}^i a_k j_k < a_i d_{i-1}/d_i$. Assign degrees as

$$\text{deg } X_k = a_k, \quad \text{deg } \lambda_{j_1, \dots, j_i}^{(i)} = a_i d_{i-1}/d_i - \sum_{k=1}^i a_k j_k.$$

EXAMPLE 1. $A_2 = (n, s)$, $n, s \in \mathbb{N}_+$, $\text{GCD}\{n, s\} = 1$.

Since $A_2 = (n, s)$ is telescopic, from Proposition 2.3 (ii), we have $V(A_2) = \{(0, n)\}$. Therefore we have

$$F_2(X_1, X_2) = X_2^n - X_1^s - \sum_{n_{j_1} + s_{j_2} < ns} \lambda_{j_1, j_2}^{(2)} X_1^{j_1} X_2^{j_2},$$

which is the (n, s) -curve introduced in [2]. In particular we obtain the elliptic curves if $n = 2$ and $s = 3$ and the hyperelliptic curves of genus g if $n = 2$ and $s = 2g + 1$.

EXAMPLE 2. $A_3 = (4, 6, 5)$.

Since $A_3 = (4, 6, 5)$ is telescopic, from Proposition 2.3 (ii), we have $V(A_3) = \{(0, 2, 0), (0, 0, 2)\}$. Therefore we have

$$F_2(X_1, X_2, X_3) = X_2^2 - X_1^3 - \lambda_{0,1,1}^{(2)} X_2 X_3 - \lambda_{1,1,0}^{(2)} X_1 X_2 - \lambda_{1,0,1}^{(2)} X_1 X_3 - \lambda_{2,0,0}^{(2)} X_1^2 - \lambda_{0,1,0}^{(2)} X_2 - \lambda_{0,0,1}^{(2)} X_3 - \lambda_{1,0,0}^{(2)} X_1 - \lambda_{0,0,0}^{(2)}$$

and

$$F_3(X_1, X_2, X_3) = X_3^2 - X_1 X_2 - \lambda_{1,0,1}^{(3)} X_1 X_3 - \lambda_{2,0,0}^{(3)} X_1^2 - \lambda_{0,1,0}^{(3)} X_2 - \lambda_{0,0,1}^{(3)} X_3 - \lambda_{1,0,0}^{(3)} X_1 - \lambda_{0,0,0}^{(3)}.$$

3. Holomorphic 1-forms for telescopic curves

Let C be a telescopic (a_1, \dots, a_t) -curve and $\Gamma(C, \Omega_C^1)$ the linear space consisting of holomorphic 1-forms on C . In this section we construct a basis of $\Gamma(C, \Omega_C^1)$. Let G be the matrix defined by

$$G := \begin{pmatrix} \frac{\partial F_2}{\partial X_1} & \dots & \frac{\partial F_2}{\partial X_t} \\ \dots & \dots & \dots \\ \frac{\partial F_t}{\partial X_1} & \dots & \frac{\partial F_t}{\partial X_t} \end{pmatrix}$$

and G_i the matrix obtained by removing the i -th column from G . Then we have the following theorem.

Theorem 3.1.

$$P := \left\{ \frac{x_1^{k_1} \dots x_t^{k_t}}{\det G_1(x)} dx_1 \mid (k_1, \dots, k_t) \in B(A_t), 0 \leq \sum_{i=1}^t a_i k_i \leq 2g - 2 \right\}$$

is a basis of $\Gamma(C, \Omega_C^1)$ over \mathbb{C} , where $\det G_1(x)$ denotes $\det G_1(X = x)$.

We order the elements of P in the ascending order with respect to the order at ∞ and write $\{du_1, \dots, du_g\}$.

In order to prove Theorem 3.1, we need some lemmas.

Lemma 3.1. *If $\det G_i(p) \neq 0$ for $p = (p_1, \dots, p_t) \in C^{\text{aff}}$ and $1 \leq i \leq t$, then $v_p(x_i - p_i) = 1$.*

Proof. Without loss of generality, we assume $i = 1$. Suppose $v_p(x_1 - p_1) \geq 2$. Then there exists k ($2 \leq k \leq t$) such that $v_p(x_k - p_k) = 1$. In fact, if $v_p(x_k - p_k) \geq 2$ for any k , then $v_p(f) \geq 2$ or $v_p(f) = 0$ for any $f \in R$. Then $v_p(g) \geq 2$ or $v_p(g) = 0$ for any $g \in R_p$, where R_p is the localization of R at p . This contradicts that R_p is a discrete valuation ring.

There exist $\{\gamma_{ij}, \delta_{j_1, \dots, j_t}^{(i)}\} \in \mathbb{C}$ such that for $2 \leq i \leq t$

$$F_i(X_1, \dots, X_t) = \sum_{j=1}^t \gamma_{ij}(X_j - p_j) + \sum_{j_1 + \dots + j_t \geq 2} \delta_{j_1, \dots, j_t}^{(i)}(X_1 - p_1)^{j_1} \cdots (X_t - p_t)^{j_t},$$

where $\gamma_{ij} = (\partial F_i / \partial X_j)(p)$. Since $F_i(x_1, \dots, x_t) = 0$ and $v_p(x_1 - p_1) \geq 2$, we have $v_p(\sum_{j=2}^t \gamma_{ij}(x_j - p_j)) = v_p((x_k - p_k)(\sum_{j=2}^t \gamma_{ij}(x_j - p_j)/(x_k - p_k))) \geq 2$. Since $v_p(x_k - p_k) = 1$, we have $\sum_{j=2}^t \gamma_{ij} b_j = 0$, where $b_j = ((x_j - p_j)/(x_k - p_k))(p)$. Therefore we obtain

$$G_1(p) \begin{pmatrix} b_2 \\ \cdot \\ \cdot \\ b_t \end{pmatrix} = \begin{pmatrix} 0 \\ \cdot \\ \cdot \\ 0 \end{pmatrix}.$$

Since $b_k = 1 (\neq 0)$, we have $\det G_1(p) = 0$. This contradicts the assumption of Lemma 3.1. Therefore we obtain $v_p(x_1 - p_1) = 1$. □

Lemma 3.2. (i) *As an element of K , we have $\det G_1(x) \neq 0$.*
 (ii) $\text{div}(dx_1 / \det G_1(x)) = (2g - 2)\infty$.

Proof. Since the differential $d(F_i(x_1, \dots, x_t)) = 0$ for any i , we have

$$G(x) \begin{pmatrix} dx_1 \\ \cdot \\ \cdot \\ dx_t \end{pmatrix} = \begin{pmatrix} 0 \\ \cdot \\ \cdot \\ 0 \end{pmatrix}.$$

By multiplying some elementary matrices on the left, the above equation becomes

$$\begin{pmatrix} w_2 & z_{22} & z_{23} & \cdots & z_{2t} \\ w_3 & 0 & z_{33} & \cdots & z_{3t} \\ & & \cdots & & \\ w_t & 0 & \cdots & & z_{tt} \end{pmatrix} \begin{pmatrix} dx_1 \\ \cdot \\ \cdot \\ dx_t \end{pmatrix} = \begin{pmatrix} 0 \\ \cdot \\ \cdot \\ 0 \end{pmatrix}.$$

Since C^{aff} is non-singular, for any $p \in C^{\text{aff}}$ there exists i such that $\det G_i(p) \neq 0$. Therefore we have $w_t \neq 0$ or $z_{tt} \neq 0$ as elements of K . Since $v_\infty(x_j) = -a_j$, we have $x_j \notin \mathbb{C}$, therefore $dx_j \neq 0$ for any j . Since $w_t dx_1 = z_{tt} dx_t$, we have $w_t \neq 0$ and $z_{tt} \neq 0$. Therefore, by multiplying some elementary matrices on the left, the above equation becomes

$$\begin{pmatrix} w'_2 & z_{22} & z_{23} & \cdots & 0 \\ w'_3 & 0 & z_{33} & \cdots & 0 \\ & & \cdots & & \\ w_t & 0 & \cdots & & z_{tt} \end{pmatrix} \begin{pmatrix} dx_1 \\ \cdot \\ \cdot \\ dx_t \end{pmatrix} = \begin{pmatrix} 0 \\ \cdot \\ \cdot \\ 0 \end{pmatrix}.$$

Similarly we obtain

$$\begin{pmatrix} w''_2 & z_{22} & 0 & \cdots & 0 \\ w''_3 & 0 & z_{33} & \cdots & 0 \\ & & \cdots & & \\ w''_t & 0 & \cdots & & z_{tt} \end{pmatrix} \begin{pmatrix} dx_1 \\ \cdot \\ \cdot \\ dx_t \end{pmatrix} = \begin{pmatrix} 0 \\ \cdot \\ \cdot \\ 0 \end{pmatrix},$$

where $w''_2, \dots, w''_t, z_{22}, \dots, z_{tt} \in K$ are non-zero. Therefore we obtain $\det G_1(x) = \pm z_{22} \cdots z_{tt} \neq 0$, which complete the proof of (i).

Next we prove that the 1-form $dx_1/\det G_1(x)$ is both holomorphic and non-vanishing on C^{aff} . When $\det G_1(p) \neq 0$ for $p \in C^{\text{aff}}$, from Lemma 3.1, $dx_1/\det G_1(x)$ is both holomorphic and non-vanishing at p . Suppose $\det G_1(p) = 0$ for $p \in C^{\text{aff}}$. Since C^{aff} is non-singular, there exists i ($2 \leq i \leq t$) such that $\det G_i(p) \neq 0$. Since $w'_i dx_1 + z_{ii} dx_i = 0$, we have $w'_i z_{22} \cdots \widehat{z_{ii}} \cdots z_{tt} dx_1 + z_{22} \cdots z_{tt} dx_i = 0$, where $\widehat{z_{ii}}$ denotes to remove z_{ii} . Therefore we obtain

$$(-1)^{i-2} \det G_i(x) dx_1 + \det G_1(x) dx_i = 0.$$

Since $\det G_1(x) \neq 0$ and $\det G_i(x) \neq 0$, we have

$$\frac{dx_1}{\det G_1(x)} = (-1)^{i-1} \frac{dx_i}{\det G_i(x)}.$$

Therefore, from $\det G_i(p) \neq 0$ and Lemma 3.1, $dx_1/\det G_1(x)$ is holomorphic and non-vanishing at p . On the other hand, by Riemann–Roch’s theorem, we have $\deg \operatorname{div}(dx_1/\det G_1(x)) = 2g - 2$, which complete the proof of (ii). □

Proof of Theorem 3.1. From Lemma 3.2 and Proposition 2.1 (i), we have $P \subset \Gamma(C, \Omega_C^1)$ and the elements of P are linearly independent. Since $\dim_{\mathbb{C}} \Gamma(C, \Omega_C^1) = g$, it is sufficient to prove $\#P = g$. It is well-known that there are g gap values at ∞ from 0 to $2g - 1$. Since $\dim_{\mathbb{C}} L((2g - 1)v_{\infty}) = \dim_{\mathbb{C}} L((2g - 2)v_{\infty}) = g$ (Riemann–Roch’s theorem), $2g - 1$ is a gap value at ∞ . Therefore, from Proposition 2.1 (i) and Proposition 2.2 (i), we have $\#\{(k_1, \dots, k_t) \in B(A_t) \mid 0 \leq \sum_{i=1}^t a_i k_i \leq 2g - 2\} = g$, which complete the proof of Theorem 3.1.

4. Second kind differentials for telescopic curves

In this section we construct dr_i for a telescopic (a_1, \dots, a_t) -curve C . For $2 \leq i \leq t$ and $1 \leq j \leq t$, let

$$h_{ij} = \frac{F_i(Y_1, \dots, Y_{j-1}, X_j, X_{j+1}, \dots, X_t) - F_i(Y_1, \dots, Y_{j-1}, Y_j, X_{j+1}, \dots, X_t)}{X_j - Y_j}$$

and

$$H = \begin{pmatrix} h_{22} & \cdots & h_{2t} \\ \dots\dots\dots\dots\dots\dots \\ h_{t2} & \cdots & h_{tt} \end{pmatrix}.$$

We consider the 1-form

$$\Omega(x, y) := \frac{\det H(x, y)}{(x_1 - y_1) \det G_1(x)} dx_1$$

and the bilinear form (cf. [15], p.181, 2.4)

$$(4) \quad \hat{\omega}(x, y) := d_y \Omega(x, y) + \sum c_{i_1, \dots, i_t; j_1, \dots, j_t} \frac{x_1^{i_1} \cdots x_t^{i_t} y_1^{j_1} \cdots y_t^{j_t}}{\det G_1(x) \det G_1(y)} dx_1 dy_1$$

on $C \times C$, where $x = (x_1, \dots, x_t)$, $y = (y_1, \dots, y_t)$, $c_{i_1, \dots, i_t; j_1, \dots, j_t} \in \mathbb{C}$, $(i_1, \dots, i_t) \in B(A_t)$ satisfying $0 \leq \sum_{k=1}^t a_k i_k \leq 2g - 2$, and $(j_1, \dots, j_t) \in B(A_t)$.

We take a basis $\{\alpha_i, \beta_i\}_{i=1}^g$ of the homology group $H_1(C, \mathbb{Z})$ such that their intersection numbers are $\alpha_i \circ \alpha_j = \beta_i \circ \beta_j = 0$ and $\alpha_i \circ \beta_j = \delta_{ij}$.

DEFINITION 4.1 (cf. [15], p.181, 2.4). Let $\Delta = \{(p, p) \mid p \in C\}$. A meromorphic symmetric bilinear form $\omega(x, y)$ on $C \times C$ is called a normalized fundamental form if the following conditions are satisfied.

(i) $\omega(x, y)$ is holomorphic except Δ where it has a double pole. For $p \in C$ take a local coordinate s around p . Then the expansion in $s(x)$ at $s(y)$ is of the form

$$\omega(x, y) = \left(\frac{1}{(s(x) - s(y))^2} + \text{regular} \right) ds(x) ds(y).$$

(ii) $\int_{\alpha_i} \omega = 0$ for any i , where the integration is with respect to any one of the variables.

Normalized fundamental form exists and unique (cf. [15] p.182). Then we have the following theorem.

Theorem 4.1. (i) *There exists a set of $c_{i_1, \dots, i_t; j_1, \dots, j_t}$ such that $\hat{\omega}(x, y) = \hat{\omega}(y, x)$, non-zero $c_{i_1, \dots, i_t; j_1, \dots, j_t}$ is a homogeneous polynomial of $\{\lambda_{i_1, \dots, i_t}^{(i)}\}$ of degree*

$$2 \sum_{k=2}^t \frac{d_{k-1}}{d_k} a_k - \sum_{k=1}^t (i_k + j_k + 2) a_k,$$

and $c_{i_1, \dots, i_t; j_1, \dots, j_t} = 0$ if $2 \sum_{k=2}^t (d_{k-1}/d_k) a_k - \sum_{k=1}^t (i_k + j_k + 2) a_k < 0$.

For a set of $c_{i_1, \dots, i_t; j_1, \dots, j_t}$ such that $\hat{\omega}(x, y) = \hat{\omega}(y, x)$, we have the following properties.

(ii) *The bilinear form $\hat{\omega}$ satisfies the condition (i) of Definition 4.1.*

(iii) *For $du_i := (x_1^{k_{i1}} \cdots x_t^{k_{it}} / \det G_1(x)) dx_1$, we define*

$$dr_i = \sum_{j_1, \dots, j_t} c_{k_{i1}, \dots, k_{it}; j_1, \dots, j_t} \frac{y_1^{j_1} \cdots y_t^{j_t}}{\det G_1(y)} dy_1.$$

Then dr_i is a second kind differential for any i , and the set $\{du_i, dr_i\}_{i=1}^g$ is a symplectic basis of $H^1(C, \mathbb{C})$.

Let B be the set of branch points for the map $x_1: C \rightarrow \mathbb{P}^1, (x_1, \dots, x_t) \rightarrow [x_1 : 1]$ (cf. [17], p.24, Example 2.2). Since the ramification index of the map x_1 at ∞ is a_1 , we have $\deg x_1 = a_1$ (cf. [17], p.28, Proposition 2.6). For $p \in C$ we set $x_1^{-1}(x_1(p)) = \{p^{(0)}, p^{(1)}, \dots, p^{(a_1-1)}\}$ with $p = p^{(0)}$, where the same $p^{(i)}$ is listed according to its ramification index.

Lemma 4.1. *Let U be a domain in \mathbb{C} , $f(z_1, z_2)$ a holomorphic function on $U \times U$, and $g(z) = f(z, z)$. If $g \equiv 0$ on U , then there exists a holomorphic function $h(z_1, z_2)$ on $U \times U$ such that $f(z_1, z_2) = (z_1 - z_2)h(z_1, z_2)$.*

Proof. Let $h(z_1, z_2) = f(z_1, z_2)/(z_1 - z_2)$. Given z_1 , $h(z_1, \cdot)$ has a singularity only at z_1 , where its singularity is removable. Therefore $h(z_1, \cdot)$ is holomorphic on U . Similarly $h(\cdot, z_2)$ is holomorphic on U . Therefore h is holomorphic on $U \times U$. \square

Lemma 4.2. *The 1-form $\Omega(x, y)$ is holomorphic except $\Delta \cup \{(p^{(i)}, p) \mid i \neq 0, p \in B \text{ or } p^{(i)} \in B\} \cup C \times \{\infty\} \cup \{\infty\} \times C$.*

Proof. Since $dx_1/\det G_1(x)$ is holomorphic on C (cf. Lemma 3.2), $\Omega(x, y)$ is holomorphic except $\Delta \cup \{(p^{(i)}, p) \mid p \in C, i \neq 0\} \cup C \times \{\infty\} \cup \{\infty\} \times C$. We prove that $\Omega(x, y)$ is holomorphic on $\{(p^{(i)}, p) \mid i \neq 0, p \notin B, p^{(i)} \notin B\}$. We have

$$(5) \quad F_i(X_1, \dots, X_t) = \sum_{j=1}^t h_{ij} \cdot (X_j - Y_j) + F_i(Y_1, \dots, Y_t).$$

Set $X = x$ and $Y = y$, then we have

$$\sum_{j=1}^t h_{ij}(x, y) \cdot (x_j - y_j) = 0.$$

Take $(p^{(i)}, p) \in C \times C$ such that $i \neq 0, p \notin B$, and $p^{(i)} \notin B$, then we have

$$\begin{pmatrix} h_{21} & \cdots & h_{2t} \\ \dots\dots\dots \\ h_{t1} & \cdots & h_{tt} \end{pmatrix}_{X=p^{(i)}, Y=p} \begin{pmatrix} p_1^{(i)} - p_1 \\ \cdot \\ p_t^{(i)} - p_t \end{pmatrix} = \begin{pmatrix} 0 \\ \cdot \\ 0 \end{pmatrix}.$$

Since $p_1^{(i)} - p_1 = 0$, we have

$$H(p^{(i)}, p) \begin{pmatrix} p_2^{(i)} - p_2 \\ \cdot \\ p_t^{(i)} - p_t \end{pmatrix} = \begin{pmatrix} 0 \\ \cdot \\ 0 \end{pmatrix}.$$

Since $(p_2^{(i)} - p_2, \dots, p_t^{(i)} - p_t) \neq (0, \dots, 0)$, we have $\det H(p^{(i)}, p) = 0$. Since $p \notin B$ and $p^{(i)} \notin B$, we can take (x_1, y_1) as a local coordinate around $(p^{(i)}, p)$. Therefore, from Lemma 4.1, there exists a holomorphic function $h(x_1, y_1)$ around $(p^{(i)}, p)$ such that $\det H(x, y) = (x_1 - y_1)h(x_1, y_1)$. Therefore $\Omega(x, y)$ is holomorphic at $(p^{(i)}, p)$. \square

Lemma 4.3. *Let $p \notin B$, s a local coordinate around p . Then the expansion of $\Omega(x, y)$ in $s(y)$ at $s(x)$ is of the form*

$$\Omega(x, y) = \left(\frac{-1}{s(y) - s(x)} + \text{regular} \right) ds(x).$$

Proof. Set $Y = y$ in (5), then we have

$$F_i(X_1, \dots, X_t) = \sum_{j=1}^t h_{ij}(X, y) \cdot (X_j - y_j).$$

Therefore we obtain

$$\frac{\partial F_i}{\partial X_k}(x_1, \dots, x_t) = \sum_{j=1}^t \frac{\partial h_{ij}}{\partial X_k}(x, y) \cdot (x_j - y_j) + h_{ik}(x, y).$$

Set $x = y$, then we have

$$\frac{\partial F_i}{\partial X_k}(x_1, \dots, x_t) = h_{ik}(x, x).$$

Therefore we obtain $\det G_1(x) = \det H(x, x)$. On the other hand, since $p \notin B$, we can take (x_1, y_1) as a local coordinate around (p, p) . Since $p \notin B$, we have $\det G_1(p) \neq 0$. In fact, if $\det G_1(p) = 0$, then $dx_1/\det G_1(x)$ is not holomorphic at p , which contradicts Lemma 3.2 (ii). Therefore $\det H(x, y)/\det G_1(x)$ is holomorphic at (p, p) . Therefore, from Lemma 4.1, there exists a holomorphic function $\tilde{h}(x_1, y_1)$ around (p, p) such that $\det H(x, y)/\det G_1(x) = 1 + (x_1 - y_1)\tilde{h}(x_1, y_1)$. Therefore we obtain Lemma 4.3. \square

Lemma 4.4. *When we express*

$$\det H(X, Y) = \sum \epsilon_{m_1, \dots, m_t, n_1, \dots, n_t} X_1^{m_1} \cdots X_t^{m_t} Y_1^{n_1} \cdots Y_t^{n_t},$$

we have $\sum_{k=1}^t a_k(m_k + n_k) \leq \sum_{k=2}^t a_k((d_{k-1}/d_k) - 1)$.

Proof. When we express

$$F_i(X_1, \dots, X_t) = \sum_{k=0}^m \tilde{F}_{ik}^{(j)}(X_1, \dots, X_{j-1}, X_{j+1}, \dots, X_t) X_j^k,$$

we have $h_{ij} = \sum_{k=1}^m \tilde{F}_{ik}^{(j)}(Y_1, \dots, Y_{j-1}, X_{j+1}, \dots, X_t) \sum_{l=0}^{k-1} X_j^l Y_j^{k-l-1}$. Assign degrees as $\deg Y_k = a_k$, then h_{ij} is a homogeneous polynomial of $\{\lambda_{j_1, \dots, j_t}^{(i)}, X_k, Y_k\}$ of degree $a_i d_{i-1}/d_i - a_j$. Therefore we obtain Lemma 4.4. \square

Lemma 4.5. *The meromorphic bilinear form $d_y \Omega(x, y)$ is holomorphic except $\Delta \cup \{(p^{(i)}, p) \mid i \neq 0, p \in B \text{ or } p^{(i)} \in B\} \cup C \times \{\infty\}$.*

Proof. It is sufficient to prove that $d_y \Omega(x, y)$ is holomorphic at (∞, y) , $y \neq \infty$. From Lemma 4.4, with respect to x , we obtain

$$v_\infty(\det H(x, y)) \geq - \sum_{k=2}^t a_k((d_{k-1}/d_k) - 1).$$

If $v_\infty(\det H(x, y)) > -\sum_{k=2}^t a_k((d_{k-1}/d_k) - 1)$, then from Lemma 3.2 (ii) and Proposition 2.3 (iii) we obtain $v_\infty(\Omega(x, y)) \geq 0$. Therefore $d_y\Omega(x, y)$ is holomorphic at (∞, y) . If $v_\infty(\det H(x, y)) = -\sum_{k=2}^t a_k((d_{k-1}/d_k) - 1)$, then $v_\infty(\Omega(x, y)) = -1$. Let s be a local coordinate around ∞ , then from Lemma 4.4 there exists a constant e (which does not depend on y) such that

$$\Omega(x, y) = \left(\frac{e}{s} + \text{regular}\right) ds.$$

Therefore $d_y\Omega(x, y)$ is holomorphic at (∞, y) , $y \neq \infty$. □

Lemma 4.6. *Let ω be the normalized fundamental form. Then there exist second kind differentials $d\hat{r}_i$ ($1 \leq i \leq g$) which are holomorphic except $\{\infty\}$ and satisfy the equation*

$$\omega(x, y) - d_y\Omega(x, y) = \sum_{i=1}^g du_i(x) d\hat{r}_i(y).$$

Proof. Set $B_2 = \{(p^{(i)}, p) \mid p \in B \setminus \{\infty\} \text{ or } p^{(i)} \in B \setminus \{\infty\}\}$ in the proof of [15] Lemma 5, then proof of Lemma 4.6 is similar to that of [15] Lemma 5. □

Lemma 4.7. *Let Q be the linear space consisting of meromorphic differentials on C which are singular only at ∞ and*

$$S = \{(x_1^{i_1} \cdots x_t^{i_t} / \det G_1(x)) dx_1 \mid (i_1, \dots, i_t) \in B(A_t)\}.$$

Then S is a basis of Q .

Proof. For $\eta \in Q$ we consider the meromorphic function $\eta/(dx_1/\det G_1(x))$. From Lemma 3.2 (ii), it may have a pole only at ∞ . From Proposition 2.1 (i) and Proposition 2.2 (i), $\eta/(dx_1/\det G_1(x))$ is a linear combination of $x_1^{i_1} \cdots x_t^{i_t}$ with $(i_1, \dots, i_t) \in B(A_t)$ and the elements of S are linearly independent. □

Proof of Theorem 4.1 (i). We have

$$\begin{aligned} d_y\Omega(x, y) &= \frac{\left\{ \sum_{k=1}^t (-1)^{k+1} (x_1 - y_1) (\partial \det H / \partial Y_k)(x, y) \det G_k(y) \right\} + \det G_1(y) \det H(x, y)}{(x_1 - y_1)^2 \det G_1(x) \det G_1(y)} dx_1 dy_1. \end{aligned}$$

Then, $\det G_k$, $\det H$, and $(\partial \det H / \partial Y_k)$ are homogeneous polynomials of $\{\lambda_{j_1, \dots, j_t}^{(i)}, X_j, Y_j\}$ of degree $\sum_{i=2}^t (d_{i-1}/d_i) a_i - \sum_{i \neq k} a_i$, $\sum_{i=2}^t ((d_{i-1}/d_i) - 1) a_i$, and $\{\sum_{i=2}^t ((d_{i-1}/d_i) - 1) a_i\} - a_k$, respectively. Let us write

$$d_y\Omega(x, y) = \frac{\sum q_{i_1, \dots, i_t; j_1, \dots, j_t} x_1^{i_1} \cdots x_t^{i_t} y_1^{j_1} \cdots y_t^{j_t}}{(x_1 - y_1)^2 \det G_1(x) \det G_1(y)} dx_1 dy_1,$$

where $(i_1, \dots, i_t), (j_1, \dots, j_t) \in B(A_t)$, and $q_{i_1, \dots, i_t; j_1, \dots, j_t} \in \mathbb{C}$. Then $q_{i_1, \dots, i_t; j_1, \dots, j_t} \in \mathbb{Z}[\{\lambda_{i_1, \dots, i_t}^{(i)}\}]$ and $q_{i_1, \dots, i_t; j_1, \dots, j_t}$ is homogeneous of degree $2 \sum_{k=2}^t ((d_{k-1}/d_k) - 1)a_k - \sum_{k=1}^t (i_k + j_k)a_k$. Note that if $(m_1, \dots, m_t) \in B(A_t)$, then $(m_1 + m, m_2, \dots, m_t) \in B(A_t)$ for $m \in \mathbb{N}$. Therefore we obtain

$$\begin{aligned} & \sum c_{i_1, \dots, i_t; j_1, \dots, j_t} \frac{x_1^{i_1} \cdots x_t^{i_t} y_1^{j_1} \cdots y_t^{j_t}}{\det G_1(x) \det G_1(y)} \\ &= \frac{\sum (c_{i_1-2, \dots, i_t; j_1, \dots, j_t} - 2c_{i_1-1, \dots, i_t; j_1-1, \dots, j_t} + c_{i_1, \dots, i_t; j_1-2, \dots, j_t}) x_1^{i_1} \cdots x_t^{i_t} y_1^{j_1} \cdots y_t^{j_t}}{(x_1 - y_1)^2 \det G_1(x) \det G_1(y)}, \end{aligned}$$

where $(i_1, \dots, i_t), (j_1, \dots, j_t) \in B(A_t)$. Therefore $\hat{\omega}(x, y) = \hat{\omega}(y, x)$ is equivalent to

$$\begin{aligned} & c_{i_1-2, \dots, i_t; j_1, \dots, j_t} - 2c_{i_1-1, \dots, i_t; j_1-1, \dots, j_t} + c_{i_1, \dots, i_t; j_1-2, \dots, j_t} - c_{j_1-2, \dots, j_t; i_1, \dots, i_t} \\ &+ 2c_{j_1-1, \dots, j_t; i_1-1, \dots, i_t} - c_{j_1, \dots, j_t; i_1-2, \dots, i_t} \\ &= q_{j_1, \dots, j_t; i_1, \dots, i_t} - q_{i_1, \dots, i_t; j_1, \dots, j_t}. \end{aligned}$$

By Lemma 4.6, 4.7, the system of the above linear equations has a solution. Moreover it has a solution such that each $c_{i_1, \dots, i_t; j_1, \dots, j_t}$ is a linear combination of $q_{i'_1, \dots, i'_t; j'_1, \dots, j'_t}$ satisfying $i'_1 + j'_1 = i_1 + j_1 + 2$, $(i'_k, j'_k) = (i_k, j_k)$ or $(i'_k, j'_k) = (j_k, i_k)$ for $k = 2, \dots, t$. In particular one can take $c_{i_1, \dots, i_t; j_1, \dots, j_t}$ such that $c_{i_1, \dots, i_t; j_1, \dots, j_t} = 0$ if $2 \sum_{k=2}^t (d_{k-1}/d_k)a_k - \sum_{k=1}^t (i_k + j_k + 2)a_k < 0$ and

$$\text{deg } c_{i_1, \dots, i_t; j_1, \dots, j_t} = 2 \sum_{k=2}^t \frac{d_{k-1}}{d_k} a_k - \sum_{k=1}^t (i_k + j_k + 2)a_k$$

if $c_{i_1, \dots, i_t; j_1, \dots, j_t} \neq 0$. □

Proof of Theorem 4.1 (ii). From Lemma 4.6, $d_y \Omega(x, y)$ is holomorphic except $\Delta \cup C \times \{\infty\}$ and so is $\hat{\omega}$. Since $\hat{\omega}(x, y) = \hat{\omega}(y, x)$, $\hat{\omega}$ is holomorphic except Δ . From the definition of dr_i , we obtain

$$\hat{\omega} - \omega = \sum_{i=1}^g du_i(x)(dr_i(y) - d\hat{r}_i(y)).$$

On the other hand $\hat{\omega} - \omega$ is holomorphic except Δ and $\sum_{i=1}^g du_i(x)(dr_i(y) - d\hat{r}_i(y))$ is holomorphic except $C \times \{\infty\}$. Therefore $\hat{\omega} - \omega$ is holomorphic except $\{\infty\} \times \{\infty\}$. Therefore $\hat{\omega} - \omega$ and $dr_i - d\hat{r}_i$ are holomorphic on $C \times C$ and C respectively, which complete the proof of Theorem 4.1 (ii). □

Proof of Theorem 4.1 (iii). The 1-form dr_i is a second kind differential. In fact $dr_i - d\hat{r}_i$ is holomorphic 1-form as is just proved in the proof of Theorem 4.1 (ii) and $d\hat{r}_i$ is a second kind differential from Lemma 4.6. Proof of Theorem 4.1 (iii) is similar to the case of the (n, s) -curves (cf. [15] Lemmas 7, 8, Proposition 3). □

5. Sigma functions for telescopic curves

In this section we construct the sigma function for a telescopic (a_1, \dots, a_t) -curve C . First we take the following data.

1. A basis $\{\alpha_i, \beta_i\}_{i=1}^g$ of the homology group $H_1(C, \mathbb{Z})$ such that their intersection numbers are $\alpha_i \circ \alpha_j = \beta_i \circ \beta_j = 0$ and $\alpha_i \circ \beta_j = \delta_{ij}$.
2. The symplectic basis $\{du_i, dr_i\}_{i=1}^g$ of the first cohomology group $H^1(C, \mathbb{C})$ constructed in Sections 3 and 4.

We define the period matrices by

$$2\omega_1 = \left(\int_{\alpha_j} du_i \right), \quad 2\omega_2 = \left(\int_{\beta_j} du_i \right), \quad -2\eta_1 = \left(\int_{\alpha_j} dr_i \right), \quad -2\eta_2 = \left(\int_{\beta_j} dr_i \right).$$

Then ω_1 is invertible. Set $\tau = \omega_1^{-1}\omega_2$, then τ is symmetric and $\text{Im } \tau > 0$. By the Riemann’s bilinear relation

$$2\pi i \eta \circ \eta' = \sum_{i=1}^g \left(\int_{\alpha_i} \eta \int_{\beta_i} \eta' - \int_{\alpha_i} \eta' \int_{\beta_i} \eta \right),$$

the matrix

$$M := \begin{pmatrix} \omega_1 & \omega_2 \\ \eta_1 & \eta_2 \end{pmatrix}$$

satisfies

$$M \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix} {}^t M = -\frac{\pi \sqrt{-1}}{2} \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix},$$

where I_g denotes the unit matrix of degree g . Since $\eta_1 \omega_1^{-1}$ is symmetric (cf. [15] Lemma 8), we obtain the following proposition.

Proposition 5.1 (generalized Legendre relation).

$${}^t M \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix} M = -\frac{\pi \sqrt{-1}}{2} \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix}.$$

Let $\delta = \tau \delta' + \delta''$ be the Riemann’s constant of C with respect to our choice $(\infty, \{\alpha_i, \beta_i\}_{i=1}^g)$. Since the divisor of the holomorphic 1-form du_g is $(2g-2)\infty$, the Riemann’s constant δ becomes a half period. Then the sigma function $\sigma(u)$ associated with C is defined as follows.

DEFINITION 5.1 (Sigma function). For $u \in \mathbb{C}^s$

$$\begin{aligned} \sigma(u) &= \sigma(u; M) = c \cdot \exp\left(\frac{1}{2} {}^t u \eta_1 \omega_1^{-1} u\right) \theta \begin{bmatrix} \delta' \\ \delta'' \end{bmatrix} ((2\omega_1)^{-1} u, \tau) \\ &= c \cdot \exp\left(\frac{1}{2} {}^t u \eta_1 \omega_1^{-1} u\right) \\ &\quad \times \sum_{n \in \mathbb{Z}^s} \exp\{\pi \sqrt{-1} {}^t (n + \delta') \tau (n + \delta') + 2\pi \sqrt{-1} {}^t (n + \delta') ((2\omega_1)^{-1} u + \delta'')\}, \end{aligned}$$

where c is a constant.

By Proposition 5.1 we obtain the following proposition.

Proposition 5.2. For any $m_1, m_2 \in \mathbb{Z}^s$ and $u \in \mathbb{C}^s$, we have

$$\begin{aligned} \sigma(u + 2\omega_1 m_1 + 2\omega_2 m_2) / \sigma(u) &= \exp(\pi \sqrt{-1} ({}^t m_1 m_2 + 2 {}^t \delta' m_1 - 2 {}^t \delta'' m_2)) \\ &\quad \times \exp({}^t (2\eta_1 m_1 + 2\eta_2 m_2)(u + \omega_1 m_1 + \omega_2 m_2)). \end{aligned}$$

REMARK. In this paper we have constructed sigma functions explicitly for telescopic curves. On the other hand Nakayashiki [15] showed that the first term of the series expansion around the origin of the sigma function for an (n, s) -curve becomes Schur function corresponding to the partition determined from the gap sequence at infinity and the expansion coefficients are homogeneous polynomials of the coefficients of the defining equation of the curve. One will be able to extend these results to telescopic curves.

A. Proof of Proposition 2.1

Lemma A.1. $V(A_t) + \mathbb{N}^t = \mathbb{N}^t \setminus B(A_t)$.

Proof. If $M \notin B(A_t)$ and $N \in \mathbb{N}^t$, then $M + N \notin B(A_t)$. Therefore we have $V(A_t) + \mathbb{N}^t \subset \mathbb{N}^t \setminus B(A_t)$. Suppose $V(A_t) + \mathbb{N}^t \subsetneq \mathbb{N}^t \setminus B(A_t)$. Take $M_1 \in \mathbb{N}^t \setminus B(A_t)$ satisfying $M_1 \notin V(A_t) + \mathbb{N}^t$. Since $M_1 \notin V(A_t)$ and $M_1 \notin B(A_t)$, there exist $M_2 \in \mathbb{N}^t \setminus B(A_t)$ and $(0, \dots, 0) \neq N_1 \in \mathbb{N}^t$ such that $M_1 = M_2 + N_1$. Since $M_1 \notin V(A_t) + \mathbb{N}^t$, we have $M_2 \notin V(A_t) + \mathbb{N}^t$. Similarly, for the element $M_i \in \mathbb{N}^t \setminus B(A_t)$ satisfying $M_i \notin V(A_t) + \mathbb{N}^t$, there exist M_{i+1} and N_i such that $M_{i+1} \in \mathbb{N}^t \setminus B(A_t)$, $M_{i+1} \notin V(A_t) + \mathbb{N}^t$, $(0, \dots, 0) \neq N_i \in \mathbb{N}^t$, and $M_i = M_{i+1} + N_i$. Therefore there exists a infinite sequence $\Psi(M_1) > \Psi(M_2) > \dots > \Psi(M_i) > \dots$. This is contradiction. \square

Proof of Proposition 2.1 (i). From (2) it is sufficient to prove

$$\text{Span}\{X^N \mid N \in B(A_t)\} + (\{F_M \mid M \in V(A_t)\}) = \mathbb{C}[X].$$

We prove that for any $T \in \mathbb{N}^t$

$$X^T \in \text{Span}\{X^N \mid N \in B(A_t)\} + (\{F_M \mid M \in V(A_t)\})$$

by transfinite induction with respect to the well-order $<$ in \mathbb{N}^t . The statement is correct for the minimal element $T = (0, \dots, 0)$. Suppose that it is correct for any $U \in \mathbb{N}^t$ satisfying $U < T$. Since it is correct for $T \in B(A_t)$, we assume $T \notin B(A_t)$. From Lemma A.1, there exist $M \in V(A_t)$ and $Z \in \mathbb{N}^t$ such that $T = M + Z$. Then we have $X^T = X^M X^Z = (X^M - F_M)X^Z + F_M X^Z$. For any monomial X^U in $(X^M - F_M)X^Z$, we have $U < T$. Therefore, by the assumption of transfinite induction, the statement is correct for $T \notin B(A_t)$. \square

We define the function $o: R \rightarrow \mathbb{N} \cup \{-\infty\}$ by

$$o(f) = \begin{cases} -\infty & \text{for } f = 0, \\ \max\{\Psi(N) \mid \lambda_N \neq 0\} & \text{for } f \neq 0, \end{cases}$$

where for $f \neq 0$ we express $f = \sum_N \lambda_N x^N$ with $\lambda_N \in \mathbb{C}$ and $N \in B(A_t)$.

Lemma A.2. $o(x^T) = \Psi(T)$ for any $T \in \mathbb{N}^t$.

Proof. We prove the statement by transfinite induction with respect to the well-order $<$ in \mathbb{N}^t . It is correct for the minimal element $T = (0, \dots, 0) \in \mathbb{N}^t$. Suppose that it is correct for any $U \in \mathbb{N}^t$ satisfying $U < T$. Since it is correct for $T \in B(A_t)$, we assume $T \notin B(A_t)$. From Lemma A.1, there exist $M \in V(A_t)$ and $Z \in \mathbb{N}^t$ such that $T = M + Z$. Then we have $X^T = X^M X^Z = (X^M - F_M)X^Z + F_M X^Z$. Since $X^M - F_M = X^L + \sum_N \lambda_N X^N$ from (1), we have $x^T = (x^L + \sum_N \lambda_N x^N)x^Z = x^{L+Z} + \sum_N \lambda_N x^{N+Z}$. Since $N + Z < L + Z < T$, by the assumption of transfinite induction, we have $o(x^{L+Z}) = \Psi(L + Z)$ and $o(x^{N+Z}) = \Psi(N + Z)$. Since $o(f + g) = \max\{o(f), o(g)\}$ for $f, g \in R$ satisfying $o(f) \neq o(g)$, we have $o(x^T) = o(x^{L+Z} + \sum_N \lambda_N x^{N+Z}) = o(x^{L+Z}) = \Psi(L + Z) = \Psi(T)$. \square

Lemma A.3. *The function o satisfies the following properties:*

- (i) $o(f) = -\infty$ if and only if $f = 0$,
- (ii) $o(fg) = o(f) + o(g)$ for any $f, g \in R$, where we define $-\infty + (-\infty) = a + (-\infty) = (-\infty) + a = -\infty$ for $a \in \mathbb{N}$,
- (iii) $o(f + g) \leq \max\{o(f), o(g)\}$,
- (iv) $o(R \setminus \{0\}) = \langle A_t \rangle$, in particular $\mathbb{N} \setminus o(R \setminus \{0\})$ is a finite set, and
- (v) $o(a) = 0$ for any $0 \neq a \in \mathbb{C}$.

Proof. (i), (iii), (v), and $o(R \setminus \{0\}) = \langle A_t \rangle$ are trivial. Since $\text{GCD}\{a_1, \dots, a_t\} = 1$, $\mathbb{N} \setminus \langle A_t \rangle$ is a finite set (cf. [16], Theorem 5). We prove (ii). If $f = 0$ or $g = 0$, then

$o(fg) = o(f) + o(g) = -\infty$. Suppose $f \neq 0$ and $g \neq 0$. Then we can express

$$f = \lambda_M x^M + \sum_T \lambda_T x^T \quad \text{and} \quad g = \tilde{\lambda}_N x^N + \sum_Z \tilde{\lambda}_Z x^Z,$$

where $\lambda_M, \lambda_T, \tilde{\lambda}_N, \tilde{\lambda}_Z \in \mathbb{C}$, $\lambda_M \neq 0$, $\tilde{\lambda}_N \neq 0$, $M, T, N, Z \in B(A_t)$, $\Psi(T) < \Psi(M)$, and $\Psi(Z) < \Psi(N)$. From Lemma A.2, we have $o(fg) = o(\lambda_M \tilde{\lambda}_N x^{M+N}) = \Psi(M + N) = \Psi(M) + \Psi(N) = o(f) + o(g)$. □

Proof of Proposition (ii). Take $f, g \in R$ satisfying $fg = 0$. Then, since $-\infty = o(fg) = o(f) + o(g)$, we have $o(f) = -\infty$ or $o(g) = -\infty$. Therefore we obtain $f = 0$ or $g = 0$. □

Lemma A.4. *$B \subset \mathbb{N}^t$ be a set such that the restriction map of $\Psi: \mathbb{N}^t \rightarrow \langle A_t \rangle$ on B is bijective. Then the set $\{x^M \mid M \in B\} \subset R$ is a basis of R over \mathbb{C} .*

Proof. Since $o(x^T) = \Psi(T)$ for $T \in \mathbb{N}^t$ and $o(f + g) = \max\{o(f), o(g)\}$ for $f, g \in R$ satisfying $o(f) \neq o(g)$, the elements of the set $\{x^M \mid M \in B\}$ are linearly independent. Since $R = \text{Span}\{x^N \mid N \in B(A_t)\}$, in order to prove $R = \text{Span}\{x^M \mid M \in B\}$, it is sufficient to prove $\text{Span}\{x^N \mid N \in B(A_t)\} \subset \text{Span}\{x^M \mid M \in B\}$. We prove $\text{Span}\{x^N \mid N \in B(A_t), \Psi(N) \leq m\} \subset \text{Span}\{x^M \mid M \in B, \Psi(M) \leq m\}$ for any $m \in \mathbb{N}$ by induction. For $m = 0$ the statement is trivial. Suppose that the statement is correct for any i with $0 \leq i \leq m - 1$. If $m \notin \langle A_t \rangle$, then since $\text{Span}\{x^M \mid M \in B, \Psi(M) \leq m\} = \text{Span}\{x^M \mid M \in B, \Psi(M) \leq m - 1\}$ and $\text{Span}\{x^N \mid N \in B(A_t), \Psi(N) \leq m\} = \text{Span}\{x^N \mid N \in B(A_t), \Psi(N) \leq m - 1\}$, the statement is correct. Suppose $m \in \langle A_t \rangle$. Take $T \in B$ satisfying $\Psi(T) = m$. If $T \in B(A_t)$, then since $\text{Span}\{x^M \mid M \in B, \Psi(M) \leq m\} = \text{Span}\{x^M \mid M \in B, \Psi(M) \leq m - 1\} \cup \mathbb{C}\{x^T\}$ and $\text{Span}\{x^N \mid N \in B(A_t), \Psi(N) \leq m\} = \text{Span}\{x^N \mid N \in B(A_t), \Psi(N) \leq m - 1\} \cup \mathbb{C}\{x^T\}$, the statement is correct. Suppose $T \notin B(A_t)$. Then we can express $x^T = \lambda_L x^L + \sum_N \lambda_N x^N$, where $0 \neq \lambda_L, \lambda_N \in \mathbb{C}$, $L, N \in B(A_t)$, $\Psi(L) = m$, and $\Psi(N) \leq m - 1$. Since $x^L = \lambda_L^{-1}(x^T - \sum_N \lambda_N x^N) \in \text{Span}\{x^N \mid N \in B(A_t), \Psi(N) \leq m - 1\} \cup \mathbb{C}\{x^T\} \subset \text{Span}\{x^M \mid M \in B, \Psi(M) \leq m - 1\} \cup \mathbb{C}\{x^T\} \subset \text{Span}\{x^M \mid M \in B, \Psi(M) \leq m\}$, we have $\text{Span}\{x^N \mid N \in B(A_t), \Psi(N) \leq m\} \subset \text{Span}\{x^M \mid M \in B, \Psi(M) \leq m\}$. □

Lemma A.5. *Given i , there exists a set $T_i \subset \mathbb{N}^{i-1} \times \{0\} \times \mathbb{N}^{t-i}$ such that $\#T_i = a_i$ and for the set $B_i := T_i + \{0\}^{i-1} \times \mathbb{N} \times \{0\}^{t-i}$ the restriction map of $\Psi: \mathbb{N}^t \rightarrow \langle A_t \rangle$ on B_i is bijective.*

Proof. Since $\text{GCD}\{a_1, \dots, a_t\} = 1$, the set $\{c \in a_1\mathbb{N} + \dots + a_{i-1}\mathbb{N} + a_{i+1}\mathbb{N} + \dots + a_t\mathbb{N} \mid c \equiv j \pmod{a_i}\}$ is not empty for any j with $0 \leq j \leq a_i - 1$. Let $c_j = \min\{c \in a_1\mathbb{N} + \dots + a_{i-1}\mathbb{N} + a_{i+1}\mathbb{N} + \dots + a_t\mathbb{N} \mid c \equiv j \pmod{a_i}\}$. Take $N_j \in \mathbb{N}^{i-1} \times \{0\} \times \mathbb{N}^{t-i}$ satisfying $\Psi(N_j) = c_j$. Let $T_i = \{N_j \mid 0 \leq j \leq a_i - 1\}$. Then T_i satisfies the conditions of Lemma A.5. □

Proof of Proposition 2.1 (iii). Since $o(x^T) = \Psi(T)$ for $T \in \mathbb{N}^t$ and $o(f + g) = \max\{o(f), o(g)\}$ for $f, g \in R$ satisfying $o(f) \neq o(g)$, the elements of the set $\{x^M \mid M \in \{0\}^{i-1} \times \mathbb{N} \times \{0\}^{t-i}\} \subset \mathbb{C}[x_i]$ are linearly independent. Therefore the extension of field $\mathbb{C}(x_i)/\mathbb{C}$ is a simple transcendental extension for any i . Next we prove $[K : \mathbb{C}(x_i)] \leq a_i$ for any i . From Lemma A.4 and Lemma A.5, we have $R = \mathbb{C}[x_1, \dots, x_t] = \text{Span}\{x^M \mid M \in T_i + \{0\}^{i-1} \times \mathbb{N} \times \{0\}^{t-i}\}$. Therefore $\mathbb{C}[x_1, \dots, x_t] = \mathbb{C}[x_i]f_0 + \dots + \mathbb{C}[x_i]f_{a_i-1}$, where $f_j = x^{N_j}$ (see the proof of Lemma A.5 for N_j). Since $f_0 = 1$, we obtain the finite extension of integral domain $\mathbb{C}(x_i) \subset \mathbb{C}(x_i)f_0 + \dots + \mathbb{C}(x_i)f_{a_i-1}$. Since $\mathbb{C}(x_i)$ is a field, $\mathbb{C}(x_i)f_0 + \dots + \mathbb{C}(x_i)f_{a_i-1}$ is also a field. Therefore we obtain $\mathbb{C}(x_i)f_0 + \dots + \mathbb{C}(x_i)f_{a_i-1} = K$ and $[K : \mathbb{C}(x_i)] \leq a_i$. \square

Proof of Proposition 2.1 (iv). We define the function $v_\infty : K \rightarrow \mathbb{Z} \cup \{\infty\}$ by

$$v_\infty(f) = \begin{cases} \infty & \text{for } f = 0, \\ -o(f_1) + o(f_2) & \text{for } f \neq 0, \end{cases}$$

where for $f \neq 0$ we express $f = f_1/f_2$ with $f_1, f_2 \in R$. The definition of v_∞ is well-defined. In fact, if $0 \neq f = f_1/f_2 = g_1/g_2$, then since $f_1g_2 = g_1f_2 \in R$, we have $o(f_1) + o(g_2) = o(f_1g_2) = o(g_1f_2) = o(g_1) + o(f_2)$. From Lemma A.3, one can check that the function v_∞ is a discrete valuation of K . From Lemma A.2, we obtain $v_\infty(x_i) = -a_i$. From [19] p.19 Theorem 1.4.11, we obtain $[K : \mathbb{C}(x_i)] = \deg(x_i)_\infty \geq \deg(a_i v_\infty) = a_i$. On the other hand, in the proof of Proposition 2.1 (iii), we proved $[K : \mathbb{C}(x_i)] \leq a_i$. Therefore we obtain $(x_i)_\infty = a_i v_\infty$. \square

B. Proof of Proposition 2.2

Proof of Proposition 2.2 (i). It is trivial that $R \subset \bigcup_{k=0}^\infty L(kv_\infty)$. On the other hand we have

$$\bigcup_{k=0}^\infty L(kv_\infty) \subset \bigcap_{v \neq v_\infty} \mathcal{O}_v \subset \bigcap_{p \in C^{\text{aff}}} \mathcal{O}_p = R,$$

where $\mathcal{O}_v = \{f \in K \mid v(f) \geq 0\}$ and $\mathcal{O}_p = \{f \in K \mid v_p(f) \geq 0\}$ (see Proposition 2.2 (ii) for v_p). \square

Proof of Proposition 2.2 (ii). It is trivial that the map ϕ is injective. We prove that the map ϕ is surjective. Let v be a discrete valuation such that $v \neq v_\infty$. Since $v(x_i) \geq 0$ for any i , we have $R \subset \mathcal{O}_v$. Let P be the maximal ideal of \mathcal{O}_v and $m := P \cap R$. Then we have

$$\mathbb{C} \hookrightarrow R/m \hookrightarrow \mathcal{O}_v/P.$$

Since $[\mathcal{O}_v/P : \mathbb{C}] = 1$, we have $\mathbb{C} \simeq R/m \simeq \mathcal{O}_v/P$. Therefore m is a maximal ideal. Let R_m be the localization of R with respect to m . Then R_m and \mathcal{O}_v are discrete val-

uation rings satisfying $R_m \subset \mathcal{O}_v$ and $P \cap R_m = mR_m$. Therefore, from [8] p.40 Theorem 6.1A, we obtain $R_m = \mathcal{O}_v$. Since there exists $p \in C^{\text{aff}}$ such that $\mathcal{O}_p = R_m$, we have $\mathcal{O}_p = \mathcal{O}_v$. Therefore we obtain $v_p = v$ and the map ϕ is surjective. \square

C. Proof of Proposition 2.3

Let $T(A_t) = B(A_t) \cap (\{0\} \times \mathbb{N}^{t-1})$.

- Lemma C.1.** (i) $T(A_t) = \{M(b_i) \in B(A_t) \mid i = 0, \dots, a_1 - 1\}$, where $b_i = \min\{b \in a_2\mathbb{N} + \dots + a_t\mathbb{N} \mid b \equiv i \pmod{a_1}\}$. In particular $\#T(A_t) = a_1$.
 (ii) $B(A_t) = T(A_t) + \mathbb{N} \times \{0\}^{t-1}$.
 (iii) $V(A_t) \subset \{T(A_t) + \mathbf{e}_i \mid i = 2, \dots, t\} \setminus T(A_t) \subset \{0\} \times \mathbb{N}^{t-1}$.
 (iv) The set $\{0\}^{i-1} \times \mathbb{N} \times \{0\}^{t-i} \cap V(A_t)$ consists of only one element for any i ($2 \leq i \leq t$).

Proof. We have $M(b_i) = (m_1, \dots, m_t) \in \{0\} \times \mathbb{N}^{t-1}$. In fact, if $m_1 \neq 0$, then we have $\Psi((0, m_2, \dots, m_t)) \equiv b_i \equiv i \pmod{a_1}$ and $\Psi((0, m_2, \dots, m_t)) < b_i$, which contradicts the definition of b_i . Therefore we have $M(b_i) \in T(A_t)$. For $M, N \in \{0\} \times \mathbb{N}^{t-1}$ satisfying $\Psi(M) > \Psi(N)$ and $\Psi(M) - \Psi(N) = ea_1$ for some $e \in \mathbb{N}_+$, we have $M \notin T(A_t)$. In fact, for $N' := (e, 0, \dots, 0) + N$, we have $M > N'$ and $\Psi(M) = \Psi(N')$, which means $M \notin B(A_t)$. Therefore we obtain (i).

Next we prove $B(A_t) \subset T(A_t) + \mathbb{N} \times \{0\}^{t-1}$. Let $M = (m_1, \dots, m_t) \in B(A_t)$, $M_1 = (0, m_2, \dots, m_t)$, and $M_2 = (m_1, 0, \dots, 0)$. Since $M_1 + M_2 \in B(A_t)$, we have $M_1, M_2 \in B(A_t)$. Since $M_1 \in B(A_t) \cap (\{0\} \times \mathbb{N}^{t-1}) = T(A_t)$, we have $M \in T(A_t) + \mathbb{N} \times \{0\}^{t-1}$. Suppose $B(A_t) \subsetneq T(A_t) + \mathbb{N} \times \{0\}^{t-1}$. Then from (i) there exist i ($0 \leq i \leq a_1 - 1$) and $M_3 \in \mathbb{N} \times \{0\}^{t-1}$ such that $M(b_i) + M_3 \notin B(A_t)$. Take $N \in B(A_t)$ satisfying $\Psi(M(b_i) + M_3) = \Psi(N)$. Since $N \in B(A_t) \subset T(A_t) + \mathbb{N} \times \{0\}^{t-1}$ and $\Psi(N) \equiv i \pmod{a_1}$, there exists $M_4 \in \mathbb{N} \times \{0\}^{t-1}$ such that $N = M(b_i) + M_4$. Therefore $M_3 > M_4$, $M_3, M_4 \in \mathbb{N} \times \{0\}^{t-1}$, and $\Psi(M_3) = \Psi(M_4)$, which is contradiction. Therefore we obtain $B(A_t) = T(A_t) + \mathbb{N} \times \{0\}^{t-1}$.

Next we prove $V(A_t) \subset \{0\} \times \mathbb{N}^{t-1}$. Let $M = (m_1, \dots, m_t) \in V(A_t)$, $M_1 = (0, m_2, \dots, m_t)$, and $M_2 = (m_1, 0, \dots, 0)$. Since $M \notin B(A_t)$ and $M_2 \in B(A_t)$, we have $M_1 \notin B(A_t)$. From the definition of $V(A_t)$, we obtain $M_2 = (0, \dots, 0)$. Therefore we obtain $V(A_t) \subset \{0\} \times \mathbb{N}^{t-1}$.

Let $M \in V(A_t) \subset \{0\} \times \mathbb{N}^{t-1}$. Since $M \neq (0, \dots, 0)$, there exist i ($2 \leq i \leq t$) and $M_1 \in \{0\} \times \mathbb{N}^{t-1}$ such that $M = M_1 + \mathbf{e}_i$. Since $M_1 \in B(A_t)$ from the definition of $V(A_t)$, we have $M_1 \in B(A_t) \cap (\{0\} \times \mathbb{N}^{t-1}) = T(A_t)$. Therefore we obtain (iii).

For $2 \leq i \leq t$, the set $\{0\}^{i-1} \times \mathbb{N} \times \{0\}^{t-i} \cap \{\mathbb{N}^t \setminus B(A_t)\}$ is not empty. In fact, since

$$\Psi((0, \dots, 0, a_1, 0, \dots, 0)) = \Psi((a_i, 0, \dots, 0)) = a_1 a_i,$$

we have $(0, \dots, 0, a_1, 0, \dots, 0) > (a_i, 0, \dots, 0)$. Let N_i be the minimal element of $\{0\}^{i-1} \times \mathbb{N} \times \{0\}^{t-i} \cap \{\mathbb{N}^t \setminus B(A_t)\}$. Then we obtain $\{0\}^{i-1} \times \mathbb{N} \times \{0\}^{t-i} \cap V(A_t) = \{N_i\}$. Therefore we obtain (iv). \square

Let $SV(A_t) = \{N_i \mid 2 \leq i \leq t\}$ (see the proof of Lemma C.1 (iv) for N_i). For $F = \sum \lambda_N X^N \in \mathbb{C}[X]$, we define *multideg* of F by

$$\text{multideg}(F) = \begin{cases} -\infty & \text{for } F = 0, \\ \max_{<} \{N \in \mathbb{N}^t \mid \lambda_N \neq 0\} & \text{for } F \neq 0. \end{cases}$$

Also we define *leading term* of F by

$$LT(F) = \begin{cases} 0 & \text{for } F = 0, \\ \lambda_T X^T & \text{for } F \neq 0, \text{ where } T = \text{multideg}(F). \end{cases}$$

For a ideal $J \subset \mathbb{C}[X]$, we define

$$\Delta(J) = \mathbb{N}^t \setminus \bigcup_{F \in J \setminus \{0\}} \{\text{multideg}(F) + \mathbb{N}^t\}.$$

Then we have

(6) $\text{Span}\{X^M \mid M \in \Delta(J)\} \cap J = \{0\}.$

Lemma C.2. (i) $\{F_M \mid M \in SV(A_t)\}$ is a Gröbner basis of the ideal $J := (\{F_M \mid M \in SV(A_t)\})$ with respect to the order $<$ in \mathbb{N}^t , i.e., $(\{LT(F) \mid F \in J\}) = (\{LT(F_M) \mid M \in SV(A_t)\})$.

(ii) $\text{Span}\{X^N \mid N \in B(A_t)\} \cap (\{F_M \mid M \in SV(A_t)\}) = \{0\}.$

Proof. For $M, N \in SV(A_t)$ ($M \neq N$), we have $\text{L.C.M.}\{LT(F_M), LT(F_N)\} = LT(F_M)LT(F_N)$. Therefore, from [6] p.102 Theorem 3 and p.103 Proposition 4, we obtain (i). From (i) we obtain $\Delta(\{F_M \mid M \in SV(A_t)\}) = \mathbb{N}^t \setminus \{SV(A_t) + \mathbb{N}^t\} \supset \mathbb{N}^t \setminus \{V(A_t) + \mathbb{N}^t\} = B(A_t)$, where the last equality is due to Lemma A.1. Since $\text{Span}\{X^N \mid N \in \Delta(\{F_M \mid M \in SV(A_t)\})\} \cap (\{F_M \mid M \in SV(A_t)\}) = \{0\}$ from (6), we have $\text{Span}\{X^N \mid N \in B(A_t)\} \cap (\{F_M \mid M \in SV(A_t)\}) = \{0\}$. □

Lemma C.3. If A_t is telescopic, then the following properties are satisfied.

(i) $T(A_t) = \{(0, m_2, \dots, m_t) \in \mathbb{N}^t \mid 0 \leq m_i \leq d_{i-1}/d_i - 1, i = 2, \dots, t\}.$

(ii) $SV(A_t) = V(A_t) = \{(d_{i-1}/d_i)\mathbf{e}_i \mid 2 \leq i \leq t\}.$

Proof. Let $U = \{(0, m_2, \dots, m_t) \in \mathbb{N}^t \mid 0 \leq m_i \leq d_{i-1}/d_i - 1, i = 2, \dots, t\}$. Take $u = (0, u_2, \dots, u_t) \in U$ and $v = (0, v_2, \dots, v_t) \in U$ satisfying $u \neq v$. First we prove $\Psi(u) \not\equiv \Psi(v) \pmod{a_1}$. Suppose that there exists an integer w such that $\Psi(u) - \Psi(v) = wa_1$. Let ρ be the positive integer such that $u_\rho \neq v_\rho, u_{\rho+1} = v_{\rho+1}, \dots, u_t = v_t$. Without loss of generality we assume $u_\rho > v_\rho$. Then we have $(u_\rho - v_\rho)a_\rho = wa_1 - \sum_{k=2}^{\rho-1} (u_k - v_k)a_k$ and $0 < u_\rho - v_\rho < d_{\rho-1}/d_\rho$, which is contradiction. Therefore we obtain $\Psi(u) \not\equiv \Psi(v) \pmod{a_1}$. Since A_t is telescopic, for any $u = (0, u_2, \dots, u_t) \in \mathbb{N}^t$, there exists $u' \in U$

such that $\Psi(u) \equiv \Psi(u') \pmod{a_1}$. Since $\Psi(u) \geq \Psi(u')$ and $\#U = a_1$, we have $\{\Psi(u) \mid u \in U\} = \{b_0, \dots, b_{a_1-1}\}$, where $b_i = \min\{b \in a_2\mathbb{N} + \dots + a_t\mathbb{N} \mid b \equiv i \pmod{a_1}\}$. Finally we prove $u \in B(A_t)$ for any $u \in U$. Take $u \in U$, then there exists $u'' = (u''_1, \dots, u''_t) \in B(A_t)$ such that $\Psi(u) = \Psi(u'')$. Since A_t is telescopic, we have $0 \leq u''_j < d_{j-1}/d_j$ for $2 \leq j \leq t$. Since $u''_1 = 0$ from the definition of b_i , we obtain $u'' \in U$. Therefore we obtain $u = u'' \in B(A_t)$. From Lemma C.1 (i), we obtain (i). From Lemma C.1 (iii) (iv) and the definition of $V(A_t)$, we obtain (ii). \square

Proof of Proposition 2.3. From Lemma C.2 (ii) and Lemma C.3 (ii), the condition (2) is satisfied. From Lemma C.1 (ii) and Lemma C.3 (i), we obtain Proposition 2.3 (i). From Lemma C.3 (ii), we obtain Proposition 2.3 (ii). From Proposition 2.1 (i) and Proposition 2.2 (i), the gap values at ∞ are $\mathbb{N} \setminus \langle A_t \rangle$. Therefore, from [16] Theorem 5, we obtain Proposition 2.3 (iii). \square

ACKNOWLEDGEMENT. The author would like to thank his supervisor Prof. Joe Suzuki for suggesting extending the sigma functions by using the Miura canonical form. The author would like to thank Prof. Yoshihiro Ônishi for his warm encouragements and valuable discussions. The author would like to thank Prof. Ryuichi Harasawa for valuable comments for the Miura canonical form. The author would like to thank the referee for the helpful comments. This research was supported by Grant-in-Aid for JSPS Fellows (22-2421) from Japan Society for the Promotion of Science.

References

- [1] V.M. Bukhshtaber, D.V. Leĭkin and V.Z. Ènol'skiĭ: *Kleinian functions, hyperelliptic Jacobians and applications*, Reviews in Math. and Math. Phys. **10**, Gordon and Breach, London, (1997), 1–125.
- [2] V.M. Bukhshtaber, D.V. Leĭkin and V.Z. Ènol'skiĭ: *Rational analogs of Abelian functions*, Funct. Anal. Appl. **33** (1999), 83–94.
- [3] V.M. Bukhshtaber, D.V. Leĭkin and V.Z. Ènol'skiĭ: *Uniformization of Jacobi varieties of trigonal curves and nonlinear differential equations*, Funct. Anal. Appl. **34** (2000), 159–171.
- [4] V.M. Bukhshtaber, D.V. Leĭkin and V.Z. Ènol'skiĭ: *σ -functions of (n, s) -curves*, Russian Math. Surveys **54** (1999), 628–629.
- [5] V.M. Buchstaber and D.V. Leĭkin: *Addition laws on Jacobian varieties of plane algebraic curves*, Proc. Steklov Inst. of Math. **251**, (2005), 49–120.
- [6] D. Cox, J. Little and D. O'Shea: *Ideals, Varieties, and Algorithms*, Undergraduate Texts in Mathematics, Springer, New York, 1992.
- [7] J.D. Fay: *Theta Functions on Riemann Surfaces*, Lecture Notes in Mathematics **352**, Springer, Berlin, 1973.
- [8] R. Hartshorne: *Algebraic Geometry*, Springer, New York, 1977.
- [9] F. Klein: *Ueber hyperelliptische Sigmafunctionen*, Math. Ann. **27** (1886), 431–464.
- [10] F. Klein: *Ueber hyperelliptische Sigmafunctionen*, Math. Ann. **32** (1888), 351–380.
- [11] S. Lang: *Introduction to Algebraic and Abelian Functions*, second edition, Springer, New York, 1982.

- [12] S. Matsutani: *Sigma functions for a space curve (3, 4, 5) type with an appendix by J. Komeda*, arXiv:1111.2413v1 [math-ph] 18 Dec 2011.
- [13] S. Miura: *Linear codes on affine algebraic curves*, IEICE Trans. **J81-A**, (1998), 1398–1421, in Japanese.
- [14] D. Mumford: *Tata Lectures on Theta, I, II*, Birkhäuser Boston, Boston, MA, 1983.
- [15] A. Nakayashiki: *On algebraic expressions of sigma functions for (n, s) curves*, Asian J. Math. **14** (2010), 175–211.
- [16] A. Nijenhuis and H.S. Wilf: *Representations of integers by linear forms in nonnegative integers*, J. Number Theory **4** (1972), 98–106.
- [17] J.H. Silverman: *The Arithmetic of Elliptic Curves*, Springer, New York, 1986.
- [18] J. Suzuki: *Miura conjecture on affine curves*, Osaka J. Math. **44** (2007), 187–196.
- [19] H. Stichtenoth: *Algebraic Function Fields and Codes*, Universitext, Springer, Berlin, 1993.

Department of Mathematics
Graduate School of Science
Osaka University
Machikaneyama-chou 1-1
Toyonaka-si, Osaka 560-0043
Japan
e-mail: t-ayano@cr.math.sci.osaka-u.ac.jp