

Title	One-pot electrochemical formation of meso,meso-linked porphyrin arrays
Author(s)	Ogawa, Takuji; Nishimoto, Yoshihiro; Ono, Noboru et al.
Citation	CHEMICAL COMMUNICATIONS. 1998, 3, p. 337-338
Version Type	VoR
URL	https://hdl.handle.net/11094/3271
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

One-pot electrochemical formation of *meso,meso*-linked porphyrin arrays

Takuji Ogawa,^{*a} Yoshihiro Nishimoto,^b Naoya Yoshida,^c Noboru Ono^b and Atsuhiko Osukua^{*c}

^a Institute for Fundamental Research of Organic Chemistry, Kyushu University, Hakozaki, Higashi-ku, Fukuoka 812-81, Japan

^b Department of Chemistry, Faculty of Science, Ehime University, Matsuyama 790, Japan

^c Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-01, Japan

Electrochemical oxidation of [5,15-bis(3,5-di-*tert*-butylphenyl)porphyrinato]zinc(II) **1**, [5,15-bis(4-methoxycarbonylphenyl)porphyrinato]zinc(II), [5,15-bis(pentafluorophenyl)porphyrinato]zinc(II) or [10-bromo-5,15-diphenylporphyrinato]zinc(II) in benzonitrile using a platinum net as the working electrode afforded the corresponding *meso,meso*-linked oligomer porphyrin arrays, and up to the octamer porphyrin array could be isolated for **1**.

Polynuclear porphyrins have been attracting considerable attention as biomimetic models of photosynthetic systems as well as photonic materials and functional molecular devices.^{1–7}

In a previous communication we reported the oxidative coupling reaction of [5,15-bis(3,5-di-*tert*-butylphenyl)porphyrinato]zinc(II) **1** with silver(I) salts to give the porphyrin dimer **2**, the porphyrin trimer **3** and the porphyrin tetramer **4**.⁸ The possible mechanism proposed is an initial one-electron oxidation of zinc porphyrin with Ag^I followed by nucleophilic attack of a neutral zinc porphyrin. This mechanism suggests that electrochemical oxidation also may lead to the formation of the *meso,meso*-coupled porphyrin arrays. Here we report an electrochemical preparation of the porphyrin arrays, by which the usage of expensive silver salts is not required, the oxidizing potentials can be best tuned to the porphyrin used, and the long porphyrin arrays can be prepared in a one-step procedure through prolonged electrolysis.

Porphyrin **1** ($n = 1$) (219 mg, 0.293 mmol) was electrolyzed in dry benzonitrile (210 ml) in a single cell with NBu₄ClO₄ (0.1 M) as the electrolyte, a platinum net (1.8 × 2.4 cm) as the counter electrode, and a platinum net (3 × 4 cm) as the working electrode; the electricity used was 0.732 mF. The porphyrin products were purified by being passed through an alumina and then a silica gel chromatography column, and isolated by preparative scale HPLC using a gel permeation column (GPC–HPLC). The structures were identified through comparison with the authentic samples:⁸ **1** (44.0%), **2** ($n = 2$) (20.7%), **3** ($n = 3$) (2.9%) and **4** ($n = 4$) (0.9%). When porphyrin **1** (211 mg, 0.282 mmol) was electrolyzed under similar conditions using electricity of 1.65 mF, the higher homologous pentamer **5** ($n = 5$), hexamer **6** ($n = 6$), heptamer **7** ($n = 7$) and octamer **8** ($n = 8$)


could be isolated at yields of 0.5, 0.4, 0.2 and 0.05%, respectively, together with **1** (6.7%), **2** (10.2%), **3** (3.2%) and **4** (1.6%) by GPC–HPLC. The new product was characterized by ¹H NMR spectroscopy (500 or 600 MHz) and FAB mass spectrometry or MALDI-TOF mass spectrometry.†

Treatment of [5,15-bis(4-methoxycarbonylphenyl)porphyrinato]zinc(II) **9** at higher oxidation potential (+0.68 V vs. Ag/AgNO₃) than **1** (+0.54 V vs. Ag/AgNO₃) with AgClO₄ under the conditions reported previously⁸ did not afford the corresponding *meso,meso*-linked porphyrin arrays, although an unidentified complex mixture together with ca. 60% of the recovered starting material were obtained. However, when **9** (51 mg, 79 μmol) was electrolyzed using electricity of 134 μF, it afforded the *meso,meso*-linked porphyrin dimer **10** (34.3%), porphyrin trimer **11** (6.9%) and the starting porphyrin **9** (47.4%). Similarly, [5,15-bis(pentafluorophenyl)porphyrinato]zinc(II) **12** (+0.77 V vs. Ag/AgNO₃) and (10-bromo-5,15-diphenylporphyrinato)zinc(II) **13** (+0.56 V vs. Ag/AgNO₃), which did not give the corresponding *meso,meso*-linked porphyrin arrays by the silver salt method, afforded the corresponding dimers **14** and **15** under similar conditions by electrolysis in isolated yields of 11.6 and 47.5%, respectively. Thus the electrochemical method has a wider scope for the substrates than the silver salt method.

These *meso,meso*-linked porphyrin arrays so far isolated were quite soluble in CHCl₃. Of particular note is the fact that even the arrays of porphyrin **9**, **12** and **13**, which have little sterical hindrance to prevent π–π stacking of the porphyrin planes, also had high solubility. This is probably because the porphyrin planes connect nearly perpendicular to each other due to their sterical requirement, and hence their intermolecular π–π stacking was largely prevented.

These results indicate that the present electrochemical method is a convenient way of preparing *meso,meso*-linked porphyrin arrays which should be good candidates for long ‘molecular wires’^{9,10} having a linear rigid structure and high solubility.

Work at Kyoto was partly supported by Grants-in Aid for Scientific Research (No. 08874074) from the Ministry of Education, Science, Sports and Culture of Japan and by the Tokuyama Science Foundation.



Scheme 1 Reagents and conditions: i, Anodic oxidation, Pt net as working electrode, Pt net as counter electrode, in benzonitrile, 0.1 M NBu₄ClO₄

Notes and References

* E-mail: ogawat@ms.ifoc.kyushu-u.ac.jp

† **5** (Ar = 3,5-di-*tert*-butylphenyl, R = H, $n = 5$): δ_{H} (CDCl₃) 1.37 (s, Bu^t), 1.40 (s, Bu^t), 1.50 (s, Bu^t), 7.62 (t, J 2.0, 4 H, Ar-4-H), 7.65 (t, J 1.5, 2 H, Ar-4-H), 7.76 (t, J 2.0, 4 H, Ar-4-H), 8.12 (d, J 2.0, 8 H, Ar-2,6-H), 8.15 (d, J 1.5, 4 H, Ar-2,6-H), 8.16 (d, J 2, 8 H, Ar-2,6-H) 8.19 (d, J 4.5, 4 H, β protons), 8.32 (d, J 4.5, 4 H, β protons), 8.35 (d, J 5.0, 4 H, β protons), 8.36 (d, J 5.0, 4 H, β protons), 8.75 (d, J 5.0, 4 H, β protons), 8.82 (d, J 4.5, 4 H, β protons), 8.84 (d, J 4.0, 4 H, β protons), 8.85 (d, J 4.5, 4 H, β protons), 9.22 (d, J 4.5, 4 H, β protons), 9.53 (d, J 4.5, 4 H, β protons) and 10.42 (s, 2 H); m/z (FAB) 3742.1 (C₂₄₀H₂₅₂N₂₀Zn₅ requires 3743.7). **6** (Ar = 3,5-di-*tert*-butylphenyl, R = H, $n = 6$): m/z (TOF) 4563 (C₂₈₈H₃₀₂N₂₄Zn₆ requires 4492). **7** (Ar = 3,5-di-*tert*-butylphenyl, R = H, $n = 7$): m/z (TOF) 5237 (C₃₃₆H₃₅₂N₂₈Zn₇ requires 5240). **8** (Ar = 3,5-di-*tert*-butylphenyl, R = H, $n = 8$): m/z (TOF) 6053 (C₃₈₄H₄₀₂N₃₂Zn₈ requires 5989). **10** (Ar = 4-methoxycarbonylphenyl, R = H, $n = 2$): δ_{H} (CDCl₃) 3.89 (s, 12 H), 8.03 (d, 4 H, J 5.0, β protons), 8.21 (d, 8 H, J 8.5, phenyl), 8.25 (d, 8 H, J 8.5, phenyl), 8.57 (d, 4 H, J 4.5, β protons), 9.03 (d, 4 H, J 4.5, β protons), 9.45 (d, 4 H, J 4.5, β protons) and 10.35 (s, 2 H, *meso* protons); m/z (FAB) 1281.1 (C₇₂H₄₆N₈O₈Zn₂ requires 1282.0). **11** (Ar = 4-methoxycarbonylphenyl, R = H, $n = 3$): δ_{H} ((CD₃)₂CO) 3.90 (s, 6 H, inner CH₃OCOC₆H₄), 4.04 (s, 12 H, outer CH₃OCOC₆H₄), 8.11 (d, 4 H, J 4.5, β protons), 8.23 (d, 4 H, J 4.5, β protons), 8.25–8.45 (m, 24 H, CH₃OCOC₆H₄), 8.57 (d, 4 H, J 5.0, β protons), 8.65 (d, 4 H, J 4.5, β protons), 9.05 (d, 4 H, J 4.5, β protons), 9.58 (d, 4 H, J 4.0, β protons) and 10.48 (s, 2 H, *meso* protons); m/z (FAB) 1921.7 (C₁₀₈H₆₈N₁₂O₁₂Zn₃ requires 1921.9). **14** (Ar = pentafluorophenyl, R = H, $n = 2$): δ_{H} (CDCl₃) 8.18 (d, 4 H, J 4.5, β protons), 8.65 (d, 4 H, J 5, β protons), 9.11 (d, 4 H, J 4.5, β protons), 9.61 (d, 4 H, J 5, β protons) and 10.50 (s, 2 H, *meso* protons); m/z (FAB) 1410 (C₆₄H₁₈F₂₀N₈Zn₂ requires

1410). **15** (Ar = Ph, R = Br, $n = 2$): δ_{H} (CDCl₃) 7.5–7.7 (m, 12 H, Ph), 8.06 (d, 4 H, J 5, β protons), 8.19 (dd, 8 H, J 8 and 2, Ph), 8.60 (d, 4 H, J 4.5, β protons), 9.04 (d, 4 H, J 5, protons) and 9.85 (d, 4 H, J 4.5, β protons); m/z (FAB) 1207.7 (C₆₄H₃₆Br₂N₈Zn₂ requires 1208.0).

- 1 M. O. Senge, W. W. Kalisch and K. Ruhlandt-Senge, *Chem. Commun.*, 1996, 2149.
- 2 L. Ruhlmann and A. Giraudeau, *Chem. Commun.*, 1996, 2007.
- 3 A. Giraudeau, L. Ruhlmann, L. E. Kahf and M. Gross, *J. Am. Chem. Soc.*, 1996, **118**, 2969.
- 4 A. Osuka, S. Marumo, N. Mataga, S. Taniguchi, T. Okada, I. Yamazaki, Y. Nishimura, T. Ohno and K. Nozaki, *J. Am. Chem. Soc.*, 1996, **118**, 155.
- 5 A. P. H. Schenning, F. B. G. Benneker, H. P. M. Geurts, X. Y. Liu and R. J. M. Nolte, *J. Am. Chem. Soc.*, 1996, **118**, 8549.
- 6 K. Susumu, T. Shimidzu, K. Tanaka and H. Segawa, *Tetrahedron Lett.*, 1996, **37**, 8399.
- 7 A. Osuka, S. Nakajima, T. Okada, S. Taniguchi, K. Nozaki, T. Ohno, I. Yamazaki, Y. Nishimura and N. Mataga, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 92.
- 8 A. Osuka and H. Shimidzu, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 135. Similar *meso,meso*-coupled porphyrin arrays have been prepared independently by different methods. Ref. 6 and R. G. Khoury, L. Jaquinod and K. M. Smith, *Chem. Commun.*, 1997, 1057.
- 9 J. S. Schumm, D. L. Pearson and J. M. Tour, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1360.
- 10 J. M. Tour, *Chem. Rev.*, 1996, **96**, 537.

Received in Cambridge, UK, 23rd October 1997; 7/07653C