

N-Tosyltriarylbi-muthimines. Synthesis and Reactions with Some Electrophiles


Hitomi SUZUKI,* Chie NAKAYA, Yoshihiro MATANO,* and Takuji OGAWA*†

Department of Chemistry, Faculty of Science, Kyoto University, Kitashirakawa, Sakyo-ku, Kyoto 606

†Department of Chemistry, Faculty of Science, Ehime University, Bunkyo-cho, Matsuyama 790

N-Tosyltriarylbi-muthimines were prepared by the reaction of triarylbi-muthines with chloramine-T and their properties were characterized. These bi-muthimines reacted with aromatic aldehydes, acid chlorides, and isocyanates to form N-arylidene-tosylamides, N-aryltosylamides, and N-aryl-N'-tosylureas, respectively. In the presence of copper catalyst, they readily decomposed into N-aryl- and N,N-diaryltosylamides, tosylamide, and triarylbi-muthine.

Of all imino compounds derived from stable 15th and 16th Group elements, bi-muthimine is the least known and the literature so far contains only one report,¹⁾ in which N-tosyltri-phenylbi-muthimine (**1a**) was obtained as a moisture sensitive solid by the reaction of tri-phenylbi-muthine with chloramine-T in dry acetonitrile under reflux. In the report, little or no properties of this compound were mentioned. In connection with our work on bi-muthonium ylides,²⁻⁶⁾ we became interested in this type of compounds and tried to prepare **1a** according to the reported procedure.⁷⁾ In spite of many attempts, however, the isolation of bi-muthimine **1a** as a crystalline solid was unsuccessful. The product obtained was either a pale yellow thick oil or a polymeric powder which was nearly insoluble in common organic solvents. Similar treatments of tri-*p*-tolylbi-muthine and tri-mesitylbi-muthine with chloramine-T⁸⁾ gave the corresponding bi-muthimines **1b** and **1c** as oily products, which gradually turned into a glassy mass on storage under vacuum. Bi-muthimine **1d** from tris(4-chlorophenyl)bi-muthine was less stable and gave no satisfactory elemental analysis. Electron-withdrawing substituent on aromatic ring attached to the bi-muth atom apparently lowers the stability of bi-muthimine **1d**. Some physical properties of four bi-muthimines **1a-d** are summarized in Table 1.



When methanol was substituted for acetonitrile, methyl diarylbismuthinates were obtained in moderate yields.⁹⁾

Bismuthimines **1a-c** are moderately stable under dry nitrogen, but readily decompose on standing in open air. They are soluble in acetonitrile, benzene, and dichloromethane; almost insoluble in hexane, and decompose in hydroxylic solvents. In solution bismuthimines were less stable and gradually separated a polymeric substance.

Table 1. Some physical properties of N-tosylbismuthimines (**1**)

Compound		¹ H NMR δ , ppm vs. TMS ^{e)}	IR ν /cm ⁻¹
1a ^{a)}	Pale orange oil	2.37(3H,s), 7.2-8.0(19H,m)	1460, 1325, 1155, 1080 920 <NaCl>
1b ^{b)}	Pale yellow glass	2.30(9H,s), 2.39(3H,s) 7.1-7.9(16H,m)	2920, 1450, 1335, 1160 1090, 920 <KBr>
1c ^{c)}	Pale yellow glass	1.95(9H,s), 2.00(3H,s), 2.55 (18H,s), 6.64(6H,s), 6.75 (2H,d, <i>J</i> =7.9 Hz), 7.68(2H,d, <i>J</i> =7.9 Hz)	2920, 1450, 1335, 1160 1070, 940 <KBr>
1d ^{d)}	Pale yellow oil	2.41(3H,s), 7.1-7.8(16H,m)	1480, 1335, 1160, 1090 925 <NaCl>

a) Found : C, 49.45 ; H, 3.92 ; N, 2.76%. Calcd for C₂₅H₂₂NO₂SBi : C, 49.27 ; H, 3.64 ; N, 2.30%.

b) New compound : Found : C, 51.71 ; H, 4.55 ; N, 2.55%. Calcd for C₂₈H₂₈NO₂SBi : C, 51.61 ; H, 4.33 ; N, 2.15%. The glassy mass decomposed up to ca. 50 °C without showing any definite melting range.

c) New compound ; no satisfactory analysis was available. The glassy mass decomposed up to ca. 75 °C without showing any definite melting range.

d) New compound ; bismuthimine **1d** was quite unstable and gave no satisfactory analysis.

e) **1a,b,d** : in CDCl₃. **1c** : in C₆D₆. Deuterated solvents were dried with molecular sieves 4A before use.

Measurements were carried out immediately after the preparation of each sample.


Table 2. N-Arylidene-tosylamides (**2**) obtained

R ¹	R ²	R ³	Solvent	Yield/% ^{a)}
H	H	H	CH ₃ CN	23
H	H	H	C ₆ H ₆	31
H	H	CH ₃	C ₆ H ₆	80 ^{b)}
H	H	Cl	C ₆ H ₆	72 ^{b)}
CH ₃	H	H	C ₆ H ₆	22
Cl	H	H	C ₆ H ₆	16


a) Yields refer to the isolated compounds and were not optimized. All products are known. Equimolar bismuthimine and aldehyde were heated for 1 h under reflux.

b) Bismuthimine / aldehyde ratio was 2 to 1.


When heated with aromatic aldehydes in benzene under reflux, bismuthimine **1** underwent a reaction analogous to the Wittig reaction, affording N-arylidenetosylamides (**2**) in yields shown in Table 2. Compound **1** failed to react with ketones.



Reaction of bismuthimine **1a** with benzoyl chloride proceeded slowly in hot benzene and gave after 24 h triphenylbismuth dichloride (**3**, 10%) and N-benzoyltosylamide (**4**, 13%) along with a significant amount of bismuth-containing polymeric substance. Benzoyl chloride itself lacks ability to acylate tosylamide under the conditions employed.



Bismuthimine **1a** reacted with phenylisocyanate instantaneously at room temperature and after aqueous work-up N-phenyl-N'-tosylurea (**5**) was obtained in 45% yield along with N,N'-diphenylurea (**6**, 38%). Carbon disulfide was unreactive toward **1**. Anticipating in vain a possible generation of a nitrenoid species, bismuthimine **1** was decomposed under various conditions in the presence of unsaturated compounds. However, copper powder as well as copper salt were found to promote self-decomposition of **1** into N-aryl- or N,N-diaryltosylamides, triarylbismuthine, and tosylamide.



Compound **1a**, previously described as a moisture-sensitive solid,¹⁾ failed to appear as such in spite of many attempts. Our syrupy product, when left to stand in air, gradually turned into a solid, major components of which were triphenylbismuthine, its oxide, tosylamide, and bismuth-containing polymeric powder. Thus, a "bismuthimine **1a**" reported by Wittig and coworker¹⁾ was likely to be a mixture of triphenylbismuth oxide and some other decomposition products from **1a**. Triphenylbismuth oxide was converted into pentaphenylbismuth by treatment with excess phenyllithium and this behavior is in accord with the reported property of "**1a**", on which structural assignment was made by the previous workers. On a similar treatment, our product failed to produce pentaphenylbismuth.

References

- 1) G. Wittig and D. Hellwinkel, *Chem. Ber.*, **97**, 789 (1964).
- 2) H. Suzuki, T. Murafuji, and T. Ogawa, *Chem. Lett.*, **1988**, 847.
- 3) T. Ogawa, T. Murafuji, and H. Suzuki, *Chem. Lett.*, **1988**, 849.
- 4) T. Ogawa, T. Murafuji, K. Iwata, and H. Suzuki, *Chem. Lett.*, **1989**, 325.
- 5) T. Ogawa, T. Murafuji, and H. Suzuki, *J. Chem. Soc., Chem. Commun.*, **1989**, 1749.
- 6) M. Yasui, T. Kikuchi, F. Iwasaki, H. Suzuki, T. Murafuji, and T. Ogawa, *J. Chem. Soc., Perkin Trans. I*, in press.
- 7) A representative procedure is as follows. To a suspension of chloramine-T (0.32 g, 1.4 mmol) in dry acetonitrile (20 ml) was added triphenylbismuthine (0.44 g, 1 mmol) and the mixture was heated at 75-80 °C with stirring under argon. The color of the suspension gradually turned yellow. After 3 h the mixture was cooled to room temperature and freed from the precipitated NaCl and unchanged chloramine-T. A yellowish filtrate was concentrated under reduced pressure and the residue was left in vacuo for 3 h to give a pale orange syrupy oil **1a** almost in a quantitative yield. Similarly, bismuthimines **1b-d** were obtained as a glassy mass (**1b,c**) or a syrupy oil (**1d**) in nearly quantitative yields. Prolonged heating of the reaction mixture led to decrease in yields of bismuthimines and extensive formation of bismuth-containing polymeric powder. Neither attempts to crystallize the bismuthimines from a variety of organic solvents nor to purify them by column chromatography were successful.
- 8) Commercial chloramine-T was dehydrated and dried under vacuum prior to use. Hydrated reagent gave the same product, although the formation of side products became rather prominent.
- 9) T. Ogawa, T. Murafuji, K. Iwata, and H. Suzuki, *Chem. Lett.*, **1988**, 2021.

(Received September 13, 1990)