

Title	一側性肺障害における肺血流分布の変化および低酸素呼吸の影響
Author(s)	竹田, 清
Citation	大阪大学, 1988, 博士論文
Version Type	VoR
URL	https://hdl.handle.net/11094/35694
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University



**HYPOXIA ENHANCES UNILATERAL LUNG INJURY
BY INCREASING BLOOD FLOW TO THE INJURED LUNG**

Kiyoshi Takeda, Mark J. Knapp, Walter G. Wolfe and James D. Crapo

**Departments of Medicine and Surgery
Duke University Medical Center
Durham, North Carolina 27710**

Abstract presented at the American Thoracic Society Meetings, May
11-14, 1986.

Dr. Knapp is a recipient of the Parker B. Francis grant.

Address Correspondence to:

James D. Crapo, M.D.

Box 3177

Duke University Medical Center

Durham, North Carolina 27710

(919) 684-6266

Abbreviated Title: Hypoxia enhances unilateral lung injury

ABSTRACT

We hypothesized that in unilateral lung injury, bilateral hypoxic ventilation would induce vasoconstriction in the normal lung, redirect blood flow to the injured lung, and cause enhanced edema formation. Unilateral left lung injury was induced by intrabronchial instillation of 1.5 ml/kg of 0.1 N HCl. After HCl injury, blood flow to the injured left lung decreased progressively from $.70 \pm .04$ to $.37 \pm .05$ L/min and percent of flow to the injured left lung (Q_L/Q_T) decreased from 37.7 ± 2.2 to $23.6 \pm 2.2\%$ at 240 min. Exposure to hypoxia (12% O₂), for three 10-minute episodes did not affect Q_L/Q_T in normal animals, but after unilateral HCl injury, it caused blood flow to the injured left lung to increase significantly. A concomitant decrease in blood flow occurred to the non-injured right lung, resulting in a significant increase in Q_L/Q_T . The enhanced blood flow to the injured lung was associated with a significant increase in the wet-to-dry lung weight ratio in the dependent regions of the injured lung. These findings demonstrate that in unilateral HCl-induced lung injury, transient hypoxia can enhance blood flow to the areas of injury and increase lung edema formation.

Index Terms: Hypoxia, acid aspiration, unilateral lung injury, lung water, hypoxic pulmonary vasoconstriction.

INTRODUCTION

In experimental unilateral lung injury, blood flow to the injured lung has been reported to decrease (1). This physiological response is important in maintaining arterial oxygenation by shunting blood flow away from the injured to the normal lung. One of the mechanisms for this response is postulated to be hypoxic pulmonary vasoconstriction (HPV). HPV has been demonstrated to exist in areas of atelectasis as well as during hypoxic ventilation in animals and humans (2,3). In addition to maintaining arterial oxygenation, HPV may lessen the severity of lung injury by reducing blood flow to the lung with increased capillary permeability. Most experimental models of acute lung injury, including acid aspiration, involve endothelial damage and are characterized by an enhanced permeability type of edema (4,5). Bishop *et al.* reported that minoxidil, a potent HPV inhibitor, further increased the production of extravascular lung water during the early phase of acute lung injury in rabbits when it was given prior to oleic acid administration (6). These findings suggest that when blood flow increases to areas of injury and increased capillary permeability, edema formation increases. Clinically, patients with acute lung injury may be exposed to transient hypoxia during procedures such as endotracheal suctioning (7) and bronchoscopic examination (8). During the hypoxic period, the pulmonary vasculature in the normal lung may elicit HPV and shunt blood flow to injured areas of the lung, resulting in aggravation of injury.

The purposes of this study were 1) to observe serial physiological changes, including blood flow distribution, after unilateral lung injury; 2) to determine the effects of brief periods of bilateral hypoxic

ventilation on blood flow distribution; and 3) to measure the extent of edema formation in the injured lung following hypoxic ventilation by following wet-to-dry lung weight ratios (W/D).

METHODS AND MATERIALS

Animal Preparation

Twenty adult, male mongrel dogs screened for heart worm infestation and weighing 17 to 25 kg were anesthetized with intravenous pentobarbital sodium (30 mg/kg) and intubated with a cuffed endotracheal tube (9.0 mm ID). Mechanical ventilation with 100% oxygen was maintained with a Harvard animal ventilator to achieve a PaCO₂ of 32 to 38 torr by adjusting the tidal volume (20 ml/kg) and respiratory rate (10-13 cycles/min). Before sternotomy, 7.5 mg/kg of pentobarbital sodium and 0.15 mg/kg of pancuronium bromide were administered intravenously. After this, anesthesia was maintained with hourly administration of pentobarbital sodium (3 mg/kg) and pancuronium bromide (0.05 mg/kg). By using this regimen, no sign of light anesthesia was observed. To minimize the effects of these drugs, they were given 45 minutes prior to measurements.

After induction of anesthesia, a balloon-tipped Swan-Ganz catheter (7.5 Fr.) was advanced into the main pulmonary artery through the right external jugular vein to measure pulmonary arterial pressure (PAP). The position of the catheter was assured by direct palpation after thoracotomy. By cutdown of the right groin, a 14-G polyethylene catheter was placed in the femoral artery to measure arterial pressure (AP). Another catheter was advanced into the right atrium through the right femoral vein to measure right atrial pressure (RAP). After placement of catheters, the chest was opened via a median sternotomy and ventilation was maintained with an application of 5 cm H₂O end-expiratory pressure. A catheter was placed by a direct left atriotomy

for the purpose of left atrial pressure (LAP) measurement. A segmental branch of pulmonary vein draining from each lower lobe was cannulated with a thin polyethylene tube. The tube was advanced from the segmental branch into the lobar vein to obtain mixed pulmonary venous blood. An electromagnetic flow probe (5 or 6 mm in diameter, Micron Co.) with a nonocclusive zero function was placed around the left main pulmonary artery, and the mean blood flow to the left lung was recorded on a strip chart recorder (Hewlett-Packard Co., Model 7758A). The flowmeter was calibrated in vivo in every experiment. Blood flow to the right lung (Q_R) was calculated as the difference between Q_T and Q_L . A 36 Fr. Carlen's tube, designed for left endobronchial intubation, was placed via a tracheostomy to facilitate unilateral instillation of hydrochloric acid (HCl) and was kept in place during the rest of the study.

All surgical procedures were done under sterile conditions. Normal saline was infused at a rate of 50 ml/hr via a catheter in a peripheral vein during the study period. Intravascular pressures were measured with Statham pressure transducers (P 23 Db) and recorded on a strip chart recorder. The mid-point of the left atrium was used as zero reference. Q_T was determined using the thermal dilution technique and was done in triplicate using 10 ml of 5% dextrose solution at room temperature. Blood samples were withdrawn from femoral and pulmonary arteries and pulmonary veins and pH, PO_2 and PCO_2 were measured immediately using a blood gas analyzer (Radiometer, ABL-30). Total, left and right pulmonary vascular resistances (PVR_T , PVR_L and PVR_R , respectively) were calculated from the perfusion pressure (PAP-LAP) divided by the respective lung blood flow. Systemic

vascular resistance (SVR) was also calculated from systemic perfusion pressure (AP-RAP) divided by Q_T . Both PVR and SVR were expressed as mmHg/l/min. Body temperature was maintained at 37.0°C by using a heating pad.

Study Design (Figure 1)

After stabilization of hemodynamics following surgery, baseline hemodynamic and blood gas data were collected. Dogs were divided into four groups. A control group consisted of five dogs who underwent the same surgical procedure and placement of catheters as the injured animals. In the HCl group, five dogs had HCl instillation into the left lung after baseline measurements. A 10-Fr polyvinyl chloride tube was advanced into the left main bronchus through a left-sided orifice of a Carlen's double-lumen catheter and 1.5 ml/kg body weight of 0.1 N HCl was instilled slowly. In the supine position, intrabronchially injected HCl went preferentially into the lower lobe. To induce a more homogeneous injury, HCl was instilled at four positions; supine, head down, head down + left side dependent recumbent and left side dependent recumbent positions. Distribution of HCl was confirmed by adding a small amount of 1% methylene blue as an indicator. Immediately after instillation of the HCl, intrabronchial suction of the right lung was performed to make sure that unilateral instillation into the left lung was achieved. After HCl administration, data were collected hourly for four hours. In the hypoxia group, 5 dogs were ventilated with a hypoxic gas mixture (12% O₂ + 88% N₂) for 10 minutes periodically at 60, 120 and 180 minutes after the baseline measurements. Hemodynamic and blood gas data were obtained at each pre- and post-hypoxic period.

The HCl + hypoxia group consisted of five dogs. They had both HCl instillation and hypoxic ventilation at the same times as the hypoxia group. After four hours of observation, the dogs were sacrificed with a fatal dose of pentobarbital sodium, and zero blood flow to the left lung was recorded in order to complete the in vivo calibration of the flow probe. Thereafter, both lower lobes were removed.

After the animals were sacrificed, the lungs and heart were removed en bloc by cutting the great vessels and allowing blood to drain freely into the chest. Next, the pericardium and heart were removed, and the lobes were individually separated. The identical procedure was carried out in all animals, so the wet-to-dry lung weight ratios (W/D) were not biased. After the lower lobes were removed, the dorsal (dependent) and ventral (non-dependent) parts of the lower lobes were identified in relation to the supine position of the dog over the six-hour experiment. The dorsal and ventral portions were separated by placing the lobes on a table in the exact anatomic relationship to the position of the animal and horizontally cutting the lobes in half.

Statistics

Analysis of variance followed by Duncan's multiple range test was used to compare physiologic data between and among groups. A two-sided, paired Student's t test was used to analyze data between pre- and post-hypoxia. A two-sided, unpaired Student's t test was used for comparison of W/D. Differences were considered statistically significant at P values of less than 0.05.

RESULTS

Effects of Acid Aspiration on Physiological Parameters

Hemodynamic changes following unilateral HCl administration are summarized in Table 1. In the control group, hemodynamics were stable during the study period. In the HCl-treated group, total pulmonary blood flow (Q_T) showed a tendency to decrease after HCl administration. At baseline blood flow to the left lung (Q_L) was $.70 \pm .04$ L/min and percent of flow to the left lung (Q_L/Q_T) was $37.7 \pm 2.2\%$. However, following HCl administration, Q_L decreased progressively to $.37 \pm .05$ L/min and Q_L/Q_T decreased to $23.6 \pm 2.2\%$ at the end of a 4 hour study (Figure 2). On the other hand, blood flow to the right lung (Q_R) was stable during the study. A slight decrease was seen in mean systemic artery pressure; however, mean pulmonary artery pressure (PAP) did not change during the four-hour study period. Those changes were reflected in the vascular resistances. Total pulmonary vascular resistance (PVR_T) showed a tendency to increase after 120 minutes. Pulmonary vascular resistance in the right lung (PVR_R) did not change; however, pulmonary vascular resistance in the left lung (PVR_L) increased significantly and at the end of the experiment, it increased to 184% of the baseline (Figure 3). Systemic vascular resistance (SVR) increased significantly 180 minutes after HCl administration and stayed elevated thereafter. Heart rate (HR), left atrial pressure (LAP) and right atrial pressure (RAP) did not change throughout the study.

Table 2 summarizes blood gas data. In the control group, PO_2 , PCO_2 and pH did not change during the study. Following HCl administration, the pulmonary venous PO_2 in the injured left lung decreased

significantly but the pulmonary venous PO_2 in the non-injured right lung was not affected. Arterial PO_2 decreased significantly during the first 2 hours after HCl; however, it showed a tendency to increase toward the baseline during the last 2 hours. Mixed venous PO_2 showed a time-related profile similar to the arterial PO_2 . An increase of PCO_2 and a decrease of pH were observed in left pulmonary venous blood after HCl administration. On the other hand, right pulmonary venous PCO_2 decreased markedly after HCl administration. PCO_2 and pH did not change in arterial and mixed venous blood.

Effects of Bilateral Hypoxic Ventilation on Unilateral Lung Injury.

Hemodynamic and blood gas changes following bilateral hypoxic ventilation are summarized in Tables 3 and 4, respectively.

In both the hypoxia and HCl + hypoxia groups, bilateral hypoxic ventilation caused a significant elevation in pulmonary artery pressure (PAP). The PAP increase was greater after the second and third hypoxic ventilation than after the first hypoxic ventilation as previously reported (9). In the hypoxia group, total blood flow (Q_T) and blood flow to both the left (Q_L) and right (Q_R) lungs increased significantly following hypoxic ventilation (Table 3). Because the increase in blood flow was uniformly distributed, percent of flow to the left lung (Q_L/Q_T) did not change (open symbols, Fig. 4). In the presence of HCl induced unilateral lung injury, exposure to hypoxia caused an increase in blood flow to the injured left lung in conjunction with a decrease in blood flow to the normal right lung (Table 3). The result was a consistent increase in percent blood flow to the left lung (solid symbols, Fig. 4). In Figure 5, changes in pulmonary vascular resistance are depicted. In the hypoxia alone group, pulmonary vascular

resistances in both left and right lungs increased significantly after the second and third hypoxic ventilation, suggesting that the non-injured lung had a vasoconstrictive response to hypoxia. In the HCl + hypoxia group, the non-injured right lung had a vasoconstrictive response to hypoxia and PVR_R increased significantly. However, in the injured left lung, no vasoconstrictive response to hypoxia was observed (Fig. 5).

Hypoxic ventilation caused a significant decrease in PO_2 of the blood obtained from every sampling site in both groups. In the hypoxia group, the PO_2 decreased significantly; however, there was no difference between right and left pulmonary venous blood. Due to the low PO_2 caused by HCl injury, left pulmonary venous PO_2 was consistently lower than right pulmonary venous PO_2 after hypoxic ventilation in the HCl + hypoxia group. In general, PCO_2 and pH were not affected by 10 minutes of exposure to hypoxia.

Lung Water (Table 5)

In the control and hypoxia groups, there were no statistically significant difference in either the W/D ratios of the right and left lungs or in the W/D ratios of the dependent and non-dependent aspects of the right and left lung. In the HCL group, the W/D ratios of both the dependent and non-dependent portions of the injured left lower lobe increased significantly from that of the uninjured right lower lobe: $4.96 \pm .13$ to $6.83 \pm .33$, and $5.16 \pm .07$ to $7.08 \pm .16$ respectively. In the HCL-hypoxia group, the W/D ratios of the dependent and non-dependent regions of the injured left lower lobe also increased significantly over the corresponding right lower lobe: $5.02 \pm .07$ to $6.88 \pm .22$, and $5.12 \pm .09$ to $7.74 \pm .17$ respectively.

Next, we compared the W/D ratios of the HCl group to the HCl-hypoxia group. There was no statistical difference between the W/D ratios in the ventral or non-dependent regions of the injured left lower lobes in the two groups: $6.83 \pm .33$ to $6.88 \pm .22$. However, there was a statistically significant increase in the W/D ratio in the dorsal or dependent regions of the injured left lower lobe in the HCl-hypoxia group versus the HCl group: $7.08 \pm .16$ to $7.74 \pm .17$ (Table 5).

DISCUSSION

Acid aspiration is one of the major causes of acute lung injury and is characterized by increased alveolar-capillary membrane permeability (4,5). Clinically, it is often recognized as a localized injury such as unilateral lung injury. The physiological sequences of pulmonary acid aspiration have been extensively studied (10-14). In general, there are decreases in heart rate, blood pressure and cardiac index immediately after HCl administration followed by a progressive decrease in cardiac index and a progressive increase in systemic and pulmonary vascular resistances with or without an elevation of pulmonary artery pressure. Fabre et al. (11) compared these sequences to an early parasympathetic and successive hypovolemic syndrome. Following unilateral HCl administration, we observed a progressive decrease in cardiac output; however, pulmonary artery pressure did not change during the study. These findings are consistent with the study of Cameron et al. (13). They also found a progressive increase in hematocrit and suggested that the animals were sequestering fluid into the injured lung and were becoming hypovolemic even though they were hemodynamically stable.

We demonstrated that blood flow to the injured left lung progressively decreased with a concomitant increase in pulmonary vascular resistance in the injured lung following HCl administration. We also found a marked decrease in the pulmonary venous PO_2 in the injured lung from 552 ± 22 to 70 ± 8 torr; however, arterial PO_2 was affected less. Using differentially labeled microspheres, Fisher and Wood (15) investigated the effects of lobar acid injury on lobar and sublobar pulmonary perfusion. They sectioned the lobe into visibly injured and non-injured

segments and demonstrated that percent blood flow to the injured region decreased significantly three hours after HCl administration. However, percent flow to the non-injured region was not altered. They also studied the effect of their blood sampling technique on lobar venous PO_2 and demonstrated that blood from the ventral and dorsal part of pulmonary vein had oxygen tensions of 400 torr and 60 - 70 torr, respectively and mixed pulmonary venous blood had an intermediate PO_2 . We collected mixed pulmonary venous blood with special attention to their findings; however, PO_2 was comparable to the PO_2 of blood from the dorsal (injured) part in their study. The lower PO_2 in our study may be attributed to a more severe injury since we used a larger amount of HCl. It is noteworthy that the percent decrease of flow to the injured segment in their study (-26.3%) is quite similar to our study (-27.1%) 3 hours after HCl administration.

Flow to the injured lung decreased because of a pulmonary vasoconstrictive response to acid injury. Local hypoxia induced by acid injury may elicit hypoxic pulmonary vasoconstriction (HPV) which, in turn, shunts blood to the normal region and maintains arterial oxygenation. The presence of HPV in acute lung injury is suggested by the fact that vasodilators can worsen gas exchange (16); however, this is still controversial. Vascular reactivity is reported to be increased in rat lungs injured with α -naphthylthiourea (17). On the other hand, HPV is reported to be altered or impaired in some lung injuries, such as bacterial pneumonia (18), endotoxemia (19,20), pulmonary oxygen toxicity (21,22), etc. Stephenson *et al.* (1) reported that in ethylchlorvynol-induced unilateral lung injury, the percent flow to the injured right lung decreased by 10%; however, this reduction was

inadequate to prevent a decrease in arterial PO_2 . It was suggested that, following acute lung injury, the reactivity of pulmonary vasculature was altered such that blood flow to hypoxic alveoli was maintained. In our study, percent flow to the injured left lung decreased by 37% 4 hours after HCl administration. Their study might be affected by a presence of systemic hypoxemia (PaO_2 60.7 torr). We found that arterial PO_2 decreased significantly, reaching its trough at 2 hours after HCl administration; however, it returned to pre-HCl level at 4 hours. This finding might be consistent with the study in which HPV is reported to be absent at one and two hours, but return by four hours after acid instillation in goats (23). Whatever the mechanism(s) is(are) to reduce the blood flow to the injured lung, this phenomenon is important to decrease the shunt and also to lessen the severity of injury which is characterized by permeability edema.

Another mechanism by which the flow to the injured lung decreases is mechanical. It has been suggested that perivascular cuffing during the early stage of pulmonary edema may cause a compression of small vasculature, resulting in a decreased blood flow to the edematous region (24). However, extensive edema is reported to be required to cause blood redistribution (25). By raising the pulmonary arterial and venous pressures, Bhattacharya *et al.* (26) induced edema in isolated perfused canine lobe. In edematous lobes, blood flow remained constant until lobe weight had doubled, suggesting that extensive edema is essential to reduce the blood flow to the edematous region. We observed alveolar flooding pathologically and a wet-to-dry lung weight ratio of 6.96, both of which suggest that severe pulmonary edema occurred. In addition, our preliminary study revealed that, at 4 hours

after HCl administration, the injured left lower lobe tripled its weight from a mean of 30.2 g to 87.6 g. If the HCl-induced edema occurred in a very short period of time, the blood flow decrease observed in our study might be explained by this mechanical factor.

In HCl-treated animals exposed to bilateral hypoxia, blood flow to the right non-injured lung decreased, but blood flow to the injured left lung increased (Fig. 4). This is explained by an increase in pulmonary vascular resistance in the non-injured lung without an increase in pulmonary vascular resistance in the injured lung which implies that either an impaired vascular reactivity to hypoxia or on-going vasoconstriction blocked the further response to hypoxia in the injured lung. It is plausible to postulate that an augmented blood flow to the injured lung can increase pulmonary shunt and aggravate the lung water accumulation. Bishop and Cheney (16) investigated effects of vasodilator treatment on gas exchange and lung water accumulation over a five-hour period in oleic acid-induced pulmonary edema. They found that minoxidil, which is a potent HPV inhibitor (27), increased venous admixture by 38%, whereas animals treated with hydralazine, which does not inhibit HPV (28), gave a smaller increase of 21%. Lung water accumulation was significantly accelerated by minoxidil; however, a smaller and non-significant increase was seen in the hydralazine group. Moreover, they observed a significantly higher mortality in minoxidil-treated dogs and concluded that vasodilators with a minimal effect on HPV might be preferable in the face of respiratory failure.

We hypothesized that the increase in blood flow secondary to bilateral hypoxia would increase intravascular pressure and, therefore, increase pulmonary edema in the injured lung. In our original protocol,

we sought to determine W/D ratios from both the dependent and non-dependent regions of the lung because we thought that injury may be increased in the dependent areas of the lung from conditions of this experiment. We found no increase in lung water accumulation in either the control group or in the hypoxia group alone, indicating no injury occurred secondary to manipulation and instrumentation of the animals. In both the HCl group and the HCl-hypoxia group, there were significant increases in W/D ratios in the injured lung. Also, in both of these injured groups, there was an increase in the W/D ratios in the dependent region versus the non-dependent region of the injured left lower lobe with the values in the HCl-hypoxia group achieving statistical significance. This is not surprising since gravity even in the normal human lung increases blood flow to the dependent areas of the lung in the upright position. In an injured capillary bed, this increase in blood flow would alter Starling's forces and increase edema formation. In the HCl-hypoxia group where blood flow was redirected from the normal lung to the injured lung by bilateral hypoxia, the W/D ratio from the dependent region was significantly greater than the corresponding region of the HCl group. These data indicate that local increases in pulmonary intravascular pressure either secondary to gravity or bilateral hypoxia will further increase W/D ratios and presumably lung water in an injured lung.

In clinical settings, patients with acute lung injury may be exposed to hypoxia during routine airway management procedures such as endotracheal suctioning (7), diagnostic bronchoscopy (8), ventilator circuit change (31), etc. Desaturation of arterial blood and hypoxia-induced arrhythmias have been reported during these procedures (32,

33). Our findings showed that hypoxia elicited a vasoconstrictive response in the normal lung and redistributed the blood flow to the injured lung. If this effect is repetitive or prolonged, it may cause significant increases in lung water accumulation and aggravate the acute lung injury.

In summary, following unilateral HCl administration, blood flow to the injured lung decreased without any effect on the contralateral lung. This decrease in flow to the injured lung is a protective mechanism to maintain arterial oxygenation and lessen the severity of lung injury. If blood flow to the injured lung is increased either by blockade of HPV or by an elevation of pulmonary vascular pressure, intrapulmonary shunting and lung water accumulation may be accelerated, resulting in more severe injury, especially in the acute phase of lung injury.

Acknowledgement

The writers wish to thank Dr. Vito A. Mantese for technical support. This work was supported in part by NIH Grant HL31992 and by a grant from RJR-Nabisco. Dr. Mark Knapp was a Parker B. Francis Fellow in pulmonary research.

REFERENCES

1. Stephenson, A.H., R.S. Sprague, T.E. Dahms, and A.J. Lonigro. Unilateral acute lung injury induced by ethylchlorvynol in anesthetized dogs. J Appl Physiol 56: 1252-1259, 1984.
2. Glasser, S.A., K.B. Domino, L. Lindgren, P. Parcella, C. Marshall, and B.E. Marshall. Pulmonary blood pressure and flow during atelectasis in the dog. Anesthesiology 58: 225-231, 1983.
3. Bindslev L., A. Jolin, G. Hedenstierna, S. Baehrendtz, and J. Santesson. Hypoxic pulmonary vasoconstriction in the human lung: Effect of repeated hypoxic challenges during anesthesia. Anesthesiology 62: 621-625, 1985.
4. Jones, J.G., R.F. Grossman, M. Berry, G. Slavin, G.H. Hulands, and B. Minty. Alveolar-capillary membrane permeability. Correlation with functional, radiographic and postmortem changes after fluid aspiration. Am Rev Respir Dis 120: 399-410, 1979.
5. Toung, J.K., D. Bordos, D.W. Benson, D. Carter, G.D. Zuidema, S. Permutt, and J.L. Cameron. Aspiration pneumonia: Experimental evaluation of albumin and steroid therapy. Ann Surg 183: 179-184, 1976.
6. Bishop, M.J., T. Huang, and F.W. Cheney. Effect of vasodilator treatment on the resolution of oleic acid injury in dogs. Am Rev Respir Dis 131: 421-425, 1985.
7. Adlkofer, S.R.M., and M.M. Powaser. The effect of endotracheal suctioning on arterial blood gases in patients after cardiac surgery. Heart Lung 7: 1011-1014, 1978.

8. Albertini, R.E., J.H. Harrell, N. Kurihara, and K.M. Moser. Arterial hypoxemia induced by fiberoptic bronchoscopy. JAMA 230: 1666-1667, 1974.
9. Unger, M., M. Atkins, W.A. Briscoe, and T.K.C. King. Potentiation of pulmonary vasoconstriction response with repeated intermittent hypoxia. J Appl Physiol 43: 662-667, 1977.
10. Greenfield, L.J., R.P. Singleton, D.R. McCaffree, and J.J. Coalson. Pulmonary effects of experimental graded aspiration of hydrochloric acid. Ann Surg 170: 74-86, 1969.
11. Fabre, J., C. Martin, C. Pomane, P. Long, and G. Francois. The early period of experimental Mendelson's syndrome in dogs with PaCO₂ and pH in a normal range. Resuscitation 11: 103-110, 1984.
12. Winn, R., J. Stothert, B. Nadir, and J. Hildebrandt. Lung mechanics following aspiration of 0.1 N hydrochloric acid. J Appl Physiol 55: 1051-1056, 1983.
13. Cameron, J.L., P. Caldini, J.K. Toung, and G.D. Zuidema. Aspiration pneumonia: Physiologic data following experimental aspiration. Surgery 72: 238-245, 1972.
14. Broe, P.J., J.K. Toung, S. Permutt, and J.L. Cameron. Aspiration pneumonia: Treatment with pulmonary vasodilators. Surgery 94: 95-99, 1983.
15. Fisher, C.J., Jr, and L.D.H. Wood. Effect of lobar acid injury on pulmonary perfusion and gas exchange in dogs. J Appl Physiol 49: 150-156, 1980.
16. Bishop, M.J., and F.W. Cheney. Vasodilators worsen gas exchange in dog oleic-acid lung injury. Anesthesiology 64: 435-439, 1986.

17. Hill, N.S., and S. Rounds. Vascular reactivity is increased in rat lungs injured with β -naphthylthiourea. J Appl Physiol 54: 1693-1701, 1983.
18. Lampron, N., F. Lemaire, B. Teisseire, A. Harf, M. Palot, D. Matamis, and A.M. Lorino. Mechanical ventilation with 100% oxygen does not increase intrapulmonary shunt in patients with severe bacterial pneumonia. Am Rev Respir Dis 131: 409-413, 1985.
19. Weir, E.K., J. Mlczoch, J.T. Reeves, and R.F. Grover. Endotoxin and prevention of hypoxic pulmonary vasoconstriction. J Lab Clin Med 88: 975-983, 1976.
20. Hutchison, A.A., M.L. Ogletree, J.R. Snapper, and K.L. Brigham. Effect of endotoxemia on hypoxic pulmonary vasoconstriction in unanesthetized sheep. J Appl Physiol 58: 1463-1468, 1985.
21. Newman, J.H., I.F. McMurty, and J.T. Reeves. Blunted pulmonary pressor responses to hypoxia in blood perfused, ventilated lungs isolated from oxygen toxic rats.: Possible role of prostaglandins. Prostaglandins 22: 11-20, 1981.
22. Newman, J.H., J.E. Loyd, D.K. English, M.L. Ogletree, W.J. Fulkerson, and K.L. Brigham. Effects of 100% oxygen on lung vascular function in awake sheep. J Appl Physiol 54: 1379-1386, 1983.
23. Pavlin, E., L. Reed, and R. Winn. Hypoxic pulmonary vasoconstriction (HPV) is absent at 1 and 2 hours after aspiration of 0.1 N HCl but returns at 4 and 24 hours. Am Rev Respir Dis 133: A225, 1986.

24. Hughes, J.M.B. *Regional Differences in the Lung*. New York, Academic Press; 1977, 381-418.
25. Muir, A.L., D.L. Hall, P.D. Despas, and J.C. Hogg. Distribution of blood flow in the lungs in acute pulmonary edema in dogs. J Appl Physiol 33: 763-769, 1972.
26. Bhattacharya, J., K. Nakahara, and N.C. Staub. Effect of edema on pulmonary blood flow in the isolated perfused dog lung lobe. J Appl Physiol 48: 444-449, 1980.
27. Bishop, M.J., and F.W. Cheney. Comparison of the effects of minoxidil and nifedipine on hypoxic pulmonary vasoconstriction in dogs. J Cardiovasc Pharmacol 5: 184-189, 1983.
28. Bishop, M.J., S. Kennard, L.D. Artman, and F.W. Cheney. Hydralazine does not inhibit canine hypoxic pulmonary vasoconstriction. Am Rev Respir Dis 128: 998-1001, 1983.
29. Julien, M., M.R. Flick, J.M. Hoeffel, and J.F. Murray. Accurate reference measurement for postmortem lung water. J Appl Physiol 56: 248-253, 1984.
30. Breen, P.H., P.T. Schumacker, J. Sandoval, I. Mayers, L. Oppenheimer, and L.D.H. Wood. Increased cardiac output increases shunt: role of pulmonary edema and perfusion. J Appl Physiol 59: 1313-1321, 1985.
31. Keller, C.A., and G.L. Ruppel. Oxygen desaturation during ventilator circuit changes. Heart Lung 14: 359-362, 1985.
32. Shrader, D.L., and S. Lakshminarayan. The effect of fiberoptic bronchoscopy on cardiac rhythm. Chest 73: 821-824, 1978.
33. Shim, C., N. Fine, R. Fernandez, and M.H. Williams, Jr. Cardiac arrhythmias resulting from tracheal suctioning. Ann Int Med 71: 1149-1153, 1969.

FIGURE LEGENDS

Figure 1: Scheme of experimental design. Arrows on the time line show when anesthetic drugs were given. Animals were ventilated with 100% O₂ after anesthesia was initiated except for the three 10-minute periods of hypoxia shown by the cross hatched areas on the third and fourth groups. The open circles with a solid center represent the times when physiologic data were collected.

Figure 2: Effect of HCl administration on percent of flow to the left lung (Q_L/Q_T).

In the control group (open circles), Q_L/Q_T did not change during the study. On the other hand, in the HCl group (solid circles), Q_L/Q_T decreased progressively.

* : P < 0.05 vs baseline

Figure 3: Effect of HCl administration on left and right pulmonary vascular resistances (PVR_L and PVR_R , respectively).

PVR_R in the non-injured right lung did not change, however, PVR_L in the injured left lung increased progressively after HCl administration.

* : P < 0.05 vs baseline

Figure 4: Effect of bilateral hypoxic ventilation on percent of flow to the left lung (Q_L/Q_T). The cross hatched regions represent the periods of hypoxic ventilation. There was no change in

Q_L/Q_T after bilateral hypoxic ventilation in the hypoxia group (open symbols). However, in the HCl + hypoxia group (solid symbols), hypoxic ventilation caused a significant increase in Q_L/Q_T , suggesting that a shift in blood flow toward the injured left lung occurred after hypoxia.

* : $P < 0.05$ vs comparable pre-hypoxia value.

Figure 5: Effects of bilateral hypoxic ventilation on left and right pulmonary vascular resistance (PVR_L and PVR_R , respectively). A. In the hypoxia group, bilateral hypoxic ventilation caused significant increases in PVR_L and PVR_R . B. In the HCl + hypoxia group, PVR_R in the non-injured right lung increased after hypoxic ventilation, however, in the injured left lung there was no increase in PVR_L after hypoxic ventilation.

In both A and B, the cross hatched regions represent the periods of hypoxic ventilation.

* : $P < 0.05$ vs the value immediately before that period of hypoxia.

TABLE 1. HEMODYNAMIC CHANGES AFTER HCl

	CONTROL				HCl			
	Baseline	240 min	60 min	120 min	180 min	240 min		
Q_T L/min	2.02 ± .10	2.12 ± .06	1.82 ± .17	1.59 ± .10	1.53 ± .09	1.56 ± .12		
Q_R L/min	1.33 ± .08	1.42 ± .06	1.21 ± .11	1.09 ± .07	1.10 ± .04	1.19 ± .09		
Q_L L/min	.69 ± .02	.70 ± .03	.61 ± .07	.50 ± .06*	.43 ± .05*	.37 ± .05*		
Q_L/Q_T %	34.1 ± .8	33.0 ± 1.4	33.5 ± 1.8	31.2 ± 2.8	27.5 ± 2.3*	23.6 ± 2.2*		
HR /min	162 ± 9	163 ± 10	148 ± 6	155 ± 6	151 ± 7	159 ± 5		
SAP mmHg	117 ± 7	126 ± 2	95 ± 8	98 ± 7	98 ± 7	99 ± 7		
CVP mmHg	3 ± 0	3 ± 1	2 ± 1	2 ± 1	1 ± 1	1 ± 1		
SVR mmHg/L/min	57.2 ± 4.7	57.9 ± 1.9	51.8 ± 3.3	60.3 ± 1.6	64.1 ± 4.2*	62.9 ± 2.5		
PAP mmHg	15 ± 1	15 ± 1	15 ± 1	14 ± 1	15 ± 1	14 ± 1		
LAP mmHg	5 ± 1	4 ± 1	4 ± 1	3 ± 1	4 ± 1	4 ± 1		
PVR_T mmHg/L/min	4.9 ± .4	5.4 ± .3	5.7 ± 1.0	6.6 ± .6	7.1 ± .3	6.4 ± .7		
PVR_R mmHg/L/min	7.5 ± .6	8.0 ± .6	8.6 ± 1.4	8.8 ± .9	9.8 ± .5	8.4 ± 1.0		
PVR_L mmHg/L/min	14.5 ± 1.3	16.1 ± 1.1	17.5 ± 3.4	20.4 ± 3.4	26.4 ± 2.2*	27.2 ± 2.1*		

Q_T : Total pulmonary blood flow
 Q_R : Blood flow to right lung
 Q_L : Blood flow to left lung
 Q_L/Q_T : Percent of flow to the left lung
HR : Heart rate

SAP : Mean systemic artery pressure
PAP : Mean pulmonary artery pressure
CVP : Central venous pressure
LAP : Left atrial pressure

SVR : Systemic vascular resistance
 PVR_T : Total pulmonary vascular resistance
 PVR_R : Right pulmonary vascular resistance
 PVR_L : Left pulmonary vascular resistance

Data are presented as mean ± SE.

* : P < 0.05 vs Baseline

TABLE 2. EFFECTS OF HCl ON GAS EXCHANGE

	CONTROL		HCl				
	Baseline	240 min	Baseline	60 min	120 min	180 min	240 min
Femoral							
PO ₂ †	500 ± 33	487 ± 42	558 ± 19	419 ± 40*	367 ± 50*	470 ± 22	471 ± 30
PCO ₂ †	35 ± 1	36 ± 1	38 ± 1	37 ± 2	35 ± 2	34 ± 2	33 ± 1
Artery							
pH	7.38 ± .01	7.38 ± .03	7.35 ± .03	7.36 ± .03	7.36 ± .03	7.37 ± .02	7.38 ± .02
Pulmonary							
PO ₂	46 ± 4	46 ± 3	44 ± 5	40 ± 3	40 ± 3	42 ± 3	41 ± 3
PCO ₂	43 ± 2	46 ± 2	48 ± 2	48 ± 1	45 ± 2	46 ± 3	48 ± 2
Artery							
pH	7.30 ± .01	7.31 ± .03	7.29 ± .01	7.29 ± .01	7.29 ± .01	7.27 ± .02	7.28 ± .02
Left							
PO ₂	494 ± 14	517 ± 32	552 ± 22	110 ± 31*	129 ± 39*	70 ± 8*	106 ± 28*
PCO ₂	26 ± 2	31 ± 2	27 ± 2	39 ± 4	39 ± 6	42 ± 5	38 ± 6
pH	7.44 ± .03	7.42 ± .03	7.44 ± .02	7.32 ± .03	7.32 ± .06	7.31 ± .04*	7.33 ± .05
Right							
PO ₂	519 ± 46	543 ± 29	558 ± 27	524 ± 10	556 ± 14	570 ± 17	532 ± 29
PCO ₂	29 ± 2	31 ± 1	32 ± 4	26 ± 4	28 ± 4	28 ± 4	29 ± 3
pH	7.44 ± .03	7.44 ± .05	7.40 ± .04	7.41 ± .04	7.40 ± .04	7.40 ± .04	7.38 ± .02

Data are presented as mean ± SE.

† : torr

* : P < 0.05 vs Baseline

TABLE 3 EFFECTS OF BILATERAL HYPOXIC VENTILATION ON HEMODYNAMIC PARAMETERS

	Hypoxia Group						HCl + Hypoxia Group					
	1st		2nd		3rd		1st		2nd		3rd	
	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia
Q_T l/min	2.23±.12	2.64±.12*	2.16±.10	2.59±.14**	2.17±.09	2.42±.11*	1.94±.14	2.06±.16*	1.73±.17	1.84±.16	1.63±.20	1.71±.16
Q_R l/min	1.42±.04	1.64±.18	1.43±.04	1.60±.09*	1.41±.04	1.52±.05*	1.34±.14	1.20±.14*	1.19±.17	.97±.09	1.15±.18	1.00±.09
Q_L l/min	.79±.09	1.00±.15*	.73±.09	.99±.11**	.76±.10	.89±.08*	.61±.05	.86±.08**	.54±.05	.87±.08*	.48±.06	.71±.10*
Q_L/Q_T %	34.9±2.1	38.0±5.6	34.0±2.7	37.5±2.9	34.7±3.2	36.8±1.7	31.7±3.2	42.3±4.1**	32.3±3.8	47.4±1.6**	30.2±3.9	41.2±3.0**
HR /min	158±7	160±9	158±7	160±7	155±6	162±8	148±9	149±10	150±7	152±7	150±7	150±9
SAP mmHg	130±10	130±10	128±10	131±11	126±10	129±10	105±7	102±8	99±7	93±9	96±9	90±13
CVP mmHg	0±1	0±1	0±1	1±1	0±1	0±1	1±1	2±1	1±1	2±1	0±1	2±1
PAP mmHg	15±1	23±1**	15±1	27±2**	15±1	27±2**	14±1	19±1**	13±0	21±1**	13±0	22±1**
LAP mmHg	1±1	2±1	1±1	1±1	1±1	1±1	1±0	2±1	2±0	3±1	2±0	2±1

Data are presented as mean ± SE.

Abbreviations used are the same as Table 1

* : p < 0.05 vs comparable pre-hypoxia value

** : p < 0.01 vs comparable pre-hypoxia value

TABLE 4 BLOOD GAS CHANGES AFTER BILATERAL HYPOXIC VENTILATION

	Hypoxia Group						HCl + Hypoxia Group					
	1st		2nd		3rd		1st		2nd		3rd	
	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia	Pre-Hypoxia	Post-Hypoxia
Femoral PO ₂ †	523±30	40±3**	490±35	39±2**	505±34	38±1**	301±45	40±3**	333±58	37±3**	389±51	37±2**
Femoral PCO ₂ †	35±1	32±1	31±1	31±2	33±1	31±2	34±2	34±1	34±2	33±2	33±2	34±3
Artery pH	7.35±.02	7.40±.03	7.38±.03	7.41±.04	7.37±.04	7.40±.03	7.39±.02	7.39±.01	7.39±.02	7.41±.03	7.39±.02	7.38±.03
Pulmonary PO ₂	53±4	27±2**	51±3	26±2**	50±3	24±2**	39±1	24±1**	38±1	23±1**	37±1	20±2**
Pulmonary Artery PCO ₂	40±1	38±1	40±1	35±3	41±2	37±2*	45±2	44±2	46±2	40±2	43±3	37±1
Pulmonary Artery pH	7.29±.03	7.34±.02	7.33±.02	7.4±.02	7.31±.02	7.34±.02*	7.31±.02	7.32±.01	7.30±.02	7.33±.01	7.29±.03	7.31±.02
Left PO ₂	478±31	47±5**	470±19	44±3**	485±13	43±4**	66±16	31±4*	70±21	27±2	60±15	27±4
Pulmonary PCO ₂	27±2	27±1	26±2	27±1	27±3	26±2	39±2	34±3*	39±4	38±3	40±5	41±3
Vein pH	7.40±.04	7.41±.03	7.43±.04	7.42±.03	7.41±.04	7.41±.04	7.35±.03	7.39±.04	7.35±.04	7.37±.03	7.34±.03	7.35±.03
Right PO ₂	534±29	49±2**	539±23	45±1**	547±33	43±3**	550±9	52±2**	533±22	49±4**	546±43	44±3**
Pulmonary PCO ₂	26±1	26±1	26±1	25±1	27±1	27±1	27±2	26±1	27±2	25±0	29±4	26±1
Vein pH	7.40±.02	7.42±.03	7.42±.02	7.43±.04	7.41±.02	7.43±.03	7.45±.04	7.44±.02	7.45±.02	7.45±.02	7.43±.01	7.43±.03

Data are presented as mean ± SE.

† ; torr

* ; p < 0.05 vs comparable pre-hypoxia value

** ; p < 0.01 vs comparable pre-hypoxia value

TABLE 5

REGIONAL WET/DRY WEIGHT RATIOS

	<u>Control Group</u>	<u>Hypoxia Group</u>	<u>HCl Group</u>	<u>HCl-Hypoxia Group</u>
<u>Left Lung</u>				
Ventral or Non-Dependent Region	4.91 ± .11	4.99 ± .06	6.83 ± .33*	6.88 ± .22 *
Dorsal or Dependent Region	4.98 ± .11	5.03 ± .09	7.08 ± .16*	7.74 ± .17 * † ‡
<u>Right Lung</u>				
Ventral or Non-Dependent Region	5.11 ± .13	5.11 ± .15	4.96 ± .13	5.02 ± .07
Dorsal or Dependent Region	5.15 ± .23	4.95 ± .07	5.16 ± .07	5.12 ± .09

All data are mean ± SE

* Significantly different from control (p < .05)

† Significantly different from HCl group (p < .05)

‡ Significantly different from HCl-hypoxia group, ventral or non-dependent region (p < .05)

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

