Title	Primary structure of human pancreatic α -amylase gene : its comparison with human salivary α -amylase gene
Author(s)	Horii, Akira
Citation	
Issue Date	
oaire:version	VoR
URL	https://hdl.handle.net/11094/36665
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

GEN 02187

Primary structure of human pancreatic α -amylase gene: its comparison with human salivary α -amylase gene

(Recombinant DNA; gene conversion; cosmid; evolutionary mouse and human divergence; exon-intron junctions)

Akira Horii, Mitsuru Emi *, Naohiro Tomita, Takahiro Nishide, Michio Ogawa ^a, Takesada Mori ^a and Kenichi Matsubara

Institute for Molecular and Cellular Biology, Osaka University, Yamadaoka, Suita 565 (Japan) and ^a Second Department of Surgery, Osaka University Medical School, Fukushima-ku, Osaka 553 (Japan) Tel. (06)451-0051

Received 18 June 1987

Revised 11 August 1987

Accepted 18 August 1987

SUMMARY

We have determined the entire structure of the human pancreatic α -amylase (Amy2) gene. It is approx. 9 kb long and is separated into ten exons. This gene (*amy2*) has a structure very similar to that of human salivary α -amylase (Amy1) gene [Nishide et al. Gene 41 (1986a) 299–304] in the nucleotide sequence and the size and location of the exons. The major difference lies in the fact that *amy1* has one extra exon on the 5' side. Other differences are at the 5' border of exon 1 and the 3' border of exon 10. The close similarity of these two genes, as compared with mouse pancreatic and salivary amylase genes, suggests that during evolution, the divergence into the two amylase genes may have occurred after the divergence of mice and man.

INTRODUCTION

Human α -amylases which hydrolyze α -1,4 glucosidic bonds, consist of two major isozymes, i.e., salivary type and pancreatic type. These two isozymes differ in molecular size (Matsuura et al., 1978; Stiefel et al., 1973), isoelectric point (Matsuura et al.,

1978) and antigenic properties (Boehm-Truitt et al., 1978). Our previous studies on cDNAs of salivary and pancreatic α -amylases (Nakamura et al., 1984; Nishide et al., 1986b) have shown that at least two different, but closely related genes are transcribed in tissue-specific fashion. In addition, salivary α -amylase gene (*amy1*), but not pancreatic α -amylase gene

Correspondence to: Dr. A. Horii, Institute for Molecular and Cellular Biology, Osaka University, Yamadaoka, Suita 565 (Japan) Tel. (06)877-5244.

* Present address: Howard Hughes Medical Institute, University of Utah, Salt Lake City, UT 84132 (U.S.A.) Tel. (801)581-3741.

Abbreviations: aa, amino acid(s); Amy1, human salivary α -amylase; *amy1*, gene, cDNA or mRNA coding for Amy1; Amy2, human pancreatic α -amylase; *amy2*, gene, cDNA or mRNA coding for Amy2; bp, base pair(s); cDNA, DNA complementary to RNA; cos, cosmid; kb, kilobase(s) or 1000 bp; N, any nucleoside; nt, nucleotide(s); p, plasmid; PolIk, Klenow (large) fragment of *E. coli* DNA polymerase I.

(*amy2*), is expressed in some cancers (Y. Nakamura, N.T., T.N., M.E., A.H., M.O., T.M., G. Kosaki, T. Okabe, M. Fujisawa, N. Osawa, T. Kameya and K.M., manuscript in prep.; N.T., N. Matsuura, A.H., M.E., T.N., M.O., T.M., O. Doi, and K.M., ms. in prep.). Thus, it seems to be of interest to clarify the mechanisms of the specific expression control, and therefore, we started to analyze these genes. This paper reports the structure of human *amy2* and compares it with the structure of *amy1* (Nishide et al., 1986a). The results show that these two genes are remarkably similar, albeit they express differently, suggesting that they separated very recently, or that there has been a very efficient gene conversion process to conserve the sequence during evolution.

MATERIALS AND METHODS

(a) Enzymes and reagents

Restriction endonucleases were purchased from Takara Shuzo (Kyoto, Japan) and Toyobo (Osaka Japan). Polk and *Escherichia coli* T4 ligase were from Takara Shuzo.

(b) Library screening

The human genomic cosmid library carrying approx. 40–45-kb inserts was kindly provided by Dr. Y. Nakamura (Lau et al., 1983). Recombinant colonies carrying human *amy1* and *amy2* were screened using the mixture of ³²P-labeled human *amy1* and *amy2* cDNAs (Nishide et al., 1986b) as the probe. Selection of a clone (cosP2) that carries the *amy2* is described in the text.

(c) Mapping of cosmid clones

Each cosmid clone was digested with some restriction endonucleases, electrophoresed in 0.5% or 0.35% agarose gels (Maniatis et al., 1982), transferred and hybridized to ³²P-labeled cDNA, or to ³²P-labeled fragment of the cosmid vector DNA, according to Southern (1975). We also used double digestion or partial digestion for fine mapping.

(d) Sequence analysis

DNA fragments were extracted from the cosmid clone (cosP2), subcloned in pUC8 (Vieira et al., 1982) or in Bluescribe (Vector Cloning Systems), and in M13 vectors mp10, mp11 (Messing, 1983), mp18 or mp19 (Yanisch-Perron et al., 1985), and sequenced by the dideoxy method (Sanger et al., 1980; Hattori et al., 1986).

(e) Poly(A)⁺ RNA preparation

A resected human pancreas was frozen in liquid nitrogen and stored at –70°C. RNA was isolated as described (Nakamura et al., 1984). Poly(A)⁺ RNA was purified by oligo(dT)-cellulose column chromatography.

(f) Primer extension analysis

Primer extension was carried out using pancreatic poly(A)⁺ RNA as a template and a 5' end-labeled synthetic oligodeoxynucleotide primer (5'–CCAG-CAGAACCCAATGGT–3'). This primer is complementary to a sequence of pancreatic α -amylase cDNA covering codons 9 through 14. Analysis was carried out in a 6% polyacrylamide–8 M urea gel along with appropriate dideoxy sequencing samples as size markers.

RESULTS AND DISCUSSION

(a) A cosmid carrying the human pancreatic α -amylase gene

The human genomic cosmid library was screened with human *amy1* and *amy2* cDNA probes (Nishide et al., 1986b). Out of 1.2×10^5 colonies screened, nine clones carrying full-length DNA covering an amylase gene were obtained. Eight of them gave an identical restriction cleavage pattern with that of salivary α -amylase gene (Nishide et al., 1986a). One, which we named cosP2, gave a different cleavage map. As will be shown below, the cosP2 carries the DNA that covers the entire human pancreatic α -amylase gene.

Fig. 1. Restriction map of human genes *amy2* and *amy1*. Relative positions of exons, deduced from the nucleotide sequence, are indicated by solid boxes. B, *Bam*HI; E, *Eco*RI; G, *Bgl*II; H, *Hind*III; Hp, *Hpa*I; N, *Ban*III; P, *Pst*I; S, *Sac*I; X, *Xba*I. Only relevant restriction sites are displayed. A few errors in the restriction map previously published (Nishide et al., 1986a) have been corrected. The numbering of the exons in *amy1* has been revised so as to unify the nomenclatures for the two genes. The primer used for extension analysis is also shown, with its 5'–3' orientation, above the exon 1 of *amy2*.

(b) The structure of *amy2*

The insert in the cosP2 was analyzed by restriction endonucleases. The fragments were monitored by their ability to hybridize to *amy1* and *amy2* cDNA regional probes (Nakamura et al., 1984; Nishide et al., 1986b). The fragments were subcloned for fine restriction mapping and nucleotide sequencing. The restriction map and location of exons (see below) are shown in Fig. 1, and the nucleotide sequence, along with the predicted amino acid sequence, is shown in Fig. 2. The nucleotide sequence of the amino acid coding region, along with its 5'- and 3'-flanking regions which were regarded to be exons, completely agrees with that of human pancreatic α -amylase cDNA (Nishide et al., 1986b). These sequences also agree with the cDNA sequence reported by Wise et al. (1984), except for 1 nt (thymine instead of cytosine) in exon 8 at aa position 370. These results indicate that the gene we analyzed in cosP2 is actually the human pancreatic α -amylase gene.

Fig. 1 also shows the map and location of exons in *amy1*. The two genes are very similar in structure including the locations and lengths of exons as well as introns. One of the major differences lies in the fact that *amy1* has an extra exon, referred to as exon S*, which is lacking from *amy2* (see below). The 5'

border of exon 1 and the 3' border of exon 10, each differ to some extent between *amy2* and *amy1*. The difference in the 5' border of exon 1 (an extra 32 nt is transcribed in the exon 1 of the *amy1*) can be used to monitor the presence of a small amount of *amy2* mRNA among large amounts of *amy1* mRNA (Y. Nakamura, N.T., T.N., M.E., A.H., M.O., T.M., G. Kosaki, T. Okabe, M. Fujisawa, N. Osawa, T. Kameya and K.M., ms. in prep.)

In addition to the overall homologies between transcribed sequences in *amy2* and *amy1*, we see striking similarities in the size of introns and the sequences that cover the 5' and 3' flanking regions of the two genes. The 5' untranscribed region of *amy2* which corresponds to the exon S of *amy1* is 93% homologous to the exon S. This raised the question as to whether *amy2* is transcribed in this region. Accordingly, we determined the transcription start point of *amy2* by primer extension analysis, as shown in Fig. 3, using the mRNA from human pancreas. The result showed unambiguously that the transcription starts at 14 bp upstream from the start codon in the exon 1. S1 nuclease mapping confirmed this result (not shown). Therefore, the homologous sequence to the exon S in *amy2* is not transcribed. The TATA box of *amy2* lies 29–23 nt upstream from the transcription start point which lies in the exon 1 in *amy1*. The TATA box of *amy1* lies farther upstream from the exon S (34–28 nt upstream from the cap site of *amy1*).

* The exons in *amy1* were renumbered so as to unify the nomenclature of corresponding exons in the two genes (viz., *amy1* and *amy2*). Thus, *amy1* consists of exons S, 1, 2, 3, ... 9, 10 and *amy2* consists of exons 1, 2, 3, ... 9, 10.

[illegible]

Fig. 2. The nucleotide and predicted amino acid sequences of human *amy2* and *amy1*. The entire sequence for *amy2* is shown. Only the differing nucleotides or amino acids in *amy1* are presented beneath *amy2* sequence. Exons are shaded and boxed with dashed lines, and the putative TATA boxes and poly(A) signals are boxed with solid lines. A few errors in the sequences of nucleotides and predicted amino acids in *amy1* previously published (Nishide et al., 1986a,b) have been corrected. Dashes in the nucleotide sequences indicate deletions.

Fig. 3. Primer extension analysis. Lanes A, G, C and T show the known nucleotide sequence ladders employed as size standards (30–70 nt). Lane P shows the cDNA primed with a ^{32}P -5'-end-labeled 18-nt synthetic oligodeoxynucleotide (see MATERIALS AND METHODS, section f, and Fig. 1) using pancreatic poly(A)⁺ RNA as a template. The arrow indicates the primer extension product, 56 nt in size, containing a 42-nt coding region and a 14-nt 5'-nontranslated region.

(c) Comparison of amylase genes between man and mouse

Schibler et al. (1982) published structures of α -amylase genes in mouse. They showed that in this animal the *amy2* gene consists of ten exons and the *amy1* gene of eleven. In addition to the pancreas and salivary gland, α -amylase can be produced in the liver, whose gene consists of eleven exons. The structures of mouse and human α -amylase genes are similar from exon 1 through exon 10, which is equivalent to *amy2* (Fig. 4). These four genes show close similarity in sizes of exons 1 through 10. In this region, the nucleotide sequences of mRNAs representing *amy2* exons are 85% homologous between man and mice, and 81% homologous, with respect to *amy1* exons (Hagenbüchle et al., 1980). Fig. 4 shows that the sizes of introns in human and mouse amylase genes differ to some extent. To our surprise, introns in human pancreatic and salivary amylase

genes also show very close similarities. As far as we sequenced, the nucleotide sequences are more than 90% homologous. The extra exon (exon S) in the salivary amylase genes in both man and mice have no sequence homology. Moreover, their size and location show no similarities. Thus, the size of exon S in man is 154 bp and is located 357 bp upstream from the 5' end of exon 1, whereas in mice, the size is 50 bp and the location is 7.77 kb upstream from the 5' end of exon 1 (Schibler et al., 1982). It thus seems likely that at some stage of evolution, the exon S in both species was created by mutation in the upstream region to the basal ten exons which may represent the primordial amylase gene and is similar to *amy2*. The expression of amylase in mouse liver (Schibler et al., 1980; 1982; 1983; Young et al., 1981) is also mediated through an additional exon, called exon L, that is located at the 5' side of the basal ten exons. Acquisition of such an extra exon for expression in the salivary gland (or in the liver) in both mammalian species is interesting for the tissue-specific expression of a gene, and suggests that similar mechanisms may be found in some other genes, particularly those coding isozymes that are expressed in tissue-specific fashion.

The nucleotide sequence homology between the basal ten exons in *amy1* and *amy2* is 98% in humans. Not only are the lengths and locations of the exons similar, but also the introns and flanking regions. Close similarity between the basal ten exons is also observed in *amy1* and *amy2* in mice (89% homologous). On the other hand, the overall nucleotide sequence homology between corresponding exons from the two mammals is 81% (salivary type) and 85% (pancreas type). These data suggest that during evolution, the divergence into the two amylase genes (viz. *amy1* and *amy2*) occurred after the divergence of mice and man. Alternatively, the highly conserved sequence homologies within a species, may be accounted for by gene conversion among the members of the amylase gene family.

Swift et al. (1984), Boulet et al. (1986) and Osborn et al. (1987) found a conserved sequence, possibly a part of the enhancer sequence that plays a role in pancreas-specific gene expression. The corresponding 5' region of human *amy2* has the almost identical sequence (AGGTCATTTAGATGATTTCCATGAGAGACTT) at 153–183 bp upstream from the cap site of *amy2*. Interestingly, the human

amyl has this sequence, too, except for substitution of 9th T by G in intron S. The 5'-flanking region of *amy2* is 94% homologous to the exon S of *amyl*, as shown in Fig. 2. Our unpublished data show that up to at least 0.6 kb from the cap site of human *amyl*, the sequences for *amyl* and *amy2* are very similar. Thus, from sequencing studies alone, it is not possible to point out the *cis*-acting control regions for tissue-specific expression. Further analyses are needed to clarify these problems.

ACKNOWLEDGEMENTS

We thank Dr. Y. Nakamura for providing us with the human DNA cosmid library, Drs. T. Yamamoto and Y. Takahashi for their advice and discussions, and Mr. T. Umekage for his technical assistance.

REFERENCES

- Boehm-Truitt, M., Harrison, E., Wolf, R.O. and Notkins, A.L.: Radioimmunoassay for human salivary amylase. *Anal. Biochem.* 85 (1978) 476-487.
- Boulet, A.M., Erwin, C.R. and Rutter, W.J.: Cell-specific enhancers in the rat exocrine pancreas. *Proc. Natl. Acad. Sci. USA* 83 (1986) 3599-3603.
- Hagenbüchle, O., Bovey, R. and Young, R.A.: Tissue-specific expression of mouse α -amylase genes: nucleotide sequence of isozyme mRNAs from pancreas and salivary gland. *Cell* 21 (1980) 179-187.
- Hattori, M. and Sakaki, Y.: Dideoxy sequencing method using denatured plasmid templates. *Anal. Biochem.* 152 (1986) 232-238.
- Lau, Y.-F. and Kan, Y.W.: Versatile cosmid vectors for the isolation, expression, and rescue of gene cluster. *Proc. Natl. Acad. Sci. USA* 80 (1983) 5225-5229.
- Maniatis, T., Fritsch, E.F. and Sambrook, J.: *Molecular Cloning. A Laboratory Manual.* Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1982.
- Matsuura, K., Ogawa, M., Kosaki, G., Minamimura, N. and Yamamoto, T.: α -Amylase from human pancreatic juice as an electrophoretically pure isozyme. *J. Biochem.* 83 (1978) 329-332.
- Messing, J.: New M13 vectors for cloning. *Methods Enzymol.* 101 (1983) 20-78.
- Nakamura, Y., Ogawa, M., Nishide, T., Emi, M., Kosaki, G., Himeno, S. and Matsubara, K.: Sequences of cDNAs for human salivary and pancreatic α -amylases. *Gene* 28 (1984) 263-270.
- Nishide, T., Nakamura, Y., Emi, M., Yamamoto, T., Ogawa, M., Mori, T. and Matsubara, K.: Primary structure of human salivary α -amylase gene. *Gene* 41 (1986a) 299-304.
- Nishide, T., Emi, M., Nakamura, Y. and Matsubara, K.: Corrected sequences of cDNAs for human salivary and pancreatic α -amylases. *Gene* 50 (1986b) 371-372.
- Osborn, L., Rosenberg, M.P., Keller, S.A. and Meisler, M.H.: Tissue specific and insulin-dependent expression of a pancreatic amylase gene in transgenic mice. *Mol. Cell. Biol.* 7 (1987) 326-334.
- Sanger, F., Coulson, A.R., Barrell, B.G., Smith, A.J.H. and Roe, B.A.: Cloning in single-stranded bacteriophage as an aid to rapid DNA sequencing. *J. Mol. Biol.* 143 (1980) 161-178.
- Schibler, U., Tosi, M., Pittet, A.-C., Fabiani, L. and Wellauer, P.K.: Tissue-specific expression of mouse α -amylase genes. *J. Mol. Biol.* 142 (1980) 93-116.
- Schibler, U., Pittet, A.-C., Young, R.A., Hagenbüchle, O., Tosi, M., Gellman, S. and Wellauer, P.K.: The mouse α -amylase multigene family: Sequence organization of members expressed in the pancreas, salivary gland and liver. *J. Mol. Biol.* 155 (1982) 247-266.
- Schibler, U., Hagenbüchle, O., Wellauer, P.K. and Pittet, A.C.: Two promoters of different strengths control the transcription of the mouse alpha-amylase gene *Amy-1^a* in the parotid gland and the liver. *Cell* 33 (1983) 501-508.
- Southern, E.M.: Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98 (1975) 503-517.
- Stiefel, D.J. and Keller, P.J.: Preparation and some properties of human pancreatic amylase including a comparison with human parotid amylase. *Biochim. Biophys. Acta* 302 (1973) 345-361.
- Swift, G.H., Craik, C.S., Stary, S.J., Quinto, C., Lahaie, R.G., Rutter, W.J. and MacDonald, R.J.: Structure of the two related elastase genes expressed in the rat pancreas. *J. Biol. Chem.* 259 (1984) 14271-14278.
- Vieira, J. and Messing, J.: The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers. *Gene* 19 (1982) 259-268.
- Wise, R.J., Karn, R.C., Larsen, S.H., Hodes, M.E., Gardell, S.J. and Rutter, W.J.: A complementary DNA sequence that predicts a human pancreatic amylase primary structure consistent with the electrophoretic mobility of the common isozymes, Amy2 A. *Mol. Biol. Med.* 2 (1984) 307-322.
- Yanisch-Perron, C., Vieira, J. and Messing, J.: Improved M13 phage cloning vectors and host strains: nucleotide sequence of the M13mp18 and pUC19 vectors. *Gene* 33 (1985) 103-119.
- Young, R.A., Hagenbüchle, O. and Schibler, U.: A single mouse α -amylase gene specifies two different tissue-specific mRNAs. *Cell* 23 (1981) 451-458.

Communicated by H. Yoshikawa.