

Title	A note on Azumaya's theorem
Author(s)	Sugano, Kozo
Citation	Osaka Journal of Mathematics. 4(1) P.157-P.160
Issue Date	1967
Text Version	publisher
URL	<a href="https://doi.org/10.18910/3729">https://doi.org/10.18910/3729</a>
DOI	10.18910/3729
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## A NOTE ON AZUMAYA'S THEOREM

Kozo SUGANO

(Received May 4, 1967)

We say that a left  $\Lambda$ -module  $M$  is a cogenerator of the category of left  $\Lambda$ -modules if for every submodule  $N_1$  of a left  $\Lambda$ -module  $N$  there exists a  $\Lambda$ -homomorphism  $f$  from  $N$  to  $M$  such that  $f(N_1) \neq 0$ . Let  $\{I_\alpha\}$  be the full set of non isomorphic irreducible  $\Lambda$ -modules, and  $\{E_\alpha\}$  be the set of their injective hulls. Then a left  $\Lambda$ -module  $M$  is a cogenerator if and only if  $M$  contains every  $E_\alpha$ . In this case the sum of all  $E_\alpha$ 's is a direct sum by Zorn's Lemma (see Lemma 1[3]). A cogenerator is a faithful module (Lemma 2). The aim of this paper is to compare the ring for which every faithful module is a cogenerator with the ring for which every faithful module is generator (see G. Azumaya [1]). We assume every ring has units and every module is unitary.

**Lemma 1.** *Let  $M$  be a left  $\Lambda$ -module and  $A$  be an arbitrary set of index. Then the followings are equivalent.*

- (1)  $M$  is a cogenerator
- (2)  $\sum_{\nu \in A}^\oplus M_\nu$  ( $M_\nu \cong M$ ) is a cogenerator
- (3)  $\prod_{\nu \in A} M_\nu$  ( $M_\nu \cong M$ ) is a generator

Proof. (1) $\Rightarrow$ (2) $\Rightarrow$ (3) is clear by Lemma 1 [3]. So we shall prove (3) $\Rightarrow$ (1).

Choose any  $E_\alpha$ , then we have  $\Lambda$ -maps  $E_\alpha \xrightarrow{\tau} \prod M_\nu \xrightarrow{\pi_\nu} M_\nu$ , where  $\tau$  is a monomorphism and  $\pi_\nu$  are the canonical maps. Let  $f_\nu = \pi_\nu \cdot \tau$  then we see  $\bigcap \ker f_\nu = 0$ . If  $\ker f_\nu \neq 0$  for every  $\nu \in A$ , then  $I_\alpha \subseteq \bigcap \ker f_\nu \neq 0$  since  $I_\alpha$  is irreducible and  $E_\alpha$  is an essential extension of  $I_\alpha$ . Hence  $M_\nu$  has an isomorphic image of  $E_\alpha$  and  $M$  is a cogenerator.

**Corollary 1.** *A ring  $\Lambda$  is a self-cogenerator ring if and only if every  $E_\alpha$ , i.e.,  $\sum^\oplus E_\alpha$  is projective.*

Proof. If  $\sum^\oplus E_\alpha$  is projective,  $\sum^\oplus E_\alpha < \oplus \sum^\oplus \Lambda$ , and  $\sum^\oplus \Lambda$  is a cogenerator. Hence  $\Lambda$  is a cogenerator by Lemma 1. Conversely if  $\Lambda$  is a cogenerator,  $\Lambda \oplus > E_\alpha$ , and  $E_\alpha$  is projective.

**Lemma 2.** *If  $M$  is a cogenerator and  $\mathfrak{I}$  is a left ideal of  $\Lambda$ , then  $\mathfrak{I} = l_\Lambda(r_M(\mathfrak{I}))$ . Hence every cogenerator is faithful. Conversely assume that  $\Lambda$  is a left self-cogenerator ring. Then every faithful module is a cogenerator.*

Proof. In general we have  $l_\Lambda(r_M(I)) \supseteq I$ . If  $l_\Lambda(r_M(I)) \neq I$ , we have a  $\Lambda$ -homomorphism  $f$  from  $\Lambda/I$  to  $M$  such that  $f(l_\Lambda(r_M(I)/I)) \neq 0$ . This means that there exists  $m \in M$  such that  $Im = 0$  and  $l_\Lambda(r_M(I)) m \neq 0$ , which is a contradiction since  $r_M(l_\Lambda(r_M(I))) = r_M(I)$ . Thus  $I = l_\Lambda(r_M(I))$ . Since  $0 = l_\Lambda(r_M(0)) = l(M)$ ,  $M$  is faithful. If  $\Lambda$  is a left self cogenerator ring and  $M$  is a faithful left module, we have a monomorphism  $f$  from  $\Lambda$  to  $\Pi M$  such that  $f(\lambda) = (\lambda m)_{m \in M}$ . Hence  $\Pi M$  is a cogenerator, consequently  $M$  is a cogenerator by Lemma 1.

**Lemma 3.** *If  $\Lambda$  is a self cogenerator ring,  $l(x)$  is 0 or a minimal left ideal for every maximal right ideal of  $\Lambda$  and  $r(I)$  is an essential extension of a minimal right ideal for every maximal left ideal. Consequently every right ideal (resp. if  $\Lambda$  is right lower distinguished, every left ideal) contains a minimal right (resp. left) ideal of  $\Lambda$ .*

Proof. If  $l(x) \neq 0$ ,  $r = r(l(x)) = r(I')$ , and  $l(x) = l(r(I')) = I'$ , where  $I'$  is an arbitrary left subideal of  $l(x)$ . Thus  $l(x)$  is a minimal left ideal. Let  $I$  be a maximal left ideal. Then there exists a minimal left ideal  $\Lambda x$  such that  $\Lambda/I \cong \Lambda x$  and  $E(\Lambda x) \subseteq \Lambda$ . Let  $y \in r(I)$ , then  $\Lambda y \cong \Lambda/I \cong \Lambda x$ . Hence there exists  $z \in E(\Lambda x)$  such that  $yz = x$ , since  $E(\Lambda x)$  is injective. Therefore  $y\Lambda \supseteq x\Lambda$  for any  $y \in r(I)$ , and we see  $x\Lambda$  is minimal and  $r(I)$  is an essential extension of  $x\Lambda$ . Let  $y$  be an arbitrary element of  $\Lambda$  and  $I$  a maximal left ideal such that  $I \supseteq l(y)$ . Then there is a  $\Lambda$ -homomorphism  $\Lambda y \cong \Lambda/I \rightarrow \Lambda/I \cong \Lambda x$ , where  $E(\Lambda x) \subseteq \Lambda$ . Then  $yz = x$  for some  $z \in E(\Lambda x)$ , and  $x\Lambda \subseteq y\Lambda$ . Since  $x \in r(I)$ ,  $x\Lambda$ , consequently,  $y\Lambda$  contains a minimal right ideal. Thus every right ideal contains a minimal right ideal. If  $\Lambda$  is right lower distinguished,  $l(x) \neq 0$  for every maximal right ideal. Let  $I$  be an arbitrary left ideal. Then  $I = l(r(I)) \supseteq l(x)$ , where  $r$  is a maximal right ideal such that  $r \supseteq r(I)$ . Therefore  $I$  contains a minimal left ideal.

**Proposition 1.** *Let  $\Lambda$  be a left self cogenerator ring and  $\{E_\alpha\}, \{I_\alpha\}$  be as before. If  $E_\alpha = \Lambda e_\alpha$ , then  $e_\alpha \Lambda$  is an essential extension of a minimal right ideal  $x_\alpha \Lambda$ .  $x_\alpha \Lambda$  and  $x_\beta \Lambda$  are not isomorphic for any  $e_\alpha \neq e_\beta$ .*

Proof.  $l(e_\alpha) = \Lambda(1 - e_\alpha)$ . By Lemma 3 [3]  $\Lambda(1 - e_\alpha) \oplus Ne_\alpha$  is a maximal ideal. Hence  $\Lambda/\Lambda(1 - e_\alpha) \oplus Ne_\alpha \cong \Lambda e_\alpha/Ne_\alpha \cong \Lambda x_\alpha$  where  $E(\Lambda x_\alpha) \subseteq \Lambda$ . Then the same argument as the proof of Lemma 3 shows that  $x_\alpha \Lambda$  is a minimal right ideal and  $e_\alpha \Lambda$  is an essential extension of  $x_\alpha \Lambda$ . If  $x_\alpha \Lambda \cong x_\beta \Lambda$ , then  $r(x_\alpha) = r(x_\beta z)$  for some  $z \in \Lambda$ . Then  $\Lambda x_\alpha = l(r(\Lambda x_\alpha)) = l(r(\Lambda x_\beta z)) = \Lambda x_\beta z \cong \Lambda x_\beta$ , since  $\Lambda x_\beta \cong \Lambda e_\beta/Ne_\beta$  is minimal. This is a contradiction since  $\Lambda e_\alpha/Ne_\alpha \not\cong \Lambda e_\beta/Ne_\beta$ . Hence  $e_\alpha \Lambda \not\cong e_\beta \Lambda$ .

The following lemma is well known.

**Lemma 4.** *Let  $P_1, P_2$  be finitely generated projective module. Then  $P_1 \cong P_2$  if and only if  $P_1/NP_1 \cong P_2/NP_2$ .*

**Proposition 2.** *If  $\Lambda/N$  is a semi-simple ring where  $N$  is the radical of  $\Lambda$ ,*

and  $\Lambda$  is a left self cogenerator ring, then  $\Lambda$  is a finite direct sum of injective hulls of irreducible left ideals. Every commutative self cogenerator ring is self-injective and a direct sum of a finite number of injective hulls of irreducible ideals such that any two of which are not isomorphic.

Proof. If  $\Lambda/N$  is semisimple,  $\Lambda$  has only a finite number of non isomorphic irreducible modules, say  $I_1, I_2, \dots, I_n$ . Let  $E_i (i=1, \dots, n)$  be their injective hulls. Then  $E_i/NE_i$ 's are all the non isomorphic irreducible modules by Lemma 3 [3] and Lemma 4. Since  $\Lambda/N \cong I_1^{\alpha_1} + I_2^{\alpha_2} + \dots + I_n^{\alpha_n} \cong (E_1/NE_1)^{\alpha_1} + (E_2/NE_2)^{\alpha_2} + \dots + (E_n/NE_n)^{\alpha_n}$ ,  $\Lambda \cong E_1^{\alpha_1} + E_2^{\alpha_2} + \dots + E_n^{\alpha_n}$  by Lemma 4. Let  $R$  be a commutative self cogenerator ring. Then  $R \supseteq \sum^{\oplus} E_{\alpha}$ . Since  $\sum^{\oplus} E_{\alpha}$  is faithful  $\sum^{\oplus} E_{\alpha} = l(r(\sum^{\oplus} E_{\alpha})) = l(0) = R$ . Then the number of  $E_{\alpha}$ 's is finite and any two of which are not isomorphic.

Now we consider following two properties of a ring  $\Lambda$ .

- (\*) Every faithful left module is a generator
- (\*\*) Every faithful left module is a cogenerator.

Then we have

**Theorem 1.**  $\Lambda$  has Property (\*) if and only if  $\Lambda/N$  is semisimple and  $\Lambda$  has Property (\*\*). In case  $\Lambda$  is commutative Property (\*) and Property (\*\*) are equivalent.

Proof. By Theorem 6 G. Azumaya [1] we see that a ring  $\Lambda$  has Property (\*) if and only if  $\Lambda$  is left self injective and a finite direct sum of indecomposable left ideals each of which is an essential extension of a minimal left ideal. In this case  $\Lambda/N$  is a semisimple ring (see Theorem 7 [7].) Hence the proof is straightforward by Proposition 2 and Lemma 2.

REMARK 1. Proposition 1 means that a left self cogenerator ring contains at least the same number of the isomorphism types of irreducible right ideals as that of irreducible left ideals. Hence if in addition  $\Lambda/N$  is a semisimple ring  $\Lambda$  is right lower distinguished, since  $\Lambda/N$ , consequently  $\Lambda$  has the same number of isomorphism types of irreducible right ideals as that of irreducible left ideals (see Theorem 7 [1]).

REMARK 2. Lemma 3 [3] can be stated more generally as follows.

**Lemma 5.** If  $M$  is a projective, injective and indecomposable  $\Lambda$ -module, then  $M/NM$  is an irreducible  $\Lambda$ -module, and  $M$  is isomorphic to a principal left ideal of  $\Lambda$ .

Proof. We need only to prove that  $M \cong \Lambda e, e^2 = e$ , and  $\Omega = \text{Hom}(\Lambda e, \Lambda e)$  is a local ring. Since  $M$  is  $\Lambda$ -projective,  $M$  is contained in a free  $\Lambda$ -module,  $F = \sum \Lambda u_{\alpha}$ . Let  $m$  be a non zero element of  $M$  and  $m = \sum_{i=1}^m \lambda_i u_{\alpha_i}, \lambda_i \in \Lambda$ .

If  $\lambda \lambda_i = 0$  implies  $\lambda m = 0$  for any  $\lambda \in \Lambda$  and some  $i$ , then  $\Lambda m \cong \Lambda \lambda_i \subseteq \Lambda$ . If we consider the projection  $\pi_{\alpha_i}$  of  $F$  to  $\Lambda$ , then  $M \cong \pi_{\alpha_i}(M)$  since  $M$  is an essential extension of  $\Lambda m$ . If  $\lambda m \neq 0$  and  $\lambda \lambda_i = 0$  for some  $\lambda \in \Lambda$ , then replacing  $m$  by  $\lambda m$  we may assume  $m = \sum_{i=2}^n \lambda_i u_{\alpha_i}$ . Repeating this argument we can show as above that  $M \cong \Lambda e$ ,  $e^2 = e$ . Since  $\Lambda e$  is an essential extension of every subideal, we can easily show that  $\text{Hom}(\Lambda e, \Lambda e) \cong e \Lambda e$  is a local ring. Then the same proof as Lemma 3 [3] shows that  $M/NM$  is irreducible.

The author rewrote this paper after Osofsky's paper appeared, and he gives thanks to M. Harada for helpful suggestions.

OSAKA CITY UNIVERSITY

### References

- [1] G. Azumaya: *Completely faithful modules and self-injective rings*, Nagoya Math. J. **27** (1966), 697–708.
- [2] E. Matlis: *Injective modules over Noetherian rings*, Pacific J. Math. **8** (1958), 511–528.
- [3] B.L. Osofsky: *A generalization of quasi-Frobenius rings*, J. Algebra **4** (1966), 373–387.