



Title	Flexible boundaries in deformations of hyperbolic 3-manifolds
Author(s)	Fujii, Michihiko; Kojima, Sadayoshi
Citation	Osaka Journal of Mathematics. 1997, 34(3), p. 541-551
Version Type	VoR
URL	<a href="https://doi.org/10.18910/3840">https://doi.org/10.18910/3840</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## FLEXIBLE BOUNDARIES IN DEFORMATIONS OF HYPERBOLIC 3-MANIFOLDS

MICHIHIKO FUJII and SADAYOSHI KOJIMA

(Received March 11, 1996)

### 0. Introduction

Let  $M$  be a cusped hyperbolic 3-manifold with non-empty geodesic boundary. A small Dehn filling deformation of  $M$  on the cusps can be performed so that the boundary is kept to be geodesic. Then assigning to each deformation a hyperbolic structure on the boundary, we get a map  $B_M$  from the space of such deformations to the Teichmüller space of  $\partial M$ . More precise argument for this fact will be given in §1.

Motivated by the conjectures posed in Cooper-Long [1] and Kapovich [5], Neumann and Reid [7] discovered many examples of  $M$  such that  $B_M$  is a constant map. Fujii [3] also obtained another concrete family of small deformations for some  $M$  such that  $B_M$  maps his family to a constant structure. These examples at first contrasted with our naive intuition that the deformation of hyperbolic structure affects everywhere. But the fact itself would not be too surprising once we realized that the dimension of the source can be bigger than that of the target and in that case  $B_M$  can never be injective. A more pertinent problem to set up for the moment would be what the map  $B_M$  looks like. In fact, we have known only a little about it so far.

Under the circumstances above, it would be worth finding examples for which we can convert this rather difficult problem to something we can do by hand. In this paper, we will construct infinitely many one-cusped examples of  $M$  so that  $B_M$  is a local embedding at the complete structure. The polyhedral construction will be discussed in §2. Then by using its polyhedral structure, we will compute the derivative of  $B_M$  at the complete structure by hand in the later sections.

The authors are indebted to Professor T. Soma for the argument in §1, and we would like to thank him.

### 1. The map $B_M$

We briefly review Dehn filling deformations of cusped hyperbolic 3-manifolds. Let  $N$  be a noncompact, orientable, complete hyperbolic 3-manifold of finite volume, and  $\bar{\rho}_0 : \pi_1(N) \rightarrow \mathrm{PSL}_2(\mathbb{C})$  its holonomy representation. According to

Thurston [9] (cf. [2]),  $\bar{\rho}_0$  has a lift  $\rho_0 : \pi_1(N) \rightarrow \mathrm{SL}_2(\mathbb{C})$ . Since  $\mathrm{SL}_2(\mathbb{C})$  is an algebraic set, the space of representations  $\mathrm{Hom}(\pi_1(N), \mathrm{SL}_2(\mathbb{C}))$  is also an algebraic set. To each representation  $\rho$ , associated is its character  $\chi_\rho$ . Culler and Shalen [2] showed that the irreducible component of  $\mathrm{Hom}(\pi_1(N), \mathrm{SL}_2(\mathbb{C}))$  containing  $\rho_0$  is mapped by this correspondence onto the space of characters  $X$ , which is an affine variety. The preimage of a character  $\chi_\rho$  near  $\chi_{\rho_0}$  consists of representations conjugate to  $\rho$ . Thus a small neighborhood of  $\chi_{\rho_0}$  in  $X$  is bijectively identified with the set of conjugacy classes of  $\mathrm{SL}_2(\mathbb{C})$ -representations near the conjugacy class of  $\rho_0$ .

This small neighborhood is also identified with the set of conjugacy classes of  $\mathrm{PSL}_2(\mathbb{C})$ -representations near the conjugacy class of  $\bar{\rho}_0$ . This is because any path in  $\mathrm{Hom}(\pi_1(N), \mathrm{PSL}_2(\mathbb{C}))$  based at  $\bar{\rho}_0$  lifts to a path in  $\mathrm{Hom}(\pi_1(N), \mathrm{SL}_2(\mathbb{C}))$  based at  $\rho_0$  and hence the covering projection  $\mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{PSL}_2(\mathbb{C})$  induces a local homeomorphism of  $X$  at  $\chi_{\rho_0}$ . Through this correspondence, we will identify the set of equivalence classes of  $\mathrm{PSL}_2(\mathbb{C})$ -representations near the conjugacy class of  $\bar{\rho}_0$  with a small neighborhood of  $\chi_{\rho_0}$  in  $X$ .

It has been known by Thurston [9] and Neumann-Zagier [8] that the complex dimension of  $X$  is equal to the number of cusps of  $N$  and that the character of  $\rho_0$  is a smooth point. If we choose a set of meridional elements  $\{m_j\}$  for all cusps of  $N$ , then the traces of these elements turn out to be a local coordinate of  $X$  at the conjugacy class of  $\rho_0$ .

Thurston originally introduced another parameter which is the set of complex lengths of the  $\rho(m_j)$ 's. The complex length of  $\rho(m_j)$  is well-defined with sign by orienting the axis of  $\rho(m_j)$ . Their squares turn out to be a local coordinate at  $\chi_{\rho_0}$  in  $X$ . To each representation  $\rho$  near  $\rho_0$ , he assigned a hyperbolic manifold  $N_\rho$ , see [9]. It can be interpreted as a small deformation of  $N$  and is called a Dehn filling deformation of  $N$ . The topological type of the deformation may be different from the original. Thurston's parameter can be converted to the set of pairs of real numbers which is a generalization of Dehn surgery coefficients in the classical knot theory.

Now, suppose that  $M$  is an orientable complete hyperbolic 3-manifold of finite volume with both cusps and compact geodesic boundaries. Let  $DM$  be the double of  $M$  along the boundary and  $\rho_0$  be a holonomy representation of  $DM$ . The manifold  $DM$  admits an obvious involution  $\tau$  switching the sides.

**DEFINITION.** Fix a set of meridians  $m_j$  closed under  $\tau$ , and choose a small neighborhood  $U$  of  $\chi_{\rho_0}$  so that the traces of the  $m_j$ 's become a local coordinate of the space of characters at  $\chi_{\rho_0}$ . Let  $\mathcal{D}_M$  be a diagonal set in  $U$  fixed by an involution on  $U$  which is induced by  $\tau$ . It is a smooth submanifold of real dimension  $= \#\{\text{cusps of } DM\}$ .

**Lemma 1.** *The restriction of a representation  $\rho$  near  $\rho_0$  whose conjugacy class*

is in  $\mathcal{D}_M$  to  $\pi_1(\partial_0 M)$  is fuchsian (i.e., a discrete faithful representation to  $\mathrm{SL}_2(\mathbf{R})$ ), where  $\partial_0 M$  is a component of the boundary  $\partial M$ .

**Proof.** Choose  $\rho$  as above, and denote the associated Dehn filling deformation by  $DM_\rho$ . There is also a topological involution  $\tau_\rho$  on  $DM_\rho$  switching sides. Then since  $\rho$  is close to  $\rho_0$ ,  $\tau_\rho$  induces a nearby deformation of  $DM$  and the induced structure is the same as  $DM_\rho$  by the local parametrization around the complete structure  $\rho_0$  (see [8] [9]). Hence  $\tau_\rho$  can be deformed to an isometry  $T_\rho$  by a tiny isotopy. We want to show that  $T_\rho$  is an involution fixing a geodesic surface isotopic to  $\partial M$ .

There is a dense subset  $Y$  in  $\mathcal{D}_M$  near  $[\rho_0]$ , corresponding to a set of rational rays in the Dehn filling coefficient space, so that the deformation  $DM_y$  for  $y \in Y$  is a cone manifold with cone angle more than  $2\pi$ . Then by changing the metric near the cusps as in [4],  $DM_y$  can be modified to a compact negatively curved manifold  $\overline{DM}_y$  with obvious involutive symmetry so that  $T_y$  is also modified near the cusps to an isometry  $\overline{T}_y$  of  $\overline{DM}_y$ . Then  $\overline{T}_y^2$  is an isometry on the compact negatively curved manifold which is isotopic to the identity. Thus  $\overline{T}_y^2$  must be the identity and  $\overline{T}_y$  is an isometric involution. Since  $\overline{T}_y$  and  $T_y$  are the same map away from neighborhoods of the cusps, the fact implies that  $T_y$  itself is an involution.

Since  $T_y$  is deformed to  $\tau_y$  by a tiny isotopy,  $\partial M$  admits an equivariant collar neighborhood with respect to  $\tau_y$  which separates  $DM_y$  into two parts. We may choose one of them  $A$  such that  $T_y(A) \cap A = \emptyset$  and  $DM_y - (T_y(A) \cup A)$  is homeomorphic to  $\partial M \times I$ . This implies by Kim and Tollefson [6] that  $T_y$  fixes a surface isotopic to  $\partial M$ . Since  $T_y$  is an isometry, the fixed surface must be totally geodesic.

Let  $\rho_y$  be a corresponding representation. Then since  $\partial M$  is realized as the geodesic surface, the restriction of  $\rho_y$  to  $\pi_1(\partial_0 M)$  is fuchsian. Since  $Y$  is dense in  $\mathcal{D}_M$ , and since the set of fuchsian representations is closed, we are done.  $\square$

Assigning the hyperbolic structure of the boundary to such a deformation  $DM_\rho$  where  $\rho \in \mathcal{D}_M$ , we get a map

$$B_M : \mathcal{D}_M \rightarrow \mathcal{T}(\partial M)$$

where  $\mathcal{T}$  is the Teichmüller space of  $\partial M$ .

## 2. Examples

Consider the Whitehead link  $L = K_1 \cup K_2$  in  $S^3$ . Removing a thin tubular neighborhood of  $K_2$  from the complement of  $L$ , we obtain a manifold  $W$  with one compact toral boundary and one toral end. Choose an arc  $\Sigma$  connecting two points on  $\partial W$  as in Figure 1.




Fig. 1.

To give hyperbolic orbifold structures  $O_n$  on  $W$  with the singular set  $\Sigma$  indexed by natural numbers  $n \geq 2$ , we recall the fact, for instance in [9], that the regular ideal octahedron is a fundamental domain of the hyperbolic manifold homeomorphic to the Whitehead link complement. Replace the regular ideal octahedron by the truncated octahedron as in Figure 2, where the dihedral angle along each edge connecting truncated faces is  $\pi/2n$  and that of each edge through  $\infty$  is  $\pi/2$ . Then the faces topologically identified to create the Whitehead link complement are still isometric and the identification gives a hyperbolic orbifold  $O_n$  whose underlying space is  $W$  where the singular set is  $\Sigma$  with rotation angle  $2\pi/n$ .

Decompose the truncated octahedron into 4 truncated tetrahedra and we will deform them to obtain deformations of  $O_n$  in the next section. The edge lengths of the truncated tetrahedra determine their shapes. As we will see later, they are subject to the relations required by the gluing consistency to give nonsingular hyperbolic orbifold structures with geodesic boundary on  $W$ . Thus we obtain a space  $\mathcal{W}$  of the deformations of  $O_n$  parameterized by the edge lengths of the truncated tetrahedra.

Let  $\pi_1^{\text{orb}}(W)$  be the fundamental group of the orbifold  $O_n$  and  $\rho_0$  a lift of the holonomy representation of  $\pi_1^{\text{orb}}(W)$  in  $\text{SL}_2(\mathbb{C})$ . Take a meridional element  $m$  for the cusp  $K_1$ . To each deformation of  $O_n$  represented by an element  $w \in \mathcal{W}$ , we have two data. One is a representation  $\rho$  near  $\rho_0$  and the other is a canonical direction of the axis of  $\rho(m)$  to which the end of truncated tetrahedra spiral. Assigning to each  $w \in \mathcal{W}$  the complex length of  $\rho(m)$  with respect to this orientation, we get a map  $G : \mathcal{W} \rightarrow \mathbb{C}$ . It will be shown in Lemma 3 that  $G$  is a local diffeomorphism at  $O_n$ .

On the other hand, each representation corresponds to a pair of complex lengths




Fig. 2.

which differ only in sign. Hence by assigning to each  $w \in \mathcal{W}$  the character  $\chi_\rho$  of  $\rho$ , we get a map  $\pi$  of  $\mathcal{W}$  to the space of characters of the representations of  $\pi_1^{\text{orb}}(W)$  in  $\text{SL}_2(\mathbb{C})$ . The space of characters at  $\chi_{\rho_0}$  has complex dimension  $> 1$  since  $W$  has a boundary with negative Euler characteristic, and  $\pi$  doubly covers the image  $\mathcal{D}_{O_n} = \pi(\mathcal{W})$  of complex dimension  $= 1$  branched at  $\chi_{\rho_0}$ .

The orbifold  $O_n$  has a toral boundary with two cone points of rotation angle  $2\pi/n$ . Let  $\mathcal{T}(\partial O_n)$  be the Teichmüller space of the orbifold  $\partial O_n$ . It is homeomorphic to  $\mathbb{R}^4$ . Assigning the hyperbolic structure of the boundary to each  $\chi_\rho$ , we get a map

$$B_{O_n} : \mathcal{D}_{O_n} \rightarrow \mathcal{T}(\partial O_n).$$

Our goal is

**Theorem.** *The derivative of the map  $B_{O_n}$  at  $\chi_{\rho_0}$  has rank 2.*

Now let  $P_n$  be an  $n$ -fold cyclic covering of  $W$  branched along  $\Sigma$ . The manifold  $P_n$  supports a complete hyperbolic structure  $M_n$  which covers  $O_n$ . Pulling back the hyperbolic structure of the geodesic boundary associated to each  $w \in \mathcal{W}$  by the covering  $P_n \rightarrow W$ , we get an embedding of  $\mathcal{T}(\partial O_n)$  into  $\mathcal{T}(\partial M_n)$ . Also the covering induces an injective homomorphism  $\pi_1(P_n) \rightarrow \pi_1^{\text{orb}}(W)$ , and hence a map  $\mathcal{D}_{O_n} \rightarrow \mathcal{D}_{M_n}$  on the characters by the restriction. It is a local diffeomorphism onto the image at  $\chi_{\rho_0}$  because  $m^n$  lies in  $\pi_1(P_n)$ , and actually a local diffeomorphism




Fig. 3.

by dimension count. Therefore, by Theorem,  $B_{M_n} : \mathcal{D}_{M_n} \rightarrow \mathcal{T}(\partial M_n)$  is a local embedding at  $M_n$ . Varying  $n$ , we have

**Corollary.** *There are infinitely many hyperbolic 3-manifolds  $M$  with both a cusp and a boundary such that the map  $B_M : \mathcal{D}_M \rightarrow \mathcal{T}(\partial M)$  is a local embedding near the complete structure.*

### 3. Truncated Tetrahedra and Gluing Consistency

The truncated octahedron to create  $O_n$  is decomposed into four congruent truncated tetrahedra as in Figure 3. We will parametrize the deformations of  $O_n$  in terms of the deformations of the shapes of these blocks.

First of all, label the triangular faces by  $A$ ,  $B$ , and their edges by  $A_i$ ,  $B_i$  ( $i = 1, 2, 3$ ) as in Figure 4. We call each of these edges *an external edge*, and denote the length of  $A_j$  and  $B_j$  by  $a_j$  and  $b_j$  respectively. These lengths are subject to two equations.

One is the following. If we let  $l$  be the length of the edge shared by two pentagonal faces, then regarding it as the bottom of the left pentagon, we obtain an expression of  $l$  in terms of  $a_1$  and  $b_2$ ,

$$\cosh l = \frac{\cosh a_1 \cosh b_2 + 1}{\sinh a_1 \sinh b_2}.$$

Simultaneously, if we regard it as the bottom of the right pentagon, we obtain an expression of  $l$  in terms of  $a_2$  and  $b_1$ ,

$$\cosh l = \frac{\cosh a_2 \cosh b_1 + 1}{\sinh a_2 \sinh b_1}.$$




Fig. 4.

Then since these two are the same quantity, we obtain one identity involving edge lengths, which we call the relation of type (1).

The other is concerned with angles. By the hyperbolic cosine rule for the top triangle, we have

$$\cos \theta_{\text{top}} = \frac{\cosh a_1 \cosh a_2 - \cosh a_3}{\sinh a_1 \sinh a_2},$$

where  $\theta_{\text{top}}$  is the angle between  $A_1$  and  $A_2$ . If we look at the bottom triangle, then the corresponding angle  $\theta_{\text{bottom}}$  has an expression in terms of the  $b_j$ 's. They represent the same dihedral angle, and we obtain another relation,

$$(2) \quad \theta_{\text{top}} = \theta_{\text{bottom}}.$$

It is not hard to verify that the set of six length variables subject to the relations (1) and (2) parametrizes isometry classes of labelled truncated tetrahedra.

To create a nonsingular but not necessarily complete hyperbolic orbifold structure on  $W$ , it is sufficient to verify gluing consistency which consists of the isometricity conditions for faces to be identified, and the cone angle conditions along edges. We will see when these are satisfied.

If the external edges to be identified have the same length, then the isometricity condition for face identification is satisfied. Since there are twelve such pairs, there are twelve simple identities in  $\{a_j, b_j\}$  we must obviously require. For simplicity, we just assign the same variable to each pair to be identified from the beginning and reduce the number of the variables to the half.

Then the relations of type (1) and (2) for the four truncated tetrahedra become dependent after gluing. In fact, reading off the lengths of the bottom edges of the pentagonal faces in order, we can see that one of the four equations of type (1) becomes a consequence of the other three.



To compute the cone angle conditions along edges, we label them by  $E_1, E_2, E_3$  and  $\Sigma$  as in Figure 1. The dihedral angle of each edge is described in terms of the lengths of external edges as the above expression of  $\theta_{\text{top}}$ . To obtain a nonsingular orbifold structure, the total sum of dihedral angles around the first three edges must be  $2\pi$  and the last  $2\pi/n$ . These constraints give four identities. The last one is independent from the others, however one of the first three identities is a consequence of the other two. To see this, recall that a toral section of the end always admits a similarity structure. Then the total sum of angles of triangles appeared in the horospherical triangulation is  $4 \times 2\pi$ . It is equal to the sum of the total sum of dihedral angles along  $E_1$  and  $E_2$  and the double of that of  $E_3$ .

We thus have obtained ten relations with twelve variables from gluing consistency. These relations define a map

$$f : \mathbf{R}^{12} \longrightarrow \mathbf{R}^{10},$$

such that its zero set  $f^{-1}(0)$  consists of the points in  $\mathbf{R}^{12}$  satisfying the gluing consistency.

Let  $w_0 \in \mathbf{R}^{12}$  be the point corresponding to the complete hyperbolic structure. Denote by  $x$  and  $y$  the two variables indicated in Figure 2, and by  $z_1, \dots, z_{10}$  the other 10 variables. It is not hard to see that the rank of the matrix  $(\partial f_i / \partial z_j(w_0))$  is 10 and also find the unique solutions of the following two linear equations in terms of  $u$  and  $v$  respectively:

$$\begin{aligned} \left( \frac{\partial f_i}{\partial z_j}(w_0) \right) u &= - \left( \frac{\partial f_i}{\partial x}(w_0) \right), \\ \left( \frac{\partial f_i}{\partial z_j}(w_0) \right) v &= - \left( \frac{\partial f_i}{\partial y}(w_0) \right). \end{aligned}$$

Denote the unique solutions by  $u_0$  and  $v_0$  respectively. Then we have obtained

**Lemma 2.** *A neighborhood  $\mathcal{W}$  of  $w_0$  in  $f^{-1}(0)$  is a 2-dimensional smooth manifold and we have two paths on  $\mathcal{W} \subset \mathbf{R}^{12}$ ,*

$$\begin{aligned} \xi(t) &= w_0 + xt + (\text{higher order}), \\ \eta(t) &= w_0 + yt + (\text{higher order}), \end{aligned}$$

such that

$$x = \begin{pmatrix} 1 \\ 0 \\ u_0 \end{pmatrix}, \quad y = \begin{pmatrix} 0 \\ 1 \\ v_0 \end{pmatrix},$$

where the  $y$ -component of  $\xi(t)$  and the  $x$ -component of  $\eta(t)$  are constant, and the  $x$ -component of  $\xi(t)$  and the  $y$ -component of  $\eta(t)$  have no terms of degrees  $n$  ( $n \geq 2$ ).




Fig. 5.

#### 4. Dehn Filling Space and Computation

**Lemma 3.** *The map  $G : \mathcal{W} \rightarrow \mathbb{C}$  defined in §2 is a local diffeomorphism at  $w_0 \in \mathcal{W}$ .*

*Proof.* The complex length of a meridional element can be read off from the triangulation by horospherical sections as in [8]. By direct computation using triangulation induced from the polyhedral decomposition of  $W$ , we can verify that the rank of the Jacobian of  $G$  is 2.  $\square$

Recall that  $\pi : \mathcal{W} \rightarrow \mathcal{D}_{O_n}$  is a 2-fold covering branched at  $\pi(w_0)$ . In fact  $\pi$  is a map which locally looks like a square function  $z \rightarrow z^2$ .

The lengths  $L_i$  of geodesic segments  $S_i$  ( $i = 1, \dots, 4$ ) which are illustrated by thick lines in Figure 5 give rise to a quadruple  $(L_1, L_2, L_3, L_4)$  which defines a global coordinate of  $T(\partial O_n)$ .

Let  $\tilde{B}$  be a map assigning to the element of  $\mathcal{W}$  the corresponding hyperbolic structure of the boundary. Then its induced map from  $\mathcal{D}_{O_n}$  is  $B_{O_n}$ . Let  $\tilde{B}_i$  (resp.  $B_i$ ) be the composition of  $\tilde{B}$  (resp.  $B_{O_n}$ ) with  $L_i$ .

$$\begin{array}{ccccc}
 \mathcal{W} & & \tilde{B} & & \\
 \pi \downarrow & \searrow & & & \\
 \mathcal{D}_{O_n} & \xrightarrow{B_{O_n}} & T(\partial O_n) & \xrightarrow{(L_1, L_2, L_3, L_4)} & \mathbb{R}^4.
 \end{array}$$

Now consider a quadrilateral in general. If the lengths of four sides and one

of diagonals are known, then the length of the other diagonal can be expressed in terms of them by hyperbolic trigonometry. Applying this to the quadrilateral in Figure 5 which is made of two triangular faces, we have an expression of  $\tilde{B}_i$  as a function of our length parameters.

Because of the local picture of  $\pi$ , letting  $\bar{\xi}(t) = \pi \circ \xi(\sqrt{t})$  and  $\bar{\eta}(t) = \pi \circ \eta(\sqrt{t})$ , we obtain smooth paths on  $\mathcal{D}_{O_n}$  such that its tangent vectors

$$v = \frac{d}{dt} \bar{\xi}(t)|_{t=0},$$

$$w = \frac{d}{dt} \bar{\eta}(t)|_{t=0}$$

are nontrivial. The images of these vectors by the derivative  $dB_i$  are now expressed by

$$dB_i(v) = \left. \frac{dB_i(\bar{\xi}(t))}{dt} \right|_{t=0} = \left. \frac{d\tilde{B}_i(\xi(\sqrt{t}))}{dt} \right|_{t=0},$$

$$dB_i(w) = \left. \frac{dB_i(\bar{\eta}(t))}{dt} \right|_{t=0} = \left. \frac{d\tilde{B}_i(\eta(\sqrt{t}))}{dt} \right|_{t=0}.$$

To carry out the actual computation of the right hand sides, we used the Taylor expansions of  $\xi(t)$  and  $\eta(t)$  up to the second degree, which can be derived from the formula,

$$\frac{d^2 f_i(\xi)}{dt^2}(0) = \sum_{j,k} \frac{\partial^2 f_i}{\partial z_j \partial z_k}(\mathbf{w}_0) \frac{d\xi_k}{dt}(0) \frac{d\xi_j}{dt}(0) + \sum_j \frac{\partial f_i}{\partial z_j}(\mathbf{w}_0) \frac{d^2 \xi_j}{dt^2}(0).$$

By performing this rather lengthy but direct computations by hand, we verified the following:

**Lemma 4.**

$$dB_1(v) = -\frac{1}{\sqrt{c}} \quad (< 0), \quad dB_2(v) = \frac{1}{\sqrt{c}} \quad (> 0),$$

$$dB_1(w) = \frac{s^2(1-c)}{4\sqrt{c}} \quad (> 0), \quad dB_2(w) = \frac{s+1}{8s\sqrt{c}} \quad (> 0),$$

where  $c = \cos \frac{\pi}{2n}$  and  $s = \sin \frac{\pi}{2n}$ .

Lemma 4 shows that these tangent vectors on  $\mathcal{D}_{O_n}$  go to a linearly independent pair in the tangent space of  $\mathcal{T}(\partial O_n)$  at the original structure and we complete the proof of Theorem.

Finally, we would like to mention that our naive method works equally well to verify the vanishing of the derivative of  $B_M$  for Neumann and Reid's example. The details were worked out by T. Hayakawa at Tokyo Institute of Technology.

---

### References

- [1] D. Cooper and D. Long: *An undetected slope in a knot manifold*, In : Topology '90, Proceedings of the Research Semester in Low Dimensional Topology at Ohio State University, 111–121, Berlin New York : de Gruyter 1992.
- [2] M. Culler and P.B. Shalen: *Varieties of group representations and splittings of 3-manifolds*, Ann. Math. **117** (1983), 109–146.
- [3] M. Fujii: *Deformations of a hyperbolic 3-manifold not affecting its totally geodesic boundary*, Kodai Math. J. **16** (1993), 441–454.
- [4] M. Gromov and W.P. Thurston: *Pinching constants for hyperbolic manifolds*, Invent. Math. **89** (1987), 1–12.
- [5] M. Kapovich: *Eisenstein series and Dehn surgery*. (Preprint)
- [6] P.K. Kim and J.L. Tollefson: *PL involutions of fibered 3-manifolds*, Trans. A.M.S. **232** (1977), 221–237.
- [7] W.D. Neumann and A.W. Reid: *Rigidity of cusps in deformations of hyperbolic 3-orbifolds*, Math. Ann. **295** (1993), 223–237.
- [8] W.D. Neumann and D. Zagier: *Volumes of hyperbolic 3-manifolds*, Topology **24** (1985), 307–332.
- [9] W.P. Thurston: *The geometry and topology of 3-manifolds*, Lect. Notes, Princeton Univ., 1978.

M. Fujii  
 Department of Mathematical Sciences  
 Yokohama City University  
 22-2 Seto, Kanazawa-ku, Yokohama  
 Kanagawa-ken 236, Japan  
 e-mail: mfujii@yokohama-cu.ac.jp

S. Kojima  
 Department of Mathematical  
 and Computing Sciences  
 Tokyo Institute of Technology  
 Ohokayama, Meguro, Tokyo 152, Japan  
 e-mail: sadayosi@is.titech.ac.jp

