



Title	On the Robertello invariants of proper links
Author(s)	Kawauchi, Akio
Citation	Osaka Journal of Mathematics. 1984, 21(1), p. 81-90
Version Type	VoR
URL	<a href="https://doi.org/10.18910/4202">https://doi.org/10.18910/4202</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Kawauchi, A.  
Osaka J. Math.  
21 (1984), 81-90

## ON THE ROBERTELLO INVARIANTS OF PROPER LINKS

Dedicated to Professor Minoru Nakaoka on his 60th birthday

AKIO KAWAUCHI

(Received September 14, 1982)

Robertello's invariant of a classical knot in [9] was generalized by Gordon in [2] to an invariant of a knot in a  $Z$ -homology 3-sphere, and by the author in [5] to an invariant,  $\delta(k \subset S)$ , of a knot  $k$  in a  $Z_2$ -homology 3-sphere  $S$ . In this paper, we shall generalize this invariant to two mutually related invariants,  $\delta_0(L \subset S)$  and  $\delta(L \subset S)$ , of a proper link  $L$  in a  $Z_2$ -homology 3-sphere  $S$ . In the case of a classical proper link, this  $\delta_0$ -invariant can be considered as an invariant suggested by Robertello in [9, Theorem 2]. A difference between  $\delta_0(L \subset S)$  and  $\delta(L \subset S)$  is that  $\delta_0(L \subset S)$  is generally an oriented link type invariant, but  $\delta(L \subset S)$  is an unoriented link type invariant. A proper link in a  $Z_2$ -homology 3-sphere (which is not a  $Z$ -homology 3-sphere) naturally occurs when considering a branched cyclic covering of a 3-sphere, branched along a certain proper link. (If the number of the components of the link is  $\geq 2$ , the branched covering space can not be a  $Z$ -homology 3-sphere by the Smith theory.) So, we consider a proper link  $\tilde{L}$  in a  $Z_2$ -homology 3-sphere  $\tilde{S}$ , obtained from a proper link  $L$  in a  $Z_2$ -homology 3-sphere  $S$  by taking a branched cyclic covering, branched along  $L$ . When the covering degree is prime, we shall establish a relationship between  $\delta(\tilde{L} \subset \tilde{S})$  and  $\delta(L \subset S)$  and then a relationship between  $\delta_0(\tilde{L} \subset \tilde{S})$  and  $\delta_0(L \subset S)$ .

In Section 1 we define and discuss the slope of a link in a 3-manifold as a generalization of the slope of a knot in a 3-manifold, introduced in [5]. In Section 2 the  $\delta_0$ -invariant and the  $\delta$ -invariant are defined and studied. Section 3 deals with relationships between  $\delta(\tilde{L} \subset \tilde{S})$  and  $\delta(L \subset S)$  and between  $\delta_0(\tilde{L} \subset \tilde{S})$  and  $\delta_0(L \subset S)$ .

Throughout this paper spaces and maps will be considered in the piecewise linear category, and notations and conventions will be the same as those of [5] unless otherwise stated.

### 1. The slope of a link in a 3-manifold

Let  $M$  be a connected oriented 3-manifold. Let  $L$  be an oriented link with  $r$  components in the interior of  $M$ . Let  $o(L)$  denote the order ( $\geq 1$ ) of

the homology class  $[L] \in H_1(M; \mathbb{Z})$ . Let  $\tau H_1(M)$  be the torsion part of  $H_1(M; \mathbb{Z})$ . Let  $\phi: \tau H_1(M) \times \tau H_1(M) \rightarrow \mathbb{Q}/\mathbb{Z}$  be the linking pairing.

**DEFINITION 1.1.** The *slope* of the link  $L$ , denoted by  $s(L) = s(L \subset M)$  is defined by the identity

$$s(L) = \begin{cases} -\phi([L], [L]) & (o(L) < +\infty), \\ \infty & (o(L) = +\infty). \end{cases}$$

If  $s(L) = 0$ , then we say that the link  $L$  is *flat*.

When  $r = 1$ ,  $s(L)$  is the same as the slope defined in [5, Definition 1.4] by [5, Lemma 1.8]. Let  $r \geq 2$ . Let  $B_1, B_2, \dots, B_{r-1}$  be mutually disjoint oriented bands in the interior of  $M$  attaching to  $L$  as 1-handles. If we obtain a knot  $k$  from  $L$  by surgery along such  $B_1, B_2, \dots, B_{r-1}$ , then we say that the knot  $k$  is *obtained from  $L$  by a fusion*.

**Lemma 1.2.** *Let  $k$  be a knot obtained from a link  $L$  by a fusion. Then  $s(L) = s(k)$ .*

**Proof.** Clearly,  $[L] = [k]$  in  $H_1(M; \mathbb{Z})$ . The result follows from Definition 1.1.

Assume that each component  $k_i$  of  $L$  is a knot of finite order, i.e.,  $o(k_i) < +\infty$ ,  $i = 1, 2, \dots, r$ . Then the total  $\mathbb{Q}$ -linking number  $\lambda(L) = \lambda(L \subset M) \in \mathbb{Q}$  of the link  $L \subset M$  is defined by  $\lambda(L) = \sum_{i>j} \text{Link}_M(k_i, k_j)$ . When  $r = 1$ , we understand that  $\lambda(L) = 0$ .

**Lemma 1.3.** *In  $\mathbb{Q}/\mathbb{Z}$   $s(L) = \sum_{i=1}^r s(k_i) - 2\lambda(L)$ .*

**Proof.** Since  $o(L) < +\infty$  and  $[L] = \sum_{i=1}^r [k_i]$ ,  $s(L) = -\phi([L], [L]) = \sum_{i=1}^r -\phi([k_i], [k_i]) - 2 \sum_{i>j} \phi([k_i], [k_j])$ . Using that  $\phi([k_i], [k_j]) \equiv \text{Link}_M(k_i, k_j) \pmod{1}$  for  $i \neq j$  and  $s(k_i) = -\phi([k_i], [k_i])$ , we have a desired congruence.

For each element  $s \in \mathbb{Q}/\mathbb{Z}$  we can have coprime positive integers  $a, b$  such that  $s \equiv a/b \pmod{1}$ . This fraction  $a/b$  and the denominator  $b$  are called a *normal presentation* and the *denominator* of the element  $s \in \mathbb{Q}/\mathbb{Z}$ , respectively. Now we assume that the denominator of the slope  $s(k_i)$  is odd,  $i = 1, 2, \dots, r$ . Then  $s(k_i)$  has a normal presentation of type  $2a_i/b_i$ ,  $i = 1, 2, \dots, r$ .

**DEFINITION 1.4.** We define

$$s^*(L) = \sum_{i=1}^r a_i/b_i - \lambda(L)$$

in  $\mathbb{Q}/\mathbb{Z}$  and call it the *half-slope* of the link  $L \subset S$ .

The following is easily proved.

**Lemma 1.5.** *In  $Q/Z$   $2s^*(L)=s(L)$ , and if  $s(L)=0$ , then  $s^*(L)$  is 0 or  $1/2$  according as the denominator of  $\lambda(L) \in Q/Z$  is odd or even.*

## 2. The $\delta_0$ -invariant and the $\delta$ -invariant

We consider an oriented link  $L$  with components  $k_i$ ,  $i=1, 2, \dots, r$ , in an oriented  $Z_2$ -homology 3-sphere  $S$ .

**DEFINITION 2.1.** The link  $L$  is *proper* if the mod 2 linking number,  $\text{Link}_S(k_i, L-k_i)_2=0$  for all  $i$ ,  $1 \leq i \leq r$ . (We understand a knot to be a proper link.)

Let  $W$  be a compact oriented 4-manifold. Let  $F$  be a locally flat, oriented (possibly disconnected) surface of (total) genus 0 in  $W$ . We say that such a pair  $F \subset W$  is *admissible* for a link  $L \subset S$ , if  $S$  is a component of  $\partial W$ ,  $\partial F=L$ ,  $H_1(\partial W; Z_2)=0$  and  $[F_2^+] \in H_2(W; Z_2)$  is characteristic, i.e.,  $[F_2^+] \cdot x = x^2$  for all  $x \in H_2(W; Z_2)$ , where  $F_2^+$  is a (mod 2) cycle obtained from  $F$  by attaching (mod 2) 2-chains  $c_i$  in  $S$  with  $\partial c_i = -k_i$ ,  $i=1, 2, \dots, r$ .

**Lemma 2.2.** *For any proper link  $L \subset S$  there exists an admissible pair  $F \subset W$ .*

**Proof.** Let  $T(L)=\bigcup_{i=1}^r T(k_i)$  be a tubular neighborhood of  $L=\bigcup_{i=1}^r k_i$  in  $S$ . Construct a 4-manifold  $W=(-S) \times [-1, 1] \cup D^2 \times D_1^2 \cup \dots \cup D^2 \times D_r^2$  identifying  $T(k_i) \times 1$  with  $(\partial D^2) \times D_i^2$ ,  $i=1, \dots, r$ , so that  $H_1(\partial W; Z_2)=0$ . Let  $D_i=(-k_i) \times [-1, 1] \cup D^2 \times 0_i$  be a disk. Let  $F=\bigcup_{i=1}^r D_i$ . To show that  $F \subset W$  is admissible for  $L \subset S$ , it suffices to check that  $[F_2^+] \in H_2(W; Z_2)$  is characteristic. Note that  $[D_{i2}^+]$ ,  $i=1, \dots, r$ , form a basis for  $H_2(W; Z_2)$ . Since  $[F_2^+] = \sum_{i=1}^r [D_{i2}^+]$ , we have

$$\begin{aligned} [F_2^+] \cdot [D_{i2}^+] &= [D_{i2}^+]^2 + \sum_{j \neq i} [D_{j2}^+] \cdot [D_{i2}^+] \\ &= [D_{i2}^+]^2 + \text{Link}_S(L-k_i, k_i)_2 \\ &= [D_{i2}^+]^2, \quad i=1, \dots, r. \end{aligned}$$

This implies that  $[F_2^+]$  is characteristic. This completes the proof.

The pair  $F \subset W$ , constructed in the proof of Lemma 2.2 is called a *standard admissible pair* for the proper link  $L \subset S$ .

**DEFINITION 2.3.** Let  $L \subset S$  be a proper link. Then we define

$$\delta_0(L) = \delta_0(L \subset S) = ([F_2^+]^2 - \text{sign } W)/16 - \mu(\partial W)$$

in  $Q/Z$  for any admissible pair  $F \subset W$  for  $L \subset S$ , where  $F_2^+$  is a rational 2-cycle obtained from  $F$  by attaching rational 2-chains  $c_i^q$  in  $S$  with  $\partial c_i^q = -k_i$ ,  $i=1, \dots, r$ .

**REMARK 2.4.** We can define the invariant  $\delta_0(L \subset S)$  by using a more gen-

eral pair  $F \subset W$ , where the (total) genus of  $F$  may be positive or  $F$  may be non-orientable (cf. Freedman-Kirby [1], Guillou-Marin [3], Matsumoto [7]).

To see the well-definedness of  $\delta_0(L)$ , consider a standard admissible pair  $F^* \subset W^*$  for  $L \subset S$ . Construct an oriented 4-manifold  $\bar{W} = W \cup -W^*$  identifying two copies of  $S$ . Then  $\Sigma = F \cup -F^*$  is the disjoint union of 2-spheres. Since  $[F_2^+]$  and  $[F_2^{*+}]$  are characteristic, we see that the mod 2 homology class  $[\Sigma]_2 \in H_2(\bar{W}; \mathbb{Z}_2)$  is characteristic. By the Rochlin theorem ([6], [10]),  $\mu(\partial \bar{W}) = ([\Sigma]^2 - \text{sign } \bar{W})/16$  in  $Q/Z$ . But,  $\mu(\partial \bar{W}) = \mu(\partial W) - \mu(\partial W^*)$ ,  $[\Sigma]^2 = [F_2^+]^2 - [-F_2^{*+}]^2 = [F_2^+]^2 - [F_2^{*+}]^2$  and  $\text{sign } \bar{W} = \text{sign } W - \text{sign } W^*$ , where we count  $[-F_2^{*+}]^2, [F_2^{*+}]^2$  in  $W^*$ . It follows that

$$([F_2^+]^2 - \text{sign } W)/16 - \mu(\partial W) = ([F_2^{*+}]^2 - \text{sign } W^*)/16 - \mu(\partial W^*)$$

in  $Q/Z$ , showing the well-definedness of  $\delta_0(L)$ .

**DEFINITION 2.5.** Two links  $L_i \subset S_i$ ,  $i=0, 1$ , are said to be *cobordant in the weak sense* if:

- (1) There exists a compact oriented 4-manifold  $W$  such that  $\partial W = -S_0 \cup S_1$  and  $H_*(W, S_i; \mathbb{Z}_2) = 0$ ,  $i=0, 1$ ,
- (2) There exists a locally flat, compact oriented (possibly disconnected) surface  $F$  of (total) genus 0 in  $W$  such that  $\partial F = -L_0 \cup L_1$  (See Figure 1).




Figure 1.

**Theorem 2.6.** If proper links  $L_i \subset S_i$ ,  $i=0, 1$ , are cobordant in the weak sense, then  $\delta_0(L_0) = \delta_0(L_1)$ .

**Proof.** Let  $F \subset W$  be a cobordism pair for  $L_i \subset S_i$ ,  $i=0, 1$ , stated in Definition 2.5. Construct an oriented 4-manifold  $W' = W \cup D^3 \times [0, 1]$  identifying a 3-cell in  $S_i - L_i$  with  $D^3 \times i$  for each  $i$ ,  $i=0, 1$ . Then  $\partial W'$  is a connected sum  $(-S_0) \# S_1$ , which is a  $\mathbb{Z}_2$ -homology 3-sphere containing a proper link  $L'$ , regarded as the union  $-L_0 \cup L_1$ . Clearly,  $\delta_0(L' \subset (-S_0) \# S_1) = -\delta_0(L_0 \subset S_0) +$

$\delta_0(L_1 \subset S_1)$ . Note that  $H_2(W'; Z_2) = 0$ . Then  $F \subset W'$  is admissible for  $L' \subset (-S_0) \# S_1$ , and hence

$$\delta_0(L' \subset (-S_0) \# S_1) = ([F_Q^+]^2 - \text{sign } W')/16 - \mu((-S_0) \# S_1) = 0,$$

because  $W'$  is spin and  $H_2(W'; Q) = 0$ . Thus,  $\delta_0(L_0 \subset S_0) = \delta_0(L_1 \subset S_1)$ . This completes the proof.

In [5, Definition 2.1] the  $\delta$ -invariant  $\delta(k)$  of a knot  $k$  in  $S$  was defined so as to be  $\delta(k) = \delta_0(k)$ .

**Corollary 2.7.** *Let  $k \subset S$  be a knot obtained from a proper link  $L \subset S$  by a fusion. Then  $\delta_0(L) = \delta_0(k) = \delta(k)$ .*

Proof. The knot  $k \subset S$  and the link  $L \subset S$  are cobordant in the weak sense. The result follows from Theorem 2.6.

By a Dehn surgery we obtain from a knot  $k \subset S$  a unique (up to homeomorphism), closed, connected, oriented 3-manifold  $M$  such that  $H_1(M; Z)/\text{odd torsion} \cong Z$ , called a  $Z_2$ -homology handle (cf. [5, Remark 1.6 and Corollary 1.7]). In [4] we defined an invariant  $\in(M)$ , being 0 or 1, of  $M$ , calculable from the  $Z_2$ -Alexander polynomial of  $M$ .

**Corollary 2.8.** *Let  $L \subset S$  be a proper link. Let  $M$  be the  $Z_2$ -homology handle of a knot  $k \subset S$ , obtained from  $L$  by a fusion. Let  $a/b$  be a normal presentation of the slope  $s(L \subset S)$  with  $a$  odd. Then we have*

$$\delta_0(L) = \in(M)/2 + (a/b - ab)/16$$

in  $Q/Z$ .

Proof. By Lemma 1.2  $s(L) = s(k)$ . By Corollary 2.7  $\delta_0(L) = \delta(k)$ . Then the desired congruence follows from [5, Theorem 2.7 and Corollary 3.6].

**DEFINITION 2.9.** For a proper link  $L$  in  $S$  we define

$$\delta(L) = \delta(L \subset S) = \delta_0(L \subset S) + \lambda(L \subset S)/8$$

in  $Q/Z$ .

**REMARK 2.10.** Definition 2.9 is analogous to Murasugi's definition of the unoriented link type signature in [7] (cf. [5, Remark 4.8]).

**Theorem 2.11.** *The invariant  $\delta(L \subset S)$  is an unoriented link type invariant of a proper link  $L \subset S$ . That is,  $\delta(L \subset S) = \delta(L' \subset S')$  for any link  $L' \subset S'$  with an orientation-preserving homeomorphism  $S \rightarrow S'$  sending  $L$  to  $L'$  setwise.*

Proof. It suffices to show that  $\delta(L)$  does not depend on any particular orientations of the components,  $k_i$ , of  $L$ . Let  $F = \bigcup_{i=1}^r D_i \subset W$  be a standard admissible pair for  $L = \bigcup_{i=1}^r k_i \subset S$ . Note that  $[F_Q^+]^2 = \sum_{i=1}^r [D_{iQ}^+]^2 + 2 \sum_{i>j} [D_{iQ}^+] [D_{jQ}^+]$ .

$[D_{iQ}^+] = \sum_{i=1}^r [D_{iQ}^+]^2 - 2\lambda(L)$ . Then

$$\delta(L) = \delta_0(L) + \lambda(L)/8 = (\sum_{i=1}^r [D_{iQ}^+]^2 - \text{sign } W)/16 - \mu(\partial W).$$

Since  $[D_{iQ}^+]^2$  is not altered by changing the orientation of  $D_i$  (that is,  $k_i$ ), we have a desired result.

A link  $L \subset S$  is *amphicheiral* if there is an orientation-preserving homeomorphism  $S \rightarrow -S$  sending  $L$  to itself setwise. The following is direct from Theorem 2.11.

**Corollary 2.12.** *If a proper link  $L \subset S$  is amphicheiral, then  $2\delta(L)=0$  in  $Q/Z$ .*

Here is an example of a classical proper link.

**EXAMPLE 2.13.** Let  $L_r$  be an  $r$ -component link in a 3-sphere  $S^3$ , illustrated in Figure 2, where  $r \geq 2$ . The link




Figure 2.

$L_r$  is clearly proper. Choosing a suitable orientation of  $S^3$ ,  $\lambda(L_r)=r$ . Since we can have a trivial knot from  $L_r$  by a fusion, we see that  $\delta_0(L_r)=0$ . Therefore,  $\delta(L_r)=r/8$  in  $Q/Z$ .

### 3. Branched cyclic coverings and the $\delta$ - and $\delta_0$ -invariants

We consider a link  $\tilde{L} \subset \tilde{S}$  obtained from a link  $L \subset S$  by taking an  $n$ -fold cyclic branched covering of  $S$ , branched along  $L$ . Namely,  $\tilde{S}$  is the branched covering space over  $S$ , associated with an epimorphism  $H_1(S-L; Z) \rightarrow Z_n$  sending each meridian of  $L$  to a unit of  $Z_n$ , and  $\tilde{L}$  is the lift of  $L$ . We assume that  $\tilde{S}$  is a  $Z_2$ -homology 3-sphere.

First we consider the case  $n=2$ . Then  $L$  and  $\tilde{L}$  are knots by the Smith theory. Let  $L=k$  and  $\tilde{L}=\tilde{k}$ .

**Theorem 3.1.** *Let  $2a/b$  be a normal presentation of the slope  $s(\tilde{k})$ . Then*

$$\delta(\tilde{k}) = \delta(k) - (a/b - ab)/8$$

in  $Q/Z$ . In particular, if  $\tilde{k}$  is flat, then  $\delta(\tilde{k}) = \delta(k)$ .

Proof. By [5, Lemma 4.5]  $s(k) = 2s(\tilde{k}) = 4a/b$ . Since  $(2a+b)/b$  and  $(4a+b)/b$  are normal presentations of  $s(\tilde{k})$  and  $s(k)$ , respectively, we see from Corollary 2.8 that

$$\delta(\tilde{k}) = \in(\tilde{M})/2 + \{(2a+b)/b - (2a+b)b\}/16 = \in(\tilde{M})/2 + (a/b - ab)/8 + (1 - b^2)/16,$$

and

$$\delta(k) = \in(M)/2 + \{(4a+b)/b - (4a+b)b\}/16 = \in(M)/2 + (a/b - ab)/4 + (1 - b^2)/16,$$

where  $\tilde{M}$  and  $M$  are the  $Z_2$ -homology handles of  $\tilde{k} \subset \tilde{S}$  and  $k \subset S$ , respectively. Since  $\tilde{M}$  is a 2-fold covering space of  $M$ , it follows from [4, Lemma 4.2] that  $\in(\tilde{M}) = \in(M)$ . Now we have a desired congruence. This completes the proof.

Next, to consider the case that the covering degree  $n$  is an odd prime  $p$ , we remark the following:

**Lemma 3.2.**  $\lambda(\tilde{L}) = \lambda(L)/n$ .

**Corollary 3.3.**  $\tilde{L}$  is proper if and only if  $L$  is proper.

Proof of Lemma 3.2. It suffices to show that for  $i \neq j$   $\text{Link}_{\tilde{S}}(\tilde{k}_i, \tilde{k}_j) = \text{Link}_S(k_i, k_j)/n$ . Let  $F_i$  be a characteristic surface (cf. [5]) of  $k_i$  in  $S$  such that  $L - k_i$  intersects  $F_i$  transversally. Write  $[\partial F_i] = a_i r_i [m_i] + b_i r_i [l_i]$  in  $H_1(\partial T(k_i); Z)$  for a meridian-longitude pair  $(m_i, l_i)$  of  $T(k_i)$  such that the lift of  $l_i$  has  $n$  components, where  $(a_i, b_i) = 1$  and  $r_i$  is an integer  $> 0$ . Let  $\tilde{l}_i$  be a component of the lift of  $l_i$ . For the lift  $\tilde{m}_i$  of  $m_i$ , the pair  $(\tilde{m}_i, \tilde{l}_i)$  forms an  $m.l.$  pair of a tubular neighborhood  $T(\tilde{k}_i)$  of  $\tilde{k}_i$  which is the lift of  $T(k_i)$ . Note that the lift  $\tilde{F}_i$  of  $F_i$  is an oriented surface which is a branched  $Z_n$ -covering space of  $F_i$  branched over the set  $F_i \cap (L - k_i)$ . Clearly  $[\partial \tilde{F}_i] = a_i r_i [\tilde{m}_i] + b_i r_i n [\tilde{l}_i]$  in  $H_1(\partial T(\tilde{k}_i); Z)$ . Since the intersection numbers,  $\text{Int}(\tilde{F}_i, \tilde{k}_j)$  and  $\text{Int}(F_i, k_j)$  are equal, we have

$$\text{Link}_{\tilde{S}}(\tilde{k}_i, \tilde{k}_j) = \text{Int}(\tilde{F}_i, \tilde{k}_j)/b_i r_i n = \text{Int}(F_i, k_j)/b_i r_i n = \text{Link}_S(k_i, k_j)/n.$$

This completes the proof.

Proof of Corollary 3.3. When  $n$  is even,  $L$  and  $\tilde{L}$  are knots by the Smith theory. So, assume  $n$  is odd. It suffices to show that  $\text{Link}_{\tilde{S}}(\tilde{k}_i, \tilde{k}_j)_2 = \text{Link}_S(k_i, k_j)_2$  for  $i \neq j$ . This is obtained by a mod 2 version of the proof of Lemma 3.2, since  $b_i r_i n$  is odd. This completes the proof.

We shall show the following theorem, where note that  $(p^2 - 1)/8$  is an in-

teger.

**Theorem 3.4.** *Let  $\tilde{L} = \cup_{i=1}^r \tilde{k}_i \subset \tilde{S}$  be a proper link and assume that the covering degree is an odd prime  $p$ . Let  $2a_i/b_i$  be a normal presentation of the slope  $s(\tilde{k}_i)$ ,  $i=1, 2, \dots, r$ . Then*

$$\delta(\tilde{L}) = p\delta(L) - \{(p^2-1)/8\} \sum_{i=1}^r a_i/b_i$$

in  $Q/Z$ .

Proof. Let  $F \subset W$  and  $\tilde{F} \subset \tilde{W}$  be standard admissible pairs for  $L \subset S$  and  $\tilde{L} \subset \tilde{S}$ , respectively, such that  $\tilde{F} \subset \tilde{W}$  is obtained from  $F \subset W$  by taking a  $Z_p$ -covering branched along  $F$ . [One can see directly or by a transfer argument that such pairs exist.] Let  $\partial W - S = S^*$  and  $\partial \tilde{W} - \tilde{S} = \tilde{S}^*$ . By the proof of Theorem 2.11,

$$\begin{aligned} \delta(L) &= (\sum_{i=1}^r [D_{iQ}^+]^2 - \text{sign } W)/16 - \mu(S) - \mu(S^*), \text{ and} \\ \delta(\tilde{L}) &= (\sum_{i=1}^r [\tilde{D}_{iQ}^+]^2 - \text{sign } \tilde{W})/16 - \mu(\tilde{S}) - \mu(\tilde{S}^*), \end{aligned}$$

where  $F = \cup_{i=1}^r D_i$ ,  $\tilde{F} = \cup_{i=1}^r \tilde{D}_i$ , and  $\tilde{D}_i$  corresponds to  $D_i$ . Then since  $[D_{iQ}^+]^2/p = [\tilde{D}_{iQ}^+]^2$  (cf. [5, the proof of Lemma 4.9]),

$$\begin{aligned} \delta(\tilde{L}) - p\delta(L) &= (1-p^2) \sum_{i=1}^r [\tilde{D}_{iQ}^+]^2/16 + (-\text{sign } \tilde{W} + p \text{ sign } W)/16 \\ &\quad - (\mu(\tilde{S}) - p\mu(S)) - (\mu(\tilde{S}^*) - p\mu(S^*)). \end{aligned}$$

By the definition of  $\alpha$ -invariant in [5, Section 4],

$$\alpha(Z_p, \tilde{S}) + \alpha(Z_p, \tilde{S}^*) = -\text{sign } \tilde{W} + p \text{ sign } W - (\sum_{i=1}^r [\tilde{D}_{iQ}^+]^2) (p^2-1)/3.$$

Therefore,

$$\begin{aligned} \delta(\tilde{L}) - p\delta(L) &= \{1-p^2+(p^2-1)/3\} (\sum_{i=1}^r [\tilde{D}_{iQ}^+]^2)/16 - (\mu(\tilde{S}) - p\mu(S) \\ &\quad - \alpha(Z_p, \tilde{S})/16) - (\mu(\tilde{S}^*) - p\mu(S^*) - \alpha(Z_p, \tilde{S}^*)/16). \end{aligned}$$

First, let  $p > 3$ . Then by [5, Theorems 11.1 and 12.1],

$$\begin{aligned} \mu(\tilde{S}^*) &= p\mu(S^*) + \alpha(Z_p, \tilde{S}^*)/16, \text{ and} \\ \mu(\tilde{S}) &= p\mu(S) + \alpha(Z_p, \tilde{S})/16 + \{(p^2-1)/24\} \sum_{i=1}^r a_i/b_i, \end{aligned}$$

where note that  $(p^2-1)/24$  is an integer. Since  $[\tilde{D}_{iQ}^+]^2 \equiv 2a_i/b_i \pmod{1}$  (cf. [5, Lemma 2.6]), it follows that

$$\begin{aligned} \delta(\tilde{L}) - p\delta(L) &= -\{(p^2-1)/24\} \sum_{i=1}^r [\tilde{D}_{iQ}^+]^2 - \{(p^2-1)/24\} \sum_{i=1}^r a_i/b_i \\ &= -\{(p^2-1)/8\} \sum_{i=1}^r a_i/b_i. \end{aligned}$$

Now let  $p=3$ . By [5, Theorems 11.1 and 12.1],

$$\mu(\tilde{S}^*) = 3\mu(S^*) + 9\alpha(Z_3, \tilde{S}^*)/16, \text{ and}$$

$$\mu(\tilde{S}) = 3\mu(S) + 9\alpha(Z_3, \tilde{S})/16 + 3\sum_{i=1}^r a_i/b_i.$$

Directly or by a transfer argument,  $\text{sign } \tilde{W} = \text{sign } W$ . Then

$$\begin{aligned} \alpha(Z_3, \tilde{S})/2 + \alpha(Z_3, \tilde{S}^*)/2 &= -\text{sign } \tilde{W}/2 + 3 \text{ sign } W/2 - (\sum_{i=1}^r [\tilde{D}_{iQ}^+]^2)4/3 \\ &\equiv -(\sum_{i=1}^r [\tilde{D}_{iQ}^+]^2)4/3 \pmod{1}. \end{aligned}$$

Then,

$$\begin{aligned} \delta(\tilde{L}) - 3\delta(L) &= -(\sum_{i=1}^r [\tilde{D}_{iQ}^+]^2)/3 - \alpha(Z_3, \tilde{S})/2 - \alpha(Z_3, \tilde{S}^*)/2 \\ &\quad - 3\sum_{i=1}^r a_i/b_i = \sum_{i=1}^r [\tilde{D}_{iQ}^+]^2 - 3\sum_{i=1}^r a_i/b_i = -\sum_{i=1}^r a_i/b_i \end{aligned}$$

in  $Q/Z$ , because  $[\tilde{D}_{iQ}^+]^2 \equiv 2a_i/b_i \pmod{1}$ . This completes the proof.

**Theorem 3.5.** *Let  $\tilde{L} \subset \tilde{S}$  be proper and assume that the covering degree is an odd prime  $p$ . Then we have*

$$\delta_0(\tilde{L}) = p\delta_0(L) - \{(p^2-1)/8\} s^*(\tilde{L})$$

in  $Q/Z$ , where  $s^*(\tilde{L})$  is the half-slope of the link  $\tilde{L} \subset \tilde{S}$ .

Proof. By Theorem 3.4,

$$\delta_0(\tilde{L}) + \lambda(\tilde{L})/8 = p\delta_0(L) + p\lambda(L)/8 - \{(p^2-1)/8\} \sum_{i=1}^r a_i/b_i.$$

Since  $\lambda(L)/p = \lambda(\tilde{L})$  by Lemma 3.2, we have

$$\delta_0(\tilde{L}) - p\delta_0(L) = -\{(p^2-1)/8\} (\sum_{i=1}^r a_i/b_i - \lambda(\tilde{L})) = -\{(p^2-1)/8\} s^*(\tilde{L}).$$

This completes the proof.

**Corollary 3.6.** *If  $\tilde{L}$  is flat, then  $\delta_0(\tilde{L}) = \delta_0(L)$ .*

Proof. By Lemma 1.5  $s(\tilde{L}) = 0$  implies  $s^*(\tilde{L}) = 0$ . So, by Theorem 3.5  $\delta_0(\tilde{L}) = p\delta_0(L)$ . By Lemmas 1.3, 3.2 and [5, Lemma 4.5],  $s(L) = ps(\tilde{L})$ , so that  $s(L) = 0$ . By Corollary 2.8  $2\delta_0(L) = 0$ . Using that  $p$  is odd, the proof is completed.

## References

- [1] M. Freedman and R. Kirby: *A geometric proof of Rohlin's theorem*, Proc. Sympos. Pure Math. 32, Part 2, Amer. Math. Soc., Providence, R.I., 1978, 85–97.
- [2] C. McA. Gordon: *Knots, homology spheres, and contractible 4-manifolds*, Topology **14** (1975), 151–172.
- [3] L. Guillou and A. Marin: *Une extension d'un théorème de Rohlin sur la signature*, C.R. Acad. Sci. Paris **285**, Serie A (1977), 95–98.
- [4] A. Kawauchi: *On 3-manifolds admitting orientation-reversing involutions*, J. Math. Soc. Japan **33** (1981), 571–589.

- [5] A. Kawauchi: *On the Rochlin invariants of  $Z_2$ -homology 3-spheres with cyclic actions*, Japan. J. Math. **8** (1982), 217–258.
- [6] M.A. Kervaire and J.W. Milnor: *On 2-spheres in a 4-manifold*, Proc. Nat. Acad. Sci. U.S.A. **49** (1961), 1651–1657.
- [7] Y. Matsumoto: *An elementary proof of Rochlin's signature theorem and its extension by Guillou and Marin* (preprint).
- [8] K. Murasugi: *On the signature of links*, Topology **9** (1970), 283–298.
- [9] R. Robertello: *An invariant of knot cobordism*, Comm. Pure Appl. Math. **18** (1965), 543–555.
- [10] V.A. Rochlin: *New results in the theory of 4-dimensional manifolds*, Dokl. Akad. Nauk. SSSR **84** (1952), 221–224 (Russian).

Department of Mathematics  
Osaka City University  
Sumiyoshi-ku, Osaka 558  
Japan