

Title	An application on Nagao's lemma
Author(s)	Ikeda, Tadashi; Sasaki, Hiroki; Yoshida, Tomoyuki
Citation	Osaka Journal of Mathematics. 22(2) P.391-P.392
Issue Date	1985
Text Version	publisher
URL	https://doi.org/10.18910/4258
DOI	10.18910/4258
rights	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/repo/ouka/all/>

AN APPLICATION ON NAGAO'S LEMMA

Dedicated to Professor Hiroshi Nagao on his sixtieth birthday

TADASI IKEDA, HIROKI SASAKI AND TOMOYUKI YOSHIDA

(Received July 5, 1984)

With time, the importance of Nagao's lemma has grown in modular representation theory of finite groups. In this note, we add another application.

Let G be a finite group, and let F be a field of characteristic $p > 0$.

For a subgroup H of G and a (right) FG -module V , we denote V^H the fixed-point-set of H in V , so that V^H is an $FN_G(H)$ -module. The trace map $Tr_H^G: V^H \rightarrow V^G$ is defined by $Tr_H^G(v) = \sum_g vg$, where g runs over a complete set of representatives of $H \backslash G$.

Main Theorem. *Let V be an indecomposable FG -module in a block B , and let P be a p -subgroup of G . Then each composition factor of the $FN_G(P)$ -module*

$$V(P) := V^P / \sum_{A < P} Tr_A^P(V^A),$$

where A runs over proper subgroups of P , belongs to a block b such that $b^G = B$.

REMARK. If $V(P) \neq 0$, then P is contained in a defect group of B .

Proof. of the theorem. Set $N = N_G(P)$. Let e be the centrally primitive idempotent of FG corresponding to B . Let $s: Z(FG) \rightarrow Z(FN)$ be the Brauer homomorphism. Then Nagao's lemma ([2], Chapter III, Theorem 7.5) states that

$$V_N = V_N s(e) \oplus W_1 \oplus \cdots \oplus W_n$$

as FN -modules, where each W_i is Q_i -projective FN -module for some p -subgroup Q_i of N with $P \not\cong Q_i$. Thus in order to prove the theorem, it will suffice to show that

$$W_i^P \subseteq \sum_{A < P} Tr_A^P(V^A),$$

where A runs over proper subgroups of P . But this follows directly from the following lemma, and so the theorem is proved.

Lemma. *Let N be a finite group with a normal p -subgroup P . Let W be a Q -projective FN -module, where $Q \cong P$. Then*

$$W^P = \sum_{A < P} Tr_A^P(W^A),$$

where A runs over proper subgroups of P .

Proof. In order to prove this lemma, we may assume that for some FQ -module U ,

$$W = \text{Ind}_Q^N(U).$$

Then by Mackey decomposition, we have that

$$W_P = \bigoplus_n \text{Ind}_P^P \cap Q^n(U_P^n \cap Q^n),$$

where n runs over a complete set of representatives of $Q \backslash N/P$ and $Q^n = n^{-1}Qn$. Let n be an element of N and set $R = P \cap Q^n$, $X = U_R^n$. Since P is normal in N and Q is not contained in P , we have that R is a proper subgroup of P . Thus, in order to prove the lemma, it will suffice to show that

$$(\text{Ind}_R^P(X))^P \subseteq Tr_R^P(\text{Ind}_R^P(X)^R).$$

But this follows directly from an easy calculation (eq. [2] Chapter II Lemma 3.4). The lemma is proved.

REMARK. The main theorem can be proved also by the Brauer homomorphism of modules, which is defined by Broué and Puig [1]. Let B be a block of G and e a corresponding central primitive idempotent of FG . We define the Brauer homomorphism Br_P^V with respect to P by the canonical homomorphism $V^P \rightarrow V(P)$. Now let $s_P: Z(FG) \rightarrow Z(FC_G(P))$ be the classical Brauer homomorphism with respect to P . Then we can prove that $Br_P^V(ve) = Br_P^V(v)s_P(e)$ for the element v of V^P . The main theorem is immediate from this fact.

References

- [1] M. Broué: *On Scott modules and p -permutation modules: an elementary approach through the Brauer morphism, following remarks of L. Puig*, Proc. Amer. Math. Soc. **93** (1985), 401–408.
- [2] W. Feit: *The representation theory of finite groups*, North-Holland, 1982.

Tadasi Ikeda and Tomoyuki Yoshida
 Department of Mathematics
 Faculty of Science
 Hokkaido University
 Sapporo 060, Japan

Hiroki Sasaki
 Department of Mathematics
 Faculty of Education
 Yamaguchi University
 Yamaguchi 753, Japan