



| | |
|--------------|---|
| Title | On an Isomorphism between Specht module and Left cell of $S(n)$ |
| Author(s) | Naruse, Hiroshi |
| Citation | 大阪大学, 1999, 博士論文 |
| Version Type | VoR |
| URL | https://doi.org/10.11501/3169544 |
| rights | |
| Note | |

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

On an Isomorphism between Specht Module and Left Cell of \mathfrak{S}_n

Hiroshi NARUSE

Okayama University

(Communicated by Y. Kawada)

Dedicated to Professor Nagayoshi Iwahori on his 60th birthday

Introduction.

The irreducible representations of the symmetric group \mathfrak{S}_n are classically known and they are parametrized by the partitions of n . One of their realizations is so called Specht module which is defined over \mathbb{Z} and has a natural base (Young's natural base). On the other hand, Kazhdan and Lusztig [3] constructed another realization, the W -graph representation which is obtained from a left cell of \mathfrak{S}_n and it also has a natural \mathbb{Z} -base (the vertices of the W -graph). We give in this paper an explicit isomorphism between the above two modules and show that the base change matrix is uni-triangular for some ordering of the base. Using this isomorphism, if a partition λ satisfies certain condition, we can construct the W -graph of the left cell of \mathfrak{S}_n (without some edges which connect vertices having the same I -set) corresponding to λ not using the Kazhdan-Lusztig polynomials $P_{y,w}$ but using the relations in Specht module (the Garnir relations) inductively.

§1. The λ -diagram.

1.1. Notations. Let $P(n)$ be the set of partitions $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$, where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r > 0$ and $\sum_{i=1}^r \lambda_i = n$. For a partition λ , the Young frame of λ is the arrangement of n squares; the first row λ_1 , the second row λ_2 , \dots , the last row λ_r parts, and line up to the left. Young tableau of shape λ has the frame λ and each square is numbered from 1 to n . A Young tableau of shape λ is called standard if its numbering is increasing from left to right in each row and from top to bottom in

Received October 17, 1986

Revised December 24, 1988

each column. We write Tab_λ the set of all Young tableaux of shape λ and STab_λ the set of all standard Young tableaux of shape λ . The transposition of the partition λ is denoted by ${}^t\lambda$. For a Young tableau D of shape λ , tD denotes the transposition of D whose shape is ${}^t\lambda$. We denote by $D_{\lambda, \text{TOP}}$ the standard Young tableau of shape λ , whose numbering is 1 to λ_1 for the first row, λ_1+1 to $\lambda_1+\lambda_2$ for the second row and so on. We set $D_{\lambda, \text{BOT}} = {}^t(D_{\lambda, \text{TOP}})$. For a Young tableau D , we put

$$\begin{aligned} I(D) &= \{i \mid 1 \leq i \leq n-1, i+1 \text{ is in a lower position than } i \text{ in } D\}, \\ I_0(D) &= \{i \in I(D) \mid i+1 \text{ is in the left side of } i \text{ in } D\}, \\ I_1(D) &= \{i \in I(D) \mid i+1 \text{ is directly below } i \text{ in } D\}. \end{aligned}$$

Then we have

LEMMA 1. For a standard Young tableau D of shape λ , $\lambda \in P(n)$,

- (1) $I(D) = I_0(D) \sqcup I_1(D)$ (disjoint union).
- (2) $I(D) \sqcup I({}^tD) = \{1, 2, \dots, n-1\}$.
- (3) $I_0(D) = \emptyset$ if and only if $D = D_{\lambda, \text{BOT}}$.
- (4) $I_1({}^tD) = \emptyset$ if and only if $D = D_{\lambda, \text{TOP}}$.

PROOF. (1) The equality follows immediately from the definitions.

(2) We see easily that $i+1$ is either in A or in B (see Fig. 1). Hence $i+1 \in A$ iff $i \in I({}^tD)$ and $i+1 \in B$ iff $i \in I(D)$.




FIGURE 1

(3) The if part clearly follows. On the other hand, if $I_0(D) = \emptyset$, $i+1$ must be directly below i or in the right side of i , hence D must have the form $D_{\lambda, \text{BOT}}$.

(4) Note that ${}^tD_{\lambda, \text{TOP}} = D_{\lambda, \text{BOT}}$. Then (4) follows from (3).

1.2. The Robinson-Schensted correspondence. The following fact is well known.

PROPOSITION 1 (cf. [4]). There is a bijection from \mathfrak{S}_n to $\bigsqcup_{\lambda \in P(n)} (\text{STab}_\lambda \times \text{STab}_\lambda)$ which is called Robinson-Schensted's correspondence. By this, if x corresponds to $(P(x), Q(x))$, then $P(x^{-1}) = Q(x)$.

We denote by s_i the transposition $(i, i+1) \in \mathfrak{S}_n$. The set $S = \{s_i\}_{1 \leq i \leq n-1}$ generates \mathfrak{S}_n as a Coxeter group. Let $l(x)$ be the length of $x \in \mathfrak{S}_n$ with respect to this generator set S .

LEMMA 2. For x in \mathfrak{S}_n , we have

- (1) $l(s_i x) = l(x) - 1$ if and only if $i \in I(P(x))$,
- (2) $l(x s_i) = l(x) - 1$ if and only if $i \in I(Q(x))$.

PROOF. (1) Suppose $x \in \mathfrak{S}_n$ has a form $x = \begin{pmatrix} 1 & 2 & \cdots & n \\ a_1 & a_2 & \cdots & a_n \end{pmatrix}$. Then $l(s_i x) = l(x) - 1$ iff i appears after $i+1$ in a_1, a_2, \dots, a_n , which is equivalent to the condition $i \in I(P(x))$ by the construction of $P(x)$.

(2) As $l(x^{-1}) = l(x)$, (2) follows from (1).

1.3. For $D \in \text{STab}_\lambda$, we define $x(D) \in \mathfrak{S}_n$ by

$$x(D) = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ a_1 & a_2 & a_3 & \cdots & a_n \end{pmatrix} \quad \text{if } D = \begin{array}{|c|c|} \hline a_s & \\ \hline \vdots & \\ \hline a_2 & \vdots \\ \hline a_1 & a_{s+1} \\ \hline \end{array}$$

LEMMA 3. For $D \in \text{STab}_\lambda$, we have

- (1) By the Robinson-Schensted correspondence, $x(D)$ corresponds to $(D, D_{\lambda, \text{Bot}})$.
- (2) $l(s_i x(D)) = l(x(D)) - 1$ if and only if $i \in I(D)$.

PROOF. (1) It is clear from the definition of $x(D)$ and the construction of the Robinson-Schensted correspondence.

(2) Clear from (1) and Lemma 2.

LEMMA 4.

- (1) $l(x(D_{\lambda, \text{Top}})) = \frac{n^2 - \sum \lambda_i^2}{2} - \frac{\sum (i-1)\lambda_i(\lambda_i - 1)}{2}$.
- (2) $l(x(D_{\lambda, \text{Bot}})) = \sum (i-1)\lambda_i$.

1.4. For $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r) \in P(n)$, we put

$$N = l(x(D_{\lambda, \text{Top}})) - l(x(D_{\lambda, \text{Bot}})) = \frac{n(n+1)}{2} - \frac{\sum i\lambda_i(\lambda_i + 1)}{2}.$$

For $0 \leq k \leq N$, we define a subset \mathcal{D}_k^λ of STab_λ inductively from N to 0. We put

$$\mathcal{D}_N^\lambda = \{D_{\lambda, \text{TOP}}\},$$

and for $0 \leq k < N$, we put

$$\mathcal{D}_k^\lambda = \{D \in \text{STab}_\lambda \mid D = s_j D' \text{ for some } D' \in \mathcal{D}_{k+1}^\lambda, j \in I_0(D')\},$$

where $s_j D'$ is the standard tableau interchanging j and $j+1$ in D .

PROPOSITION 2.

- (1) $\coprod_{0 \leq k \leq N} \mathcal{D}_k^\lambda = \text{STab}_\lambda$ (disjoint union).
- (2) $\mathcal{D}_0^\lambda = \{D_{\lambda, \text{BOT}}\}$.

PROOF. For $D \in \mathcal{D}_k^\lambda$, we can show inductively that

$$l(x(D)) = l(x(D_{\lambda, \text{TOP}})) - (N - k).$$

Hence if $k \neq l$, $\mathcal{D}_k^\lambda \cap \mathcal{D}_l^\lambda = \emptyset$ i.e. disjoint. For $D \in \text{STab}_\lambda$, compare it with $D_{\lambda, \text{TOP}}$ for the top row from left to right then for the second row in the same manner and so on. If the first difference is a in D , we must have $a-1 \in I_0({}^t D)$ and $D' = s_{a-1} D \in \text{STab}_\lambda$. For this D' , instead of D , do the same operation. Continuing this, we finally get $D_{\lambda, \text{TOP}}$. If it takes m times, then $D \in \mathcal{D}_{N-m}^\lambda$. If we do the same operation on ${}^t D$ (compare with ${}^t D_{\lambda, \text{BOT}}$), we get ${}^t D_{\lambda, \text{BOT}}$ after k times interchange. Transposing all of them, if $D \in \mathcal{D}_k^\lambda$ then $D_{\lambda, \text{BOT}} \in \mathcal{D}_{N-k}^\lambda$. Therefore we get (1) and (2).

For $D \in \text{STab}_\lambda$, we define $\text{ht}(D) = k$ if $D \in \mathcal{D}_k^\lambda$.

1.5. Partial orders on STab_λ . We define three partial orders on STab_λ .

(1) For $D, D' \in \text{STab}_\lambda$, if $\text{ht}(D') = \text{ht}(D) + 1$ and $D' = (i, j)D$, for some transposition (i, j) , we define $D < D'$. Let \leq be the transitive closure of this relation, thus \leq is a partial order on STab_λ .

(2) We define a partial order \trianglelefteq on STab_λ as follows. For $D, D' \in \text{STab}_\lambda$,

$$D \trianglelefteq D' \text{ if and only if } m_{ir}(D) \leq m_{ir}(D') \text{ for all } i \text{ and all } r,$$

where $m_{ir}(D) = \#\{j \mid j \leq i \text{ and } j \text{ appears in the first } r \text{ rows of } D\}$ (cf. James [2; 3.10]). This order can be interpreted as follows. For $D \in \text{STab}_\lambda$ and $1 \leq i \leq n$, we define $D_{\leq i}$ the sub-diagram of D whose entries are less than or equal to i . Then this is also a standard Young tableau of size i . We also define for a Young tableau D , $\lambda(D) = (\lambda_1(D), \lambda_2(D), \dots, \lambda_r(D))$ where $\lambda_i(D)$ is the length of i -th row of D , hence D has a Young frame of shape $\lambda(D)$. For two partitions $\lambda, \mu \in P(n)$, we define $\lambda \leq \mu$ iff

$\lambda_1 \leq \mu_1, \lambda_1 + \lambda_2 \leq \mu_1 + \mu_2, \dots, \lambda_1 + \lambda_2 + \dots + \lambda_r \leq \mu_1 + \mu_2 + \dots + \mu_s$ and $r \geq s$ ($\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r), \mu = (\mu_1, \mu_2, \dots, \mu_s)$). Then $D \leq D'$ iff $\lambda(D_{\leq i}) \leq \lambda(D'_{\leq i})$ for all $1 \leq i \leq n$.

(3) We define the third partial order \leq on STab_λ . $D \leq D'$ if and only if $x(D) \leq x(D')$, where \leq is the Bruhat order on \mathfrak{S}_n . Then we have

PROPOSITION 3. For $D, D' \in \text{STab}_\lambda$,

$D \leq D'$ if and only if $D \trianglelefteq D'$, and these imply $D \leq D'$.

PROOF. Clearly $D \leq D'$ implies $D \trianglelefteq D'$. The partial order \leq is generated by the relation $D_1 < D_2, D_2 = (p, q)D_1$. If p is in the k -th row and q is in the l -th row in D_1 , and if $p > q$, then $k < l$. It is easy to see that

$$m_{ir}(D_1) = m_{ir}(D_2) \quad \text{for } i < q \text{ or } i \geq p \text{ or } r < k \text{ or } r \geq l,$$

and

$$m_{ir}(D_1) + 1 = m_{ir}(D_2) \quad \text{for } q \leq i < p \text{ and } k \leq r < l,$$

therefore $D_1 \triangleleft D_2$. Hence $D \leq D'$ implies $D \trianglelefteq D'$. On the other hand, using the lemma below inductively, we can prove that $D \trianglelefteq D'$ implies $D \leq D'$.

LEMMA 5. For $D, D' \in \text{STab}_\lambda$ such that $D \triangleleft D'$, there exists $D'' \in \text{STab}_\lambda$ such that $\text{ht}(D'') = \text{ht}(D) + 1$ and $D < D'' \trianglelefteq D'$.

PROOF. Compare D and D' for the first row from left to right then for the second row in the same manner and so on. If the first difference is a in D and b in D' in the p -th row, then $a > b$ by $D \triangleleft D'$. A corner of a Young tableau is a square that there is no square attached to the right of and below it. We put

$$C_D = \{j \mid b \leq j < a, j \text{ is in a corner of } D_{\leq a-1}, \text{ and } j \text{ is a lower square than } a \text{ in } D\}.$$

As b is in the lower position than a in D , $C_D \neq \emptyset$. Let $c \in C_D$ be in the highest position among C_D . Then $b \leq c < a$, and if we put $D'' = (a, c)D$ then $D'' \in \text{STab}_\lambda$ and $\text{ht}(D'') = \text{ht}(D) + 1$ by the definition of c , therefore $D < D''$. We must show $D'' \trianglelefteq D'$. If c is in the q -th row of D ,

$$m_{ir}(D'') = m_{ir}(D) \leq m_{ir}(D') \quad \text{if } i < c \text{ or } i \geq a \text{ or } r < p \text{ or } r \geq q,$$

and

$$m_{i,r}(D'') = m_{i,r}(D) + 1 \quad \text{if } c \leq i < a \text{ and } p \leq r < q.$$

Therefore it is enough to show that

$$(1.5.1) \quad m_{i,r}(D) < m_{i,r}(D') \quad \text{if } c \leq i < a \text{ and } p \leq r < q.$$

We fix i , $c \leq i < a$, and move r from p to $q-1$. If the first negation of (1.5.1) occurs for $r=r_0$, then $m_{i,r_0}(D) = m_{i,r_0}(D')$. We define s to be the smallest j such that $\lambda_j(D_{\leq a-1}) = \lambda_j(D'_{\leq a-1})$. For $p \leq k < s$,

$$m_{i,k}(D) = m_{b-1,k}(D) \leq m_{b-1,k}(D') < m_{b,k}(D') \leq m_{i,k}(D').$$


Therefore r_0 must satisfy $s \leq r_0 < q$. But then,

$$\lambda_{r_0+1}(D_{\leq i}) = \lambda_{r_0}(D_{\leq i}) > \lambda_{r_0}(D'_{\leq i}) \geq \lambda_{r_0+1}(D'_{\leq i}),$$

hence $m_{i,r_0+1}(D) > m_{i,r_0+1}(D')$ which contradicts the fact $D \triangleleft D'$. Therefore we have (1.5.1) and finally we get $D'' \trianglelefteq D'$.

1.6. We define the λ -diagram to be the Hasse diagram of $(\text{STab}_\lambda, \leq)$ whose vertices have two data D and $I(D)$. If $D \in \mathcal{D}_k^\lambda$, $D' \in \mathcal{D}_{k+1}^\lambda$ and $s_j D' = D$ then we label the edge $\{D, D'\}$ as s_j . An edge may have no label (cf. Example 5.1).

EXAMPLE. $\lambda = (3, 2)$.



§2. The left cells of \mathcal{S}_n .

By Kazhdan and Lusztig [3], the left cells of \mathcal{S}_n are determined as follows. For $D \in \text{STab}_\lambda$, we define

$X_D = \{x \in \mathfrak{S}_n \mid Q(x) = D \text{ for the Robinson-Schensted correspondence}\}.$

We denote by Γ_D the W -graph corresponding to X_D with edge and multiplicity defined by $\mu(x, y)$ (cf. [3]). Then $\{\Gamma_D \mid D \in \text{STab}_\lambda, \lambda \in P(n)\}$ is the set of all left cells of \mathfrak{S}_n and if $D, D' \in \text{STab}_\lambda$, then Γ_D and $\Gamma_{D'}$ are isomorphic as W -graphs ([3; Theorem 1.4]). We particularly consider $\Gamma^\lambda = \Gamma_{D_{\lambda, \text{Bot}}}$ on $X^\lambda = X_{D_{\lambda, \text{Bot}}}$.

PROPOSITION 4. *The λ -diagram can be identified with the induced subgraph which is obtained from the Hasse diagram of the Bruhat order of \mathfrak{S}_n by restricting vertices to X^λ .*

PROOF. For $D \in \text{STab}_\lambda$, $x(D)$ corresponds to $(D, D_{\lambda, \text{Bot}})$. For $D, D' \in \text{STab}_\lambda$, $l(x(D')) = l(x(D)) + 1$ if and only if $\text{ht}(D') = \text{ht}(D) + 1$, and in this case $x(D') > x(D)$ if and only if $D' > D$ (cf. Hiller [1; Definition 6.0]).

LEMMA 6. *If $x \leq y$ and $l(y) - l(x) \geq 2$, then*

$$\mu(x, y) \neq 0 \text{ implies } \mathcal{L}(y) \subset \mathcal{L}(x),$$

where $\mathcal{L}(x) = \{s \in S \mid sx < x\}$.

PROOF. By [3; (2.3.g)], for $s \in S$,

$$P_{x, y} = P_{sx, y} \text{ if } x < y, sy < y, sx > x,$$

where $P_{x, y}$ is the Kazhdan-Lusztig polynomial.

$$\begin{aligned} \text{The degree of } P_{sx, y} &\leq \frac{1}{2}(l(y) - l(sx) - 1) \\ &= \frac{1}{2}(l(y) - l(x) - 2) \\ &< \frac{1}{2}(l(y) - l(x) - 1). \end{aligned}$$

Therefore if $\mu(x, y) \neq 0$ and $sy < y$ then we must have $sx < x$. Hence $\mathcal{L}(y) \subset \mathcal{L}(x)$.

COROLLARY 1. Γ^λ is obtained from λ -diagram by adding some edges $\{D, D'\}$ for $D \in \mathcal{D}_k^\lambda, D' \in \mathcal{D}_l^\lambda, l - k = \text{odd} \geq 3, I(D') \subset I(D)$.

PROOF. For $D \in \text{STab}_\lambda, I(D) = \{i \mid s_i \in \mathcal{L}(x(D))\}$, thus it is clear from the lemma above.

For $x \in X^\lambda$ we denote by g_D the base element of Γ^λ corresponding

to x , where $D=P(x)$. Thus $\{g_D\}_{D \in \text{STab}_\lambda}$ is the natural \mathbb{Z} -base for Γ^λ .

§3. Specht modules.

In this section we will quote some results of James [2].

3.1. We define an equivalence relation \sim on Tab_λ by $D \sim D'$ if and only if i -th rows of D and D' are equal as sets for all i . We define

$$\text{Tabloid}_\lambda = \text{Tab}_\lambda / \sim \quad (\text{the set of equivalence classes under } \sim).$$

We denote by $[D]$ the equivalence class containing D . And we put

$$M_\lambda = \text{the free } \mathbb{Z}\text{-module on } \text{Tabloid}_\lambda.$$

Then \mathfrak{S}_n acts naturally on M_λ . We also define for $D \in \text{Tab}_\lambda$

$$V_D = \{x \in \mathfrak{S}_n \mid x \text{ stabilizes each column of } D\},$$

$$k_D = \sum_{x \in V_D} \text{sgn}(x)x \in \mathbb{Z}[\mathfrak{S}_n],$$

$$e_D = k_D[D] \in M_\lambda,$$

$$S^\lambda = \langle e_D \rangle_{D \in \text{Tab}_\lambda} \subset M_\lambda.$$

S^λ is called the Specht module corresponding to the partition λ .

LEMMA 7. *The $\mathbb{Z}[\mathfrak{S}_n]$ module S^λ is a cyclic module generated by any e_D , $D \in \text{Tab}_\lambda$.*

It is also known that

PROPOSITION 5 ([2; Theorem 8.4]). *$\{e_D\}_{D \in \text{STab}_\lambda}$ is a \mathbb{Z} -base of S^λ .*

The action of \mathfrak{S}_n on S^λ with respect to this base $\{e_D\}_{D \in \text{STab}_\lambda}$ is determined by the action of $s_i = (i, i+1)$ on e_D for $D \in \text{STab}_\lambda$.

PROPOSITION 6 (James [2; 25.1]). *The action of s_i on e_D is calculated as follows ($D \in \text{STab}_\lambda$):*

(1) *If $i \in I_1(D)$, then $s_i e_D = -e_D$.*

(2) *If $i \in I_0(D) \cup I_0({}^t D)$, then $s_i e_D = e_{s_i D}$.*

(3) *If $i \in I_1({}^t D)$, then $s_i e_D = e_D + \sum_{D' < D} a_{D'} D'$.*

The case (3) is calculated by the Garnir relations.

3.2. **The Garnir relations.** For $D \in \text{Tab}_\lambda$, let X be a subset of the i -th column of D and Y be a subset of the $(i+1)$ -th column of D . We define, for $Z \subset \{1, 2, \dots, n\}$, \mathfrak{S}_Z to be the permutation group on the set Z . We fix a coset representative of $\mathfrak{S}_{X \cup Y} / \mathfrak{S}_X \times \mathfrak{S}_Y$ and denote it by

$\{\sigma_i\}_{1 \leq i \leq r}$. Then we define

$$G_{X,Y} = \sum_{1 \leq i \leq r} \text{sgn}(\sigma_i) \sigma_i \in \mathbf{Z}[\mathfrak{S}_n].$$

PROPOSITION 7 (James [2; Theorem 7.2]). *If $|X \cup Y| >$ the number of squares in the i -th column of D , then*

$$G_{X,Y} e_D = 0$$

which is called the Garnir relation.

§ 4. The isomorphism.

4.1. Now we can state the main result of this paper. We consider two $\mathbf{Z}[\mathfrak{S}_n]$ modules with natural bases, $S^\lambda = \langle e_D \rangle_{D \in \text{STab}_\lambda}$ and $\Gamma^\lambda = \langle g_D \rangle_{D \in \text{STab}_\lambda}$.

THEOREM. *There exists a unique isomorphism φ from S^λ to Γ^λ such that*

$$\varphi(e_D) = g_D + \sum_{D' < D} a_{D',D} g_{D'} \quad (a_{D',D} \in \mathbf{Z}).$$

PROOF. Let W_I be the subgroup of \mathfrak{S}_n generated by a subset $I \subset S$. For $\mu \in P(n)$, we set $I(\mu) = I(D_{t_{\mu, \text{Bot}}})$ and regard it as a subset of S by identifying i with s_i . Then $W_{I(\mu)}$ is the horizontal subgroup of $D_{\mu, \text{Top}}$. We also set $S_Q^\lambda = S^\lambda \otimes \mathbf{Q}$ and $\Gamma_Q^\lambda = \Gamma^\lambda \otimes \mathbf{Q}$. By Stanley [5; 4.1]

$$\langle \text{Ind}_{W_{I(\mu)}}^{\mathfrak{S}_n} \mathbf{1}, S_Q^\lambda \rangle = \#\{D \in \text{STab}_\lambda \mid I(D) \subset \widehat{I(\mu)}\}$$

where $\widehat{I(\mu)} = \{1, 2, \dots, n-1\} - I(\mu)$. This multiplicity is the Kostka number $K_{\lambda, \mu}$. On the other hand

$$\begin{aligned} \langle \text{Ind}_{W_{I(\mu)}}^{\mathfrak{S}_n} \mathbf{1}, \Gamma_Q^\lambda \rangle &= \langle \mathbf{1}, \Gamma_Q^\lambda|_{W_{I(\mu)}} \rangle \\ &= \#\{D \in \text{STab}_\lambda \mid I(D) \cap I(\mu) = \emptyset\} \\ &= \#\{D \in \text{STab}_\lambda \mid I(D) \subset \widehat{I(\mu)}\}. \end{aligned}$$

As the matrix $(K_{\lambda, \mu})_{\lambda, \mu \in P(n)}$ is invertible, we get $S_Q^\lambda \simeq \Gamma_Q^\lambda$ as $\mathbf{Q}[\mathfrak{S}_n]$ modules. The -1 eigenspace of s_i in Γ_Q^λ is $\sum_{i \in I(D)} \mathbf{Q}g_D$, and it is easy to see that if $I(D) \supset I(D_{\lambda, \text{Bot}})$ for $D \in \text{STab}_\lambda$, then $D = D_{\lambda, \text{Bot}}$. Therefore the common -1 eigenspace of $I(D_{\lambda, \text{Bot}})$ in Γ_Q^λ is $\mathbf{Q}g_{D_{\lambda, \text{Bot}}}$. But since $e_{D_{\lambda, \text{Bot}}}$ is a common -1 eigenvector of $I(D_{\lambda, \text{Bot}})$ in S_Q^λ , there is an isomorphism $\tilde{\varphi}: S_Q^\lambda \simeq \Gamma_Q^\lambda$ such that $\tilde{\varphi}(e_{D_{\lambda, \text{Bot}}}) = g_{D_{\lambda, \text{Bot}}}$.

By Lemma 7, we can define $\varphi = \tilde{\varphi}|_{S^\lambda}: S^\lambda \rightarrow \Gamma^\lambda$, which is a $\mathbf{Z}[\mathfrak{S}_n]$ homo-

morphism. We will show that this homomorphism φ has the required property by induction on $\text{ht}(D)$. If $\text{ht}(D)=0$, then $D=D_{\lambda, \text{Bot}}$ and by the construction $\varphi(e_D)$ has the required form. If $\text{ht}(D)>0$, take $j \in I_0(D)$ and define $D'=s_j D$. Then by induction

$$\varphi(e_{D'}) = g_{D'} + \sum_{D'' < D'} a_{D'', D'} g_{D''} .$$

As φ is a $Z[\mathfrak{S}_n]$ homomorphism,

$$\varphi(e_D) = \varphi(s_j e_{D'}) = s_j \varphi(e_{D'}) .$$

On the other hand

$$s_j g_{D'} = g_D + g_{D'} + \sum_{D^* < D'} b_{D^*} g_{D^*}$$

and

$$s_j g_{D''} = \begin{cases} -g_{D''} & \text{if } j \in I(D'') \\ g_{D''} + \text{lower} & \text{if } j \in I_1({}^t D'') \\ g_{D^*} + g_{D''} + \text{lower} & \text{if } j \in I_0({}^t D'') \end{cases}$$

where $D^\circ = s_j D''$ and $D'' < D^\circ < D$. Therefore $\varphi(e_D)$ also has the required form. φ is an isomorphism by this form, and uniqueness is clear.

4.2. Construction of left cells. For $D \in \text{STab}_\lambda$ we define

$$B_D = \{D' \in \text{STab}_\lambda \mid \text{ht}(D) - \text{ht}(D') = \text{odd} \geq 3 \text{ and } I(D') \supset I(D)\}$$

and

$$C(D) = \left(\bigcup_{D' \in B_D} I(D') \right) - I(D) \subset \{1, 2, \dots, n-1\} .$$

For $\lambda \in P(n)$, we define a condition (C_λ) as follows:

- (C_λ) For all $D \in \text{STab}_\lambda - \{D_{\lambda, \text{Bot}}\}$, there exists $j \in I_0(D)$ such that $j \notin C(s_j D)$.

REMARK. For $n \leq 7$, all partitions $\lambda \in P(n)$ satisfy the condition (C_λ) . But there are partitions which do not satisfy the condition for $n \geq 8$, e.g. $\lambda = (2, 2, 2, 1, 1)$.

COROLLARY 2. *If a partition λ satisfies the condition (C_λ) , we can construct sub-W-graph of the left cell Γ^λ from the λ -diagram using the Garnir relations.*

PROOF. We construct Γ^λ inductively on $\text{ht}(D)$, $D \in \text{STab}_\lambda$. For $\text{ht}(D) < 3$, λ -diagram is equal to Γ^λ . For $k \geq 3$, assume that Γ^λ is already constructed for $\text{ht}(D) < k$, so the action of s_i on g_D is already defined for $\text{ht}(D) < k$ and all i . Take $D \in \mathcal{D}_k^\lambda$. If $C(D) = \emptyset$, then no edges are added and take another $D \in \mathcal{D}_k^\lambda$. For $i \in C(D)$, we have two cases.

(1) If $i \in I_0({}^t D)$, put $D' = s_i D$. By the condition (C_i) for D' , there exists $j \in I_0(D')$ such that $j \notin C(s_j D')$. Then $\varphi(e_{D'}) = s_j \varphi(e_{s_j D'})$ is already determined in terms of $g_{D'}$, $D^\circ \leq D'$. On the other hand $\varphi(e_D) = g_D + \text{lower}$ is also determined. As $\varphi(e_{D'}) = s_i \varphi(e_D)$, we get $s_i g_D = g_{D'} + g_D + \sum_{D'' < D} \alpha_{D''} g_{D''}$. Then add all the edges $\{D, D''\}$ for $\alpha_{D''} \neq 0$ and define $\mu(D'', D) = \alpha_{D''}$.

(2) If $i \in I_1({}^t D)$, then $s_i e_D = e_D + \sum_{D^* < D} \beta_{D^*} e_{D^*}$ (calculated by the Garnir relations). $\varphi(e_D)$ is already determined and as $s_i \varphi(e_D) = \varphi(s_i e_D)$, we get $s_i g_D = g_D + \sum_{D'' < D} \gamma_{D''} g_{D''}$. Then add all the edges $\{D, D''\}$ for $\gamma_{D''} \neq 0$ and define $\mu(D'', D) = \gamma_{D''}$.

Do this operation for all $i \in C(D)$. For all $D \in \mathcal{D}_k^\lambda$ do the same operation and we get Γ^λ for $\text{ht}(D) \leq k$. Thus inductively we get Γ^λ . (If there are vertices with the same I -set, some edges may be added to become the left cell.)

§ 5. Examples.


In this section we calculate W -graphs (the left cells) and transition matrices for $\lambda = (3, 2, 1)$ and for all partitions of $n = 7$ using Corollary 2.

5.1. $\lambda = (3, 2, 1)$. The λ -diagram is indicated below. We label the Young tableaux D_1, D_2, \dots, D_{16} from top to bottom. We also write $e_i = e_{D_i}$ and $g_j = g_{D_j}$. As the action of s_i on g_D is already known for $\text{ht}(D) \leq 2$, we can calculate $\varphi(e_D)$ for $\text{ht}(D) \leq 3$ inductively:

$$\begin{aligned} \varphi(e_{16}) &= g_{16} && \text{by definition of } \varphi, \\ \varphi(e_{15}) &= s_5 \varphi(e_{16}) = s_5 g_{16} = g_{15} + g_{16}, \\ \varphi(e_{14}) &= s_3 \varphi(e_{16}) = s_3 g_{16} = g_{14} + g_{16}, \\ \varphi(e_{13}) &= s_2 \varphi(e_{14}) = s_2 (g_{14} + g_{16}) = g_{13} + g_{14}, \\ &\vdots \\ \varphi(e_7) &= s_5 \varphi(e_{11}) = g_7 + g_{11} + g_{12} + g_{14}. \end{aligned}$$

We define the action of s_i on g_D for $\text{ht}(D) = 3$. $C(D_9) = \{1\}$ and this is the case (2) of Corollary 2.

$$s_1 e_9 = e_9 - e_{11} - e_{16} \quad \text{by Garnir relation,}$$



λ -diagram for $\lambda=(3, 2, 1)$

| | | | | | | |
|---|----|-----|------|-----|----|---|
| 1 | 11 | 111 | 1111 | 110 | 01 | 1 |
| 0 | 10 | 111 | 1111 | 111 | 11 | 1 |
| 0 | 01 | 001 | 0110 | 100 | 00 | 1 |
| 0 | 00 | 100 | 1100 | 110 | 10 | 1 |
| 0 | 00 | 010 | 1011 | 111 | 11 | 1 |
| 0 | 00 | 001 | 0111 | 111 | 10 | 1 |
| 0 | 00 | 000 | 1000 | 110 | 10 | 0 |
| 0 | 00 | 000 | 0100 | 110 | 11 | 1 |
| 0 | 00 | 000 | 0010 | 101 | 10 | 1 |
| 0 | 00 | 000 | 0001 | 011 | 10 | 0 |
| 0 | 00 | 000 | 0000 | 100 | 10 | 0 |
| 0 | 00 | 000 | 0000 | 010 | 11 | 1 |
| 0 | 00 | 000 | 0000 | 001 | 10 | 0 |
| 0 | 00 | 000 | 0000 | 000 | 10 | 1 |
| 0 | 00 | 000 | 0000 | 000 | 01 | 1 |
| 0 | 00 | 000 | 0000 | 000 | 00 | 1 |

transition matrix for $\lambda=(3, 2, 1)$

and

$$s_1\varphi(e_9) = \varphi(e_9 - e_{11} - e_{16}) = g_9 + g_{13} .$$

On the other hand

$$s_1\varphi(e_9) = s_1g_9 + s_1(g_{11} + e_{13} + g_{14} + g_{16}) = s_1g_9 - g_{11} + g_{13} - g_{16} .$$

Therefore $s_1g_9 = g_9 + g_{11} + g_{16}$ and we get $\mu(D_{16}, D_9) = 1$. $C(D_8) = \{2\}$ and this is the case (1) of Corollary 2. $s_2e_8 = e_6 = s_4e_{10}$ and the action of s_4 on g_{10} is known:

$$\begin{aligned} s_4\varphi(e_{10}) &= s_4(g_{10} + g_{12} + g_{13} + g_{14}) \\ &= g_6 + g_8 + g_9 + g_{10} + g_{11} + g_{12} + g_{13} + g_{14} + g_{16} . \end{aligned}$$

On the other hand

$$\begin{aligned} s_2\varphi(e_8) &= s_2g_8 + s_2(g_{11} + g_{12} + g_{14} + g_{15} + g_{16}) \\ &= s_2g_8 + g_9 + g_{10} + g_{11} + g_{12} + g_{13} + g_{14} , \end{aligned}$$

therefore $s_2g_8 = g_6 + g_8 + g_{16}$ and we get $\mu(D_{16}, D_8) = 1$. As we defined all the actions of s_i on g_D for $\text{ht}(D) \leq 3$, we can calculate $\varphi(e_D)$ for $\text{ht}(D) = 4$. In this way we get transition matrix and W -graph simultaneously. The added edges are


$$\begin{array}{ll} D_8 \text{---} D_{16} & \text{case (1),} \\ D_9 \text{---} D_{16}, \quad D_5 \text{---} D_{15}, \quad D_8 \text{---} D_{11}, \quad D_1 \text{---} D_7, \quad D_1 \text{---} D_{10} & \text{case (2),} \end{array}$$

with multiplicities 1. This W -graph is also the left cell.

5.2. $n=7$. We indicate only half of the cases. For transposed partition ${}^t\lambda$, the Young tableaux are tD with the same numbering. The edges are the same and the I -sets are $I({}^tD) = \{1, 2, \dots, 6\} - I(D)$. The transition matrix is indicated in lower triangle, e.g. for ${}^t\lambda = (2, 2, 1, 1, 1)$, $\varphi(e_8) = g_8 + g_6 + g_5 + g_3$. Dotted edges are added for the W -graph to become the left cell.

$$\textcircled{\emptyset} \begin{matrix} 1 \\ 1234567 \end{matrix}$$

$$\begin{array}{l} \lambda = (7) \\ 1 \\ {}^t\lambda = (1, 1, 1, 1, 1, 1, 1) \end{array}$$




$\lambda = (6, 1)$

```

1 1 1 1 1 1
1 1 1 1 1 1
0 1 1 1 1 1
0 0 1 1 1 1
0 0 0 1 1 1
0 0 0 0 1 1
    
```

$\epsilon \lambda = (2, 1, 1, 1, 1, 1)$




$\lambda = (5, 1, 1)$

```

1 1 10 10 100 00 00 0 0
1 1 11 11 110 10 00 0 0
0 1 10 11 111 11 10 0 0
1 1 01 01 010 10 00 0 0
0 0 10 10 110 11 11 1 0
0 1 11 01 011 11 10 0 0
0 0 00 10 100 10 10 1 1
0 0 10 11 010 11 11 1 0
0 0 01 01 001 01 10 0 0
0 0 00 10 110 10 10 1 1
0 0 00 01 011 01 11 1 0
0 0 00 00 010 11 10 1 1
0 0 00 00 001 01 01 1 0
0 0 00 00 000 01 11 1 1
0 0 00 00 000 00 01 1 1
    
```


$\epsilon \lambda = (3, 1, 1, 1, 1)$



$\lambda = (4, 1, 1, 1)$

| | | | | | | | | | | |
|---|---|----|-----|-----|-----|-----|-----|----|---|---|
| 1 | 1 | 1 | 10 | 100 | 000 | 000 | 000 | 00 | 0 | 0 |
| 1 | 1 | 11 | 110 | 100 | 000 | 000 | 00 | 0 | 0 | 0 |
| 0 | 1 | 10 | 110 | 110 | 100 | 000 | 00 | 0 | 0 | 0 |
| 1 | 1 | 01 | 011 | 101 | 010 | 000 | 00 | 0 | 0 | 0 |
| 0 | 0 | 10 | 100 | 100 | 100 | 100 | 00 | 0 | 0 | 0 |
| 0 | 1 | 11 | 010 | 111 | 111 | 010 | 00 | 0 | 0 | 0 |
| 1 | 1 | 01 | 001 | 001 | 010 | 000 | 00 | 0 | 0 | 0 |
| 0 | 0 | 10 | 110 | 100 | 110 | 110 | 10 | 0 | 0 | 0 |
| 0 | 0 | 01 | 010 | 010 | 101 | 011 | 01 | 0 | 0 | 0 |
| 0 | 1 | 11 | 011 | 001 | 011 | 010 | 00 | 0 | 0 | 0 |
| 0 | 0 | 00 | 010 | 110 | 100 | 110 | 11 | 1 | 0 | 0 |
| 0 | 0 | 10 | 110 | 101 | 010 | 010 | 10 | 0 | 0 | 0 |
| 0 | 0 | 01 | 011 | 011 | 001 | 011 | 01 | 0 | 0 | 0 |
| 0 | 0 | 00 | 000 | 010 | 100 | 100 | 10 | 1 | 1 | 0 |
| 0 | 0 | 00 | 010 | 111 | 111 | 010 | 11 | 1 | 0 | 0 |
| 0 | 0 | 00 | 001 | 001 | 001 | 001 | 01 | 0 | 0 | 0 |
| 0 | 0 | 00 | 000 | 010 | 101 | 110 | 10 | 1 | 1 | 0 |
| 0 | 0 | 00 | 000 | 001 | 011 | 011 | 01 | 1 | 0 | 0 |
| 0 | 0 | 00 | 000 | 000 | 001 | 011 | 11 | 1 | 1 | 0 |
| 0 | 0 | 00 | 000 | 000 | 000 | 001 | 01 | 1 | 1 | 1 |

$\lambda = (4, 1, 1, 1)$




$\lambda = (5, 2)$

```

1 1 11 12 121 12 11 1
1 1 11 11 111 12 11 1
0 1 10 11 111 11 11 1
0 1 01 01 011 12 11 1
0 0 10 10 110 11 11 1
1 1 11 01 011 11 11 1
0 0 00 10 100 10 10 1
0 0 10 11 010 11 11 1
1 0 00 01 001 01 11 1
0 0 00 10 110 10 10 1
0 0 01 01 011 01 11 1
0 0 00 00 010 11 10 1
0 0 01 00 000 01 01 1
0 0 00 00 001 01 11 1
    
```


$\lambda = (2, 2, 1, 1, 1)$



$\lambda = (4, 3)$

| | | | | | | |
|---|---|----|-----|-----|-----|---|
| 1 | 1 | 11 | 121 | 111 | 111 | 2 |
| 1 | 1 | 11 | 111 | 111 | 111 | 2 |
| 0 | 1 | 10 | 110 | 111 | 111 | 1 |
| 0 | 1 | 01 | 011 | 111 | 111 | 1 |
| 0 | 0 | 10 | 100 | 100 | 110 | 1 |
| 1 | 1 | 11 | 010 | 111 | 111 | 1 |
| 0 | 0 | 01 | 001 | 001 | 011 | 1 |
| 0 | 0 | 10 | 110 | 100 | 110 | 1 |
| 1 | 0 | 00 | 010 | 010 | 101 | 1 |
| 0 | 0 | 01 | 011 | 001 | 011 | 1 |
| 0 | 0 | 01 | 010 | 110 | 100 | 1 |
| 1 | 0 | 00 | 010 | 101 | 010 | 1 |
| 0 | 0 | 10 | 010 | 011 | 001 | 1 |
| 1 | 1 | 11 | 111 | 111 | 111 | 1 |

$\mu = (2, 2, 2, 1)$




$\lambda=(3, 3, 1)$

```

1 11 1111 1111 1111 211 11 1
1 10 1110 1111 1111 110 01 1
1 01 0011 0111 1111 110 01 1
0 10 1000 1100 1100 110 00 0
0 10 0100 1010 1001 010 01 0
1 11 0010 0111 1111 111 11 1
0 01 0001 0001 0110 100 00 1
1 10 1100 1000 1000 010 00 0
0 10 1010 0100 1100 110 10 1
0 10 0110 0010 1011 111 11 1
0 01 0011 0001 0111 111 10 1
1 11 1110 1110 1000 110 10 0
0 00 0010 0101 0100 110 11 1
0 10 0110 0011 0010 101 10 1
0 01 0011 0011 0001 011 10 0
1 11 1111 1111 1110 100 10 0
0 01 0010 0111 1101 010 11 1
1 11 0010 0011 0011 001 10 0
1 11 1011 0111 1111 111 10 1
0 01 0000 0010 1001 010 01 1
1 01 0001 0011 1111 111 11 1
    
```

$\lambda=(3, 2, 2)$



$\lambda=(4, 2, 1)$

```

1 11 111 12111 111210 111101 10101 0110 01 0
1 10 111 11111 111111 111111 11111 1110 01 1
1 01 001 00110 010200 010100 10001 0100 01 0
0 10 100 11100 111110 111111 11112 1211 11 1
0 10 010 01011 101111 111111 11111 1110 01 1
1 11 001 00111 010111 011110 11011 1100 01 1
0 00 100 10000 110000 111000 11100 1200 11 1
1 10 110 01000 101110 111111 11111 1111 11 1
0 10 101 00100 010110 011111 11112 1201 11 1
0 10 011 00010 000101 010110 11011 1100 01 1
1 11 011 00001 000011 001010 01010 1000 00 1
0 00 100 11000 100000 111000 11100 1100 10 1
0 00 100 10100 010000 011000 11100 1210 11 1
1 00 000 01000 001000 100101 10111 1111 11 1
1 11 111 01110 000100 010110 11011 1101 11 1
1 10 111 01101 000010 001011 01111 1101 10 1
1 10 011 00011 000001 000010 01010 1000 00 1
0 00 010 01000 101000 100000 10100 1100 10 0
0 00 100 11100 110100 010000 11000 1100 10 1
0 00 100 11100 110010 001000 01100 1110 11 1
1 01 000 01000 001100 000100 10011 1101 11 1
1 11 111 01111 000111 000010 01011 1101 10 1
1 00 010 01001 001010 000001 00101 0101 10 0
0 00 010 01010 101100 110100 10000 1100 10 0
0 00 100 11100 110110 011010 01000 1100 11 1
0 00 010 01000 101010 101001 00100 0110 11 1
0 01 000 00000 000100 000110 00010 1001 10 1
1 01 011 01111 001111 000111 00001 0101 10 0
0 00 000 00010 000101 010110 11010 1000 10 0
0 00 010 01110 111110 111111 11101 0100 11 1
0 00 010 00000 001000 100001 00100 0010 01 1
0 01 001 00110 000100 000110 00011 0001 10 0
0 00 000 00110 010101 010110 11011 1101 10 1
0 00 010 00010 001000 100101 10101 0110 01 1
0 00 000 00010 000001 000110 11011 1101 11 1

```

 $\lambda=(3, 2, 1, 1)$

5.3. Remark. The algorithm of Corollary 2 can be ameliorated and we get W -graphs for all partitions of $n \leq 9$. We have a microcomputer program to calculate these W -graphs. In these cases, the transition matrices are all nonnegative.

ACKNOWLEDGEMENTS. The author is grateful to the referee for useful suggestions.

References

- [1] H. HILLER, *Geometry of Coxeter Groups*, Research Notes in Math., **54** (1982), Pitman.
- [2] G. D. JAMES, *The Representation Theory of the Symmetric Groups*, Lecture Notes in Math., **682** (1978), Springer-Verlag.
- [3] D. KAZHDAN and G. LUSZTIG, Representations of Coxeter groups and Hecke algebras, *Invent. Math.*, **53** (1979), 165-184.
- [4] D. E. KNUTH, *The Art of Computer Programming*, vol. 3, 5.1.4, Addison-Wesley, 1973.
- [5] R. P. STANLEY, Some aspects of groups acting on finite posets, *J. Combin. Theory Ser. A*, **32** (1982), 132-161.

Present Address:

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, OKAYAMA UNIVERSITY
TSUSHIMA-NAKA, OKAYAMA 700, JAPAN