Title	Characterization of mouse $lre1\alpha$: cloning, mRNA localization in the brain and functional analysis in a neural cell line
Author(s)	Miyoshi, Ko
Citation	
Issue Date	
oaire:version	VoR
URL	https://hdl.handle.net/11094/43697
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Research report

Characterization of mouse Ire1 α : cloning, mRNA localization in the brain and functional analysis in a neural cell line

Ko Miyoshi^{a,c,*}, Taiichi Katayama^{a,b,c}, Kazunori Imaizumi^{a,b,c}, Manabu Taniguchi^{a,c},
Yasutake Mori^{a,c}, Junichi Hitomi^{a,c}, Daishi Yui^{a,c}, Takayuki Manabe^{a,c}, Fumi Gomi^{a,c},
Takunari Yoneda^{a,c}, Masaya Tohyama^{a,c}

^aDepartment of Anatomy and Neuroscience, Graduate School of Medicine, Osaka University, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

^bTanabe Seiyaku Company Limited, Yodogawaku, Osaka 532-0031, Japan

^cCREST of Japan Science and Technology Corporation (JST), Kawaguchi, Saitama 332-0012, Japan

Accepted 26 September 2000

Abstract

In yeast, an endoplasmic reticulum (ER)-associated protein, Ire1p, is believed to initiate the unfolded protein response (UPR), that is responsible for protein folding in the ER under stressed conditions. Two mammalian homologs of Ire1p have been identified, Ire1 α and Ire1 β . We have previously reported that familial Alzheimer's disease linked presenilin-1 variants downregulate the signaling pathway of the UPR by affecting the phosphorylation of Ire1 α . In the present study, we cloned the mouse homolog of Ire1 α for generating genetically modified mice. Ire1 α was ubiquitously expressed in all mouse tissues examined, and was expressed preferentially in neuronal cells in mouse brain. This led us to investigate the effects of the downregulation of the UPR on the survival of neuronal cells under conditions of ER stress. Morphological and biochemical studies using a dominant-negative form of mouse Ire1 α have revealed that cell death caused by ER stress can be attributed to apoptosis, and that the downregulation of the UPR enhances the apoptotic process in the mouse neuroblastoma cell line, Neuro2a. Our results indicate that genetically modified mice such as transgenic mice with a dominant-negative form of Ire1 α might provide further understanding of the pathogenic mechanisms of Alzheimer's disease and other neurodegenerative disorders. © 2000 Elsevier Science B.V. All rights reserved.

Theme: Cellular and molecular biology

Topic: Gene structure and function: general

Keywords: Ire1 α ; ER stress; UPR; Alzheimer's disease; Neuro2a

1. Introduction

Various cell stresses such as perturbation of calcium homeostasis, deprivation of trophic factors and oxidative insults cause accumulation of unfolded proteins in the endoplasmic reticulum (ER). Cells respond to these ER stresses by increasing transcription of genes encoding ER-resident chaperones such as GRP78/BiP, GRP94, calreticulin and protein disulfide isomerase (PDI), which facilitate protein folding in the ER. This induction system is termed the unfolded protein response (UPR; reviewed in

Ref. [14]). In yeast, the UPR signaling is initiated by Ire1p/Ern1p, an ER-associated type I transmembrane protein that contains both serine/threonine kinase and ribonuclease domains in its cytosolic portion [1,8,13,18]. The ER-luminal portion of Ire1p is thought to sense the perturbed environment in the ER, followed by oligomerization and trans-phosphorylation of the kinase domain [12,18], and downstream signaling through the activation of a sequence-specific ribonuclease activity [13]. Two mammalian homologs of yeast Ire1p have been identified, Ire1 α [16] and Ire1 β [17]. The expression of Ire1 β is limited to the lung and the intestine, where its expression level is very low (our unpublished data). Recently we showed that familial Alzheimer's disease (FAD) linked presenilin-1 variants downregulate the signaling pathway

*Corresponding author. Tel.: +81-6-6879-3221; fax: +81-6-6879-3229.

E-mail address: miyoshi@anat2.med.osaka-u.ac.jp (K. Miyoshi).

of the UPR by affecting the phosphorylation of Ire1 α [4], suggesting that Ire1 α might be one of the components that participate in pathogenic mechanisms in brains of FAD patients.

In mammals, little has been understood about in vivo expression of Ire1 α and the mechanism by which Ire1 α senses ER stress and activates the downstream pathway of the UPR. Here, we cloned the mouse homolog of Ire1 α for generating genetically modified mice, and investigated the expression of Ire1 α mRNA in mouse tissues. In situ hybridization analysis has revealed that in mouse brain Ire1 α mRNA is expressed preferentially in neuronal cells, which led us to investigate the effects of the downregulation of the UPR in a mouse neuroblastoma cell line, Neuro2a, using a dominant-negative form of mouse Ire1 α . Morphological and biochemical studies have shown that cell death caused by ER stress can be attributed to apoptosis, and that the downregulation of the UPR enhances the apoptotic process.

2. Materials and methods

2.1. Animals

Adult ddY mice (8 weeks old) weighing ~25 g were anesthetized with ether inhalation and decapitated. Several tissues were then promptly removed. Brains were also removed from embryonic 14- and 18-day, and postnatal 1-, 7- and 14-day ddY mice.

2.2. cDNA cloning

Approximately one million colonies from a unidirectional oligo (dT) primed mouse brain cDNA library (C57 Black/6, female, 19 weeks; Stratagene) were screened by hybridizing to the insert of a mouse expressed sequence tag (EST) clone J0729C04 (Genbank accession #AU016754). The library screening was performed by standard methods.

2.3. RT-PCR

Total RNA was isolated from mouse tissues by extraction with Rneasy Mini Kit (QIAGEN) according to the manufacturer's instructions. Five micrograms of each total RNA were reverse-transcribed into first strand cDNA using Ready-To-Go You-Prime First-Strand Beads (Amersham Pharmacia) with the reaction volume of 33 μ l according to the manufacturer's instructions. One microlitre of the reaction mixture was then amplified by PCR using Taq DNA polymerase (TaKaRa) and primer pairs designed for the specific detection of mouse Ire1 α and β -Actin (as an internal control) cDNA target sequences. PCR was performed using the following conditions: after an initial denaturation of 2 min at 95°C, the amplification profile

consisted of 30 s at 95°C, 1 min annealing, 30 s at 72°C for 29 cycles (for the tissue distribution of Ire1 α), 31 cycles (for the developmental changes of Ire1 α in brain) or 18 cycles (for β -Actin). The annealing temperatures were 68°C for Ire1 α and 60°C for β -Actin. The primers used in this experiment were 5'-CCCTGATAGGTTGAATCCTGGCTATGTG-3' (sense) and 5'-AATCTATGCGCTAATCTGCTGGCCTCTG-3' (antisense) for Ire1 α and 5'-AGCTGAGAGGGAAATCGTGC-3' (sense) and 5'-ACCAGACAGCACTGTGTTGG-3' (antisense) for β -Actin. The PCR products were separated by 4% acrylamide gel electrophoresis and visualized by staining with ethidium bromide.

2.4. In situ hybridization

Frozen sections (15 μ m thick) of adult mouse brain (8 weeks old) were cut in various planes and thaw-mounted onto poly-L-lysine-coated slides. Digoxigenin-labeled cRNA probes (antisense and sense) were generated by in vitro transcription using a cDNA fragment (2980–3426 nt) of mouse Ire1 α as a template in the presence of digoxigenin-labeled dUTP (Boehringer Mannheim). Hybridization and posthybridization procedures were performed as described previously [3].

2.5. Expression plasmids

The coding region of mouse Ire1 α cDNA was cloned into pcDNA3.1(+) at *Kpn*I and *Apa*I sites. A truncated form of mouse Ire1 α cDNA, Δ Ire1 α , that lacked the sequence of kinase and ribonuclease domains, was generated and tagged with a Flag-epitope sequence at its C-terminal by PCR using wild-type cDNA as a template, and also cloned into pcDNA3.1(+) at the same sites.

2.6. Cell culture and assessment of apoptosis by Hoechst 33258 staining

Neuro2a cells, a mouse neuroblastoma cell line, were stably transfected with mock (empty plasmid), wild-type mouse Ire1 α cDNA or a truncated form, Δ Ire1 α . Two lines were selected from the stable lines transfected with mock (#3, #8), wild-type (#3, #10) and Δ Ire1 α (#2, #5), respectively. They were replated on 10-cm dishes at a density of 2×10^3 cells/cm², maintained in α -MEM (Life Technologies) containing 10% fetal calf serum at 37°C for 48 h and then treated with 2 μ g/ml tunicamycin (Sigma) or 3 μ M A23187 (Sigma) in the same medium. At 10, 20, 30 or 40 h after treatment, cells were stained with 5 μ g/ml Hoechst 33258 (Molecular Probes) in PBS for 15 min. The average percentages of cells with chromatin condensation in randomly selected fields were scored as an indicator of apoptosis under UV illumination.

A

mouse 1 MPARWLLLLLALLPPPGPSFGRTSTVTLPETLLFVSTLDGSLHAVSKRTGSIKWLKEDPVLQVPTHVEEPAFLPDPN
human 1 MPARWLLLLLALL—PGGIFGRTSTVTLPETLLFVSTLDGSLHAVSKRTGSIKWLKEDPVLQVPTHVEEPAFLPDPN

mouse 81 DGSlyTLGGKNEGLTKLPFTIPELVQASPCRSSDGILYMGKKQDIWYVIDLLTGEKQQLSSAFADSLCPSTSLLYLGR
human 79 DGSlyTLGSKNEGLTKLPFTIPELVQASPCRSSDGILYMGKKQDIWYVIDLLTGEKQQLSSAFADSLCPSTSLLYLGR

mouse 161 TEYTITMYDTKTRELWNATYFDYAASLPEDDGDYKMSHFVSNQDGLVVTVDSESGDVLWIQNYASPVVAFYVWQGEVLR
human 159 TEYTITMYDTKTRELWNATYFDYAASLPEDDGDYKMSHFVSNQDGLVVTVDSESGDVLWIQNYASPVVAFYVWQGEVLR

mouse 241 KVVHINVAVETLRYLTFMSGVGRITKWYFPKETEAKSKLTPTLYVGKYSTSLYASPMVEGVAVVPRGSTLPLLEG
human 239 KVVHINVAVETLRYLTFMSGVGRITKWYFPKETEAKSKLTPTLYVGKYSTSLYASPMVEGVAVVPRGSTLPLLEG

mouse 321 PQTDGVTIGDKGECVITPSTDLKFDPLGKSKNLNRYWLLIGHHETPLSASTKMLERFPNNLPKHRENVIPADSEKR
human 319 PQTDGVTIGDKGECVITPSTDLKFDPLGKSKNLNRYWLLIGHHETPLSASTKMLERFPNNLPKHRENVIPADSEKR

mouse 401 SFEEVINIVGQTSNDTPPTVSQDVEEKLARAPAKPEAPVDSMLKDMATIILSTFLLVGWVAFIITYPLSVHQQLQHQHQQ
human 399 SFEEVINIVGQTSNDTPPTVSQDVEEKLARAPAKPEAPVDSMLKDMATIILSTFLLVGWVAFIITYPLSVHQQLQHQHQQ

mouse 481 FQKELEKIQLL—QQQLPFHPHGLTDQPEFLDSSGPFSESSGTSSPSPRASNHSHPSSASRAGTSPSLEQDDED
human 479 FQKELEKIQLLQQQQQLPFHPHGLTDQPEFLDSSGPFSESSGTSSPSPRASNHSHPSSASRAGTSPSLEQDDED

mouse 559 EETRMVIVGKISFCPKDVLGHGAEGTIVYKGMFNDNRDVAVKRILPECFSFADREVQLLRESDEHPNVIRYFCTEKDRQFQ
human 559 EETRMVIVGKISFCPKDVLGHGAEGTIVYKGMFNDNRDVAVKRILPECFSFADREVQLLRESDEHPNVIRYFCTEKDRQFQ

mouse 639 YIAIELCAATLQEYVEQKDFAHGLPEITLLHQTTSGLAHLHSLNIVHRDLKPHNILLSMPNAHGRIKAMISDFGLCKKL
human 639 YIAIELCAATLQEYVEQKDFAHGLPEITLLHQTTSGLAHLHSLNIVHRDLKPHNILLSMPNAHGRIKAMISDFGLCKKL

mouse 719 AVGRHSFRRSGVPGTEGWIAPEMLSEDCKDNPTYTVDIFSAGCVFYVYISEGNHFPFGKSLQRQANILLGACNLDCFHSD
human 719 AVGRHSFRRSGVPGTEGWIAPEMLSEDCKDNPTYTVDIFSAGCVFYVYISEGNHFPFGKSLQRQANILLGACNLDCFHSD

mouse 799 KHEDVIARELIEKMIAMDPOQRPSAKHVLKHPFFWSLEKQLQFFQDVSDRIEALDGPVIRQLERGGRAVVKMDWRENI
human 799 KHEDVIARELIEKMIAMDPOQRPSAKHVLKHPFFWSLEKQLQFFQDVSDRIEALDGPVIRQLERGGRAVVKMDWRENI

mouse 879 TVPLQTDLRKFRTYKGSVRDLLRAMRNKKHYRELPEVEQETLGSIPDDFVRYFTSRFPHLLSHTYQAMELCRHERLFQ
human 879 TVPLQTDLRKFRTYKGSVRDLLRAMRNKKHYRELPEVEQETLGSIPDDFVRYFTSRFPHLLSHTYQAMELCRHERLFQ

mouse 959 TYYWHEPTEPQPPVIPAL 977
human 959 TYYWHEPTEPQPPVIPAL 977

B

Fig. 1. Amino acid sequence analysis and structure of mouse Ire1 α . (A) Alignment of mouse and human Ire1 α amino acid sequences. Black boxes and shaded boxes represent nonconserved and conserved amino acids, respectively. All other amino acids are identical. Mouse and human sequences share a 92% identity. GenBank accession number for the mouse cDNA sequence is AB031332. (B) Domain organization of mouse Ire1 α . SP and TM refer to the potential signal peptide and the putative transmembrane domain, respectively.

Fig. 2. The expression of mouse *Ire1α* mRNA was investigated by RT-PCR. Total RNA was isolated from different tissues of adult mice (A) and brains of mice at several developmental stages (B), and reverse-transcribed into cDNA, that was then amplified by PCR using a primer pair designed for the specific detection of mouse *Ire1α*. β -Actin was also examined as an internal control. E14, E18, P1, P7, P14 and 8W refer to embryonic 14 and 18 days, and postnatal 1, 7 and 14 days, and 8 weeks old, respectively.

2.7. Electron microscopy

Neuro2a cells expressing Δ *Ire1α* were treated with 2 μ g/ml tunicamycin for 40 h. PBS containing 2% glutaraldehyde and 4% paraformaldehyde was then employed in fixing the cells at 4°C for 45 min. They were post-fixed in 1% OsO_4 at 25°C for 30 min, dehydrated in the graded ethanol solutions and embedded in Quetol 812 (NISSHIN EM Co.). Areas containing cells with apoptotic morphology were block-mounted and cut into 80-nm sections. The sections were stained with uranyl acetate (10% in 50% ethanol) and lead citrate, and then examined with a Hitachi H-7100 electron microscope.

2.8. Measurement of caspase-3-like activity

Stable Neuro2a lines treated with 2 μ g/ml tunicamycin for 40 h were washed with PBS, harvested and lysed by ICE buffer (50 mM Tris-HCl (pH 7.4), 1 mM EDTA and 10 mM EGTA) containing 10 μ M digitonin. The lysates were then incubated at 37°C for 10 min and centrifuged at 15,000 rev./min for 10 min. Each sample (50 μ g protein) was incubated with 50 μ M of Ac-DEVD-MCA (Peptide Institute, Inc.), the substrate for caspase-3, in ICE buffer with the reaction volume of 1 ml at 37°C for 30 min. Levels of released 7-amino-4-methylcoumarin (AMC) were measured using a spectrofluorometer with excitation at 380 nm and emission at 460 nm.

2.9. Western blot analysis

The expression levels of wild-type *Ire1α* and Δ *Ire1α* in stable Neuro2a lines were demonstrated by Western blot analysis. Cells were washed with PBS, harvested and lysed in NP-40 lysis buffer (1% NP-40, 10 mM Tris-HCl (pH 7.8), 150 mM NaCl, 1 mM EDTA, 1 mM phenylmethylsulphonyl fluoride (PMSF) and 10 μ g/ml aprotinin). The lysates were then incubated on ice for 50 min, centrifuged at 15,000 $\times g$ for 20 min and boiled for 5 min with SDS sample buffer. Equal amounts of protein (30 μ g) were subjected to 10% SDS-PAGE, transferred to PVDF mem-

brane and blotted with rabbit polyclonal anti-*Ire1α* antibody, which was made by injecting rabbits with the cytosolic portion of *Ire1α*-GST fusion protein. The membrane was then incubated with alkaline phosphatase-conjugated sheep anti-rabbit IgG antibody (Boehringer Mannheim). For Western blot analysis of caspase-3, Neuro2a lines which were treated with 2 μ g/ml tunicamycin for 40 h were lysed in NP-40 lysis buffer. The samples were prepared by the same procedures as the analysis of *Ire1α*. Equal amounts of protein (30 μ g) were subjected to 12% SDS-PAGE, and Western blot analysis was performed using rabbit polyclonal anti-caspase-3 antibody (Santa Cruz), which recognizes pro-caspase-3 and active subunits, as the primary antibody, and anti-rabbit IgG antibody as the secondary antibody. For Western blot analysis of cytochrome *c*, Neuro2a cells expressing Δ *Ire1α* which were treated with 2 μ g/ml tunicamycin for 12, 24 or 40 h were harvested by centrifugation, and the cell pellets then washed with PBS and resuspended with buffer A (50 mM Tris-HCl pH 8.0, 1 mM EDTA, 0.32 M sucrose and 0.1 mM PMSF). The cells were homogenized with 10 strokes of a Teflon homogenizer, and the homogenates then separated as cytosol and membrane fractions by ultracentrifugation at 105,000 $\times g$ for 60 min. Equal amounts of protein (6 μ g) from the cytosol fraction were subjected to 12% SDS-PAGE, and Western blot analysis was performed using mouse monoclonal anti-cytochrome *c* antibody (PharMingen) as the primary antibody, and goat anti-mouse IgG antibody (Sigma) as the secondary antibody.

3. Results

3.1. Isolation of complementary DNA encoding mouse *Ire1α*

A mouse expressed sequence tag (EST) clone J0729C04 was found to have significant sequence homology to a portion of human *Ire1α* cDNA. The clone was used as the probe to screen a mouse brain cDNA library (C57 Black/

6, female, 19 weeks; Stratagene) and an ~4.0-kb cDNA that contains an ORF encoding 977 amino acid residues was obtained from several overlapping clones and deposited in the DDBJ/EMBL/GenBank (accession #AB031332). The amino acid sequence had a 92% identity to that of human Ire1 α , indicating that the cDNA encodes a mouse homolog of Ire1 α (Fig. 1A). Domain organization of mouse Ire1 α is shown in Fig. 1B. The encoded protein has an N-terminal hydrophobic stretch of 20 residues, thought to function as a signal peptide, followed by the sensor domain. A 21-residue hydrophobic stretch, predicted to serve as a transmembrane domain, locates approximately in the middle of the molecule. The C-terminal half of the protein contains a linker region, the serine/threonine kinase domain and the ribonuclease domain. The amino acid sequence identity between mouse Ire1 α and β was 49%.

3.2. The expression of mouse Ire1 α mRNA

The distribution of Ire1 α mRNA was analyzed by RT-PCR on different tissues from adult mouse (Fig. 2A). Ire1 α was ubiquitously expressed in all tissues examined; it was expressed at relatively high levels in thymus, liver and lung. To examine the changes of Ire1 α mRNA expression during brain development, RT-PCR was performed on the brains of embryonic 14 (E14) and 18 (E18) day, and postnatal 1 (P1), 7 (P7) and 14 (P14) day, and 8-week-old (8W) mice (Fig. 2B). A gradual increase in the expression of Ire1 α mRNA along brain development was observed before P1, while the expression level was almost equivalent through the postnatal and adult stages.

3.3. In situ localization of Ire1 mRNA in mouse brain

In situ hybridization using cRNA probes was carried out to analyze the distribution of cells expressing Ire1 α mRNA in mouse brain (Fig. 3). Fresh-frozen sections of brains from adult mice (8 weeks old) were hybridized with digoxigenin-labeled cRNA antisense and sense probes. Ire1 α mRNA was expressed in cortical neurons (Fig. 3A and B), hippocampal neurons (Fig. 3A and C) and olfactory neurons (Fig. 3D and E). Signal intensity on cortical neurons was equivalent in all sections examined. In hippocampus, Ire1 α mRNA was expressed in pyramidal cells of CA1-3 and granule cells of dentate gyrus, with the expression level in these hippocampal neurons being equivalent. In olfactory bulb, signals with relatively weak intensity were detected in mitral and granule cells. Very weak signals for Ire1 α mRNA were diffusely detected in other neuronal cells in mouse brain containing Purkinje cells and brainstem neurons. Signals were undetectable in oligodendrocytes and astrocytes.

3.4. The involvement of apoptosis in cell death induced by the downregulation of the UPR

The preferential expression of Ire1 α in neuronal cells in mouse brain led us to perform function analysis of Ire1 α in a mouse neural cell line. In yeast, overexpression of a truncated form of Ire1p that lacks an effector domain can downregulate the activation of the pathway that leads to the induction of *KAR2/GRP78* [8,12]. In mammalian cells, derivatives of Ire1 α and β that lack the cytosolic portion have been shown to demonstrate similar dominant-negative effects on the UPR [8,10]. In the present study, a truncated form of mouse Ire1 α , Δ Ire1 α , that lacked kinase and ribonuclease domains (Fig. 4A) was used to interfere with the activity of the endogenous inducers of the UPR in Neuro2a cells, a mouse neuroblastoma cell line. We confirmed that the levels of GRP78 mRNA induction in Neuro2a cells expressing Δ Ire1 α were lower than those in cells expressing wild-type Ire1 α or mock (empty plasmid) after treatment with tunicamycin or A23187 (data not shown), indicating dominant-negative effects of Δ Ire1 α on the UPR; tunicamycin and A23187 are known to provoke ER stress by inhibition of protein glycosylation or perturbation of calcium homeostasis, respectively. To analyze the effects of the downregulation of the UPR, we examined the time-course of cell viability under conditions of ER stress in Neuro2a cells by investigating nuclear morphology with Hoechst 33258 staining (Fig. 4C and D). Neuro2a cells expressing a truncated form of Ire1 α , Δ Ire1 α , displayed chromatin condensation, that is the morphological feature of apoptosis, at a significantly higher rate than cells expressing wild-type or mock at 20 h or later after treatment with 2 μ g/ml tunicamycin (Fig. 4C). A significantly increased rate of apoptotic morphology of cells expressing Δ Ire1 α was also observed after treatment with 3 μ M A23187 (Fig. 4D), suggesting that the downregulation of the UPR enhances the apoptotic process under conditions of ER stress in Neuro2a cells. The morphological change described above that was revealed by Hoechst 33258 staining is shown in Fig. 5A. Neuro2a cells expressing Δ Ire1 α treated with tunicamycin were also examined under electron microscopy (Fig. 5B). They exhibited changes consistent with apoptosis, such as loss of cytoplasm and nuclear fragmentation. Enlarged ER lumens were also observed, indicating that the cells were under conditions of ER stress.

In mammalian cells, a family of cysteine proteases termed caspases are known to be activated during the apoptotic process (reviewed in Ref. [15]). To examine the biochemical features of cell death induced by the downregulation of the UPR, we measured caspase-3-like protease activity in ER-stress exposed Neuro2a cells (Fig. 6A). At 40 h after treatment with tunicamycin, caspase-3-like activity in Neuro2a cells expressing Δ Ire1 α was significantly higher than that in cells expressing wild-type

Fig. 3. In situ hybridization analysis of Ire1 α mRNA in mouse brain. Signals were detected in neuronal cells of the cortex and hippocampus (A), and in the olfactory bulb (D). High magnification views are shown for cortical neurons (B), hippocampal CA1 pyramidal cells (C), and olfactory mitral and granule cells (E). DG, dentate gyrus; py, pyramidal cell layer; gr, granule cell layer; mi, mitral cell layer. Scale bars: 100 μ m for (A), (D); 10 μ m for (B), (C); 20 μ m for (E).

or mock. Caspase-3 is synthesized as a 32-kDa precursor (pro-caspase-3), that is activated proteolytically into 20, 19 or 17 kDa subunits when cells are undergoing the apoptotic process [2]. To confirm the involvement of caspase-3 in cell death induced by the downregulation of the UPR, we examined the cleavage of caspase-3 in ER-stress exposed Neuro2a cells by Western blot analysis using an anti-caspase-3 antibody that can detect the proform and active subunits (Fig. 6B). The processing of caspase-3 was observed in Neuro2a cells treated with tunicamycin for 40 h. Active fragments of caspase-3 were more abundant in cells expressing Δ Ire1 α than in cells expressing wild-type or mock, suggesting that caspases contribute to cell death enhanced by the downregulation of the UPR in Neuro2a cells.

When cells are exposed to stimuli that trigger apoptosis,

cytochrome *c* is rapidly released from mitochondria into the cytosol where it activates caspases [5–7,19]. We therefore examined cytosolic cytochrome *c* in ER-stress exposed Neuro2a cells expressing Δ Ire1 α by Western blot analysis (Fig. 6C). Cytochrome *c* was detected in the cytosol fraction at 12 h after treatment with tunicamycin, and became more abundant at 24 and 40 h. This cytochrome *c* release into the cytosol from mitochondria preceded the morphological changes of cells demonstrated in Fig. 4, suggesting that cytochrome *c* might play a key role in the course of cell death enhanced by the downregulation of the UPR in Neuro2a cells.

These morphological and biochemical findings suggest that cell death caused by ER stress in Neuro2a cells can be attributed to apoptosis and that the downregulation of the UPR enhances the apoptotic process.

Fig. 4. The time-course of cell viability under conditions of ER stress was examined by investigating nuclear morphology. A truncated form of mouse Ire1 α , Δ Ire1 α , which lacks kinase and ribonuclease domains, was used for its dominant-negative effects on the UPR. The structures of Δ Ire1 α and wild-type Ire1 α are shown in (A). SP and TM refer to the potential signal peptide and the putative transmembrane domain, respectively. Neuro2a cells, a mouse neuroblastoma cell line, were stably transfected with mock (empty plasmid), wild-type Ire1 α or Δ Ire1 α . Two lines were selected from the stable lines transfected with mock (#3, #8), wild-type (#3, #10) and Δ Ire1 α (#2, #5), respectively. The expression levels of the transgenes in these lines were demonstrated by Western blot analysis (B). Arrows indicate wild-type Ire1 α and Δ Ire1 α . Size markers (in kilodaltons) are on the left. At indicated time points after treatment with 2 μ g/ml tunicamycin (C) or 3 μ M A23187 (D), the stable lines were stained with Hoechst 33258, and the average percentages of cells with chromatin condensation in randomly selected fields were scored as an indicator of apoptosis. Data are means $\pm$ S.D. of four independent experiments.

4. Discussion

In the present study, we cloned the mouse homolog of Ire1 α for generating genetically modified mice. From the 92% identity of the amino acid sequence between mouse and human Ire1 α , and the ubiquitous expression of Ire1 α mRNA in mouse tissues, it is likely that Ire1 α protein is a main stress sensor that plays an essential role in cell survival against ER stress in mammalian organs. Relatively high expression levels of Ire1 α were observed in thymus, liver and lung, suggesting the possibility that in these organs Ire1 α might participate in organ-specific stress response systems as well as the UPR. The low oxygen environment in embryonic circulation might be connected with the lower expression levels of Ire1 α mRNA in mouse brain at embryonic stages compared to postnatal and adult stages.

Like many types of nonneuronal cells, neurons respond to extracellular stimuli that provoke stressed conditions in the ER. For example, transient cerebral ischemia is known to induce a stress response which is in many respects

identical to that triggered by the depletion of ER calcium pools, implying the possibility that disturbance of the ER calcium homeostasis may play a major role in ischemia-induced cell injury of neurons [9–11]. In the present study, in situ hybridization analysis has revealed that in mouse brain Ire1 α mRNA is preferentially expressed in cortical, hippocampal and olfactory neurons and in other neuronal cells containing Purkinje cells and brainstem neurons, suggesting that Ire1 α is the essential proximal sensor of the UPR pathway in neuronal cells.

We have previously reported that familial Alzheimer's disease (FAD) linked *presenilin-1* mutations downregulate the signaling pathway of the UPR by affecting the phosphorylation of Ire1 α [4]. However, the character of the resulting vulnerability to ER stress has been unclear. Our present study, using a dominant-negative form of Ire1 α , has revealed that the downregulation of the UPR enhances the apoptotic process under conditions of ER stress in a mouse neural cell line, by investigating the morphological changes of cells, activation of caspase-3 and cytochrome *c* release from mitochondria into the cytosol. Together with

Fig. 5. Morphological study of Neuro2a cells expressing Δ Ire1 α treated with 2 μ g/ml tunicamycin for 40 h. (A) Hoechst 33258 staining of tunicamycin-treated cells (right panel) and untreated Neuro2a cells (left panel). Arrows indicate the cells with chromatin condensation, which is the morphological feature of apoptosis. (B) Examination under electron microscopy of treated cells (right panel) and untreated cells (left panel). Enlarged ER lumens are indicated by arrows. Scale bars: 2 μ m.

the preferential expression of Ire1 α in neuronal cells in mouse brain, this finding raises the possibility that the perturbed ER-stress response might be involved in the neuron-specific apoptosis observed in brains of Alzheimer's disease patients containing FAD ones. Enhanced apoptosis by the downregulation of the UPR in a mouse neural cell line suggests that transgenic mice with a dominant-negative form of Ire1 α targeted to neurons would be a good animal model for evaluating the effects of the disturbed stress response on the mammalian nervous system, and that generating genetically modified mice on Ire1 α including dominant-negative form transgenic mice might contribute to further understanding of the pathogenic mechanisms of Alzheimer's disease and other neurodegenerative disorders.

In our analysis by in situ hybridization for Ire1 α mRNA,

signals were undetectable in oligodendrocytes and astrocytes, implying the possibility that the major ER-stress sensing molecule in these types of cells might not be Ire1 α , or that they might mediate ER stress by a mechanism different to the UPR.

In the present study, we have shown that the downregulation of the UPR enhances the apoptotic process in a neural cell line. This apoptotic process is thought to be triggered by the accumulation of unfolded proteins in the ER lumen, which cannot undergo folding and maturation due to the inhibition of the ER-resident chaperones' induction. However, the signaling pathway by which the stress in the ER lumen activates caspases which reside in the cytosol is unclear, even though our results suggest the involvement of cytochrome *c* release from mitochondria into the cytosol. Further studies should be performed to

Fig. 6. The activation of caspase-3 and the release of cytochrome *c* from mitochondria into the cytosol were analyzed as indicators of apoptosis to investigate the effects of the downregulation of the UPR. (A) Caspase-3-like protease activity in ER-stress exposed Neuro2a lines stably transfected with mock (#3), wild-type Ire1 α (#3) or Δ Ire1 α (#2). At 40 h after treatment with 2 μ g/ml tunicamycin, Ac-DEVD-MCA cleaving activity in the cell extracts was measured. Values represent caspase-3-like activities relative to untreated Neuro2a cells expressing mock, and are shown as means $\pm$ S.D. of three independent experiments. (B) The cleavage of caspase-3 in ER-stress exposed Neuro2a cells expressing mock, wild-type Ire1 α or Δ Ire1 α . Following treatment with 2 μ g/ml tunicamycin for 40 h, the cell lysates which contained equal amounts of protein (30 μ g) were subjected to Western blot analysis using an anti-caspase-3 antibody that can detect both pro-caspase-3 and active subunits. Arrows indicate pro-caspase-3 (p32) and active subunits (p19, p17). Size markers (in kilodaltons) are on the left. (C) The release of cytochrome *c* from mitochondria into the cytosol in ER-stress exposed cells. At indicated time points after treatment with 2 μ g/ml tunicamycin, Neuro2a cells expressing Δ Ire1 α were harvested, homogenized and separated as cytosol and membrane fractions by ultracentrifugation. Equal amounts of protein (6 μ g) from the cytosol fraction were subjected to Western blot analysis using an anti-cytochrome *c* antibody.

investigate the mechanisms by which the accumulation of unfolded proteins in the ER provokes downstream apoptotic signaling in neuronal cells.

References

- [1] J.S. Cox, C.E. Shamu, P. Walter, Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase, *Cell* 73 (1993) 1197–1206.
- [2] T. Fernandes-Alnemri, R.C. Armstrong, J. Krebs, S.M. Srinivasula, L. Wang, F. Bullrich, L.C. Fritz, J.A. Trapani, K.J. Tomaselli, G. Litwack, E.S. Alnemri, In vitro activation of CPP32 and Mch3 by Mch4, a novel human apoptotic cysteine protease containing two FADD-like domains, *Proc. Natl. Acad. Sci. USA* 93 (1996) 7464–7469.
- [3] K. Imaizumi, M. Tsuda, A. Wanaka, M. Tohyama, T. Takagi, Differential expression of sgk mRNA, a member of the Ser/Thr protein kinase gene family, in rat brain after CNS injury, *Mol. Brain Res.* 26 (1994) 189–196.
- [4] T. Katayama, K. Imaizumi, N. Sato, K. Miyoshi, T. Kudo, J. Hitomi, T. Morihara, T. Yoneda, F. Gomi, Y. Mori, Y. Nakano, J. Takeda, T. Tsuda, Y. Itoyama, O. Murayama, A. Takashima, P. St George-Hyslop, M. Takeda, M. Tohyama, Presenilin-1 mutations downregulate the signalling pathway of the unfolded-protein response, *Nat. Cell Biol.* 1 (1999) 479–485.
- [5] R.M. Kluck, E. Bossy-Wetzel, D.R. Green, D.D. Newmeyer, The release of cytochrome *c* from mitochondria: a primary site for Bcl-2 regulation of apoptosis, *Science* 275 (1997) 1132–1136.
- [6] R.M. Kluck, S.J. Martin, B.M. Hoffman, J.S. Zhou, D.R. Green, D.D. Newmeyer, Cytochrome *c* activation of CPP32-like proteolysis plays a critical role in a *Xenopus* cell-free apoptosis system, *EMBO J.* 16 (1997) 4639–4649.
- [7] X. Liu, C.N. Kim, J. Yang, R. Jemmerson, X. Wang, Induction of apoptotic program in cell-free extracts: requirement for dATP and cytochrome *c*, *Cell* 86 (1996) 147–157.
- [8] K. Mori, W. Ma, M.J. Gething, J. Sambrook, A transmembrane protein with a cdc2+/CDC28-related kinase activity is required for signaling from the ER to the nucleus, *Cell* 74 (1993) 743–756.
- [9] W. Paschen, Disturbances of calcium homeostasis within the endoplasmic reticulum may contribute to the development of ischemic-cell damage, *Med. Hypotheses* 47 (1996) 283–288.
- [10] W. Paschen, J. Doutheil, C. Gissel, M. Treiman, Depletion of neuronal endoplasmic reticulum calcium stores by thapsigargin: effect on protein synthesis, *J. Neurochem.* 67 (1996) 1735–1743.
- [11] W. Paschen, J. Doutheil, Disturbance of endoplasmic reticulum functions: a key mechanism underlying cell damage?, *Acta. Neurochir. Suppl.* 73 (1999) 1–5.
- [12] C.E. Shamu, P. Walter, Oligomerization and phosphorylation of the Ire1p kinase during intracellular signaling from the endoplasmic reticulum to the nucleus, *EMBO J.* 15 (1996) 3028–3039.
- [13] C. Sidrauski, P. Walter, The transmembrane kinase Ire1p is a site-specific endonuclease that initiates mRNA splicing in the unfolded protein response, *Cell* 90 (1997) 1031–1039.
- [14] C. Sidrauski, R. Chapman, P. Walter, The unfolded protein response: an intracellular signalling pathway with many surprising features, *Trends Cell Biol.* 8 (1998) 245–249.
- [15] N.A. Thornberry, Y. Lazebnik, Caspases: enemies within, *Science* 281 (1998) 1312–1316.
- [16] W. Tirasophon, A.A. Welihinda, R.J. Kaufman, A stress response pathway from the endoplasmic reticulum to the nucleus requires a novel bifunctional protein kinase/endoribonuclease (Ire1p) in mammalian cells, *Genes Dev.* 12 (1998) 1812–1824.
- [17] X.Z. Wang, H.P. Harding, Y. Zhang, E.M. Jolicoeur, M. Kuroda, D. Ron, Cloning of mammalian Ire1 reveals diversity in the ER stress responses, *EMBO J.* 17 (1998) 5708–5717.
- [18] A.A. Welihinda, R.J. Kaufman, The unfolded protein response pathway in *Saccharomyces cerevisiae*. Oligomerization and trans-autophosphorylation of Ire1p(En1p) are required for kinase activation, *J. Biol. Chem.* 271 (1996) 18181–18187.
- [19] J. Yang, X. Liu, K. Bhalla, C.N. Kim, A.M. Ibrado, J. Cai, T.I. Peng, D.P. Jones, X. Wang, Prevention of apoptosis by Bcl-2: release of cytochrome *c* from mitochondria blocked, *Science* 275 (1997) 1129–1132.