

Title	Lifespan for radially symmetric solutions to systems of semilinear wave equations with multiple speeds
Author(s)	Katayama, Soichiro
Citation	Osaka Journal of Mathematics. 2008, 45(3), p. 691-717
Version Type	VoR
URL	https://doi.org/10.18910/4620
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

LIFESPAN FOR RADially SYMMETRIC SOLUTIONS TO SYSTEMS OF SEMILINEAR WAVE EQUATIONS WITH MULTIPLE SPEEDS

SOICHIRO KATAYAMA

(Received June 13, 2007)

Abstract

We consider the Cauchy problem for a system of semilinear wave equations with multiple propagation speeds in three space dimensions. We obtain the sharp lower bound for the lifespan of radially symmetric solutions to a class of these systems. We also show global existence of radially symmetric solutions to another class of systems with small initial data.

1. Introduction and the main results

For $c > 0$, we define

$$\square_c = \partial_t^2 - c^2 \Delta_x = \partial_0^2 - c^2 \sum_{j=1}^3 \partial_j^2,$$

where $\partial_0 = \partial_t = \partial/\partial t$, and $\partial_j = \partial/\partial x_j$ for $j = 1, 2, 3$. The above constant c is called the propagation speed. We simply write \square for $\square_1 = \partial_t^2 - \Delta_x$.

This paper is devoted to a study on the Cauchy problem for systems of semilinear wave equations in three space dimensions of the type

$$(1.1) \quad \square_{c_i} u_i = F_i(u, \partial u) \quad \text{for } (t, x) \in (0, \infty) \times \mathbb{R}^3 \quad (i = 1, \dots, m)$$

with initial data

$$(1.2) \quad u_i(0, x) = \varepsilon f_i(x), \quad (\partial_t u_i)(0, x) = \varepsilon g_i(x)$$

for $x \in \mathbb{R}^3$ ($i = 1, \dots, m$), where c_i ($1 \leq i \leq m$) are given positive constants, $u = (u_j)_{1 \leq j \leq m}$, and $\partial u = (\partial_a u_j)_{1 \leq j \leq m, 0 \leq a \leq 3}$, while ε is a small positive parameter. In the following, we assume that $F(u, v) = (F_j(u, v))_{1 \leq j \leq m}$ is a smooth function of $(u, v) \in \mathbb{R}^m \times \mathbb{R}^{4m}$, vanishing together with its first derivatives at $(u, v) = (0, 0)$. We suppose

2000 Mathematics Subject Classification. Primary 35L70; Secondary 35L05, 35L15, 35L55.

This research is partially supported by Grant-in-Aid for Young Scientists (B) (No. 16740094), MEXT.

$f = (f_j)_{1 \leq j \leq m}$, $g = (g_j)_{1 \leq j \leq m} \in C_0^\infty(\mathbb{R}^3; \mathbb{R}^m)$. For simplicity of exposition, we also suppose that the propagation speeds c_i ($1 \leq i \leq m$) are distinct.

Let $T_\varepsilon = T_\varepsilon(f, g, F)$ be the supremum of all T such that the Cauchy problem (1.1)–(1.2) admits a C^∞ -solution $u = (u_j)_{1 \leq j \leq m}$ for $(t, x) \in [0, T) \times \mathbb{R}^3$. T_ε is called the lifespan of the classical solution to the Cauchy problem (1.1)–(1.2). We say that *small data global existence* (or (SDGE)) holds for (1.1)–(1.2) if for any $f, g \in C_0^\infty(\mathbb{R}^3; \mathbb{R}^m)$, there exists a positive constant ε_0 such that $T_\varepsilon(f, g, F) = \infty$ for all $\varepsilon \in (0, \varepsilon_0]$. When $T_\varepsilon(f, g, F) < \infty$, we say that the solution blows up in finite time.

For the following single wave equation

$$(1.3) \quad \begin{cases} \square u = (\partial_t u)^2 \text{ (or } u(\partial_t u)) & \text{in } (0, \infty) \times \mathbb{R}^3, \\ u(0, x) = \varepsilon f(x), \quad (\partial_t u)(0, x) = \varepsilon g(x) & \text{for } x \in \mathbb{R}^3, \end{cases}$$

it is known that there exist $f, g \in C_0^\infty(\mathbb{R}^3)$ and two positive constants C_1 and ε_1 such that

$$(1.4) \quad T_\varepsilon \leq \exp(C_1 \varepsilon^{-1})$$

for any $\varepsilon \in (0, \varepsilon_1]$ (see John [3], Sideris [18], and Kubo [14]). In other words, for such f and g , the solution to (1.3) blows up in finite time no matter how small ε is. The above upper bound for the lifespan T_ε is sharp in the sense that for any $f, g \in C_0^\infty(\mathbb{R}^3)$, there exist two positive constants C_2 and ε_2 such that

$$(1.5) \quad T_\varepsilon \geq \exp(C_2 \varepsilon^{-1})$$

for any $\varepsilon \in (0, \varepsilon_2]$ (see John-Klainerman [4], and Klainerman [10]; see also Lindblad [16] for the case $m = 1$ and $F(u, 0) = O(|u|^3)$ for small u , and the author [6] for the case $m \geq 2$ and $F(u, \partial u) = O(|u|^3 + |\partial u|^2)$ around $(u, \partial u) = (0, 0)$).

The above example (1.3) shows that some restriction on F is necessary for (SDGE). To recall known results for (SDGE), we introduce several types of nonlinearities. Let $\phi = (\phi_i)_{1 \leq i \leq m}$ and $\psi = (\psi_i)_{1 \leq i \leq m}$ be C^2 -functions. In the following, $\alpha_{N,i}^j$, $\alpha_{I,i}^{jk,ab}$, $\alpha_{II,i}^{j,ab}$, $\alpha_{III,i}^{jk,a}$ and $\alpha_{IV,i}^{j,a}$ are arbitrary constants. First of all, we introduce *null terms*

$$(1.6) \quad N_i(\partial\phi, \partial\psi) = \sum_{j=1}^m \alpha_{N,i}^j Q_0(\phi_j, \psi_j; c_j),$$

where $Q_0(v, w; c)$ is the null form defined by

$$Q_0(v, w; c) = (\partial_t v)(\partial_t w) - c^2 \sum_{k=1}^3 (\partial_k v)(\partial_k w)$$

(see Klainerman [11]; note that another type of the null form

$$Q_{ab}(v, w) = (\partial_a v)(\partial_b w) - (\partial_b v)(\partial_a w),$$

which was also introduced in [11], does not appear here, just because we have restricted our attention to the simplified situation of semilinear systems with distinct speeds). Next we introduce

$$(1.7) \quad R_i^I(\partial\phi, \partial\psi) = \sum_{\substack{1 \leq j, k \leq m \\ j \neq k}} \sum_{0 \leq a, b \leq 3} \alpha_{I,i}^{jk,ab} (\partial_a \phi_j)(\partial_b \psi_k),$$

$$(1.8) \quad R_i^{II}(\partial\phi, \partial\psi) = \sum_{\substack{1 \leq j \leq m \\ j \neq i}} \sum_{0 \leq a, b \leq 3} \alpha_{II,i}^{j,ab} (\partial_a \phi_j)(\partial_b \psi_j),$$

which we call *nonresonant terms of types (I) and (II)*, respectively. Similarly we define *nonresonant terms of types (III) and (IV)* by

$$(1.9) \quad R_i^{III}(\phi, \partial\psi) = \sum_{\substack{1 \leq j, k \leq m \\ j \neq k}} \sum_{0 \leq a \leq 3} \alpha_{III,i}^{jk,a} \phi_j (\partial_a \psi_k),$$

$$(1.10) \quad R_i^{IV}(\phi, \partial\psi) = \sum_{\substack{1 \leq j \leq m \\ j \neq i}} \sum_{0 \leq a \leq 3} \alpha_{IV,i}^{j,a} \phi_j (\partial_a \psi_j),$$

respectively. Finally, let H_i be a smooth function of $(u, \partial u)$, satisfying

$$(1.11) \quad H_i(u, \partial u) = O(|u|^3 + |\partial u|^3) \quad \text{near } (u, \partial u) = (0, 0).$$

Now the known results for (SDGE) can be summarized as follows. If F_i has the form

$$(1.12) \quad F_i(u, \partial u) = N_i(\partial u, \partial u) + R_i^I(\partial u, \partial u) + R_i^{II}(\partial u, \partial u) + H_i(u, \partial u) \\ \text{for all } i \in \{1, \dots, m\},$$

then (SDGE) holds for (1.1)–(1.2) (see the author [6]; see also Klainerman [11], Christodoulou [2], Kovalyov [13], Yokoyama [21], Sideris-Tu [19], Kubota-Yokoyama [15], and Sogge [20]). Note that in (1.12), quadratic terms of F_i depend only on ∂u . On the other hand, even if u is involved in quadratic terms, we also have (SDGE) for (1.1)–(1.2), if F_i can be written as

$$(1.13) \quad F_i(u, \partial u) = N_i(\partial u, \partial u) + R_i^I(\partial u, \partial u) + R_i^{III}(u, \partial u) + R_i^{IV}(u, \partial u) + H_i(u, \partial u) \\ \text{for all } i \in \{1, \dots, m\}$$

(see Katayama-Yokoyama [9]; see also the author [5, 7]).

From (1.12) and (1.13), it seems reasonable to conjecture that if F_i can be written as

$$(1.14) \quad F_i(u, \partial u) = N_i(\partial u, \partial u) + R_i^I(\partial u, \partial u) + R_i^{II}(\partial u, \partial u) \\ + R_i^{III}(u, \partial u) + R_i^{IV}(u, \partial u) + H_i(u, \partial u)$$

for all $i \in \{1, \dots, m\}$, then (SDGE) holds. But this conjecture turns out to be false because of the following counterexample by Ohta [17]:

$$(1.15) \quad \begin{cases} \square_{c_1} u_1 = F_1(u, \partial u) := u_2(\partial_t u_1), \\ \square_{c_2} u_2 = F_2(u, \partial u) := (\partial_t u_1)^2. \end{cases}$$

Note that F_1 and F_2 in (1.15) are the nonresonant terms of types (III) and (II), respectively. Hence (1.14) holds for $F = (F_1, F_2)$ in (1.15), but neither (1.12) nor (1.13) is satisfied. In [17], it was proved that (SDGE) does not hold for the above system (1.15) in general. More precisely, for the system (1.15) with $c_1 < c_2$, it was shown that there exist radially symmetric data $f, g \in C_0^\infty(\mathbb{R}^3; \mathbb{R}^2)$ and two positive constants C_3 and ε_3 such that

$$(1.16) \quad T_\varepsilon \leq \exp(C_3 \varepsilon^{-2})$$

for all $\varepsilon \in (0, \varepsilon_3]$.

Since the upper bound of the lifespan obtained in (1.16) is somewhat longer than (1.4), it is interesting to investigate sharpness of (1.16). Our first aim in this paper is to get the lower bound of the lifespan for (1.15). Unfortunately, because it is difficult to obtain energy estimates for (1.15) in large time, we restrict our consideration to radial solutions. Note that the upper bound (1.16) was also obtained for radial solutions.

Before stating our results, we introduce some notation. We say that ϕ is a radially symmetric C_0^∞ -function if ϕ belongs to $C_0^\infty(\mathbb{R}^3)$ and there exists a function $\tilde{\phi} \in C^\infty([0, \infty))$ such that $\phi(x) = \tilde{\phi}(|x|)$ for any $x \in \mathbb{R}^3$. We say $F = F(u, \partial u)$ is rotationally invariant if

$$F(u_O(t, x), \partial u_O(t, x)) = F(u(t, O(x)), (\partial u)(t, O(x)))$$

holds for any C^1 -function $u = u(t, x)$ and any orthogonal transformation $O = O(x)$ on \mathbb{R}^3 , where u_O is defined by $u_O(t, x) = u(t, O(x))$. It is easy to see that if $F = F(u, \partial u)$ is rotationally invariant, and the initial data f and g are radially symmetric C_0^∞ -functions, then the solution u to (1.1)–(1.2) is radial, namely $u(t, x) = \tilde{u}(t, |x|)$ with some function $\tilde{u} = \tilde{u}(t, r)$.

For $\phi = (\phi_i)_{1 \leq i \leq m}$ and $\psi = (\psi_i)_{1 \leq i \leq m}$, we define

$$\begin{aligned} r_i^I(\partial\phi, \partial\psi) &= \sum_{\{(j,k):j \neq k\}} (\beta_{I,i}^{jk,0}(\partial_t \phi_j)(\partial_t \psi_k) + \beta_{I,i}^{jk,1}(\nabla_x \phi_j) \cdot (\nabla_x \psi_k)), \\ r_i^{II}(\partial\phi, \partial\psi) &= \sum_{\{j:j \neq i\}} (\beta_{II,i}^{j,0}(\partial_t \phi_j)(\partial_t \psi_j) + \beta_{II,i}^{j,1}(\nabla_x \phi_j) \cdot (\nabla_x \psi_j)), \\ r_i^{III}(\phi, \partial_t \psi) &= \sum_{\{(j,k):j \neq k\}} \beta_{III,i}^{jk} \phi_j(\partial_t \psi_k), \\ r_i^{IV}(\phi, \partial_t \psi) &= \sum_{\{j:j \neq i\}} \beta_{IV,i}^j \phi_j(\partial_t \psi_j) \end{aligned}$$

for $i = 1, \dots, m$, where $\beta_{I,i}^{jk,a}$, $\beta_{II,i}^{j,a}$ ($a = 0, 1$), $\beta_{III,i}^{jk}$ and $\beta_{IV,i}^j$ are arbitrary constants. Here $\nabla_x \phi = (\partial_1 \phi, \partial_2 \phi, \partial_3 \phi)$ for a C^1 -function ϕ , and \cdot denotes the inner product of \mathbb{R}^3 . Note that $r_i^I(\partial u, \partial u)$, $r_i^{II}(\partial u, \partial u)$, $r_i^{III}(u, \partial u)$, and $r_i^{IV}(u, \partial u)$ are rotationally invariant nonresonant terms of types (I), (II), (III), and (IV), respectively. It is easy to see that the null terms $N_i(\partial u, \partial u)$ are also rotationally invariant.

Theorem 1.1. *Let the propagation speeds c_1, \dots, c_m be distinct. Assume that for each $i \in \{1, \dots, m\}$, F_i has the form*

$$(1.17) \quad \begin{aligned} F_i(u, \partial u) &= N_i(\partial u, \partial u) + r_i^I(\partial u, \partial u) + r_i^{II}(\partial u, \partial u) \\ &\quad + r_i^{III}(u, \partial_i u) + H_i(u, \partial u), \end{aligned}$$

where H_i is rotationally invariant, and satisfies (1.11).

Then, for any radially symmetric C_0^∞ -functions f and g , there exist two positive constants ε_0 and C such that the lifespan T_ε for (1.1)–(1.2) satisfies

$$(1.18) \quad T_\varepsilon(f, g, F) \geq \exp(C\varepsilon^{-2})$$

for any $\varepsilon \in (0, \varepsilon_0]$.

Note that (1.17) contains the null terms, nonresonant terms of types (I), (II) and (III), and terms of higher order. Since F in (1.15) has the form (1.17), the upper bound (1.16) and the lower bound (1.18) guarantee the sharpness of one another, as far as radially symmetric solutions are considered.

To get (1.18), we follow a similar strategy to that in Katayama-Matsumura [8], where the sharp lower bound of the lifespan for the system

$$\begin{cases} \square_{c_1} u_1 = u_1 u_2 & \text{in } (0, \infty) \times \mathbb{R}^3, \\ \square_{c_2} u_2 = u_1^3 & \text{in } (0, \infty) \times \mathbb{R}^3 \end{cases}$$

with $c_1 \neq c_2$ was obtained. The proof of Theorem 1.1 will be given in Section 3.

Now we turn our attention to another problem. Ohta’s counterexample (1.15) says that (SDGE) does not hold for (1.14), especially for a combination of nonresonant terms of types (II) and (III). Our next question is what happens for other combinations. Here we give an example which suggests (SDGE) may hold in general for a combination of null terms, nonresonant terms of types (I), (II) and (IV).

Theorem 1.2. *Let $m = 2$, and consider the Cauchy problem (1.1)–(1.2) with*

$$(1.19) \quad F_1(u, \partial u) := N_1(\partial u, \partial u) + r_1^I(\partial u, \partial u) + r_1^{II}(\partial u, \partial u) + r_1^{IV}(u, \partial_i u),$$

$$(1.20) \quad F_2(u, \partial u) := N_2(\partial u, \partial u) + r_2^I(\partial u, \partial u) + r_2^{II}(\partial u, \partial u).$$

Assume $c_1 \neq c_2$. Then, for any radially symmetric C_0^∞ -functions $f = (f_1, f_2)$ and $g = (g_1, g_2)$, there exists a positive constant ε_0 such that

$$(1.21) \quad T_\varepsilon(f, g, F) = \infty \quad \text{for any } \varepsilon \in (0, \varepsilon_0].$$

Nonresonant terms of types (II) and (IV) are involved in (1.19)–(1.20). Hence neither (1.12) nor (1.13) holds for this $F = (F_1, F_2)$. Nonetheless (SDGE) holds for (1.1) with this F as far as we consider radial solutions. Theorem 1.2 suggests that there may be a certain sufficient condition for (SDGE) other than (1.12) and (1.13). Of course, even for (1.19)–(1.20), it may possibly happen that (SDGE) does not hold for general C_0^∞ data. This problem is still open. The proof of Theorem 1.2 will be given in Section 4.

Throughout this paper, various positive constants, which may change line by line, are denoted just by the same letter C .

2. Basic decay estimates

In this section, we derive basic L^∞ - L^∞ decay estimates.

For $\phi, \psi \in C_0^\infty(\mathbb{R}^3)$ and a positive constant c , we write $U_c^*[\phi, \psi]$ for the solution to the Cauchy problem for

$$\begin{cases} \square_c U_c^*[\phi, \psi](t, x) = 0 & \text{for } (t, x) \in (0, \infty) \times \mathbb{R}^3, \\ U_c^*[\phi, \psi](0, x) = \phi(x), (\partial_t U_c^*[\phi, \psi])(0, x) = \psi(x) & \text{for } x \in \mathbb{R}^3. \end{cases}$$

Similarly, for a continuous function $G = G(t, x)$ on $(0, \infty) \times \mathbb{R}^3$, we write $U_c[G]$ for the solution to the Cauchy problem for

$$\begin{cases} \square_c U_c[G](t, x) = G(t, x) & \text{for } (t, x) \in (0, \infty) \times \mathbb{R}^3, \\ U_c[G](0, x) = (\partial_t U_c[G])(0, x) = 0 & \text{for } x \in \mathbb{R}^3. \end{cases}$$

For $\rho \in \mathbb{R}$, we write $\langle \rho \rangle = \sqrt{1 + \rho^2}$. For a continuous function ϕ , a non-negative constant ν , and $t, r \in [0, \infty)$, $\|\phi\|_{\nu, t, r}$ is defined by

$$\|\phi\|_{\nu, t, r} = \sup_{y \in \mathbb{R}^3 \text{ with } |t-r| \leq |y| \leq t+r} \langle |y| \rangle^\nu |\phi(y)|.$$

For $U_c^*[\phi, \psi]$, we have the following.

Lemma 2.1. *Let $c > 0$ and $\kappa > 0$. Then there exists a positive constant C such that we have*

$$(2.1) \quad \begin{aligned} & \langle t + |x| \rangle \langle ct - |x| \rangle^\kappa |U_c^*[\phi, \psi](t, x)| \\ & \leq C(\|\phi\|_{2+\kappa, ct, |x|} + \|\nabla_x \phi\|_{2+\kappa, ct, |x|} + \|\psi\|_{2+\kappa, ct, |x|}) \end{aligned}$$

for $(t, x) \in [0, \infty) \times \mathbb{R}^3$.

For the proof, see Kubota-Yokoyama [15]. More precisely, the estimate stated in [15] is not exactly (2.1), but we can easily obtain (2.1) by investigating their proof (see also Asakura [1]).

Let c_1, \dots, c_m be given positive constants, and let $c_0 = 0$. We define

$$w(t, r) = \min_{0 \leq j \leq m} \langle c_j t - r \rangle$$

for $(t, r) \in [0, \infty) \times [0, \infty)$. We also define

$$(2.2) \quad \Phi_\kappa(t, r) = \begin{cases} \langle t+r \rangle^{-\kappa} & \text{if } \kappa < 0, \\ \log\left(2 + \frac{\langle t+r \rangle}{\langle t-r \rangle}\right) & \text{if } \kappa = 0, \\ \langle t-r \rangle^{-\kappa} & \text{if } \kappa > 0, \end{cases}$$

$$(2.3) \quad \Psi_\mu(t) = \begin{cases} \log(2+t) & \text{if } \mu = 0, \\ 1 & \text{if } \mu > 0. \end{cases}$$

Note that we have $\Phi_0(t, r) \leq C \log(2+t)$ for any $(t, r) \in [0, \infty) \times [0, \infty)$. For $1 \leq i \leq m$, $(t, x) \in [0, \infty) \times \mathbb{R}^3$ and $(t, r) \in [0, \infty) \times [0, \infty)$, we put

$$\begin{aligned} \Theta_i(t, x) &= \{(\tau, y) \in [0, t] \times \mathbb{R}^3; |x| - c_i(t - \tau) \leq |y| \leq |x| + c_i(t - \tau)\}, \\ \Theta_i^*(t, r) &= \{(\tau, \lambda) \in [0, t] \times [0, \infty); |r - c_i(t - \tau)| \leq \lambda \leq r + c_i(t - \tau)\}. \end{aligned}$$

Then, for $U_{c_i}[G]$ we have

Lemma 2.2. *Let $i \in \{1, \dots, m\}$. For $\rho > 0$ and $\mu \geq 0$, there exists a positive constant C such that*

$$(2.4) \quad \begin{aligned} & (t + |x|) \Phi_{\rho-1}(c_i t, |x|)^{-1} |U_{c_i}[G](t, x)| \\ & \leq C \Psi_\mu(t) \sup_{(\tau, y) \in \Theta_i(t, x)} \langle \tau + |y| \rangle^\rho w(\tau, |y|)^{1+\mu} |y| |G(\tau, y)| \end{aligned}$$

for $(t, x) \in [0, \infty) \times \mathbb{R}^3$.

Proof. The case where $\rho > 1$ and $\mu > 0$ was proved by Katayama-Yokoyama [9, Section 8]. Other cases can be proved by apparent modifications of the proof in [9], and we only give a sketch of the proof here (see also the author [6] for the case $\rho = 1$ and $\mu = 0$, and Katayama-Matsumura [8] for the case $\rho > 1$ and $\mu = 0$).

Without loss of generality, we may assume $c_i = 1$. Then, following [9, Section 8], we find

$$(2.5) \quad \begin{aligned} |U_{c_i}[G](t, x)| & \leq C r^{-1} \int_0^t \int_{|r-t+\tau|}^{t-\tau+r} \mathcal{W}(\tau, \lambda)^{-1} d\lambda d\tau \\ & \quad \times \sup_{(\tau, y) \in \Theta_i(t, x)} \mathcal{W}(\tau, |y|) |y| |G(\tau, y)| \end{aligned}$$

for any positive function $\mathcal{W} = \mathcal{W}(\tau, \lambda)$, where $r = |x|$. Therefore our task is to show

$$(2.6) \quad r^{-1} \int_0^t \int_{|r-t+\tau|}^{t-\tau+r} \mathcal{W}(\tau, \lambda)^{-1} d\lambda d\tau \leq C \langle t+r \rangle^{-1} \Phi_{\rho-1}(t, r) \Psi_\mu(t)$$

with $\mathcal{W}(t, r) = \langle t+r \rangle^\rho w(t, r)^{1+\mu}$. Since $w(t, r)^{-1} \leq \sum_{j=0}^m \langle c_j t - r \rangle^{-1}$, it suffices to evaluate the integral

$$(2.7) \quad J_{\rho, \mu, a}(t, r) := r^{-1} \int_0^t \int_{|r-t+\tau|}^{t-\tau+r} (1+\tau+\lambda)^{-\rho} (1+|a\tau-\lambda|)^{-1-\mu} d\lambda d\tau$$

with $a \geq 0$. Performing a change of variables $\alpha = \tau + \lambda$ and $\beta = \lambda - a\tau$, we obtain

$$(2.8) \quad J_{\rho, \mu, a}(t, r) = \frac{1}{(a+1)r} \int_{|t-r|}^{t+r} (1+\alpha)^{-\rho} d\alpha \int_{\hat{\beta}}^\alpha (1+|\beta|)^{-1-\mu} d\beta$$

with $\hat{\beta} = \{(1-a)\alpha + (1+a)(r-t)\}/2$ (see [9, (8.6) and (8.7)]).

For example, if $\mu > 0$, it is easy to see

$$(2.9) \quad J_{1, \mu, a}(t, r) \leq Cr^{-1} \int_{|t-r|}^{t+r} (1+\alpha)^{-1} d\alpha.$$

By a direct calculation, we get

$$(2.10) \quad r^{-1} \int_{|t-r|}^{t+r} (1+\alpha)^{-1} d\alpha = Cr^{-1} \log \left(\frac{1+t+r}{1+|t-r|} \right) \leq C \langle t+r \rangle^{-1} \Phi_0(t, r)$$

for $r \geq (t+1)/2$. On the other hand, we also have

$$(2.11) \quad r^{-1} \int_{|t-r|}^{t+r} (1+\alpha)^{-1} d\alpha \leq C(1+|t-r|)^{-1} \leq C(1+t+r)^{-1}$$

for $r \leq (t+1)/2$. (2.9), (2.10) and (2.11) imply (2.6) for the case $\rho = 1$ and $\mu > 0$ immediately. Other cases can be treated similarly. □

By (2.5) and similar lines to (2.6)–(2.11), we also have

Corollary 2.3. *Let $i \in \{1, \dots, m\}$ and*

$$\mathcal{W}(t, r)^{-1} = A_1 \langle t+r \rangle^{-1} w(t, r)^{-1} + A_2 \langle t+r \rangle^{-1} w(t, r)^{-2}$$

with some positive constants A_1 and A_2 . Then we have

$$\begin{aligned} & \langle t+|x| \rangle \Phi_0(c_i t, |x|)^{-1} |U_{c_i}[G](t, x)| \\ & \leq C(A_1 \log(2+t) + A_2) \sup_{(\tau, y) \in \Theta_i(t, x)} \mathcal{W}(\tau, |y|) |y| |G(\tau, y)|. \end{aligned}$$

For $c > 0$, $(t, r) \in [0, \infty) \times [0, \infty)$, and a C^1 -function $G = G(t, r)$ on $[0, \infty) \times [0, \infty)$, we define

$$(2.12) \quad L_c[G](t, r) = \frac{1}{2c} \int_0^t \left(\int_{\lambda_c^-(\tau; t, r)}^{\lambda_c^+(\tau; t, r)} \check{G}(\tau, \lambda) d\lambda \right) d\tau,$$

where

$$(2.13) \quad \lambda_c^\pm(\tau; t, r) = r \pm c(t - \tau),$$

and \check{G} is defined by $\check{G}(t, r) = rG(t, |r|)$ for $(t, r) \in [0, \infty) \times \mathbb{R}$. Then easy calculations lead to

$$(2.14) \quad (\partial_t \pm c\partial_r)L_c[G](t, r) = I_c^\pm[\check{G}](t, r),$$

$$(2.15) \quad \partial_t(\partial_t \pm c\partial_r)L_c[G](t, r) = \check{G}(t, r) \pm cI_c^\pm[\partial_r\check{G}](t, r),$$

$$(2.16) \quad \partial_r(\partial_t \pm c\partial_r)L_c[G](t, r) = I_c^\pm[\partial_r\check{G}](t, r),$$

where $I_c^\pm[H](t, r)$ is defined by

$$I_c^\pm[H](t, r) = \int_0^t H(\tau, \lambda_c^\pm(\tau; t, r)) d\tau, \quad (t, r) \in [0, \infty) \times \mathbb{R}$$

for a function $H = H(t, r)$. Note that we have $(\partial_r\check{G})(t, r) = G(t, |r|) + |r|(\partial_r G)(t, |r|)$ for $r \in \mathbb{R}$. It is also easy to verify that a classical solution v to

$$(2.17) \quad \begin{cases} \square_c v(t, x) = G(t, |x|) & \text{in } (0, \infty) \times \mathbb{R}^3, \\ v(0, x) = \partial_t v(0, x) = 0 & \text{for } x \in \mathbb{R}^3 \end{cases}$$

can be written as

$$(2.18) \quad v(t, x) = |x|^{-1}L_c[G](t, |x|) \quad \text{for } (t, x) \in [0, \infty) \times (\mathbb{R}^3 \setminus \{0\}).$$

Before we proceed to estimate derivatives of solutions to wave equations, we give two technical lemmas.

Lemma 2.4. *Let $c > 0$, $\alpha \neq 0$ and $p \geq 0$. Then we have*

$$(2.19) \quad \int_0^t (1 + |\alpha\tau - |r \pm ct||)^{-(1+p)} d\tau \leq C\Psi_p(t)$$

for $(t, r) \in [0, \infty) \times [0, \infty)$, where Ψ_μ is defined by (2.3).

Proof. It is very easy to treat the case $p > 0$, and we only consider the case $p = 0$ here. Suppose $r \geq (|\alpha| + c)t$. Then we get

$$(2.20) \quad \int_0^t (1 + |\alpha\tau - |r \pm ct||)^{-1} d\tau = \int_0^t (1 + r \pm ct - \alpha\tau)^{-1} d\tau \\ = \frac{1}{\alpha} \log\left(\frac{1 + r \pm ct}{1 + r \pm ct - \alpha t}\right).$$

(2.20) implies (2.19), because we have

$$(1 + |\alpha|t)^{-1} \leq \frac{1 + r \pm ct}{1 + r \pm ct - \alpha t} \leq 1 + |\alpha|t$$

for $r \geq (|\alpha| + c)t$.

On the other hand, if $r < (|\alpha| + c)t$, it is easy to see

$$\int_0^t (1 + |\alpha\tau - |r \pm ct||)^{-1} d\tau \leq C \log(2 + t + r) \leq C \log(2 + t).$$

This completes the proof. □

For $c > 0$, $a \geq 0$, $\rho \geq 1$ and $\mu \geq 0$, we define

$$K_{c,a,\rho,\mu}^{\pm}(t, r) = \int_0^t \langle \tau + |\lambda_c^{\pm}(\tau; t, r)| \rangle^{-\rho} \langle a\tau - |\lambda_c^{\pm}(\tau; t, r)| \rangle^{-(1+\mu)} d\tau.$$

Lemma 2.5. *Let $c > 0$.*

(i) *For $\mu \geq 0$ and $\rho \geq 1$, we have*

$$(2.21) \quad K_{c,c,\rho,\mu}^+(t, r) \leq C \Psi_{\mu}(t) \langle ct + r \rangle^{-\rho},$$

$$(2.22) \quad K_{c,c,\rho,\mu}^-(t, r) \leq C \Psi_{\mu}(t) \langle ct - r \rangle^{-\rho} + C \Phi_{\rho-1}(ct, r) \langle ct - r \rangle^{-(1+\mu)}$$

for $(t, r) \in [0, \infty) \times [0, \infty)$, where $\Phi_{\rho-1}$ and Ψ_{μ} are from (2.2) and (2.3), respectively.

(ii) *Let $a \geq 0$, and suppose $a \neq c$. Then, for $\mu \geq 0$ and $\rho > 0$, we have*

$$(2.23) \quad K_{c,a,\rho,\mu}^{\pm}(t, r) \leq C \langle ct \pm r \rangle^{-\rho} \Psi_{\mu}(t)$$

for $(t, r) \in [0, \infty) \times [0, \infty)$, where Ψ_{μ} is given by (2.3).

Proof. First we note that $K_{c,a,\rho,\mu}^{\pm}(t, r)$ is bounded by

$$C \int_0^t (1 + c\tau + |\lambda_c^{\pm}(\tau; t, r)|)^{-\rho} (1 + |a\tau - |\lambda_c^{\pm}(\tau; t, r)||)^{-(1+\mu)} d\tau.$$

Since $c\tau + \lambda_c^+(\tau; t, r) = ct + r$, and $c\tau - \lambda_c^+(\tau; t, r) = 2c\tau - (r + ct)$, Lemma 2.4 implies

$$\begin{aligned} K_{c,c,\rho,\mu}^+(t, r) &\leq C\langle ct + r \rangle^{-\rho} \int_0^t (1 + |2c\tau - (r + ct)|)^{-(1+\mu)} d\tau \\ &\leq C\langle ct + r \rangle^{-\rho} \Psi_\mu(t). \end{aligned}$$

Suppose $r < ct$. Observing that we have

$$|\lambda_c^-(\tau; t, r)| = \begin{cases} c\tau - (ct - r) & \text{if } \tau \geq \frac{ct - r}{c}, \\ -c\tau + ct - r & \text{if } \tau < \frac{ct - r}{c}, \end{cases}$$

we get

$$\begin{aligned} K_{c,c,\rho,\mu}^-(t, r) &\leq C\langle ct - r \rangle^{-\rho} \int_0^{t \wedge (ct-r)/c} (1 + |2c\tau - (ct - r)|)^{-(1+\mu)} d\tau \\ &\quad + C\langle ct - r \rangle^{-(1+\mu)} \int_{t \wedge (ct-r)/c}^t (1 + 2c\tau - (ct - r))^{-\rho} d\tau, \end{aligned}$$

where $\alpha \wedge \beta = \min\{\alpha, \beta\}$. By Lemma 2.4, we see that the first term on the right-hand side of the above is bounded by $C\langle ct - r \rangle^{-\rho} \Psi_\mu(t)$. We also see that the second term is bounded by $C\langle ct - r \rangle^{-(1+\mu)} \Phi_{\rho-1}(ct, r)$, since we have

$$\int_{(ct-r)/c}^t (1 + 2c\tau - (ct - r))^{-1} d\tau = \frac{1}{2c} \log \frac{1 + ct + r}{1 + ct - r}$$

and

$$\int_{(ct-r)/c}^t (1 + 2c\tau - (ct - r))^{-\rho} d\tau \leq \frac{1}{2c(\rho - 1)} (1 + ct - r)^{-\rho+1}$$

for $\rho > 1$, provided that $(ct - r)/c < t$.

Now suppose $r > ct$. Then we have

$$K_{c,c,\rho,\mu}^-(t, r) \leq C\langle ct - r \rangle^{-(1+\mu)} \int_0^t (1 + 2c\tau + (r - ct))^{-\rho} d\tau,$$

and a similar argument to the above leads to

$$K_{c,c,\rho,\mu}^-(t, r) \leq C\langle ct - r \rangle^{-(1+\mu)} \Phi_{\rho-1}(ct, r).$$

Finally we are going to prove (2.23). Let $a \geq 0$ and $a \neq c$. Since we have $c\tau + |\lambda_c^\pm(\tau; t, r)| \geq C|c\tau \pm r|$ for $\tau \geq 0$, $a\tau - \lambda_c^+(\tau; t, r) = (a+c)\tau - (r+ct)$, and

$$a\tau - |\lambda_c^-(\tau; t, r)| = \begin{cases} (a-c)\tau - (r-ct) & \text{if } \tau \geq \frac{ct-r}{c}, \\ (a+c)\tau - (ct-r) & \text{if } \tau < \frac{ct-r}{c}, \end{cases}$$

Lemma 2.4 implies

$$\begin{aligned} K_{c,a,\rho,\mu}^\pm(t, r) &\leq C\langle ct \pm r \rangle^{-\rho} \int_0^t (1 + |a\tau - |\lambda_c^\pm(\tau; t, r)||)^{-(1+\mu)} d\tau \\ &\leq C\langle ct \pm r \rangle^{-\rho} \Psi_\mu(t). \end{aligned}$$

This completes the proof. □

Let c_1, \dots, c_m be given positive constants, and $c_0 = 0$ as before. For $i \in \{1, 2, \dots, m\}$, we define

$$w_i(t, r) = \min_{\substack{0 \leq j \leq m \\ j \neq i}} \langle c_j t - r \rangle.$$

Lemma 2.6. *Let $i \in \{1, \dots, m\}$. Suppose $G \in C^1([0, \infty) \times [0, \infty))$, and let v be a classical solution to*

$$\square_{c_i} v(t, x) = G(t, |x|) \quad \text{for } (t, x) \in (0, \infty) \times \mathbb{R}^3$$

with $v = \partial_t v = 0$ at $t = 0$. Set

$$\mathcal{D}[v](t, x) = \langle r \rangle \sum_{|\alpha|=1} |\partial_{t,r}^\alpha v(t, x)| + r \sum_{|\alpha|=2} |\partial_{t,r}^\alpha v(t, x)|,$$

where $r = |x|$, $\partial_r = \sum_{j=1}^3 (x_j/|x|)\partial_j$, and $\partial_{t,r}^\alpha$ denotes $\partial_t^{\alpha_1} \partial_r^{\alpha_2}$ for a multi-index $\alpha = (\alpha_1, \alpha_2)$. We define $D_{+,c} = \partial_t + c\partial_r$ for $c > 0$, and

$$\mathcal{D}_i^+[v](t, x) = \langle r \rangle |D_{+,c_i} v(t, x)| + r \sum_{|\alpha|=1} |\partial_{t,r}^\alpha D_{+,c_i} v(t, x)|.$$

We also set

$$\mathcal{M}[G](t, r) = \langle r \rangle |G(t, r)| + r \sum_{|\alpha|=1} |\partial_{t,r}^\alpha G(t, r)|.$$

Then we have the following estimates:

(i) For $\rho \geq 1$ and $\mu \geq 0$, we have

$$\begin{aligned} & (\Psi_\mu(t)\langle c_i t - r \rangle^{-\rho} + \Phi_{\rho-1}(c_i t, r)\langle c_i t - r \rangle^{-(1+\mu)})^{-1} \mathcal{D}[v](t, x) \\ & \leq C \sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \langle \tau + \lambda \rangle^\rho w(\tau, \lambda)^{1+\mu} \mathcal{M}[G](\tau, \lambda). \end{aligned}$$

(ii) For $0 < \rho \leq 1$ and $\mu > 0$, we have

$$\langle c_i t - r \rangle^\rho \mathcal{D}[v](t, x) \leq C \sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \langle \tau + \lambda \rangle^\rho w_i(\tau, \lambda)^{1+\mu} \mathcal{M}[G](\tau, \lambda).$$

(iii) For $\rho \geq 1$ and $\mu \geq 0$, we have

$$\begin{aligned} & (t+r)\Phi_{\rho-1}(c_i t, r)^{-1} \mathcal{D}_i^+[v](t, x) \\ & \leq C \Psi_\mu(t) \sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \langle \tau + \lambda \rangle^\rho w(\tau, \lambda)^{1+\mu} \mathcal{M}[G](\tau, \lambda). \end{aligned}$$

Proof. For $\rho > 0$, Lemma 2.1 implies

$$\begin{aligned} & (t+r)\langle c_i t - r \rangle^\rho |U_{c_i}^*[0, G(0, |\cdot|)](t, x)| \\ (2.24) \quad & \leq C \|G(0, |\cdot|)\|_{2+\rho, c_i t, r} \\ & \leq C \sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \langle \tau + \lambda \rangle^\rho w(\tau, \lambda) \mathcal{M}[G](\tau, \lambda), \end{aligned}$$

since we have $\langle \lambda \rangle^{2+\rho} = \langle 0+\lambda \rangle^\rho w(0, \lambda) \langle \lambda \rangle$. For $\rho > 0$, $\mu \geq 0$, and $0 \leq a \leq 3$, Lemma 2.2 leads to

$$\begin{aligned} & (t+r)\Phi_{\rho-1}(c_i t, r)^{-1} |U_{c_i}[\partial_a \square_{c_i} v](t, x)| \\ (2.25) \quad & \leq C \Psi_\mu(t) \sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \langle \tau + \lambda \rangle^\rho w(\tau, \lambda)^{1+\mu} \lambda |(\partial_a G)(\tau, \lambda)|. \end{aligned}$$

Since we have $\partial_a v = U_{c_i}[\partial_a \square_{c_i} v] + \delta_{a0} U_{c_i}^*[0, G(0, |\cdot|)]$ with the Kronecker delta δ_{ab} , and $(t+r)^{-1} \Phi_{\rho-1}(c_i t, r) \leq C \langle c_i t - r \rangle^{-\rho}$ for $\rho > 0$, from Lemma 2.2, (2.24) and (2.25) we get

$$\begin{aligned} & \langle c_i t - r \rangle^\rho (|v(t, x)| + |\partial_t v(t, x)| + |\partial_r v(t, x)|) \\ (2.26) \quad & \leq C \Psi_\mu(t) \sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \langle \tau + \lambda \rangle^\rho w(\tau, \lambda)^{1+\mu} \mathcal{M}[G](\tau, \lambda) \end{aligned}$$

for $\rho > 0$ and $\mu \geq 0$.

For $\rho > 0$, it is easy to see

$$(2.27) \quad \langle t+r \rangle^\rho r |G(t, r)| \leq C \sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \langle \tau + \lambda \rangle^\rho \mathcal{M}[G](\tau, \lambda).$$

Now we set

$$\begin{aligned} \tilde{\mathcal{D}}[v](t, x) &= |\partial_t(rv(t, x))| + |\partial_r(rv(t, x))| + |\partial_r \partial_t(rv(t, x))| \\ &\quad + |\partial_t^2(rv(t, x)) - rG(t, r)| + |\partial_r^2(rv(t, x))|. \end{aligned}$$

Since we have

$$\mathcal{D}[v](t, x) \leq C \left(\tilde{\mathcal{D}}[v](t, x) + \sum_{|\alpha| \leq 1} |\partial_{t,r}^\alpha v(t, x)| + r|G(t, r)| \right),$$

in view of (2.26) and (2.27) we only have to prove (i) and (ii) with $\mathcal{D}[v]$ replaced by $\tilde{\mathcal{D}}[v]$. As we have mentioned before, we have $rv(t, x) = L_{c_i}[G](t, r)$. Therefore, from (2.14), (2.15) and (2.16) we get

$$(2.28) \quad \tilde{\mathcal{D}}[v](t, x) \leq C \sum_{s=+,-} (|I_{c_i}^s[\check{G}](t, r)| + |I_{c_i}^s[\partial_r \check{G}](t, r)|),$$

and we find

$$(2.29) \quad \begin{aligned} \tilde{\mathcal{D}}[v](t, x) &\leq C \sum_{s=+,-} \int_0^t \mathcal{W}(\tau, |\lambda_{c_i}^s(\tau; t, r)|)^{-1} d\tau \\ &\quad \times \sup_{(\tau, \lambda) \in \Theta_i^s(t, r)} \mathcal{W}(\tau, \lambda) \mathcal{M}[G](\tau, \lambda) \end{aligned}$$

for any positive function $\mathcal{W} = \mathcal{W}(\tau, \lambda)$.

We use (2.29) with

$$\mathcal{W}(\tau, \lambda) = \langle \tau + \lambda \rangle^\rho w(\tau, \lambda)^{1+\mu} \quad \text{and} \quad \mathcal{W}(\tau, \lambda) = \langle \tau + \lambda \rangle^\rho w_i(\tau, \lambda)^{1+\mu}$$

to obtain (i) and (ii), respectively. Noting that we have

$$\begin{aligned} w(\tau, \lambda)^{-(1+\mu)} &\leq \sum_{0 \leq j \leq m} \langle c_j \tau - \lambda \rangle^{-(1+\mu)}, \\ w_i(\tau, \lambda)^{-(1+\mu)} &\leq \sum_{\substack{0 \leq j \leq m \\ j \neq i}} \langle c_j \tau - \lambda \rangle^{-(1+\mu)} \end{aligned}$$

for $\mu \geq 0$, and using Lemma 2.5 to estimate $\int_0^t \mathcal{W}(\tau, |\lambda_{c_i}^\pm(\tau; t, r)|)^{-1} d\tau$, we obtain (i) and (ii) with $\mathcal{D}[v]$ replaced by $\tilde{\mathcal{D}}[v]$.

Concerning (iii), our task is to estimate $\sum_{|\alpha| \leq 1} |r \partial_{t,r}^\alpha D_{+,c_i} v(t, x)|$, because (2.24) and (2.25) imply the desired bound for $|D_{+,c_i} v(t, x)|$.

We have

$$\begin{aligned} r \partial_t^j D_{+,c_i} v(t, x) &= \partial_t^j D_{+,c_i}(rv(t, x)) - c_i \partial_t^j v(t, x) \quad \text{for } j = 0, 1, \\ r \partial_r D_{+,c_i} v(t, x) &= \partial_r D_{+,c_i}(rv(t, x)) - D_{+,c_i} v(t, x) - c_i \partial_r v(t, x), \end{aligned}$$

and by (2.14)–(2.16) we also have

$$\sum_{|\alpha| \leq 1} |\partial_{t,r}^\alpha D_{+,c_i}(rv(t, x))| \leq C(I_{c_i}^+[\check{G}] + I_{c_i}^+[\partial_r \check{G}]) + C|\check{G}(t, r)|.$$

Hence we obtain the desired estimate for $\sum_{|\alpha| \leq 1} r|\partial_{t,r}^\alpha D_{+,c_i}v(t, x)|$ from Lemmas 2.2 and 2.5 together with (2.24), (2.25) and (2.27) (note that we have $\langle t+r \rangle \Phi_{\rho-1}^{-1}(t, r) \leq \langle t+r \rangle^\rho$ for $\rho \geq 1$). This completes the proof. \square

From the proof of Lemma 2.6, with using Corollary 2.3 in place of Lemma 2.2 and choosing \mathcal{W} as in Corollary 2.3, we also have

Corollary 2.7. *Let v and G be as in Lemma 2.6, and*

$$\mathcal{W}(t, r)^{-1} = A_1 \langle t+r \rangle^{-1} w(t, r)^{-1} + A_2 \langle t+r \rangle^{-1} w(t, r)^{-2}$$

with some positive constants A_1 and A_2 . Then we have

$$\begin{aligned} & (\langle c_i t - r \rangle^{-1} + \Phi_0(c_i t, r) \langle c_i t - r \rangle^{-2})^{-1} \mathcal{D}[v](t, x) \\ & + \langle t+r \rangle \Phi_0(c_i t, r)^{-1} \mathcal{D}_i^+[v](t, x) \\ & \leq C(A_1 \log(2+t) + A_2) \sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \mathcal{W}(\tau, \lambda) \mathcal{M}[G](\tau, \lambda), \end{aligned}$$

where $r = |x|$.

We conclude this section with a decay estimate for $\mathcal{D}_i^+[U_{c_i}^*]$.

Lemma 2.8. *Let $i \in \{1, \dots, m\}$, $\kappa > 0$, and $v = U_{c_i}^*[\phi, \psi]$. Suppose that ϕ and ψ are radially symmetric functions. Then we have*

$$\begin{aligned} & (t + |x|) \langle c_i t - |x| \rangle^\kappa \mathcal{D}_i^+[v](t, x) \\ & \leq C \left(\sum_{0 \leq k \leq 2} \|\partial_r^k \phi\|_{2+\kappa, c_i t, |x|} + \sum_{0 \leq k \leq 1} \|\partial_r^k \psi\|_{2+\kappa, c_i t, |x|} \right). \end{aligned}$$

Proof. Since ϕ and ψ are radially symmetric, we see that v also is radially symmetric. Set $w(t, r) = rv(t, (|r|, 0, 0))$, $\check{\phi}(r) = r\phi(|r|, 0, 0)$, and $\check{\psi}(r) = r\psi(|r|, 0, 0)$. Then we get $(\partial_t^2 - c_i^2 \partial_r^2)w(t, r) = 0$ for $(t, r) \in [0, \infty) \times \mathbb{R}$, with $w = \check{\phi}$ and $\partial_t w = \check{\psi}$ at $t = 0$. It is easy to check that we have

$$\mathcal{D}_i^+[v](t, x) \leq C \sum_{|\alpha| \leq 1} |(D_{+,c_i} \partial_{t,r}^\alpha w)(t, |x|)| + \sum_{|\alpha| \leq 1} |\partial_{t,r}^\alpha v(t, x)|.$$

By Lemma 2.1, we obtain

$$\begin{aligned}
 & \langle t + |x| \rangle \langle c_i t - |x| \rangle^k \sum_{|\alpha| \leq 1} |\partial_{t,r}^\alpha v(t, x)| \\
 (2.30) \quad & \leq C \left(\sum_{0 \leq k \leq 2} \|\partial_r^k \phi\|_{2+\kappa, c_i t, r} + \sum_{0 \leq k \leq 1} \|\partial_r^k \psi\|_{2+\kappa, c_i t, r} \right).
 \end{aligned}$$

Since $(\partial_t - c_i \partial_r)(D_{+,c_i} \partial_{t,r}^\alpha w)(t, r) = 0$, we get

$$(2.31) \quad D_{+,c_i} \partial_{t,r}^\alpha w(t, r) = (D_{+,c_i} \partial_{t,r}^\alpha w)(0, c_i t + r).$$

Now it is easy to see

$$\begin{aligned}
 & \langle t + r \rangle^{1+\kappa} \sum_{|\alpha| \leq 1} |(D_{+,c_i} \partial_{t,r}^\alpha w)(0, c_i t + r)| \\
 (2.32) \quad & \leq \sum_{|\alpha| \leq 1} |\langle c_i t + r \rangle^{1+\kappa} (D_{+,c_i} \partial_{t,r}^\alpha w)(0, c_i t + r)| \\
 & \leq C \left(\sum_{0 \leq k \leq 2} \|\partial_r^k \phi\|_{2+\kappa, c_i t, r} + \sum_{0 \leq k \leq 1} \|\partial_r^k \psi\|_{2+\kappa, c_i t, r} \right).
 \end{aligned}$$

(2.30), (2.31), and (2.32) imply the desired result. □

3. Proof of Theorem 1.1

For brevity, when $v = v(t, x)$ is radially symmetric, we write sometimes $v = v(t, x)$ and sometimes $v = v(t, r)$ with $r = |x|$ in the following. In other words, if there exists $\tilde{v} = \tilde{v}(t, r)$ such that $v(t, x) = \tilde{v}(t, |x|)$, we do not distinguish v from \tilde{v} .

Suppose that all the assumptions in Theorem 1.1 are fulfilled. Let $(u^{(k)})_{1 \leq k \leq 3} = ((u_i^{(k)})_{1 \leq i \leq m})_{1 \leq k \leq 3}$ be a solution to

$$\begin{aligned}
 (3.1) \quad \square_{c_i} u_i^{(1)} &= N_i(\partial u^{(1)}, \partial u^{(1)}) + r_i^I(\partial u^{(1)}, \partial u^{(1)}) + r_i^{III}(u^{(1)}, \partial_t u^{(1)}) \\
 &+ H_i(u^{(1)}, \partial u^{(1)}),
 \end{aligned}$$

$$\begin{aligned}
 (3.2) \quad \square_{c_i} u_i^{(2)} &= 2N_i(\partial u^{(1)}, \partial u^{(2)+(3)}) + 2r_i^I(\partial u^{(1)}, \partial u^{(2)+(3)}) \\
 &+ N_i(\partial u^{(2)+(3)}, \partial u^{(2)+(3)}) + r_i^I(\partial u^{(2)+(3)}, \partial u^{(2)+(3)}) \\
 &+ r_i^{II}(\partial u^{(1)+(2)+(3)}, \partial u^{(1)+(2)+(3)}) \\
 &+ r_i^{III}(u^{(1)+(2)+(3)}, \partial_t u^{(2)}) \\
 &+ H_i(u^{(1)+(2)+(3)}, \partial u^{(1)+(2)+(3)}) - H_i(u^{(1)}, \partial u^{(1)}),
 \end{aligned}$$

$$(3.3) \quad \square_{c_i} u_i^{(3)} = r_i^{III}(u^{(2)+(3)}, \partial_t u^{(1)}) + r_i^{III}(u^{(1)+(2)+(3)}, \partial_t u^{(3)})$$

for $1 \leq i \leq m$, with initial data

$$(3.4) \quad u^{(1)} = \varepsilon f, \quad \partial_t u^{(1)} = \varepsilon g, \quad u^{(2)} = \partial_t u^{(2)} = u^{(3)} = \partial_t u^{(3)} = 0$$

at $t = 0$, where $u^{(j)+(k)} = u^{(j)} + u^{(k)}$ for $1 \leq j, k \leq 3$, and $u^{(1)+(2)+(3)} = u^{(1)} + u^{(2)} + u^{(3)}$. Since f and g are radially symmetric, and nonlinearity is rotationally invariant, we see that $u^{(k)}$ ($k = 1, 2, 3$) are radial functions. Note that we have $(\nabla_x v) \cdot (\nabla_x w) = (\partial_r v)(\partial_r w)$ for radial functions v and w . Set $u = u^{(1)} + u^{(2)} + u^{(3)}$, and we find that u satisfies (1.1)–(1.2). Hence our task is to solve the Cauchy problem (3.1)–(3.3) with (3.4).

From the classical local existence theorem, the Cauchy problem (3.1)–(3.4) admits a unique solution $(u^{(k)})_{1 \leq k \leq 3}$ for $0 \leq t < T$ with some $T > 0$. Let T_ε be the supremum of such T .

We define

$$\begin{aligned} e_1[u^{(1)}](t, r) &= \sum_{i=1}^m \{ (t+r) \langle c_i t - r \rangle (|u_i^{(1)}(t, r)| + \mathcal{D}_i^+[u_i^{(1)}](t, r)) \\ &\quad + \langle c_i t - r \rangle^2 \mathcal{D}[u_i^{(1)}](t, r) \}, \\ e_2[u^{(2)}](t, r) &= \sum_{i=1}^m \{ (t+r) \Phi_0(c_i t, r)^{-1} (|u_i^{(2)}(t, r)| + \mathcal{D}_i^+[u_i^{(2)}](t, r)) \\ &\quad + \langle c_i t - r \rangle \mathcal{D}[u_i^{(2)}](t, r) \}, \\ e_3[u^{(3)}](t, r) &= \sum_{i=1}^m \{ (t+r) \Phi_0(c_i t, r)^{-1} (|u_i^{(3)}(t, r)| + \mathcal{D}_i^+[u_i^{(3)}](t, r)) \\ &\quad + (\langle c_i t - r \rangle^{-1} + \langle c_i t - r \rangle^{-2} \Phi_0(c_i t, r))^{-1} \mathcal{D}[u_i^{(3)}](t, r) \}, \end{aligned}$$

where $\Phi_0(c_i t, r)$ is given by (2.2), and $\mathcal{D}[v](t, r)$ and $\mathcal{D}_i^+[v](t, r)$ are from Lemma 2.6. We also define

$$E_k(T) = \sup_{(t,r) \in [0,T) \times [0,\infty)} e_k[u^{(k)}](t, r) \quad (k = 1, 2, 3)$$

for $0 < T \leq T_\varepsilon$, and $E_k(0) = \sup_{r \in [0,\infty)} e_k[u^{(k)}](0, r)$.

Proposition 3.1. *Assume $0 < T < T_\varepsilon$, and let M_k ($k = 1, 2, 3$) be positive constants. Suppose that ε is a positive constant satisfying*

$$M_3 \varepsilon^3 \leq M_2 \varepsilon^2 \leq M_1 \varepsilon \leq 1$$

and $\varepsilon \leq 1$. Then there exist three positive constants C_1, C_2 and C_3 (which are in-

dependent of $T, T_\varepsilon, \varepsilon,$ and M_k ($k = 1, 2, 3$) such that

$$(3.5) \quad E_k(T) \leq M_k \varepsilon^k \quad (k = 1, 2, 3)$$

implies

$$(3.6) \quad E_1(T) \leq C_1(\varepsilon + M_1^2 \varepsilon^2),$$

$$(3.7) \quad E_2(T) \leq C_2(M_1^2 \varepsilon^2 + (M_1^2 + M_2)M_2 \varepsilon^4 \log(2 + T) + M_3^2 \varepsilon^6 (\log(2 + T))^2),$$

$$(3.8) \quad E_3(T) \leq C_3(M_1 M_2 \varepsilon^3 + M_2 M_3 \varepsilon^5 \log(2 + T)).$$

Before proving Proposition 3.1, we show that Theorem 1.1 follows from it.

Proof of Theorem 1.1. Set

$$(3.9) \quad \begin{aligned} M_1 &= \max \left\{ \frac{2E_1(0)}{\varepsilon}, 4C_1 \right\}, & M_2 &= \max \left\{ \frac{2E_2(0)}{\varepsilon^2}, 6C_2 M_1^2 \right\}, \\ M_3 &= \max \left\{ \frac{2E_3(0)}{\varepsilon^3}, 4C_3 M_1 M_2 \right\}. \end{aligned}$$

Choose a positive constant ε_1 (≤ 1) which is small enough to satisfy

$$(3.10) \quad M_1 \varepsilon_1 \leq \frac{1}{4C_1} \quad \text{and} \quad M_3 \varepsilon_1^3 \leq M_2 \varepsilon_1^2 \leq M_1 \varepsilon_1 \leq 1.$$

Let $0 < \varepsilon \leq \varepsilon_1$, and assume $\varepsilon^2 \log(2 + T_\varepsilon) \leq C_4$, where

$$(3.11) \quad C_4 = \min \left\{ \frac{1}{6C_2(M_1^2 + M_2)}, \sqrt{\frac{M_2}{6C_2 M_3^2}}, \frac{1}{4C_3 M_2} \right\}.$$

Define

$$T = \sup\{S \in [0, T_\varepsilon]; E_k(S) \leq M_k \varepsilon^k \quad (k = 1, 2, 3)\}.$$

Since (3.9) implies $E_k(0) \leq M_k \varepsilon^k / 2$ ($k = 1, 2, 3$), the continuity of E_k implies $E_k(S) \leq M_k \varepsilon^k$ ($k = 1, 2, 3$) for some $S \in (0, T_\varepsilon]$. Hence T is positive.

Now suppose $T < T_\varepsilon$. Then, Proposition 3.1, (3.9), (3.10), and (3.11) lead to $E_k(T) \leq M_k \varepsilon^k / 2$ ($k = 1, 2, 3$), which contradict the definition of T because E_k ($k = 1, 2, 3$) are continuous functions. Hence we conclude $T = T_\varepsilon$, and $E_k(T_\varepsilon) \leq M_k \varepsilon^k$ for $1 \leq k \leq 3$.

From these *a priori* estimates, we see that $\sum_{|\alpha| \leq 1} \sum_{k=1}^3 |\partial_{t,r}^\alpha u^{(k)}(t, r)|$ is bounded for $(t, r) \in [0, T_\varepsilon] \times [0, \infty)$. Then, from the system (3.1)–(3.3), it is easy to show that $\sum_{|\alpha|=2} \sum_{k=1}^3 |\partial_{t,r}^\alpha u^{(k)}(t, r)|$ is also bounded for $(t, r) \in [0, T_\varepsilon] \times [0, \infty)$. Now the classical local existence theorem assures that we can extend the solution $(u^{(k)})_{1 \leq k \leq 3}$ beyond

T_ε . This contradicts the definition of T_ε . Accordingly we find $\varepsilon^2 \log(2 + T_\varepsilon) > C_4$ for $0 < \varepsilon \leq \varepsilon_1$, which immediately implies $T_\varepsilon \geq \exp(C_5 \varepsilon^{-2})$ for $\varepsilon \leq \varepsilon_0$ with appropriately chosen positive constants C_5 and ε_0 . This completes the proof. \square

Now we are going to prove Proposition 3.1.

Proof of Proposition 3.1. Let ϕ and ψ are radially symmetric C^2 -functions. First we observe that we have

$$(3.12) \quad \mathcal{M}[\phi(\partial_{t,r}^\gamma \psi)](t, r) \leq C(|\phi(t, r)| + \langle r \rangle^{-1} \mathcal{D}[\phi](t, r)) \mathcal{D}[\psi](t, r),$$

$$(3.13) \quad \mathcal{M}[(\partial_{t,r}^\beta \phi)(\partial_{t,r}^\gamma \psi)](t, r) \leq C \langle r \rangle^{-1} \mathcal{D}[\phi](t, r) \mathcal{D}[\psi](t, r)$$

for $|\beta| = |\gamma| = 1$, where \mathcal{M} is defined as in Lemma 2.6. Since

$$2Q_0(\phi, \psi; c_i) = (\partial_t \phi - c_i \partial_r \phi)(D_{+,c_i} \psi) + (\partial_t \psi - c_i \partial_r \psi)(D_{+,c_i} \phi)$$

for any radially symmetric functions ϕ and ψ , we also obtain

$$(3.14) \quad \begin{aligned} & \mathcal{M}[Q_0(\phi, \psi; c_i)](t, r) \\ & \leq C \langle r \rangle^{-1} (\mathcal{D}[\phi](t, r) \mathcal{D}_i^+[\psi](t, r) + \mathcal{D}[\psi](t, r) \mathcal{D}_i^+[\phi](t, r)). \end{aligned}$$

For $0 \leq j \leq m$, we define

$$\Lambda_j = \{(t, r) \in [0, \infty) \times [0, \infty); |c_j t - r| < \delta t\}$$

with some small $\delta > 0$, where $c_0 = 0$ as before. Note that $\langle c_j t - r \rangle$ is equivalent to $\langle t + r \rangle$ outside Λ_j . If δ is chosen sufficiently small, then there is no intersection between Λ_j and Λ_k for $j \neq k$. Hence we have

$$(3.15) \quad \langle c_j t - r \rangle^{-1} \langle c_k t - r \rangle^{-1} \leq C \langle t + r \rangle^{-1} w(t, r)^{-1}$$

for $j \neq k$. Moreover, if we have $j \neq i$ and $k \neq i$ in addition, we get

$$(3.16) \quad \langle c_j t - r \rangle^{-1} \langle c_k t - r \rangle^{-1} \leq C \langle t + r \rangle^{-1} w_i(t, r)^{-1}.$$

Similarly, for $\kappa \geq 0$ and $j \neq k$, we have

$$(3.17) \quad \langle c_j t - r \rangle^{-\kappa} \Phi_0(c_k t, r) \leq C(\langle c_j t - r \rangle^{-\kappa} + \langle t + r \rangle^{-\kappa} \Phi_0(c_k t, r)),$$

$$(3.18) \quad \Phi_0(c_j t, r) \Phi_0(c_k t, r) \leq C(\Phi_0(c_j t, r) + \Phi_0(c_k t, r)).$$

From (3.15), we especially have $\langle r \rangle \langle c_j t - r \rangle^{-1} \leq C \langle t + r \rangle^{-1}$ for $1 \leq j \leq m$, and we obtain

$$(3.19) \quad |u_j^{(1)}(t, r)| + \langle r \rangle^{-1} \mathcal{D}[u_j^{(1)}](t, r) \leq C \langle t + r \rangle^{-1} \langle c_j t - r \rangle^{-1} M_1 \varepsilon,$$

$$(3.20) \quad |u_j^{(k)}(t, r)| + \langle r \rangle^{-1} \mathcal{D}[u_j^{(k)}](t, r) \leq C \langle t + r \rangle^{-1} \Phi_0(c_j t, r) M_k \varepsilon^k$$

for $k = 2, 3$. Having these estimates in mind, we are going to evaluate each non-linearity.

First we estimate nonlinear terms contained in (3.1). We have

$$\begin{aligned} \mathcal{M}[N_i(\partial u^{(1)}, \partial u^{(1)})](t, r) &\leq C \sum_j \langle r \rangle^{-1} \langle t+r \rangle^{-1} \langle c_j t - r \rangle^{-3} M_1^2 \varepsilon^2 \\ &\leq C \langle t+r \rangle^{-2} w(t, r)^{-3} M_1^2 \varepsilon^2, \\ \mathcal{M}[r_i^I(\partial u^{(1)}, \partial u^{(1)})](t, r) &\leq C \sum_{j \neq k} \langle r \rangle^{-1} \langle c_j t - r \rangle^{-2} \langle c_k t - r \rangle^{-2} M_1^2 \varepsilon^2 \\ &\leq C \langle t+r \rangle^{-3} w(t, r)^{-2} M_1^2 \varepsilon^2, \\ \mathcal{M}[r_i^{III}(u^{(1)}, \partial_t u^{(1)})](t, r) &\leq C \sum_{j \neq k} \langle t+r \rangle^{-1} \langle c_j t - r \rangle^{-1} \langle c_k t - r \rangle^{-2} M_1^2 \varepsilon^2 \\ &\leq C \langle t+r \rangle^{-2} w(t, r)^{-2} M_1^2 \varepsilon^2. \end{aligned}$$

On the other hand, since H_i is a rotationally invariant function of cubic order, we have

$$\begin{aligned} \mathcal{M}[H_i(u^{(1)}, \partial u^{(1)})](t, r) &\leq C \{ \langle r \rangle |u^{(1)}|^3 + (|u^{(1)}|^2 + |\partial_{t,r} u^{(1)}|^2) \mathcal{D}[u^{(1)}] \} \\ &\leq C \langle t+r \rangle^{-2} w(t, r)^{-3} M_1^3 \varepsilon^3. \end{aligned}$$

Summing up, we get

$$(3.21) \quad \mathcal{M}[\square_{c_i} u_i^{(1)}](t, r) \leq C \langle t+r \rangle^{-2} w(t, r)^{-2} M_1^2 \varepsilon^2.$$

Hence Lemmas 2.1, 2.8, 2.2, and Lemma 2.6-(i), (iii) with $(\rho, \mu) = (2, 1)$ lead to

$$(3.22) \quad E_1(T) \leq C_1(\varepsilon + M_1^2 \varepsilon^2).$$

Next we turn our attention to (3.2). Let ν be a positive and small constant. Then we have

$$(3.23) \quad \Phi_0(c_i t, r) \leq C \langle t+r \rangle^\nu \langle c_i t - r \rangle^{-\nu}.$$

Using this inequality, we start with

$$\begin{aligned} &\mathcal{M}[N_i(\partial u^{(2)+(3)}, \partial u^{(2)+(3)})](t, r) \\ &\leq C \sum_j \langle r \rangle^{-1} \langle t+r \rangle^{-1} \langle c_j t - r \rangle^{-1} \Phi_0(c_j t, r)^2 (M_2 \varepsilon^2 + M_3 \varepsilon^3)^2 \\ &\leq C \langle t+r \rangle^{-2+2\nu} w(t, r)^{-1-2\nu} M_2^2 \varepsilon^4, \\ &\mathcal{M}[r_i^I(\partial u^{(2)+(3)}, \partial u^{(2)+(3)})](t, r) \\ &\leq C \sum_{j \neq k} \langle r \rangle^{-1} \langle c_j t - r \rangle^{-1} \langle c_k t - r \rangle^{-1} \Phi_0(c_j t, r) \Phi_0(c_k t, r) (M_2 \varepsilon^2 + M_3 \varepsilon^3)^2 \\ &\leq C \langle t+r \rangle^{-2+\nu} w(t, r)^{-1-\nu} M_2^2 \varepsilon^4. \end{aligned}$$

Since $\partial u^{(1)}$ enjoys a better estimate than $\partial u^{(2)+(3)}$, it is easy to obtain

$$\begin{aligned} \mathcal{M}[N_i(\partial u^{(1)}, \partial u^{(2)+(3)})](t, r) &\leq C \langle t+r \rangle^{-2+\nu} w(t, r)^{-2-\nu} M_1 M_2 \varepsilon^3, \\ \mathcal{M}[r_i^I(\partial u^{(1)}, \partial u^{(2)+(3)})](t, r) &\leq C \langle t+r \rangle^{-2+\nu} w(t, r)^{-2-\nu} M_1 M_2 \varepsilon^3. \end{aligned}$$

Now we are proceeding to rather delicate parts. For simplicity of exposition, we set $u = u^{(1)+(2)+(3)}$. Then we have

$$\begin{aligned} &\mathcal{M}[r_i^{II}(\partial u, \partial u)](t, r) \\ &\leq C \sum_{j \neq i} \langle r \rangle^{-1} \{ \langle c_j t - r \rangle^{-4} M_1^2 \varepsilon^2 + \langle c_j t - r \rangle^{-2} (M_2^2 \varepsilon^4 + M_3^2 \varepsilon^6) \\ &\quad + \langle c_j t - r \rangle^{-4} \Phi_0(c_j t, r)^2 M_3^2 \varepsilon^6 \} \\ &\leq C \langle t+r \rangle^{-1} w_i(t, r)^{-2} (M_1^2 \varepsilon^2 + M_3^2 \varepsilon^6 \{\log(2+T)\}^2) \end{aligned}$$

for $0 \leq t < T$. Here we have used $\Phi_0(c_j t, r) \leq C \log(2+t)$. We also get

$$\begin{aligned} &\mathcal{M}[r_i^{III}(u^{(1)}, \partial_t u^{(2)})](t, r) \\ &\leq C \sum_{j \neq k} \langle t+r \rangle^{-1} \langle c_j t - r \rangle^{-1} \langle c_k t - r \rangle^{-1} M_1 M_2 \varepsilon^3 \\ &\leq C \langle t+r \rangle^{-2} w(t, r)^{-1} M_1 M_2 \varepsilon^3 \leq C \langle t+r \rangle^{-2+\nu} w(t, r)^{-1-\nu} M_1 M_2 \varepsilon^3, \\ &\mathcal{M}[r_i^{III}(u^{(2)+(3)}, \partial_t u^{(2)})](t, r) \\ &\leq C \sum_{j \neq k} \langle t+r \rangle^{-1} \Phi_0(c_j t, r) \langle c_k t - r \rangle^{-1} (M_2 \varepsilon^2 + M_3 \varepsilon^3) M_2 \varepsilon^2 \\ &\leq C (\langle t+r \rangle^{-1} w(t, r)^{-1} + \langle t+r \rangle^{-2+\nu} w(t, r)^{-\nu}) M_2^2 \varepsilon^4 \\ &\leq C \langle t+r \rangle^{-1} w(t, r)^{-1} M_2^2 \varepsilon^4. \end{aligned}$$

Setting $\tilde{H}_i = H_i(u, \partial u) - H_i(u^{(1)}, \partial u^{(1)})$, we obtain

$$\begin{aligned} \mathcal{M}[\tilde{H}_i](t, r) &\leq C \left\{ \sum_{k=1}^3 (|u^{(k)}| + \langle r \rangle^{-1} \mathcal{D}[u^{(k)}]) \right\}^2 \sum_{k=2}^3 (\langle r \rangle |u^{(k)}| + \mathcal{D}[u^{(k)}]) \\ &\leq C \langle t+r \rangle^{-2+3\nu} w(t, r)^{-3\nu} M_1^2 M_2 \varepsilon^4 \\ &\leq C \langle t+r \rangle^{-1} w(t, r)^{-1} M_1^2 M_2 \varepsilon^4. \end{aligned}$$

Now we set

$$\begin{aligned} G_{i,1} &= r_i^{II}(\partial u, \partial u), \quad G_{i,2} = r_i^{III}(u^{(2)+(3)}, \partial_t u^{(2)}) + \tilde{H}_i, \\ G_{i,3} &= \square_{c_i} u_i^{(2)} - G_{i,1} - G_{i,2}. \end{aligned}$$

Since we have shown

$$\mathcal{M}[G_{i,3}](t, r) \leq C \langle t+r \rangle^{-2+2\nu} w(t, r)^{-1-\nu} M_1 M_2 \varepsilon^3,$$

from Lemma 2.2 and Lemma 2.6-(i), (iii) with $(\rho, \mu) = (2 - 2\nu, \nu)$ (note that we may assume $2 - 2\nu > 1$), we get

$$(3.24) \quad e_2[U_{c_i}[G_{i,3}]](t, r) \leq C M_1 M_2 \varepsilon^3 \leq C M_1^2 \varepsilon^2.$$

Also these lemmas with $(\rho, \mu) = (1, 0)$ yield

$$(3.25) \quad e_2[U_{c_i}[G_{i,2}]](t, r) \leq C(M_1^2 + M_2)M_2 \varepsilon^4 \log(2+t).$$

On the other hand, by Lemmas 2.2 and 2.6-(ii), (iii) with $(\rho, \mu) = (1, 1)$, we get

$$(3.26) \quad e_2[U_{c_i}[G_{i,1}]](t, r) \leq C(M_1^2 \varepsilon^2 + M_3^2 \varepsilon^6 \{\log(2+T)\}^2).$$

Now (3.24), (3.25) and (3.26) imply

$$(3.27) \quad E_2(T) \leq C_2(M_1^2 \varepsilon^2 + (M_1^2 + M_2)M_2 \varepsilon^4 \log(2+T) + M_3^2 \varepsilon^6 \{\log(2+T)\}^2)$$

with an appropriate constant C_2 .

Finally we consider (3.3). We get

$$\begin{aligned} & \mathcal{M}[r_i^{\text{III}}(u^{(2)+(3)}, \partial_t u^{(1)})] \\ & \leq C \sum_{j \neq k} \langle t+r \rangle^{-1} \Phi_0(c_j t, r) \langle c_k t - r \rangle^{-2} M_1 \varepsilon (M_2 \varepsilon^2 + M_3 \varepsilon^3) \\ & \leq C(\langle t+r \rangle^{-1} w(t, r)^{-2} + \langle t+r \rangle^{-3+\nu} w(t, r)^{-\nu}) M_1 M_2 \varepsilon^3 \\ & \leq C \langle t+r \rangle^{-1} w(t, r)^{-2} M_1 M_2 \varepsilon^3, \end{aligned}$$

where $\nu (> 0)$ is a small constant. We also obtain

$$\begin{aligned} \mathcal{M}[r_i^{\text{III}}(u^{(1)}, \partial_t u^{(3)})] & \leq C \sum_{j \neq k} \langle t+r \rangle^{-1} \langle c_j t - r \rangle^{-1} \langle c_k t - r \rangle^{-1} \\ & \quad \times (1 + \langle c_k t - r \rangle^{-1} \Phi_0(c_k t, r)) M_1 M_3 \varepsilon^4 \\ & \leq C(\langle t+r \rangle^{-2} w(t, r)^{-1} + \langle t+r \rangle^{-2+\nu} w(t, r)^{-2-\nu}) M_1 M_3 \varepsilon^4 \\ & \leq C \langle t+r \rangle^{-1} w(t, r)^{-2} M_1 M_3 \varepsilon^4. \end{aligned}$$

From Lemma 2.2 and Lemma 2.6-(i), (iii) with $(\rho, \mu) = (1, 1)$, we obtain

$$(3.28) \quad e_3[U_{c_i}[r_i^{\text{III}}(u^{(2)+(3)}, \partial_t u^{(1)}) + r_i^{\text{III}}(u^{(1)}, \partial_t u^{(3)})]](t, r) \leq C M_1 M_2 \varepsilon^3.$$

On the other hand, for $j \neq k$ and small $\nu (> 0)$, we have

$$\begin{aligned} & \Phi_0(c_j t, r) \langle c_k t - r \rangle^{-1} (1 + \langle c_k t - r \rangle^{-1} \Phi_0(c_k t, r)) \\ & \leq C (\langle t + r \rangle^{-1} \Phi_0(c_j t, r) + w(t, r)^{-1} + w(t, r)^{-2} \Phi_0(c_k t, r)) \\ & \leq C (\langle t + r \rangle^{-1+\nu} w(t, r)^{-\nu} + w(t, r)^{-1} + w(t, r)^{-2} \log(2 + T)) \\ & \leq C (w(t, r)^{-1} + w(t, r)^{-2} \log(2 + T)). \end{aligned}$$

Hence we get

$$\begin{aligned} & \mathcal{M}[r_i^{\text{III}}(u^{(2)+(3)}, \partial_t u^{(3)})](t, r) \\ & \leq C \sum_{j \neq k} \langle t + r \rangle^{-1} \Phi_0(c_j t, r) \langle c_k t - r \rangle^{-1} \\ & \quad \times (1 + \langle c_k t - r \rangle^{-1} \Phi_0(c_k t, r)) (M_2 \varepsilon^2 + M_3 \varepsilon^3) M_3 \varepsilon^3 \\ & \leq C \langle t + r \rangle^{-1} (w(t, r)^{-1} + w(t, r)^{-2} \log(2 + T)) M_2 M_3 \varepsilon^5. \end{aligned}$$

Therefore, by Corollaries 2.3 and 2.7, we obtain

$$(3.29) \quad e_3[U_i[r_i^{\text{III}}(u^{(2)+(3)}, \partial_t u^{(3)})]](t, r) \leq C M_2 M_3 \varepsilon^5 \log(2 + T).$$

Finally (3.28) and (3.29) imply

$$(3.30) \quad E_3(T) \leq C_3 \{M_1 M_2 \varepsilon^3 + M_2 M_3 \varepsilon^5 \log(2 + T)\}.$$

This completes the proof. □

4. Proof of Theorem 1.2

Suppose that all the assumptions in Theorem 1.2 are fulfilled. Let $u = (u_1, u_2)$ be a solution to (1.1)–(1.2) (with (1.19) and (1.20)) for $0 \leq t < T_\varepsilon$. Fix κ satisfying $1/2 < \kappa < 1$. We put

$$\begin{aligned} e_1^*[u_1](t, r) &= \langle t + r \rangle^\kappa (|u_1(t, r)| + \mathcal{D}_1^+[u_1](t, r)) \\ & \quad + \langle c_1 t - r \rangle^\kappa \mathcal{D}[u_1](t, r), \\ e_2^*[u_2](t, r) &= \langle t + r \rangle \Phi_0(c_2 t, r)^{-1} (|u_2(t, r)| + \mathcal{D}_2^+[u_2](t, r)) \\ & \quad + \langle c_2 t - r \rangle \mathcal{D}[u_2](t, r), \end{aligned}$$

and

$$(4.1) \quad E(T) = \sup_{(t,r) \in [0,T] \times [0,\infty)} (e_1^*[u_1](t, r) + e_2^*[u_2](t, r))$$

for $0 < T \leq T_\varepsilon$, with $E(0) = \sup_{r \in [0,\infty)} (e_1^*[u_1](0, r) + e_2^*[u_2](0, r))$.

Similarly to the proof of Theorem 1.1, what we need for the proof of Theorem 1.2 is the following.

Proposition 4.1. *Assume $0 < T < T_\varepsilon$, and let M_0 be a positive constant. Suppose that ε is a positive constant satisfying $M_0\varepsilon \leq 1$ and $\varepsilon \leq 1$. Then there exists a positive constant C_0 , which is independent of T , T_ε , ε , and M_0 , such that $E(T) \leq M_0\varepsilon$ implies*

$$(4.2) \quad E(T) \leq C_0(\varepsilon + M_0^2\varepsilon^2).$$

From Proposition 4.1, following a similar argument in the proof of Theorem 1.1, we see that $E(T)$ stays bounded as far as the solution u exists for $0 \leq t < T$, and the local existence theorem implies Theorem 1.2.

Proof of Proposition 4.1. Lemmas 2.1 and 2.8 yield

$$(4.3) \quad \sum_{i=1}^2 e_i^*[U_{c_i}^*[\varepsilon f_i, \varepsilon g_i]](t, r) \leq C\varepsilon.$$

As for the nonlinearities, firstly we have

$$\begin{aligned} \mathcal{M}[Q_0(u_1, u_1; c_1)](t, r) &\leq C\langle r \rangle^{-1} \langle t+r \rangle^{-\kappa} \langle c_1 t - r \rangle^{-\kappa} M_0^2 \varepsilon^2 \\ &\leq C\langle t+r \rangle^{-2\kappa} w(t, r)^{-1} M_0^2 \varepsilon^2 \\ &\leq C\langle t+r \rangle^{-1-\nu} w(t, r)^{-2\kappa+\nu} M_0^2 \varepsilon^2 \end{aligned}$$

for $\nu > 0$ satisfying $2\kappa - \nu > 1$. Similarly we get

$$\begin{aligned} \mathcal{M}[Q_0(u_2, u_2; c_2)](t, r) &\leq C\langle r \rangle^{-1} \langle t+r \rangle^{-1} \langle c_2 t - r \rangle^{-1} \Phi_0(c_2 t, r) M_0^2 \varepsilon^2 \\ &\leq C\langle t+r \rangle^{-2+\nu} w(t, r)^{-1-\nu} M_0^2 \varepsilon^2 \\ &\leq C\langle t+r \rangle^{-1-\nu} w(t, r)^{-2+\nu} M_0^2 \varepsilon^2 \end{aligned}$$

for small $\nu > 0$. Thus we obtain

$$(4.4) \quad \mathcal{M}[N_i(\partial u, \partial u)](t, r) \leq C\langle t+r \rangle^{-1-\nu} w(t, r)^{-2\kappa+\nu} M_0^2 \varepsilon^2.$$

On the other hand, we have

$$\begin{aligned} \mathcal{M}[r_i^1(\partial u, \partial u)](t, r) &\leq C\langle r \rangle^{-1} \langle c_1 t - r \rangle^{-\kappa} \langle c_2 t - r \rangle^{-1} M_0^2 \varepsilon^2 \\ &\leq C\langle t+r \rangle^{-1-\kappa} w(t, r)^{-1} M_0^2 \varepsilon^2 \\ &\leq C\langle t+r \rangle^{-1-\kappa/2} w(t, r)^{-1-\kappa/2} M_0^2 \varepsilon^2. \end{aligned}$$

Summing up, from Lemma 2.2, (i) and (iii) in Lemma 2.6, we obtain

$$(4.5) \quad e_i^*[U_{c_i}[N_i(\partial u, \partial u) + r_i^1(\partial u, \partial u)]](t, r) \leq CM_0^2\varepsilon^2$$

for $i = 1, 2$.

Now we are going to estimate the main parts. For r_1^{IV} , we have

$$\begin{aligned} \mathcal{M}[r_1^{IV}(u, \partial u)](t, r) &\leq C\langle t+r \rangle^{-1}\Phi_0(c_2t, r)\langle c_2t-r \rangle^{-1}M_0^2\varepsilon^2 \\ &\leq C\langle t+r \rangle^{-1+\nu}w_1(t, r)^{-1-\nu}M_0^2\varepsilon^2 \end{aligned}$$

for small $\nu > 0$. By (ii) in Lemma 2.6 with $(\rho, \mu) = (1 - \nu, \nu)$, we get

$$(4.6) \quad \langle c_1t-r \rangle^\kappa \mathcal{D}[U_{c_1}[r_1^{IV}]](t, r) \leq CM_0^2\varepsilon^2,$$

provided that ν is small enough to satisfy $0 < \kappa \leq 1 - \nu$.

On the other hand, since

$$\langle t+r \rangle^{-1+\nu}w_1(t, r)^{-1-\nu} \leq C\langle t+r \rangle^{1-\kappa}\langle t+r \rangle^{-2+\kappa+\nu}w_1(t, r)^{-1-\nu}$$

and $\langle \tau + \lambda \rangle \leq \langle t + r \rangle$ for $(\tau, \lambda) \in \Theta_i^*(t, r)$, we get

$$\sup_{(\tau, \lambda) \in \Theta_i^*(t, r)} \langle \tau + \lambda \rangle^{2-\kappa-\nu}w(\tau, \lambda)^{1+\nu} \mathcal{M}[r_1^{IV}](\tau, \lambda) \leq CM_0^2\varepsilon^2\langle t+r \rangle^{1-\kappa}.$$

Now, since we may assume $2 - \kappa - \nu > 1$, Lemmas 2.2 and 2.6–(iii) with $(\rho, \mu) = (2 - \kappa - \nu, \nu)$ lead to

$$(4.7) \quad |U_{c_1}[r_1^{IV}](t, r)| + \mathcal{D}_1^+[U_{c_1}[r_1^{IV}]](t, r) \leq C\langle t+r \rangle^{-\kappa}M_0^2\varepsilon^2.$$

From (4.6) and (4.7), we obtain

$$(4.8) \quad e_1^*[U_{c_1}[r_1^{IV}(u, \partial u)]](t, r) \leq CM_0^2\varepsilon^2.$$

Similarly we also obtain

$$(4.9) \quad e_1^*[U_{c_1}[r_1^{II}(\partial u, \partial u)]](t, r) \leq CM_0^2\varepsilon^2.$$

Since we have $2\kappa > 1$, we get

$$\begin{aligned} \mathcal{M}[r_2^{II}(\partial u, \partial u)](t, r) &\leq C\langle r \rangle^{-1}\langle c_1t-r \rangle^{-2\kappa}M_0^2\varepsilon^2 \\ &\leq C\langle t+r \rangle^{-1}w_2(t, r)^{-2\kappa}M_0^2\varepsilon^2. \end{aligned}$$

Hence, from Lemma 2.2, (ii) and (iii) in Lemma 2.6, we obtain

$$(4.10) \quad e_2^*[U_{c_2}[r_2^{II}(\partial u, \partial u)]](t, r) \leq C_0M_0^2\varepsilon^2.$$

Finally, (4.2) follows from (4.3), (4.5), (4.8), (4.9) and (4.10). This completes the proof. \square

References

- [1] F. Asakura: *Existence of a global solution to a semi-linear wave equation with slowly decreasing initial data in three space dimensions*, Comm. Partial Differential Equations **11** (1986), 1459–1487.
- [2] D. Christodoulou: *Global solutions of nonlinear hyperbolic equations for small initial data*, Comm. Pure Appl. Math. **39** (1986), 267–282.
- [3] F. John: *Blow-up for quasi-linear wave equations in three space-dimensions*, Comm. Pure Appl. Math. **34** (1981), 29–51.
- [4] F. John and S. Klainerman: *Almost global existence to general nonlinear wave equations in three space-dimensions*, Comm. Pure Appl. Math. **37** (1984), 235–268.
- [5] S. Katayama: *Global existence for a class of systems of nonlinear wave equations in three space dimensions*, Chinese Ann. Math. **25B** (2004), 463–482.
- [6] S. Katayama: *Global and almost-global existence for systems of nonlinear wave equations with different propagation speeds*, Diff. Integral Eqs. **17** (2004), 1043–1078.
- [7] S. Katayama: *Global existence for systems of wave equations with nonresonant nonlinearities and null forms*, J. Differential Equations **209** (2005), 140–171.
- [8] S. Katayama and A. Matsumura: *Sharp lower bound for the lifespan of the systems of semi-linear wave equations with multiple speeds*, J. Math. Kyoto Univ. **45** (2005), 391–403.
- [9] S. Katayama and K. Yokoyama: *Global small amplitude solutions to systems of nonlinear wave equations with multiple speeds*, Osaka J. Math. **43** (2006), 283–326.
- [10] S. Klainerman: *Uniform decay estimates and the Lorentz invariance of the classical wave equation*, Comm. Pure Appl. Math. **38** (1985), 321–332.
- [11] S. Klainerman: *The null condition and global existence to nonlinear wave equations*; in Non-linear Systems of Partial Differential Equations in Applied Mathematics, Lectures in Applied Math. **23**, Amer. Math. Soc., Providence, RI, 1986, 293–326.
- [12] S. Klainerman and T.C. Sideris: *On almost global existence for nonrelativistic wave equations in 3D*, Comm. Pure Appl. Math. **49** (1996), 307–321.
- [13] M. Kovalyov: *Resonance-type behaviour in a system of nonlinear wave equations*, J. Differential Equations **77** (1989), 73–83.
- [14] H. Kubo: *Blow-up of solutions to semilinear wave equations with initial data of slow decay in low space dimensions*, Diff. Integral Eqs. **7** (1994), 315–321.
- [15] K. Kubota and K. Yokoyama: *Global existence of classical solutions to systems of nonlinear wave equations with different speeds of propagation*, Japanese J. Math. **27** (2001), 113–202.
- [16] H. Lindblad: *On the lifespan of solutions of nonlinear wave equations with small initial data*, Comm. Pure Appl. Math. **43** (1990), 445–472.
- [17] M. Ohta: *Counterexample to global existence for system of nonlinear wave equations with different propagation speeds*, Funkcialaj Ekvacioj **46** (2003), 471–477.
- [18] T.C. Sideris: *Global behavior of solutions to nonlinear wave equations in three space dimensions*, Comm. Partial Differential Equations **8** (1983), 1283–1323.
- [19] T.C. Sideris and S.-Y. Tu: *Global existence for systems of nonlinear wave equations in 3D with multiple speeds*, SIAM J. Math. Anal. **33** (2001), 477–488.
- [20] C.D. Sogge: *Global existence for nonlinear wave equations with multiple speeds*; in Harmonic Analysis at Mount Holyoke, Contemp. Math. **320**, Amer. Math. Soc., Providence, RI, 2003, 353–366.
- [21] K. Yokoyama: *Global existence of classical solutions to systems of wave equations with critical nonlinearity in three space dimensions*, J. Math. Soc. Japan **52** (2000), 609–632.

Department of Mathematics
Wakayama University
930 Sakaedani, Wakayama 640-8510
Japan
e-mail: katayama@center.wakayama-u.ac.jp