



Title	On semifield planes of even order
Author(s)	Hiramine, Yutaka
Citation	Osaka Journal of Mathematics. 1983, 20(3), p. 645-658
Version Type	VoR
URL	<a href="https://doi.org/10.18910/4671">https://doi.org/10.18910/4671</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## ON SEMI-FIELD PLANES OF EVEN ORDER

YUTAKA HIRAMINE

(Received June 30, 1981)

### 1. Introduction

Let  $\pi$  be a non-Desarguesian semi-field plane with an autotopism group  $G$  and let  $u(\pi)$  denote the number of the orbits of  $G$  on the points not incident with any side of the autotopism triangles.

In their paper [9], M.J. Kallaher and R.A. Liebler have conjectured that  $u(\pi) \geq 5$  and they have proved that the conjecture is true if  $G$  is solvable and the order of  $\pi$  is not  $2^6$ .

In this paper we treat semi-field planes of even order whose autotopism groups are not necessarily solvable and prove the following.

**Theorem 1.** *Let  $\pi$  be a non-Desarguesian semi-field plane of order  $2^r$ . If  $r$  is not divisible by 4, then  $u(\pi) \geq 5$ .*

The proof requires the use of the Kallaher-Liebler's theorem mentioned above and the following lemma which we prove in section 3.

**Lemma 2.** *Let  $\pi$  be a non-Desarguesian semi-field plane of order  $2^6$  with a solvable autotopism group. Then  $u(\pi) \geq 5$ .*

### 2. Notations and preliminaries

Our notation is largely standard and taken from [3] and [6]. Let  $G$  be a permutation group on  $\Omega$ . For  $X \leq G$  and  $\Delta \subset \Omega$ , we define  $F(X) = \{\alpha \in \Omega \mid \alpha^x = \alpha \text{ for all } x \in X\}$ ,  $X(\Delta) = \{x \in X \mid \Delta^x = \Delta\}$ ,  $X_\Delta = \{x \in X \mid \alpha^x = \alpha \text{ for all } \alpha \in \Delta\}$  and  $X^\Delta = X(\Delta)/X_\Delta$ , the restriction of  $X$  on  $\Delta$ . When  $X$  is a collineation group of a projective plane, we denote by  $F(X)$  the set of fixed points and fixed lines of  $X$ .

**Lemma 2.1.** *Let  $G$  be a transitive permutation group on a finite set  $\Omega$ ,  $H$  a stabilizer of a point of  $\Omega$  and  $M$  a nonempty subset of  $G$ . Then  $|F(M)| = |N_G(M)| \times |ccl_G(M) \cap H|/|H|$ . Here  $ccl_G(M) \cap H = \{g^{-1}Mg \mid g^{-1}Mg \subset H, g \in G\}$ .*

Proof. Set  $W = \{(L, \alpha) \mid L \in ccl_G(M), \alpha \in F(L)\}$  and  $W_\alpha = \{L \mid (L, \alpha) \in W\}$ ,

$\alpha \in F(L)\}$ . By the transitivity of  $G$ ,  $|W_\alpha| = |W_\beta|$  holds for every  $\alpha, \beta \in \Omega$ . Counting the number of elements of  $W$  in two ways, we obtain  $|G: N_G(M)| \times |F(M)| = |G: H| \times |ccl_G(M) \cap H|$ . Thus we have the lemma.

**Lemma 2.2.** *Let  $PG(2, q)$  denote the Desarguesian projective plane of order  $q$  where  $q = 2^n$  and  $n \equiv 1 \pmod{2}$ . Set  $Y = PSL(3, q)$  and  $X = \langle f \rangle Y$ , where  $f$  is a field automorphism of  $Y$  of order  $n$ . Set  $G = X_{P, Q, R}$  and  $N = G \cap Y$ , where  $P = [1, 0, 0]$ ,  $Q = [0, 1, 0]$  and  $R = [0, 0, 1]$ .*

(i) *Let  $A$  be a noncyclic abelian  $p$ -subgroup of  $G$  of order  $p^2$  for a prime  $p$ . Then  $A$  is not semi-regular on the set of points contained in  $PG(2, q) - F(A)$ .*

(ii) *Let  $C$  be a cyclic subgroup of  $G$  of order  $q-1$ . Then  $C \subset N$ .*

**Proof.** Since  $A \cap N \neq 1$  and  $N \simeq Z_{q-1} \times Z_{q-1}$ ,  $p$  is an odd prime. Let  $T$  be the translation group with respect to the line  $g$  joining  $[1, 0, 0]$  and  $[0, 1, 0]$ . Deny (i) and let  $\Omega$  denote the set of points in  $F(A)$ . Then, by Theorem 5.3.6 of [3],  $T = \langle C_T(x) \mid 1 \neq x \in A \rangle$ . By the semi-regularity of  $A$ ,  $C_T(x)$  acts on  $\Omega$  for each  $x \in A - \{1\}$ . Hence  $T$  acts on  $\Omega$ .

Let  $\Delta$  denote the set of points not incident with the line  $g$ . Clearly  $[0, 0, 1] \in \Delta \cap \Omega$ . Since  $T$  is transitive on  $\Delta$ , we have (i).

Set  $D = C \cap N$  and let  $\begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix}$  be a generator of  $D$ . Then  $C \triangleright D$  and  $C/D \simeq CN/N \leq G/N \simeq Z_n$  and so  $|D| \geq (q-1)/n$ . Set  $\langle h \rangle = C_{\langle f \rangle}(D)$  and  $s = |\langle h \rangle|$ . Then  $n = r \times s$  for an integer  $r$ . It follows that  $ba^{-1}, ca^{-1} \in GF(2^r)^\times$ . Hence  $|D| \leq 2^r - 1$ . From this,  $2^r - 1 \geq |D| \geq (q-1)/n$ . We can easily verify that  $s = 1$ . Therefore  $C_{\langle f \rangle}(D) = 1$ , whence  $C \leq C_G(D) = NC_{\langle f \rangle}(D) = N$ . Thus  $C \leq N$ .

In the rest of the paper we assume the following.

**Hypothesis 2.3.** *Let  $\pi$  be a non-Desarguesian semi-field plane of order  $2^r$  coordinatized by a semi-field  $D$  with respect to the points  $U_1 = (0, 0)$ ,  $U_2 = (0)$ ,  $U_3 = (\infty)$  and let  $G$  be the autotopism group of  $\pi$  with respect to  $U_1, U_2, U_3$ . Let  $l_i$  be the line joining  $U_j$  and  $U_k$  for  $i, j, k$  with  $\{i, j, k\} = \{1, 2, 3\}$  and let  $\Phi(\pi)$  be the set of points of  $\pi$  not incident with  $l_1, l_2$  or  $l_3$ . Let  $u(\pi)$  denote the number of  $G$ -orbits on  $\Phi(\pi)$ . Set  $K_i = G_{(U_i, l_i)}$  for  $1 \leq i \leq 3$  and let  $N_1, N_2$  or  $N_3$  be the right, middle or left nucleus, respectively.*

$D$  may be considered as a right vector space over  $N_1$  or  $N_2$  and as a left vector space over  $N_2$  or  $N_3$ . The multiplicative group  $N_i^\times$  is isomorphic to  $K_i$  for each  $i$  with  $1 \leq i \leq 3$  (Chapter 8 of [6]). Set  $\bar{l}_i = l_i - \{U_j, U_k\}$  for  $i, j$ , with  $\{i, j, k\} = \{1, 2, 3\}$ .

### 3. The proof of Lemma 2.

Throughout this section  $\pi$  is a projective plane satisfying the hypothesis 2.3 and the following.

**Hypothesis 3.1.** (i) The order of  $\pi$  is  $2^6$ .

(ii) Set  $u=u(\pi)$ . Then  $u \leq 4$ .

(iii) The autotopism group  $G$  is solvable.

**Lemma 3.2.**  $|K_t|=1, 3$  or  $7$  for every  $t \in \{1, 2, 3\}$  and  $u=3$  or  $4$ .

Proof. Since  $\pi$  is non-Desarguesian,  $D$  is not a field. Hence,  $N_t$  is isomorphic to  $GF(2)$ ,  $GF(4)$  or  $GF(8)$  for  $t \in \{1, 2, 3\}$ . By Theorem 8.2 of [6],  $|K_t|=1, 3$  or  $7$ .

By Corollary 4.1.1 of [9] and Hypothesis 3.1 (ii),  $u=3$  or  $4$ .

**Lemma 3.3.** If  $G$  is transitive on  $\bar{l}_t$  for some  $t \in \{1, 2, 3\}$ , then the following hold.

(i)  $G/K_t \leq \Gamma L(1, 2^6)$  and  $G/K_t$  contains an element of order 9.

(ii) Let  $m$  be an arbitrary line through  $U_t$  such that  $m \neq l_j, l_k$  for  $\{t, j, k\} = \{1, 2, 3\}$ . Set  $A = m \cap l_t$ . Then  $G_m = G_A$ ,  $|G : G_A| = 3^2 \cdot 7$  and the number of  $G_A$ -orbits on  $m - \{U_t, A\}$  is equal to  $u$ .

(iii) Let  $\Delta_1, \Delta_2, \dots, \Delta_u$  be the orbits stated in (ii). Set  $x_s = |\Delta_s|$ ,  $1 \leq s \leq u$ , and assume that  $x_1 \leq x_2 \leq \dots \leq x_u$ . Then  $|G_A|$  is divisible by  $x_s$  for every  $s$  and  $6 \times |K_t|$  is divisible by  $|G_A|$ . Furthermore  $\sum_{s=1}^u x_s = 63$ .

Proof. By Lemma 2.1 of [9],  $G$  is a transitive linear group on  $D$ . Hence it follows from a Huppert's theorem ([7]) that  $G/K_t \leq \Gamma L(1, 2^6)$ . If  $G/K_t$  contains no element of order 9, then its Sylow 3-subgroup is an elementary abelian 3-subgroup of order at most 9. By the structure of  $\Gamma L(1, 2^6)$ ,  $G/K_t$  is not a transitive linear group, a contradiction. Thus  $G/K_t$  contains an element of order 9 and (i) holds.

Let  $m, A$  be as in (ii). Since  $G$  fixes  $U_t$  and  $l_t$ , we have  $G_m = G_A$ . Clearly  $|G : G_A| = |A^G| = |\bar{l}_t| = 2^6 - 1 = 3^2 \cdot 7$ . As any point of  $\Phi(\pi)$  lies on a line of  $[U_t] - \{l_j, l_k\}$ ,  $\Phi(\pi) \cap m (= m - \{U_t, A\})$  is a union of  $u$   $G_A$ -orbits, hence (ii) holds.

Since  $G/K_t \leq \Gamma L(1, 2^6)$ ,  $G_A/K_t \leq Z_6$ . Hence  $6 \times |K_t|$  is divisible by  $|G_A|$ . Clearly  $x_s = |\Delta_s|$  divides  $|G_A|$  and  $\sum_{s=1}^u x_s = |\Delta_1 \cup \Delta_2 \cup \dots \cup \Delta_u| = |\bar{l}_t| = 2^6 - 1 = 3^2 \cdot 7$ . Thus (iii) holds.

**Lemma 3.4.** Suppose  $u=4$ . Then there exists  $i \in \{1, 2, 3\}$  having the following properties:

- (i)  $G$  is transitive on  $\bar{l}_i$ ,
- (ii)  $K_i$  is isomorphic to  $Z_7$  and  $G$  has a normal Sylow 7-subgroup and
- (iii)  $|G:G_A|=63$ ,  $G_A/K_i$  is isomorphic to  $Z_6$  and  $C_{G_A}(K_i)=K_i$  for each  $A \in \bar{l}_i$ .

Proof. By Lemma 6.1 of [9], there exists  $i \in \{1, 2, 3\}$  such that  $G$  is transitive on  $\bar{l}_i$ . Assume that  $K_i \neq Z_7$ . Then  $K_i \leq Z_3$  by Lemma 3.2. Let  $m, A, x_s$  be as in Lemma 3.3. We have  $x_s \mid 6 \mid |K_i| = 6$  or 18 and  $x_1 + x_2 + x_3 + x_4 = 63$ , hence  $|K_i| = 3$ ,  $|G_A| = 18$  and  $(x_1, x_2, x_3, x_4) = (9, 18, 18, 18)$ .

Let  $z$  be an involution in  $G_A$ . Then  $z$  is a Baer involution and so  $|F(z) \cap (m - \{U_i, A\})| = 7$  because  $m \in F(z)$ . If  $F(z) \cap \Delta_s \neq \emptyset$ , then  $|\Delta_s| \leq \frac{1}{2} |G_A|$ . In particular  $F(z) \cap \Delta_s = \emptyset$  for  $s \geq 2$  and so  $|F(z) \cap \Delta_1| = 7$ . Since  $G_A/K_i \cong Z_6$  and  $z \notin K_i$ ,  $C_{G_A}(z) \neq \langle z \rangle$ . Hence an element of  $C_{G_A}(z)$  of order 3 acts on  $F(z) \cap \Delta_1$  and fixes at least one point on it. It follows that  $|\Delta_1| \leq \frac{1}{6} |G_A| = 3$ , a contradiction. Therefore we have  $K_i \cong Z_7$  and so  $G$  has a normal Sylow 7-subgroup by Lemma 3.3. Thus (ii) holds.

Let  $m (= U_i A)$ ,  $\Delta_s, x_i$  for  $t = i$  be as in Lemma 3.3 (ii). Since  $G_A \geq K_i \cong Z_7$  and  $K_i$  acts semi-regularly on  $m - \{U_i, A\}$ ,  $7 \mid |\Delta_s| = x_s$  for all  $s \in \{1, 2, 3, 4\}$ . Moreover, by Lemma 3.3,  $x_1 + x_2 + x_3 + x_4 = 63$ . Hence  $(x_1, x_2, x_3, x_4) = (7, 7, 7, 42)$   $(7, 14, 21, 21)$  or  $(14, 14, 14, 21)$  and so  $|G_A| = 42$ . Thus  $G_A/K_i \cong Z_6$  by the similar argument as in the proof of Lemma 3.3 (iii). Let  $y$  be an element of  $C_{G_A}(K_i)$  and assume that the order of  $y$  is 2 or 3. Since  $G_A/K_i \cong Z_6$  and  $K_i \cong Z_7$ ,  $y$  is contained in the center of  $G_A$ . Hence  $G_A$  acts on  $F(y)$  and therefore  $\Delta_s$  is contained in  $F(y)$  for each  $s$  with  $|\langle y \rangle| \nmid x_s$ . As above,  $(x_1, x_2, x_3, x_4) = (7, 7, 7, 42)$ ,  $(7, 14, 21, 21)$  or  $(14, 14, 14, 21)$  and hence  $|F(y) \cap m| \geq 21 + 2 = 23$ . Since  $F(y) \cap \Phi(\pi) \neq \emptyset$ ,  $y$  is a planar collineation. Therefore  $y = 1$ , a contradiction. Thus  $C_{G_A}(K_i) = K_i$ .

**Lemma 3.5.** Suppose  $u = 4$  and let notations be as in Lemma 3.4. Then, for some  $s \in \{1, 2, 3\} - \{i\}$   $O(G)$  has no orbit of length 7 on  $l_s$ .

Proof. Suppose false. Let  $P$  be a Sylow 7-subgroup of  $G$ . By Lemma 3.4 (ii),  $|P| = 7^2$  and  $P$  is a normal subgroup of  $G$ . Let  $s \in \{1, 2, 3\} - \{i\}$  and let  $\Omega_1$  be a  $P$ -orbit of length 7 on  $l_s$ . Then there exists another  $P$ -orbit of length 7, say  $\Omega_2$ , on  $l_s$  because  $7^2 \nmid |\bar{l}_s - \Omega_1|$ .

Let  $Q$  be a Sylow 3-subgroup of  $O(G)$ . By Lemmas 3.3 and 3.4,  $K_i \cong Z_7$  and a Sylow 3-subgroup of  $G/K_i$  is isomorphic to that of a Sylow 3-subgroup of  $\Gamma L(1, 2^6)$ . Hence  $Q = \langle a, b \mid a^9 = b^3 = 1, b^{-1}ab = a^4 \rangle$  for suitable  $a, b$  in  $Q$ . We note that  $Q' = [Q, Q] = \langle a^3 \rangle$ .

Since  $|\Omega_1| = |\Omega_2| = 7 < 9$ ,  $a^3$  acts trivially on  $\Omega_1 \cup \Omega_2$ , hence  $|F(a^3) \cap l_s| \geq 2 + |\Omega_1| + |\Omega_2| = 16$ . As  $s (\in \{1, 2, 3\} - \{i\})$  is arbitrary,  $a^3$  is planar and moreover we have  $F(a^3) = \pi$ , by Theorem 3.7 of [6], which implies that  $a^3 = 1$ . This

is a contradiction. Thus we have the lemma.

**Lemma 3.6.**  $u=3$ .

*Proof.* Assume that  $u \neq 3$ . Then, by Lemma 3.2,  $u=4$  and we can apply Lemmas 3.4 and 3.5. Let notations be as in them.

Let  $P$  be a Sylow 7-subgroup of  $G$  and  $\Gamma$  the set of  $P$ -orbits on  $\bar{l}_s$ . Set  $H=O(G)$ . Since  $P$  is a normal subgroup of  $H$  by Lemma 3.4 (ii),  $H$  induces a permutation group on  $\Gamma$ . Since  $P \geq K_i$  and  $K_i$  is semi-regular on  $\bar{l}_s$ , every  $P$ -orbit in  $\Gamma$  has length 7 or  $7^2$ . If an orbit in  $\Gamma$  has length  $7^2$ ,  $\Gamma$  contains exactly two  $P$ -orbits of length 7, which are also  $H$ -orbits of length 7, contrary to Lemma 3.5. Therefore each  $P$ -orbit in  $\Gamma$  has length 7 and so  $|\Gamma|=9$ .

If  $H$  acts transitively on  $\Gamma$ ,  $G$  is transitive on  $\bar{l}_s$  and therefore  $G/K_s \leq \Gamma L(1, 2^6)$  by Lemma 3.3 (i). It follows that  $\Gamma L(1, 2^6) \geq G_A K_s / K_s \cong G_A / G_A \cap K_s \cong G_A$ . Therefore an involution in  $G_A$  centralizes a Sylow 7-subgroup of  $G_A$  by the structure of  $\Gamma L(1, 2^6)$ , contrary to Lemma 3.4 (iii). Hence  $H$  is not transitive on  $\Gamma$ .

Let  $Q$  be a Sylow 3-subgroup of  $H$ . Then  $|Q|=27$  and  $[Q, Q]=Q' \cong Z_3$  as in the proof of Lemma 3.5. Since  $H=PQ$ ,  $\Gamma^H=\Gamma^Q$ . On the other hand  $H$  is not transitive on  $\Gamma$ . Hence  $Q^\Gamma$  is abelian and therefore  $Q'$  acts trivially on  $\Gamma$ . We note that  $G/C_G(P) \leq Z_6$  or  $G/C_G(P) \leq GL(2, 7)$  according as  $P \cong Z_{49}$  or  $Z_7 \times Z_7$ , respectively. Hence  $Q'$  is contained in  $C_G(P)$ . Since  $Q'$  acts trivially on  $\Gamma$  and each orbit  $\Delta \in \Gamma$  is of length 7,  $F(Q') \cap \Delta \neq \emptyset$ . Therefore  $Q' \leq K_s$  because  $[P, Q']=1$ . In particular  $Q'$  is semi-regular on  $\bar{l}_j$ , where  $\{j\} = \{1, 2, 3\} - \{i, s\}$ . Hence  $QK_i$  is transitive on  $\bar{l}_j$ . By Lemma 3.3 (i),  $G/K_j \cong \Gamma L(1, 2^6)$  and  $K_j \cong Z_7$ . Let  $z$  be an involution in  $G_A$ . Then  $[z, P] \leq K_i \cap K_j = 1$  and so  $z \in C_{G_A}(K_i)$ , contrary to Lemma 3.4 (iii). Thus we have  $u=3$ .

**Lemma 3.7.** Assume that there exists a line  $l$  through  $U_i$  with  $l \neq l_j, l_k$ , where  $\{i, j, k\} = \{1, 2, 3\}$ , such that  $G_l$  acts transitively on  $l - \{U_i, l \cap l_i\}$ . Then the following hold.

- (i)  $G_l$  is transitive on  $\bar{l}_i$  for  $t=j, k$ .
- (ii)  $G$  has two or three orbits on  $\bar{l}_i$ .

*Proof.* Let  $A_1, A_2 \in \bar{l}_j$  and set  $B_1 = U_j A_1 \cap l$  and  $B_2 = U_j A_2 \cap l$ . By assumption, there exists an element  $x \in G_l$  such that  $B_1^x = B_2$ . Since  $U_j A_2 \cap l = B_2 = B_1^x = U_j A_1^x \cap l$  and  $A_2, A_1^x \in \bar{l}_j$ , it follows that  $A_1^x = A_2$ . Hence  $G_l$  is transitive on  $\bar{l}_j$ . Similarly  $G_l$  is transitive on  $\bar{l}_k$ . Thus (i) holds.

Let  $d$  be the number of  $G$ -orbits on  $\bar{l}_i$ . Clearly  $d$  is at most 3. If  $d=1$ ,  $G$  acts transitively on  $\Phi(\pi)$ , contrary to  $u=3$ . Thus (ii) holds.

**Lemma 3.8.** Let  $l$  be the line satisfying the assumption in Lemma 3.7. If  $7^2 \mid |G|$  and  $7^3 \nmid |G|$ , then  $K_i \cong Z_3$  and  $|G| \mid 2 \cdot 3^2 \cdot 7^2$ .

Proof. By Lemmas 3.2, 3.3 (i) and 3.7 (i),  $K_j$  and  $K_k$  are isomorphic to  $Z_7$ ; otherwise  $7^2 \nmid |G|$ . Set  $A = I \cap I_i$ . Then  $G_i = G_A$  and so  $G_i/K_i = G_A/K_i$ . Since  $G/K_i \leq GL(6, 2)$ ,  $G_i/K_i$  is isomorphic to a subgroup of  $L$ , where

$$L = \left\{ \begin{bmatrix} 1 & a_2 \cdots a_6 \\ 0 & \\ \vdots & M \\ 0 & \end{bmatrix} \mid a_2, \dots, a_6 \in GF(2), M \in GL(5, 2) \right\}.$$

Since  $L/O_2(L) \cong GL(5, 2)$ , a Sylow 3-subgroup of  $L$  is an elementary abelian group of order 9. On the other hand, by Lemmas 3.3 and 3.7 (i),  $G_i$  contains an element of order 9. Therefore  $K_i \cong Z_3$ .

For a subgroup  $X$  of  $G$ ,  $\bar{X}$  denotes the homomorphic image of  $X$  in  $G/K_i$ . Since  $\bar{K}_j \neq \bar{K}_k$  and  $\bar{G} \leq GL(6, 2)$ ,  $\bar{K}_j \times \bar{K}_k$  is a Sylow 7-subgroup of  $\bar{G}$  and so  $\bar{K}_j \times \bar{K}_k$  has two subgroups  $\langle \bar{a} \rangle$  and  $\langle \bar{b} \rangle$  of order 7 which fix nonzero vectors on  $\bar{I}_i$ . Set  $H = O(G)$ . By Lemmas 3.3 (i) and 3.7 (i),  $G/K_i \leq \Gamma L(1, 2^6)$  for  $t \in \{j, k\}$ , so that  $|G:H| \leq 2$ . Since  $\bar{G} \triangleleft \bar{K}_i$  for  $t \in \{j, k\}$ ,  $H$  normalizes  $\bar{K}_j$ ,  $\bar{K}_k$ ,  $\langle \bar{a} \rangle$  and  $\langle \bar{b} \rangle$ . As  $K_i$  acts semi-regularly on  $\bar{I}_i$ , we have  $\bar{K}_i \neq \langle \bar{a} \rangle, \langle \bar{b} \rangle$  for  $t \in \{j, k\}$ . Without loss of generality, we can assume that  $\langle \bar{a}\bar{b} \rangle = \bar{K}_j$ . Let  $\bar{g} \in \bar{H}$ . Then  $\bar{g}^{-1}\bar{a}\bar{g} = \bar{a}^p$  and  $\bar{g}^{-1}\bar{b}\bar{g} = \bar{b}^q$  for some  $p, q$  with  $1 \leq p, q \leq 6$ , so we have  $\bar{g}^{-1}\bar{a}\bar{b}\bar{g} = \bar{a}^p\bar{b}^q \in \bar{K}_j = \langle \bar{a}\bar{b} \rangle$ . Hence  $p = q$ . From this,  $\bar{H}/C_{\bar{H}}(\langle \bar{a} \rangle \times \langle \bar{b} \rangle) \leq O(\text{Aut}(Z_7)) \cong Z_3$ . Since  $C_{GL(6, 2)}(\langle \bar{a} \rangle \times \langle \bar{b} \rangle) = \langle \bar{a} \rangle \times \langle \bar{b} \rangle$ , we have  $|\bar{H}| \mid 3|\langle \bar{a} \rangle \times \langle \bar{b} \rangle| = 3 \cdot 7^2$  and therefore  $|H| \mid 3^2 \cdot 7^2$ . Thus we obtain  $|G| \mid 2 \cdot 3^2 \cdot 7^2$ .

**Lemma 3.9.** *Let  $i \in \{1, 2, 3\}$  and set  $\{i, j, k\} = \{1, 2, 3\}$ . Then the following hold.*

- (i) *For every line  $m \in [U_i] - \{I_j, I_k\}$ ,  $G_m$  has three orbits on  $m - \{U_i, m \cap I_i\}$ .*
- (ii)  *$G$  acts transitively on  $\bar{I}_i$  and  $G/K_i \leq \Gamma L(1, 2^6)$ .*

Proof. Deny (i). Then, since  $u = u(\pi) = 3$ , there exists a line  $I \in [U_i]$  satisfying the assumption of Lemma 3.7. Let  $\{\Omega_1, \Omega_2, \dots, \Omega_p\}$  be the set of  $G$ -orbits on  $\bar{I}_i$  and set  $b_s = |\Omega_s|$  for  $1 \leq s \leq p$ . By Lemma 3.7 (ii),  $p = 2$  or  $3$ .

Assume  $p = 3$ . Set  $b = \max\{b_1, b_2, b_3\}$ ,  $b = |\Omega_p|$  and let  $A \in \Omega_p$ . Since  $u = 3$ ,  $G_A$  is transitive on  $m - \{U_i, A\}$ , where  $m = AU_i$ . Therefore  $63 \mid |G_A|$ . Hence  $63b \mid |G|$  because  $|G| = b|G_A|$ . By Lemmas 3.2, 3.3 (i) and 3.7 (i), we have  $|G| \mid 2 \cdot 3^4 \cdot 7^2$  and so  $b \mid 2 \cdot 3^2 \cdot 7$ . Since  $3b \geq b_1 + b_2 + b_3 = 63$ , it follows that  $21 \leq b < 63$ , hence  $b = 21$  or  $42$  and  $3^3 \cdot 7^2 \mid |G|$ , contrary to Lemma 3.8. Thus  $p \neq 3$ .

Assume  $p = 2$ . Let  $A \in \Omega_1$ ,  $B \in \Omega_2$  and set  $g = AU_i$ ,  $h = BU_i$ . Since  $u = 3$ , without loss of generality we may assume that  $G_A$  is transitive on  $g - \{U_i, A\}$  and that  $G_B$  has two orbits on  $h - \{U_i, B\}$ , say  $\Gamma_1, \Gamma_2$ . Similarly as in the last paragraph we obtain the following:

$$b_1, b_2 \mid |G|, |G| \mid 2 \cdot 3^4 \cdot 7^2, b_1 + b_2 = 63.$$

Hence  $\{b_1, b_2\} = \{21, 42\}$ ,  $\{14, 49\}$  or  $\{9, 54\}$ . We note that  $|G: G_g| = |G: G_A| = b_1$ ,  $|G: G_h| = |G: G_B| = b_2$  and  $63 \mid |G_A|$ .

If  $\{b_1, b_2\} = \{21, 42\}$ ,  $|G| = |G_A|b_1$  and  $21 \mid b_1$ . Hence  $3^3 \cdot 7^2 \mid |G|$ , contrary to Lemma 3.8.

If  $\{b_1, b_2\} = \{14, 49\}$ ,  $|G: G_A| = 14$  because  $7^3 \nmid |G|$ . Hence  $|G: G_h| = 49$ . By Lemma 3.8,  $|G| \mid 2 \cdot 3^2 \cdot 7^2$ . Therefore  $|G_h| \mid 18$ . Since  $h - \{U_i, B\}$  is a union of  $G_h$ -orbits  $\Gamma_1, \Gamma_2$ , we have  $|\Gamma_1| + |\Gamma_2| = 63$  and  $|\Gamma_1|, |\Gamma_2| \mid 18$ . This is a contradiction.

If  $\{b_1, b_2\} = \{9, 54\}$ , we have  $|G: G_A| = 9$  as  $3^5 \nmid |G|$ . Hence  $3^4 \mid |G|$  and so  $7^2 \nmid |G|$  by Lemma 3.8. Therefore  $|G| = 2 \cdot 3^4 \cdot 7$ . From this,  $|G_h| = |G_B| = 21$ . Hence  $|\Gamma_1|, |\Gamma_2| \mid 21$ . However,  $|\Gamma_1| + |\Gamma_2| = 63$ , a contradiction. Thus we have (i), and (ii) follows immediately from (i).

By Lemma 3.9, we can apply Lemma 3.3 for every  $t \in \{1, 2, 3\}$  and obtain the following.

**Lemma 3.10.** *Let notations be as in Lemma 3.3. Then the following hold.*

(i)  $3^2 \cdot 7 \mid |G|$ ,  $|G| \mid 2 \cdot 3^4 \cdot 7^2$  and  $3^4 \cdot 7^2 \nmid |G|$ .

(ii)  $3^3, 7^2 \nmid x_s$  for all  $s \in \{1, 2, 3\}$ .

*Proof.* By Lemmas 3.2 and 3.3 (i) (ii), we have (i). By Lemma 3.3 (ii) (iii),  $|G_A| = |G|/63$  and  $x_s \mid |G_A|$ . Hence  $x_s \mid 2 \cdot 3^2 \cdot 7$ . Thus we have (ii).

**Lemma 3.11.** *Let notations be as in Lemma 3.3 and assume that  $21 \mid x_2$ . Then the following hold.*

(i)  $K_1 \cong K_2 \cong K_3 \cong Z_7$  and  $G/K_t$  is isomorphic to a subgroup of  $\Gamma L(1, 2^6)$  of index at most 2 for each  $t \in \{1, 2, 3\}$ .

(ii) Let  $Q$  be a Sylow 3-subgroup of  $G$ . Then  $|Q| = 3^3$  and  $Q = \langle a, b \mid a^9 = b^3 = 1, b^{-1}ab = a^4 \rangle$  for suitable  $a, b$  in  $Q$ . Moreover, for any element  $v$  of order 3 in  $Q - Z(Q)$ ,  $F(v)$  is a subplane of order 4.

*Proof.* By Lemma 3.3,  $x_3$  divides  $|G_A|$  and  $|G: G_A| = 3^2 \cdot 7$ , so that  $3^3 \cdot 7^2 \mid |G|$ . It follows from Lemma 3.10 (i) that  $|G| = 3^3 \cdot 7^2$  or  $2 \cdot 3^3 \cdot 7^2$ . Therefore (i) holds.

By (i), the order of a Sylow 3-subgroup  $Q$  of  $G$  is  $3^3$ . Hence  $Q$  is of the form stated in (ii) by the structure of  $\Gamma L(1, 2^6)$ . We note that  $Q$  has exactly two conjugacy classes of subgroups of order 3. Let  $v \in Q - Z(Q)$  such that  $\langle v \rangle \cong Z_3$ . Then, as an element in  $\Gamma L(1, 2^6)$ ,  $v$  fixes three nonzero elements, that is,  $|F(v) \cap \bar{l}_t| = 3$  for all  $t \in \{1, 2, 3\}$ . Hence  $F(v)$  is a subplane of order 4.

**Lemma 3.12.** *Let notations be as in Lemma 3.3. Then  $(x_1, x_2, x_3) = (7, 14, 42)$ ,  $K_t \cong Z_7$  and  $G/K_t \cong \Gamma L(1, 2^6)$  for each  $t \in \{1, 2, 3\}$ .*

*Proof.* By Lemmas 3.3 (iii) and 3.10, we have  $x_1 \leq x_2 \leq x_3$ ,  $x_1 + x_2 + x_3 = 63$



and  $3^3, 7^2 \nmid x_s, x_s \mid |G| \mid 2 \cdot 3^4 \cdot 7^2$  for  $s \in \{1, 2, 3\}$ . Hence  $(x_1, x_2, x_3) = (21, 21, 21)$  or  $(7, 14, 42)$ . On the other hand  $K_1 \simeq K_2 \simeq K_3 \simeq Z_7$  by Lemma 3.11 (i).

Assume that  $(x_1, x_2, x_3) = (21, 21, 21)$ . Let  $\Delta_s$  be as defined in Lemma 3.3. Let  $P_s \in \Delta_s$  and let  $\Phi_s$  be the  $G$ -orbit containing  $P_s$  for  $s \in \{-1, 2, 3\}$ . Clearly  $|\Phi_s| = 63x_s$ . Let  $v$  be the element as defined in Lemma 3.11 (ii) and let  $P \in F(v) \cap \Phi$ . Then  $P \in \Phi_s$  for some  $s \in \{1, 2, 3\}$ . Therefore  $|G_P| = |G|/|\Phi_s| \mid 2 \cdot 3^3 \cdot 7^2 / 63 \cdot 21 = 2$ , contrary to  $v \in G_P$ . Thus  $(x_1, x_2, x_3) = (7, 14, 42)$  and so  $G/K_t \simeq \Gamma L(1, 2^6)$  for all  $t \in \{1, 2, 3\}$ .

**Lemma 3.13.** *Let  $\Delta_1$  be as in Lemma 3.3. Then the following hold.*

(i) *Let  $P \in \Delta_1$ . Then  $G_P = \langle x \rangle \simeq Z_6$  and a Sylow 7-subgroup of  $G$  acts on  $F(x^3) \cap \bar{L}_t$  for all  $t \in \{1, 2, 3\}$ .*

(ii)  *$F(x^c)$  is a subplane of  $\pi$  of order  $2^c$  for  $c = 2, 3$ .*

Proof. Similarly as in the proof of Lemma 3.12, we obtain  $|G_P| = |G|/|P^G| = 2 \cdot 3^3 \cdot 7^2 / 63 \cdot 7 = 6$ . Since  $G_P \leq G_A$ ,  $G_P \cap K_t = 1$  and  $G_A/K_t \simeq Z_6$ , we have  $G_P \simeq G_P K_t / K_t \leq Z_6$ . Hence  $G_P \simeq Z_6$ . Set  $\langle x \rangle = G_P$ . Clearly  $x^3$  is an involution in  $G_P$  and so by the property of  $\Gamma L(1, 2^6)$ ,  $x^3$  centralizes the Sylow 7-subgroup of  $G/K_t$  for all  $t \in \{1, 2, 3\}$ . Let  $S$  be the Sylow 7-subgroup of  $G$ . Then  $[x^3, S] \leq \bigcap_{t=1}^3 K_t = 1$  and therefore  $S$  centralizes  $x^3$ . Hence  $S$  acts on  $F(x^3) \cap \bar{L}_t$  for all  $t \in \{1, 2, 3\}$ . Thus (i) holds.

By Theorem 4.3 of [6],  $F(x^3)$  is a subplane of  $\pi$  of order  $2^3$  and by Lemmas 3.11 (ii) and 3.12,  $F(x^2)$  is a subplane of order  $2^2$ .

If we coordinatize  $\pi$  by choosing  $(0, 0)$  as  $U_1$ ,  $(0)$  as  $U_2$ ,  $(\infty)$  as  $U_3$ ,  $(1, 1)$  as  $P$  which was defined in Lemma 3.13, then we get a semi-field  $F$ . In general,  $F$  is not always isomorphic to  $D$  and since  $\pi$  is non-Desarguesian,  $F$  is not a field. Thus  $\pi$  is a semi-field plane coordinatized by  $F$  and it also satisfies Hypothesis 2.3.

**Lemma 3.14.** *Set  $F_1 = \{d \mid d \in F, (d, 0) \in F(x^3)\}$ ,  $F_2 = \{d \mid d \in F, (0, d) \in F(x^3)\}$  and  $F_3 = \{d \mid d \in F, (d, 0) \in F(x^2)\}$ . Then  $F_1 = F_2 \simeq GF(8)$  and  $F_3 \simeq GF(4)$ .*

Proof. Since  $F(x)$  contains  $(0, 0)$ ,  $(0)$ ,  $(\infty)$  and  $(1, 1)$ , it also contains  $(1)$ . By Lemma 3.13 and the definition of the coordinatization of  $\pi$ , we have the lemma.

**Lemma 3.15.** *Let  $N_1$ ,  $N_2$  or  $N_3$  be the right, middle or left nucleus, respectively. Then  $N_1 = N_2 = N_3 \simeq GF(8)$ .*

Proof. By Lemmas 3.12, we have  $N_t \simeq GF(8)$  for all  $t \in \{1, 2, 3\}$ . Furthermore, the multiplicative group  $N_t^\times = \{d \mid (d, 0) \in (1, 0)^{K_t}\}$  for  $t = 1, 2$  and  $N_3^\times = \{d \mid (0, d) \in (0, 1)^{K_3}\}$  by the proof of Theorems 7.9 and 8.2 of [6]. Since

$K_1$  and  $K_2$  are semi-regular on  $\bar{I}_3$  and  $(1, 0) \in F(x^3)$ , it follows from Lemma 3.13 that  $N_1 = N_2 = F_1$ . Similarly  $N_3 = F_2$ . By Lemma 3.14, we have  $N_1 = N_2 = N_3 \simeq GF(8)$ .

**Lemma 3.16.** Set  $N = N_1 = N_2 = N_3$  and  $F_3^\times = \langle \theta \rangle$ .

(i)  $N$  does not contain  $\theta$  and  $F$  is a right and left vector space over  $N$  with a basis  $\{1, \theta\}$ .

(ii) For any  $\xi \in F$ ,  $(\xi\theta)\theta = \xi(\theta^2)$ .

Proof. (i) follows immediately from Lemmas 3.14 and 3.15.

Set  $\xi = a + b\theta$  for  $a, b \in N$ . Then  $(\xi\theta)\theta = ((a + b\theta)\theta)\theta = (a\theta + (b\theta)\theta)\theta = (a\theta)\theta + ((b\theta)\theta)\theta = a\theta^2 + (b\theta^2)\theta = a\theta^2 + b\theta^3$  because  $a, b \in N = N_3$  and  $\langle \theta \rangle = F_3^\times$ . Hence  $(\xi\theta)\theta = a\theta^2 + (b\theta)\theta^2 = \xi(\theta^2)$ . Thus (ii) holds.

**Lemma 3.17.**  $\theta \in N$ .

Proof. Let  $\xi, \eta \in F$  and set  $\xi = a + b\theta$ ,  $\eta = c + d\theta$  for  $a, b, c, d \in N$ . Then,  $(\xi\eta)\theta = ((a + b\theta)(c + d\theta))\theta = (ac)\theta + ((b\theta)c)\theta + (a(d\theta))\theta + ((b\theta)(d\theta))\theta$ . Similarly  $\xi(\eta\theta) = a(c\theta) + (b\theta)(c\theta) + a((d\theta)\theta) + (b\theta)((d\theta)\theta)$ . Since  $a \in N = N_3$  and  $c \in N = N_2$ , we have  $(ac)\theta = a(c\theta)$ ,  $((b\theta)c)\theta = (b\theta)(c\theta)$  and  $(a(d\theta))\theta = a((d\theta)\theta)$ . Since  $d \in N = N_2$ ,  $((b\theta)(d\theta))\theta = (((b\theta)d)\theta)\theta$  and by Lemma 3.16,  $((b\theta)d)\theta = (b\theta)(d\theta)^2$ , so that  $((b\theta)(d\theta))\theta = ((b\theta)d)\theta^2 = (b\theta)(d\theta^2) = (b\theta)((d\theta)\theta)$  as  $d \in N = N_2 = N_3$ . Hence  $(\xi\eta)\theta = \xi(\eta\theta)$  and so  $\theta \in N_3 = N$ .

Proof of Lemma 2.

By Lemmas 3.16 (i) and 3.17, we obtain a contradiction and so the lemma holds.

#### 4. The proof of Theorem 1

Throughout this section  $\pi$  is a semi-field plane satisfying Hypothesis 2.3 and the following.

**Hypothesis 4.1.**  $r \not\equiv 0 \pmod{4}$  and  $u(\pi) \leq 4$ .

**Lemma 4.2.** (i)  $G$  is not solvable.

(ii)  $u(\pi) = 2, 3$  or  $4$ .

(iii) There exists  $i \in \{1, 2, 3\}$  such that  $G$  is transitive on  $\bar{I}_i$ .

Proof. By Theorem of [8], Theorem 6.3 of [9] and the lemma proved in §3, we have (i).

It follows from Kallaher's theorem [8] that  $u(\pi) \neq 1$  and so (ii) holds.

If  $u(\pi) = 2$  or  $3$ , we have (iii) by a similar argument as in the proof of Lemma 3.7. If  $u = 4$ , we can apply Lemma 6.1 of [9] and (iii) follows.

**Lemma 4.3.** *Let  $S$  be a Sylow 2-subgroup of  $G$  and set  $\pi_0 = F(S)$ ,  $H = G(\pi_0)$ ,  $\bar{G} = G/O(G)$ . Then the following hold.*

- (i)  $S \neq 1$  and  $S$  is semi-regular on  $\pi - \pi_0$ .
- (ii)  $\pi_0$  is a Baer subplane of  $\pi$ .
- (iii)  $\bar{G}' \simeq PSL(2, q)$  for some even  $q$ . Moreover  $H = O(G)N_c(S)$  and  $|G:H| = q+1$ .

*Proof.* By the Feit-Thompson theorem [2] and Lemma 4.1 (i), the order of  $G$  is even and so  $S \neq 1$ . Let  $z$  be an involution in the center of  $S$ . Then  $F(z)$  is a Baer subplane of order  $2^{r/2}$  and  $S^{F(z)}$  is a collineation group. By Hypothesis 4.1,  $2^{r/4}$  is not an integer. Therefore  $S^{F(z)} = 1$ . Hence (i) and (ii) hold.

By Lemma 4.2 (ii),  $G \neq H$  and clearly  $H \geq S$ . Hence  $H$  is a strongly embedded subgroup of  $G$ . By a Bender's theorem [1] and by Corollary 3.2 of [4], (iii) holds.

**Lemma 4.4.** *Set  $\Delta = \pi_0 \cap \bar{l}_i$  and  $\Gamma = \{\Delta^g \mid g \in G\}$ . Then the following hold.*

- (i)  $\bar{l}_i = \bigcup_{\Delta^g \in \Gamma} \Delta^g$  and  $\Delta^x \cap \Delta^y = \phi$  for distinct  $\Delta^x$  and  $\Delta^y$  in  $\Gamma$ .
- (ii) Set  $N = O(G)$ . Then  $G(\Delta) = H \geq N = G_\Gamma$  and  $G$  is doubly transitive on  $\Gamma$ .

*Proof.* By Lemma 4.3 (iii),  $H = N \cdot N_c(S)$ . Since  $G(\pi_0) \leq G(\Delta)$  and  $H$  is a maximal subgroup of  $G$ , we have  $H = G(\Delta)$ . Hence  $G$  is doubly transitive on  $\Gamma$  (See [1] §3). Since  $N$  is a normal subgroup of  $G$  and  $N \leq G(\Delta)$ ,  $N$  is contained in  $G_\Gamma$  and so  $N = G_\Gamma$  by Lemma 4.3 (iv). Thus (ii) holds.

Clearly  $\Delta^g \subset \bar{l}_i$  for all  $g \in G$ , hence  $\bar{l}_i = \bigcup_{\Delta^g \in \Gamma} \Delta^g$  by Lemma 4.2 (iii). Suppose  $\Delta^x \neq \Delta^y$  and  $\Delta^x \cap \Delta^y \neq \phi$  and set  $g = xy^{-1}$ . Then  $\Delta^g \neq \Delta$  and  $\Delta^g \cap \Delta \neq \phi$ . By Lemma 4.3 (i),  $S$  and  $S^g$  fix  $\Delta^g \cap \Delta$  pointwise. By (ii),  $G = \langle N, S, S^g, G(\Delta) \cap G(\Delta^g) \rangle$ . Hence  $G$  fixes  $\Delta^g \cap \Delta$  as a set, contrary to Lemma 4.2 (iii). Thus (i) holds.

**Lemma 4.5.**  $q^2 = 2^r$  and  $|\Delta| = q-1$ ,  $|\Gamma| = q+1$ .

*Proof.* By Lemmas 4.3 (iii) and 4.4 (ii),  $|\Gamma| = |G:H| = q+1$  and by Lemma 4.4 (i)  $|\Gamma| = |\bar{l}_i|/|\Delta| = (2^r - 1)/|\Delta|$ . On the other hand  $|\Delta| = 2^{r/2} - 1$  since  $\pi_0$  is a Baer subplane of  $\pi$ . Hence  $q^2 = 2^r$  and  $|\Delta| = q-1$ .

**Lemma 4.6.**  $\pi_0 (= F(S))$  is a Desarguesian projective plane of order  $q$  and the number of  $N_c(S)$ -orbits on  $\Phi(\pi) \cap \pi_0$  is one or three.

*Proof.* Let  $\Lambda$  be a  $G$ -orbit on  $\Phi(\pi)$  and suppose  $\Lambda \cap \pi_0 \neq \phi$ . Let  $P \in \Lambda \cap \pi_0$ . Then  $G_P \geq S$ . Hence  $|\Lambda| = |G:G_P| \equiv 1 \pmod{2}$  and moreover  $N_c(S)$  is transitive on  $\Lambda \cap \pi_0$  by Theorem 3.5 of [11]. Since  $|\Phi| \equiv 1 \pmod{2}$

and  $u=u(\pi) \leq 4$ , the number of  $G$ -orbits  $\Lambda$  on  $\Phi$  such that  $\Lambda \cap \pi_0 \neq \emptyset$  is one or three. Hence the number of  $N_G(S)$ -orbits on  $\pi_0 \cap \Phi$  is one or three.

Since the order of  $\pi_0$  is  $2^{r/2}$  and  $2^{r/4}$  is not an integer, the autotopism group of  $\pi_0$  is of odd order. By Theorem 6.3 of [9] and Theorem of [8],  $\pi_0$  is a Desarguesian plane of order  $q$ .

By Lemma 4.3,  $|G:G(\pi_0)|=q+1$ . We set  $\{\pi_0^g \mid g \in G\} = \{\pi_0, \pi_1, \dots, \pi_q\}$ . Then the following lemma holds.

**Lemma 4.7.** *Set  $N=O(G)$ . Then*

- (i)  $N_{\pi_s}$  acts faithfully on  $\pi_t$  and  $|N_{\pi_s}| \mid (q-1)^2(r/2)$  for all  $s, t$  ( $s \neq t$ ) and
- (ii)  $N_{\pi_t}$  is a normal subgroup of  $N$  and  $[N_{\pi_s}, N_{\pi_t}] = 1$  for all  $s, t$  ( $s \neq t$ ).

Proof. By Lemma 4.4 (ii),  $N$  acts on  $\pi_t$  and so  $N_{\pi_t}$  is a normal subgroup of  $N$ . By Lemma 4.3 (ii),  $\pi_s$  and  $\pi_t$  are Baer subplanes of  $\pi$ , so that  $N_{\pi_s} \cap N_{\pi_t} = 1$ . Hence  $N_{\pi_s}$  acts faithfully on  $\pi_t$  and  $[N_{\pi_s}, N_{\pi_t}] \leq N_{\pi_s} \cap N_{\pi_t} = 1$ . Moreover  $|N_{\pi_s}| \mid (q-1)^2(r/2)$  since  $\pi_t$  is a Desarguesian plane of order  $q$ .

**Lemma 4.8.** *Assume  $N_{\pi_0} \neq 1$  and let  $P$  be a minimal normal subgroup of  $N_{\pi_0}$  and let  $p$  be a prime dividing the order of  $P$ . Then a Sylow  $p$ -subgroup of  $N_{\pi_0}$  is cyclic and  $P$  is a normal subgroup of  $N$ . Moreover  $P$  is isomorphic to  $Z_p$ .*

Proof. Let  $Q$  be a Sylow  $p$ -subgroup of  $N_{\pi_0}$ . Since  $F(Q) = \pi_0$ ,  $Q$  is semi-regular on  $\pi_t - \pi_0$  for  $t \neq 0$ . By Lemma 2.2 (i) and Theorem 5.4.10 of [3],  $Q$  is cyclic. Hence, by Lemma 4.7 (ii), we have the lemma.

**Lemma 4.9.** *Let  $P$  be as in Lemma 4.8. Then the following hold.*

- (i) Set  $L = \langle P^g \mid g \in G \rangle$ . Then  $L$  is a normal subgroup of  $G$  and is an elementary abelian  $p$ -group.
- (ii)  $p \nmid r$  and  $|L| \leq p^3$ .

Proof. (i) follows immediately from Lemma 4.8. Clearly  $L \leq N$ . Set  $X = N_{\pi_0}$ . Since  $X \cap L = P$  and  $L/P \cong LX/X \leq N/X \cong N^{\pi_0}$ ,  $|L/P|$  is at most  $p^3$ . Moreover  $|L/P| \leq p^2$  if  $p \nmid r$ . Therefore it suffices to show  $p \nmid r$ . Assume  $p \mid r$ . Since  $H$  normalizes  $X$ ,  $P$  is a normal subgroup of  $H$  and so  $L$  contains at least  $q+1$  subgroups of order  $p$ . Hence  $q+1 \leq (p^4-1)/(p-1) = p^3+p^2+p+1$ . On the other hand  $p \mid r/2$  and  $q=2^{r/2}$ , so that  $(r/2)^3 + (r/2)^2 + r/2 + 1 \geq 2^{r/2} + 1$ . From this  $r=6$  or  $10$  and  $p=r/2$ . But  $p \nmid q-1$  for  $r=6$  or  $10$ . Therefore,  $|L/P| \leq p$  and so  $q+1 \leq (p^2-1)/(p-1) = p+1 \leq 6$ , a contradiction. Thus  $p \nmid r/2$ .

**Lemma 4.10.**  $N_{\pi_0} = 1$ .

Proof. Assume  $N_{\pi_0} \neq 1$  and let  $P, L$  be as in Lemma 4.8, 4.9, respectively.

If  $|C_G(L)|$  is even, all Sylow 2-subgroups of  $G$  are contained in  $C_G(L)$  by Lemma 4.3 (iii). Hence  $\langle S^g | g \in G \rangle$  acts on  $F(P)$  ( $=\pi_0$ ), which is contrary to  $G(\pi_0)=H$ . Therefore  $|C_G(L)|$  is odd. In particular  $S$  is isomorphic to a subgroup of  $G/C_G(L)$ .

By Lemmas 4.3 and 4.9,  $(G/C_G(L))' \leq SL(3, p)$  and  $|G/C_G(L) : (G/C_G(L))'|$  is odd. Hence  $S$  is isomorphic to a subgroup of  $SL(3, p)$ . Since a Sylow 2-subgroup of  $SL(3, p)$  is semi-dihedral or wreathed,  $S$  is an elementary abelian group of order 4 and so  $q=2^2$ . Hence  $r=4 \equiv 0 \pmod{4}$ , a contradiction.

**Lemma 4.11.** *Let  $G^{(\infty)}$  denote the last term of the derived series of  $G$ . Set  $M=G^{(\infty)}$ . Then  $M \simeq PSL(2, q)$ .*

*Proof.* Let  $X$  be a subgroup of  $G$  generated by all Sylow 2-subgroups of  $G$ . By Lemma 4.3 (iii),  $X \leq M$  and  $|M/X|$  is odd. It follows from the Feit-Thompson theorem that  $M=M^{(\infty)} \leq X$  and hence  $X=M$ . By Lemmas 4.4 (ii) and 4.10,  $[S, N] \leq N \cap G_{\pi_0} = N_{\pi_0} = 1$ , so that  $N$  centralizes  $X$  ( $=M$ ) and  $M \cap N = Z(M)$ ,  $M/Z(M) \simeq PSL(2, q)$ . By a property of  $PSL(2, q)$ ,  $M \simeq PSL(2, q)$ .

**Lemma 4.12.** (i) *Let  $t \in \{1, 2, 3\}$ ,  $P \in \bar{l}_t$  and let  $X$  be a subgroup of  $G_P$ . Then  $|F(X) \cap l_t| = 2^a + 1$  for an integer  $a \geq 1$ .*

(ii)  *$M$  is transitive on  $\bar{l}_i$  and  $|M_P| = q$  for  $P \in \bar{l}_i$ . Here  $i$  is the integer defined in Lemma 4.2 (iii).*

*Proof.* Let  $A$  be the full collineation group of  $\pi$  and set  $T_1 = A_{(U_3, l_2)}$ ,  $T_2 = A_{(U_3, l_1)}$ ,  $T_3 = A_{(U_2, l_1)}$ . Since  $U_3$  is a translation point and  $l_1$  is a translation line,  $T_1 \simeq T_2 \simeq T_3 \simeq E_{q^2}$  and  $XT_i$  is a transitive linear group on  $l_i$ . Since  $(XT_i)_P = X$ , we have (i) by Lemma 2.1.

Let  $\{\Delta_1, \dots, \Delta_m\}$  be the set of  $M$ -orbits on  $\bar{l}_i$ . Since  $G$  is transitive on  $\bar{l}_i$  and  $G \triangleright M$ ,  $|\Delta_1| = \dots = |\Delta_m| \equiv 1 \pmod{2}$ . Let  $P \in \Delta_1$  and set  $M_P = CS$  with  $C \leq Z_{q-1}$  and  $|N_M(S) : M_P| = k$ . As  $M \simeq PSL(2, q)$ ,  $k \mid q-1$  and  $F(M_P) \cap \Delta_v \neq \phi$  for each  $v \in \{1, \dots, m\}$ .

Assume  $C \neq 1$ . Then  $|N_M(C)| = 2(q-1)$  as  $M \simeq PSL(2, q)$ . By Lemma 2.1,  $|F(C) \cap \bar{l}_i| = m \times \frac{2(q-1) \times |S|}{|M_P|} = 2mk$  and applying (i), we have  $2mk = 2^a - 1$  for an integer  $a \geq 1$ , a contradiction. Thus  $C=1$  and  $|M_P| = q$ . Therefore  $|P^M| = |M : M_P| = q^2 - 1$  and (ii) follows.

**Lemma 4.13.** *Let  $j \in \{1, 2, 3\} - \{i\}$  and  $P \in \bar{l}_j$ . Then  $q \mid |M_P|$ .*

*Proof.* By Lemma 4.3 (i), it suffices to consider the case that  $|M_P| \equiv 1 \pmod{2}$ . As  $M \simeq PSL(2, q)$ ,  $M \leq Z_{q \pm 1}$ . Since  $|\bar{l}_j| = q^2 - 1 \geq |P^M| = |M : M_P|$  and  $P^M \cap F(S) = \phi$ , we have  $M_P \simeq Z_{q+1}$  and  $|l_j - P^M| = |F(S) \cap l_j| = q+1$ . Hence  $F(S) \cap l_j = F(M) \cap l_j$ . Therefore  $|F(M_P)| = q+1 + \frac{2(q+1) \times 1}{|M_P|} = q+3$  by Lem-

mma 2.1. Applying Lemma 4.12 (i),  $q+3=2^a+1$  for an integer  $a \geq 1$ . This is a contradiction.

**Lemma 4.14.**  *$M$  is transitive on  $\bar{l}_j$  and  $M_P$  is a Sylow 2-subgroup of  $M$  for each  $j \in \{1, 2, 3\}$  and  $P \in \bar{l}_j$ .*

Proof. By Lemma 4.12 (ii), we may assume  $j \in \{1, 2, 3\} - \{i\}$ . First we argue that  $F(M) \cap \bar{l}_j = \phi$ . Set  $\Delta = F(M) \cap \bar{l}_j$  and assume  $\Delta \neq \phi$ . Let  $\pi_0$  be as defined in Lemma 4.6 and set  $N_M(S) = DS$  with  $D \cong Z_{q-1}$ . By Lemma 4.12 (ii),  $D^{\pi_0} \cong D$  and  $\pi_0 \cap F(D) \supset \Delta$ . Since  $\pi_0$  is a Desarguesian plane of order  $q$ ,  $F(D) \supset \pi_0 \cap l_j$  by Lemma 2.2 (ii). Therefore, by Lemmas 2.1 and 4.13,  $|F(D) \cap \bar{l}_j| = |\Delta| + 2(q-1 - |\Delta|) = 2(q-1) - |\Delta|$ . Applying Lemma 4.12 (i),  $|\Delta| = 2^a - 1$  and  $2(q-1) - |\Delta| = 2^b - 1$  for integers  $a, b \geq 1$ , hence  $2q = 2^a + 2^b$ . However, as  $|\Delta| < |\pi_0 \cap \bar{l}_j| = q-1 < |F(D) \cap \bar{l}_j| = 2^b - 1$ , we have  $2^a < q < 2^b$ . This is a contradiction. Thus  $F(M) \cap \bar{l}_j = \phi$ .

Let  $\{\Delta_1, \dots, \Delta_m\}$  be the set of  $M$ -orbits on  $\bar{l}_j$ . By Lemma 4.3,  $|\Delta_t| \mid q^2 - 1$  for each  $t$ . Assume  $|M_P| \neq q$  for some  $P \in \bar{l}_j$ . We may assume  $P \in \Delta_1$  and  $M_P \triangleright S$ . Set  $M_P = CS$  with  $1 \neq C \leq Z_{q-1}$ . By a similar argument as in the last paragraph  $F(S) \cap l_j \subset F(C) \cap l_j$  and so  $F(C) \cap \Delta_t \neq \phi$  for each  $t$ . Hence  $|F(C) \cap \Delta_t| = 2 \times |F(S) \cap \Delta_t|$  by Lemma 2.1. Hence  $|F(C) \cap \bar{l}_j| = 2 \times |F(S) \cap \bar{l}_j| = 2(q-1)$  and so  $|F(C) \cap l_j| = 2q$ , contrary to Lemma 4.12 (i). Thus  $|M_P| = q$  and  $M$  is transitive on  $\bar{l}_j$ .

Let  $X$  be the full collineation group of  $\pi$  and set  $A = X_{(l_1, l_1)}$ ,  $B = X_{(U_3, U_3)}$  and  $T = AB$ . Since  $U_3$  is a translation point and  $l_1$  is a translation line,  $A$  and  $B$  are elementary abelian normal 2-subgroups of  $X$  of order  $q^4$ . Hence  $T$  is a normal 2-subgroup of  $X$ .

**Lemma 4.15.** (i)  *$T$  is a nonabelian normal 2-subgroup of  $X$ .*

(ii)  *$C_T(x) = 1$  for any element  $x (\neq 1)$  of  $M$  of odd order.*

Proof. If  $T$  is abelian,  $T_P = 1$  for  $P \notin l_1$  because  $A$  acts transitively on the set of points not incident with the line  $l_1$ . Hence  $|T| = |T : T_P| = q^4 + q^2 + 1 - (q^2 + 1) = q^4$  and so  $T = A = B$ , a contradiction. Thus (i) holds.

Assume  $C_T(x) \neq 1$  and let  $t$  be an involution in  $C_T(x)$ . Then, there exist element  $t_1 \in A$  and  $t_2 \in B$  such that  $t = t_1 t_2$ . By Lemma 4.14,  $F(X) = \{U_1, U_2, U_3, l_1, l_2, l_3\}$  and so  $t$  acts on  $\{U_1, U_2, U_3\}$ . Since  $F(X) = \{U_3, l_1\}$ , it follows that  $(U_2)^t \in l_1$  and  $(U_3)^t = U_3$ . Hence we have  $(U_1)^t = U_1$  and  $(U_2)^t = U_2$  and so  $F(t_2) = F(t_1 t) \supset \{U_2, U_3\}$ . Therefore  $t_2 \in X_{(U_3, l_1)} \leq X_{(l_1, l_1)}$ , which implies  $t \in X_{(l_1, l_1)}$ . However, as  $(U_1)^t = U_1$ , this is a contradiction. Thus  $C_T(x) = 1$ .

Proof of Theorem 1.

Since  $3 \mid |M| = |PSL(2, q)|$ , there exists an element  $x \in M$  of order 3.

By Lemma 4.15 (ii),  $C_T(x)=1$ . Applying Theorem 8.2 of [5] to the group  $MT$ ,  $T$  is an abelian 2-group, which is contrary to Lemma 4.15 (i). Thus we have the theorem.

---

### References

- [1] H. Bender: *Transitive Gruppen gerader Ordnung, in denen jede Involution genau einen Punkt festlässt*, J. Algebra **17** (1971), 527–554,
- [2] W. Feit and J. Thompson: *Solvability of groups of odd order*, Pacific J. Math. **13** (1963), 775–1029.
- [3] D. Gorenstein: *Finite groups*, Harper and Row, New York, 1968.
- [4] C. Hering: *On finite line transitive affine planes*, Geom. Dedicata **1** (1973), 387–393.
- [5] G. Higman: *Odd characterization of finite simple groups*, Lecture Notes, University of Michigan, 1968.
- [6] D.R. Hughes and F.C. Piper: *Projective planes*, Berlin-Heidelberg-New York, Springer-Verlag, 1973.
- [7] B. Huppert: *Zweifach transitive, auflösbare Permutationsgruppen*, Math. Z. **68** (1957), 126–150.
- [8] M.J. Kallaher: *A conjecture on semi-field planes*, Arch. Math. **26** (1975), 436–440.
- [9] M.J. Kallaher and R.A. Liebler: *A conjecture on semi-field planes II*, Geom. Dedicata **8** (1979), 13–30.
- [10] R. Steinberg: *Automorphisms of finite linear groups*, Canad. J. Math. **12** (1960), 606–615.
- [11] H. Wielandt: *Finite permutation groups*, Academic Press, New York, 1964.

Department of Mathematics  
 College of General Education  
 Osaka University  
 Toyonaka, Osaka 560, Japan