

Title	Correction to : "A construction for irregular discriminants"
Author(s)	Craig, Maurice
Citation	Osaka Journal of Mathematics. 15(2) P.461
Issue Date	1978
Text Version	publisher
URL	https://doi.org/10.18910/4838
DOI	10.18910/4838
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/repo/ouka/all/>

Craig, M.
Osaka J. Math.
15 (1978), 461

CORRECTION TO
"A CONSTRUCTION FOR IRREGULAR DISCRIMINANTS"

MAURICE CRAIG

This Journal, vol. 14 (1977), 365–402

(Received January 23, 1978)

- 365 –11 ...the ideal $(A_i, \frac{1}{2}(B_i + \sqrt{D}))$...
- 390 +11 ...once it is determined that the square-free part of $D(t)$ has degree at least three.
- 391 –12 ...we observe that $(A_i(t), B_i(t))$ divides R_i hence T . By (77), it must therefore divide $(A_i(3), B_i(3))$, which equals 1 by Part 7.
- 400 +8 $\nu=17555^2+462\cdot 389^2$

It may also be remarked that both (33) and the solution discussed in Note A can be obtained by the chord and tangent process applied to the intersections of the plane cubic curve (27) with the line $\lambda=0$.

