



Title	Quantum deformations of certain prehomogeneous vector spaces. II
Author(s)	森田, 良幸
Citation	Osaka Journal of Mathematics. 2000, 37(2), p. 385-403
Version Type	VoR
URL	https://doi.org/10.18910/4931
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

QUANTUM DEFORMATIONS OF CERTAIN PREHOMOGENEOUS VECTOR SPACES. II

YOSHIYUKI MORITA

(Received March 5, 1998)

Introduction

Let G be a reductive algebraic group over the complex number field \mathbb{C} and let \mathfrak{g} be its Lie algebra. The quantized coordinate algebra $A_q(G)$ of G is constructed as a certain dual Hopf algebra of the quantized enveloping algebra $U_q(\mathfrak{g})$ of \mathfrak{g} . The Hopf algebras $U_q(\mathfrak{g})$ and $A_q(G)$ over $\mathbb{C}(q)$ tend to the ordinary enveloping algebra $U(\mathfrak{g})$ and the coordinate algebra $A(G)$ respectively when the parameter q tends to 1 in a certain sense (Drinfeld [1], Jimbo [3]).

Let us consider what object we should regard as a quantum deformation of an affine variety X with G -action.

An affine variety X is endowed with an action of G if and only if its coordinate algebra $A(X)$ is equipped with a right $A(G)$ -comodule structure

$$\tau : A(X) \rightarrow A(X) \otimes A(G)$$

which is simultaneously an algebra homomorphism. By the duality between $U(\mathfrak{g})$ and $A(G)$ we obtain a locally finite left $U(\mathfrak{g})$ -module structure

$$(*) \quad \gamma : U(\mathfrak{g}) \otimes A(X) \rightarrow A(X)$$

given by

$$(**) \quad \tau(n) = \sum_i n_i \otimes f_i \Rightarrow \gamma(u \otimes n) = \sum_i \langle u, f_i \rangle n_i,$$

where $\langle \cdot, \cdot \rangle : U(\mathfrak{g}) \times A(G) \rightarrow \mathbb{C}$ is the dual pairing. Since τ is an algebra homomorphism, we have

$$(***) \quad u \in U(\mathfrak{g}), m, n \in A(X), \Delta(u) = \sum_i u_i \otimes v_i \Rightarrow u(mn) = \sum_i (u_i m)(v_i n),$$

where $\Delta : U(\mathfrak{g}) \rightarrow U(\mathfrak{g}) \otimes U(\mathfrak{g})$ is the coproduct. Then the action of G on X is uniquely determined by the infinitesimal action γ . Moreover, for a locally finite left

$U(\mathfrak{g})$ -module structure $(*)$ on $A(X)$ satisfying $(***)$ and a certain condition on irreducible $U(\mathfrak{g})$ -modules appearing as submodules of $A(X)$, there exists a unique action of G on X whose infinitesimal action is given by γ .

Now we define the notion of a quantum deformation of an affine variety X with G -action as follows. A (not necessarily commutative) $\mathbb{C}(q)$ -algebra $A_q(X)$ endowed with a locally finite left $U_q(\mathfrak{g})$ -module structure

$$\gamma_q : U_q(\mathfrak{g}) \otimes A_q(X) \rightarrow A_q(X)$$

is called a quantum deformation of X if $A_q(X)$ and γ_q tend to $A(X)$ and $\gamma : U(\mathfrak{g}) \otimes A(X) \rightarrow A(X)$ respectively when q tends to 1 and if it satisfies

$$u \in U_q(\mathfrak{g}), \quad m, n \in A_q(X), \quad \Delta(u) = \sum_i u_i \otimes v_i \Rightarrow u(mn) = \sum_i (u_i m)(v_i n).$$

It seems to be an interesting problem to determine in which case X admits a quantum deformation. In this paper we consider the problem when X is a prehomogeneous vector space, that is, when X is a vector space with a linear G -action containing an open G -orbit. Such a quantum deformation was intensively studied in the case where $G = GL_m(\mathbb{C}) \times GL_n(\mathbb{C})$ and $X = M_{mn}(\mathbb{C})$ (see Taft-Towber [10], Hashimoto-Hayashi [2] and Noumi-Yamada-Mimachi [7]), and also in the case where $G = GL_n(\mathbb{C})$ and X is the set of skew symmetric matrices of degree n (see Strickland [8]).


In our previous paper [4] we gave a general method to construct quantum deformations of prehomogeneous vector spaces of parabolic type. Moreover, for each non-open G -orbit C on X , we have shown that the defining ideal of the closure \overline{C} and its canonical generators admit quantum deformations inside $A_q(X)$. It includes the existence of the quantum deformation of the irreducible relative invariant when X is a regular prehomogeneous vector space. Indeed, the canonical generator of the defining ideal of the closure of the one-codimensional orbit is nothing but the irreducible relative invariant.

Quantum deformations of prehomogeneous vector spaces of commutative parabolic type associated to classical simple Lie algebras are intensively studied in Kamita [5]. In this paper we shall deal with the remaining two cases

- (I) $G = \mathbb{C}^\times \times \text{Spin}(10, \mathbb{C})$, $X = \mathbb{C}^{16}$, the scalar multiplication and the half-spin representation,
- (II) $G = \mathbb{C}^\times \times E_6$, $X = \mathbb{C}^{27}$, the scalar multiplication and the 27-dimensional irreducible representation of E_6 ,

which naturally arise from the exceptional simple Lie algebras of type E_6 and E_7 respectively using the method in our previous paper [4]. In Introduction we shall only state the results in case (II).

Let \mathfrak{g}_{E_7} be a simple Lie algebra of type E_7 over \mathbb{C} and let \mathfrak{h} be its Cartan subalgebra. We shall use the labelling of the vertices of the Dynkin diagram 1.



Dynkin diagram 1.

Set $I_0 = \{1, 2, \dots, 7\}$, $I = I_0 \setminus \{1\}$. Let $\Delta \subset \mathfrak{h}^*$ be the root system of type E_7 . We denote the set of simple roots by $\{\alpha_i\}_{i \in I_0}$ and the set of positive roots by Δ^+ . Let $(\cdot, \cdot) : \mathfrak{h}^* \times \mathfrak{h}^* \rightarrow \mathbb{C}$ be a standard symmetric bilinear form. Set $D = \Delta^+ \setminus \sum_{i \in I} \mathbb{Z}\alpha_i$. Then we have $\sharp D = 27$. Set $\Lambda = \{1, 2, \dots, 27\}$, and fix a bijection $\Lambda \ni j \mapsto \beta_j \in D$ such that $\beta_k - \beta_j \in \sum_{i \in I_0} \mathbb{Z}_{\geq 0}\alpha_i$ implies $j \leq k$, where $\mathbb{Z}_{\geq 0} = \{n \in \mathbb{Z} \mid n \geq 0\}$. Set $\delta = 3\alpha_1 + 4\alpha_2 + 5\alpha_3 + 6\alpha_4 + 3\alpha_5 + 4\alpha_6 + 2\alpha_7$. For each $n \in \Lambda$ there exist exactly five pairs $(i, j) \in \Lambda^2$ such that $\beta_i + \beta_j = \delta - \beta_n$, $i < j$. We denote them by $(i_1^n, j_1^n), (i_2^n, j_2^n), (i_3^n, j_3^n), (i_4^n, j_4^n), (i_5^n, j_5^n) \in \Lambda^2$ where $i_5^n < i_4^n < i_3^n < i_2^n < i_1^n$. Let $K_i^{\pm 1}, E_i, F_i$ ($i \in I_0$) be the canonical generators of $U_q(\mathfrak{g}_{E_7})$, and set $U_q(\mathfrak{g}) = \langle K_1^{\pm 3}, K_j^{\pm 1}, E_j, F_j \mid j \in I \rangle \subset U_q(\mathfrak{g}_{E_7})$. Then $U_q(\mathfrak{g})$ is isomorphic to the tensor product of $\mathbb{C}(q)[K, K^{-1}]$ and the quantized enveloping algebra of type E_6 , where $K = K_1^3 K_2^4 K_3^5 K_4^6 K_5^3 K_6^4 K_7^2$.

Theorem 0.1. *A quantum deformation of the 27-dimensional irreducible prehomogeneous vector space X of $G = \mathbb{C}^\times \times E_6$ is given by the following.*

(a) $A_q(X)$ is an associative $\mathbb{C}(q)$ -algebra defined by the following generators and fundamental relations:

Generators: Y_i with $i = 1, \dots, 27$.

Fundamental relations: For $i < j$

$$Y_i Y_j = \begin{cases} qY_j Y_i & \text{if } \beta_i + \beta_j \text{ does not have another decomposition } \beta + \beta', \beta, \beta' \in D, \\ Y_j Y_i + qY_b Y_a - q^{-1}Y_a Y_b & \text{if there exist } k \in I, a, b \in \Lambda \text{ such that } \beta_a = \beta_i + \alpha_k, \beta_b = \beta_j - \alpha_k, \\ Y_j Y_i & \text{otherwise.} \end{cases}$$

(b) The action $\gamma_q : U_q(\mathfrak{g}) \otimes A_q(X) \rightarrow A_q(X)$ is given by the following.

For $2 \leq k \leq 7, 1 \leq m \leq 7$

$$\gamma_q(F_k \otimes Y_i) = \begin{cases} Y_j & \text{if there exists } j \text{ such that } \beta_j = \beta_i + \alpha_k, \\ 0 & \text{otherwise,} \end{cases}$$

$$\gamma_q(E_k \otimes Y_i) = \begin{cases} Y_j & \text{if there exists } j \text{ such that } \beta_j = \beta_i - \alpha_k, \\ 0 & \text{otherwise,} \end{cases}$$

$$\gamma_q(K_m \otimes Y_i) = q^{-(\alpha_m, \beta_i)} Y_i.$$

(c) The quantum deformation of the irreducible relative invariant of X is given by

$$\varphi = \sum_{n \in \Lambda} (-q)^{|\beta_n|-1} Y_n \psi_n,$$

where $|\beta| = \sum_{i \in I_0} m_i$ ($\beta = \sum_{i \in I_0} m_i \alpha_i$), $\psi_n = Y_{i_3^n} Y_{j_3^n} - q Y_{i_4^n} Y_{j_4^n} + q^2 Y_{i_5^n} Y_{j_5^n} - q^3 Y_{i_2^n} Y_{j_2^n} + q^4 Y_{i_1^n} Y_{j_1^n}$.

The author expresses gratitude to Professor Noriaki Kawanaka and Professor Toshiyuki Tanisaki.

1. Preliminaries

Let \mathfrak{g} be a simple Lie algebra of type E_6 or E_7 over the complex number field \mathbb{C} , and let \mathfrak{h} be a Cartan subalgebra of \mathfrak{g} . Let $\Delta \subset \mathfrak{h}^*$ be the root system, and let $W \subset GL(\mathfrak{h})$ be the Weyl group. We denote the set of positive roots by Δ^+ and the set of simple roots by $\{\alpha_i\}_{i \in I_0}$, where I_0 is an index set. For $i \in I_0$ we denote the simple reflection corresponding to α_i by $s_i \in W$. Let $(\ , \) : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ be the invariant symmetric bilinear form such that $(\alpha, \alpha) = 2$ for any $\alpha \in \Delta$. Set $a_{ij} = (\alpha_i, \alpha_j)$. The matrix $(a_{ij})_{i,j \in I_0}$ is called the Cartan matrix of type E_6 or E_7 . For $\alpha \in \Delta$ we denote the corresponding root space by \mathfrak{g}_α . Set $\mathfrak{n}^+ = \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_\alpha$, $\mathfrak{n}^- = \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_{-\alpha}$. For a subset $I \subset I_0$ we define

$$\Delta_I = \Delta \cap \sum_{i \in I} \mathbb{Z} \alpha_i, \quad W_I = \langle s_i \mid i \in I \rangle.$$

We set

$$\mathfrak{l}_I = \mathfrak{h} \oplus \left(\bigoplus_{\alpha \in \Delta_I} \mathfrak{g}_\alpha \right), \quad \mathfrak{n}_I^+ = \bigoplus_{\alpha \in \Delta^+ \setminus \Delta_I} \mathfrak{g}_\alpha, \quad \mathfrak{n}_I^- = \bigoplus_{\alpha \in \Delta^+ \setminus \Delta_I} \mathfrak{g}_{-\alpha}.$$

Let G be a connected algebraic group with Lie algebra \mathfrak{g} . We denote by L_I the subgroup of G corresponding to \mathfrak{l}_I . Then L_I acts on \mathfrak{n}_I^\pm via the adjoint action.

The quantized enveloping algebra $U_q(\mathfrak{g})$ (Drinfel'd [1], Jimbo [3]) is an associative algebra over the rational function field $\mathbb{C}(q)$ generated by the elements E_i, F_i, K_i, K_i^{-1} ($i \in I_0$) satisfying the following fundamental relations:

$$\begin{aligned} K_i K_j &= K_j K_i, & K_i K_i^{-1} &= K_i^{-1} K_i = 1, \\ K_i E_j &= q^{a_{ij}} E_j K_i, & K_i F_j &= q^{-a_{ij}} F_j K_i, \\ E_i F_j - F_j E_i &= \delta_{ij} \frac{K_i - K_i^{-1}}{q - q^{-1}}, \\ E_i E_j &= E_j E_i & (i \neq j, a_{ij} = 0), \\ E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 &= 0 & (i \neq j, a_{ij} = -1), \\ F_i F_j &= F_j F_i & (i \neq j, a_{ij} = 0), \\ F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 &= 0 & (i \neq j, a_{ij} = -1). \end{aligned}$$

A Hopf algebra structure on $U_q(\mathfrak{g})$ is defined as follows. The comultiplication $\Delta : U_q(\mathfrak{g}) \rightarrow U_q(\mathfrak{g}) \otimes U_q(\mathfrak{g})$ is the algebra homomorphism satisfying

$$\Delta(K_i) = K_i \otimes K_i, \quad \Delta(E_i) = E_i \otimes K_i^{-1} + 1 \otimes E_i, \quad \Delta(F_i) = F_i \otimes 1 + K_i \otimes F_i.$$

The counit $\epsilon : U_q(\mathfrak{g}) \rightarrow \mathbb{C}(q)$ is the algebra homomorphism satisfying

$$\epsilon(K_i) = 1, \quad \epsilon(E_i) = \epsilon(F_i) = 0.$$

The antipode $S : U_q(\mathfrak{g}) \rightarrow U_q(\mathfrak{g})$ is the algebra antiautomorphism satisfying

$$S(K_i) = K_i^{-1}, \quad S(E_i) = -E_i K_i, \quad S(F_i) = -K_i^{-1} F_i.$$

Using the Hopf algebra structure, we define the adjoint action of $U_q(\mathfrak{g})$ on $U_q(\mathfrak{g})$ as follows. For $x, y \in U_q(\mathfrak{g})$ write $\Delta(x) = \sum_k x_k^1 \otimes x_k^2$ and set $\text{ad}(x)y = \sum_k x_k^1 y S(x_k^2)$. Then $\text{ad} : U_q(\mathfrak{g}) \rightarrow \text{End}_{\mathbb{C}(q)}(U_q(\mathfrak{g}))$ is an algebra homomorphism. For $x, y, z \in U_q(\mathfrak{g})$ we have $\text{ad}(x)(yz) = \sum_k (\text{ad}(x_k^1)y)(\text{ad}(x_k^2)z)$, where $\Delta(x) = \sum_k x_k^1 \otimes x_k^2$.

We define subalgebras $U_q(\mathfrak{n}^-)$ and $U_q(\mathfrak{l}_I)$ for $I \subset I_0$ by

$$U_q(\mathfrak{n}^-) = \langle F_i \mid i \in I_0 \rangle, \quad U_q(\mathfrak{l}_I) = \langle E_i, F_i, K_j, K_j^{-1} \mid i \in I, j \in I_0 \rangle.$$

For $i \in I_0$ we define an algebra automorphism T_i of $U_q(\mathfrak{g})$ by

$$\begin{aligned} T_i(K_j) &= K_j K_i^{-a_{ij}}, \\ T_i(E_j) &= \begin{cases} -F_i K_i & (i = j) \\ E_j & (i \neq j, a_{ij} = 0) \\ E_i E_j - q^{-1} E_j E_i & (i \neq j, a_{ij} = -1), \end{cases} \\ T_i(F_j) &= \begin{cases} -K_i^{-1} E_i & (i = j) \\ F_j & (i \neq j, a_{ij} = 0) \\ F_j F_i - q F_i F_j & (i \neq j, a_{ij} = -1) \end{cases} \end{aligned}$$

(see Lusztig [6]). For $w \in W$ choose a reduced expression $w = s_{i_1} \cdots s_{i_r}$ and set $T_w = T_{i_1} \cdots T_{i_r}$. It is known that T_w does not depend on the choice of a reduced expression.

We shall use the following later (see Lusztig [6]).

Lemma 1.1. *If $w(\alpha_i) = \alpha_j$ for $w \in W$ and $i, j \in I_0$, then we have $T_w(F_i) = F_j$.*

For $I \subset I_0$ let w_I be the longest element of W_I and let w_0 be the longest element of W . Choose a reduced expression $w_I w_0 = s_{i_1} \cdots s_{i_r}$ of $w_I w_0$ and set

$$\beta_j = s_{i_1} s_{i_2} \cdots s_{i_{j-1}}(\alpha_{i_j}), \quad Y_j = Y_{\beta_j} = T_{i_1} \cdots T_{i_{j-1}}(F_{i_j})$$

for $1 \leq j \leq r$. Then it is known that $\{\beta_j \mid 1 \leq j \leq r\} = \Delta^+ \setminus \Delta_I$. Set

$$U_q(\mathfrak{n}_I^-) = \sum_{d_j \geq 0} \mathbb{C}(q) Y_1^{d_1} \cdots Y_r^{d_r}.$$

Then $\{Y_1^{d_1} \cdots Y_r^{d_r} \mid d_j \in \mathbb{Z}_{\geq 0}, 1 \leq j \leq r\}$ is a basis of $U_q(\mathfrak{n}_I^-)$ and $U_q(\mathfrak{n}_I^-)$ is a subalgebra of $U_q(\mathfrak{n}^-)$. we have


$$U_q(\mathfrak{n}_I^-) = U_q(\mathfrak{n}^-) \cap T_{w_I}^{-1} U_q(\mathfrak{n}^-)$$

and $U_q(\mathfrak{n}_I^-)$ does not depend on the choice of a reduced expression of $w_I w_0$ (see Lusztig [6]).

If $\mathfrak{n}_I^+ \neq \{0\}$, $[\mathfrak{n}_I^+, \mathfrak{n}_I^+] = \{0\}$, then Y_β for $\beta \in \Delta^+ \setminus \Delta_I$ does not depend on the choice of a reduced expression of $w_I w_0$ (see [4]). In this case we denote the $\mathbb{C}(q)$ -algebra $U_q(\mathfrak{n}_I^-)$ by A_q . We can regard it as a quantum deformation of the coordinate algebra $A = \mathbb{C}[\mathfrak{n}_I^+]$ of \mathfrak{n}_I^+ as explained in [4].

2. Case of type E_6

Let \mathfrak{g} be a simple Lie algebra of type E_6 . We shall use the labelling of the vertices of the Dynkin diagram 2.



Dynkin diagram 2.

Hence we have $I_0 = \{1, 2, 3, 4, 5, 6\}$. Set $I = \{2, 3, 4, 5, 6\}$. In this case we have $\mathfrak{n}_I^+ \neq \{0\}$, $[\mathfrak{n}_I^+, \mathfrak{n}_I^+] = \{0\}$. Then \mathfrak{l}_I is isomorphic to $\mathbb{C} \oplus \mathfrak{o}(10, \mathbb{C})$ and \mathfrak{n}_I^+ is a 16-dimensional irreducible prehomogeneous vector space. There are three L_I -orbits $\{0\}, C_0, O$ on \mathfrak{n}_I^+ satisfying $\{0\} \subset \overline{C_0} \subset \overline{O}$. Let $J_{C_0} \subset \mathbb{C}[\mathfrak{n}_I^+]$ be the defining ideal of the closure of C_0 , and let $J_{C_0}^0$ denote the subspace of J_{C_0} consisting of the polynomials in J_{C_0} with homogeneous degree 2. Then $J_{C_0}^0$ is a ten-dimensional irreducible \mathfrak{l}_I -module, and it generates the ideal J_{C_0} .

We fix a reduced expression

$$w_I w_0 = s_1 s_2 s_3 s_4 s_5 s_3 s_2 s_1 s_6 s_5 s_3 s_2 s_4 s_3 s_5 s_6$$

of $w_I w_0$ and define the elements Y_i ($i \in \Lambda = \{1, 2, \dots, 16\}$) as in Section 1.

Set $I'_0 = \{1, 2, 3, 4, 5\}$, $I' = \{2, 3, 4, 5\}$, $\Lambda' = \{1, 2, \dots, 8\}$. Then $\{\alpha_i\}_{i \in I'_0}$ is a set of simple roots of type D_5 . Let \mathfrak{g}' be the simple subalgebra of \mathfrak{g} corresponding to I'_0 . We choose a reduced expression $w_{I'} w_{I'_0} = s_1 s_2 s_3 s_4 s_5 s_3 s_2 s_1$ of $w_{I'} w_{I'_0}$. The elements Y_i ($i \in \Lambda'$) can be computed inside $U_q(\mathfrak{g}')$.

Let $\beta_j = \sum_{i \in I'_0} m_i^j \alpha_i$ and set $\mathbf{m}^j = (m_1^j, \dots, m_5^j)$ for $j \in \Lambda$. Then we have

$$\begin{aligned}
 \mathbf{m}^1 &= (1, 0, 0, 0, 0, 0), & \mathbf{m}^2 &= (1, 1, 0, 0, 0, 0), & \mathbf{m}^3 &= (1, 1, 1, 0, 0, 0), \\
 \mathbf{m}^4 &= (1, 1, 1, 1, 0, 0), & \mathbf{m}^5 &= (1, 1, 1, 0, 1, 0), & \mathbf{m}^6 &= (1, 1, 1, 1, 1, 0), \\
 \mathbf{m}^7 &= (1, 1, 2, 1, 1, 0), & \mathbf{m}^8 &= (1, 2, 2, 1, 1, 0), & \mathbf{m}^9 &= (1, 1, 1, 0, 1, 1), \\
 \mathbf{m}^{10} &= (1, 1, 1, 1, 1, 1), & \mathbf{m}^{11} &= (1, 1, 2, 1, 1, 1), & \mathbf{m}^{12} &= (1, 2, 2, 1, 1, 1), \\
 \mathbf{m}^{13} &= (1, 1, 2, 1, 2, 1), & \mathbf{m}^{14} &= (1, 2, 2, 1, 2, 1), & \mathbf{m}^{15} &= (1, 2, 3, 1, 2, 1), \\
 \mathbf{m}^{16} &= (1, 2, 3, 2, 2, 1).
 \end{aligned}$$

If $(\beta_j, \alpha_k) = -1$ for $j \in \Lambda$ and $k \in I$, then $s_k(\beta_j) = \beta_j + \alpha_k \in \Delta^+$. Since $k \neq 1$ and $m_1^j = 1$, we have $\beta_j + \alpha_k \notin \Delta_I$. Therefore there exists $l \in \Lambda$ satisfying $\beta_j + \alpha_k = \beta_l$. Conversely if $\beta_j + \alpha_k = \beta_l$ ($j, l \in \Lambda, k \in I$), then we have $(\beta_j, \alpha_k) = -1, s_k(\beta_j) = \beta_l$.

There exist 20 triplets $(k, j, l) \in I \times \Lambda \times \Lambda$ satisfying $\beta_j + \alpha_k = \beta_l$. The triplets are the following: (2, 1, 2), (3, 2, 3), (4, 3, 4), (5, 3, 5), (5, 4, 6), (4, 5, 6), (3, 6, 7), (2, 7, 8), (6, 5, 9), (4, 9, 10), (3, 10, 11), (2, 11, 12), (5, 11, 13), (5, 12, 14), (2, 13, 14), (3, 14, 15), (4, 15, 16), (6, 6, 10), (6, 7, 11), (6, 8, 12).

For $k \in I, j \in \Lambda$, we have $\beta_j - 2\alpha_k, \beta_j + 2\alpha_k \notin \Delta^+ \setminus \Delta_I$.

Lemma 2.1. *Let $\beta, \beta' \in \Delta^+ \setminus \Delta_I$ satisfying $\beta + \alpha_k = \beta'$ ($k \in I$). Then we can choose a reduced expression $w_I w_0 = s_{i_1} s_{i_2} \cdots s_{i_{16}}$ and $p \in \Lambda$ satisfying*

$$\begin{aligned}
 \beta &= s_{i_1} s_{i_2} \cdots s_{i_{p-1}}(\alpha_{i_p}), & \beta' &= s_{i_1} s_{i_2} \cdots s_{i_{p-1}} s_{i_p}(\alpha_{i_{p+1}}), & (\alpha_{i_p}, \alpha_{i_{p+1}}) &= -1, \\
 \alpha_k &= s_{i_1} s_{i_2} \cdots s_{i_{p-1}}(\alpha_{i_{p+1}}).
 \end{aligned}$$

Proof. Among the 20 triplets (k, j, l) satisfying $\beta_j + \alpha_k = \beta_l$ ($k \in I, j, l \in \Lambda$), the 12 triplets satisfy $l = j + 1, (\alpha_{i_j}, \alpha_{i_{j+1}}) = -1$. Therefore it is sufficient to deal with the remaining 8 cases. In the cases $(k, j, l) = (5, 3, 5), (5, 4, 6), (5, 11, 13), (5, 12, 14)$, the reduced expression

$$w_I w_0 = s_1 s_2 s_3 s_5 s_4 s_3 s_2 s_1 s_6 s_5 s_3 s_4 s_2 s_3 s_5 s_6$$

of $w_I w_0$ with $p = 3, 5, 11, 13$ respectively satisfies the required properties. In the cases $(k, j, l) = (6, 5, 9), (6, 6, 10), (6, 7, 11), (6, 8, 12)$, the reduced expression

$$w_I w_0 = s_1 s_2 s_3 s_4 s_5 s_6 s_3 s_5 s_2 s_3 s_1 s_2 s_4 s_3 s_5 s_6$$

of $w_I w_0$ with $p = 5, 7, 9, 11$ respectively satisfies the required properties. □

It is known that $U_q(\mathfrak{n}_I^+) = \bigoplus_{\beta \in \Delta^+ \setminus \Delta_I} \mathbb{C}(q) Y_\beta$ is an irreducible $U_q(\mathfrak{l}_I)$ -module. (see [4])

Lemma 2.2. *For $k \in I, j \in \Lambda$, we have*

$$\text{ad}(F_k) Y_j = \begin{cases} Y_l & \text{if there exists } l \in \Lambda \text{ such that } \beta_l = \beta_j + \alpha_k, \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{ad}(E_k)Y_j = \begin{cases} Y_l & \text{if there exists } l \in \Lambda \text{ such that } \beta_l = \beta_j - \alpha_k, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Since $\bigoplus_{j \in \Lambda} \mathbb{C}(q)Y_j$ is a $U_q(\mathfrak{l}_I)$ -module, we have $\text{ad}(F_k)Y_j = 0$ if $\beta_j + \alpha_k \notin \Delta^+ \setminus \Delta_I$, and we have $\text{ad}(E_k)Y_j = 0$ if $\beta_j - \alpha_k \notin \Delta^+ \setminus \Delta_I$.

We shall show $\text{ad}(F_k)Y_\beta = Y_{\beta'}$ for $\beta, \beta' \in \Delta^+ \setminus \Delta_I$ and $k \in I$ satisfying $\beta' = \beta + \alpha_k$. By Lemma 2.1 we can choose a reduced expression of $w_I w_0 = s_{i_1} s_{i_2} \cdots s_{i_{16}}$ satisfying $\beta = s_{i_1} s_{i_2} \cdots s_{i_{p-1}}(\alpha_{i_p})$, $\beta' = s_{i_1} s_{i_2} \cdots s_{i_{p-1}} s_{i_p}(\alpha_{i_{p+1}})$, $(\alpha_{i_p}, \alpha_{i_{p+1}}) = -1$. Then we can write $Y_\beta = T_{i_1} T_{i_2} \cdots T_{i_{p-1}}(F_{i_p})$, $Y_{\beta'} = T_{i_1} T_{i_2} \cdots T_{i_{p-1}} T_{i_p}(F_{i_{p+1}})$. Since $(\alpha_{i_p}, \alpha_{i_{p+1}}) = -1$, we have $T_{i_p}(F_{i_{p+1}}) = F_{i_{p+1}} F_{i_p} - q F_{i_p} F_{i_{p+1}}$. Moreover, since $\alpha_k = s_{i_1} s_{i_2} \cdots s_{i_{p-1}}(\alpha_{i_{p+1}})$, we have $T_{i_1} T_{i_2} \cdots T_{i_{p-1}}(F_{i_{p+1}}) = F_k$ by Lemma 1.1, and hence

$$\begin{aligned} Y_{\beta'} &= T_{i_1} T_{i_2} \cdots T_{i_{p-1}} T_{i_p}(F_{i_{p+1}}) \\ &= T_{i_1} T_{i_2} \cdots T_{i_{p-1}}(F_{i_{p+1}} F_{i_p} - q F_{i_p} F_{i_{p+1}}) = F_k Y_\beta - q Y_\beta F_k. \end{aligned}$$

Since $(\beta, \alpha_k) = -1$, we have $\text{ad}(F_k)Y_\beta = F_k Y_\beta - q Y_\beta F_k$. Hence we have $\text{ad}(F_k)Y_\beta = Y_{\beta'}$.

Let us show $\text{ad}(E_k)Y_\beta = Y_{\beta'}$ for $\beta, \beta' \in \Delta^+ \setminus \Delta_I$ and $k \in I$ satisfying $\beta' = \beta - \alpha_k$. By the above argument we have $Y_\beta = \text{ad}(F_k)Y_{\beta'} = F_k Y_{\beta'} - q Y_{\beta'} F_k$. Since $\beta' - \alpha_k = \beta - 2\alpha_k \notin \Delta^+ \setminus \Delta_I$, we have $\text{ad}(E_k)Y_{\beta'} = 0$, and hence $E_k Y_{\beta'} = Y_{\beta'} E_k$. Since $(\beta', \alpha_k) = -1$, we have $K_k Y_{\beta'} = q Y_{\beta'} K_k$. Hence we have

$$\begin{aligned} \text{ad}(E_k)Y_\beta &= (E_k Y_\beta - Y_\beta E_k) K_k = (E_k (F_k Y_{\beta'} - q Y_{\beta'} F_k) - (F_k Y_{\beta'} - q Y_{\beta'} F_k) E_k) K_k \\ &= \left(\frac{K_k - K_k^{-1}}{q - q^{-1}} Y_{\beta'} - q Y_{\beta'} \frac{K_k - K_k^{-1}}{q - q^{-1}} \right) K_k = (Y_{\beta'} K_k^{-1}) K_k = Y_{\beta'}. \quad \square \end{aligned}$$

Next we shall consider quadratic fundamental relations among the elements Y_i . Since we have

$$\sum_{i, j \in \Lambda} \mathbb{C}(q)Y_i Y_j = \bigoplus_{s \leq t} \mathbb{C}(q)Y_s Y_t,$$

we can write

$$Y_i Y_j = \sum_{\substack{s \leq t \\ \beta_i + \beta_j = \beta_s + \beta_t}} a_{s,t}^{i,j} Y_s Y_t \quad (a_{s,t}^{i,j} \in \mathbb{C}(q))$$

for $i > j$ (see [4]). Hence if $\beta_i + \beta_j$ does not have another decomposition $\beta + \beta'$ ($\beta, \beta' \in \Delta^+ \setminus \Delta_I$, $\beta_i + \beta_j = \beta + \beta'$) then we have $Y_i Y_j = a_{i,j} Y_i Y_j$ for some $a_{i,j} \in \mathbb{C}(q)$. We denote the set of weights of the ten-dimensional irreducible highest weight \mathfrak{l}_I -module $J_{C_0}^0$ with highest weight $-\beta_1 - \beta_8$ by Γ . For $\beta, \beta' \in \Delta^+ \setminus \Delta_I$ a weight $\beta + \beta'$ has another decomposition if and only if we have $-(\beta + \beta') \in \Gamma$. We fix a bijection

$\{1, 2, \dots, 10\} \ni n \mapsto -\delta_n \in \Gamma$ such that if $\delta_m - \delta_n \in \sum_{i \in I_0} \mathbb{Z}_{\geq 0} \alpha_i$, then $n \leq m$. For each n there exist exactly four pairs $(i, j) \in \Lambda^2$ such that $i < j$, $\beta_i + \beta_j = \delta_n$. We denote them by $(i_1^n, j_1^n), (i_2^n, j_2^n), (i_3^n, j_3^n), (i_4^n, j_4^n) \in \Lambda^2$ where $i_4^n < i_3^n < i_2^n < i_1^n$. Set $\mathbf{A}(n) = (i_4^n, i_3^n, i_2^n, i_1^n, j_1^n, j_2^n, j_3^n, j_4^n) \in \Lambda^8$ ($1 \leq n \leq 10$). Then we have

$$\begin{aligned} \mathbf{A}(1) &= (1, 2, 3, 4, 5, 6, 7, 8), & \mathbf{A}(2) &= (1, 2, 3, 4, 9, 10, 11, 12), \\ \mathbf{A}(3) &= (1, 2, 5, 6, 9, 10, 13, 14), & \mathbf{A}(4) &= (1, 3, 5, 7, 9, 11, 13, 15), \\ \mathbf{A}(5) &= (2, 3, 5, 8, 9, 12, 14, 15), & \mathbf{A}(6) &= (1, 4, 6, 7, 10, 11, 13, 16), \\ \mathbf{A}(7) &= (2, 4, 6, 8, 10, 12, 14, 16), & \mathbf{A}(8) &= (3, 4, 7, 8, 11, 12, 15, 16), \\ \mathbf{A}(9) &= (5, 6, 7, 8, 13, 14, 15, 16), & \mathbf{A}(10) &= (9, 10, 11, 12, 13, 14, 15, 16). \end{aligned}$$

We denote the set $\{i_4^n, i_3^n, i_2^n, i_1^n, j_1^n, j_2^n, j_3^n, j_4^n\}$ by $|\mathbf{A}(n)|$ for $1 \leq n \leq 10$. For any $i, j \in \Lambda$ there exists n satisfying $i, j \in |\mathbf{A}(n)|$.

Set

$$\mathcal{A} = \{(k, n, n') \in I \times \Lambda \times \Lambda \mid \delta_n + \alpha_k = \delta_{n'}\}.$$

Then

$$\mathcal{A} = \{(6, 1, 2), (5, 2, 3), (3, 3, 4), (2, 4, 5), (4, 4, 6), (2, 6, 7), (4, 5, 7), (3, 7, 8), (5, 8, 9), (6, 9, 10)\}.$$

For any $n \in \{2, 3, \dots, 10\}$ we can take a sequence $((k_1, n_1, n'_1), \dots, (k_s, n_s, n'_s))$ of \mathcal{A} satisfying $n_1 = 1, n'_s = n, n'_j = n_{j+1}$ ($1 \leq j \leq s - 1$).

For $(k, n, n') \in \mathcal{A}$ and $m \in \{1, 2, 3, 4\}$, we have either

$$(P_m^+) \quad (\beta_{i_m^n}, \alpha_k) = 0, i_m^{n'} = i_m^n, (\beta_{j_m^n}, \alpha_k) = -1, \beta_{j_m^{n'}} = \beta_{j_m^n} + \alpha_k$$

or

$$(P_m^-) \quad (\beta_{i_m^n}, \alpha_k) = -1, \beta_{i_m^{n'}} = \beta_{i_m^n} + \alpha_k, (\beta_{j_m^n}, \alpha_k) = 0, j_m^{n'} = j_m^n.$$

Proposition 2.3. For any $i, j \in \Lambda$ satisfying $i < j$, we have

$$(Q6) \quad Y_i Y_j = \begin{cases} Y_j Y_i & \text{if there exists } n \text{ such that } i = i_1^n, j = j_1^n, \\ Y_{j_2^n} Y_{i_2^n} + (q - q^{-1}) Y_{i_1^n} Y_{j_1^n} & \text{if there exists } n \text{ such that } i = i_2^n, j = j_2^n, \\ Y_{j_m^n} Y_{i_m^n} + q Y_{j_{m-1}^n} Y_{i_{m-1}^n} - q^{-1} Y_{i_{m-1}^n} Y_{j_{m-1}^n} & \text{if there exist } n, m \in \{3, 4\} \text{ such that } i = i_m^n, j = j_m^n, \\ q Y_j Y_i & \text{otherwise.} \end{cases}$$

Proof. Since there exists some n satisfying $i, j \in |\mathbf{A}(n)|$ for any $i, j \in \Lambda$, it is sufficient to show that for any $1 \leq n \leq 10$ the elements $Y_{i_m^n}, Y_{j_m^n}$ ($1 \leq m \leq 4$) satisfy

the following relations.

$$(Rn) \quad \begin{cases} Y_{i_1}^n Y_{j_1}^n = Y_{j_1}^n Y_{i_1}^n & (Rn, 1) \\ Y_{i_m}^n Y_{j_m}^n = Y_{j_m}^n Y_{i_m}^n + q Y_{j_{m-1}}^n Y_{i_{m-1}}^n - q^{-1} Y_{i_{m-1}}^n Y_{j_{m-1}}^n \quad (2 \leq m \leq 4) & (Rn, 2) \\ Y_{l_1} Y_{l_2} = q Y_{l_2} Y_{l_1} \\ \quad (l_1, l_2 \in |A(n)|, l_1 < l_2, (l_1, l_2) \neq (i_m^n, j_m^n) \quad (1 \leq m \leq 4)) & (Rn, 3) \end{cases}$$

When $n = 1$, the elements Y_i ($1 \leq i \leq 8$) satisfy the same relations as those for type D_5 , hence the relations (R1) hold.

For any $m > 1$ there exists a sequence $((k_1, n_1, n'_1), \dots, (k_s, n_s, n'_s))$ of \mathcal{A} satisfying $n_1 = 1, n'_s = m, n'_j = n_{j+1}$ ($1 \leq j \leq s-1$), and hence it is sufficient to show the relations (Rn') for $(k, n, n') \in \mathcal{A}$ assuming the relations (Rn).

Let $(k, n, n') \in \mathcal{A}$. Assume that the relations (Rn) hold.

We first show that the relation (Rn',1) holds. If the condition (P_1^+) is satisfied, then we have $Y_{i_1}^{n'} = Y_{i_1}^n, F_k Y_{i_1}^n = Y_{i_1}^n F_k, Y_{j_1}^{n'} = \text{ad}(F_k) Y_{j_1}^n = F_k Y_{j_1}^n - q Y_{j_1}^n F_k$. Since $Y_{i_1}^n Y_{j_1}^n = Y_{j_1}^n Y_{i_1}^n$, we have

$$\begin{aligned} Y_{i_1}^{n'} Y_{j_1}^{n'} &= Y_{i_1}^n \text{ad}(F_k) Y_{j_1}^n = Y_{i_1}^n (F_k Y_{j_1}^n - q Y_{j_1}^n F_k) \\ &= (F_k Y_{j_1}^n - q Y_{j_1}^n F_k) Y_{i_1}^n = Y_{j_1}^{n'} Y_{i_1}^{n'}. \end{aligned}$$

If the condition (P_1^-) is satisfied, then we can prove the formula (Rn',1) similarly.

Next we prove the formula (Rn',2). Assume the condition (P_m^+) is satisfied, then we have

$$\begin{aligned} Y_{i_m}^{n'} Y_{j_m}^{n'} &= Y_{i_m}^n (F_k Y_{j_m}^n - q Y_{j_m}^n F_k) \\ &= F_k Y_{j_m}^n Y_{i_m}^n - q Y_{j_m}^n F_k Y_{i_m}^n \\ &\quad + q (F_k Y_{j_{m-1}}^n Y_{i_{m-1}}^n - q Y_{j_{m-1}}^n Y_{i_{m-1}}^n F_k) \\ &\quad - q^{-1} (F_k Y_{i_{m-1}}^n Y_{j_{m-1}}^n - q Y_{i_{m-1}}^n Y_{j_{m-1}}^n F_k). \end{aligned}$$

If the condition (P_{m-1}^+) is satisfied, then we have

$$\begin{aligned} F_k Y_{j_{m-1}}^n Y_{i_{m-1}}^n - q Y_{j_{m-1}}^n Y_{i_{m-1}}^n F_k &= Y_{j_{m-1}}^n (F_k Y_{i_{m-1}}^n - q Y_{i_{m-1}}^n F_k) = Y_{j_{m-1}}^{n'} Y_{i_{m-1}}^{n'}, \\ F_k Y_{i_{m-1}}^n Y_{j_{m-1}}^n - q Y_{i_{m-1}}^n Y_{j_{m-1}}^n F_k &= (F_k Y_{i_{m-1}}^n - q Y_{i_{m-1}}^n F_k) Y_{j_{m-1}}^n = Y_{i_{m-1}}^{n'} Y_{j_{m-1}}^{n'}, \end{aligned}$$

and if the condition (P_{m-1}^-) is satisfied, then we have

$$\begin{aligned} F_k Y_{j_{m-1}}^n Y_{i_{m-1}}^n - q Y_{j_{m-1}}^n Y_{i_{m-1}}^n F_k &= (F_k Y_{j_{m-1}}^n - q Y_{j_{m-1}}^n F_k) Y_{i_{m-1}}^n = Y_{j_{m-1}}^{n'} Y_{i_{m-1}}^{n'}, \\ F_k Y_{i_{m-1}}^n Y_{j_{m-1}}^n - q Y_{i_{m-1}}^n Y_{j_{m-1}}^n F_k &= Y_{i_{m-1}}^n (F_k Y_{j_{m-1}}^n - q Y_{j_{m-1}}^n F_k) = Y_{i_{m-1}}^{n'} Y_{j_{m-1}}^{n'}. \end{aligned}$$

Hence we have $Y_{i_m}^{n'} Y_{j_m}^{n'} = Y_{j_m}^{n'} Y_{i_m}^{n'} + q Y_{j_{m-1}}^{n'} Y_{i_{m-1}}^{n'} - q^{-1} Y_{i_{m-1}}^{n'} Y_{j_{m-1}}^{n'}$. The formula (Rn',2) is proved. When the condition (P_m^-) is satisfied, we can prove it similarly.

Finally we prove the formula (Rn',3). Let $l'_1, l'_2 \in |\mathbf{A}(n')|$ satisfying $l'_1 < l'_2$ and $(l'_1, l'_2) \neq (i_m^{n'}, j_m^{n'})$ for $1 \leq m \leq 4$. When $l'_p = i_m^{n'} \in |\mathbf{A}(n')|$ (resp. $l'_p = j_m^{n'}$), we denote $i_m^n \in |\mathbf{A}(n)|$ (resp. j_m^n) by l_p for $p = 1, 2$. Since $l_1 < l_2$ and $(l_1, l_2) \neq (i_m^n, j_m^n)$ for $1 \leq m \leq 4$, we have $Y_{l_1} Y_{l_2} = q Y_{l_2} Y_{l_1}$. We have the following possibilities:

- (1) $l'_1 = l_1, l'_2 = l_2, (\beta_{l_1}, \alpha_k) = (\beta_{l_2}, \alpha_k) = 0$,
- (2) $l'_1 = l_1, (\beta_{l_1}, \alpha_k) = 0, \beta_{l'_2} = \beta_{l_2} + \alpha_k, (\beta_{l_2}, \alpha_k) = -1$,
- (3) $\beta_{l'_1} = \beta_{l_1} + \alpha_k, (\beta_{l_1}, \alpha_k) = -1, l'_2 = l_2, (\beta_{l_2}, \alpha_k) = 0$,
- (4) $\beta_{l'_1} = \beta_{l_1} + \alpha_k, \beta_{l'_2} = \beta_{l_2} + \alpha_k, (\beta_{l_1}, \alpha_k) = (\beta_{l_2}, \alpha_k) = -1$.

In the case (1) the formula (Rn',3) is obvious.

In the case (2) we have $F_k Y_{l_1} = Y_{l_1} F_k, Y_{l'_2} = \text{ad}(F_k) Y_{l_2} = F_k Y_{l_2} - q Y_{l_2} F_k$. Hence we have

$$Y_{l'_1} Y_{l'_2} = Y_{l_1} (F_k Y_{l_2} - q Y_{l_2} F_k) = q (F_k Y_{l_2} - q Y_{l_2} F_k) Y_{l_1} = q Y_{l_2} Y_{l'_1}.$$

In the case (3) we can prove it similarly to the case (2).

In the case (4) we have $Y_{l'_p} = F_k Y_{l_p} - q Y_{l_p} F_k$ for $p = 1, 2$. Since $\beta_{l'_p} + \alpha_k = \beta_{l_p} + 2\alpha_k \notin \Delta^+ \setminus \Delta_I$ and $(\beta_{l'_p}, \alpha_k) = 1$, we have $\text{ad}(F_k) Y_{l'_p} = F_k Y_{l'_p} - q^{-1} Y_{l'_p} F_k = 0$ for $p = 1, 2$. Hence we have $F_k F_k Y_{l'_p} - (q + q^{-1}) F_k Y_{l'_p} F_k + Y_{l'_p} F_k F_k = 0, F_k Y_{l'_p} F_k = (q + q^{-1})^{-1} (F_k F_k Y_{l'_p} + Y_{l'_p} F_k F_k)$ for $p = 1, 2$. By these formulas we have

$$\begin{aligned} Y_{l'_1} Y_{l'_2} &= (F_k Y_{l_1} - q Y_{l_1} F_k)(F_k Y_{l_2} - q Y_{l_2} F_k) \\ &= F_k Y_{l_1} F_k Y_{l_2} - q F_k Y_{l_1} Y_{l_2} F_k - q Y_{l_1} F_k F_k Y_{l_2} + q^2 Y_{l_1} F_k Y_{l_2} F_k \\ &= \frac{1}{q + q^{-1}} F_k F_k Y_{l_1} Y_{l_2} + \frac{1}{q + q^{-1}} Y_{l_1} F_k F_k Y_{l_2} - q F_k Y_{l_1} Y_{l_2} F_k - q Y_{l_1} F_k F_k Y_{l_2} \\ &\quad + \frac{q^2}{q + q^{-1}} Y_{l_1} F_k F_k Y_{l_2} + \frac{q^2}{q + q^{-1}} Y_{l_1} Y_{l_2} F_k F_k \\ &= \frac{1}{q + q^{-1}} F_k F_k Y_{l_1} Y_{l_2} - q F_k Y_{l_1} Y_{l_2} F_k + \frac{q^2}{q + q^{-1}} Y_{l_1} Y_{l_2} F_k F_k. \end{aligned}$$

Similarly we have

$$Y_{l'_2} Y_{l'_1} = \frac{1}{q + q^{-1}} F_k F_k Y_{l_2} Y_{l_1} - q F_k Y_{l_2} Y_{l_1} F_k + \frac{q^2}{q + q^{-1}} Y_{l_2} Y_{l_1} F_k F_k.$$

Since $Y_{l_1} Y_{l_2} = q Y_{l_2} Y_{l_1}$, we have $Y_{l'_1} Y_{l'_2} = q Y_{l'_2} Y_{l'_1}$. □

By [4] and Proposition 2.3 we obtain the following:

Theorem 2.4. *The formulas (Q6) give fundamental relations for the generator system $\{Y_i\}_{i \in \Lambda}$ of the algebra $A_q = U_q(\mathfrak{n}_I^-)$.*

We shall construct a quantum deformation of the lowest degree part $J_{C_0}^0$ of the defining ideal J_{C_0} and we shall give canonical generators of a quantum analogue of

J_{C_0} .

Set

$$\psi_n = Y_{i_4^n} Y_{j_4^n} - q Y_{i_3^n} Y_{j_3^n} + q^2 Y_{i_2^n} Y_{j_2^n} - q^3 Y_{i_1^n} Y_{j_1^n},$$

for $1 \leq n \leq 10$. Recall that $\mathbf{A}(n) = (i_4^n, i_3^n, i_2^n, i_1^n, j_1^n, j_2^n, j_3^n, j_4^n)$. Using the formulas $(Rn,1)$, $(Rn,2)$, we can write $\psi_n = Y_{j_4^n} Y_{i_4^n} - q^{-1} Y_{j_3^n} Y_{i_3^n} + q^{-2} Y_{j_2^n} Y_{i_2^n} - q^{-3} Y_{j_1^n} Y_{i_1^n}$.

Lemma 2.5. *We have*

$$\begin{aligned} \text{ad}(F_k)\psi_n &= \begin{cases} \psi_{n'} & \text{if there exists } n' \text{ such that } \delta_n + \alpha_k = \delta_{n'}, \\ 0 & \text{otherwise,} \end{cases} \\ \text{ad}(E_k)\psi_n &= \begin{cases} \psi_{n'} & \text{if there exists } n' \text{ such that } \delta_n - \alpha_k = \delta_{n'}, \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

for $k \in I$, and

$$\text{ad}(K_k)\psi_n = q^{-(\delta_n, \alpha_k)} \psi_n$$

for $k \in I_0$.

Proof. Let $(k, n, n') \in \mathcal{A}$. We shall show $\text{ad}(F_k)\psi_n = \psi_{n'}$. If the condition (P_m^+) is satisfied, then we have $\text{ad}(F_k)Y_{i_m^n} = 0$, $Y_{i_m^{n'}} = Y_{i_m^n}$, $\text{ad}(K_k)Y_{i_m^n} = Y_{i_m^n}$, $\text{ad}(F_k)Y_{j_m^n} = Y_{j_m^{n'}}$. Hence

$$\text{ad}(F_k)(Y_{i_m^n} Y_{j_m^n}) = (\text{ad}(F_k)Y_{i_m^n})Y_{j_m^n} + (\text{ad}(K_k)Y_{i_m^n})(\text{ad}(F_k)Y_{j_m^n}) = Y_{i_m^{n'}} Y_{j_m^{n'}}.$$

If the condition (P_m^-) is satisfied, then we have $\text{ad}(F_k)Y_{i_m^n} = Y_{i_m^{n'}}$, $\text{ad}(F_k)Y_{j_m^n} = 0$. Hence $\text{ad}(F_k)(Y_{i_m^n} Y_{j_m^n}) = Y_{i_m^{n'}} Y_{j_m^{n'}}$ similarly. Therefore we have $\text{ad}(F_k)\psi_n = \psi_{n'}$.

Next we prove $\text{ad}(E_k)\psi_{n'} = \psi_n$. We have $\text{ad}(E_k)Y_{i_m^{n'}} = 0$, $\text{ad}(E_k)Y_{j_m^{n'}} = Y_{j_m^n}$ if the condition (P_m^+) is satisfied, and we have $\text{ad}(E_k)Y_{i_m^{n'}} = Y_{i_m^n}$, $\text{ad}(K_k^{-1})Y_{j_m^{n'}} = Y_{j_m^{n'}}$, $j_m^{n'} = j_m^n$, $\text{ad}(E_k)Y_{j_m^{n'}} = 0$ if the condition (P_m^-) is satisfied. Hence we have

$$\text{ad}(E_k)(Y_{i_m^{n'}} Y_{j_m^{n'}}) = (\text{ad}(E_k)Y_{i_m^{n'}})(\text{ad}(K_k^{-1})Y_{j_m^{n'}}) + Y_{i_m^{n'}}(\text{ad}(E_k)Y_{j_m^{n'}}) = Y_{i_m^n} Y_{j_m^n}$$

for $1 \leq m \leq 4$. Therefore we have $\text{ad}(E_k)\psi_{n'} = \psi_n$.

In other 50 cases, where $\delta_n + \alpha_k \notin \{\delta_l \mid 1 \leq l \leq 10\}$, we can check $\text{ad}(F_k)\psi_n = 0$ by a case-by-case consideration as follows.

In the 10 cases where there exists n' satisfying $\text{ad}(F_k)\psi_{n'} = \psi_n$, $((k, n) = (6, 2), (5, 3), (3, 4), (2, 5), (4, 6), (2, 7), (4, 7), (3, 8), (5, 9), (6, 10))$, we have $\text{ad}(F_k)Y_{i_m^n} = \text{ad}(F_k)Y_{j_m^n} = 0$ for $1 \leq m \leq 4$, and hence the assertion is obvious.

In the 8 cases $(k, n) = (5, 1), (6, 3), (6, 4), (6, 5), (6, 6), (6, 7), (6, 8), (5, 10)$, we have $\text{ad}(F_k)Y_{i_m^n} = \text{ad}(F_k)Y_{j_m^n} = 0$ for $m = 3, 4$, $\text{ad}(F_k)Y_{i_2^n} = Y_{j_1^n}$, $\text{ad}(F_k)Y_{j_2^n} = 0$,

$\text{ad}(F_k)Y_{i_1^n} = Y_{j_2^n}$, $\text{ad}(F_k)Y_{j_1^n} = 0$, and hence $\text{ad}(F_k)(Y_{i_2^n}Y_{j_2^n}) = Y_{j_1^n}Y_{j_2^n}$, $\text{ad}(F_k)(Y_{i_1^n}Y_{j_1^n}) = Y_{j_2^n}Y_{j_1^n}$. Thus we have $\text{ad}(F_k)\psi_n = q^2(Y_{j_1^n}Y_{j_2^n} - qY_{j_2^n}Y_{j_1^n}) = 0$ by Proposition 2.3.

In the remaining 32 cases there exists $m' \in \{2, 3, 4\}$ such that $\text{ad}(F_k)Y_{i_m^n} = 0$ ($m \neq m'$), $\text{ad}(F_k)Y_{j_m^n} = 0$ ($m \neq m' - 1$), $\text{ad}(F_k)Y_{i_{m'}^n} = Y_{i_{m'-1}^n}$, $\text{ad}(F_k)Y_{j_{m'-1}^n} = Y_{j_{m'}^n}$, $\text{ad}(K_k)Y_{i_{m'-1}^n} = q^{-1}Y_{i_{m'-1}^n}$. Then we have $\text{ad}(F_k)(Y_{i_m^n}Y_{j_{m'}^n}) = Y_{i_{m'-1}^n}Y_{j_{m'}^n}$, $\text{ad}(F_k)(Y_{i_{m'-1}^n}Y_{j_{m'-1}^n}) = q^{-1}Y_{i_{m'-1}^n}Y_{j_{m'-1}^n}$, $\text{ad}(F_k)\psi_n = q^{4-m'}(1 - qq^{-1})Y_{i_{m'-1}^n}Y_{j_{m'}^n} = 0$.

The weight $\beta_{i_m^n} + \beta_{j_m^n}$ does not depend on m . Hence we have $\text{ad}(K_k)\psi_n = q^{-(\delta_n, \alpha_k)}\psi_n$ where $\delta_n = \beta_{i_m^n} + \beta_{j_m^n}$.

Finally we show $\text{ad}(E_k)\psi_n = 0$ if $\delta_n - \alpha_k \notin \{\delta_l \mid 1 \leq l \leq 10\}$. We can check $\text{ad}(E_k)\psi_1 = 0$ for any $k = 2, 3, \dots, 6$ directly. It follows that $\sum_{n=1}^{10} \mathbb{C}(q)\psi_n = U_q(\mathfrak{l}_l)\psi_1$ and hence $\sum_{n=1}^{10} \mathbb{C}(q)\psi_n$ is an $\text{ad} U_q(\mathfrak{l}_l)$ -stable subspace with weights in $\{-\delta_l \mid 1 \leq l \leq 10\}$. Therefore we have $\text{ad}(E_k)\psi_n = 0$ if $\delta_n - \alpha_k \notin \{\delta_l \mid 1 \leq l \leq 10\}$. \square

Proposition 2.6. $\sum_{n=1}^{10} \mathbb{C}(q)\psi_n$ is an irreducible highest weight $U_q(\mathfrak{l}_l)$ -module with highest weight vector ψ_1 .

Proof. By Lemma 2.5 $\sum_{n=1}^{10} \mathbb{C}(q)\psi_n$ is a finite dimensional $U_q(\mathfrak{l}_l)$ -submodule generated by a highest weight vector ψ_1 with highest weight $-\delta_1$. Thus it is irreducible. \square

By [4] and Proposition 2.6 we obtain the following:

Theorem 2.7. A quantum analogue of the defining ideal J_{C_0} of the closure of the non-trivial non-open orbit C_0 is given by the two-sided ideal of A_q generated by $\{\psi_n \mid 1 \leq n \leq 10\}$.

3. Case of type E_7

Let \mathfrak{g} be a simple Lie algebra of type E_7 . We shall use the labelling of the vertices of the Dynkin diagram 1. Hence we have $I_0 = \{1, 2, 3, 4, 5, 6, 7\}$. Set $I = \{2, 3, 4, 5, 6, 7\}$. In this case we have $\mathfrak{n}_I^+ \neq \{0\}$, $[\mathfrak{n}_I^+, \mathfrak{n}_I^+] = \{0\}$. Then \mathfrak{l}_I is isomorphic to $\mathbb{C} \oplus \mathfrak{g}_{E_6}$, where \mathfrak{g}_{E_6} is a Lie algebra of type E_6 over \mathbb{C} , and \mathfrak{n}_I^+ is a 27-dimensional irreducible prehomogeneous vector space. There are four L_I -orbits $\{0\}, C_1, C_2, O$ on \mathfrak{n}_I^+ satisfying $\{0\} \subset \overline{C_1} \subset \overline{C_2} \subset \overline{O}$. Let $J_{C_1} \subset \mathbb{C}[\mathfrak{n}_I^+]$ be the defining ideal of the closure of C_1 , and let $J_{C_1}^0$ denote the subspace of J_{C_1} consisting of the polynomials in J_{C_1} with homogeneous degree 2. Then $J_{C_1}^0$ is a 27-dimensional irreducible \mathfrak{l}_I -module, and it generates the ideal J_{C_1} . Let $J_{C_2} \subset \mathbb{C}[\mathfrak{n}_I^+]$ be the defining ideal of the closure of C_2 , and let $J_{C_2}^0$ denote the subspace of J_{C_2} consisting of the polynomials in J_{C_2} with homogeneous degree 3. Then $J_{C_2}^0$ is a one-dimensional irreducible \mathfrak{l}_I -module generated by the irreducible relative invariant, and it generates the ideal J_{C_2} .

We fix a reduced expression

$$w_I w_0 = s_1 s_2 s_3 s_4 s_5 s_6 s_4 s_3 s_2 s_1 s_7 s_6 s_4 s_3 s_5 s_4 s_6 s_7 s_2 s_3 s_4 s_6 s_5 s_4 s_3 s_2 s_1$$

of $w_I w_0$ and define the elements Y_i ($i \in \Lambda = \{1, 2, \dots, 27\}$) as in Section 1.

Set $I'_0 = \{1, 2, 3, 4, 5, 6\}$, $I' = \{2, 3, 4, 5, 6\}$, $\Lambda' = \{1, 2, \dots, 10\}$. Then $\{\alpha_i\}_{i \in I'_0}$ is a set of simple roots of type D_6 . Let \mathfrak{g}' be the simple subalgebra of \mathfrak{g} corresponding to I'_0 . We choose a reduced expression $w_{I'} w_{I'_0} = s_1 s_2 s_3 s_4 s_5 s_6 s_4 s_3 s_2 s_1$ of $w_{I'} w_{I'_0}$. The elements Y_i ($i \in \Lambda'$) can be computed inside $U_q(\mathfrak{g}')$.

Let $\beta_j = \sum_{i \in I_0} m_i^j \alpha_i$ and set $\mathbf{m}^j = (m_1^j, \dots, m_7^j)$ for $j \in \Lambda$. Then we have

$$\begin{aligned} \mathbf{m}^1 &= (1, 0, 0, 0, 0, 0, 0), & \mathbf{m}^2 &= (1, 1, 0, 0, 0, 0, 0), & \mathbf{m}^3 &= (1, 1, 1, 0, 0, 0, 0), \\ \mathbf{m}^4 &= (1, 1, 1, 1, 0, 0, 0), & \mathbf{m}^5 &= (1, 1, 1, 1, 1, 0, 0), & \mathbf{m}^6 &= (1, 1, 1, 1, 0, 1, 0), \\ \mathbf{m}^7 &= (1, 1, 1, 1, 1, 1, 0), & \mathbf{m}^8 &= (1, 1, 1, 2, 1, 1, 0), & \mathbf{m}^9 &= (1, 1, 2, 2, 1, 1, 0), \\ \mathbf{m}^{10} &= (1, 2, 2, 2, 1, 1, 0), & \mathbf{m}^{11} &= (1, 1, 1, 1, 0, 1, 1), & \mathbf{m}^{12} &= (1, 1, 1, 1, 1, 1, 1), \\ \mathbf{m}^{13} &= (1, 1, 1, 2, 1, 1, 1), & \mathbf{m}^{14} &= (1, 1, 2, 2, 1, 1, 1), & \mathbf{m}^{15} &= (1, 1, 1, 2, 1, 2, 1), \\ \mathbf{m}^{16} &= (1, 1, 2, 2, 1, 2, 1), & \mathbf{m}^{17} &= (1, 1, 2, 3, 1, 2, 1), & \mathbf{m}^{18} &= (1, 1, 2, 3, 2, 2, 1), \\ \mathbf{m}^{19} &= (1, 2, 2, 2, 1, 1, 1), & \mathbf{m}^{20} &= (1, 2, 2, 2, 1, 2, 1), & \mathbf{m}^{21} &= (1, 2, 2, 3, 1, 2, 1), \\ \mathbf{m}^{22} &= (1, 2, 2, 3, 2, 2, 1), & \mathbf{m}^{23} &= (1, 2, 3, 3, 1, 2, 1), & \mathbf{m}^{24} &= (1, 2, 3, 3, 2, 2, 1), \\ \mathbf{m}^{25} &= (1, 2, 3, 4, 2, 2, 1), & \mathbf{m}^{26} &= (1, 2, 3, 4, 2, 3, 1), & \mathbf{m}^{27} &= (1, 2, 3, 4, 2, 3, 2). \end{aligned}$$

If $(\beta_j, \alpha_k) = -1$ for $j \in \Lambda$ and $k \in I$, then $s_k(\beta_j) = \beta_j + \alpha_k \in \Delta^+ \setminus \Delta_I$ and there exists $l \in \Lambda$ satisfying $\beta_j + \alpha_k = \beta_l$. Conversely if $\beta_j, \beta_l \in \Delta^+ \setminus \Delta_I$ satisfying $\beta_l - \beta_j = \alpha_k$ ($k \in I$), then we have $(\beta_j, \alpha_k) = -1$, $s_k(\beta_j) = \beta_l$.

For $k \in I$, $j \in \Lambda$, we have $\beta_j - 2\alpha_k, \beta_j + 2\alpha_k \notin \Delta^+ \setminus \Delta_I$.

Set

$$\mathcal{B} = \{(k, j, l) \in I \times \Lambda \times \Lambda \mid \beta_j + \alpha_k = \beta_l\}.$$

We have

$$\begin{aligned} \mathcal{B} = \{ & (2, 1, 2), (3, 2, 3), (4, 3, 4), (5, 4, 5), (6, 4, 6), (6, 5, 7), (5, 6, 7), (4, 7, 8), (3, 8, 9), \\ & (2, 9, 10), (7, 6, 11), (7, 7, 12), (7, 8, 13), (7, 9, 14), (7, 10, 19), (5, 11, 12), \\ & (4, 12, 13), (3, 13, 14), (6, 13, 15), (6, 14, 16), (3, 15, 16), (4, 16, 17), (5, 17, 18), \\ & (2, 14, 19), (2, 16, 20), (2, 17, 21), (2, 18, 22), (6, 19, 20), (4, 20, 21), (5, 21, 22), \\ & (3, 21, 23), (3, 22, 24), (5, 23, 24), (4, 24, 25), (6, 25, 26), (7, 26, 27)\}. \end{aligned}$$

In particular, we have $|\mathcal{B}| = 36$.

Lemma 3.1. *Let $\beta, \beta' \in \Delta^+ \setminus \Delta_I$ satisfying $\beta + \alpha_k = \beta'$ ($k \in I$). Then we can choose a reduced expression $w_I w_0 = s_{i_1} s_{i_2} \cdots s_{i_{27}}$ and $p \in \Lambda$ satisfying*

$$\begin{aligned} \beta &= s_{i_1} s_{i_2} \cdots s_{i_{p-1}}(\alpha_{i_p}), & \beta' &= s_{i_1} s_{i_2} \cdots s_{i_{p-1}} s_{i_p}(\alpha_{i_{p+1}}), & (\alpha_{i_p}, \alpha_{i_{p+1}}) &= -1, \\ \alpha_k &= s_{i_1} s_{i_2} \cdots s_{i_{p-1}}(\alpha_{i_{p+1}}). \end{aligned}$$

Proof. The 21 triplets (k, j, l) in \mathcal{B} satisfy $l = j + 1$, $(\alpha_{ij}, \alpha_{i,j+1}) = -1$. Therefore it is sufficient to deal with the remaining 15 cases. In the cases $(k, j, l) = (6, 4, 6), (6, 5, 7), (6, 13, 15), (6, 14, 16), (3, 21, 23), (3, 22, 24)$, we can take

$$w_l w_0 = s_1 s_2 s_3 s_4 s_6 s_5 s_4 s_3 s_2 s_1 s_7 s_6 s_4 s_5 s_3 s_4 s_6 s_7 s_2 s_3 s_4 s_5 s_6 s_4 s_3 s_2 s_1$$

with $p = 4, 6, 13, 15, 21, 23$, and in the cases $(k, j, l) = (7, 6, 11), (7, 7, 12), (7, 8, 13), (7, 9, 14), (7, 10, 19)$, we can take

$$w_l w_0 = s_1 s_2 s_3 s_4 s_5 s_6 s_7 s_4 s_6 s_3 s_4 s_2 s_3 s_1 s_2 s_5 s_4 s_6 s_7 s_3 s_4 s_6 s_5 s_4 s_3 s_2 s_1$$

with $p = 6, 8, 10, 12, 14$, and in the cases $(k, j, l) = (2, 14, 19), (2, 16, 20), (2, 17, 21), (2, 18, 22)$, we can take

$$w_l w_0 = s_1 s_2 s_3 s_4 s_5 s_6 s_4 s_3 s_2 s_1 s_7 s_6 s_4 s_5 s_3 s_2 s_4 s_3 s_6 s_4 s_7 s_6 s_5 s_4 s_3 s_2 s_1$$

with $p = 15, 17, 19, 21$. □

We can show the following similarly to the case E_6 . We omit the details.

Lemma 3.2. For $k \in I, j \in \Lambda$, we have

$$\begin{aligned} \text{ad}(F_k)Y_j &= \begin{cases} Y_l & \text{if there exists } (k, j, l) \in \mathcal{B}, \\ 0 & \text{otherwise,} \end{cases} \\ \text{ad}(E_k)Y_j &= \begin{cases} Y_l & \text{if there exists } (k, l, j) \in \mathcal{B}, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

The $U_q(\mathfrak{l}_I)$ -module $\bigoplus_{j \in \Lambda} \mathbb{C}(q)Y_j$ is an irreducible highest weight module with highest weight vector Y_1 and lowest weight vector Y_{27} . Hence, for any $1 \leq m \leq 26$, there exists a sequence $((k_1, n'_1, n_1), \dots, (k_s, n'_s, n_s))$ of \mathcal{B} satisfying $n_1 = 27, n'_s = m, n'_j = n_{j+1} (1 \leq j \leq s - 1)$.

Next we shall consider relations among the elements Y_i . We can write

$$Y_i Y_j = \sum_{\substack{s \leq i \\ \beta_i + \beta_j = \beta_s + \beta_t}} a_{s,t}^{i,j} Y_s Y_t \quad (a_{s,t}^{i,j} \in \mathbb{C}(q))$$

for $i > j$ (see [4]). Hence if $\beta_i + \beta_j$ does not have another decomposition $\beta + \beta'$ ($\beta, \beta' \in \Delta^+ \setminus \Delta_I, \beta_i + \beta_j = \beta + \beta'$) then we have $Y_i Y_j = a_{i,j} Y_j Y_i$ for some $a_{i,j} \in \mathbb{C}(q)$. Set $\delta = 2\varpi_1 = 3\alpha_1 + 4\alpha_2 + 5\alpha_3 + 6\alpha_4 + 3\alpha_5 + 4\alpha_6 + 2\alpha_7$, where ϖ_1 is the fundamental weight corresponding to α_1 . We denote a set of weights of the 27-dimensional irreducible highest weight \mathfrak{l}_I -module $J_{C_1}^0$ with highest weight $-\beta_1 - \beta_{10}$ by Γ . Set $\gamma_n = \delta - \beta_n (n \in \Lambda)$, and we have $\Gamma = \{-\gamma_n \mid n \in \Lambda\}$. For $\beta, \beta' \in \Delta^+ \setminus \Delta_I$ a weight $\beta + \beta'$ has another decomposition if and only if we have $-(\beta + \beta') \in \Gamma$. For each $n \in \Lambda$ there

exist exactly five pairs $(i, j) \in \Lambda^2$ such that $i < j$, $\beta_i + \beta_j = \gamma_n$. We denote them by $(i_1^n, j_1^n), (i_2^n, j_2^n), (i_3^n, j_3^n), (i_4^n, j_4^n), (i_5^n, j_5^n) \in \Lambda^2$ where $i_5^n < i_4^n < i_3^n < i_2^n < i_1^n$, $j_1^n < j_2^n < j_3^n < j_4^n < j_5^n$, and i_1^n, j_1^n satisfy the following condition (P_1^+) or (P_1^-) . Set

- $\mathbf{B}(n) = (i_5^n, i_4^n, i_3^n, i_2^n, i_1^n, j_1^n, j_2^n, j_3^n, j_4^n, j_5^n) \in \Lambda^{10}$ ($n \in \Lambda$). Then we have
- $\mathbf{B}(1) = (10, 19, 20, 21, 23, 22, 24, 25, 26, 27)$, $\mathbf{B}(2) = (9, 14, 16, 17, 23, 18, 24, 25, 26, 27)$,
 - $\mathbf{B}(3) = (8, 13, 15, 17, 21, 18, 22, 25, 26, 27)$, $\mathbf{B}(4) = (7, 12, 15, 16, 20, 18, 22, 24, 26, 27)$,
 - $\mathbf{B}(5) = (6, 11, 15, 16, 20, 17, 21, 23, 26, 27)$, $\mathbf{B}(6) = (5, 12, 13, 14, 19, 18, 22, 24, 25, 27)$,
 - $\mathbf{B}(7) = (4, 11, 13, 14, 19, 17, 21, 23, 25, 27)$, $\mathbf{B}(8) = (3, 11, 12, 14, 19, 16, 20, 23, 24, 27)$,
 - $\mathbf{B}(9) = (2, 11, 12, 13, 19, 15, 20, 21, 22, 27)$, $\mathbf{B}(10) = (1, 11, 12, 13, 14, 15, 16, 17, 18, 27)$,
 - $\mathbf{B}(11) = (5, 7, 8, 9, 10, 18, 22, 24, 25, 26)$, $\mathbf{B}(12) = (4, 6, 8, 9, 10, 17, 21, 23, 25, 26)$,
 - $\mathbf{B}(13) = (3, 6, 7, 9, 10, 16, 20, 23, 24, 26)$, $\mathbf{B}(14) = (2, 6, 7, 8, 10, 15, 20, 21, 22, 26)$,
 - $\mathbf{B}(15) = (3, 4, 5, 9, 10, 14, 19, 23, 24, 25)$, $\mathbf{B}(16) = (2, 4, 5, 8, 10, 13, 19, 21, 22, 25)$,
 - $\mathbf{B}(17) = (2, 3, 5, 7, 10, 12, 19, 20, 22, 24)$, $\mathbf{B}(18) = (2, 3, 4, 6, 10, 11, 19, 20, 21, 23)$,
 - $\mathbf{B}(19) = (1, 6, 7, 8, 9, 15, 16, 17, 18, 26)$, $\mathbf{B}(20) = (1, 4, 5, 8, 9, 13, 14, 17, 18, 25)$,
 - $\mathbf{B}(21) = (1, 3, 5, 7, 9, 12, 14, 16, 18, 24)$, $\mathbf{B}(22) = (1, 3, 4, 6, 9, 11, 14, 16, 17, 23)$,
 - $\mathbf{B}(23) = (1, 2, 5, 7, 8, 12, 13, 15, 18, 22)$, $\mathbf{B}(24) = (1, 2, 4, 6, 8, 11, 13, 15, 17, 21)$,
 - $\mathbf{B}(25) = (1, 2, 3, 6, 7, 11, 12, 15, 16, 20)$, $\mathbf{B}(26) = (1, 2, 3, 4, 5, 11, 12, 13, 14, 19)$,
 - $\mathbf{B}(27) = (1, 2, 3, 4, 5, 6, 7, 8, 9, 10)$.

For $n \in \Lambda$ we denote the set $\{i_5^n, i_4^n, i_3^n, i_2^n, i_1^n, j_1^n, j_2^n, j_3^n, j_4^n, j_5^n\}$ by $|\mathbf{B}(n)|$. For any $i, j \in \Lambda$ there exists $n \in \Lambda$ satisfying $i, j \in |\mathbf{B}(n)|$.

For $(k, n', n) \in \mathcal{B}$ and $m \in \{1, 2, 3, 4, 5\}$, we have either

$$(P_m^+) \quad (\beta_{i_m^n}, \alpha_k) = 0, i_m^{n'} = i_m^n, (\beta_{j_m^n}, \alpha_k) = -1, \beta_{j_m^{n'}} = \beta_{j_m^n} + \alpha_k$$

or

$$(P_m^-) \quad (\beta_{i_m^n}, \alpha_k) = -1, \beta_{i_m^{n'}} = \beta_{i_m^n} + \alpha_k, (\beta_{j_m^n}, \alpha_k) = 0, j_m^{n'} = j_m^n.$$

Proposition 3.3. For any $i, j \in \Lambda$ satisfying $i < j$, we have

$$(Q7) \quad Y_i Y_j = \begin{cases} Y_j Y_i & \text{if there exists } n \in \Lambda \text{ such that } \{i, j\} = \{i_1^n, j_1^n\}, \\ Y_{j_2^n} Y_{i_2^n} + (q - q^{-1}) Y_{i_1^n} Y_{j_1^n} & \text{if there exists } n \in \Lambda \text{ such that } i = i_2^n, j = j_2^n, \\ Y_{j_m^n} Y_{i_m^n} + q Y_{j_{m-1}^n} Y_{i_{m-1}^n} - q^{-1} Y_{i_{m-1}^n} Y_{j_{m-1}^n} & \text{if there exist } n \in \Lambda, m \in \{3, 4, 5\} \text{ such that } i = i_m^n, j = j_m^n, \\ q Y_j Y_i & \text{otherwise.} \end{cases}$$

Proof. Since there exists $n \in \Lambda$ satisfying $i, j \in |\mathbf{B}(n)|$ for any $i, j \in \Lambda$, it is

sufficient to show

$$\begin{aligned}
 (\mathbf{Rn}) \quad & \begin{cases} Y_{i_1^n} Y_{j_1^n} = Y_{j_1^n} Y_{i_1^n} & (\mathbf{Rn}, 1) \\ Y_{i_m^n} Y_{j_m^n} = Y_{j_m^n} Y_{i_m^n} + q Y_{j_{m-1}^n} Y_{i_{m-1}^n} - q^{-1} Y_{i_{m-1}^n} Y_{j_{m-1}^n} \quad (2 \leq m \leq 5) & (\mathbf{Rn}, 2) \\ Y_{l_1} Y_{l_2} = q Y_{l_2} Y_{l_1} & (\mathbf{Rn}, 3) \end{cases} \\
 & (l_1, l_2 \in |\mathbf{B}(n)|, l_1 < l_2, \{l_1, l_2\} \neq \{i_m^n, j_m^n\} \quad (1 \leq m \leq 5))
 \end{aligned}$$

for $n \in \Lambda$ and $1 \leq m \leq 5$.

When $n = 27$, the elements Y_i ($1 \leq i \leq 10$) satisfy the same relations as those for type D_6 , and hence relations (R27) hold.

Since there exists a sequence $((k_1, n'_1, n_1), \dots, (k_s, n'_s, n_s))$ of \mathcal{B} satisfying $n_1 = 27, n'_s = m, n'_j = n_{j+1}$ ($1 \leq j \leq s - 1$) for any $1 \leq m \leq 26$, it is sufficient to show (\mathbf{Rn}') for $(k, n', n) \in \mathcal{B}$ assuming (\mathbf{Rn}) . This is proved similarly to Proposition 2.3. Details are omitted. \square

By [4] and Proposition 3.3 we obtain the following:

Theorem 3.4. *The formulas (Q7) give fundamental relations for the generator system $\{Y_i\}_{i \in \Lambda}$ of the algebra $A_q = U_q(\mathfrak{n}^-)$.*

We shall construct a quantum deformation of the lowest degree part $J_{C_1}^0$ of the defining ideal J_{C_1} and we shall give canonical generators of a quantum deformation of J_{C_1} .

Set

$$\psi_n = Y_{i_5^n} Y_{j_5^n} - q Y_{i_4^n} Y_{j_4^n} + q^2 Y_{i_3^n} Y_{j_3^n} - q^3 Y_{i_2^n} Y_{j_2^n} + q^4 Y_{i_1^n} Y_{j_1^n},$$

for $n \in \Lambda$, where $\mathbf{B}(n) = (i_5^n, i_4^n, i_3^n, i_2^n, i_1^n, j_1^n, j_2^n, j_3^n, j_4^n, j_5^n)$. Using the formulas $(\mathbf{Rn}, 1), (\mathbf{Rn}, 2)$, we can write

$$\psi_n = Y_{j_5^n} Y_{i_5^n} - q^{-1} Y_{j_4^n} Y_{i_4^n} + q^{-2} Y_{j_3^n} Y_{i_3^n} - q^{-3} Y_{j_2^n} Y_{i_2^n} + q^{-4} Y_{j_1^n} Y_{i_1^n}.$$

Similarly to Lemma 2.5 and Proposition 2.6 we can show the following:

Lemma 3.5. *We have*

$$\begin{aligned}
 \text{ad}(F_k)\psi_n &= \begin{cases} \psi_{n'} & \text{if there exists } (k, n', n) \in \mathcal{B}, \\ 0 & \text{otherwise,} \end{cases} \\
 \text{ad}(E_k)\psi_n &= \begin{cases} \psi_{n'} & \text{if there exists } (k, n, n') \in \mathcal{B}, \\ 0 & \text{otherwise} \end{cases}
 \end{aligned}$$

for $k \in I$, and

$$\text{ad}(K_k)\psi_n = q^{-(\gamma_n, \alpha_k)}\psi_n$$

for $k \in I_0$.

Proposition 3.6. $\sum_{n \in \Lambda} \mathbb{C}(q)\psi_n$ is an irreducible highest weight $U_q(\mathfrak{l}_1)$ -module with highest weight vector ψ_{27} .

By [4] and Proposition 3.6 we obtain the following:

Theorem 3.7. A quantum deformation of the defining ideal J_{C_1} of the closure of the non-open orbit C_1 is given by the two-sided ideal of A_q generated by $\{\psi_n \mid n \in \Lambda\}$.

Set

$$\varphi = \sum_{n \in \Lambda} (-q)^{|\beta_n|-1} Y_n \psi_n,$$

where $|\beta| = \sum_{i \in I_0} m_i$ ($\beta = \sum_{i \in I_0} m_i \alpha_i$).

Proposition 3.8. $\mathbb{C}(q)\varphi$ is a one-dimensional $U_q(\mathfrak{l}_1)$ -module.

Proof. By Proposition 3.3 we can check that the coefficient $a_{1,10,27}$ of $Y_1 Y_{10} Y_{27}$ in $\varphi = \sum_{i < j < k} a_{ijk} Y_i Y_j Y_k$ is $1 + q^8 + q^{16}$. Therefore we have $\varphi \neq 0$.

Let $(k, n, n') \in \mathcal{B}$. Then we have $|\beta_{n'}| = |\beta_n| + 1$, $\text{ad}(F_k)Y_n = Y_{n'}$, $\text{ad}(F_k)Y_{n'} = 0$, $\text{ad}(F_k)\psi_{n'} = \psi_n$, $\text{ad}(F_k)\psi_n = 0$, $(\beta_{n'}, \alpha_k) = 1$. Hence $\text{ad}(F_k)(Y_n \psi_n - q Y_{n'} \psi_{n'}) = Y_{n'} \psi_n - q q^{-1} Y_{n'} \psi_n = 0$. Therefore we have $\text{ad}(F_k)\varphi = 0$ for any $k \in I$, and similarly we have $\text{ad}(E_k)\varphi = 0$ for any $k \in I$. Since $\gamma_n + \beta_n = \delta$ for any $n \in \Lambda$, we have $\text{ad}(K_k)\varphi = q^{-(\delta, \alpha_k)}\varphi$ for any $k \in I_0$. In particular, we have $\text{ad}(K_k)\varphi = \varphi$ for any $k \in I$, and $\text{ad}(K_1)\varphi = q^{-2}\varphi$. □

The element φ is a quantum deformation of the irreducible relative invariant on the prehomogeneous vector space.

Theorem 3.9. A quantum deformation of the defining ideal J_{C_2} of the closure of the non-open orbit C_2 is given by the two-sided ideal of A_q generated by φ .

References

- [1] V.G. Drinfel'd: *Hopf algebra and the Yang-Baxter equation*, Soviet Math. Dokl. **32** (1985), 254–258.
- [2] M. Hashimoto and T. Hayashi: *Quantum multilinear algebra*, Tohoku Math. J. **44** (1992), 471–521.
- [3] M. Jimbo: *A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985), 63–69.
- [4] A. Kamita, Y. Morita and T. Tanisaki: *Quantum deformations of certain prehomogeneous vector spaces I*, Hiroshima Math. J. **28** (1998), 527–540.
- [5] A. Kamita: *Quantum deformations of certain prehomogeneous vector spaces III*, Hiroshima Math. J. **30** (2000), 79–115.
- [6] G. Lusztig: *Quantum groups at roots of 1*, Geometriae Dedicata, **35** (1990), 89–114.
- [7] M. Noumi, H. Yamada and K. Mimachi: *Finite dimensional representations of the quantum group $GL_q(n; \mathbb{C})$ and the zonal spherical functions*, Japan J. Math. **19** (1993), 31–80.
- [8] E. Strickland: *Classical invariant theory for the quantum symplectic group*, Adv. Math. **123** (1996), 78–90.
- [9] T. Tanisaki: *Highest weight modules associated to parabolic subgroups with commutative unipotent radicals*, in Algebraic Groups and their Representations, R.W. Carter and J. Saxt (eds.), Proceedings of the NATO ASI conference, Kluwer Academic Publishers, Dordrecht, 1998, 73–90.
- [10] E. Taft and J. Towber: *Quantum deformation of flag schemes and Grassmann schemes. I. A q -deformation of the shape-algebra for $GL(n)$* , J. Algebra, **142** (1991), 1–36.

Department of Mathematics
Faculty of Science
Hiroshima University
Higashi-Hiroshima, 739-8526,
Japan
e-mail: morita@math.sci.hiroshima-u.ac.jp

