

Title	An estimate of the ribbon number by the Jones polynomial
Author(s)	Mizuma, Yoko
Citation	Osaka Journal of Mathematics. 2006, 43(2), p. 365-369
Version Type	VoR
URL	https://doi.org/10.18910/4957
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

AN ESTIMATE OF THE RIBBON NUMBER BY THE JONES POLYNOMIAL

Dedicated to Professor Yukio Matsumoto on the occasion of his 60th birthday

YOKO MIZUMA

(Received March 18, 2005)

Abstract

For a ribbon knot we define the notion of its ribbon number. In this paper we estimate the ribbon number for a ribbon knot by using the Jones polynomial. As a corollary we determine the ribbon number of the Kinoshita-Terasaka knot.

1. Introduction

To investigate the complexity of a ribbon knot, we define the notion of the ribbon number. This obvious measure of a ribbon knot's complexity is often hard to determine. In fact, even in a simple case of the Kinoshita-Terasaka knot, its ribbon number is hard to determine. In this paper, we estimate the ribbon number by using a formula for the first derivative at -1 of the Jones polynomial of a ribbon knot of 1-fusion in [6]. As a corollary we determine the ribbon number of the Kinoshita-Terasaka knot.

1.1. Definitions and theorems.

DEFINITION 1.1. A *ribbon disk* is an immersed 2-disk of D^2 into S^3 with only transverse double points such that the singular set consists of ribbon singularities, that is, the preimage of each ribbon singularity consists of a properly embedded arc in D^2 and an embedded arc interior to D^2 (see Fig. 1). A knot is a *ribbon knot* if it bounds a ribbon disk in S^3 (cf. [3], [4]).

For a ribbon knot we define its ribbon number as follows.

DEFINITION 1.2. The *ribbon number* of a ribbon knot is defined as the minimal number of ribbon singularities needed for a ribbon disk bounded by the ribbon knot.

Here we have some remarks of Definition 1.2.




Fig. 1. A ribbon disk with two ribbon singularities

REMARK 1.3. A ribbon knot whose ribbon number is zero is a trivial knot and there does not exist a ribbon knot whose ribbon number is one.

REMARK 1.4. The ribbon number of a ribbon knot K is greater than or equal to the genus of K ([1]).

Now we can state the following theorem.

Theorem 1.5. *Let K be a ribbon knot satisfying $\Delta_K(t) = 1$ and $J'_K(-1) \neq 0$, where $\Delta_K(t)$ is the Alexander polynomial of K and $J'_K(-1)$ is the first derivative at -1 of the Jones polynomial of K . Then the ribbon number of K is greater than or equal to three.*

To state the next theorem we review a ribbon knot of 1-fusion as follows.

DEFINITION 1.6. We call a knot K in S^3 a ribbon knot of 1-fusion, if it has a knot diagram described in Fig. 2, where n is even and each small rectangle named C_i is determined by $c_i \in \{-1, 0, 1\}$ ($i = 1, 2, \dots, n$) and there are disjointly embedded $(n+1)$ bands in the big rectangle, being knotted, twisted and mutually linked (cf. [6], [5]). The diagram is called 1-fusion diagram of K and gives a ribbon disk bounded by K .

REMARK 1.7. A ribbon knot of 1-fusion is a band sum of 2-component trivial link and vice versa.

Here we can state the following theorem.




Fig. 2. A 1-fusion diagram




Fig. 3. $(1, 1, 0, -1)$




Fig. 4. The Kinoshita-Terasaka knot

Theorem 1.8. *If K has a 1-fusion diagram with $(c_1, c_2, c_3, c_4) = (1, 1, 0, -1)$ as shown in Fig. 3, where $J'_K(-1) \neq 0$, then the ribbon number of K is three.*

1.2. An application of theorems. Now we consider the ribbon number of the Kinoshita-Terasaka knot, which has a 1-fusion diagram as in Fig. 4 and $\Delta(t) = 1$ and $J'(-1) = 48$. Hence we obtain the following theorem from Theorem 1.8.

Theorem 1.9. *The ribbon number of the Kinoshita-Terasaka knot is three.*

Note that the genus of the Kinoshita-Terasaka knot is two ([2]).

2. Proof

Now we start to prove theorems.

Proof of Theorem 1.5. Let K be a ribbon knot satisfying $\Delta_K(t) = 1$ and $J'_K(-1) \neq 0$. K is not a trivial knot, so the ribbon number of K is greater than or equal to two. Note that a ribbon disk with two ribbon singularities bounded by a non-trivial knot is bounded by a ribbon knot which has one of eight 1-fusion diagrams as shown in Fig. 5, where C_i is determined by $c_i \in \{-1, 1\}$ ($i = 1, 2, 3$). By using a formula for the Alexander polynomial of a ribbon knot of 1-fusion in [5], $\Delta(t) \neq 1$ for each knot in the left side of Fig. 5 and $\Delta(t) = 1$ for each knot in the right side of Fig. 5. By using a formula for the first derivative at -1 of the Jones polynomial of a ribbon knot of 1-fusion, $J'(-1) = 0$ for each knot in the right side of Fig. 5 (Example 1.12 in [6]). Hence the ribbon number of K is not two. This completes the proof. \square

Proof of Theorem 1.8. By using a formula for the Alexander polynomial of a ribbon knot of 1-fusion in [5], a knot K which has a 1-fusion diagram as shown in


Fig. 5. (c_1, c_2) and $(0, c_2, c_3, 0)$

Fig. 3 satisfies $\Delta_K(t) = 1$. By Theorem 1.5, the ribbon number of K is greater than or equal to three. The diagram in Fig. 3 gives a ribbon disk bounded by K which has three ribbon singularities, hence the ribbon number of K is three. \square

ACKNOWLEDGMENTS. I am grateful to Professor Akio Kawauchi for suggesting me this problem of ribbon number.

References

- [1] R.H. Fox: *Characterizations of slices and ribbons*, Osaka J. Math. **10** (1973) 69–76.
- [2] D. Gabai: *Genera of the arborescent links*, Mem. Amer. Math. Soc. **59** (1986) 1–98.
- [3] L.H. Kauffman: *On Knots*, Ann. of Math. Stud. **115**, Princeton Univ. Press, Princeton, NJ, 1987.
- [4] A. Kawauchi: *A Survey of Knot Theory*, Birkhäuser Verlag, 1996.
- [5] Y. Marumoto: *On ribbon 2-knots of 1-fusion*, Math. Sem. Notes Kobe Univ. **5** (1977), 59–68.
- [6] Y. Mizuma: *Ribbon knots of 1-fusion, the Jones polynomial and the Casson-Walker invariant*, Rev. Mat. Complut. **18** (2005), 387–425.
- [7] H. Terasaka: *On null-equivalent knots*, Osaka Math. J. **11** (1959), 95–113.

Research Institute for Mathematical Sciences
 Kyoto University
 Sakyo-ku, Kyoto 606-8502, Japan
 e-mail: mizuma@kurims.kyoto-u.ac.jp

The Fields Institute for Research in Mathematical Sciences
 222 College Street, Toronto, Ontario, M5T 3J1, Canada
 e-mail: ymizuma@fields.utoronto.ca