



Title	GENERATORS OF THE HOMOLOGICAL GOLDMAN LIE ALGEBRA
Author(s)	Kawazumi, Nariya; Kuno, Yusuke; Toda, Kazuki
Citation	Osaka Journal of Mathematics. 2014, 51(3), p. 665-671
Version Type	VoR
URL	<a href="https://doi.org/10.18910/50814">https://doi.org/10.18910/50814</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## GENERATORS OF THE HOMOLOGICAL GOLDMAN LIE ALGEBRA

NARIYA KAWAZUMI\*, YUSUKE KUNO and KAZUKI TODA

(Received November 28, 2012)

### Abstract

We determine the minimum number of generators of the homological Goldman Lie algebra of a surface consisting of elements of the first homology group of the surface.

### 1. Introduction

Let  $\Sigma$  be a compact connected oriented surface of genus  $g \geq 1$ . The first homology group  $H = H_1(\Sigma, \mathbb{Z})$  of  $\Sigma$  is equipped with a skew-symmetric bilinear form  $\langle -, - \rangle: H \times H \rightarrow \mathbb{Z}$  called the *intersection form*. We denote by  $\mathbb{Q}H$  the  $\mathbb{Q}$ -vector space with basis the set  $H$ ;

$$\mathbb{Q}H := \left\{ \sum_{i=1}^n c_i [x_i] \mid n \in \mathbb{N}, c_i \in \mathbb{Q}, x_i \in H \right\},$$

where  $[-]: H \rightarrow \mathbb{Q}H$  is the embedding as basis. Let  $[-, -]: \mathbb{Q}H \times \mathbb{Q}H \rightarrow \mathbb{Q}H$  be a bilinear form defined by  $[[x], [y]] := \langle x, y \rangle [x + y]$  for  $x, y \in H$ . It is easy to see that this bilinear form is skew-symmetric and satisfies the Jacobi identity. We call the  $\mathbb{Q}$ -vector space  $\mathbb{Q}H$  equipped with the Lie bracket  $[-, -]$  the *homological Goldman Lie algebra of  $\Sigma$* . This Lie algebra was originally introduced by Goldman [1] pp. 295–297.

The purpose of this paper is to study generators of  $\mathbb{Q}H$ . In the previous paper [3], the third-named author determined the ideals of  $\mathbb{Q}H$ . In particular, it follows that the abelianization of  $\mathbb{Q}H$  is finite dimensional if and only if the intersection form  $\langle -, - \rangle$  is non-degenerate; otherwise  $\mathbb{Q}H$  is not finitely generated as a Lie algebra. On the other hand, the abelianization of  $\mathbb{Z}H$  is not finitely generated even in the non-degenerate case.

Hereafter we assume that  $\langle -, - \rangle$  is non-degenerate, i.e.,  $\Sigma$  is closed or has one boundary component. Then there exists a  $\mathbb{Z}$ -basis  $\{A_i, B_i\}_{i=1}^g$  of  $H$ , called a *symplectic*

---

2010 Mathematics Subject Classification. 57N05, 17B65.

\*Partially supported by the Grant-in-Aid for Scientific Research (A) (No. 20244003), (A) (No. 22244005) and (B) (No. 24340010) from the Japan Society for Promotion of Sciences.

basis of  $H$ , such that

$$\langle A_i, B_j \rangle = \delta_{ij}, \quad \langle A_i, A_j \rangle = \langle B_i, B_j \rangle = 0$$

for all  $i, j \in \{1, \dots, g\}$ , where  $\delta_{ij}$  is Kronecker's delta. Throughout this paper, we fix a symplectic basis of  $H$ .

We will show that if  $\langle -, - \rangle$  is non-degenerate, then the Lie algebra  $\mathbb{Q}H$  is finitely generated. Moreover, we determine the minimum number of generators of  $\mathbb{Q}H$  consisting of elements of  $H$ . Our main theorem is the following.

**Theorem 1.1.** *Suppose the intersection form  $\langle -, - \rangle$  is non-degenerate. There is a subset  $S$  of  $H$  such that  $\{[s] \mid s \in S\}$  generates  $\mathbb{Q}H$  as a Lie algebra and  $\#S = 2g + 2$ . In particular, the Lie algebra  $\mathbb{Q}H$  is finitely generated. Moreover, if  $S$  is a subset of  $H$  and  $\{[s] \mid s \in S\}$  generates  $\mathbb{Q}H$  as a Lie algebra, we have  $\#S \geq 2g + 2$ .*

**REMARK 1.2.** As we see in Lemma 2.2 and Theorem 2.3,  $\mathbb{Q}H$  is generated by primitive elements of  $H$  and  $[0]$ . By Theorem 3.2, if  $S$  is a subset of  $H$  and  $\{[s] \mid s \in S\}$  generates  $\mathbb{Q}H$ , then  $0 \in S$ . Therefore, the quotient Lie algebra  $\mathbb{Q}H/\mathbb{Q}[0]$  is generated by  $2g + 1$  primitive elements of  $H$ , and  $2g + 1$  is the minimum number of generators of  $\mathbb{Q}H/\mathbb{Q}[0]$  consisting of elements of  $H$ . This result reminds us a result of Humphries [2] that the mapping class group of a closed oriented surface is generated by  $2g + 1$  Dehn twists, and  $2g + 1$  is the minimum number of generators consisting of Dehn twists. Although we do not see any relationship between Humphries's result and our result, this coincidence seems interesting.

On the other hand, there is another Lie algebra associated to the oriented surface  $\Sigma$  called the *Goldman Lie algebra*. This Lie algebra is denoted by  $\mathbb{Q}\hat{\pi}$ , where  $\hat{\pi}$  is the set of homotopy classes of oriented loops on  $\Sigma$ . For more details, see Goldman [1]. The natural projection  $\hat{\pi} \rightarrow H$  induces a surjective Lie algebra homomorphisms  $\mathbb{Q}\hat{\pi} \rightarrow \mathbb{Q}H$  and  $\mathbb{Q}\hat{\pi}/\mathbb{Q}1 \rightarrow \mathbb{Q}H/\mathbb{Q}[0]$ . Here  $1$  is the homotopy class of a constant loop. Simple closed curves on  $\Sigma$  are analogous to primitive elements of  $H$ . However, in contrast with the above property of  $\mathbb{Q}H$ , the quotient Lie algebra  $\mathbb{Q}\hat{\pi}/\mathbb{Q}1$  is not generated by simple closed curves. This follows from the fact that the kernel of the Turaev cobracket [4] is a proper Lie subalgebra of  $\mathbb{Q}\hat{\pi}/\mathbb{Q}1$ , and simple closed curves are in the kernel of the Turaev cobracket. It is not known whether the Goldman Lie algebra  $\mathbb{Q}\hat{\pi}$  for a surface which is closed or has one boundary component is finitely generated or not.

The rest of this paper is devoted to the proof of Theorem 1.1.

## 2. Upper bound of the number of generators

We define the  $\mathbb{Q}$ -linear map  $ad: \mathbb{Q}H \rightarrow \text{End}(\mathbb{Q}H)$  by  $ad(X)(Y) := [X, Y]$ .

**Lemma 2.1.** *Fix  $k \in \{1, \dots, g\}$ . Let  $\mathfrak{g}$  be a Lie subalgebra of  $\mathbb{Q}H$  such that  $[\pm A_k], [\pm B_k] \in \mathfrak{g}$ . If  $X \in H$  satisfies  $\langle A_k, X \rangle = 0$ ,  $\langle B_k, X \rangle = 0$  and  $[X + A_k] \in \mathfrak{g}$ , then we have  $[X + aA_k + bB_k] \in \mathfrak{g}$  for all  $(a, b) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$ .*

Proof. First of all, we have  $[X + aA_k] \in \mathfrak{g}$  for all  $a \in \mathbb{Z} \setminus \{0\}$  since

$$ad([-B_k]) ad([A_k])^{a-1} ad([B_k])([X + A_k]) = -a[X + aA_k] \quad \text{if } a > 0,$$

and

$$ad([B_k]) ad([-A_k])^{-a+1} ad([-B_k])([X + A_k]) = -a[X + aA_k] \quad \text{if } a < 0.$$

Similarly, we have  $[X + bB_k] \in \mathfrak{g}$  for all  $b \in \mathbb{Z} \setminus \{0\}$  since

$$ad([-A_k]) ad([B_k])^b([X + A_k]) = (-1)^{b+1}b[X + bB_k] \quad \text{if } b > 0,$$

and

$$ad([-A_k]) ad([-B_k])^{-b}([X + A_k]) = -b[X + bB_k] \quad \text{if } b < 0.$$

Therefore we have  $[X + aA_k + bB_k] \in \mathfrak{g}$  if  $ab = 0$  and  $(a, b) \neq (0, 0)$ .

Suppose  $ab \neq 0$ . By what we have just proved, we have  $[X + aA_k] \in \mathfrak{g}$ . Then we have  $[X + aA_k + bB_k] \in \mathfrak{g}$  since

$$ad([B_k])^b([X + aA_k]) = (-a)^b[X + aA_k + bB_k] \quad \text{if } b > 0,$$

and

$$ad([-B_k])^{-b}([X + aA_k]) = a^{-b}[X + aA_k + bB_k] \quad \text{if } b < 0.$$

This completes the proof. □

**Lemma 2.2.** *The set*

$$\{[\pm A_i] \ (1 \leq i \leq g), [\pm B_i] \ (1 \leq i \leq g), [A_i + A_j] \ (1 \leq i < j \leq g), [0]\}$$

*generates  $\mathbb{Q}H$  as a Lie algebra. In particular,  $\mathbb{Q}H$  is finitely generated.*

Proof. Let  $\mathfrak{g}$  be the Lie subalgebra generated by the above set.

**Claim 1.** *For any integer  $n > 0$  and indices  $i_1, \dots, i_n$  with  $1 \leq i_1 < \dots < i_n \leq g$ , we have  $[A_{i_1} + \dots + A_{i_n}] \in \mathfrak{g}$ .*

We prove Claim 1 by induction on  $n$ . If  $n = 1$  or  $n = 2$ , the claim follows from the assumption of the lemma. Suppose  $n > 2$  and let  $i_1, \dots, i_n$  be indices with  $1 \leq$

$i_1 < \dots < i_n \leq g$ . By the inductive assumption, we have  $[A_{i_1} + \dots + A_{i_{n-1}}] \in \mathfrak{g}$ . Since

$$\begin{aligned} & ad([-B_{i_1}]) ad([-A_{i_1}]) ad([A_{i_1} + A_{i_n}]) ad([B_{i_1}]) ([A_{i_1} + \dots + A_{i_{n-1}}]) \\ & = [A_{i_1} + \dots + A_{i_n}], \end{aligned}$$

we have  $[A_{i_1} + \dots + A_{i_n}] \in \mathfrak{g}$ . This proves Claim 1.

Let  $x \in H$  be an arbitrary element. If  $x = 0$ , we have  $[x] \in \mathfrak{g}$  by the assumption of the lemma. Suppose  $x \neq 0$ . Since  $\{A_i, B_i\}_{i=1}^g$  is a  $\mathbb{Z}$ -basis of  $H$ , we can write  $x = a_1 A_{i_1} + b_1 B_{i_1} + \dots + a_n A_{i_n} + b_n B_{i_n}$  with  $1 \leq i_1 < \dots < i_n \leq g$  and  $(a_1, b_1), \dots, (a_n, b_n) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$ .

**Claim 2.** *We have  $[a_1 A_{i_1} + b_1 B_{i_1} + \dots + a_m A_{i_m} + b_m B_{i_m} + A_{i_{m+1}} + \dots + A_{i_n}] \in \mathfrak{g}$  for all  $m = 1, \dots, n$ .*

We prove Claim 2 by induction on  $m$ . Suppose  $m = 1$ . We have  $[A_{i_2} + \dots + A_{i_n}] \in \mathfrak{g}$  by Claim 1. Applying Lemma 2.1 to  $k = i_1$  and  $X = A_{i_2} + \dots + A_{i_n}$ , we have  $[a_1 A_{i_1} + b_1 B_{i_1} + A_{i_2} + \dots + A_{i_n}] \in \mathfrak{g}$ . This proves the case  $m = 1$ .

Suppose  $m > 1$ . By the inductive assumption, we have  $[a_1 A_{i_1} + b_1 B_{i_1} + \dots + a_{m-1} A_{i_{m-1}} + b_{m-1} B_{i_{m-1}} + A_{i_{m+1}} + \dots + A_{i_n}] \in \mathfrak{g}$ . Applying Lemma 2.1 to  $k = i_m$  and  $X = a_1 A_{i_1} + b_1 B_{i_1} + \dots + a_{m-1} A_{i_{m-1}} + b_{m-1} B_{i_{m-1}} + A_{i_{m+1}} + \dots + A_{i_n}$ , we obtain  $[a_1 A_{i_1} + b_1 B_{i_1} + \dots + a_m A_{i_m} + b_m B_{i_m} + A_{i_{m+1}} + \dots + A_{i_n}] \in \mathfrak{g}$ . This proves Claim 2.

Applying Claim 2 to  $m = n$ , we have  $[x] \in \mathfrak{g}$ . Since  $\{[x] \mid x \in H\}$  is a  $\mathbb{Q}$ -basis of  $\mathbb{Q}H$ , we obtain  $\mathfrak{g} = \mathbb{Q}H$ . This completes the proof.  $\square$

Now we give generators of  $\mathbb{Q}H$  consisting of  $2g + 2$  elements of  $H$ .

**Theorem 2.3.** *The set*

$$\{[A_i] \ (1 \leq i \leq g), [B_i] \ (1 \leq i \leq g), [-A_1 - \dots - A_g - B_1 - \dots - B_g], [0]\}$$

generates  $\mathbb{Q}H$  as a Lie algebra.

**Proof.** Let  $\mathfrak{g}$  be the Lie subalgebra generated by the above set. Set  $X := [-A_1 - \dots - A_g - B_1 - \dots - B_g]$ .

First of all, we have  $[A_1 + \dots + A_g], [B_1 + \dots + B_g] \in \mathfrak{g}$  since

$$ad([B_1]) \cdots ad([B_g]) ad([A_1])^2 \cdots ad([A_g])^2 (X) = (-1)^g [A_1 + \dots + A_g],$$

and

$$ad([A_1]) \cdots ad([A_g]) ad([B_1])^2 \cdots ad([B_g])^2 (X) = [B_1 + \dots + B_g].$$

Next, we have  $[-A_i], [-B_i] \in \mathfrak{g}$  for all  $i = 1, \dots, g$  since

$$\begin{aligned} & ad([B_1 + \dots + B_g]) ad([A_1]) \dots ad([A_{i-1}]) ad([A_{i+1}]) \dots ad([A_g])(X) \\ &= (-1)^{g-1}[-A_i], \end{aligned}$$

and

$$\begin{aligned} & ad([A_1 + \dots + A_g]) ad([B_1]) \dots ad([B_{i-1}]) ad([B_{i+1}]) \dots ad([B_g])(X) \\ &= -[-B_i]. \end{aligned}$$

Finally, we have  $[A_i + A_j] \in \mathfrak{g}$  for  $1 \leq i < j \leq g$  since

$$ad([B_1 + \dots + B_g]) ad([A_1 + \dots + A_g]) ad([A_i]) ad([A_j])(X) = 2g[A_i + A_j].$$

Now the assertion follows from Lemma 2.2.  $\square$

### 3. Lower bound of the number of generators

**Lemma 3.1.** *Let  $S$  be a subset of  $H$ ,  $\mathfrak{g}$  the Lie subalgebra generated by  $\{[s] \mid s \in S\}$ , and  $M$  the submonoid in  $H$  generated by  $S$ , i.e.,*

$$M = \{s_1 + \dots + s_n \in H \mid n \in \mathbb{N}, s_i \in S\}.$$

*Then, we have  $\mathfrak{g} \subset \mathbb{Q}M$ .*

Proof. The set  $\{[s_1, [s_2, [\dots, [s_{n-1}, s_n] \dots]]] \mid n \in \mathbb{N}, s_i \in S\}$  generates  $\mathfrak{g}$  as a  $\mathbb{Q}$ -vector space. Since

$$[s_1, [s_2, [\dots, [s_{n-1}, s_n] \dots]]] = \left( \prod_{i=1}^{n-1} \left\langle s_i, \sum_{j=i+1}^n s_j \right\rangle \right) [s_1 + \dots + s_n] \in \mathbb{Q}M,$$

we obtain  $\mathfrak{g} \subset \mathbb{Q}M$ .  $\square$

**Theorem 3.2.** *Let  $S$  be a subset of  $H$ . If  $\{[s] \mid s \in S\}$  generates  $\mathbb{Q}H$  as a Lie algebra, we have  $0 \in S$  and  $\#S \geq 2g + 2$ .*

Proof. Suppose  $\mathbb{Q}H$  is generated by  $\{[s] \mid s \in S\}$  as a Lie algebra. Now, we have  $[\mathbb{Q}H, \mathbb{Q}H] \subset \mathbb{Q}(H \setminus \{0\})$  since  $\langle x, y \rangle = 0$  for  $x, y \in H$  with  $x + y = 0$ . This implies that  $0 \in S$ .

Let  $M$  be the submonoid generated by  $S$ . By Lemma 3.1 we have  $\mathbb{Q}H \subset \mathbb{Q}M$ , thus  $H \subset M$ . Since  $H \supset M$ , we obtain  $H = M$ . In other words, the set  $S$  generates  $H$  as a monoid. In particular,  $S \setminus \{0\}$  generates  $H$  as a  $\mathbb{Z}$ -module. Since  $H$  is a free  $\mathbb{Z}$ -module of rank  $2g$ , we have  $\#(S \setminus \{0\}) \geq 2g$ .

If  $\#(S \setminus \{0\}) = 2g$ ,  $S \setminus \{0\}$  is a  $\mathbb{Z}$ -basis of  $H$ . Then  $M = \{\sum_{s \in S \setminus \{0\}} a_s s \mid a_s \geq 0\} \neq H$ , which contradicts  $M = H$ . Hence  $\#(S \setminus \{0\}) > 2g$ , therefore we have  $\#S \geq 2g + 2$ .  $\square$

By Theorems 2.3 and 3.2, we obtain Theorem 1.1.

**REMARK 3.3.** Let  $R$  be a commutative ring with unit. Using  $R$  instead of  $\mathbb{Q}$ , we can similarly define the homological Goldman Lie algebra  $RH$ . If  $R$  includes  $\mathbb{Q}$ , the same result as Theorem 1.1 holds for  $RH$ .

In Theorem 3.2, we assumed  $S \subset H$ . This condition is essential. If we consider generators which are not necessarily elements of  $H$ , we can find generators of  $\mathbb{Q}H$  whose number is less than  $2g + 2$ .

**Propositon 3.4.** *The Lie algebra  $\mathbb{Q}H$  is generated by  $g + 2$  elements  $[A_1], \dots, [A_g]$ ,  $[-A_1 - \dots - A_g - B_1 - \dots - B_g] + [B_1] + \dots + [B_g] + [0]$ , and  $[B_1 + \dots + B_g]$ .*

Proof. Let  $\mathfrak{g}$  be the Lie subalgebra generated by  $[A_1], \dots, [A_g]$ ,  $X := [-A_1 - \dots - A_g - B_1 - \dots - B_g] + [B_1] + \dots + [B_g] + [0]$ , and  $Y := [B_1 + \dots + B_g]$ .

Suppose  $g = 1$ . Then  $X = [-A_1 - B_1] + [B_1] + [0]$  and  $Y = [B_1]$ . We have  $[-A_1] \in \mathfrak{g}$  since  $[X, Y] = -[-A_1]$ , and we have  $[-B_1] \in \mathfrak{g}$  since  $[X - Y, [A_1]] = [-B_1]$ . Then we have  $[-A_1 - B_1] \in \mathfrak{g}$  since  $[[[-A_1], [-B_1]] = [-A_1 - B_1]$ . Since  $[0] = X - [-A_1 - B_1] - Y$  we have  $[0] \in \mathfrak{g}$ . Therefore we obtain  $\{[A_1], [B_1], [-A_1 - B_1], [0]\} \subset \mathfrak{g}$ , which implies  $\mathfrak{g} = \mathbb{Q}H$  by Theorem 2.3.

Suppose  $g \geq 2$ . Then  $(-1)^g [-B_1 - \dots - B_g] = ad([A_1]) \dots ad([A_g])(X) \in \mathfrak{g}$ . On the other hand, we have  $-g[-A_1 - \dots - A_g] = [X, Y] \in \mathfrak{g}$ . Set  $Z = [-A_1 - \dots - A_g - B_1 - \dots - B_g]$ . Then  $Z \in \mathfrak{g}$  since

$$[[-A_1 - \dots - A_g], [-B_1 - \dots - B_g]] = gZ.$$

We have  $[-A_i] \in \mathfrak{g}$  for  $i = 1, \dots, g$  since

$$ad(Y) ad([A_1]) \dots ad([A_{i-1}]) ad([A_{i+1}]) \dots ad([A_g])(Z) = (-1)^{g-1} [-A_i],$$

and we have  $[B_i] \in \mathfrak{g}$  for  $i = 1, \dots, g$  since

$$ad([-A_i]) ad([A_i])(X) + Z = -[B_i].$$

Finally, we have  $[0] \in \mathfrak{g}$  since  $[0] = X - Z - [B_1] - \dots - [B_g]$ . Hence, we have  $\mathfrak{g} = \mathbb{Q}H$  by Theorem 2.3.  $\square$

## References

- [1] W.M. Goldman: *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*, Invent. Math. **85** (1986), 263–302.
- [2] S.P. Humphries: *Generators for the mapping class group*; in Topology of Low-Dimensional Manifolds (Proc. Second Sussex Conf., Chelwood Gate, 1977), Lecture Notes in Math. **722**, Springer, Berlin, 1979, 44–47.
- [3] K. Toda: *The ideals of the homological Goldman Lie algebra*, to appear in Kodai Math. J., available at arXiv:1112.1213.
- [4] V.G. Turaev: *Skein quantization of Poisson algebras of loops on surfaces*, Ann. Sci. École Norm. Sup. (4) **24** (1991), 635–704.

Nariya Kawazumi  
 Department of Mathematical Sciences  
 University of Tokyo  
 3-8-1 Komaba Meguro-ku Tokyo 153-8914  
 Japan  
 e-mail: kawazumi@ms.u-tokyo.ac.jp

Yusuke Kuno  
 Department of Mathematics  
 Tsuda College  
 2-1-1 Tsuda-Machi Kodaira-shi Tokyo 187-8577  
 Japan  
 e-mail: kunotti@tsuda.ac.jp

Kazuki Toda  
 Tono Frontier High School  
 1127-8 Izumi-cho Kawai Toki-shi Gifu 509-5101  
 Japan  
 e-mail: ktoda@ms.u-tokyo.ac.jp