Title	High-resolution ($^3\text{He}, t$) reaction on the double- β decaying nucleus ^{136}Xe
Author(s)	Puppe, P.; Frekers, D.; Adachi, T. et al.
Citation	Physical Review C - Nuclear Physics. 84(5) p.51305-p.51305
Issue Date	2011-11-15
oa:version	VoR
URL	https://hdl.handle.net/11094/51205
rights	©2011 American Physical Society
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

High-resolution ($^3\text{He}, t$) reaction on the double- β decaying nucleus ^{136}Xe

P. Puppe,¹ D. Frekers,¹ T. Adachi,² H. Akimune,³ N. Aoi,⁴ B. Bilgier,⁵ H. Ejiri,⁴ H. Fujita,^{4,6} Y. Fujita,⁶ M. Fujiwara,⁴ E. Ganioglu,⁵ M. N. Harakeh,⁷ K. Hatanaka,⁴ M. Holl,¹ H. C. Kozer,⁵ J. Lee,⁸ A. Lennarz,¹ H. Matsubara,⁹ K. Miki,⁴ S. E. A. Orrigo,¹⁰ T. Suzuki,⁴ A. Tamii,⁴ and J. H. Thies¹

¹*Institut für Kernphysik, Westfälische Wilhelms-Universität, D-48149 Münster, Germany*

²*Research Center for Electron and Photon Science, Tohoku University, Sendai, Miyagi 982-0826, Japan*

³*Department of Physics, Konan University, Kobe 658-8501, Japan*

⁴*Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047, Japan*

⁵*Istanbul University, Science Faculty, Department of Physics, TR-34134 Vezneciler, Istanbul, Turkey*

⁶*Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan*

⁷*Kernfysisch Versneller Instituut, University of Groningen, NL-9747 AA Groningen, The Netherlands*

⁸*RIKEN Nishina Center, Wako, Saitama 351-0198, Japan*

⁹*Center for Nuclear Study, University of Tokyo, Hirosawa 2-1, Wako, Saitama 351-0198, Japan*

¹⁰*Instituto de Física Corpuscular, CSIC-Universidad de Valencia, E-46071 Valencia, Spain*

(Received 3 September 2011; published 15 November 2011)

A ($^3\text{He}, t$) charge-exchange reaction experiment on the double-beta decaying nucleus ^{136}Xe has been performed at an incident energy of 420 MeV with the objective to measure the Gamow-Teller (GT) strength distribution in ^{136}Cs . The measurements have been carried out at the dispersion-matched WS beam line and the Grand Raiden spectrometer of the Research Center for Nuclear Physics in Osaka, where an energy resolution of 42 keV was achieved. A new gas cell with thin windows made of polyethylene naphthalate has been employed as a target. The extracted GT strength distribution is confronted with the rather long $2\nu\beta\beta$ decay half-life of ^{136}Xe .

DOI: [10.1103/PhysRevC.84.051305](https://doi.org/10.1103/PhysRevC.84.051305)

PACS number(s): 25.55.Kr, 23.40.Hc, 27.60.+j

Introduction. In the context of the $\beta\beta$ decay, ^{136}Xe is a rather special system. From systematics, i.e., by looking at neighboring $\beta\beta$ decaying nuclei, its 2ν half-life would not be expected to be significantly longer than $T_{1/2}^{2\nu} \approx 10^{20}$ yr. This is a value which is mostly supported by theory as well [1–6], although in Ref. [7] some arguments for a longer half-life relating this to the $N = 82$ neutron shell closure have been provided. The relatively high $\beta\beta$ decay Q value ($Q = 2.467$ MeV) and the high Z value ($Z = 54$) are advantageous and ought to have a significant bearing on the value of the decay time. Yet, for a long time, the $2\nu\beta\beta$ decay has evaded detection [8–12], and only recently the EXO-200 collaboration successfully measured the half-life to $T_{1/2}^{2\nu} = [2.11 \pm 0.04(\text{stat.}) \pm 0.21(\text{sys.})] \times 10^{21}$ yr (90% C.L.) [13]. Accordingly, the value for the nuclear matrix element was quoted as $M_{\text{DGT}}^{2\nu} = 0.019 \text{ MeV}^{-1}$, which is the smallest one ever measured. This means that the various matrix elements, which connect the intermediate states in the ^{136}Cs intermediate nucleus must also either be small or there is some fortuitous phase cancellation among the various components at work, which suppresses the decay rather effectively. Experimental charge-exchange data on ^{136}Xe , which could shed light on the size of these matrix elements do not exist, and even more astounding, not even a single excited state in the intermediate nucleus ^{136}Cs is known [14]. Nonetheless, counting experiments on ^{136}Xe searching for the neutrinoless ($0\nu\beta\beta$) decay are planned as well, as, e.g., by the EXO collaboration, or more recently by KamLAND-ZeN [15–19]. Assuming that the $2\nu\beta\beta$ decays and the $0\nu\beta\beta$ decays are not strongly correlated, a low $2\nu\beta\beta$ decay rate is then rather advantageous for these experiments, as it would constitute

a much reduced background in the vicinity of the endpoint energy [20].

The present Rapid Communication aims at providing a much warranted experimental input into the structure of the $A = 136$ system by using the ($^3\text{He}, t$) charge-exchange reaction at intermediate energies. Charge-exchange reactions into the β^- direction [as, e.g., ($^3\text{He}, t$) or (p, n)] or β^+ direction [as, e.g., ($t, ^3\text{He}$), ($d, ^2\text{He}$) or (n, p)] are an established tool for extracting Gamow-Teller (GT) transition strengths and are therefore rather effective to provide information about the GT^- and GT^+ parts of the nuclear matrix elements of the $2\nu\beta\beta$ decay [21–33]. The selectivity is a result of the dominance of the $\vec{\sigma}\vec{\tau}$ effective nucleon-nucleon interaction at intermediate energies and low momentum transfers [34–37].

It may be instructive to recall the connection between the $2\nu\beta\beta$ decay nuclear matrix element and the $2\nu\beta\beta$ decay half-life:

$$(T_{1/2}^{2\nu})^{-1} = G^{2\nu}(Q, Z) |M_{\text{DGT}}^{2\nu}|^2 \quad (1)$$

and

$$M_{\text{DGT}}^{2\nu} = \sum_m \frac{M_m(\text{GT}^+) \cdot M_m(\text{GT}^-)}{\frac{1}{2} Q_{\beta\beta}(0_{g.s.}^{(f)}) + E_x(1_m^+) - E_0}. \quad (2)$$

Here, $E_x(1_m^+) - E_0$ is the energy difference between the m th intermediate 1^+ state and the initial ground state, and $M_m(\text{GT}^\pm)$ are the single β -decay matrix elements in the β^+ and β^- directions, respectively. [Note that the insertion of $M_m(\text{GT}^+)$ is the result of time invariance; also note that the energy denominator is often expressed in units of the electron

FIG. 2. Angular distributions of typical 1^+ , 2^- , and 0^+ (IAS) transitions. The error bars denote the combined statistical and systematic uncertainties. The lines illustrate DWBA calculations for the various components.

of ~ 42 keV was obtained for the ^{136}Xe target. This value is largely determined by the target areal thickness and its physical extent in the direction of the beam.

Analysis. Figure 1 shows three spectra of the $^{136}\text{Xe}(^3\text{He}, t)^{136}\text{Cs}$ reaction at three different forward scattering angles. The spectra have been arranged in such a way that the most forward-angle spectrum appears in the back and the others are stacked on top of each other using different color codings (black, white, and gray). This way the angular dependence of the various components can be quickly identified; e.g., forward-peaked cross sections are likely GT transitions (colored black) with angular momentum transfer $\Delta L = 0$ and more backward peaked ones are likely $\Delta L = 1$ spin-dipole transitions (colored gray). The spectra show a moderate number of isolated states up to ~ 4.5 MeV excitation energy, followed by the strongly excited isobaric analog state (IAS) at 13.38 MeV. At ~ 15 MeV one observes the peak of the giant GT resonance and at ~ 23 MeV the rather broad ($\Gamma \simeq 10$ MeV) spin-dipole resonance.

Angular distributions were extracted for the transitions indicated in Fig. 1 and compared with distorted-wave Born approximation (DWBA) calculations performed with the codes NORMOD [46] and FOLD [47]. Appropriate optical model

parameters were adapted from those of ^{90}Zr as quoted in Ref. [48]. Angular distributions for the two strongest 1^+ states at 0.59 and 3.52 MeV, the strong state at 1.00 MeV (assumed to have $J^\pi = 2^-$), and the IAS at 13.38 MeV are shown in Fig. 2 and compared with DWBA calculations. In order to arrive at a good description of the experimental angular distributions, we find that for all 1^+ states there is a need to add a significant $\Delta L = 2$ angular momentum transfer component to the cross section, which is something that may be related to the high density of unoccupied shells near the Fermi surface. This component should add coherently to the cross section, however, due to the lack of a realistic ^{136}Xe wave function, one is simply left to add it in an incoherent way. The procedure may receive some justification by the good description one is able to achieve.

There are numerous ways to extract the GT strength from the angular distribution. All of them relate the GT part of the cross section to the weak decay strength by extrapolating the angular distribution to zero momentum transfer ($q = 0$). The functional form of this extrapolation is given by the zeroth-order Bessel function $|j_0(qR_0)|^2$, with R_0 being the interaction radius (see, e.g., Ref. [37]). This extrapolation is in accord with the standard plane-wave approach and may not be an appropriate description for the hadronic reaction, as this requires incoming distorted Coulomb waves.

In the present Rapid Communication we have used a method whereby we combine the Gamow-Teller and Fermi unit cross sections at $q = 0$ to extract the GT strength:

$$R^2(E, A) = \frac{\hat{\sigma}_{\text{GT}}}{\hat{\sigma}_F} = \frac{\sigma_{\text{GT}}(q \rightarrow 0)}{B(\text{GT}^-)} \bigg/ \frac{\sigma_F(q \rightarrow 0)}{B(F)}. \quad (4)$$

Following Refs. [49–51], R^2 takes a value of 11.0 ± 1.0 in this mass range and $B(F) = 28$ for ^{136}Xe . The $B(\text{GT}^-)$ values extracted according to this recipe are listed in Table I. Because of the uncertainty with which the extra non-GT contribution can be evaluated at $q = 0$, we suppose that 50% of the non-GT part to the cross section at $q = 0$ (see, e.g., the second columns in Table I) should enter into the error calculation for the strength values. Such a conservative error margin would then also include possible non-GT tensor contributions to the cross

TABLE I. Excitation energies of 1^+ states populated in the $^{136}\text{Xe}(^3\text{He}, t)^{136}\text{Cs}$ reaction. The second columns indicate what percentage of the cross section at $q = 0$ can be attributed to a GT transition using the extraction procedure as described in the text. The extracted $B(\text{GT})$ values are given in columns 3.

E_x (MeV)	GT fraction	$B(\text{GT}^-)$	E_x (MeV)	GT fraction	$B(\text{GT}^-)$
0.59	87%	0.149(21)	2.55	66%	0.033(11)
0.85	81%	0.082(15)	2.60	78%	0.033(7)
1.91	72%	0.017(5)	2.71	66%	0.055(18)
2.01	66%	0.024(8)	2.81	71%	0.026(7)
2.29	73%	0.042(11)	2.91	58%	0.027(12)
2.36	72%	0.059(15)	3.42	59%	0.024(10)
2.45	74%	0.015(4)	3.52	74%	0.097(23)
2.50	66%	0.024(8)	$\Sigma = 0.71(1)$

FIG. 3. (a) Distribution of $B(GT^-)$ strength in the low excitation-energy region of the $^{136}\text{Xe}(^3\text{He},t)^{136}\text{Cs}$ reaction; (b) running sum of the $B(GT^-)$ strength.

section. The $B(GT)$ strength values up to ~ 4 MeV extracted in this way are displayed in Fig. 3 together with the running sum, which yields a value of $\sum_{0-3.5 \text{ MeV}} B(GT^-) = 0.71$.

Connection to $\beta\beta$ decay. In Refs. [23,24,29–32] it was reported that the low-lying GT^- and GT^+ parts of the GT strength (up to ~ 5 MeV) contribute a substantial fraction to the $\beta\beta$ nuclear matrix elements of Eq. (2). Although the higher-lying giant GT^- resonance is the most prominent component in typical (p, n) -type spectra [as is also nicely exemplified in Fig. 1 for the present $(^3\text{He}, t)$ reaction], the resonance is conspicuously absent for a charge-exchange reaction in the GT^+ direction [29–32], thereby contributing comparatively little to the nuclear matrix elements. This absence is, of course, a rather trivial effect of Pauli blocking, since in a nucleus with a large neutron excess the neutron shells in the vicinity of the Fermi surface always find a largely unoccupied proton shell, whereas this is not the case for protons.

Therefore, the moderately strong low-lying GT^- transitions appearing in the present ^{136}Xe charge-exchange spectra do not point to any special properties of ^{136}Xe in terms of an unusually strong suppression of GT strength. Several other possible scenarios may therefore be brought to bear for an explanation of the rather long half-life. Since the present $(^3\text{He}, t)$ reaction

only defines the GT^- part, one could speculate about an exceptionally weak GT^+ part, which certainly would be something one could extract from a similar charge-exchange reaction [here (n, p) type] on the $\beta\beta$ -decay daughter ^{136}Ba . Given the quoted number for the ^{136}Xe nuclear matrix element $M_{\text{DGT}}^{2\nu} = 0.019 \text{ MeV}^{-1}$ [13], the $B(GT^+)$ values for the states listed in Table I should then be on average in the range of 10^{-3} – 10^{-2} times the $B(GT^-)$ values. Such a situation would indeed be unexpected and special.

Alternatively, there is the possibility of a rather effective phase cancellation in Eq. (2), which is something already conjectured in Ref. [29] for the ^{48}Ca case. In Fig. 3, we note a clustering of GT transitions between 0 and 1 MeV and between 2 and 3.5 MeV. If the nuclear matrix elements have different signs for these two clusters, the elements under the sum in Eq. (2) could effectively cancel, especially if the $B(GT)$ values for the GT^+ and the GT^- directions were moderately correlated. This can also easily be read off from the running sum in the lower part of Fig. 3.

Whereas the first scenario could be readily tested experimentally, the second one requires a detailed theoretical understanding of what the underlying physics is, which would cause certain matrix elements to have different signs. This may be an important aspect, since theoretical quasiparticle random phase approximation (QRPA)-based calculations [1–3,52–57] often use the $2\nu\beta\beta$ decay half-life to adjust the particle-particle interaction parameter g_{pp} for predicting the $0\nu\beta\beta$ decay nuclear matrix element, irrespective of possible cancellation effects.

Conclusion. We have used a high-resolution $(^3\text{He}, t)$ charge-exchange reaction at 420 MeV on the $\beta\beta$ decaying nucleus ^{136}Xe to identify the GT^- strength distribution in the intermediate nucleus ^{136}Cs in an attempt to understand the long $2\nu\beta\beta$ decay half-life of ^{136}Xe . We find that the GT^- strength distribution even at low excitation energies exhibits a rather normal behavior. A number of well-isolated states up to ~ 4 MeV, which are concentrated in two separate clusters, have been resolved carrying a total $B(GT^-)$ strength of 0.71. We have argued that exceptionally weak transitions from the GT^+ side would be a rather improbable cause for the long half-life, whereas phase-cancellation effects for the $\beta\beta$ -decay nuclear matrix elements seem to be a more natural and likely scenario.

Acknowledgments. We thank the RCNP accelerator staff for their fine technical support during the course of the experiment. The generous financial support from the Directorate of RCNP is gratefully acknowledged. Y.F. and A.T. were partly supported by MEXT, Japan under Grant No. 22540310. S.E.A.O. was supported under the Spanish MICINN Grant No. FPA200806419-C02-01. The work was partly financed by the Deutsche Forschungsgemeinschaft (DFG) under Grant No. FR 601/3-1.

[1] P. Vogel and M. R. Zirnbauer, *Phys. Rev. Lett.* **57**, 3148 (1986).

[2] A. Staudt, K. Muto, and H. V. Klapdor-Kleingrothaus, *Europhys. Lett.* **13**, 31 (1990).

- [3] F. Šimkovic, A. Faessler, H. Mütter, V. Rodin, and M. Stauf, *Phys. Rev. C* **79**, 055501 (2009).
- [4] E. Caurier, F. Nowacki, A. Poves, and J. Retamosa, *Nucl. Phys. A* **654**, 973c (1999).
- [5] E. Caurier, F. Nowacki, and A. Poves, *Int. J. Mod. Phys. E* **16**, 552 (2007).
- [6] E. Caurier, J. Menéndez, F. Nowacki, and A. Poves, *Phys. Rev. Lett.* **100**, 052503 (2008).
- [7] H. Ejiri, *J. Phys. Soc. Jpn.* **78**, 074201 (2009).
- [8] E. Bellotti, O. Cremonesi, E. Fiorini, G. Gervasio, S. Ragazzi, and L. Rossi, *Phys. Lett. B* **266**, 193 (1991).
- [9] R. Luescher, J. Farine, F. Boehm, J. Busto, K. Gabathuler, and G. Gervasio, *Phys. Lett. B* **434**, 407 (1998).
- [10] R. Bernabei, P. Belli, F. Cappella, R. Cerulli, F. Montecchia, and A. Incicchitti, *Phys. Lett. B* **546**, 23 (2002).
- [11] J. M. Gavriljuk, V. V. Kuzminov, N. Ya. Osetrova, and S. S. Ratkevich, *Phys. Rev. C* **61**, 035501 (2000).
- [12] J. M. Gavriljuk, A. M. Gangapshev, V. V. Kuzminov, S. I. Panasenkov, and S. S. Ratkevich, *Phys. At. Nucl.* **69**, 2129 (2006).
- [13] N. Ackerman *et al.*, arXiv:1108.4193v1.
- [14] National Nuclear Data Center, Brookhaven National Laboratory (2011) [<http://www.nndc.bnl.gov>].
- [15] M. Danilov *et al.*, *Phys. Lett. B* **480**, 12 (2000).
- [16] D. Akimov, G. Bower, M. Breidenbach, R. Conley, and E. Conti, *Nucl. Phys. Proc. Suppl. B* **138**, 224 (2005).
- [17] R. Gornea, *J. Phys. Conf. Ser.* **259**, 012039 (2010).
- [18] EXO collaboration (2010) [<http://www-project.slac.stanford.edu/exo/>].
- [19] Y. Efremenko, talk given at MEDEX'11-Conference on Matrix Elements for the Double-beta decay Experiments, Prague, June 13–16, 2011 (unpublished).
- [20] H. Ejiri, *Prog. Part. Nucl. Phys.* **40**, 307 (1998).
- [21] W. P. Alford *et al.*, *Nucl. Phys. A* **514**, 49 (1990).
- [22] H. Akimune *et al.*, *Phys. Lett. B* **394**, 23 (1997).
- [23] H. Ejiri, *Phys. Rep.* **338**, 265 (2000).
- [24] H. Ejiri, *J. Phys. Soc. Jpn.* **74**, 2101 (2005).
- [25] G. W. Hitt *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **566**, 264 (2006).
- [26] S. Rakers *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **481**, 253 (2002).
- [27] S. Rakers *et al.*, *Phys. Rev. C* **70**, 054302 (2004).
- [28] S. Rakers *et al.*, *Phys. Rev. C* **71**, 054313 (2005).
- [29] E.-W. Grewe *et al.*, *Phys. Rev. C* **76**, 054307 (2007).
- [30] E.-W. Grewe *et al.*, *Phys. Rev. C* **77**, 064303 (2008).
- [31] E.-W. Grewe *et al.*, *Phys. Rev. C* **78**, 044301 (2008).
- [32] H. Dohmann *et al.*, *Phys. Rev. C* **78**, 041602 (2008).
- [33] C. J. Guess *et al.*, *Phys. Rev. C* **83**, 064318 (2011).
- [34] W. G. Love and M. A. Franey, *Phys. Rev. C* **24**, 1073 (1981).
- [35] M. A. Franey and W. G. Love, *Phys. Rev. C* **31**, 488 (1985).
- [36] C. D. Goodman *et al.*, *Phys. Rev. Lett.* **44**, 1755 (1980).
- [37] T. N. Taddeucci *et al.*, *Nucl. Phys. A* **469**, 125 (1987).
- [38] J. Suhonen and O. Civitarese, *Phys. Rep.* **300**, 123 (1998).
- [39] T. Wasaka *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **482**, 79 (2002).
- [40] M. Fujiwara *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **422**, 484 (1999).
- [41] Y. Fujita *et al.*, *Nucl. Instrum. Methods Phys. Res. B* **126**, 274 (1997).
- [42] H. Fujita *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **469**, 55 (2001).
- [43] H. Fujita *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **484**, 17 (2002).
- [44] H. Matsubara, A. Tamii, T. Adachi, M. Dozono, H. Fujita, and K. Fujita, *Nucl. Instrum. Methods* (to be submitted, 2011).
- [45] G. Audi and M. Wang (private communication).
- [46] M. A. Hofstee *et al.*, *Nucl. Phys. A* **588**, 729 (1995).
- [47] J. Cook and J. A. Carr, computer code FOLD, Florida State University, 1982 (unpublished).
- [48] J. Kamiya *et al.*, *Phys. Rev. C* **67**, 064612 (2003).
- [49] T. Adachi, Ph.D. thesis, Department of Physics, Osaka University, 2007.
- [50] T. Adachi *et al.*, *Nucl. Phys. A* **788**, 70c (2007).
- [51] Y. Fujita *et al.*, in RCNP, Annual Report 2010.
- [52] V. A. Rodin, A. Faessler, F. Šimkovic, and P. Vogel, *Phys. Rev. C* **68**, 044302 (2003).
- [53] V. A. Rodin, A. Faessler, F. Šimkovic, and P. Vogel, *Nucl. Phys. A* **766**, 107 (2006); **793**, 213(E) (2007).
- [54] F. Šimkovic, A. Faessler, V. A. Rodin, P. Vogel, and J. Engel, *Phys. Rev. C* **77**, 045503 (2008).
- [55] J. Suhonen, *Phys. Lett. B* **607**, 87 (2005).
- [56] J. Suhonen and O. Civitarese, *Phys. Lett. B* **626**, 80 (2005).
- [57] J. Suhonen and O. Civitarese, *Nucl. Phys. A* **761**, 313 (2005).