

Title	Adams operations in the connective K-theory of compact Lie groups
Author(s)	Watanabe, Takashi
Citation	Osaka Journal of Mathematics. 1986, 23(3), p. 617-632
Version Type	VoR
URL	https://doi.org/10.18910/5180
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

ADAMS OPERATIONS IN THE CONNECTIVE K-THEORY OF COMPACT LIE GROUPS

TAKASHI WATANABE

(Received May 25, 1985)

1. Introduction

Let G be a compact, 1-connected, simple Lie group of rank 2 or 3. That is, G is one of the following:

$$SU(3), Sp(2), G_2, SU(4), Spin(7) \text{ and } Sp(3).$$

In [14], for these groups G , we have given a complete description of the Chern character ([7, §1])

$$ch: K^*(G) \rightarrow H^*(G; \mathbb{Q}).$$

Using this, one can easily compute the Adams operations ψ^r ([1]) on $K^*(G)$ for all $r \in \mathbb{Z}$ (see (2.5)).

Throughout this paper p will denote an odd prime. Let us introduce some spectra ([4, Part III]). Let $\mathbf{K}Z_{(p)}$ denote the ring spectrum representing complex K -theory localized at p . Let $\mathbf{k}Z_{(p)}$ be its (-1) -connected cover. So there is a map of ring spectra $\kappa: \mathbf{k}Z_{(p)} \rightarrow \mathbf{K}Z_{(p)}$ such that

$$\kappa_*: \pi_*(\mathbf{k}Z_{(p)}) = Z_{(p)}[u] \rightarrow \pi_*(\mathbf{K}Z_{(p)}) = Z_{(p)}[u, u^{-1}]$$

satisfies $\kappa_*(u) = u$ where $|u| = 2$. As is well known, there is a ring spectrum $\mathbf{g}(p)$ such that

$$\mathbf{k}Z_{(p)} \simeq \bigvee_{i=0}^{p-2} \Sigma^{2i} \mathbf{g}(p).$$

Here the injection $\iota: \mathbf{g}(p) \rightarrow \mathbf{k}Z_{(p)}$ is a map of ring spectra such that

$$\iota_*: \pi_*(\mathbf{g}(p)) = Z_{(p)}[v] \rightarrow \pi_*(\mathbf{k}Z_{(p)}) = Z_{(p)}[u]$$

satisfies $\iota_*(v) = u^{p-1}$ where $|v| = 2(p-1)$. For r prime to p there are maps of ring spectra

$$\begin{aligned} \psi^r: \mathbf{K}Z_{(p)} &\rightarrow \mathbf{K}Z_{(p)}, \\ \psi^r: \mathbf{k}Z_{(p)} &\rightarrow \mathbf{k}Z_{(p)}, \\ \psi^r: \mathbf{g}(p) &\rightarrow \mathbf{g}(p) \end{aligned}$$

which are called the stable Adams operations ([6], [5]). They commute with κ , ι and satisfy $\psi^r(u)=ru$. Let

$$\theta_r: \mathbf{g}(p) \rightarrow \Sigma^{2(p-1)}\mathbf{g}(p)$$

be a unique map of spectra such that $(v \cdot)\theta_r \simeq \psi^r - 1$ where $v \cdot: \Sigma^{2(p-1)}\mathbf{g}(p) \rightarrow \mathbf{g}(p)$ is multiplication by v . We denote by $\mathbf{j}(p; r)$ the fibre spectrum of θ_r . If r or r' generates the group of units of Z/p^2 , then $\mathbf{j}(p; r) \simeq \mathbf{j}(p; r')$. In this case, we may write $\mathbf{j}(p)$ for $\mathbf{j}(p; r)$ and use a suitable r to discuss it. $\mathbf{j}(p)$ is known to be a ring spectrum (see [13]).

Let $\widetilde{j(p)}_i(G)$ (resp. $\widetilde{j(p)}^i(G)$) be the i -th reduced $\mathbf{j}(p)$ -homology (resp. cohomology) group of G . One of our targets is to compute the groups $\widetilde{j(p)}_i(G)$ for all the above G and p . As will be mentioned in §3, the cases $(G, p)=(G_2, 3)$, $(Sp(3), 3)$ are most interesting. Then we obtain

Theorem 1.1. *For $i \leq 21$ and $G=G_2, Sp(3)$ the groups $\widetilde{j(3)}_i(G)$ are listed in the following table:*

$i \backslash G$	0	1	2	3	4	5	6	7	8	9
G_2	0	0	0	$Z_{(3)}$	0	0	$Z/3$	0	0	0
$Sp(3)$	0	0	0	$Z_{(3)}$	0	0	0	$Z_{(3)}$	0	0
$i \backslash G$	10	11	12	13		14		15	16	
G_2	0	$Z_{(3)}$	0	0		$Z/3^3 \oplus Z_{(3)}$		0	0	
$Sp(3)$	$Z/3 \oplus Z_{(3)}$	$Z_{(3)}$	0	$Z/3$		$Z/3 \oplus Z/3^3 \oplus Z_{(3)}$		0	0	
$i \backslash G$	17	18	19	20		21				
G_2	$Z/3$	$Z/3^2$	0	0		$Z/3$				
$Sp(3)$	$Z/3$	$Z/3 \oplus Z/3^3 \oplus Z_{(3)}$	0	0		$Z/3 \oplus Z/3^3 \oplus Z_{(3)}$				

where \oplus indicates the direct sum of the groups.

Since G is parallelizable, the Poincaré duality isomorphism

$$E_i(G) \cong E^{n-i}(G)$$

holds for any spectrum E , where $n = \dim G$ (see [4, Part III]). Therefore, to compute $\widetilde{j(p)}_i(G)$ it suffices to compute $\widetilde{j(p)}^{n-i}(G)$. Theorem 1.1 is a consequence of Theorem 4.6, in which the cup-product ring structure of $\widetilde{j(p)}^*(G)$ is described for $(G, p)=(G_2, 3)$, $(Sp(3), 3)$.

The remainder of this paper is organized as follows. In §2 we collect some results for later use. In §3 we describe the action of θ_r on $\mathbf{g}(p)^*(G)$.

In §4 we compute the rings $j(\widetilde{p})^*(G)$.

2. Preliminaries

This section is devoted to describe the rings $K^*(G; Z_{(p)})$, $k^*(G; Z_{(p)})$, $g(p)^*(G)$ and the homomorphism $ch: K^*(G) \rightarrow H^*(G; Q)$.

Notice that G is assumed to be as in §1 and p is assumed to be an odd prime. According to Borel [9], G has no p -torsion and we have

Lemma 2.1. *There exist elements $x_{2m_i-1} \in H^{2m_i-1}(G; Z_{(p)})$, for $1 \leq i \leq l$ (where $l=2$ or 3), such that*

$$H^*(G; Z_{(p)}) = \Lambda(x_{2m_1-1}, x_{2m_2-1}, \dots, x_{2m_l-1})$$

where $2=m_1 \leq m_2 \leq \dots \leq m_l$ and Λ denotes an exterior algebra (over $Z_{(p)}$).

For this lemma and the values of m_i see [8].

We need the famous result of Hodgkin [11]:

Lemma 2.2. *Let $\{\rho_1, \dots, \rho_l\}$ be a system of ring generators of the complex representation ring $R(G)$. Then there exist elements $\beta(\rho_i) \in K^{-1}(G)$, for $1 \leq i \leq l$, such that*

$$K^*(G) = \Lambda(\beta(\rho_1), \dots, \beta(\rho_l)) \otimes Z[u, u^{-1}].$$

Therefore

$$K^*(G; Z_{(p)}) = \Lambda(\beta(\rho_1), \dots, \beta(\rho_l)) \otimes Z_{(p)}[u, u^{-1}].$$

The following proposition shows that

$$\kappa: k^*(G; Z_{(p)}) \rightarrow K^*(G; Z_{(p)}),$$

$$\iota: g(p)^*(G) \rightarrow k^*(G; Z_{(p)})$$

are injective.

Proposition 2.3. *One can choose elements*

$$\xi_{2m_i-1} \in g(p)^{2m_i-1}(G), \quad \text{for } 1 \leq i \leq l,$$

such that

(i) $g(p)^*(G) = \Lambda(\xi_{2m_1-1}, \dots, \xi_{2m_l-1}) \otimes Z_{(p)}[v]$.

(ii) $k^*(G; Z_{(p)}) = \Lambda(\iota(\xi_{2m_1-1}), \dots, \iota(\xi_{2m_l-1})) \otimes Z_{(p)}[u]$.

(iii) $K^*(G; Z_{(p)}) = \Lambda(\kappa\iota(\xi_{2m_1-1}), \dots, \kappa\iota(\xi_{2m_l-1})) \otimes Z_{(p)}[u, u^{-1}]$.

(iv) *The CW-filtration degree ([7, §2]) of ξ_{2m_i-1} is $2m_i-1$; or equivalently, $\kappa\iota(\xi_{2m_i-1})$ satisfies*

$$ch(u^m; \kappa\iota(\xi_{2m_i-1})) = cu_{2m_i-1} + \text{higher terms}$$

where c is a unit of $Z_{(p)}$.

Proof. By [7, §2.4] the Atiyah-Hirzebruch spectral sequence for $K^*(G; Z_{(p)})$ collapses. Therefore it follows from the naturality with respect to κ (resp. ι) that the Atiyah-Hirzebruch spectral sequence for $k^*(G; Z_{(p)})$ (resp. $g(p)^*(G)$) collapses. Thus Lemma 2.1 yields the result; in particular, for (iv) see [7, §2.5].

We quote from [14] the following

Lemma 2.4. *For our groups G , the Chern character*

$$ch: K^{-1}(G) = \tilde{K}(\Sigma G) \rightarrow \tilde{H}^*(\Sigma G; \mathcal{Q}) \cong \tilde{H}^{*-1}(G; \mathcal{Q})$$

is given by:

(1) *If $G = SU(3)$, we have*

$$ch\beta(\lambda_1) = -x_3 + \frac{1}{2}x_5,$$

$$ch\beta(\lambda_2) = -x_3 - \frac{1}{2}x_5$$

(where $\{\lambda_1, \lambda_2\}$ generates $R(SU(3))$).

(2) *If $G = Sp(2)$, we have*

$$ch\beta(\lambda_1) = x_3 - \frac{1}{6}x_7,$$

$$ch\beta(\lambda_2) = 2x_3 + \frac{2}{3}x_7.$$

(3) *If $G = G_2$, we have*

$$ch\beta(\rho_1) = 2x_3 + \frac{1}{60}x_{11},$$

$$ch\beta(\Lambda^2\rho_1) = 10x_3 - \frac{5}{12}x_{11}.$$

(4) *If $G = SU(4)$, we have*

$$ch\beta(\lambda_1) = -x_3 + \frac{1}{2}x_5 - \frac{1}{6}x_7,$$

$$ch\beta(\lambda_2) = -2x_3 + \frac{2}{3}x_7,$$

$$ch\beta(\lambda_3) = -x_3 - \frac{1}{2}x_5 - \frac{1}{6}x_7.$$

(5) *If $G = Spin(7)$, we have*

$$ch\beta(\lambda'_1) = 2x_3 - \frac{2}{3}x_7 + \frac{1}{60}x_{11},$$

$$ch\beta(\lambda'_2) = 10x_3 + \frac{2}{3}x_7 - \frac{5}{12}x_{11},$$

$$ch\beta(\Delta_7) = 2x_3 + \frac{1}{3}x_7 + \frac{1}{60}x_{11}.$$

(6) If $G=Sp(3)$, we have

$$\begin{aligned} ch\beta(\lambda_1) &= x_3 - \frac{1}{6}x_7 + \frac{1}{120}x_{11}, \\ ch\beta(\lambda_2) &= 4x_3 + \frac{1}{3}x_7 - \frac{13}{60}x_{11}, \\ ch\beta(\lambda_3) &= 6x_3 + x_7 + \frac{11}{20}x_{11}. \end{aligned}$$

An application of this result is a quick calculation of the operation ψ^r on $K^*(G)$. For example, in $K^{-1}(SU(3))$ we have

$$\begin{aligned} (2.5) \quad \psi^r(\beta(\lambda_1)) &= \frac{r^2(r+1)}{2}\beta(\lambda_1) + \frac{r^2(-r+1)}{2}\beta(\lambda_2), \\ \psi^r(\beta(\lambda_2)) &= \frac{r^2(-r+1)}{2}\beta(\lambda_1) + \frac{r^2(r+1)}{2}\beta(\lambda_2) \end{aligned}$$

(cf. the proof of Proposition 3.3).

3. The operation θ_r on $g(p)^*(G)$

In this section we first recall the facts we need about the p -localization of G . With this as a background, we shall describe the action of θ_r on $g(p)^*(G)$.

Let $B_n(p)$, for $n \geq 1$, be the S^{2n+1} -bundle over $S^{2n+2p-1}$ such that

$$H^*(B_n(p); Z/p) = \Lambda(x_{2n+1}, \mathcal{P}^1 x_{2n+1}),$$

It has a cell structure:

$$(3.1) \quad B_n(p) \simeq S^{2n+1} \cup e^{2n+1+2(p-1)} \cup e^{4n+2+2(p-1)}.$$

Then G is called p -regular if and only if it is homotopy equivalent to a product of spheres when localized at p , and G is called quasi p -regular if and only if it is homotopy equivalent to a product of spaces $B_n(p)$ and spheres when localized at p .

The following result is due to Mimura and Toda [12].

Lemma 3.2. *We have*

- (1) $SU(3) \underset{p}{\simeq} S^3 \times S^5$ for $p \geq 3$.
- (2) $Sp(2) \underset{p}{\simeq} S^3 \times S^7$ for $p \geq 5$;
 $Sp(2) \underset{3}{\simeq} B_1(3)$.
- (3) $G_2 \underset{p}{\simeq} S^3 \times S^{11}$ for $p \geq 7$;
 $G_2 \underset{5}{\simeq} B_1(5)$.

$$(4) \quad SU(4) \underset{p}{\cong} S^3 \times S^5 \times S^7 \quad \text{for } p \geq 5;$$

$$SU(4) \underset{3}{\cong} B_1(3) \times S^5.$$

$$(5) \quad Spin(7) \underset{p}{\cong} S^3 \times S^7 \times S^{11} \quad \text{for } p \geq 7;$$

$$Spin(7) \underset{5}{\cong} B_1(5) \times S^7.$$

$$(6) \quad Sp(3) \underset{p}{\cong} S^3 \times S^7 \times S^{11} \quad \text{for } p \geq 7;$$

$$Sp(3) \underset{5}{\cong} B_1(5) \times S^7.$$

We first consider the cases in which G is p -regular.

Proposition 3.3. *In the following cases there are elements $\xi_{2m_i-1} \in g(p)^{2m_i-1}(G)$, for $1 \leq i \leq l$, as in Proposition 2.3, which satisfy:*

$$(1) \quad G = SU(3), \quad p \geq 3.$$

$$(a) \quad \begin{aligned} u^2 \kappa \iota(\xi_3) &= -\frac{1}{2} \beta(\lambda_1) - \frac{1}{2} \beta(\lambda_2) && \xrightarrow{ch_1} && x_3 \\ u^3 \kappa \iota(\xi_5) &= \beta(\lambda_1) - \beta(\lambda_2) && && x_5. \end{aligned}$$

$$(b) \quad \theta_r(\xi_3) = 0, \quad \theta_r(\xi_5) = 0.$$

$$(2) \quad G = Sp(2), \quad p \geq 5.$$

$$(a) \quad \begin{aligned} u^2 \kappa \iota(\xi_3) &= \frac{2}{3} \beta(\lambda_1) + \frac{1}{6} \beta(\lambda_2) && \xrightarrow{ch} && x_3 \\ u^4 \kappa \iota(\xi_7) &= -2\beta(\lambda_1) + \beta(\lambda_2) && && x_7. \end{aligned}$$

$$(b) \quad \theta_r(\xi_3) = 0, \quad \theta_r(\xi_7) = 0.$$

$$(3) \quad G = G_2, \quad p \geq 7.$$

$$(a) \quad \begin{aligned} u^2 \kappa \iota(\xi_3) &= \frac{5}{6} \beta(\rho_1) + \frac{1}{30} \beta(\Lambda^2 \rho_1) && \xrightarrow{ch} && 2x_3 \\ u^6 \kappa \iota(\xi_{11}) &= 5\beta(\rho_1) - \beta(\Lambda^2 \rho_1) && && \frac{1}{2} x_{11}. \end{aligned}$$

$$(b) \quad \theta_r(\xi_3) = 0, \quad \theta_r(\xi_{11}) = 0.$$

$$(4) \quad G = SU(4), \quad p \geq 5.$$

$$(a) \quad \begin{aligned} u^2 \kappa \iota(\xi_3) &= -\frac{1}{3} \beta(\lambda_1) - \frac{1}{6} \beta(\lambda_2) - \frac{1}{3} \beta(\lambda_3) && && x_3 \\ u^3 \kappa \iota(\xi_5) &= \beta(\lambda_1) && - \beta(\lambda_3) && \xrightarrow{ch} && x_5 \\ u^4 \kappa \iota(\xi_7) &= -\beta(\lambda_1) + \beta(\lambda_2) - \beta(\lambda_3) && && && x_7. \end{aligned}$$

$$(b) \quad \theta_r(\xi_3) = 0, \quad \theta_r(\xi_5) = 0, \quad \theta_r(\xi_7) = 0.$$

$$(5) \quad G = Spin(7), \quad p \geq 7.$$

- (a) $u^2\kappa\iota(\xi_3) = \frac{3}{10}\beta(\lambda'_1) + \frac{1}{30}\beta(\lambda'_2) + \frac{8}{15}\beta(\Delta_7) \quad 2x_3$
 $u^4\kappa\iota(\xi_7) = -\beta(\lambda'_1) + \beta(\Delta_7) \xrightarrow{ch} x_7$
 $u^6\kappa\iota(\xi_{11}) = \beta(\lambda'_1) - \beta(\lambda'_2) + 4\beta(\Delta_7) \quad \frac{1}{2}x_{11}.$
- (b) $\theta_r(\xi_3) = 0, \theta_r(\xi_7) = 0, \theta_r(\xi_{11}) = 0.$
- (6) $G = Sp(3), p \geq 7.$
- (a) $u^2\kappa\iota(\xi_3) = \frac{2}{5}\beta(\lambda_1) + \frac{1}{10}\beta(\lambda_2) + \frac{1}{30}\beta(\lambda_3) \quad x_3$
 $u^4\kappa\iota(\xi_7) = -\frac{7}{2}\beta(\lambda_1) + \frac{1}{2}\beta(\lambda_2) + \frac{1}{4}\beta(\lambda_3) \xrightarrow{ch} x_7$
 $u^6\kappa\iota(\xi_{11}) = \beta(\lambda_1) - 2\beta(\lambda_2) + \beta(\lambda_3) \quad x_{11}.$
- (b) $\theta_r(\xi_3) = 0, \theta_r(\xi_7) = 0, \theta_r(\xi_{11}) = 0.$

Proof. We show (1) only, because the others can be shown quite similarly. Since $\{\beta(\lambda_1), \beta(\lambda_2)\}$ forms a Z -basis for $K^{-1}(SU(3))$ by Lemma 2.2 (and [14, §2]), it is easy to see that $\{-\frac{1}{2}\beta(\lambda_1) - \frac{1}{2}\beta(\lambda_2), \beta(\lambda_1) - \beta(\lambda_2)\}$ forms a $Z_{(p)}$ -basis for $K^{-1}(SU(3); Z_{(p)})$; their images under ch are as required by Lemma 2.4. On the other hand, by Proposition 2.3 $\{u^2\kappa\iota(\xi_3), u^3\kappa\iota(\xi_5)\}$ is a $Z_{(p)}$ -basis for $K^{-1}(SU(3); Z_{(p)})$. These (together with (b)) permit us to conclude that there exist $\xi_i \in g(p)^i(SU(3)), i=3, 5$, satisfying (a).

To prove (b) we compute $\psi^r(u^2\kappa\iota(\xi_3))$ and $\psi^r(u^3\kappa\iota(\xi_5))$ in $\tilde{K}(\Sigma SU(3))$. By use of the formula $ch^q\psi^r = r^q ch^q$ [1, Theorem 5.1 (vi)] where ch^q is the composition

$$\tilde{K}(\Sigma G) \xrightarrow{ch} \tilde{H}^*(\Sigma G; Q) \xrightarrow{\pi_{2q}} \tilde{H}^{2q}(\Sigma G; Q)$$

(where π_{2q} is the projection to the $2q$ -dimensional component), we have

$$ch\psi^r(u^2\kappa\iota(\xi_3)) = r^2x_3 = ch(r^2u^2\kappa\iota(\xi_3)).$$

Since $ch: \tilde{K}(\Sigma G) \rightarrow \tilde{H}^*(\Sigma G; Q)$ is injective, it follows that

$$\psi^r(u^2\kappa\iota(\xi_3)) = r^2u^2\kappa\iota(\xi_3).$$

Since $\psi^r(u^2) = r^2u^2$, it follows that

$$\psi^r(\kappa\iota(\xi_3)) = \kappa\iota(\xi_3).$$

Since ψ^r commutes with κ, ι and κ, ι are injective, it follows that

$$\psi^r(\xi_3) = \xi_3.$$

Similarly we have $\psi^r(\xi_5) = \xi_5$. So (b) follows by the definition of θ_r .

In view of Lemma 3.2, all statements in Proposition 3.3 except (a) are clear. But, if one wants to discuss a homomorphism $f^*:g(p)^*(G')\rightarrow g(p)^*(G)$ which is induced by a homomorphism of compact Lie groups $f:G\rightarrow G'$, it seems to us that (a) is necessary.

Before considering the cases in which G is quasi p -regular, we describe $g(p)^*(B_1(p))$ and the θ_r -action on it. Since θ_r detects \mathcal{L}^1 (see [13, Lemma 1.1]), it follows from the Atiyah-Hirzebruch spectral sequence argument using (3.1) that

(3.4) *There exist $\xi_i \in g(p)^i(B_1(p))$, for $i=3, 2p+1$, such that*

- (i) $g(p)^*(B_1(p)) = \Lambda(\xi_3, \xi_{2p+1}) \otimes Z_{(p)}[v]$.
- (ii) *The operation θ_r is given by*

$$\theta_r(\xi_3) = \xi_{2p+1}, \theta_r(\xi_{2p+1}) = 0.$$

Proposition 3.5. *In the following cases there are elements $\xi_{2m_i-1} \in g(p)^{2m_i-1}(G)$, for $1 \leq i \leq l$, as in Proposition 2.3, which satisfy:*

- (1) $G=Sp(2), p=3$.

- (a)
$$\begin{aligned} u^2\kappa u(\xi_3) &= \frac{1}{2}\beta(\lambda_2) \quad ch \quad x_3 + \frac{1}{3}x_7 \\ u^4\kappa u(\xi_7) &= -2\beta(\lambda_1) + \beta(\lambda_2) \longrightarrow x_7. \end{aligned}$$
- (b) $\theta_2(\xi_3) = \xi_7, \theta_2(\xi_7) = 0.$

- (2) $G=G_2, p=5$.

- (a)
$$\begin{aligned} u^2\kappa u(\xi_3) &= \beta(\rho_1) \quad 2x_3 + \frac{1}{60}x_{11} \\ u^6\kappa u(\xi_{11}) &= 5\beta(\rho_1) - \beta(\Lambda^2\rho_1) \xrightarrow{ch} \frac{1}{2}x_{11}. \end{aligned}$$
- (b) $\theta_2(\xi_3) = \frac{1}{2}\xi_{11}, \theta_2(\xi_{11}) = 0.$

- (3) $G=SU(4), p=3$.

- (a)
$$\begin{aligned} u^2\kappa u(\xi_3) &= -\frac{1}{2}\beta(\lambda_1) \quad -\frac{1}{2}\beta(\lambda_3) \quad x_3 + \frac{1}{6}x_7 \\ u^3\kappa u(\xi_5) &= \beta(\lambda_1) \quad -\beta(\lambda_3) \xrightarrow{ch} x_5 \\ u^4\kappa u(\xi_7) &= -\beta(\lambda_1) + \beta(\lambda_2) - \beta(\lambda_3) \quad x_7. \end{aligned}$$
- (b) $\theta_2(\xi_3) = \frac{1}{2}\xi_7, \theta_2(\xi_5) = 0, \theta_2(\xi_7) = 0.$

- (4) $G=Spin(7), p=5$.

- (a)
$$\begin{aligned} u^2\kappa u(\xi_3) &= \frac{1}{3}\beta(\lambda'_1) \quad +\frac{2}{3}\beta(\Delta_7) \quad 2x_3 + \frac{1}{60}x_{11} \\ u^4\kappa u(\xi_7) &= -\beta(\lambda'_1) \quad +\beta(\Delta_7) \xrightarrow{ch} x_7 \\ u^6\kappa u(\xi_{11}) &= \beta(\lambda'_1) - \beta(\lambda'_2) + 4\beta(\Delta_7) \quad \frac{1}{2}x_{11}. \end{aligned}$$

$$(b) \quad \theta_2(\xi_3) = \frac{1}{2} \xi_{11}, \quad \theta_2(\xi_7) = 0, \quad \theta_2(\xi_{11}) = 0.$$

$$(5) \quad G = Sp(3), \quad p = 5.$$

$$(a) \quad u^2\kappa(\xi_3) = \frac{5}{12}\beta(\lambda_1) + \frac{1}{12}\beta(\lambda_2) + \frac{1}{24}\beta(\lambda_3) \quad x_3 + \frac{1}{120}x_{11}$$

$$u^4\kappa(\xi_7) = -\frac{7}{2}\beta(\lambda_1) + \frac{1}{2}\beta(\lambda_2) + \frac{1}{4}\beta(\lambda_3) \xrightarrow{ch} x_7$$

$$u^6\kappa(\xi_{11}) = 2\beta(\lambda_1) - 2\beta(\lambda_2) + \beta(\lambda_3) \quad x_{11}.$$

$$(b) \quad \theta_2(\xi_3) = \frac{1}{8} \xi_{11}, \quad \theta_2(\xi_7) = 0, \quad \theta_2(\xi_{11}) = 0.$$

Proof. We prove (1) only; the proof for the others is similar. First, (a) follows from Proposition 2.3 and Lemma 2.4 as in the proof of Proposition 3.3. To prove (b) we compute $\psi^2(u^2\kappa(\xi_3))$. In $\tilde{K}(\Sigma Sp(2))$ we have

$$\begin{aligned} ch \psi^2(u^2\kappa(\xi_3)) &= 2^2x_3 + \frac{2^4}{3}x_7 \\ &= 2^2(x_3 + \frac{1}{3}x_7) + 2^2x_7 \\ &= 2^2chu^2\kappa(\xi_3) + 2^2chu^4\kappa(\xi_7). \end{aligned}$$

Therefore

$$\psi^2(u^2\kappa(\xi_3)) = 2^2u^2\kappa(\xi_3) + 2^2u^4\kappa(\xi_7).$$

Since $\iota(v) = u^2$ (where $p = 3$), it follows that

$$\psi^2(\xi_3) = \xi_3 + v\xi_7.$$

Similarly we have

$$\psi^2(\xi_{11}) = \xi_{11}.$$

These imply the result.

There remain the cases in which G is neither p -regular nor quasi p -regular.

Proposition 3.6. *In the following cases there are elements $\xi_{2m_i-1} \in g(p)^{2m_i-1}(G)$, for $1 \leq i \leq l$, as in Proposition 2.3, which satisfy:*

$$(1) \quad G = G_2, \quad p = 3.$$

$$(a) \quad \begin{array}{l} u^2\kappa(\xi_3) = \beta(\rho_1) \\ u^6\kappa(\xi_{11}) = 5\beta(\rho_1) - \beta(\Lambda^2\rho_1) \end{array} \xrightarrow{ch} \begin{array}{l} 2x_3 + \frac{1}{60}x_{11} \\ \frac{1}{2}x_{11}. \end{array}$$

$$(b) \quad \theta_2(\xi_3) = \frac{1}{2}v\xi_{11}, \quad \theta_2(\xi_{11}) = 0.$$

(2) $G = Spin(7)$, $p = 3$.

$$\begin{aligned}
 \text{(a)} \quad u^2\kappa(\xi_3) &= \beta(\lambda'_1) && 2x_3 - \frac{2}{3}x_7 + \frac{1}{60}x_{11} \\
 u^4\kappa(\xi_7) &= -\beta(\lambda'_1) + \beta(\Delta_7) \xrightarrow{ch} && x_7 \\
 u^6\kappa(\xi_{11}) &= \beta(\lambda'_1) - \beta(\lambda'_2) + 4\beta(\Delta_7) && \frac{1}{2}x_{11}.
 \end{aligned}$$

$$\text{(b)} \quad \theta_2(\xi_3) = -2\xi_7 + \frac{1}{2}v\xi_{11}, \quad \theta_2(\xi_7) = 0, \quad \theta_2(\xi_{11}) = 0.$$

(3) $G = Sp(3)$, $p = 3$.

$$\begin{aligned}
 \text{(a)} \quad u^2\kappa(\xi_3) &= \beta(\lambda_1) && x_3 - \frac{1}{6}x_7 + \frac{1}{120}x_{11} \\
 u^4\kappa(\xi_7) &= -4\beta(\lambda_1) + \beta(\lambda_2) \xrightarrow{ch} && x_7 - \frac{1}{4}x_{11} \\
 u^6\kappa(\xi_{11}) &= 2\beta(\lambda_1) - 2\beta(\lambda_2) + \beta(\lambda_3) && x_{11}. \\
 \text{(b)} \quad \theta_2(\xi_3) &= -\frac{1}{2}\xi_7, \quad \theta_2(\xi_7) = -\frac{3}{4}\xi_{11}, \quad \theta_2(\xi_{11}) = 0.
 \end{aligned}$$

This proposition follows from the calculation similar to that in the proof of Proposition 3.3. We omit the details of the proof.

It is known [10] that

$$Spin(7) \underset{p}{\cong} Sp(3).$$

Therefore $j(3)_*(Spin(7)) \cong j(3)_*(Sp(3))$. Henceforth we exclude to consider the former.

4. The $j(p)$ -cohomology of G

In Lemma 4.2 we present formulas on the multiplicative structure of $\widetilde{j(p)}^*(X)$ (where X satisfies a certain condition). In the rest of this section we compute $\widetilde{j(p)}^*(G)$ for all pairs (G, p) . Finally we comment on $\widetilde{j(p)}_*(G)$.

Throughout this section, the letters X and Y will stand for finite connected CW -complexes.

Consider the fibration sequence

$$\Sigma^{2p-3}g(p) \xrightarrow{\delta} j(p) \xrightarrow{\eta} g(p) \xrightarrow{\theta} \Sigma^{2p-2}g(p).$$

It leads to a short exact sequence

$$\begin{aligned}
 \text{(4.1)} \quad 0 \rightarrow \text{Coker}(\theta: \widetilde{g(p)}^{i-1}(X) \rightarrow \widetilde{g(p)}^{i+2p-3}(X)) \xrightarrow{\delta} \\
 \widetilde{j(p)}^i(X) \xrightarrow{\eta} \text{Ker}(\theta: \widetilde{g(p)}^i(X) \rightarrow \widetilde{g(p)}^{i+2p-2}(X)) \rightarrow 0
 \end{aligned}$$

for any $i \in \mathbb{Z}$. In this situation we shall use the following notation. For any $x \in \widetilde{g(p)^*(X)}$ we write \bar{x} for $\delta(x) \in \widetilde{j(p)^*-2p+3(X)}$; therefore, if $x \in \text{Im}(\theta)$, we have $\bar{x} = 0$. Suppose now that $x \in \text{Ker}(\theta)$. Then we denote by \bar{x} an element such that $\eta(\bar{x}) = x$; it is unique if $\widetilde{g(p)^*(X)}$ is (p) -torsion free. This condition is satisfied for $X = G$ by Proposition 2.3.

Lemma 4.2. *Suppose that $\widetilde{g(p)^*(X)}$ is torsion free. Then, with the above notations, for any $x, y \in \widetilde{g(p)^*(X)}$, the following formulas hold in $\widetilde{j(p)^*(X)}$:*


- (i) $\bar{x} \cup \bar{y} = \overline{x \cup y}$.
- (ii) $\bar{x} \cup \bar{y} = \overline{x \cup y}$.
- (iii) $\bar{x} \cup \bar{y} = \overline{x \cup y}$.
- (iv) $\bar{x} \cup \bar{y} = 0$.

Proof. Parts (i), (ii) and (iii) are proved by using the same technique as in [13, §4]; we refer to it for the details. In this proof we will use the facts which are shown there, without specific reference.

It remains to prove part (iv). Since η is a map of ring spectra and $\eta\delta \simeq 0$, we have

$$\eta(\bar{x} \cup \bar{y}) = \eta(\delta(x) \cup \delta(y)) = \eta\delta(x) \cup \eta\delta(y) = 0 \cup 0 = 0.$$

Hence there exists a $z \in g(p)^*(X)$ such that $\bar{x} \cup \bar{y} = \bar{z}$. This equality implies that, in the following diagram, the outer square is commutative:



where d is the diagonal map; $x \in g(p)^{m+2p-3}(X)$, $y \in g(p)^{n+2p-3}(X)$; $g = g(p)$, $j = j(p)$; μ_g and μ_j are multiplications in $g(p)$ and $j(p)$ respectively. The commutativity of square I is obvious and that of square II was shown in [13, Lemma 4.4]. Thus we have

$$\begin{aligned} \bar{z} &= \bar{x} \cup \bar{y} = \mu_j(\bar{x} \wedge \bar{y})d \\ &= \mu_j(\delta \wedge 1)(1 \wedge \delta)(x \wedge y)d \\ &= \delta \mu_g(1 \wedge \eta)(1 \wedge \delta)(x \wedge y)d \\ &= 0. \end{aligned}$$

By virtue of this lemma, if one computes $\widetilde{j(p)^*(X)}$ by using (4.1), then its ring structure is automatically known.

We now record some basic data for $\mathbf{j}(p)$. Since $\psi^r(v) = r^{p-1}v$, the coefficient ring of $\mathbf{j}(p)$ is given by

$$(4.3) \quad \pi_*(\mathbf{j}(p)) = Z_{(p)}\{\tilde{1}\} \oplus \bigoplus_{i \geq 1} Z/p^{1+\nu_p(i)}\{\overline{v^{i-1}}\}$$

where the formula

$$\nu_p(r^{i(p-1)} - 1) = 1 + \nu_p(i)$$

([2, Lemma (2.12)]) is essential. We also have the Cartan formula for θ_r : for any $x, y \in g(p)^*(X)$,

$$(4.4) \quad \theta_r(x \cup y) = \theta_r(x) \cup y + x \cup \theta_r(y) + v \cdot \theta_r(x) \cup \theta_r(y)$$

(cf. [13, Lemma 4.1]).

Let us enter into a computation of $\widetilde{j(p)^*(G)}$. As is well known, the cofibration

$$X \vee Y \rightarrow X \times Y \rightarrow X \wedge Y$$

leads to a split short exact sequence

$$0 \rightarrow \widetilde{j(p)^i(X \wedge Y)} \rightarrow \widetilde{j(p)^i(X \times Y)} \rightarrow \widetilde{j(p)^i(X)} \oplus \widetilde{j(p)^i(Y)} \rightarrow 0$$

for any $i \in \mathbb{Z}$. Therefore by Lemma 3.2, in order to compute $\widetilde{j(p)^*(G)}$ when G is p -regular or quasi p -regular, it suffices to determine $\widetilde{j(p)^*(B_1(p))}$. From (3.4) we deduce

Proposition 4.5. *The ring $\widetilde{j(p)^*(B_1(p))}$ is given by:*

$$\begin{aligned} \widetilde{j(p)^*(B_1(p))} &= \widetilde{j(p)^*(S^0)}\{\widetilde{\xi_3 \xi_{2p+1}}\} \oplus Z_{(p)}\{\widetilde{\xi_{2p+1}}\} \\ &\quad \oplus Z_{(p)}\{\overline{(r^{p-1} - 1)\xi_3 - v\xi_{2p+1}}\} \\ &\quad \oplus \bigoplus_{i \geq 1} Z/p^{2+\nu_p(i)+\nu_p(i+1)}\{\overline{v^{i-1}\xi_3}\} \end{aligned}$$

where the relations

$$\begin{aligned} \overline{\xi_{2p+1}} &= 0, \\ \overline{v^i \xi_{2p+1}} &= (r^{-i(p-1)} - 1) \overline{v^{i-1} \xi_3} \quad (\text{for } i \geq 1) \end{aligned}$$

hold.

Proof. By using (4.4), in $g(p)^*(B_1(p))$ we have

$$\begin{aligned} \theta_r(v^i \xi_3 \xi_{2p+1}) &= (r^{i(p-1)} - 1) v^{i-1} \xi_3 \xi_{2p+1}, \\ \theta_r(v^i \xi_{2p+1}) &= (r^{i(p-1)} - 1) v^{i-1} \xi_{2p+1}, \\ \theta_r(v^i \xi_3) &= (r^{i(p-1)} - 1) v^{i-1} \xi_3 + r^{i(p-1)} v^i \xi_{2p+1}. \end{aligned}$$

So the kernel and cokernel of θ , are easily calculated and the result follows.

In this way, if G is p -regular or quasi p -regular, the ring $j(p)^*(G)$ can be described. For the remaining cases, from parts (1) and (3) of Proposition 3.6 we deduce

Theorem 4.6. *With the notation as in Lemma 4.2, the ring $\widetilde{j(3)^*(G)}$ for $G=G_2, Sp(3)$ is given by:*

- (1) $G=G_2$.

i	$\widetilde{j(3)^*(G_2)}$
14	$Z_{(3)} \overline{\{\xi_3 \xi_{11}\}}$
13	0
12	0
11	$Z/3 \overline{\{\xi_3 \xi_{11}\}} \oplus Z_{(3)} \overline{\{\xi_{11}\}}$
10	0
9	0
8	$Z/3 \overline{\{\xi_{11}\}}$
7	$Z/3 \overline{\{v \xi_3 \xi_{11}\}}$
6	0
5	0
4	0
3	$Z/3^2 \overline{\{v^2 \xi_3 \xi_{11}\}} \oplus Z_{(3)} \overline{\{3\xi_3 - \frac{1}{10} v^2 \xi_{11}\}}$
2	0
1	0
0	$Z/3^3 \overline{\{\xi_3\}}$
-1	$Z/3 \overline{\{v^3 \xi_3 \xi_{11}\}}$
-2	0
-3	0
-4	$Z/3^2 \overline{\{v \xi_3\}}$
-5	$Z/3 \overline{\{v^4 \xi_3 \xi_{11}\}}$
-6	0
-7	0

(2) $G=Sp(3)$.

i	$j(3)^i(Sp(3))$
21	$Z_{(3)} \{ \widetilde{\xi_3 \xi_7 \xi_{11}} \}$
20	0
19	0
18	$Z/3 \{ \xi_3 \xi_7 \xi_{11} \} \oplus Z_{(3)} \{ \widetilde{\xi_7 \xi_{11}} \}$
17	0
16	0
15	0
14	$Z/3 \{ \overline{v \xi_3 \xi_7 \xi_{11}} \} \oplus Z_{(3)} \{ 3 \xi_3 \xi_{11} + \frac{1}{2} v \xi_7 \xi_{11} \}$
13	0
12	0
11	$Z/3 \{ \overline{\xi_3 \xi_{11}} \} \oplus Z_{(3)} \{ \widetilde{\xi_{11}} \}$
10	$Z/3^2 \{ \overline{v^2 \xi_3 \xi_7 \xi_{11}} \} \oplus Z_{(3)} \{ 3 \xi_3 \xi_7 + \frac{3}{4} v \xi_3 \xi_{11} + \frac{1}{40} v^2 \xi_7 \xi_{11} \}$
9	0
8	$Z/3 \{ \overline{\xi_{11}} \}$
7	$Z/3 \{ \xi_3 \xi_7 - \frac{1}{16} v \xi_3 \xi_{11} \} \oplus Z/3^3 \{ \overline{v \xi_3 \xi_{11}} \} \oplus Z_{(3)} \{ \xi_7 + \frac{1}{4} v \xi_{11} \}$
6	$Z/3 \{ \overline{v^3 \xi_3 \xi_7 \xi_{11}} \}$
5	0
4	$Z/3 \{ \overline{v \xi_{11}} \}$
3	$Z/3 \{ v \xi_3 \xi_7 - \frac{4}{5} v^2 \xi_3 \xi_{11} \} \oplus Z/3^3 \{ \overline{v^2 \xi_3 \xi_{11}} \} \oplus Z_{(3)} \{ 3 \xi_3 + \frac{1}{2} v \xi_7 + \frac{1}{10} v^2 \xi_{11} \}$
2	$Z/3 \{ \overline{v^4 \xi_3 \xi_7 \xi_{11}} \}$
1	0
0	$Z/3 \{ 3 \xi_3 - \frac{8}{5} v^2 \xi_{11} \} \oplus Z/3^3 \{ \overline{\xi_3} \}$

Proof. For (1) we have

$$\begin{aligned} \theta_2(v^i \xi_3 \xi_{11}) &= (2^{2i} - 1)v^{i-1} \xi_3 \xi_{11}, \\ \theta_2(v^i \xi_{11}) &= (2^{2i} - 1)v^{i-1} \xi_{11}, \\ \theta_2(v^i \xi_3) &= (2^{2i} - 1)v^{i-1} \xi_3 + 2^{2i-1} v^{i+1} \xi_{11}. \end{aligned}$$

For (2) we have

$$\begin{aligned} \theta_2(v^i \xi_3 \xi_7 \xi_{11}) &= (2^{2i} - 1)v^{i-1} \xi_3 \xi_7 \xi_{11}, \\ \theta_2(v^i \xi_7 \xi_{11}) &= (2^{2i} - 1)v^{i-1} \xi_7 \xi_{11}, \\ \theta_2(v^i \xi_3 \xi_{11}) &= (2^{2i} - 1)v^{i-1} \xi_3 \xi_{11} - 2^{2i-1} v^i \xi_7 \xi_{11}, \\ \theta_2(v^i \xi_{11}) &= (2^{2i} - 1)v^{i-1} \xi_{11}, \\ \theta_2(v^i \xi_3 \xi_7) &= (2^{2i} - 1)v^{i-1} \xi_3 \xi_7 - 2^{2i-2} v^i \xi_3 \xi_{11} + 2^{2i-3} v^{i+1} \xi_7 \xi_{11}, \\ \theta_2(v^i \xi_7) &= (2^{2i} - 1)v^{i-1} \xi_7 - 2^{2i-2} v^i \xi_{11}, \\ \theta_2(v^i \xi_3) &= (2^{2i} - 1)v^{i-1} \xi_3 - 2^{2i-1} v^i \xi_7. \end{aligned}$$

So the result follows from elementary calculations of the kernel and cokernel of θ_2 .

Proof of Theorem 1.1.

By using the Poincaré duality isomorphism

$$\begin{aligned} j(p)_i(G) &= \widetilde{j(p)}_i(G) \oplus \widetilde{j(p)}_i(S^0) \\ &\cong \widetilde{j(p)}^{n-i}(G) \oplus \widetilde{j(p)}^{n-i}(S^0) = j(p)^{n-i}(G) \end{aligned}$$

where $n = \dim G$, Theorem 1.1 follows from Theorem 4.6 and (4.3).

Finally we talk about the Pontrjagin ring structure of $\widetilde{j(p)}_*(G)$. Since in Lemma 2.2 each $\beta(\rho_i)$ is primitive (see [11]), the ring structure of $K_*(G)$ can be determined. Furthermore, the ψ^r -action on $K_*(G)$ can be determined by using the formula

$$\psi^r(a \cap \alpha) = \psi^r(a) \cap \psi^r(\alpha)$$

where $a \in K^*(G)$, $\alpha \in K_*(G)$ and \cap denotes the cap product. Therefore the ring structure of $\widetilde{j(p)}_*(G)$ will be obtained by using the homology instead of the cohomology and taking the same course as in this paper.

References

- [1] J.F. Adams: *Vector fields on spheres*, Ann. of Math. **75** (1962), 603–632.
- [2] J.F. Adams: *On the groups $J(X)$ -II*, Topology **3** (1965), 137–171.
- [3] J.F. Adams: *Lectures on generalised cohomology*, Lecture Notes in Math. Vol. 99, Springer, 1969.
- [4] J.F. Adams: *Stable homotopy and generalised homology*, Chicago Lectures in Math., Univ. of Chicago Press, 1974.
- [5] J.F. Adams: *Infinite loop spaces*, Ann. of Math. Studies No. 90, Princeton Univ. Press, 1978.
- [6] J.F. Adams, A.S. Harris and R.M. Switzer: *Hopf algebras of cooperations for real and complex K -theory*, Proc. London Math. Soc. (3) **23** (1971), 385–408.

- [7] M.F. Atiyah and F. Hirzebruch: Vector bundles and homogeneous spaces, Proc. Sympos. Pure Math. Vol. 3, Amer. Math. Soc., 1961.
- [8] A. Borel: *Topology of Lie groups and characteristic classes*, Bull. Amer. Math. Soc. **61** (1955), 397–432.
- [9] A. Borel: *Sous-groupes commutatifs et torsion des groupes de Lie compacts connexes*, Tôhoku Math. J. (2) **13** (1961), 216–240.
- [10] B. Harris: *On the homotopy groups of the classical groups*, Ann. of Math. **74** (1961), 407–413.
- [11] L. Hodgkin: *On the K-theory of Lie groups*, Topology **6** (1967), 1–36.
- [12] M. Mimura and H. Toda: *Cohomology operations and the homotopy of compact Lie groups-I*, Topology **9** (1970), 317–336.
- [13] T. Watanabe: *On the spectrum representing algebraic K-theory for a finite field*, Osaka J. Math. **22** (1985), 447–462.
- [14] T. Watanabe: *Chern characters on compact Lie groups of low rank*, Osaka J. Math. **22** (1985), 463–488.

Department of Mathematics
Osaka Women's University
Daisen-cho, Sakai
Osaka 590, Japan