

Title	T-cadherin Is Involved in Neuronal Circuit Formation in the Mammalian Neocortex
Author(s)	早野, 祐紀
Citation	大阪大学, 2015, 博士論文
Version Type	VoR
URL	<a href="https://doi.org/10.18910/52238">https://doi.org/10.18910/52238</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

---

# T-Cadherin Is Involved in Neuronal Circuit Formation in the Mammalian Neocortex

---

(大脳皮質神経回路形成における T-カドヘリンの役割)

Yuki Hayano

早野 祐紀

大阪大学大学院・生命機能研究科・細胞分子神経生物学研究室

2015 年 3 月

## Summary

Cortical efferent and afferent fibers are arranged in a stereotyped pattern in the intermediate zone (IZ). Here, we studied the mechanism of axonal pathway formation by identifying a molecule that is expressed in a subset of cortical axons in the rat. I found that T-cadherin (T-cad), a member of the cadherin family, is expressed in deep-layer cell axons projecting to subcortical structures, but not in upper layer callosal axons projecting to the contralateral cortex. Ectopic expression of T-cad in upper layer cells induced axons to project toward subcortical structures via the upper part of the IZ. Moreover, the axons of deep-layer cells in which T-cad expression was suppressed by RNAi projected towards the contralateral cortex via an aberrant route. These results suggest that T-cad is involved in axonal pathway formation in the developing cortex.

## Table of Contents

Summary .....	1
General introduction.....	3
References .....	9
Introduction.....	13
Materials and Methods.....	16
Generation of monoclonal antibodies .....	16
Immunohistochemistry .....	17
Purification and identification of antigen recognized by monoclonal antibody .....	18
In situ hybridization.....	20
Western blotting analysis.....	21
In utero electroporation.....	21
Coculture of cortical neurons with L-cells .....	23
Quantitative analysis of axon growth .....	24
Results .....	25
mAb 6C9 recognizes a subset of axons in the developing brain .....	25
Expression of T-cad in specific axonal pathways .....	28
Distribution of T-cad-expressing axons in the IZ.....	33
T-cad expression during development.....	35
Ectopic expression of T-cad in upper layer cells alters the projection pattern .....	37
Knockdown of T-cad in deep-layer cells causes an aberrant projection .....	46
Discussion.....	50
T-cad expression pattern in the developing cortex .....	50
Axonal adhesion by T-cad .....	52
Role of T-cad in axonal sorting during development .....	54
References.....	57
Acknowledgment.....	66
Publications .....	68
Presentations.....	69

## General introduction

The mammalian neocortex is composed of numerous types of neurons, which construct complex and elaborate neuronal networks. It is not too much to say that our lives are based on the brain architecture. Importance of the cortical architecture has been implicated from an evolutionary point of view. The cerebral cortex is the most diverse structure of the brain across species. Concomitantly with number of cortical neurons, size and folding of the cerebral cortex are increased in higher mammals (Nonaka-Kinoshita *et al.* 2013). This supports the view that the expansion of cortical circuitry is the structural basis to the higher brain functions. Thereby, when cortical network was disrupted by some reason (e.g., brain injury, drugs and congenital anomaly), the brain could not work normally, resulting in various aberrant symptoms of behavior or cognition. Thus, it is important to understand how these complex and sophisticated circuitries are established during development.

The developmental process of the cortical circuitry has been studied in detail. Generation and differentiation of multiple types of neurons are primary process of the cortical development. Subsequently, axonal elongation and synapse formation occurred to establish neuronal connections. To be more precise, excitatory neurons are born in the germinal zone of the cerebral cortex and migrate into specific layers of the cortical plate (CP; gray matter) in order of birth so that six-layered structure, each of which

contains distinct types of neurons, is constructed in the CP (McConnell and Kaznowski 1991; Frotscher 1997). During migration, each type of the projection neurons start to grow axonal process into the intermediate zone (IZ; white matter) that lies beneath the CP and the subplate (SP). Corticofugal axons of deep layer neurons is known to project to the subcortical structures, such as the brainstem and the spinal cord (Gilbert and Kelly 1975; Wise and Jones 1977; Sefton *et al.* 1981). These axons orient ventrolaterally in the IZ, then entered the brainstem through the internal capsule (IC). Layer VI cortico-thalamic axons (CTAs) terminates in the thalamic nuclei and cortico-subcortical projections from layer V orient to the hindbrain and the spinal cord through the caudate putamen (Cp). In contrast, callosal axons that originate from upper layer neurons and another population of the deep layer neurons extend medially in the IZ and cross the corpus callosum (CC) to project the contralateral hemisphere (Jacobson and Trojanowski 1974; Wise and Jones 1976; Ivy and Killackey 1981; O'Leary *et al.* 1981). In the same time of corticofugal projections, cortical afferent fibers that originate from the thalamic nuclei grow along the IC and the IZ, then entered the CP to connect cortical layer IV (Kageyama and Robertson 1993).

As seen above, the cortical circuits are consisted by numerous types of efferent and afferent fibers which connect the cortical layers and other brain regions. Although

each type of neurons projects to the different targets, there are common rules in the processes for making each pathway. Newly born neurons first migrate and grow axonal processes on preexisting scaffold provided by surrounding cells. Axons of these neurons select its direction sequentially at multiple choice points (in other words ‘intermediate targets’) on the ways to the final targets. During axonal elongation, neurons usually form axonal fasciculation with same type of neurons, and each fascicle gather and form thick bundle of the white matter such as the IZ and the CC.

Several molecular mechanisms have been reported to be involved in such processes. A well-known mechanism is controlling axonal orientation by guidance molecules, which are called attractants and repellents. Given that an attractive molecule is presented by target cells, axons expressing its receptor could be attracted to the target. On the contrary, axons with the receptor for a repulsive molecule could avoid the cells expressing the repellent. In both cases, diffusible molecules are able to produce long-range effects making molecular gradient originate from secreting cells, and membrane bound molecules provide short-range effects through contact-mediated fashions. Indeed, during cortical development, the corticofugal axons have been shown to be forced into the IZ by members of the semaphorin (Sema) superfamily (Bagnard *et al.* 1998), then orient ventrolaterally by an attractant cue netrin-1 (Metin *et al.* 1997;

Richards *et al.* 1997). Callosal axons also guided into the IZ by repulsive cue Sema3A expressed in the most superficial layer of the CP (Polleux *et al.* 1998) and forced toward midline by the Sema3A gradient (Zhao *et al.* 2011), and attracted by netrin-1 gradient from midline cells. Midline crossing in the CC is also controlled by repulsive cues Slit2 and Draxin that repel axons before the crossing (Islam *et al.* 2009) and by Wnt5a that repel axons after the crossing (Keeble *et al.* 2006). Slit proteins has also a role in preventing midline crossing of thalamocortical axons (TCAs) (Bagri *et al.* 2002). TCAs have been shown to be attracted by Neuregulin1 expressed in corridor cells, which are settled in the IC (Lopez-Bendito *et al.* 2006). Thus, combinations of multiple guidance molecules in the circumstances and receptors on the growth cones make it possible to select axonal pathway at the intermediate targets.

Other types of guidance events are axonal fasciculation. Some mechanisms have been suggested in previous studies. Most simple fashion is forming axon-axon interaction through adhesive molecules localized on the surface of axons. In the invertebrate nervous system, one of the immunoglobulin family cell adhesion molecules (IgCAMs), fasciclin II has demonstrated to be expressed on a subset of fasciculating longitudinal axons in insect embryos (Harrelson and Goodman 1988). In the absence of fasciclin II, the axons fail to form axonal bundle (Grenningloh *et al.* 1991), suggesting



that fasciclin II can function in axon-axon recognition (selective fasciculation). In the developing chick spinal cord, antibodies against neuron-glia cell adhesion molecule (NgCAM) caused defasciculation of commissural neurites (Stoeckli and Landmesser 1995). In the mammalian brains, NgCAM homologue L1 has suggested to contribute axonal fasciculation by being expressed in a specific pathways such as olfactory nerve (Gong and Shipley 1996) and TCAs. In addition to the homotypic axonal fasciculation mechanism, different types of neurons could be associated with each other. In the cerebral cortex, when callosal axons cross the midline in the CC, pioneer axons, which originate from cingulate cortex are suggested to navigate later crossing axons that have different response to a guidance cue (Piper *et al.* 2009; Fothergill *et al.* 2014). Furthermore, it has been hypothesized that TCAs and CTAs meet in the IC during development and associate with each other (Molnar and Blakemore 1995; Molnar *et al.* 1998). In both cases, homophilic and/or heterophilic adhesion might be involved in selective fasciculation.

In addition to adhesion molecules on axons, extracellular matrix molecules (ECMs) have been shown to provide favorable substrate for axonal elongation. For examples, neurocan, one of the chondroitin sulfate proteoglycans, expressed in the SP cells is suggested to be involved in the distribution of TCAs in the IZ and the SP

(Fukuda *et al.* 1997). Furthermore, surrounding repulsive cues have also been implicated to drive fasciculation by pushing axons together into narrow corridor. Examples are provided by semaphorins, Slit-1 and AL-1, a glycosylphosphatidylinositol (GPI)-linked ligand for the Eph-related tyrosine kinase receptor (Meima *et al.* 1997; Taniguchi *et al.* 1997; Raper 2000; Cloutier *et al.* 2004).

Interestingly, cortical efferents and afferent fibers once closely gather in the same path on the way to the different targets; they extend long distance forming axonal fasciculation with same types of neurons and forming thick bundle of the white matter together with different types of fascicles. They switch from one fascicles to another at the specific intermediate target points toward the final target. This ‘selective fasciculation’ strategy is considered to be beneficial for extending astonishingly long distance to the final targets. However, molecular mechanisms and functional significance of the strategy is not yet fully understood.

In this thesis, I studied the molecular mechanism for the cortical circuit formation especially focusing on selective fasciculation in the white matter. Cell surface proteins that are expressed in specific pathways are plausible candidates for affecting axonal sorting through contact-mediated mechanisms because these molecules are capable of making more confined and strict expression patterns in the white matter than

diffusible molecules. As well as specificity of the expression of such molecules, timing and/or cellular localization could be crucial for creating the accurate wiring. Since such selective axonal assembly is also observed in other nervous systems, this study will provide a general rules for neural circuit formation.

## References

Bagnard, D., M. Lohrum, D. Uzeil, A. W. Puschel and J. Bolz (1998). "Semaphorins act as attractive and repulsive guidance signals during the development of cortical projections." Development **125**(24): 5043-5053.

Bagri, A., O. Marin, A. S. Plump, J. Mak, S. J. Pleasure, J. L. R. Rubenstein and M. Tessier-Lavigne (2002). "Slit proteins prevent midline crossing and determine the dorsoventral position of major axonal pathways in the mammalian forebrain." Neuron **33**(2): 233-248.

Cloutier, J. F., A. Sahay, E. C. Chang, M. Tessier-Lavigne, C. Dulac, A. L. Kolodkin and D. D. Ginty (2004). "Differential requirements for semaphorin 3F and Slit-1 in axonal targeting, fasciculation, and segregation of olfactory sensory neuron projections." J Neurosci **24**(41): 9087-96.

Fothergill, T., A. L. Donahoo, A. Douglass, O. Zalucki, J. Yuan, T. Shu, G. J. Goodhill and L. J. Richards (2014). "Netrin-DCC signaling regulates corpus callosum formation through attraction of pioneering axons and by modulating Slit2-mediated repulsion." Cereb Cortex **24**(5): 1138-51.

Frotscher, M. (1997). "Dual role of Cajal-Retzius cells and reelin in cortical development." Cell Tissue Res **290**(2): 315-22.

Fukuda, T., H. Kawano, K. Ohyama, H. P. Li, Y. Takeda, A. Oohira and K. Kawamura (1997). "Immunohistochemical localization of neurocan and L1 in the formation of thalamocortical pathway of developing rats." J Comp Neurol **382**(2): 141-52.

Gilbert, C. D. and J. P. Kelly (1975). "The projections of cells in different layers of the cat's visual cortex." J Comp Neurol **163**(1): 81-105.

Gong, Q. and M. T. Shipley (1996). "Expression of extracellular matrix molecules and cell surface molecules in the olfactory nerve pathway during early development." J Comp Neurol **366**(1): 1-14.

Grenningloh, G., E. J. Rehm and C. S. Goodman (1991). "Genetic analysis of growth cone guidance in *Drosophila*: fasciclin II functions as a neuronal recognition molecule." Cell **67**(1): 45-57.

Harrelson, A. L. and C. S. Goodman (1988). "Growth cone guidance in insects: fasciclin II is a member of the immunoglobulin superfamily." Science **242**(4879): 700-8.

Islam, S. M., Y. Shinmyo, T. Okafuji, Y. Su, I. B. Naser, G. Ahmed, S. Zhang, S. Chen, K. Ohta, H. Kiyonari, T. Abe, S. Tanaka, R. Nishinakamura, T. Terashima, T. Kitamura and H. Tanaka (2009). "Draxin, a repulsive guidance protein for spinal cord and forebrain commissures." Science **323**(5912): 388-93.

Ivy, G. O. and H. P. Killackey (1981). "The ontogeny of the distribution of callosal projection neurons in the rat parietal cortex." J Comp Neurol **195**(3): 367-89.

Jacobson, S. and J. Q. Trojanowski (1974). "The cells of origin of the corpus callosum in rat, cat and rhesus monkey." Brain Res **74**(1): 149-55.

Kageyama, G. H. and R. T. Robertson (1993). "Development of geniculocortical projections to visual cortex in rat: evidence early ingrowth and synaptogenesis." J Comp Neurol **335**(1): 123-48.

Keeble, T. R., M. M. Halford, C. Seaman, N. Kee, M. Macheda, R. B. Anderson, S. A. Stacker and H. M. Cooper (2006). "The Wnt receptor Ryk is required for Wnt5a-mediated axon guidance on the contralateral side of the corpus callosum." J Neurosci **26**(21): 5840-8.

Lopez-Bendito, G., A. Cautinat, J. A. Sanchez, F. Bielle, N. Flames, A. N. Garratt, D. A. Talmage, L. W. Role, P. Charnay, O. Marin and S. Garel (2006). "Tangential neuronal migration controls axon guidance: A role for neuregulin-1 in thalamocortical axon navigation." Cell **125**(1): 127-142.

McConnell, S. K. and C. E. Kaznowski (1991). "Cell cycle dependence of laminar determination in developing neocortex." Science **254**(5029): 282-5.

Meima, L., I. J. Kljavin, P. Moran, A. Shih, J. W. Winslow and I. W. Caras (1997). "AL-1-induced growth cone collapse of rat cortical neurons is correlated with REK7 expression and rearrangement of the actin cytoskeleton." Eur J Neurosci **9**(1): 177-88.

Metin, C., D. Deleglise, T. Serafini and T. E. Kennedy (1997). "A role for netrin-1 in the guidance of cortical efferents." Development **124**(24): 5063-5074.

Molnar, Z., R. Adams and C. Blakemore (1998). "Mechanisms underlying the early establishment of thalamocortical connections in the rat." J Neurosci **18**(15): 5723-45.

Molnar, Z. and C. Blakemore (1995). "Guidance of thalamocortical innervation." Development of the Cerebral Cortex **193**: 127-140.

Nonaka-Kinoshita, M., I. Reillo, B. Artegiani, M. A. Martinez-Martinez, M. Nelson, V. Borrell and F. Calegari (2013). "Regulation of cerebral cortex size and folding by expansion of basal progenitors." EMBO J **32**(13): 1817-28.

O'Leary, D. D., B. B. Stanfield and W. M. Cowan (1981). "Evidence that the early postnatal restriction of the cells of origin of the callosal projection is due to the elimination of axonal collaterals rather than to the death of neurons." Brain Res **227**(4): 607-17.

Piper, M., C. Plachez, O. Zalucki, T. Fothergill, G. Goudreau, R. Erzurumlu, C. Gu and L. J. Richards (2009). "Neuropilin 1-Sema signaling regulates crossing of cingulate pioneering axons during development of the corpus callosum." Cereb Cor **19 Suppl 1**: i11-21.

Polleux, F., R. J. Giger, D. D. Ginty, A. L. Kolodkin and A. Ghosh (1998). "Patterning of cortical efferent projections by semaphorin-neuropilin interactions." Science **282**(5395): 1904-6.

Raper, J. A. (2000). "Semaphorins and their receptors in vertebrates and invertebrates." Current opinion in neurobiology **10**(1): 88-94.

Richards, L. J., S. E. Koester, R. Tuttle and D. D. O'Leary (1997). "Directed growth of early cortical axons is influenced by a chemoattractant released from an intermediate target." J Neurosci **17**(7): 2445-58.

Sefton, A. J., A. Mackay-Sim, L. A. Baur and L. J. Cottee (1981). "Cortical projections to visual centres in the rat: an HRP study." Brain Res **215**(1-2): 1-13.

Stoeckli, E. T. and L. T. Landmesser (1995). "Axonin-1, Nr-CAM, and Ng-CAM play different roles in the in vivo guidance of chick commissural neurons." Neuron **14**(6): 1165-79.

Taniguchi, M., S. Yuasa, H. Fujisawa, I. Naruse, S. Saga, M. Mishina and T. Yagi (1997). "Disruption of Semaphorin III/D Gene Causes Severe Abnormality in Peripheral Nerve Projection." Neuron **19**(3): 519-530.

Wise, S. P. and E. G. Jones (1976). "The organization and postnatal development of the commissural projection of the rat somatic sensory cortex." J Comp Neurol **168**(3): 313-43.

Wise, S. P. and E. G. Jones (1977). "Cells of origin and terminal distribution of descending projections of the rat somatic sensory cortex." J Comp Neurol **175**(2): 129-57.

Zhao, H., T. Maruyama, Y. Hattori, N. Sugo, H. Takamatsu, A. Kumanogoh, R. Shirasaki and N. Yamamoto (2011). "A molecular mechanism that regulates medially oriented axonal growth of upper layer neurons in the developing neocortex." J Comp Neurol **519**(5): 834-48.

## Introduction

During development, axons with similar destinations form bundles and pathways in the CNS. In the mammalian neocortex, afferent and efferent fibers, which connect the subcortical and cortical regions, form layers in the IZ. Thalamic afferents run in the most superficial layer of the IZ, whereas efferent fibers occupy the deeper layers (Woodward *et al.* 1990; Catalano *et al.* 1991; Henke-Fahle *et al.* 1996; Molnar *et al.* 1998). Efferent fibers are further segregated within the IZ according to their destinations (Woodward *et al.* 1990; Bicknese *et al.* 1994; Niquille *et al.* 2009); callosal axons travel in the lower layers toward the contralateral cortex, while corticofugal axons oriented to the brainstem and spinal cord occupy the upper layers (Bicknese *et al.* 1994). This pathway segregation in the IZ originates during the onset of cortical development and is thought to be a fundamental mechanism for neural circuit formation.

In general, pioneer axons first navigate toward their targets during cell migration (Schwartz *et al.* 1991; Noctor *et al.* 2004; Lickiss *et al.* 2012), and subsequent axons then grow along preexisting axons to form fiber bundles (Bentley and Keshishian 1982; Taghert *et al.* 1982; McConnell *et al.* 1989; Pike *et al.* 1992). Preexisting non-neuronal cells are also involved in guiding a particular population of axons (Bentley and Caudy 1983; Sato *et al.* 1998). In both cases, cell-cell adhesion is thought to play an important

role in the formation of selective axon pathways (Dodd and Jessell 1988; Van Vactor 1998; Wang and Marquardt 2013). Indeed, several adhesion molecules are known to contribute to axonal fasciculation in the CNS by being expressed in a subset of axons (Bastiani *et al.* 1987; Patel *et al.* 1987; Chedotal *et al.* 1995; Cohen *et al.* 1998; Weiner *et al.* 2004; Maness and Schachner 2007). In cortical circuits, cell surface and extracellular matrix molecules, which may be involved in axonal fasciculation, are expressed in distinct axonal populations in the IZ. For instance, GPI-anchored netrin (netrin-G1) is specifically expressed in thalamocortical axons (Nakashiba *et al.* 2002). Neurocan, a chondroitin sulfate proteoglycan family member, and L1 are also strongly expressed in thalamocortical axons (Oohira *et al.* 1994; Fukuda *et al.* 1997; Wiencken-Barger *et al.* 2004). On the other hand, corticocortical and corticofugal axons express Tag-1 (Cntn2), a member of the immunoglobulin superfamily (Fukuda *et al.* 1997). More recently, neuropilin-1 and nectin have also been shown to be expressed in callosal axons (Hatanaka *et al.* 2009; Niquille *et al.* 2009; Piper *et al.* 2009). However, the molecules that are expressed in a subset of afferent or efferent fibers are not fully characterized. Moreover, the molecular mechanisms underlying pathway guidance in the IZ are largely unknown.



To address this issue, I searched for molecules that are involved in cortical circuit formation using a monoclonal antibody (mAb) technique. I found that a cell adhesion molecule, T-cadherin (T-cad), one of the cadherin superfamily members is expressed in a specific pathway during developmental stages.

Cadherins are membrane proteins expressed in various tissues, contain highly conserved cadherin motifs in the extracellular regions (EC domain), and constitute the superfamily. It has been known that most of the family members promote cell adhesion through the EC domains in homophilic and/or heterophilic binding manner. Several reports have implicate that cadherin molecules are involved in the development of neuronal circuitry. In the most cases, the cell adhesions by cadherins are correlated with association of its intracellular domain and cytoskeletal components. Among the family, however, T-cad (the meaning of “truncate”-type of cadherin) has exceptional structure that lacks transmembrane domain and intracellular domain, and binds to plasma membrane by GPI-anchor. Regardless of its unique structure, T-cad has been shown to have a cell adhesion property (Vestal and Ranscht 1992; Sacristan *et al.* 1993; Ciatto *et al.* 2010), although it also act as a receptor component in vasculature tissues (Hug *et al.* 2004; Denzel *et al.* 2010).

Recently it has been reported that T-cad is expressed in cortical layers and hippocampus after developmental stages. However, little is known about precise expression and function of T-cad during cortical development. In this study, I found that T-cad is expressed in axons of corticorecipient and corticofugal neurons during specific developmental stages. *In vivo* functional analyses including ectopic expression and knockdown further revealed that T-cad contributes to axonal pathfinding by cortical projection neurons.

## **Materials and Methods**

### ***Generation of monoclonal antibodies***

mAbs were produced from rat brain homogenate according to a standard procedure (Fujita *et al.* 1982). Brain homogenate prepared from P0 Sprague-Dawley (SD) rats (Nippon Animal Company, Japan) was administered to an adult mouse (BALB/c, Nippon Animal Company). Two months later, the spleen was dissociated and fused with myeloma cells. The hybridomas were plated in 96-well dishes and selected on RPMI medium (Invitrogen) containing HAT (hypoxanthine, aminopterin, and thymidine) and 10% fetal bovine serum (FBS). Each supernatant was subjected to immunohistochemical screening (see below).

### ***Immunohistochemistry***

Newborn rat brains were used for immunohistochemical screening. To study developmental changes in staining patterns for the selected mAb (6C9), brain sections were prepared from E15 to P4 rats. Brain sections of shRNA-subjected rats and T-cad-overexpressing mice were prepared at the indicated stages. All animals were anesthetized and sacrificed in accordance with the guidelines laid down by animal welfare committees of Osaka University and Japan Neuroscience Society.

Whole brains were fixed in 4% paraformaldehyde/0.1 M phosphate buffer (PB). The tissue was further immersed sequentially in 10, 20 and 30% sucrose solution, and 10 or 20  $\mu$ m-thick sections were cut by cryostat. The sections were incubated with primary antibodies at 4°C overnight, then incubated with secondary antibodies for 2 h at room temperature, and finally mounted in an aqueous buffer (50% glycerol, 0.05 M PBS) containing diamidino-2-phenylindole (DAPI). The sections were observed with a microscope (Axiophot, Zeiss, Germany), and the images were captured with a CCD camera (DP70, Olympus, Tokyo, Japan).

The following antibodies were used against the indicated proteins: T-cadherin (mouse, culture supernatant of hybridoma 6C9); Netrin-G1 (rabbit, 1:200; a generous gift from Dr. Nakashiba, RIKEN-MIT Center for Neural Circuit Genetics); Tag-1 (rabbit,

1:2000; generated against Ig-fusion protein); Ct1p2 (Abcam, ab18465; 1:100); Tbr1 (Abcam, ab31940; 1:100); EYFP [for rat sections (Invitrogen, A6455; 1:500); for mouse sections (Nacalai tesque, 04404-84; 1:1000)]. The secondary antibodies used were Alexa Fluor 488-conjugated goat anti-mouse IgG (Invitrogen, A11029), Cy3-conjugated donkey anti-rabbit IgG (Millipore, AP182C), and Cy3-conjugated goat anti-rat IgG (Millipore, AP183C).

#### ***Purification and identification of antigen recognized by monoclonal antibody***

To identify the antigen recognized by mAb 6C9, an immunoaffinity column was generated. First, the supernatant of cultured hybridoma was collected, and the antibody was purified using a protein A column (Affi-Gel Protein A MAPS II kit, Affi-Gel protein A agarose, BioRad). The purified antibody (2 mg) was then immobilized on 1 ml of Affi-Gel Hz hydrazide gel (Affi-Gel Hz immunoaffinity kit, Bio-Rad) according to the manufacturer's instructions.

To obtain the antigen, brain homogenate was applied to the affinity column. P0 rat brains were homogenized in phosphate buffer (0.05 M PB, 0.5 M NaCl, pH 7.4) and centrifuged at 20,000 g, 4°C for 20 min. One gram of the pellet was ground in chilled acetone and then dried into powder. Two milliliters of an extraction buffer (0.05 M PB, 0.5 M NaCl, 1 mM EDTA, 10 mM CHAPS, protease inhibitor mix, pH 7.4) was added

to the acetone powder, and it was then incubated at 4°C for 1.5 h. The extract was centrifuged at 100,000 g, 4°C for 30 min, and the collected supernatant was then applied to the affinity column. The column was washed 5 times with 1 ml of loading buffer (0.05 M PB, 0.5 M NaCl, 1 mM EDTA, 5 mM CHAPS, pH 7.4). The antigen was eluted with 0.15 M citrate buffer (pH 3.0) containing 5 mM CHAPS. Fractions containing the antigen were determined by Coomassie Brilliant Blue (CBB) staining of an SDS-polyacrylamide gel.

After SDS-PAGE, the antigen was blotted onto a nitrocellulose membrane with semi-dry blotting equipment. After blocking in 5% skim milk/TBS (20 mM Tris-HCl, 0.5 M NaCl, pH 7.5), the membrane was incubated with the primary antibody at 4°C overnight. Alkaline phosphatase-conjugated anti-mouse IgG secondary antibody (Nacalai Tesque, 01800-61) was used at 1:4000 dilution in 1% skim milk/TBST. The color reaction was developed in BCIP-NBT solution (Nacalai Tesque).

For amino acid sequence analysis, antigen eluted from the affinity column was concentrated 10-fold with an Ultrafree-MC filter (5,000 NMWL, UFC3LCC00, Millipore) and applied to SDS-PAGE. The protein was transferred onto a PVDF membrane (Immobilon-PSQ, Millipore) immersed in blotting buffer (25 mM Tris, 20% methanol, 0.05% SDS, 40 mM gamma-amino-n-caproic acid) using semi-dry blotting

equipment. The antigen on the membrane was detected by CBB staining, and the visualized band was excised and subjected to N-terminal Edman sequencing.

### ***In situ hybridization***

A fragment containing the full-length rat T-cad open reading frame (ORF) was obtained by RT-PCR from P0 rat brain RNA, and ligated into pGEM-T Easy vector (Promega) (pGEM/T-cad). A fragment (nucleotides 543-1611 of rat T-cad, GenBank accession number: NM\_138889) was subcloned to produce a template for *in vitro* transcription. The template was amplified by PCR using SP6 and T7 primers. A digoxigenin (DIG)-labeled RNA probe was synthesized using SP6 polymerase to produce the antisense probe (DIG-RNA Synthesis kit, Roche). The probe was purified with gel filtration columns (Quick Spin, Roche) and stored at  $-80^{\circ}\text{C}$ .

*In situ* hybridization was performed according to the previous work (Zhong *et al.* 2004). In brief, 10- $\mu\text{m}$ -thick cryosections were prepared from newborn rat brains. After refixation and acetylation, the sections were hybridized in a buffer containing 1  $\mu\text{g}/\text{ml}$  DIG-labeled RNA probe, at  $60^{\circ}\text{C}$  overnight. After high-stringency washes and blocking, the sections were incubated with alkaline phosphatase-conjugated anti-DIG antibody. Signals were visualized with BM Purple (Roche). Nissl staining was performed on adjacent sections to determine neocortical layer structure.

### ***Western blotting analysis***

Cortical tissue was collected from E15, E18, P0, P2, P4, P7, P14 and P35 rat brains and homogenized with ice-cold PBS, which contained protease inhibitors (cOmplete protease inhibitor cocktail, Roche). After centrifugation (~20,000 g) for 30 min at 4°C, the supernatant was discarded. The pellet was resuspended with PBS that contained the protein inhibitor. Protein concentrations were measured in a part of each sample (BCA protein Assay kit, Pierce), then each sample was added to SDS-PAGE after 2× sample buffer was added. The electrophoresis proteins were transferred to a PVDF membrane. The membrane was incubated with the monoclonal antibody 6C9 followed by incubation with HRP-conjugated secondary antibody (Nacalai tesque). Finally, T-cad bands were detected by a chemiluminescent method (ECL Prime Western Blotting Detection System, GE Healthcare).

### ***In utero electroporation***

To overexpress T-cad in cortical cells *in vivo*, a mammalian expression vector encoding rat T-cad was constructed. The full-length T-cad ORF was excised from pGEM/T-cad and ligated into the *Eco*RI site of pCAGGS vector to yield pCAGGS/T-cad. For generating T-cad-overexpressing animals, pCAGGS/T-cad was mixed with an expression vector encoding the fluorescent reporter EYFP (pCAGGS/EYFP) in order to

trace transfected cells. The reporter construct alone was electroporated into control animals.

*In utero* electroporation was performed on E14.5 mouse embryos, according to previous studies (Fukuchi-Shimogori and Grove 2001; Saito and Nakatsuji 2001; Tabata and Nakajima 2001). A pregnant ICR mouse (CLEA Japan, Inc.) was deeply anesthetized, and plasmid solutions were injected into the ventricular zone. Electroporation was performed at 30 V with five 50 ms pulses at intervals of 95 ms, using an electroporator (BEX, CUY-21).

To knockdown endogenous T-cad in rats, RNAi expression plasmids were constructed. Four target sequences were designed for generating short hairpin-type RNAs (shRNAs) and were then inserted into piGEMEmU6 vector (Clontech) in order to be transcribed under the mouse U6 promoter. Three of the plasmids (piGEMEmU6/Tcad1631, piGEMEmU6/Tcad689 and pCAG-EYFP/mU6-Tcad689), which contained the shRNA sequence corresponding to nucleotides 1631-1651 (shRNA1631) and 689-707 (shRNA689) of rat T-cad mRNA (GenBank accession number NM\_138889), were found by immunocytochemistry (Fig. 13A) and western blotting in order to suppress T-cad expression in cultured cortical cells. The target sequence for shRNA689 is exactly the same as the shRNA, which was found to be free



of an off-target effect in a previous study (Paradis *et al.* 2007). Scrambled shRNA sequence was designed by randomly shuffling the nucleotides of T-cad 689-707.

The shRNA constructs were introduced into cells destined to become deep-layer neurons (E15.0 SD rat embryos) by *in utero* electroporation, as described above. The expression of fluorescent reporter was achieved by co-electroporation with pCAGGS/EYFP or by using dual promoter vector (pCAG-EYFP/mU6-Tcad689) to trace shRNA-transfected cells. Another fluorescent reporter construct (pCAGGS/DsRedII) was used for a control.

#### ***Coculture of cortical neurons with L-cells***

To examine the role of T-cad in cortical axon growth, dissociated cortical cells expressing T-cad were cocultured with T-cad-expressing L-cells. L-cells were plated on poly-ornithine-coated wells (~10,000 cells per 10 mm-diameter well). Twelve to twenty four hours after plating, L-cells were transfected with the pCAGGS/T-cad and pCAGGS/DsRedII expression plasmids by lipofection (Lipofectamine 2000, Invitrogen). For dissociated cortical cells, E15 rat cortex was dissected and dissociated with trypsinization. These cortical cells were incubated for an hour in a solution containing pCAGGS/T-cad and pCAGGS/EYFP expression plasmids with lipofectamine. Then, these transfected cortical cells were added to the transfected L-cell cultures (~50,000

cells per well). After two days the cultures were fixed and subjected to immunohistochemistry with anti-GFP and anti-DsRed (rabbit anti-RFP, 1:500; PM005, Medical & Biological Laboratories). For the control experiment, only pCAGGS/EYFP and pCAGGS/DsRedII were used to transfect cortical cells and L cells, respectively.

### ***Quantitative analysis of axon growth***

To analyze cortical cells that overexpress T-cad, 20  $\mu\text{m}$ -thick cryosections were prepared from animals that had been subjected to *in vivo* electroporation. Ectopic T-cad protein was detected by immunohistochemistry using the anti-T-cad antibody (6C9 mAb), which does not react with endogenous mouse T-cad. In the knockdown study, 6C9 mAb was used to detect endogenous rat T-cad.

To quantify axonal projection of T-cad-overexpressing cells or shRNA-transfected cells, the axonal projection in a given brain region was categorized into five groups for P0-P6 brain sections, according to the following criteria. Mark 0: virtually no axonal segments, Mark 1: up to 10 segments; Mark 2: up to 20 segments; Mark 3: up to 30 segments; Mark 4: fasciculation of numerous axons. The mark value given for each animal was determined from 1-4 sections.


## Results

### **mAb 6C9 recognizes a subset of axons in the developing brain**

To search for molecules that are expressed in subsets of axons in cortical circuits, mAbs were generated by immunizing a mouse with homogenates of newborn rat brains. After immunohistochemical screening of a thousand clones, one of the antibodies, mAb 6C9, showed a unique staining pattern in the developing brain. As shown in Fig. 1A, strong immunoreactivity was found in the IZ of the developing rat cortex. In particular, fiber-like structures were strongly stained in the IZ, but not in the ventricular zone (VZ) (Fig. 1B). This staining pattern was found primarily in embryonic and early postnatal stages (see below). Thus, the antigen for mAb 6C9 was identified as being expressed in afferent and/or efferent fibers in the developing rat cortex, although no fiber was recognized by mAb 6C9 in the mouse brain.

To identify the 6C9 antigen, I purified it by immunoaffinity chromatography (see Materials and Methods). The purified antigen was detected as a single ~90-kDa band by SDS-PAGE (Fig. 1C, left). Western blots confirmed that this band corresponded to the 6C9 antigen (Fig. 1C, right), and an N-terminal amino acid sequencing of the purified polypeptide yielded thirteen residues, SIVVSPILIPENQ. This sequence matches a portion of T-cad (Cadherin-13 or H-cadherin), which possesses five cadherin EC domains and is bound to the plasma membrane via a C-terminal GPI anchor (Fig. 1D).

Moreover, since the 13-residue amino acid sequence was detected in an affinity-purified protein from a brain homogenate, I conclude that a full-length T-cad precursor is cleaved on the N-terminal side of the first cadherin domain (Fig. 1D, arrow) to become the mature form *in vivo*.



**Fig. 1. mAb 6C9 antigen distribution in the developing brain.**

One of the monoclonal antibodies (mAb 6C9) against developing rat brain showed a unique distribution of the antigen, identified as T-cad. Immunostaining of sagittal (A) and coronal sections (B) of E18 rat brain revealed T-cad expression in the IZ throughout the neocortex (NCX). Owing to the size of the sample, the image in A is a composite of several fields of view. (C) A protein purified by affinity chromatography with mAb 6C9 was visualized as a single band by CBB staining after SDS-PAGE and western blotting. T-cad structure is illustrated diagrammatically in D. The mAb recognizes the mature form of the protein, which is generated by cleavage of the precursor form on the N-terminal side of the first cadherin repeat. Scale bars: 100  $\mu\text{m}$  (A), 250  $\mu\text{m}$  (B). A, anterior; P, posterior; D, dorsal; V, ventral; M, medial; L, lateral. OB, olfactory bulb; ac, anterior commissure; Th, thalamus; SC, superior colliculus; CP, cortical plate; IZ, intermediate zone; VZ, ventricular zone; LV, lateral ventricle; LGE, lateral ganglionic eminence; CPu, caudate putamen.


### **Expression of T-cad in specific axonal pathways**

Since T-cad expression had only been previously described in the spinal cord (Fredette and Ranscht 1994; Takeuchi *et al.* 2000) and in adult mRNA, (Matsunaga *et al.* 2013; Rivero *et al.* 2013), I needed to thoroughly examine the protein expression pattern in the newborn rat cortex . Immunohistochemistry with mAb 6C9 demonstrated that T-cad was strongly expressed in the IZ (Figs 1B, 2A). Strong immunoreactivity was also found in the internal capsule (IC) (Figs 2C), which contains thalamocortical and corticothalamic fibers, and in the cerebral peduncle (Cp), which is composed of corticospinal axons (Fig. 2D). By contrast, there was no immunoreactivity in the corpus callosum (CC), which contains cortical fibers connecting both hemispheres (Fig. 2B). In addition, T-cad expression was also detected in the anterior commissure (Fig. 1A) and the habenular commissure (data not shown). These observations indicate that T-cad is expressed in a subpopulation of corticorecipient and corticofugal axons but is not expressed in callosal axons.

To examine which cortical neuron types express *T-cad*, *in situ* hybridization was performed in the developing cortex. Strong expression of *T-cad* was found in cortical plate (CP) cells of embryonic day (E) 15.5 rat brain, but not in the IZ or VZ (Fig. 3A,D). Robust expression of T-cad was maintained in the CP, and dispersed signals were

observed in the subplate (SP) at E18.5 (Fig. 3B,E). At postnatal day (P) 1, the signal was intense in a subpopulation of layer V cells and weak in layer VI (Fig. 3C,F), but undetectable in the upper layers, which consist of layer II/III and IV cells. Moreover, *T-cad* was expressed in a few thalamic nuclei, such as the lateral geniculate nucleus (DLG and VLG in Fig. 3C) and the reticular thalamic nucleus (Rt in Fig. 3C). Thus, *T-cad* is expressed in a subpopulation of deep-layer neurons and in a small fraction of thalamic neurons.


The laminar property of T-cad-positive neurons was further studied by comparing it with expression of layer specific markers. Since endogenous T-cad protein is present on the surface of the cell bodies only at earlier stages (Fig. 4), immunohistochemistry for COUP-TF interacting protein 2 (*Ctip2*) (Arlotta *et al.* 2005; Chen *et al.* 2005; Molyneaux *et al.* 2005), a marker protein for subcerebral projection neurons in layer V, was performed on E15.5 rat neocortex. Roughly half of T-cad-positive cells in the CP expressed *Ctip2* (Fig. 4A-D). Expression of T-brain-1 (*Tbr1*) (Hevner *et al.* 2001; Han *et al.* 2011; McKenna *et al.* 2011), a marker protein for layer VI corticothalamic projection neurons, was also detected in T-cad-positive cells in the CP and the SP of E18.5 rat neocortex (Fig. 4E-H). Thus, T-cad-expressing cells mostly consist of *Ctip2*-positive (layer V) and *Tbr1*-positive (layer VI) subcerebral projection neurons.



**Fig. 2. Distribution of T-cad-expressing axons in various brain regions**


(A-D) Immunostaining of T-cad with mAb 6C9 (green) and counter staining with DAPI (blue) were performed on coronal sections of P0 rat brain. Fiber-like staining is shown in the IZ (A), IC (C) and Cp (D), but not in the CC (B). Boxed areas on Nissl staining sections (E,F) correspond to A-D. Scale bars: 250  $\mu$ m (A-D), 1 mm (E,F). HIP, hippocampus; CC, corpus callosum; IC internal capsule; Cp, cerebral peduncle.





**Fig. 3. T-cad expression in a subset of deep-layer cells.**

(A-C) *In situ* hybridization revealed strong expression of T-cad mRNA in the cortex and thalamic nuclei at late embryonic stages (E15.5 and E18.5) and a neonatal stage (P1). Dashed line in C indicates the pial surface of the neocortex. (D) At E15.5, *T-cad* expression is found in the CP and SP, but not in the IZ, SVZ and VZ. (E) Robust expression of *T-cad* was maintained in the deep layers, and dispersed signals were observed in the SP at E18.5. (F) *T-cad* was expressed exclusively in a subpopulation of layer V and VI neurons at P1. The adjacent Nissl-stained section shows the laminar structure (right). Scale bars: 500  $\mu\text{m}$  (A-C), 100  $\mu\text{m}$  (D-F). DLG, dorsal nucleus of lateral geniculate body; VLG, ventral nucleus of lateral geniculate body; Rt, Reticular thalamus; MHb, medial habenular nucleus; LHb, lateral habenular nucleus; PV, paraventricular thalamic nucleus. MZ, marginal zone; SVZ, subventricular zone; V, layer V; VI, layer VI; SP, subplate.




**Fig. 4. Expression of deep layer markers in T-cad-positive cells**

Double staining of T-cad and marker proteins was performed on coronal sections (10  $\mu\text{m}$ -thick) prepared from E15.5 (A-D) and E18.5 (E-H) rat embryos. T-cad expression was found on the surface of cell bodies in the SP and the CP only in the early stages. (A-D) Ctip2, a marker for subcerebral projection neurons in layer V was expressed in the T-cad-positive cells (arrowheads). (E-F) Tbr1, a marker protein for layer VI corticothalamic projection neurons, was also detected in T-cad-positive cells (arrowheads). Scale bars: 50  $\mu\text{m}$  (A,E), 20  $\mu\text{m}$  (B,F)

### **Distribution of T-cad-expressing axons in the IZ**

The distribution of T-cad-expressing axons in the IZ was subsequently compared to Netrin-G1- and Tag-1-positive axons, which correspond to thalamocortical and corticofugal axons, respectively (Wolfer *et al.* 1994; Lebrand *et al.* 1996; Fukuda *et al.* 1997; Nakashiba *et al.* 2002). T-cad-expressing axons were found in the upper part of the IZ at E20, when corticofugal and thalamocortical neurons extend axons to the IZ, (Fig. 5B). In the same section, Tag-1 expression was localized in the lower part of the IZ, but overlapped partly with T-cad-expressing axons (Fig. 5B-D). T-cad-positive axons also overlapped with Netrin-G1-positive axons in the upper layer of the IZ (Fig. 5H). The two populations were also partly colocalized in the IC (Fig. 5I,J). Therefore, T-cad-expressing fibers, composed of subcerebral projection fibers and a subpopulation of thalamocortical axons, occupy the upper part of IZ.




**Fig. 5. Distribution of T-cad-expressing axons in the IZ.**

Immunostaining for T-cad and marker proteins were performed on serial sections of E20 rat brain. Nissl staining shows the layer structure of the cortex (A,E). (B,F) T-cad protein was localized in the axons at E20, and cell bodies were hardly immunostained with 6C9 mAb. T-cad-expressing fibers were distributed in the upper part of the IZ. (C) Cortical efferent fibers, visualized by immunostaining for Tag-1, occupy the lower part of the IZ. (D) T-cad and Tag-1 expression are merged. T-cad positive fibers overlap partly with Tag-1-positive fibers in the middle part of the IZ (white). (G) Thalamocortical fibers, visualized by immunostaining for Netrin-G1 extend the upper part of the IZ and the SP. (H) The two populations overlap partly in the uppermost layer of the IZ. (I,J) Partial overlap is also observed in the IC. Scale bars: 100  $\mu$ m (A,B,F,J), 500  $\mu$ m (E), 200  $\mu$ m (I).

### **T-cad expression during development**

The developmental time course of T-cad expression was analyzed by western blotting of E15-P34 rat neocortex (Fig. 6A). T-cad expression was first noticeable at E15, peaked at E18 and P0, sharply decreased, and then increased until P14. The first peak in T-cad expression coincided with the time of afferent and efferent growth, and the protein was mainly localized in axons (Figs 1, 2, 6). Based on these characteristics, I focused on T-cad function in cortical circuit formation. The time course of T-cad expression in rat neocortex was examined by immunohistochemistry with mAb 6C9 (Fig. 6B). T-cad expression was first observed in the CP and IZ at E15, around the time that deep-layer neurons are migrating radially. Expression in cell bodies was observed only at the early stages (E15-18). Consistent with the western blot results, T-cad expression in the IZ was prominent between E18 and P2 though the expression increased at the late developmental stages.



**Fig. 6. Developmental changes of T-cad expression in the cortex.**

(A) A developmental time course of T-cad expression was confirmed by western blotting analysis of rat brain lysate. Expression of T-cad was first observed at E15, and became prominent from E18 to neonatal stages. Although it is decreased after P2, the intense signal appeared again at P14. Each sample (20  $\mu$ g) was applied to SDS-PAGE except for lane 1 (100  $\mu$ g, E15). (B) T-cad expression in the rat neocortex was visualized with mAb 6C9 during the first peak of the expression. The protein was localized mainly in the axons during late embryonic stages and postnatal stages. T-cad signals are clearly defined until P0 on the axons in the IZ. The signal becomes weaker after P2. Layer structure was defined by Nissl staining of the adjacent sections. Scale bar: 100  $\mu$ m. III, layer III; IV, layer IV.

### **Ectopic expression of T-cad in upper layer cells alters the projection pattern**


In order to investigate the function of T-cad in axonal projections, the protein was ectopically expressed in upper layer cells, which otherwise do not express T-cad. A vector encoding rat T-cad was co-transfected with an *eyfp* expression vector by *in utero* electroporation into E14.5 mouse VZ cells, which are destined to become layer II/III cells. In this experiment, the mouse was used to detect ectopically expressing T-cad using immunohistochemistry with mAb 6C9 since this antibody recognizes rat, but not mouse T-cad. As a control, an *eyfp*- or a *dsred*-encoding vector was transfected into the VZ cells of the rest of the littermates. First, I confirmed that cortical laminar organization was not influenced by ectopic expression. T-cad-overexpressing cells settled correctly in the upper layers, although migration was delayed and occasionally accumulated in the SVZ and IZ (Figs 7A, 8A,E, 9A,B). These electroporated cells expressed a marker protein for upper layer cells and did not express a deep-layer cell marker, indicating that cell fate is not changed by ectopically expressed T-cad. (Fig. 8).

Shortly after birth (P3-P6), axonal projections from T-cad-overexpressing cells were compared to those from control cells. In the controls, most axons were found to extend medially in the IZ (Fig. 10A, arrowhead), and crossed the midline along the CC. Although some labeled axons were occasionally found in the striatum (McGeorge and

Faulk 1989), such projection was negligible compared to the contralateral projection (Fig. 10D). By contrast, in the *T-cad*-transfected cortex, a substantial number of labeled fibers oriented laterally and projected into the IC (Fig. 10E). The abnormal projections originated from cells which migrated into the CP (Fig. 9C,D) and accumulated in the SVZ and IZ (Fig. 9A,B, asterisk). The axons from the CP turned abnormally to the lateral direction, extended in the upper part of the IZ, which is the pathway of T-cad-positive subcortical projection neurons (Fig. 9C,D, arrowheads), and entered into the IC. Ectopic T-cad expression in these axons was confirmed by immunohistochemistry with mAb 6C9 (Fig. 10C). These abnormally projecting fibers seem to gather tightly in the IC (Fig. 10C,E). Aberrant projections were further observed in the Cp of the *T-cad*-transfected brain (Fig. 10H,I), whereas there were almost none in the Cp of the control brain (Fig. 10G). The frequencies of these abnormal projections are quantified in Fig. 11. Medially projecting axons in the CC and IZ were observed in both T-cad-overexpressing and control brains (Fig. 11A,B). By contrast, there were more axons in the IC of the T-cad-overexpressing brain than in the control (Fig. 11C). Furthermore, a moderate number of axons were found in the Cp while there were almost none in the controls (Fig. 11D).




The possibility that the T-cad-overexpression phenotype is due to its homophilic binding was studied using cocultures of cortical cells and L-cells (fibroblast cell line). I found that neurites from T-cad-overexpressing cortical cells tended to grow on adjacent T-cad-overexpressing L-cells (Fig. 12A,B). By contrast, when cortical neurons were simply transfected with *eyfp* expression vector, neurites did not grow preferentially on L-cells regardless of whether or not they express T-cad (Fig. 12C,D,E). Thus, it is likely that homophilic binding is responsible for preferential axonal extension.




**Fig. 7. Final laminar location is not altered by T-cad-overexpression.**

*T-cad* and *eyfp* expression vectors were coelectroporated into cells destined to become upper layer cells (E14.5 mouse brain). (A) In the T-cad-transfected brain (right), numerous cells were found to be detained in the deep layers at P1, whereas most labeled cells had migrated to the upper layers in the control (left). Flanking images show DAPI-staining. Scale bar: 100  $\mu$ m. (B) The cortical depth was divided into 10 equal segments from segment 1 (SP) to segment 10 (pial surface). Percentage of fluorescent labeled cells in each segment is shown cumulatively in Y-axes. After the birth (P0), 37% of labeled cells were located in the deep segments (segment #1-6) in T-cad-overexpressing brain, whereas only 23% of labeled cells were found in the control (T-cad, 300 cells from 3 brains vs control, 300 cells from 3 brains; Mann-Whitney *U* test,  $p < 0.0001$ ). However, the difference was much smaller at P6, indicating that final laminar positions are not affected by the overexpression (T-cad, 200 cells from 2 brains vs control, 200 cells from 2 brains;  $p > 0.1$ ). (C) T-cad shRNA construct (piGENEmU6/Tcad1631) and *eyfp* expression vector were cotransfected into deep-layer cells of the rat neocortex. The migration patterns of shRNA-transfected cells were not significantly different from those in the control at P0–1 (shRNA, 734 cells from 2 brains vs control, 616 cells from 3 brains; Mann-Whitney *U* test,  $p > 0.1$ ) and P6–7 (shRNA, 1132 cells from 3 brains vs control, 458 cells from 2 brains;  $p > 0.05$ ).



**Fig. 8. Cellular identity persists in T-cad-overexpressing upper layer cells.**


(A) *In situ* hybridization for *mef-2c*, an upper-layer-specific transcription factor, was performed on sections of normal and T-cad-overexpressing mouse brains at P1. *mef-2c* was expressed in the upper layers of the control cortex (left). *mef-2c* was also expressed in T-cad-overexpressing cells, which accumulated in the IZ (bordered by dashed lines, right), suggesting that T-cad-expressing cells display normal cellular identity. (B-E) Immunostaining for Ctip2, a marker for subcortical projection neurons in layer V, was performed on a section of T-cad-overexpressing mouse brain at P3. No Ctip2 signal (magenta) was detected in T-cad-overexpressing cells (green), which were settled down in the upper layer (C), detained in the layer V (D), and accumulated in the IZ or the SVZ (E). Scale bars: 100  $\mu$ m (A), 250  $\mu$ m (B), 50  $\mu$ m (C-E).



**Fig. 9. The origins of T-cad-overexpressing upper layer cell axons which project to the abnormal direction.**


*T-cad* and *eyfp* were coelectroporated into mouse cortex at E14.5, and the brain was sectioned at P3. (A) T-cad-overexpressing upper layer cells occasionally accumulated in the IZ or SVZ (asterisk). (B) Higher magnification view of the boxed area in (A). The abnormal projections originated from cells which accumulated in the SVZ and IZ (A,B) and migrated into the CP (C,D). From the accumulated cells, abnormally oriented axons directly entered the IC (B), although callosal projections were observed as well (A, arrowheads). (C,D) The aberrant projections also occurred from the cells which migrated into the CP (arrowheads), although the majority of labeled axons were oriented normally to the medial direction (arrows). These axons turned abnormally to the lateral direction and misrouted in the upper part of the IZ, which is the primary pathway of T-cad-positive subcortical projection neurons (arrowheads in C,D). Scale bars: 250  $\mu$ m (A), 100  $\mu$ m (B-D).






**Fig. 10. Abnormal projection from T-cad-overexpressing upper layer cell axons.**

(A) In control P3 mice, almost all EYFP-labeled upper layer cell axons projected medially toward the contralateral cortex (arrowhead), but few projected into the IC (D). (G) No labeled axons were detected in the Cp. By contrast, T-cad-overexpressing upper layer cell axons projected both medially (arrowhead in B) and laterally (E); a few were also found in the Cp (H). (C) A higher magnification of the IC in T-cad-overexpressing brain at P6 confirms that EYFP-positive fibers express T-cad. F and I show DAPI staining of E and H, respectively. "Fig (D,E) Panels are montages of several images at different focal planes to allow axonal processes to be seen in focus throughout the sample. Scale bars: 300  $\mu$ m (A), 100  $\mu$ m (C,G), 200  $\mu$ m (D).



**Fig. 11. Quantification of abnormal projections of T-cad-overexpressing axons.**

The number of labeled axonal segments in each brain region was classified into 5 categories based on the amount of axons labeled, with 0 indicating virtually no visible segments, and 4 indicating fasciculation of numerous axons (see Materials and Methods). The frequency distributions were studied in CC (A), IZ (B), IC (C) and Cp (D). White and black histograms represent the distributions in control and T-cad-overexpressing cortex, respectively. The numbers of samples (animals) used for control and T-cad-overexpressing cortex are indicated in parentheses.



**Fig. 12. Homophilic interactions by T-cad contribute to axonal extension.**

(A, B) L-cells were plated and transfected with *dsred* and *T-cad* expression plasmids. After 12-24 hours, dissociated cortical cells from E15 rat were transfected with *eyfp* and *T-cad* expression plasmids and were plated onto the transfected L-cell cultures. After two days in culture, neurites from cortical cells (green) tended to grow on adjacent L-cells (magenta). (C, D) Neurites from cortical cells transfected with *eyfp* alone (green) scarcely grew on L-cells (magenta) transfected with *dsred* alone. Scale bars: 50  $\mu$ m. (E) Preferential growth of cortical neurons on L-cells was quantified and compared between the three groups: (1) co-culture of T-cad-negative cortical neurons and T-cad-negative L-cells, (2) T-cad-negative neurons and T-cad-transfected L-cells, (3) T-cad-transfected neurons and T-cad-transfected L-cells. Axon growth of EYFP-labeled cortical cells on the adjacent L-cells was classified into three categories in Grow along: axon grows along L-cell, Pass through: axon partly grows on L-cell, and Grow away: axon hardly grows on L-cell. The distribution of neurites is significantly different between the groups (1) and (3), and between the groups (2) and (3). Pairwise distribution comparison *p* values were calculated using Pearson's chi-square test.


### **Knockdown of T-cad in deep-layer cells causes an aberrant projection**

The role of T-cad was further studied by knocking down endogenous T-cad in deep-layer cells. Rat tissue was used because mAb 6C9 efficiently shows expression of endogenous T-cad. One of the effective constructs, shRNA689 (Fig. 13A), was electroporated into the cells destined to be deep-layer cells. Scrambled shRNA or *dsred* vectors were transfected into cortical cells of the rest of the littermates. Brain sections were immunostained with 6C9 antibody in order to detect endogenous T-cad (Fig. 13C,F,I).

Deep-layer cells are fundamentally composed of callosal neurons, which project to the contralateral hemisphere, and ipsilaterally subcortical projection neurons. When the *dsred* plasmid was electroporated in E15.0 rat cortex, 63% of DsRed-positive cells expressed T-cad (Fig. 14A,C). This indicates that 63% of the transfected cells are subcortical projection neurons expressing T-cad, while the remaining 37%, originally T-cad-negative cells, are callosal neurons. On the other hand, only 20% of the shRNA-transfected cells were obviously T-cad-positive (Fig. 14C), indicating that T-cad expression is completely diminished *in vivo* in approximately 70% of T-cad-positive cells by the present knockdown method.




At P0, labeled deep-layer cell axons projected both medially and laterally in the IZ of both the control (Fig. 13B) and the shRNA-transfected brains. In the control and scrambled shRNA-transfected brains (boxed area in Fig. 13B), medially oriented fibers that project to the contralateral cortex were observed in the deep IZ (Fig. 13C-H). In the shRNA-transfected brain, surprisingly, part of labeled deep-layer cell axons traveled in the SP toward the contralateral cortex (arrows in Fig. 13J,K), although most labeled axons traveled in the deep IZ (Fig. 13I-K). This abnormal projection corresponds to the T-cad-negative thalamocortical axon pathway (Fig. 5F-H). As callosal axons never travel in this pathway, the abnormal projection must be due to T-cad-downregulated subcerebral fibers. The frequency distributions of such abnormal projections were compared to control and shRNA-subjected brains (Fig. 13L). As a result, the abnormality was found in all of the shRNA-transfected brains, although the extent was varied among the samples. In the control and the scrambled shRNA brains, abnormal projection was not observed. Thus, the knockdown of T-cad altered both the direction and the pathway of originally T-cad-positive subcerebral axons.



**Fig. 13. Axonal projections of T-cad-downregulated deep-layer neurons.**

(A) Suppression of T-cad by shRNA was confirmed in a primary cell culture of rat embryonic neocortical neurons. Cells were cotransfected with one of the shRNA vectors (pCAG-EYFP/mU6-Tcad689) and T-cad expression vector (pCAGGS/T-cad). (B) Deep-layer cell axons (magenta) projected both medially and laterally in normal brain. The section was counterstained with DAPI. (C-K) Higher magnification views of medially oriented deep-layer axons in IZ (same positions as boxed area in B). Endogenous T-cad-expressing axons, immunostained by 6C9 mAb (green in C-K), occupy the upper part of the IZ, whereas medially oriented deep-layer cell axons (magenta) run in the lower part in control brain (C-E) and scrambled shRNA brain (F-H). (J) Part of axons originating from deep-layer cells transfected with the shRNA vector (piGENEmU6/Tcad689) abnormally traveled in the SP (arrows). As callosal axons never travel in this pathway, the abnormal projection must be due to T-cad downregulated subcerebral fibers. Sections shown were prepared from P0 rat brains. (L) The amount of abnormal projections into the SP was classified into 4 categories (Mark0-3). The numbers of animals are indicated in parentheses. Scale bars: 50  $\mu$ m (A), 250  $\mu$ m (B), 100  $\mu$ m (C).



**Fig. 14. *In vivo* knockdown of T-cad by shRNA.**

One of the shRNA constructs (piGENE-mU6/T-cad689) and fluorescent reporter plasmid (*dsred*) were coelectroporated into cells which were destined to become deep-layer cells (E15.0 rat). Ten micrometer-thick cryosections were prepared at E18.5, when endogenous T-cad is still detectable on cell bodies of cortical cells. (A) In the control, T-cad expression was observed in the CP and SP. Higher magnification views show DsRed and T-cad double positive cells (arrows). (B) Endogenous T-cad was hardly detected in the shRNA-transfected cells migrating in the CP (arrows). (C) The proportion of T-cad-expressing cells to the DsRed-labeled cells was calculated in the CP. Endogenous T-cad was detected only in 20% of the shRNA-transfected cells, whereas it was found in 63% of DsRed-labeled cells in the control, indicating that T-cad expression was effectively down-regulated by the shRNA in approximately 70% of the originally T-cad-expressing population. The data were collected from 2 embryos each. (D) Immunostaining for Tbr1 (magenta), a marker for layer VI corticothalamic projection neurons, shows that shRNA-transfected cells (green) express Tbr1, suggesting that the cellular property had not been changed. Scale bars: 50  $\mu$ m (upper panel in A,B), 10  $\mu$ m (lower panel in A,B), 10  $\mu$ m,(D).

## **Discussion**

In the present study, I identified the cell-surface molecule T-cad in specific pathways of the developing neocortex. In cortical efferent projections, T-cad was expressed specifically in axons originating from subcerebral projection neurons in deep layers. Functional analysis demonstrated that ectopic expression of T-cad in upper layer cells resulted in abnormal projections toward subcortical structures. Moreover, the deep-layer cell axons in which T-cad expression was suppressed altered their trajectory. These results suggest that T-cad plays a role in pathway formation of corticofugal axons.

### **T-cad expression pattern in the developing cortex**

To date, some classical cadherins, such as cadherin-6 and cadherin-8, are known to be expressed in the CNS (Suzuki *et al.* 1997; Inoue *et al.* 1998; Nakagawa *et al.* 1999), although their functions are still vague. On the other hand, expression of T-cad in the spinal cord repels axons of motor neurons (Fredette and Ranscht 1994; Fredette *et al.* 1996). The complete expression pattern and function of T-cad in the CNS are not well characterized. In the present study, I first showed that T-cad is expressed in subsets of cortical and subcortical neurons in the developing brain. T-cad is primarily expressed in axons, but it is also expressed in cell bodies or dendrites at early developmental stages. Other adhesion molecules have also been shown to be expressed specifically in axons.

In the developing cortex, L1 and TAG-1, immunoglobulin superfamily members, are expressed strongly in thalamic and corticofugal fibers, respectively (Wolfer *et al.* 1994; Fukuda *et al.* 1997; Fujimori *et al.* 2000; Ohyama *et al.* 2004). Several studies have demonstrated more specific relationships between axon subtypes and molecular expression. Npn1, a receptor for semaphorin 3A (Sema3A), is expressed in callosal axons in the upper layers, which project to the contralateral hemisphere (Hatanaka *et al.* 2009; Niquille *et al.* 2009; Piper *et al.* 2009). Moreover, the adhesion molecule nectin is expressed exclusively in callosal axons that originate from deep-layer cells (Prandovszky *et al.* 2008; Molyneaux *et al.* 2009). Our present results reveal a new feature: T-cad, a member of the cadherin superfamily, is expressed in a subset of subcortically projecting axons, but not in callosal axons (Figs. 2, 4). Thus, multiple cell surface molecules, including T-cad, are expressed differentially in callosal and non-callosal axons.

T-cad expression is regulated not only spatially, but also temporally; it ceases at early postnatal stages in rat (Fig. 6), when major cortical circuits are established. In particular, T-cad is strongly expressed in late embryonic stages, when efferent and afferent fibers are accumulating in the IZ. Together with its spatial distribution, the temporal expression pattern implies that T-cad may be involved in pathway guidance in

the IZ (see below). I also found that T-cad expression in the cortex increases again at late developmental stages (Fig. 6A), which is consistent with previous reports (Matsunaga *et al.* 2013). Previous studies also suggest that T-cad might play a distinct role in neural circuit formation, such as synaptic rearrangement (Zhong *et al.* 2004; Rivero *et al.* 2013).

### **Axonal adhesion by T-cad**

To date, axon guidance molecules have been shown to aid in axonal fasciculation and pathway finding via attractive and repulsive properties (Satoda *et al.* 1995; Kawasaki *et al.* 2002; Cloutier *et al.* 2004; Imai *et al.* 2009; Jaworski and Tessier-Lavigne 2012). Cell adhesion molecules play a more direct role in axon bundling through axon-axon interactions (Buttiglione *et al.* 1998; Marthiens *et al.* 2005; Schmitz *et al.* 2008). In chick CNS, several cadherins have been shown to affect pathway guidance (Treubert-Zimmermann *et al.* 2002). Chick tectofugal pathways contain several axonal populations, each of which projects to a distinct target and expresses particular cadherin family members. Moreover, ectopic cadherin expression in a given fiber type can cause an abnormal projection, indicating that cadherin family members contribute to the formation of axon bundles. Together with these previous findings, the present study

strongly suggests that T-cad is involved in pathway formation of mammalian cortical circuits. This action of T-cad can be explained by its adhesion property (see below).

T-cad lacks a cytoplasmic domain that is present in other cadherin family members for connecting with the cytoskeleton; however, a calcium-dependent homophilic interaction was suggested for T-cad in previous studies (Vestal and Ranscht 1992; Sacristan *et al.* 1993; Ciatto *et al.* 2010). T-cad might, therefore, contribute to axonal fasciculation via this homophilic binding property. The result in cocultures of cortical neurons with L-cells supports this view (Fig. 12). Moreover, such a homophilic binding property may also contribute to afferent and efferent pathway formation *in vivo* (Molnar *et al.* 1998; Molnar *et al.* 2012). However, I cannot exclude the possibility that ligand-receptor interactions between T-cad-expressing axons and adjacent cells could be involved in axonal pathway formation, since T-cad is known to act as a receptor molecule in the metabolic system (Hug *et al.* 2004; Denzel *et al.* 2010),

In the overexpression study described here, ectopically expressed T-cad caused a migration delay of the transfected cells (Fig. 7). The exogenous expression was detected not only in the axons (Fig. 10C), but also in the cell bodies of delayed and aggregated cells (data not shown). The migration delay of T-cad-overexpressing cells may also be due to cell adhesion caused by ectopically expressing T-cad on cell bodies, as the

expression on the cell bodies is more intense in the delayed cells than in the normally migrating upper layer cells.

### **Role of T-cad in axonal sorting during development**

A prominent aspect of cortical axonal formation is the selection of mediolateral orientation. Callosal axons originating in the upper layers project medially in the IZ, whereas subcortically projecting axons, which express T-cad, are directed laterally towards the IC and the Cp. The present findings indicate that T-cad contributes to pathway formation of deep-layer cell axons. In addition, T-cad may also be involved in directed axon growth of these cells. Indeed, a proportion of T-cad-overexpressing upper layer cell axons turned laterally, ran in the upper part of the IZ (Fig. 9C), then traveled in the IC and Cp (Fig. 10), which is the pathway for the corticospinal tract. Furthermore, originally subcerebral cortical axons were oriented toward the contralateral cortex by the knockdown (Fig. 13). The trajectory was also distinct from the original pathway of subcerebral cortical axons (Fig. 13I-K). Thus, it is likely that T-cad plays a role in both direction and pathway of subcortically projecting axons.

However, it is unlikely that T-cad is the only determinant of this laterally oriented axon growth because most T-cad-overexpressing upper layer cell axons extended



medially in the IZ (arrowhead in Figs 9A, 10B). Taken together, our observations suggest that endogenous T-cad is involved in laterally oriented axon growth of subcerebral projection neurons. It has been shown that netrin-1 released by IC cells attracts axons of subcerebral projection neurons (Richards *et al.* 1997), and one of our previous studies has also demonstrated that graded expression of Sema3A contributes to medially directed axon growth of upper layer cells (Zhao *et al.* 2011). Such diffusible molecules may therefore cooperate with cell surface molecules such as adhesion proteins.

Does ectopic expression of T-cad alter the identity of upper layer cells? Our data indicate that this is unlikely, as upper layer cells overexpressing T-cad still exhibited expression of an upper-layer-specific gene, but not a deep-layer-specific marker (Fig. 8). Moreover, T-cad knockdown did not change the laminar property (Fig. 14D). This contrasts with the occurrence of cell fate conversion in response to forced expression of transcription factors (Arlotta *et al.* 2005; Lai *et al.* 2008). Moreover, the final laminar locations of cells that are destined for the upper layers were also unaffected by overexpression of T-cad (Fig. 7B). T-cad may therefore be more important for fasciculation of subcortically projecting axons or axonal pathway guidance, but not differentiation of neuronal subtypes.

In summary, T-cad is expressed specifically in subcerebral projection neurons and is most likely to be involved in pathway formation during development. Differential expression of cell-surface molecules such as T-cad might thus be an integral process in the construction of axonal pathways in the mammalian brain.

## References

Arlotta, P., B. J. Molyneaux, J. Chen, J. Inoue, R. Kominami and J. D. Macklis (2005). "Neuronal subtype-specific genes that control corticospinal motor neuron development in vivo." Neuron 45(2): 207-21.

Bastiani, M. J., A. L. Harrelson, P. M. Snow and C. S. Goodman (1987). "Expression of fasciclin I and II glycoproteins on subsets of axon pathways during neuronal development in the grasshopper." Cell 48(5): 745-55.

Bentley, D. and M. Caudy (1983). "Pioneer axons lose directed growth after selective killing of guidepost cells." Nature 304(5921): 62-5.

Bentley, D. and H. Keshishian (1982). "Pathfinding by Peripheral Pioneer Neurons in Grasshoppers." Science 218(4577): 1082-1088.

Bicknese, A. R., A. M. Sheppard, D. D. O'Leary and A. L. Pearlman (1994). "Thalamocortical axons extend along a chondroitin sulfate proteoglycan-enriched pathway coincident with the neocortical subplate and distinct from the efferent path." J Neurosci 14(6): 3500-10.

Buttiglione, M., J. M. Revest, O. Pavlou, D. Karagogeos, A. Furley, G. Rougon and C. Faivre-Sarrailh (1998). "A functional interaction between the neuronal adhesion molecules TAG-1 and F3 modulates neurite outgrowth and fasciculation of cerebellar granule cells." J Neurosci 18(17): 6853-70.

Catalano, S. M., R. T. Robertson and H. P. Killackey (1991). "Early ingrowth of thalamocortical afferents to the neocortex of the prenatal rat." Proc Natl Acad Sci U S A 88(8): 2999-3003.

Chedotal, A., O. Pourquie and C. Sotelo (1995). "Initial tract formation in the brain of the chick embryo: selective expression of the BEN/SC1/DM-GRASP cell adhesion molecule." Eur J Neurosci 7(2): 198-212.

Chen, B., L. R. Schaevitz and S. K. McConnell (2005). "Fezl regulates the differentiation and axon targeting of layer 5 subcortical projection neurons in cerebral cortex." Proc Natl Acad Sci U S A 102(47): 17184-9.

Ciatto, C., F. Bahna, N. Zampieri, H. C. VanSteenhouse, P. S. Katsamba, G. Ahlsen, O. J. Harrison, J. Brasch, X. Jin, S. Posy, J. Vendome, B. Ranscht, T. M. Jessell, B. Honig and L. Shapiro (2010). "T-cadherin structures reveal a novel adhesive binding mechanism." Nat Struct Mol Biol 17(3): 339-47.

Cloutier, J. F., A. Sahay, E. C. Chang, M. Tessier-Lavigne, C. Dulac, A. L. Kolodkin and D. D. Ginty (2004). "Differential requirements for semaphorin 3F and Slit-1 in axonal targeting, fasciculation, and segregation of olfactory sensory neuron projections." J Neurosci 24(41): 9087-96.

Cohen, N. R., J. S. Taylor, L. B. Scott, R. W. Guillery, P. Soriano and A. J. Furley (1998). "Errors in corticospinal axon guidance in mice lacking the neural cell adhesion molecule L1." Current biology : CB 8(1): 26-33.

Denzel, M. S., M. C. Scimia, P. M. Zumstein, K. Walsh, P. Ruiz-Lozano and B. Ranscht (2010). "T-cadherin is critical for adiponectin-mediated cardioprotection in mice." J Clin Invest 120(12): 4342-52.

Dodd, J. and T. M. Jessell (1988). "Axon guidance and the patterning of neuronal projections in vertebrates." Science 242(4879): 692-9.

Fredette, B. J., J. Miller and B. Ranscht (1996). "Inhibition of motor axon growth by T-cadherin substrata." Development 122(10): 3163-71.

Fredette, B. J. and B. Ranscht (1994). "T-cadherin expression delineates specific regions of the developing motor axon-hindlimb projection pathway." J Neurosci 14(12): 7331-46.

Frotscher, M. (1997). "Dual role of Cajal-Retzius cells and reelin in cortical development." Cell Tissue Res 290(2): 315-22.

Fujimori, K. E., K. Takeuchi, T. Yazaki, K. Uyemura, Y. Nojyo and N. Tamamki (2000). "Expression of L1 and TAG-1 in the corticospinal, callosal, and hippocampal commissural neurons in the developing rat telencephalon as revealed by retrograde and in situ hybridization double labeling." J Comp Neurol 417(3): 275-88.

Fujita, S. C., S. L. Zipursky, S. Benzer, A. Ferrus and S. L. Shotwell (1982). "Monoclonal antibodies against the Drosophila nervous system." Proc Natl Acad Sci U S A 79(24): 7929-33.

Fukuchi-Shimogori, T. and E. A. Grove (2001). "Neocortex patterning by the secreted signaling molecule FGF8." Science 294(5544): 1071-4.

Fukuda, T., H. Kawano, K. Ohya, H. P. Li, Y. Takeda, A. Oohira and K. Kawamura (1997). "Immunohistochemical localization of neurocan and L1 in the formation of thalamocortical pathway of developing rats." J Comp Neurol 382(2): 141-52.

Han, W., K. Y. Kwan, S. Shim, M. M. Lam, Y. Shin, X. Xu, Y. Zhu, M. Li and N. Sestan (2011). "TBR1 directly represses Fezf2 to control the laminar origin and development of the corticospinal tract." Proc Natl Acad Sci U S A 108(7): 3041-6.

Hatanaka, Y., T. Matsumoto, Y. Yanagawa, H. Fujisawa, F. Murakami and M. Masu (2009). "Distinct roles of neuropilin 1 signaling for radial and tangential extension of callosal axons." J Comp Neurol 514(3): 215-25.

Henke-Fahle, S., F. Mann, M. Gotz, K. Wild and J. Bolz (1996). "Dual action of a carbohydrate epitope on afferent and efferent axons in cortical development." J Neurosci 16(13): 4195-206.

Hevner, R. F., L. Shi, N. Justice, Y. Hsueh, M. Sheng, S. Smiga, A. Bulfone, A. M. Goffinet, A. T. Campagnoni and J. L. Rubenstein (2001). "Tbr1 regulates differentiation of the preplate and layer 6." Neuron 29(2): 353-66.

Hug, C., J. Wang, N. S. Ahmad, J. S. Bogan, T. S. Tsao and H. F. Lodish (2004). "T-cadherin is a receptor for hexameric and high-molecular-weight forms of Acrp30/adiponectin." Proc Natl Acad Sci U S A 101(28): 10308-13.

Imai, T., T. Yamazaki, R. Kobayakawa, K. Kobayakawa, T. Abe, M. Suzuki and H. Sakano (2009). "Pre-target axon sorting establishes the neural map topography." Science 325(5940): 585-90.

Inoue, T., T. Tanaka, S. C. Suzuki and M. Takeichi (1998). "Cadherin-6 in the developing mouse brain: expression along restricted connection systems and synaptic localization suggest a potential role in neuronal circuitry." Dev Dyn 211(4): 338-51.

Jaworski, A. and M. Tessier-Lavigne (2012). "Autocrine/juxtacrine regulation of axon fasciculation by Slit-Robo signaling." Nature neuroscience 15(3): 367-9.

Kawasaki, T., Y. Bekku, F. Suto, T. Kitsukawa, M. Taniguchi, I. Nagatsu, T. Nagatsu, K. Itoh, T. Yagi and H. Fujisawa (2002). "Requirement of neuropilin 1-mediated Sema3A signals in patterning of the sympathetic nervous system." Development 129(3): 671-80.

Lai, T., D. Jabaudon, B. J. Molyneaux, E. Azim, P. Arlotta, J. R. Menezes and J. D. Macklis (2008). "SOX5 controls the sequential generation of distinct corticofugal neuron subtypes." Neuron 57(2): 232-47.

Lebrand, C., O. Cases, C. Adelbrecht, A. Doye, C. Alvarez, S. El Mestikawy, I. Seif and P. Gaspar (1996). "Transient uptake and storage of serotonin in developing thalamic neurons." Neuron 17(5): 823-35.

Lickiss, T., A. F. Cheung, C. E. Hutchinson, J. S. Taylor and Z. Molnar (2012). "Examining the relationship between early axon growth and transcription factor expression in the developing cerebral cortex." J Anat 220(3): 201-11.

Maness, P. F. and M. Schachner (2007). "Neural recognition molecules of the immunoglobulin superfamily: signaling transducers of axon guidance and neuronal migration." Nature Neurosci 10(1): 19-26.

Marthiens, V., J. Gavard, F. Padilla, C. Monnet, V. Castellani, M. Lambert and R. M. Mege (2005). "A novel function for cadherin-11 in the regulation of motor axon elongation and fasciculation." Mol Cell Neurosci 28(4): 715-26.

Matsunaga, E., S. Nambu, M. Oka and A. Iriki (2013). "Differential cadherin expression in the developing postnatal telencephalon of a New World monkey." J Comp Neurol 521(17): 4027-60.

McConnell, S. K., A. Ghosh and C. J. Shatz (1989). "Subplate neurons pioneer the first axon pathway from the cerebral cortex." Science 245(4921): 978-82.

McGeorge, A. J. and R. L. Faulkner (1989). "The organization of the projection from the cerebral cortex to the striatum in the rat." Neuroscience 29(3): 503-37.

McKenna, W. L., J. Betancourt, K. A. Larkin, B. Abrams, C. Guo, J. L. Rubenstein and B. Chen (2011). "Tbr1 and Fezf2 regulate alternate corticofugal neuronal identities during neocortical development." The Journal of neuroscience : the official journal of the Society for Neuroscience 31(2): 549-64.

Molnar, Z., R. Adams and C. Blakemore (1998). "Mechanisms underlying the early establishment of thalamocortical connections in the rat." J Neurosci 18(15): 5723-45.

Molnar, Z., R. Adams, A. M. Goffinet and C. Blakemore (1998). "The role of the first postmitotic cortical cells in the development of thalamocortical innervation in the reeler mouse." J Neurosci 18(15): 5746-65.

Molnar, Z., S. Garel, G. Lopez-Bendito, P. Maness and D. J. Price (2012). "Mechanisms controlling the guidance of thalamocortical axons through the embryonic forebrain." Eur J Neurosci 35(10): 1573-85.

Molyneaux, B. J., P. Arlotta, R. M. Fame, J. L. MacDonald, K. L. MacQuarrie and J. D. Macklis (2009). "Novel subtype-specific genes identify distinct subpopulations of callosal projection neurons." J Neurosci 29(39): 12343-54.

Molyneaux, B. J., P. Arlotta, T. Hirata, M. Hibi and J. D. Macklis (2005). "Fezl is required for the birth and specification of corticospinal motor neurons." Neuron 47(6): 817-31.

Nakagawa, Y., J. E. Johnson and D. D. O'Leary (1999). "Graded and areal expression patterns of regulatory genes and cadherins in embryonic neocortex independent of thalamocortical input." J Neurosci 19(24): 10877-85.

Nakashiba, T., S. Nishimura, T. Ikeda and S. Itohara (2002). "Complementary expression and neurite outgrowth activity of netrin-G subfamily members." Mech Dev 111(1-2): 47-60.

Niquille, M., S. Garel, F. Mann, J. P. Hornung, B. Otsmane, S. Chevalley, C. Parras, F. Guillemot, P. Gaspar, Y. Yanagawa and C. Lebrand (2009). "Transient neuronal populations are required to guide callosal axons: a role for semaphorin 3C." PLoS Biol 7(10): e1000230.

Noctor, S. C., V. Martinez-Cerdeno, L. Ivic and A. R. Kriegstein (2004). "Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases." Nat Neurosci 7(2): 136-44.

Ohyama, K., K. Tan-Takeuchi, M. Kutsche, M. Schachner, K. Uyemura and K. Kawamura (2004). "Neural cell adhesion molecule L1 is required for fasciculation and routing of thalamocortical fibres and corticothalamic fibres." Neurosci Res 48(4): 471-5.

Oohira, A., F. Matsui, E. Watanabe, Y. Kushima and N. Maeda (1994). "Developmentally regulated expression of a brain specific species of chondroitin sulfate proteoglycan, neurocan, identified with a monoclonal antibody IG2 in the rat cerebrum." Neuroscience 60(1): 145-57.

Paradis, S., D. B. Harrar, Y. Lin, A. C. Koon, J. L. Hauser, E. C. Griffith, L. Zhu, L. F. Brass, C. Chen and M. E. Greenberg (2007). "An RNAi-based approach identifies molecules required for glutamatergic and GABAergic synapse development." Neuron 53(2): 217-32.

Patel, N. H., P. M. Snow and C. S. Goodman (1987). "Characterization and cloning of fasciclin III: a glycoprotein expressed on a subset of neurons and axon pathways in *Drosophila*." Cell 48(6): 975-88.

Pike, S. H., E. F. Melancon and J. S. Eisen (1992). "Pathfinding by zebrafish motoneurons in the absence of normal pioneer axons." Development 114(4): 825-31.

Piper, M., C. Plachez, O. Zalucki, T. Fothergill, G. Goudreau, R. Erzurumlu, C. Gu and L. J. Richards (2009). "Neuropilin 1-Sema signaling regulates crossing of cingulate pioneering axons during development of the corpus callosum." Cereb Cor 19 Suppl 1: i11-21.

Prandovszky, E., S. Horvath, L. Gellert, S. K. Kovacs, Z. Janka, J. Toldi, D. Shukla and T. Valyi-Nagy (2008). "Nectin-1 (HveC) is expressed at high levels in neural subtypes that regulate radial migration of cortical and cerebellar neurons of the developing human and murine brain." J Neurovirol 14(2): 164-72.



Richards, L. J., S. E. Koester, R. Tuttle and D. D. O'Leary (1997). "Directed growth of early cortical axons is influenced by a chemoattractant released from an intermediate target." J Neurosci 17(7): 2445-58.

Rivero, O., S. Sich, S. Popp, A. Schmitt, B. Franke and K. P. Lesch (2013). "Impact of the ADHD-susceptibility gene CDH13 on development and function of brain networks." Eur Neuropsychopharmacol 23(6): 492-507.

Sacristan, M. P., D. J. Vestal, M. T. Dours-Zimmermann and B. Ranscht (1993). "T-cadherin 2: molecular characterization, function in cell adhesion, and coexpression with T-cadherin and N-cadherin." J Neurosci Res 34(6): 664-80.

Saito, T. and N. Nakatsuji (2001). "Efficient gene transfer into the embryonic mouse brain using in vivo electroporation." Dev Biol 240(1): 237-46.

Sato, Y., T. Hirata, M. Ogawa and H. Fujisawa (1998). "Requirement for early-generated neurons recognized by monoclonal antibody lot1 in the formation of lateral olfactory tract." J Neurosci 18(19): 7800-10.

Satoda, M., S. Takagi, K. Ohta, T. Hirata and H. Fujisawa (1995). "Differential expression of two cell surface proteins, neuropilin and plexin, in *Xenopus* olfactory axon subclasses." J Neurosci 15(1 Pt 2): 942-55.

Schmitz, C., I. Wacker and H. Hutter (2008). "The Fat-like cadherin CDH-4 controls axon fasciculation, cell migration and hypodermis and pharynx development in *Caenorhabditis elegans*." Dev Biol 316(2): 249-59.

Schwartz, M. L., P. Rakic and P. S. Goldman-Rakic (1991). "Early phenotype expression of cortical neurons: evidence that a subclass of migrating neurons have callosal axons." Proceedings of the National Academy of Sciences of the United States of America 88(4): 1354-8.

Suzuki, S. C., T. Inoue, Y. Kimura, T. Tanaka and M. Takeichi (1997). "Neuronal circuits are subdivided by differential expression of type-II classic cadherins in postnatal mouse brains." Mol Cell Neurosci 9(5-6): 433-47.

Tabata, H. and K. Nakajima (2001). "Efficient in utero gene transfer system to the developing mouse brain using electroporation: visualization of neuronal migration in the developing cortex." Neuroscience 103(4): 865-72.

Taghert, P. H., M. J. Bastiani, R. K. Ho and C. S. Goodman (1982). "Guidance of pioneer growth cones: filopodial contacts and coupling revealed with an antibody to Lucifer Yellow." Dev Biol 94(2): 391-9.

Takeuchi, T., A. Misaki, S. B. Liang, A. Tachibana, N. Hayashi, H. Sonobe and Y. Ohtsuki (2000). "Expression of T-cadherin (CDH13, H-Cadherin) in human brain and its characteristics as a negative growth regulator of epidermal growth factor in neuroblastoma cells." J Neurochem 74(4): 1489-97.

Treubert-Zimmermann, U., D. Heyers and C. Redies (2002). "Targeting axons to specific fiber tracts in vivo by altering cadherin expression." J Neurosci 22(17): 7617-26.

Van Vactor, D. (1998). "Adhesion and signaling in axonal fasciculation." Curr Opin Neurobiol 8(1): 80-6.

Vestal, D. J. and B. Ranscht (1992). "Glycosyl phosphatidylinositol--anchored T-cadherin mediates calcium-dependent, homophilic cell adhesion." J Cell Biol 119(2): 451-61.

Wang, L. and T. Marquardt (2013). "What axons tell each other: axon-axon signaling in nerve and circuit assembly." Curr Opin Neurobiol 23(6): 974-82.

Weiner, J. A., S. J. Koo, S. Nicolas, S. Fraboulet, S. L. Pfaff, O. Pourquie and J. R. Sanes (2004). "Axon fasciculation defects and retinal dysplasias in mice lacking the immunoglobulin superfamily adhesion molecule BEN/ALCAM/SC1." Mol Cell Neurosci 27(1): 59-69.

Wiencken-Barger, A. E., J. Mavity-Hudson, U. Bartsch, M. Schachner and V. A. Casagrande (2004). "The role of L1 in axon pathfinding and fasciculation." Cereb Cortex 14(2): 121-31.

Wolfer, D. P., A. Henahan-Beatty, E. T. Stoekli, P. Sonderegger and H. P. Lipp (1994). "Distribution of TAG-1/axonin-1 in fibre tracts and migratory streams of the developing mouse nervous system." J Comp Neurol 345(1): 1-32.

Woodward, W. R., N. Chiaia, T. J. Teyler, L. Leong and B. M. Coull (1990). "Organization of cortical afferent and efferent pathways in the white matter of the rat visual system." Neuroscience 36(2): 393-401.

Zhao, H., T. Maruyama, Y. Hattori, N. Sugo, H. Takamatsu, A. Kumanogoh, R. Shirasaki and N. Yamamoto (2011). "A molecular mechanism that regulates medially oriented axonal growth of upper layer neurons in the developing neocortex." J Comp Neurol 519(5): 834-48.

Zhong, Y., M. Takemoto, T. Fukuda, Y. Hattori, F. Murakami, D. Nakajima, M. Nakayama and N. Yamamoto (2004). "Identification of the genes that are expressed in the upper layers of the neocortex." Cereb Cortex 14(10): 1144-52.

## **Acknowledgment**

I would like to express my deepest gratitude to my advisor, Dr. Nobuhiko Yamamoto, you have been a tremendous mentor for me. I would like to thank you for encouraging my research and for allowing me to grow as a research scientist. Your advice on both research as well as on my career have been priceless. I would also like to extend my gratitude to my committee members, Dr. Takeshi Yagi, Dr. Fujio Murakami and Dr. Sachiko Tsukita for your insightful comments and hard questions. My sincere thanks also goes to Dr. Shigemi Norioka and Dr. Kosei Takeuchi for providing a lot of helpful suggestions and technical supports.

I would especially like to thank Dr. Ryuichi Shirasaki for giving me so many sound advises and constructive remarks. I would like to thank Dr. Noriyuki Sugo and Dr. Hong Zhao for supporting me when I collected data for my thesis, and for offering encouraging words. I thank my labmates in Laboratory of Cellular and Molecular Neurobiology for working together, and for all the fun we have had in the last three years. I also thank all members of Laboratory of Immune Response Dynamics, Immunology Frontier Research Center (IFReC) for encouraging me a lot. In particular, I am grateful to Dr. Kazuhiro Suzuki for giving me a good deal of time to study in graduate school.

A special thanks to my family. Words cannot express how grateful I am to my mother-in-law, father-in-law, my father and my husband for all of the sacrifices that you've made on my behalf. At the end, I would like express special appreciation to my beloved mother supporting me spiritually throughout my life.

## Publications

1. Hayano, Y., H. Zhao, H. Kobayashi, K. Takeuchi, S. Norioka and N. Yamamoto (2014). "The role of T-cadherin in axonal pathway formation in neocortical circuits." *Development* **141**(24): 4784-93.
2. Nakai, A., Y. Hayano, F. Furuta, M. Noda and K. Suzuki (2014). "Control of lymphocyte egress from lymph nodes through beta2-adrenergic receptors." *Journal of Experimental Medicine* **211**(13): 2583-98.
3. Takemoto, M., Y. Hattori, H. Zhao, H. Sato, A. Tamada, S. Sasaki, K. Nakajima and N. Yamamoto (2011). "Laminar and areal expression of unc5d and its role in cortical cell survival." *Cerebral Cortex* **21**(8): 1925-34.
4. Zhao, H., T. Maruyama, Y. Hattori, N. Sugo, H. Takamatsu, A. Kumanogoh, R. Shirasaki and N. Yamamoto (2011). "A molecular mechanism that regulates medially oriented axonal growth of upper layer neurons in the developing neocortex." *Journal of Comparative Neurology* **519**(5): 834-48.
5. Yamamoto, N., M. Takemoto, Y. Hattori and K. Hanamura (2006). Molecular Basis for the Formation of Lamina-Specific Thalamocortical Projection. In R. Erzurumlu, W. Guido and Z. Molnár (Eds.) *Development and plasticity in sensory thalamus and cortex*. Springer US: pp.79-91.
6. Zhong, Y., M. Takemoto, T. Fukuda, Y. Hattori, F. Murakami, D. Nakajima, M. Nakayama and N. Yamamoto (2004). "Identification of the genes that are expressed in the upper layers of the neocortex." *Cerebral Cortex* **14**(10): 1144-52.

## Presentations

1. Hayano, Y. and N. Yamamoto (2013) "The Role of T-cadherin in Axonal Pathway Formation in Neocortical Circuits." The 36<sup>th</sup> Annual Meeting of the Japan Neuroscience Society.
2. Hattori, Y., S. Matsuoka and S. Suematsu (2009) "Analysis of Antigen specific T cell Function of Artificial Lymphoid Tissue." The 39<sup>th</sup> Annual Meeting of the Japanese Society for Immunology.
3. Hattori, Y., S. Matsuoka, C. Takeshita and S. Suematsu (2009) "Immune Function of the Artificial Lymphoid Tissue." The 8<sup>th</sup> Congress of the Japanese Society for Regenerative Medicine.
4. Hattori, Y. and S. Suematsu (2007) "Functional Lymphatic Vessel Formation in Tissue-Engineered Secondary Lymphoid Tissue-Like Organoids in Mice." The 37<sup>th</sup> Annual Meeting of the Japanese Society for Immunology.
5. Hattori, Y., T. Watanabe and S. Suematsu (2006) "Vascular Structure in Tissue-Engineered Secondary Lymphoid Tissue-Like Organoid in Mice." The 36<sup>th</sup> Annual Meeting of the Japanese Society for Immunology.
6. Hattori, Y. and Yamamoto N. (2005) "Role of T-cadherin in cellular migration and axonal pathfinding of neocortical neurons." The 28<sup>th</sup> Annual Meeting of the Japan Neuroscience Society.
7. Hattori, Y., K. Inui, K. Kobayashi, H. Kobayashi, K. Takeuchi, S. Norioka and N. Yamamoto (2004) "T-cadherin expression in cortical circuit formation." The 27<sup>th</sup> Annual Meeting of the Japan Neuroscience Society.
8. Hattori, Y., M. Saito, N. Matsushita and K. Kobayashi (1998) "Visualization of axonal guidance and patterning in transgenic mice expressing green fluorescent protein." The 21<sup>th</sup> Annual Meeting of the Japan Neuroscience Society.