

Title	Two-sided bounds for the complexity of cyclic branched coverings of two-bridge links
Author(s)	Petronio, Carlo; Vesnin, Andrei
Citation	Osaka Journal of Mathematics. 46(4) P.1077-P.1095
Issue Date	2009-12
Text Version	publisher
URL	https://doi.org/10.18910/5247
DOI	10.18910/5247
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

TWO-SIDED BOUNDS FOR THE COMPLEXITY OF CYCLIC BRANCHED COVERINGS OF TWO-BRIDGE LINKS

CARLO PETRONIO and ANDREI VESNIN

(Received March 5, 2007, revised August 14, 2008)

Abstract

We consider closed orientable 3-dimensional hyperbolic manifolds which are cyclic branched coverings of the 3-sphere, with branching set being a two-bridge knot (or link). We establish two-sided linear bounds depending on the order of the covering for the Matveev complexity of the covering manifold. The lower estimate uses the hyperbolic volume and results of Cao-Meyerhoff, Guéritaud-Futer (who recently improved previous work of Lackenby), and Futer-Kalfagianni-Purcell, and it comes in two versions: a weaker general form and a sharper form. The upper estimate is based on an explicit triangulation, which also allows us to give a bound on the Delzant T-invariant of the fundamental group of the manifold.

1. Definitions, motivations and statements

Complexity. Using simple spines (a technical notion from piecewise linear topology that we will not need to recall in this paper), Matveev [23] introduced a notion of *complexity* for compact 3-dimensional manifolds. If M is such an object, its complexity $c(M) \in \mathbb{N}$ is a very efficient measure of “how complicated” M is, because:

- every 3-manifold can be uniquely expressed as a connected sum of prime ones (this is an old and well-known fact, see [15]);
- c is additive under connected sum;
- if M is closed and prime, $c(M)$ is precisely the minimal number of tetrahedra needed to triangulate M .

In the last item the notion of triangulation is only meant in a loose sense, namely just as a gluing of tetrahedra along faces, and an exception has to be made for the four prime M 's for which $c(M) = 0$, that is S^3 , $\mathbb{R}P^3$, $S^2 \times S^1$, and $L(3, 1)$.

Computing exactly the complexity $c(M)$ of any given 3-manifold M is theoretically very difficult, even if quite easy experimentally, using computers [25]. In the closed prime case the state of the art is as follows:

- A computer-aided tabulation of the closed M 's with $c(M) \leq 12$ has been completed in various steps [21, 25, 26] (see also [24]);

2000 Mathematics Subject Classification. Primary 57M27; Secondary 57M50.

This work is the result of a collaboration carried out in the framework of the INTAS project “CalcoMet-GT” 03-51-3663. The second author was also partially supported by the grant NSH-8526.2006.1, by a grant of the RFBR, and by a grant of the Siberian Branch of RAN.

- A general lower bound for $c(M)$ in terms of the homology of M was established in [27];
- Asymptotic two-sided bounds for the complexity of some specific infinite series of manifolds were obtained in [28, 29, 32];
- A conjectural formula for the complexity of any Seifert fibred space and torus bundle over the circle was proposed (and proved to be an upper bound) in [22].

Several other results, including exact computations for infinite series, have been obtained in the case of manifolds with non-empty boundary, see [4, 9, 10, 11]. Since we will stick in this paper to the closed case, we do not review them here.

Using the hyperbolic volume and deep results of Lackenby [20] improved recently for the case of hyperbolic two-bridge links in [14], and of Cao-Meyerhoff [7], together with explicit triangulation methods to be found [30, 31], we will analyze in this paper the complexity of cyclic coverings of the 3-sphere branched along two-bridge knots and links. More specifically, we will prove asymptotic two-sided linear estimates for the complexity in terms of the order of the covering. Exploiting some results of [12] we will also provide a sharper lower estimate in a restricted context. Before giving our statements we need to recall some terminology.

Two-bridge knots and links. If p, q are coprime integers with $p \geq 2$ we denote by $K(p, q)$ the two-bridge link in the 3-sphere S^3 determined by p and q , see [6, 17, 31]. It is well-known that $K(p, q)$ does not change if a multiple of $2p$ is added to q , so one can assume that $|q| < p$. In addition $K(p, -q)$ is the mirror image of $K(p, q)$. Therefore, since we will not care in the sequel about orientation, we can assume $q > 0$. Summing up, from now on our assumption will always be that the following happens:

$$(1) \quad p, q \in \mathbb{Z}, \quad p \geq 2, \quad 0 < q < p, \quad (p, q) = 1.$$

We recall that if p is odd then $K(p, q)$ is a knot, otherwise it is a 2-component link; moreover, two-bridge knots and links are alternating [6, p. 189]. Planar alternating diagrams of $K(p, q)$ will be shown below. Since we are only interested in the topology of the branched coverings of $K(p, q)$, we regard it as an unoriented knot (or link), and we define it to be equivalent to some other $K(p', q')$ if there is an automorphism of S^3 , possibly an orientation-reversing one, that maps $K(p, q)$ to $K(p', q')$. It is well-known (see [6, p. 185]) that $K(p', q')$ and $K(p, q)$ are equivalent if and only if $p' = p$ and $q' \equiv \pm q^{\pm 1} \pmod{p}$.

Under the current assumption (1), the two-bridge knot (or link) $K(p, q)$ is a torus knot (or link) precisely when q is 1 or $p - 1$, and it is hyperbolic otherwise. The simplest non-hyperbolic examples are the Hopf link $K(2, 1)$, the left-handed trefoil knot $K(3, 1)$ and its mirror image $K(3, 2)$, the right-handed trefoil (but we are considering a knot to be equivalent to its mirror image, as just explained). The easiest hyperbolic $K(p, q)$ is the figure-eight knot $K(5, 2)$.

Branched coverings. If $K(p, q)$ is a knot (i.e. p is odd) and $n \geq 2$ is an integer, the n -fold cyclic covering of S^3 branched along $K(p, q)$ is a well-defined closed

orientable 3-manifold that we will denote by $M_n(p, q)$. One way of defining it is as the metric completion of the quotient of the universal covering of $S^3 \setminus K(p, q)$ under the action of the kernel of the homomorphism $\pi_1(S^3 \setminus K(p, q)) \rightarrow \mathbb{Z}/n\mathbb{Z}$ which factors through the Abelianization $\pi_1(S^3 \setminus K(p, q)) \rightarrow H_1(S^3 \setminus K(p, q))$ and sends a meridian of $K(p, q)$, which generates $H_1(S^3 \setminus K(p, q))$, to $[1] \in \mathbb{Z}/n\mathbb{Z}$.

If $K(p, q)$ is a link and a generator $[m]$ of $\mathbb{Z}/n\mathbb{Z}$ is given, a similar construction defines the *meridian-cyclic* branched covering $M_{n,m}(p, q)$ of S^3 along $K(p, q)$, by requiring the meridians of the two components of $K(p, q)$ to be sent to $[1]$ and $[m] \in \mathbb{Z}/n\mathbb{Z}$ respectively. Note that meridian-cyclic coverings are also called *strongly cyclic* in [36], and that the two components of $K(p, q)$ can be switched, therefore we do not need to specify which meridian is mapped to $[1]$ and which to $[m]$. Since in the sequel we will prove estimates on the complexity of $M_{n,m}(p, q)$ which depend on n only and apply to every $M_{n,m}(p, q)$, with a slight abuse we will simplify the notation and indicate by $M_n(p, q)$ an arbitrary meridian-cyclic n -fold covering of S^3 branched along $K(p, q)$. This will allow us to give a unified statement for knots and links. We recall that $M_2(p, q)$ is the lens space $L(p, q)$.

Continued fractions. In the sequel we will employ continued fractions, that we define as follows:

$$[a_1, a_2, \dots, a_{k-1}, a_k] = a_1 + \frac{1}{a_2 + \dots + \frac{1}{a_{k-1} + \frac{1}{a_k}}}$$

Given p, q satisfying (1), we now recall [17, p.25] that there is a unique minimized expansion of p/q as a continued fraction with positive entries, namely an expression as $p/q = [a_1, \dots, a_k]$ with $a_1, \dots, a_{k-1} > 0$ and $a_k > 1$. (The expansion is called *minimized* because if $a_k = 1$ then $[a_1, \dots, a_{k-1}, 1] = [a_1, \dots, a_{k-1} + 1]$, as one easily sees.) We then define $l(p, q)$ to be k if $a_1 > 1$ and $k - 1$ if $a_1 = 1$.

This apparently original definition of $l(p, q)$ is explained by the following result established below (see also the proof of Proposition 2.4):

Proposition 1.1. *$l(p, q)$ is the minimum of the lengths of positive continued fraction expansions of rational numbers p'/q' such that $K(p', q')$ is equivalent to $K(p, q)$.*

REMARK 1.2. $l(p/q) = 1$ if and only if $K(p, q)$ is a torus knot (or link).

Main statements. The following will be established below:

Theorem 1.3. *Let $K(p, q)$ be a given two-bridge knot (or link) and let $(M_n(p, q))_{n=2}^\infty$ be a sequence of meridian-cyclic n -fold branched coverings of S^3 , branched along $K(p, q)$. Then:*

(2)
$$c(M_n(p, q)) \leq n(p - 1), \quad \forall n.$$

If in addition $K(p, q)$ is hyperbolic then the following inequality holds for $n \geq 7$ with $\kappa = 4$:

$$(3) \quad c(M_n(p, q)) > n \cdot \left(1 - \frac{\kappa \pi^2}{n^2}\right)^{3/2} \cdot \max\{2, 2l(p, q) - 2.6667 \dots\};$$

moreover, if $K(p, q)$ is neither $K(5, 2)$ nor $K(7, 3)$, then the inequality holds for $n \geq 6$ with $\kappa = 2\sqrt{2}$.

REMARK 1.4. Combining the inequalities (2) and (3), and letting n tend to infinity, one gets the qualitative result that the complexity of $M_n(p, q)$ is asymptotically equal to n up to a multiplicative constant.

Inequality (3) holds in vast generality but it does not appear to be numerically very effective. The next result gives a substantial improvement of the multiplicative constant appearing in the inequality. To state it, let us denote by $a_{\min}(p, q)$ be the minimal coefficient a_j appearing in the expansion $p/q = [a_1, \dots, a_k]$.

Theorem 1.5. *Let (p, q) be a pair of integers satisfying (1), such that $l(p, q) \geq 2$ and $a_{\min}(p, q) \geq 5$. For $n \in \mathbb{N}$ let $M_n(p, q)$ be an n -fold meridian-cyclic branched covering of S^3 , branched along the two-bridge knot (or link) $K(p, q)$. Then the following inequality holds:*

$$(4) \quad c(M_n(p, q)) > n \cdot \left(1 - \frac{\kappa \pi^2}{n^2}\right)^{3/2} \cdot \left(1 - \frac{2\pi^2}{1 + a_{\min}^2(p, q)^2}\right)^{3/2} \cdot 7.21985 \dots \cdot (l(p, q) - 1)$$

where $\kappa = 4$ for $n \geq 7$ and $\kappa = 2\sqrt{2}$ for $n \geq 6$.

Before stating our next result, we recall that the T-invariant $T(G)$ of a finitely presented group G was defined in [8] as the minimal number t such that G admits a presentation with t relations of length 3 and an arbitrary number of relations of length at most 2. A presentation with this property will be called *triangular*.

Proposition 1.6. *For $n \geq 2$ let $M_n(p, q)$ be a meridian-cyclic n -fold branched covering of S^3 , branched along a two-bridge knot (or link) $K(p, q)$. Then:*

$$T(\pi_1(M_n(p, q))) \leq n(p - 1).$$

We note that some connections between the Matveev complexity of a closed 3-manifold and the T-invariant of its fundamental group were already discussed in [32].

The proofs of the upper and lower complexity estimates are completely independent of each other. We will first prove the general lower inequality (3) in Section 2. Then we

will establish the upper inequality (2) (together with Proposition 1.6, which follows from the same argument) in Section 3. Next we will prove the sharper lower inequality (4) in Section 4. To conclude we will discuss in Section 5 sharper lower and upper complexity estimates for coverings of some specific knots $K(p, q)$ with $l(p, q) = 2$.

2. Hyperbolic volume and the twist number: The lower estimate

We begin by recalling that a manifold is hyperbolic if it has a Riemannian metric of constant sectional curvature -1 . We will use in the sequel many facts from hyperbolic geometry without explicit reference, see for instance [2, 5, 34].

The two versions of inequality (3) are readily deduced by combining the following three propositions. Here and always in the sequel $v_3 = 1.01494\dots$ denotes the volume of the regular ideal tetrahedron in hyperbolic 3-space \mathbb{H}^3 , and “vol(M)” is the hyperbolic volume of a manifold M . We will also need below the volume $v_8 = 3.66386\dots$ of the regular ideal octahedron in \mathbb{H}^3 .

Proposition 2.1. *If M is a closed orientable hyperbolic manifold then*

$$\text{vol}(M) < c(M) \cdot v_3.$$

Proposition 2.2. *If $K(p, q)$ is hyperbolic then $M_n(p, q)$, as defined in the statement of Theorem 1.3, is hyperbolic for $n \geq 4$. Moreover the following inequality holds for $n \geq 7$ with $\kappa = 4$:*

$$(5) \quad \text{vol}(M_n(p, q)) \geq n \cdot \left(1 - \frac{\kappa \pi^2}{n^2}\right)^{3/2} \cdot \text{vol}(S^3 \setminus K(p, q)),$$

and, if $K(p, q)$ is neither $K(5, 2)$ nor $K(7, 3)$, then the inequality holds for $n \geq 6$ with $\kappa = 2\sqrt{2}$.

Proposition 2.3. *If $K(p, q)$ is hyperbolic then*

$$(6) \quad \text{vol}(S^3 \setminus K(p, q)) \geq v_3 \cdot \max\{2, 2l(p, q) - 2.6667\dots\}.$$

We begin proofs by establishing the general connection between complexity and the hyperbolic volume:

Proof of Proposition 2.1. Set $k = c(M)$. Being hyperbolic, M is prime and not one of the exceptional manifolds S^3 , $\mathbb{R}P^3$, $S^2 \times S^1$, or $L(3, 1)$, so there exists a realization of M as a gluing of k tetrahedra. If Δ denotes the abstract tetrahedron, this realization induces continuous maps $\sigma_i: \Delta \rightarrow M$ for $i = 1, \dots, k$ given by the restrictions to the various tetrahedra of the projection from the disjoint union of the tetrahedra to M . Note that each σ_i is injective on the interior of Δ but maybe not on the boundary.

Since the gluings used to pair the faces of the tetrahedra in the construction of M are simplicial, it follows that $\sum_{i=1}^k \sigma_i$ is a singular 3-cycle, which of course represents the fundamental class $[M] \in H_3(M; \mathbb{Z})$.

We consider now the universal covering $\mathbb{H}^3 \rightarrow M$. Since Δ is simply connected, it is possible to lift σ_i to a map $\tilde{\sigma}_i: \Delta \rightarrow \mathbb{H}^3$. We then define the simplicial map $\tilde{\tau}_i: \Delta \rightarrow \mathbb{H}^3$ which agrees with $\tilde{\sigma}_i$ on the vertices, where geodesic convex combinations are used in \mathbb{H}^3 to define the notion of ‘‘simplicial’’. We also denote by $\tau_i: \Delta \rightarrow M$ the composition of $\tilde{\tau}_i$ with the projection $\mathbb{H}^3 \rightarrow M$. It is immediate to see that $\sum_{i=1}^k \tau_i$ is again a singular 3-cycle in M . Using this and taking convex combinations in \mathbb{H}^3 , one can actually check that the cycles $\sum_{i=1}^k \sigma_i$ and $\sum_{i=1}^k \tau_i$ are homotopic to each other. Therefore, since the first cycle represents $[M]$, the latter also does, which implies that $\bigcup_{i=1}^k \tau_i(\Delta)$ is equal to M , otherwise $\sum_{i=1}^k \tau_i$ would be homotopic to a map with 2-dimensional image.

Next we note that $\tilde{\tau}_i(\Delta)$ is a compact geodesic tetrahedron in \mathbb{H}^3 , so its volume is less than v_3 , see [5]. Moreover the volume of $\tau_i(\Delta)$ is at most equal to the volume of $\tilde{\tau}_i(\Delta)$, because the projection $\mathbb{H}^3 \rightarrow M$ is a local isometry, and the volume of M is at most the sum of the volumes of the $\tau_i(\Delta)$'s, because we have shown above that M is covered by the $\tau_i(\Delta)$'s (perhaps with some overlapping). This establishes the proposition. \square

Proof of Proposition 2.2. This is actually a direct application of Theorem 3.5 of [12]. We only need to note that in [12] the result is stated for hyperbolic (not necessarily two-bridge) knots (rather than links), but it is easy to see that the proof (based on [3] and Theorem 1.1 of [12]) works well also for hyperbolic two-bridge links and their meridian-cyclic coverings. \square

Before getting to the proof of Proposition 2.3 we establish the characterization of $l(p, q)$ stated in the first section:

Proof of Proposition 1.1. Under assumption (1), we know that the relevant pairs (p', q') are those with p' equal to p and q' equal to either $p - q$ or r or $p - r$, where $1 \leq r \leq p - 1$ and $q \cdot r \equiv 1 \pmod{p}$.

We begin by noting that if we take positive continued fraction expansions of p/q and $p/(p - q)$ we find 1 as the first coefficient in one case and a number greater than 1 in the other case. Supposing first that $p/q = [1, a_2, a_3, \dots, a_k]$ it is now easy to see that $p/(p - q) = [a_2 + 1, a_3, \dots, a_k]$, so the minimized positive expansion of $p/(p - q)$ has length $k - 1$. The same argument with switched roles shows that if the first coefficient a_1 of the minimized positive expansion of p/q is larger than 1 then the length of the expansion of $p/(p - q)$ is $k + 1$. Therefore the minimal length we can obtain using q and $p - q$ is indeed $l(p, q)$.

Supposing $p/q = [a_1, \dots, a_k]$, we next choose s with $1 \leq s \leq p - 1$ and $q \cdot s \equiv (-1)^{k-1} \pmod{p}$, and we note that $\{s, p - s\} = \{r, p - r\}$. Now it is not difficult to see

that p/s has a positive continued fraction expansion $p/s = [a_k, a_{k-1}, \dots, a_2, a_1]$. Note that this may or not be a minimized expansion, depending on whether a_1 is greater than 1 or equal to 1, but the length of the minimized version is $l(p, q)$ anyway, thanks to the definition we have given. By the above argument, since $a_k > 1$, the length of the minimized positive expansion of $p/(p - s)$ is 1 more than that of p/s , and the proposition is established. \square

Proof of Proposition 2.3. This will be based on results of Cao-Meyerhoff [7] and Guéritaud-Futer [14]. Note that (6) is equivalent to the two inequalities

$$(7) \quad \text{vol}(S^3 \setminus K(p, q)) \geq 2v_3$$

$$(8) \quad \text{vol}(S^3 \setminus K(p, q)) \geq v_3 \cdot (2l(p, q) - 2.6667 \dots).$$

Now, Cao and Meyerhoff have proved in [7] that the figure-eight knot complement (namely $S^3 \setminus K(5, 2)$ in our notation) and its sibling manifold (which can be described as the $(5, 1)$ -Dehn surgery on the right-handed Whitehead link) are the orientable cusped hyperbolic 3-manifolds of minimal volume, and they are the only such 3-manifolds. Each has volume equal to $2v_3 = 2.02988 \dots$, which implies inequality (7) directly.

To establish (8) we need to recall some terminology introduced by Lackenby in [20]. A *twist* in a link diagram $D \subset \mathbb{R}^2$ is either a maximal collection of bigonal regions of $\mathbb{R}^2 \setminus D$ arranged in a row, or a single crossing with no incident bigonal regions. The *twist number* $t(D)$ of D is the total number of twists in D . Moreover D is called *twist-reduced* if it is alternating and whenever $\gamma \subset \mathbb{R}^2$ is a simple closed curve meeting D transversely at two crossing only, one of the two portions into which γ separates D is contained in a twist. (This is not quite the definition in [20], but it is easily recognized to be equivalent to it for alternating diagrams.)

Lackenby proved in [20] that if D is a prime twist-reduced diagram of a hyperbolic link L in S^3 then

$$v_3 \cdot (t(D) - 2) \leq \text{vol}(S^3 \setminus L) \leq 10 \cdot v_3 \cdot (t(D) - 1),$$

where v_3 is the volume of the regular ideal tetrahedron. These estimates were improved for the case of hyperbolic two-bridge links by Guéritaud and Futer [14]. More exactly, if D is a reduced alternating diagram of a hyperbolic two-bridge link L , then by [14, Theorem B.3]

$$(9) \quad 2v_3 \cdot t(D) - 2.7066 \dots < \text{vol}(S^3 \setminus L) < 2v_8 \cdot (t(D) - 1).$$

Using the first inequality in (9), the next result implies (8), which completes the proof of Proposition 2.3 and hence of inequality (3) in Theorem 1.3:

Proposition 2.4. *The link $K(p, q)$ has a twist-reduced diagram with twist number $l(p, q)$.*




Fig. 1. The Conway normal form of a two-bridge link. The number of half-twists of the appropriate type in the j -th portion of the diagram is given by the positive integer a_j . The upper picture refers to the case of even k and the lower picture to the case of odd k .



Fig. 2. Conway diagrams of $K(23, 13)$ and $K(12, 5)$. Note that the required expansions are $23/13 = [1, 1, 3, 3]$ and $12/5 = [2, 2, 2]$.

Proof. The required diagram D is simply given by the so-called Conway normal form of $K(p, q)$ associated to the minimized positive continued fraction expansion $[a_1, \dots, a_k]$ of p/q . The definition of the Conway normal form differs for even and odd k , and it is described in Fig. 1. Two specific examples are also shown in Fig. 2.

Since the a_j 's are positive, it is quite obvious that the Conway normal diagram D always gives an alternating diagram, besides being of course prime. The twists of this diagram are almost always the obvious ones obtained by grouping together the first a_1 half-twists, then the next a_2 , and so on. An exception has to be made, however, when a_1 equals 1, because in this case the first half-twist can be grouped with the next a_2 to give a single twist (as in Fig. 2-left). Note that $a_k > 1$ by assumption, so no such phenomenon appears at the other end. Since our definition of $l(p, q)$ is precisely k if $a_1 > 1$ and $k - 1$ if $a_1 = 1$, we see that indeed the diagram always has $l(p, q)$ twists.

Before proceeding we note that if $a_1 = 1$ then the Conway normal form for $K(p, q)$ is actually the same, as a diagram, as the mirror image of the Conway normal form for $K(p, p - q)$. The picture showing this assertion gives a geometric proof of the fact that if $p/q = [1, a_2, \dots, a_k]$ then $p/(p - q) = [a_2 + 1, a_3, \dots, a_k]$, used in Proposition 1.1. So we can proceed assuming that $a_1 > 1$. In particular, each bigonal region of $S^2 \setminus D$ is one of the $(a_1 - 1) + (a_2 - 1) + \dots + (a_k - 1)$ created when inserting the a_1, a_2, \dots, a_k




Fig. 3. Labels for the non-bigonal regions of the complement of a Conway normal diagram; again k is even in the upper part of the figure and odd in the lower part.

half-twists of the normal form.

To prove that D is twist-reduced, let us look for a curve γ as in the definition, namely one that intersects D transversely at two crossings. Near each such intersection, γ must be either horizontal or vertical (see Fig. 1). Let us first show that if it meets some crossing c of D horizontally then c is the crossing arising from the single half-twist that corresponds to some coefficient a_j equal to 1. If this is not the case, then either to the left or to the right of c there is a bigonal region of $S^2 \setminus D$. Then γ must meet horizontally the crossing at the other end of this bigonal region, which readily implies that γ cannot meet the diagram in two points only.

Having shown that γ can only be vertical when it intersects vertices, except at the vertices arising from the a_j 's with $a_j = 1$, let us give labels $R_0, R_1, \dots, R_k, R_{k+1}$ to the non-bigonal regions of $S^2 \setminus D$, as in Fig. 3, and let us construct a graph Γ with vertices $R_0, R_1, \dots, R_k, R_{k+1}$ and an edge joining R_i to R_j for each segment through a crossing of D going from R_i to R_j . By assumption γ must correspond to a length-2 cycle in Γ . Now for odd k the connections existing in Γ are precisely as follows:

- an a_{2j-1} -fold connection between R_0 and R_{2j} for $j = 1, \dots, (k + 1)/2$;
- an a_{2j} -fold connection between R_1 and R_{2j+1} for $j = 1, \dots, (k - 1)/2$;
- a single connection between R_j and R_{j+2} if $2 \leq j \leq k - 1$ and $a_j = 1$.

Then the only length-2 cycles are the evident ones either between R_0 and some R_{2j} or between R_1 and some R_{2j+1} , and the curve γ corresponding to one of these cycles does bound a portion of a twist of D , as required by the definition of twist-reduced diagram.

A similar analysis for even k completes the proof. □

The proof of Proposition 2.3 is complete. □

Proof of inequality (3). Combining (7) and the first inequality in (9) with Proposition 2.4 we get

$$v_3 \cdot \max\{2, 2l(p, q) - 2.6667\dots\} < \text{vol}(S^3 \setminus K(p, q)).$$

Together with (5), this formula implies that

$$\left(1 - \frac{\kappa\pi^2}{n^2}\right)^{3/2} \cdot v_3 \cdot \max\{2, 2l(p, q) - 2.6667\dots\} \cdot n < \text{vol}(M_n(p, q))$$

with $\kappa = 4$ and $n \geq 7$ in general, and with $\kappa = 2\sqrt{2}$ and $n \geq 6$ whenever $K(p, q)$ is neither $K(5, 2)$ nor $K(7, 3)$. The conclusion now readily follows from Proposition 2.1. □

REMARK 2.5. It was pointed out by Guéritaud and Futer in [14] that the lower bound in (9) is asymptotically sharp. But it is numerically not very effective in some cases. As an example we will discuss below the case $p/q = k + 1/m$, where the lower bound given by (9), which translates into our (8), is worse than the Cao-Meyerhoff lower bound given by (7), since $l(p, q) = 2$.

REMARK 2.6. On the basis of some computer experiments, we conjecture that the Whitehead link complement (namely $S^3 \setminus K(8, 3)$, with $\text{vol}(S^3 \setminus K(8, 3)) = v_8 = 3.66386\dots$) has the smallest volume among all two-bridge two-component links.

3. Minkus polyhedral schemes and triangulations: The upper estimate

The proof of (2) and Proposition 1.6 is based on a realization of $M_n(p, q)$ as the quotient of a certain polyhedron under a gluing of its faces. This construction extends one that applies to lens spaces and it is originally due to Minkus [30]. We will briefly review it here following [31].

Let us begin from the case where $K(p, q)$ is a knot, i.e. p is odd, whence $M_n(p, q)$ is uniquely defined by p, q, n . Recall that by the assumption (1), $0 < q < p$. Then we consider the 3-ball

$$B^3 = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 \leq 1\}$$

and we draw on its boundary n equally spaced great semicircles joining the north pole $N = (0, 0, 1)$ to the south pole $S = (0, 0, -1)$. This decomposes ∂B^3 into n cyclically arranged congruent lunes L_0, \dots, L_{n-1} . Now we insert $p - 1$ equally spaced vertices on each semicircle, thus subdividing it into p identical segments, which allows us to view each lune L_i as a curvilinear polygon with $2p$ edges. Next, we denote by P_i the vertex on the semicircle $L_i \cap L_{i-1}$ which is q segments down from N , and by P'_i the vertex which is q segments up from S (indices are always meant modulo n). We then




Fig. 4. The Minkus polyhedral scheme for $M_n(p, q)$.

draw inside L_i an arc of great semicircle joining P_i to P'_{i+1} , thus bisecting L_i into two regions that we denote by R_i and R'_{i+1} , with R_i incident to N and R'_{i+1} incident to S . Fig. 4 illustrates the resulting decomposition of ∂B^3 , which is represented as $\mathbb{R}^2 \cup \{\infty\}$ with $S = \infty$. In the picture we assume $q > p/2$.

Summing up, we have subdivided ∂B^3 into $2n$ curvilinear polygons R_i, R'_i for $i = 0, \dots, n - 1$, each having $p + 1$ edges. The polygons R_i are around N and the polygons R'_i are around S , and there is a marked vertex P_i shared by R_i and R'_{i+1} (we will not need to use P'_i again). It is now possible to show that the manifold $M_n(p, q)$ is obtained from B^3 by identifying R_i with R'_i on ∂B^3 for $i = 0, \dots, n - 1$ through an orientation-reversing simplicial homeomorphism which matches the vertex P_i of R_i with the vertex P_{i-1} of R'_i .

As an example, Fig. 5 shows the Minkus polyhedral construction of the Hantzsche-Wendt manifold, that is $M_3(5, 3)$ in our notation.

Proof of inequality (2) for odd p . Referring to the above polyhedral construction of $M_n(p, q)$, we subdivide each R_i into $p - 1$ triangles by taking diagonals from the north pole N , and each R'_i so that the gluing between R_i and R'_i matches the subdivision. Note that the “diagonals” are only meant in a combinatorial sense, they cannot be taken as arcs of great circles. Since we have subdivided the R_i ’s taking diagonals from N , we can now take (combinatorial) cones with vertex at N and bases at the triangles contained in the R'_i . Note that the “lateral faces” of these cones are the triangles contained in the R_i ’s, together with some triangles in the interior of B^3 . Being based




Fig. 5. The Minkus polyhedral scheme for $M_3(5, 3)$.

on a triangle, each cone is a tetrahedron, so we have a subdivision of B^3 into $n(p - 1)$ tetrahedra. By construction the gluings on ∂B^3 restrict to gluings of the faces of these tetrahedra, therefore $M_n(p, q)$ has a (loose) triangulation made of $n(p - 1)$ tetrahedra, and the proof is now complete. \square

Proof of inequality (2) for even p . To establish (2) for even p , i.e. for 2-component two-bridge links, we extend to this case the Minkus polyhedral construction, see [31]. The way to do this is actually straight-forward: to realize the meridian-cyclic covering $M_{n,m}(p, q)$ of S^3 branched along $K(p, q)$ we subdivide ∂B^3 precisely as above, but we denote by R_i and R'_{i+m} the two regions into which the lune L_i is bisected. Then we glue R_i to R'_i by an orientation-reversing simplicial homeomorphism matching the vertex P_i of R_i with the vertex P_{i-m} of R'_i . This construction is illustrated in Fig. 6. This realization of $M_{n,m}(p, q)$ again induces a triangulation with $n(p - 1)$ tetrahedra, which proves (2) also in this case. \square

Proof of Proposition 1.6. Let us carry out only the “first half” of the subdivision we did above of the Minkus polyhedral realization of $M_n(p, q)$. Namely, we subdivide the regions R_i, R'_i on ∂B^3 into triangles, but then we do not add anything inside B^3 . This yields a cellularization of $M_n(p, q)$ with 2-cells being triangles and with a single 3-cell. Therefore there is a triangular presentation of $\pi_1(M_n(p, q))$ with precisely the same number of relations as the number of triangles in this cellularization. And this number is $n(p - 1)$, because there are $2n(p - 1)$ triangles on ∂B^3 , but they get glued in pairs. \square




Fig. 6. The Minkus polyhedral scheme for $M_{n,m}(p, q)$.

4. A sharper lower estimate

As already noticed, the lower bound on the volume given by (8) does not seem to provide very effective estimates in some instances. For this reason we discuss here a sharper lower bound, which will lead to Theorem 1.5. Its proof is based on Proposition 2.1 together with a result of Futer-Kalfagianni-Purcell [12]. To state it we associate to any two-bridge knot or link $K(p, q)$ with $\mu \in \{1, 2\}$ components a link having $l(p, q) + \mu$ components, denoted by $K_{\text{aug}}(p, q)$ and called the *augmentation* of $K(p, q)$. We only define $K_{\text{aug}}(p, q)$ for $l(p, q) \geq 2$ and to do so we change (if necessary) the pair (p, q) , without changing $K(p, q)$, so that the first coefficient a_1 in the expansion $p/q = [a_1, \dots, a_t]$ is larger than 1. This implies that $t = l(p, q)$ is the twist number of the Conway normal form of $K(p, q)$. Then we define $K_{\text{aug}}(p, q)$ by modifying $K(p, q)$ as follows:

- For all $j = 1, \dots, l(p, q)$ we encircle the two strands of $K(p, q)$ participating in the j -th sequence of half-twists of $K(p, q)$ by a small unknotted knot;
- For all $j = 1, \dots, l(p, q)$ we remove from the j -th sequence of half-twists of $K(p, q)$ as many full twists as possible.

To illustrate the definition of $K_{\text{aug}}(p, q)$ we consider the case $l(p, q) = 2$, so $p/q = k + 1/m$. Depending on the parity of k and m we get the links shown in Figs. 7 to 9. We include Rolfsen’s [33] notation and note that \mathcal{B} is the Borromean rings, a well-known hyperbolic 3-component link with volume $2v_8 = 7.32772 \dots$. It already follows




Fig. 7. $K(p, q)$ for $p/q = k + 1/m$ and $k = 2i$, $m = 2j$, and its augmentation $\mathcal{B} = 6_2^3$.




Fig. 8. $K(p, q)$ for $p/q = k + 1/m$ and $k = 2i + 1$, $m = 2j$, and its augmentation $\mathcal{B}' = 8_9^3$. Taking $k = 2i$ and $m = 2j + 1$ leads to \mathcal{B}' again.




Fig. 9. $K(p, q)$ for $p/q = k + 1/m$ and $k = 2i + 1$, $m = 2j + 1$, and its augmentation \mathcal{B}'' .

from [1] (see also [5, p.269–270]) that

$$\text{vol}(S^3 \setminus \mathcal{B}) = \text{vol}(S^3 \setminus \mathcal{B}') = \text{vol}(S^3 \setminus \mathcal{B}'')$$

(but see below for more on volume).

We are eventually ready to state [12, Proposition 3.1]:

Proposition 4.1. *If $l(p, q) \geq 2$ then $S^3 \setminus K_{\text{aug}}(p, q)$ is hyperbolic and*

$$\text{vol}(S^3 \setminus K_{\text{aug}}(p, q)) = 2v_8(l(p, q) - 1).$$

Proof of Theorem 1.5. This is now just a combination of Propositions 2.1, 2.2 and 4.1, Theorem 1.1 of [12], and the following facts:

- $K(p, q)$ is obtained by Dehn surgery on $K_{\text{aug}}(p, q)$ along the small unknotted circles, with coefficients

$$-\frac{1}{[a_1/2]}, +\frac{1}{[a_2/2]}, \dots, (-1)^l \frac{1}{[a_l(p, q)/2]};$$

- The links $K_{\text{aug}}(p, q)$ can be obtained as belted sums of the Borromean rings \mathcal{B} and their two variants \mathcal{B}' and \mathcal{B}'' , as investigated by Adams in [3];
- It follows from the results in [1] that while performing the belted sums, the sizes and shapes of the cusps relevant to our surgeries do not change;
- The geometric size and shape of a cusp in a hyperbolic link complement is determined by two linearly independent elements λ and μ of \mathbb{R}^2 , where the cusp is obtained as the quotient of \mathbb{R}^2 under the action of the lattice generated by λ and μ . Moreover λ is the holonomy of the longitude of the link component corresponding to the cusp, whereas μ is the holonomy of the meridian. Any slope on the cusp can be expressed as k times the longitude plus h times the meridian for some $k, h \in \mathbb{Z}$, and its length in the geometric cusp is the Euclidean norm of $k \cdot \lambda + h \cdot \mu$;
- One can see using SnapPea [35] that taking maximal disjoint cusps at the two “small circles” in \mathcal{B} , \mathcal{B}' and \mathcal{B}'' the holonomy of the longitude is always $(2\sqrt{2}, 0)$, while the holonomies of the meridians are given by:

$$\begin{aligned} \mu_1 &= (0, \sqrt{2}), & \mu'_1 &= (-\sqrt{2}, \sqrt{2}), & \mu''_1 &= (-\sqrt{2}, \sqrt{2}), \\ \mu_2 &= (0, \sqrt{2}), & \mu'_2 &= (0, \sqrt{2}), & \mu''_2 &= (\sqrt{2}, \sqrt{2}); \end{aligned}$$

- Even if originally obtained using numerical approximation, the information provided by SnapPea is completely reliable, having been checked using exact arithmetic in algebraic number fields with the program Snap [13]; alternatively one can work out the cusp shapes for \mathcal{B} by hand, using the fact that its hyperbolic structure is obtained by a suitable gluing of two regular ideal octahedra in hyperbolic 3-space [34], and then use the analysis in [1] for \mathcal{B}' and \mathcal{B}'' to see how the cusps change;
- Taking into account the parity of a_i and the cusp on which surgery must be performed one easily sees that the length of the surgery slope is always $\sqrt{2(1+a_i^2)}$, which is larger than 2π if $a_i \geq 5$;
- By the previous point, the lower volume estimate of [12, Theorem 1.1] applies;
- $2v_8/v_3 = 7.21985\dots$ □

5. Sharper estimates for some examples

Our upper and lower estimates for the complexity of $M_n(p, q)$ hold in general but they can be improved for specific cases. An improvement of the lower estimate was al-

ready discussed in the previous section, but it only holds asymptotically, whereas here consider definite instances.

We begin by showing that the upper bound $n(p - 1)$ for the complexity of $M_n(p, q)$ given by (2) can be significantly improved for odd p in the special case $l(p, q) = 2$ using a more specific fundamental polyhedron instead of the Minkus polyhedron. Note that if $l(p, q) = 2$ then $p/q = k + 1/m$, so $(p, q) = (km + 1, m)$.

Proposition 5.1. *Let $k, m \geq 2$ be integers. Suppose they are not both odd, so $K(km + 1, m)$ is a knot. Then, with the usual notation,*

$$(10) \quad c(M_n(km + 1, m)) \leq n \cdot (\min\{k, m\} + k + m - 3), \quad \forall n.$$

Proof. It follows from [18] that $M_n(mk + 1, m)$ can be realized by gluing together in pairs the faces of a polyhedron with $4n$ faces, half being $(k + 1)$ -edged and half being $(m + 1)$ -edged polygons. More precisely, this polyhedron is obtained by taking n polygons with $k + 1$ (respectively, $m + 1$) edges cyclically arranged around the north (respectively, south) pole of the sphere, and $2n$ polygons (n with $k + 1$ and n with $m + 1$ edges) in the remaining equatorial belt. In addition, each polygon incident to a pole is glued to one in the equatorial belt.¹ Just as in Lemma 3.1 of [29], we can now triangulate the polygons incident to the poles by taking diagonals emanating from the poles, and the polygons in the equatorial belt so that the triangulations are matched under the gluing. If we now subdivide the whole polyhedron by taking cones from the north pole, the number of tetrahedra we obtain is given by the number of triangles not incident to this pole, which is

$$n \cdot (k + 1 - 2) + 2n \cdot (m + 1 - 2) = n \cdot (k + 2m - 3).$$

Similarly, if we take cones from the south pole we get $n \cdot (2k + m - 3)$ tetrahedra, and the conclusion readily follows. □

¹As a minor fact we note that there are misprints in Figs. 1 and 2 of [18] for the case where the integers involved have different parity, and in fact the boundary patterns of F_i and \overline{F}_i do not match. Using the notation of [18], so that the integers involved are $m = 2k + 1$ and $s = 2l$, one way of fixing these figures is as follows. Keep calling $\dots, F_i, F_{i+1}, \dots$ from left to right the m -gons incident to the north pole N , so that F_i has the edges x_i on its left and x_{i+1} on its right, both emanating from N . Similarly, call $\dots, K_i, K_{i+1}, \dots$ from left to right the s -gons incident to the south pole S , so that K_i has the edges y_i on its left and y_{i+1} on its right, both emanating from S . Then the only m -gon adjacent to both F_i and K_i should be \overline{F}_{i+2} , not \overline{F}_i , while the only s -gon adjacent to both K_i and F_{i+1} should be \overline{K}_i , as in [18]. Now the boundary pattern of F_i should be given, starting from N and proceeding counterclockwise, by the word

$$x_i y_{i-1}^{-1} x_{i+2k-1}^{-1} x_{i+2k-2} \cdots x_{i+3}^{-1} x_{i+2} x_{i+1}^{-1},$$

while the boundary pattern for K_i should be given, starting from S and proceeding clockwise, by the word

$$y_i x_{i+2k-1} (y_i y_{i+1}^{-1})^{-1},$$

which allows to reconstruct the edge labelling completely.

Turning to the lower bounds, we suppose again $l(p, q) = 2$, so $(p, q) = (km + 1, m)$. Proposition 4.1 implies that

$$(11) \quad \lim_{k,m \rightarrow \infty} \text{vol}(S^3 \setminus K(km + 1, m)) = \text{vol}(S^3 \setminus \mathcal{B}) = 2v_8 = 7.32772. \dots$$

But fixing small k and m , and using the computer program SnapPea [35] to calculate the volume of $S^3 \setminus K(km + 1, m)$, one gets more specific values, and hence one can employ the usual machinery to deduce better complexity estimates. For instance, let us consider $K(5, 2) = 4_1$, the figure-eight knot, $K(7, 3) = 5_2$, and $K(9, 4) = 6_1$, where notation is again taken from [33]. Note that

$$\begin{aligned} \text{vol}(S^3 \setminus 4_1) &= 2v_3, \\ \text{vol}(S^3 \setminus 5_2) &= 2.81812. \dots, \\ \text{vol}(S^3 \setminus 6_1) &= 3.16396. \dots \end{aligned}$$

Then Propositions 2.1 and 2.2 imply the lower estimates contained in the following result, which also includes the upper estimates coming from Proposition 5.1:

Corollary 5.2. *The following bounds hold for $n \geq 7$:*

$$(12) \quad \left(1 - \frac{4\pi^2}{n^2}\right)^{3/2} \cdot 2n < c(M_n(5, 2)) \leq 3n;$$

$$(13) \quad \left(1 - \frac{4\pi^2}{n^2}\right)^{3/2} \cdot 2.77664. \dots \cdot n < c(M_n(7, 3)) \leq 4n;$$

and for $n \geq 6$:

$$(14) \quad \left(1 - \frac{2\sqrt{2}\pi^2}{n^2}\right)^{3/2} \cdot 3.11739. \dots \cdot n < c(M_n(9, 4)) \leq 5n.$$

As a matter of fact, using an explicit formula for $\text{vol}(M_n(5, 2))$ and a fundamental polyhedron with triangular faces, it was already shown in [29] that for sufficiently large n one has $2n < c(M_n(5, 2)) \leq 3n$.

Note that the general formula (2) gives $n \cdot km$ as an upper estimate for $c(M_n(km + 1, m))$, whence $4n$, $6n$ and $8n$, respectively, for the cases considered in the previous Corollary. Therefore the upper bounds $3n$, $4n$ and $5n$ in (12)–(14) are indeed stronger than those arising from (2).

ACKNOWLEDGMENT. We are grateful to Dave Futer for very useful information and discussion on the results recently obtained in [12] and [14], and to Jeff Weeks for the help he provided. We also thank the referee for suggesting us the sharper lower

bound (4) and for informing us about an independent interesting approach to lower estimates on volumes of 2-bridge knot complements announced by Ichihara [16] at a conference in February 2007, shortly after the first version of this paper was posted.

References

- [1] C.C. Adams: *Thrice-punctured spheres in hyperbolic 3-manifolds*, Trans. Amer. Math. Soc. **287** (1985), 645–656.
- [2] C.C. Adams: *Hyper Knots*, Freeman, New York, 1994.
- [3] C.C. Adams: *Waist size for cusps in hyperbolic 3-manifolds*, Topology **41** (2002), 257–270.
- [4] S. Anisov: *Exact values of complexity for an infinite number of 3-manifolds*, Mosc. Math. J. **5** (2005), 305–310, 493.
- [5] R. Benedetti and C. Petronio: *Lectures on Hyperbolic Geometry*, Springer, Berlin, 1992.
- [6] G. Burde and H. Zieschang: *Knots*, de Gruyter Studies in Mathematics **5**, de Gruyter, Berlin, 1985.
- [7] C. Cao and G.R. Meyerhoff: *The orientable cusped hyperbolic 3-manifolds of minimum volume*, Invent. Math. **146** (2001), 451–478.
- [8] T. Delzant: *Décomposition d'un groupe en produit libre ou somme amalgamée*, J. Reine Angew. Math. **470** (1996), 153–180.
- [9] R. Frigerio, B. Martelli and C. Petronio: *Complexity and Heegaard genus of an infinite class of compact 3-manifolds*, Pacific J. Math. **210** (2003), 283–297.
- [10] R. Frigerio, B. Martelli and C. Petronio: *Dehn filling of cusped hyperbolic 3-manifolds with geodesic boundary*, J. Differential Geom. **64** (2003), 425–455.
- [11] R. Frigerio, B. Martelli and C. Petronio: *Small hyperbolic 3-manifolds with geodesic boundary*, Experiment. Math. **13** (2004), 171–184.
- [12] D. Futer, E. Kalfagianni and J.S. Purcell: *Dehn filling, volume, and the Jones polynomial*, J. Differential Geom. **78** (2008), 429–464.
- [13] O. Goodman: *Snap: The computer program for studying arithmetic invariants of hyperbolic 3-manifolds*, available from <http://www.ms.unimelb.edu.au/~snap/> and from <http://sourceforge.net/projects/snap-pari>.
- [14] F. Guéritaud: *On canonical triangulations of once-punctured torus bundles and two-bridge link complements*, Geom. Topol. **10** (2006), 1239–1284.
- [15] J. Hempel: *3-Manifolds*, Ann. of Math. Studies **86**, Princeton Univ. Press, Princeton, N.J., University of Tokyo Press, Tokyo, 1976.
- [16] K. Ichihara: *Hyperbolic volumes and pant distances for two-bridge knots*, Talk at the Conference “Third East Asian School of Knots and Related Topics” held in Osaka City University, February 2007.
- [17] A. Kawachi: *A Survey of Knot Theory*, Birkhäuser, Basel, 1996.
- [18] A.C. Kim and Y. Kim: *A polyhedral description of 3-manifolds*; in *Advances in Algebra*, World Sci. Publ., River Edge, NJ., 2003, 157–162.
- [19] G. Kim, Y. Kim and A. Vesnin: *The knot S_2 and cyclically presented groups*, J. Korean Math. Soc. **35** (1998), 961–980.
- [20] M. Lackenby: *The volume of hyperbolic alternating link complements*, Proc. London Math. Soc. (3) **88** (2004), 204–224.
- [21] B. Martelli and C. Petronio: *Three-manifolds having complexity at most 9*, Experiment. Math. **10** (2001), 207–236.
- [22] B. Martelli and C. Petronio: *Complexity of geometric three-manifolds*, Geom. Dedicata **108** (2004), 15–69.
- [23] S.V. Matveev: *Complexity theory of three-dimensional manifolds*, Acta Appl. Math. **19** (1990), 101–130.

- [24] S.V. Matveev: *Algorithmic Topology and Classification of 3-Manifolds*, Algorithms and Computation in Mathematics **9**, Springer, Berlin, 2003.
- [25] S.V. Matveev: *Recognition and tabulation of three-dimensional manifolds*, Dokl. Math. **71** (2005), 20–22.
- [26] S.V. Matveev: *Tabulation of 3-manifolds up to copmlexity 12*, available from www.topology.kb.csu.ru/~recognizer.
- [27] S.V. Matveev and E.L. Pervova: *Lower bounds for the complexity of three-dimensional manifolds*, Dokl. Math. **63** (2001), 314–315.
- [28] S. Matveev, C. Petronio and A. Vesnin: *Two-sided complexity bounds for Löbell manifolds*, Dokl. Math. **76** (2007), 689–691.
- [29] S. Matveev, C. Petronio and A. Vesnin: *Two-sided asymptotic bounds for the complexity of some closed hyperbolic three-manifolds*, to appear in J. Australian Math. Soc.
- [30] J. Minkus: *The branched cyclic coverings of 2 bridge knots and links*, Mem. Amer. Math. Soc. **35** (1982).
- [31] M. Mulazzani and A. Vesnin: *The many faces of cyclic branched coverings of 2-bridge knots and links*, Atti Sem. Mat. Fis. Univ. Modena **49** (2001), 177–215.
- [32] E. Pervova and C. Petronio: *Complexity and T-invariant of Abelian and Milnor groups, and complexity of 3-manifolds*, preprint, math.GT/0412187, to appear in Math. Nachr.
- [33] D. Rolfsen: *Knots and Links*, Publish or Perish, Berkeley, CA, 1976.
- [34] W.P. Thurston: *The Geometry and Topology of 3-Manifolds*, Princeton, 1978.
- [35] J. Weeks: *SnapPea: a computer program for creating and studying hyperbolic 3-manifolds*, available from www.geometrygames.org/SnapPea/.
- [36] B. Zimmermann: *Determining knots and links by cyclic branched coverings*, Geom. Dedicata **66** (1997), 149–157.

Carlo Petronio
Dipartimento di Matematica Applicata
Università di Pisa
Via Filippo Buonarroti 1C
56127, Pisa
Italy
e-mail: petronio@dm.unipi.it

Andrei Vesnin
Sobolev Institute of Mathematics
Novosibirsk 630090
Russia
and
Department of Mathematics
Omsk Technical University
pr. Mira 11
Omsk 644050
Russia
e-mail: vesnin@math.nsc.ru