

Title	Smooth metrics on jet bundles and applications
Author(s)	Diverio, Simone
Citation	Osaka Journal of Mathematics. 2009, 46(4), p. 1019-1045
Version Type	VoR
URL	<a href="https://doi.org/10.18910/5266">https://doi.org/10.18910/5266</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

# SMOOTH METRICS ON JET BUNDLES AND APPLICATIONS

SIMONE DIVERIO

(Received July 29, 2008)

## Abstract

Following a suggestion made by J.-P. Demailly, for each  $k \geq 1$ , we endow, by an induction process, the  $k$ -th (anti)tautological line bundle  $\mathcal{O}_{X_k}(1)$  of an arbitrary complex directed manifold  $(X, V)$  with a natural smooth Hermitian metric. Then, we compute recursively the Chern curvature form for this metric, and we show that it depends (asymptotically—in a sense to be specified later) only on the curvature of  $V$  and on the structure of the fibration  $X_k \rightarrow X$ . When  $X$  is a surface and  $V = T_X$ , we give explicit formulae to write down the above curvature as a product of matrices. As an application, we obtain a new proof of the existence of global invariant jet differentials vanishing on an ample divisor, for  $X$  a minimal surface of general type whose Chern classes satisfy certain inequalities, without using a strong vanishing theorem [1] of Bogomolov.

## Contents

1. Introduction .....	1019
1.1. Main ideas and statement of the results. ....	1020
2. Projectivized jet bundles .....	1023
3. From $(X, V)$ to $(\tilde{X}, \tilde{V})$ .....	1024
4. A special choice of coordinates and local frames .....	1028
5. Curvature of $\mathcal{O}_{X_k}(1)$ and proof of Theorem 1.1 .....	1030
5.1. The case of surfaces. ....	1030
6. Holomorphic Morse inequalities for jets .....	1031
6.1. The Kähler-Einstein assumption. ....	1031
6.2. A “negative” example: quotients of the ball. ....	1033
6.3. Minimal surfaces of general type. ....	1035
7. Proof of Theorem 1.2 .....	1036
7.1. Curvature of weighted line bundles. ....	1036
7.2. End of the proof. ....	1041
References .....	1044

## 1. Introduction

In [7], Green and Griffiths showed, among other things, that if  $X$  is an algebraic surface of general type, then there exist  $m \gg k \gg 1$ , such that  $H^0(X, \mathcal{J}_{k,m}T^*X) \neq 0$ , where  $\mathcal{J}_{k,m}T^*X$  is the bundle of jet differentials of order  $k$  and degree  $m$ . Their proof relies on an asymptotic computation of the Euler characteristic  $\chi(\mathcal{J}_{k,m}T^*X)$  (which has

been possible thanks to the full knowledge of the composition series of this bundle) together with a powerful vanishing theorem of Bogomolov [1].

More precisely, if  $X$  is an  $n$ -dimensional smooth projective variety, and  $\mathcal{J}_{k,m}T^*X \rightarrow X$  is the bundle of jet differentials of order  $k$  and weighted degree  $m$ , they get the following asymptotic estimate for the holomorphic Euler characteristic:

$$\begin{aligned} \chi(\mathcal{J}_{k,m}T_X^*) &= \frac{m^{(k+1)n-1}}{(k!)^n((k+1)n-1)!} \\ &\quad \times \left( \frac{(-1)^n}{n!} c_1(X)^n (\log k)^n + O((\log k)^{n-1}) \right) + O(m^{(k+1)n-2}). \end{aligned}$$

In particular, if  $X$  is a surface of general type, then the Bogomolov vanishing theorem applies and, having cancelled the  $h^2$  term by Serre’s duality, they get a positive lower bound for  $h^0(X, \mathcal{J}_{k,m}T_X^*)$  when  $m \gg k \gg 1$ .

Nowadays, there are no general results about the existence of global *invariant* jet differentials on a surface of general type neither, of course, for varieties of general type in arbitrary dimension.

Nevertheless, thanks to a beautiful and relatively simple argument of Demailly [4], their existence should potentially lead to solve the following celebrated conjecture.

**Conjecture** (Green and Griffiths [7], Lang). *Let  $X$  be an algebraic variety of general type. Then there exist a proper algebraic sub-variety  $Y \subsetneq X$  such that every non-constant holomorphic entire curve  $f: \mathbb{C} \rightarrow X$ , has image  $f(\mathbb{C})$  contained in  $Y$ .*

A positive answer to this conjecture in dimension 2 has been given by McQuillan in [8], when the second Segre number  $c_1(X)^2 - c_2(X)$  of  $X$  is positive (this hypothesis ensures the existence of an algebraic (multi)foliation on  $X$ , whose parabolic leaves are shown to be algebraically degenerate: this is the very deep and difficult part of the proof).

**1.1. Main ideas and statement of the results.** Let  $(X, V)$  be a complex directed manifold (for precise definitions see next section) with  $\dim X = n$  and  $2 \leq \text{rank } V = r \leq n$ . Let  $\omega$  be a Hermitian metric on  $V$ . Such a metric naturally induces a smooth Hermitian metric on the tautological line bundle  $\mathcal{O}_{\tilde{X}}(-1)$  on the projectivized bundle of line of  $V$ .

Now, the Chern curvature of its dual  $\mathcal{O}_{\tilde{X}}(1)$ , is a  $(1, 1)$ -form on  $\tilde{X}$  whose restriction to the fiber over a point  $x \in X$  coincides with the Fubini-Study metric of  $P(V_x)$  with respect to  $\omega|_{V_x}$ . Thus, it is positive in the fibers direction. Next, consider the pullback  $\pi^*\omega$  on  $\tilde{X}$ : this is a  $(1, 1)$ -form which is zero in the fibers direction and, of course, positive in the base direction.

If  $X$  is compact so is  $\tilde{X}$  and hence, for all  $\varepsilon > 0$  small enough, the restriction to  $\tilde{V}$  of the  $(1, 1)$ -form given by

$$\pi^*\omega + \varepsilon^2 \Theta(\mathcal{O}_{\tilde{X}}(1))$$

gives rise to a Hermitian metric on  $\tilde{V}$ . Moreover, this metric depends on two derivatives of the metric  $\omega$ .

Of course, we can repeat this process for the compact directed manifold  $(\tilde{X}, \tilde{V})$ , and by induction, for each  $k \geq 1$  for the tower of projectivized bundles  $(X_k, V_k)$ . A priori, the Hermitian metric we obtain in this fashion on  $\mathcal{O}_{X_k}(-1)$ , depends on  $2k$  derivatives of the starting metric  $\omega$  and on the choice of  $\varepsilon^{(k)} = (\varepsilon_1, \dots, \varepsilon_{k-1})$ .

However, from a philosophical point of view, we would like to avoid the dependence on the last  $2k - 2$  derivatives of  $\omega$ , since the relevant geometrical data for  $X$  lies in the first two derivatives of  $\omega$ , namely on its Chern curvature. Here comes Demailly's suggestion: as  $\varepsilon^{(k)}$  has to be small enough, it is quite natural to look for an asymptotic expression of the Chern curvature of the metric on  $\mathcal{O}_{X_k}(-1)$  we have constructed, when  $\varepsilon^{(k)}$  tends to zero: this idea is developed in our first theorem.

**Theorem 1.1.** *The vector bundle  $V_k$  can be endowed inductively with a smooth Hermitian metric*

$$\omega^{(k)} = (\pi_k^* \omega^{(k-1)} + \varepsilon_k^2 \Theta(\mathcal{O}_{X_k}(1)))|_{V_k},$$

where the metric on  $\mathcal{O}_{X_k}(1)$  is induced by  $\omega^{(k-1)}$ , depending on  $k - 1$  positive real numbers  $\varepsilon^{(k)} = (\varepsilon_1, \dots, \varepsilon_{k-1})$ , such that the asymptotic of its Chern curvature with respect to this metric depends only on the curvature of  $V$  and on the (universal) structure of the fibration  $X_k \rightarrow X$ , as  $\varepsilon^{(k)} \rightarrow 0$ .

As a byproduct of the proof of the above theorem, we also obtain induction formulae for an explicit expression of the curvature in terms of the curvature coefficients of  $V$ . These formulae, which are quite difficult to handle in higher dimension, are reasonably simple for  $X$  a smooth surface: in this case, it turns out that the curvature coefficients of  $\mathcal{O}_{X_k}(-1)$  are given by a sequence of products of  $2 \times 2$  real matrices.

A general remark in analytic geometry is that the existence of global sections of a Hermitian line bundle is strictly correlated with the positivity properties of its Chern curvature form. One of the countless correlations, is given by the theory of Demailly's holomorphic Morse inequalities [2]. We summarize his main result here below.

**1.1.1. Holomorphic Morse inequalities.** Let  $X$  be a compact Kähler manifold of dimension  $n$ ,  $E$  a holomorphic vector bundle of rank  $r$  and  $L$  a line bundle over  $X$ . If  $L$  is equipped with a smooth metric of curvature form  $\Theta(L)$ , we define the  $q$ -index set of  $L$  to be the open subset

$$X(q, L) = \left\{ x \in X \mid i\Theta(L) \text{ has } \begin{array}{l} q \text{ negative eigenvalues} \\ n - q \text{ positive eigenvalues} \end{array} \right\},$$

for  $q = 0, \dots, n$ . Hence  $X$  admits a partition  $X = \Delta \cup \bigcup_{q=0}^n X(q, L)$ , where  $\Delta = \{x \in X \mid \det(i\Theta(L)) = 0\}$  is the degeneracy set. We also introduce

$$X(\leq q, L) \stackrel{\text{def}}{=} \bigcup_{j=0}^q X(j, L).$$

It was shown by Demailly in [2], that the partial alternating sums of the dimension of the cohomology groups of tensor powers of  $L$  with values in  $E$  satisfy the following asymptotic *strong Morse inequalities* as  $k \rightarrow +\infty$ :

$$\sum_{j=0}^q (-1)^{q-j} h^j(X, L^{\otimes k} \otimes E) \leq r \frac{k^n}{n!} \int_{X(\leq q, L)} (-1)^q \left( \frac{i}{2\pi} \Theta(L) \right)^n + O(k^{n-1}).$$

In particular, if

$$\int_{X(\leq 1, L)} \left( \frac{i}{2\pi} \Theta(L) \right)^n > 0,$$

then some high power of  $L$  twisted by  $E$  has a (many, in fact) nonzero section.

The idea is now to apply holomorphic Morse inequalities to the anti-tautological line bundle  $\mathcal{O}_{X_k}(1)$  together with the asymptotic Hermitian metric constructed above, to find global sections of invariant  $k$ -jet differentials on a surface  $X$ : we shall deal with the absolute case  $V = T_X$ . Our first geometrical hypothesis is to suppose  $X$  to be Kähler-Einstein, that is with ample canonical bundle. Nevertheless, standard arguments coming from the theory of Monge-Ampère equations, will show that we just need to assume  $X$  to be minimal and of general type, that is  $K_X$  big and numerically effective. Finally, once sections are found, we can drop the hypothesis of nefness, since the dimension of the space of global section of jet differentials is a birational invariant (see, for instance, [7] and [3]).

For each  $k \geq 1$ , in  $\mathbb{R}^k$  define the closed convex cone  $\mathfrak{N} = \{\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{R}^k \mid a_j \geq 2 \sum_{l=j+1}^k a_l \text{ for all } j = 1, \dots, k-1 \text{ and } a_k \geq 0\}$ . For  $X$  a smooth compact surface, set

$$\mathcal{O}_{X_k}(\mathbf{a})^{k+2} = F_k(\mathbf{a})c_1(X)^2 - G_k(\mathbf{a})c_2(X)$$

(see next section for the definition of the weighted line bundle  $\mathcal{O}_{X_k}(\mathbf{a})$ ) and

$$m_k = \sup_{\mathbf{a} \in \mathfrak{N} \setminus \Sigma_k} \frac{F_k(\mathbf{a})}{G_k(\mathbf{a})},$$

where  $\Sigma_k$  is the zero locus of  $G_k$ . Finally, call  $m_\infty$  the supremum of the sequence  $\{m_k\}$ .

**Theorem 1.2.** *Notations as above, the two following facts can occur: either*

- *there exists a  $k_0 \geq 1$  such that for every surface  $X$  of general type,  $\mathcal{O}_{X_{k_0}}(1)$  is big, or*
- *the sequence  $\{m_k\}$  is positive non-decreasing and for  $X$  a surface of general type, there exists a positive integer  $k$  such that  $\mathcal{O}_{X_k}(1)$  is big as soon as  $m_\infty > c_2(\hat{X})/c_1(\hat{X})^2$ , where  $\hat{X}$  is the minimal model of  $X$ .*

As a corollary, we obtain the existence of low order jet differentials, for  $X$  a minimal surface of general type whose Chern classes satisfy certain inequalities. This will be done in §7.2.1.

**2. Projectivized jet bundles**

Let  $(X, V)$  be a complex directed manifold, that is a pair consisting in a smooth complex manifold  $X$  and a holomorphic (non necessarily integrable) subbundle  $V \subset T_X$  of the tangent bundle. Set  $\tilde{X} = P(V)$ . Here,  $P(V)$  is the projectivized bundle of lines of  $V$  and there is a natural projection  $\pi: \tilde{X} \rightarrow X$ ; moreover, if  $\dim X = n$ , then  $\dim \tilde{X} = n+r-1$ , if  $\text{rank } V = r$ . On  $\tilde{X}$ , we consider the tautological line bundle  $\mathcal{O}_{\tilde{X}}(-1) \subset \pi^*V$  which is defined fiberwise as

$$\mathcal{O}_{\tilde{X}}(-1)_{(x,[v])} \stackrel{\text{def}}{=} \mathbb{C}v,$$

for  $(x, [v]) \in \tilde{X}$ , with  $x \in X$  and  $v \in V_x \setminus \{0\}$ . Next, set  $\tilde{V} = \pi_*^{-1}\mathcal{O}_{\tilde{X}}(-1)$ , where  $\pi_*: T_{\tilde{X}} \rightarrow \pi^*T_X$  is the differential of the projection: this is a holomorphic subbundle of  $T_{\tilde{X}}$  of rank  $r$ , so that we obtain in this way a new directed manifold  $(\tilde{X}, \tilde{V})$ .

Now, we start the inductive process in the directed manifold category by setting

$$(X_0, V_0) = (X, V), \quad (X_k, V_k) = (\tilde{X}_{k-1}, \tilde{V}_{k-1}).$$

In other words,  $(X_k, V_k)$  is obtained from  $(X, V)$  by iterating  $k$  times the projectivization construction  $(X, V) \mapsto (\tilde{X}, \tilde{V})$  described above.

In this process, the rank of  $V_k$  remains constantly equal to  $r$  while the dimension of  $X_k$  grows linearly with  $k$ :  $\dim X_k = n + k(r - 1)$ . Let us call  $\pi_k: X_k \rightarrow X_{k-1}$  the natural projection. Then we have, as before, a tautological line bundle  $\mathcal{O}_{X_k}(-1) \subset \pi_k^*V_{k-1}$  over  $X_k$  which fits into short exact sequences

$$(1) \quad 0 \rightarrow T_{X_k/X_{k-1}} \rightarrow V_k \xrightarrow{(\pi_k)_*} \mathcal{O}_{X_k}(-1) \rightarrow 0$$

and

$$(2) \quad 0 \rightarrow \mathcal{O}_{X_k} \rightarrow \pi_k^*V_{k-1} \otimes \mathcal{O}_{X_k}(1) \rightarrow T_{X_k/X_{k-1}} \rightarrow 0,$$

where  $T_{X_k/X_{k-1}} = \ker(\pi_k)_*$  is the relative tangent bundle.

More generally, if  $\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}^k$  is a weight, we can form the line bundle  $\mathcal{O}_{X_k}(\mathbf{a})$  by setting

$$\mathcal{O}_{X_k}(\mathbf{a}) = \bigotimes_{j=1}^k \pi_{j,k}^* \mathcal{O}_{X_j}(a_j).$$

We shall see later how, for appropriate choices of  $\mathbf{a}$ , one can obtain relatively positive line bundles  $\mathcal{O}_{X_k}(\mathbf{a})$  which, moreover, admit a non-trivial morphism to  $\mathcal{O}_{X_k}(a_1 + \dots + a_k)$  (for the last assertion see for example [3]). In particular, sections of  $\mathcal{O}_{X_k}(\mathbf{a})$  for a suitable choice of  $\mathbf{a}$  give rise to sections of  $\mathcal{O}_{X_k}(m)$  for some large  $m$ .

### 3. From $(X, V)$ to $(\tilde{X}, \tilde{V})$

Let  $(X, V)$  be a compact directed manifold of complex dimension  $n$  and  $\text{rank } V = r$ . In this section, given a Hermitian metric  $\omega$  on  $V$ , we construct a (family of) metric on  $\tilde{V}$  depending on a “small” positive constant  $\varepsilon$ , and we compute the curvature of  $\tilde{V}$  with respect to this metric, letting  $\varepsilon$  tend to zero.

So, fix a Hermitian metric  $\omega$  on  $V$ , a point  $x_0 \in X$  and a unit vector  $v_0 \in V_{x_0}$  with respect to  $\omega$ . Then there exist coordinates  $(z_1, \dots, z_n)$  centered at  $x_0$  and a holomorphic normal local frame  $e_1, \dots, e_r$  for  $V$  such that  $e_r(x_0) = v_0$  and

$$\omega(e_\lambda, e_\mu) = \delta_{\lambda,\mu} - \sum_{j,k=1}^n c_{jk\lambda\mu} z_j \bar{z}_k + O(|z|^3).$$

Remark that, as  $V$  is a holomorphic subbundle of the holomorphic tangent space of  $X$ , then there exists a holomorphic matrix  $(g_{i\lambda}(z))$  such that  $e_\lambda(z) = \sum_{i=1}^n g_{i\lambda}(z) \partial/\partial z_i$ .

Moreover, the Chern curvature at  $x_0$  of  $V$  is expressed by

$$\Theta(V)_{x_0} = \sum_{j,k=1}^n \sum_{\lambda,\mu=1}^r c_{jk\lambda\mu} dz_j \wedge d\bar{z}_k \otimes e_\lambda^* \otimes e_\mu.$$

Now consider the projectivized bundle  $\pi: P(V) = \tilde{X} \rightarrow X$  of lines in  $V$ : its points can be seen as pairs  $(x, [v])$  where  $x \in X$ ,  $v \in V_x \setminus \{0\}$  and  $[v] = \mathbb{C}v$ . In a neighborhood of  $(x_0, [v_0]) \in \tilde{X}$  we have local holomorphic coordinates given by  $(z, \xi_1, \dots, \xi_{r-1})$  where  $\xi$  corresponds to the direction  $[\xi_1 e_1(z) + \dots + \xi_{r-1} e_{r-1}(z) + e_r(z)]$  in  $V_x$ .

On  $\tilde{X}$  we have a tautological line bundle  $\mathcal{O}_{\tilde{X}}(-1) \subset \pi^* V$  such that the fiber over  $(x, [v])$  is simply  $[v]$ : then  $\mathcal{O}_{\tilde{X}}(-1) \subset \pi^* V$  inherits a metric from  $V$  in such a way that its local non vanishing section  $\eta(z, \xi) = \xi_1 e_1(z) + \dots + \xi_{r-1} e_{r-1}(z) + e_r(z)$  has squared length

$$\begin{aligned} |\eta|_\omega^2 &= 1 + |\xi|^2 - \sum_{j,k,\lambda,\mu} c_{jk\lambda\mu} z_j \bar{z}_k \xi_\lambda \bar{\xi}_\mu - \sum_{j,k,\lambda} c_{jk\lambda r} z_j \bar{z}_k \xi_\lambda \\ &\quad - \sum_{j,k,\mu} c_{jkr\mu} z_j \bar{z}_k \bar{\xi}_\mu - \sum_{j,k} c_{jkr r} z_j \bar{z}_k + O(|z|^3). \end{aligned}$$

So we have

$$\begin{aligned} \bar{\partial}|\eta|_{\omega}^2 &= \sum_{\mu} \xi_{\mu} d\bar{\xi}_{\mu} - \sum_{j,k,\lambda,\mu} c_{jk\lambda,\mu} z_j \bar{z}_k \xi_{\lambda} d\bar{\xi}_{\mu} \\ &\quad - \sum_{j,k} c_{jkrr} z_j d\bar{z}_k + O((|z| + |\xi|)^2 |dz| + |z|^2 |d\xi|), \\ \partial|\eta|_{\omega}^2 &= \sum_{\lambda} \bar{\xi}_{\lambda} d\xi_{\lambda} - \sum_{j,k,\lambda,\mu} c_{jk\lambda,\mu} z_j \bar{z}_k \bar{\xi}_{\mu} d\xi_{\lambda} \\ &\quad - \sum_{j,k} c_{jkrr} \bar{z}_k dz_j + O((|z| + |\xi|)^2 |dz| + |z|^2 |d\xi|), \\ \partial\bar{\partial}|\eta|_{\omega}^2 &= \sum_{\lambda} d\xi_{\lambda} \wedge d\bar{\xi}_{\lambda} - \sum_{j,k,\lambda,\mu} c_{jk\lambda,\mu} z_j \bar{z}_k d\xi_{\lambda} \wedge d\bar{\xi}_{\mu} - \sum_{j,k=1}^n c_{jkrr} dz_j \wedge d\bar{z}_k \\ &\quad + O((|z| + |\xi|)|dz|^2 + |z| |dz| |d\xi| + (|z| + |\xi|)^3 |d\xi|^2), \end{aligned}$$

where all the summations here are taken with  $j, k = 1, \dots, n$  and  $\lambda, \mu = 1, \dots, r - 1$ . We remark that inside the  $O$ 's there are hidden terms which are useless for our further computations. We finally obtain

$$\begin{aligned} \Theta(\mathcal{O}_{\tilde{X}}(1)) &= \partial\bar{\partial} \log |\eta|_{\omega}^2 = -\frac{1}{|\eta|_{\omega}^4} \partial|\eta|_{\omega}^2 \wedge \bar{\partial}|\eta|_{\omega}^2 + \frac{1}{|\eta|_{\omega}^2} \partial\bar{\partial}|\eta|_{\omega}^2 \\ &= \sum_{\lambda,\mu} \left( -\xi_{\mu} \bar{\xi}_{\lambda} - \sum_{j,k} c_{jk\lambda,\mu} z_j \bar{z}_k \right. \\ &\quad \left. + \delta_{\lambda,\mu} \left( 1 - |\xi|^2 + \sum_{j,k} c_{jkrr} z_j \bar{z}_k \right) \right) d\xi_{\lambda} \wedge d\bar{\xi}_{\mu} \\ &\quad - \sum_{j,k} c_{jkrr} dz_j \wedge d\bar{z}_k \\ &\quad + O((|z| + |\xi|)|dz|^2 + |z| |dz| |d\xi| + (|z| + |\xi|)^3 |d\xi|^2). \end{aligned}$$

So we get in particular

$$\Theta(\mathcal{O}_{\tilde{X}}(1))_{(x_0, [v_0])} = \sum_{\lambda=1}^{r-1} d\xi_{\lambda} \wedge d\bar{\xi}_{\lambda} - \sum_{j,k=1}^n c_{jkrr} dz_j \wedge d\bar{z}_k,$$

which shows the well known fact that

$$\Theta(\mathcal{O}_{\tilde{X}}(1))_{(x_0, [v_0])} = |\bullet|_{\mathbb{F}\mathbb{S}}^2 - \theta_{V, x_0}(\bullet \otimes v_0, \bullet \otimes v_0),$$

where FS denotes the Fubini-Study metric along the vertical tangent space  $\ker \pi_*$  and  $\theta_{V, x_0}$  is the natural Hermitian form on  $T_X \otimes V$  corresponding to  $i\Theta(V)$ , at the point  $x_0$ .



Now consider the rank  $r$  holomorphic subbundle  $\tilde{V}$  of  $T_{\tilde{X}}$  whose fiber over a point  $(x, [v])$  is given by

$$\tilde{V}_{(x,[v])} = \{ \tau \in T_{\tilde{X}} \mid \pi_* \tau \in \mathbb{C}v \}.$$

To start with, let's consider the holomorphic local frame of  $\tilde{V}$  given by  $\partial/\partial\xi_1, \dots, \partial/\partial\xi_{r-1}, \tilde{\eta}$ , where

$$\tilde{\eta}(z, \xi) = \sum_{i=1}^n \left( g_{ir}(z) + \sum_{\lambda=1}^{r-1} g_{i\lambda}(z)\xi_\lambda \right) \frac{\partial}{\partial z_i},$$

so that  $\tilde{\eta}$  formally is equal to  $\eta$  but here, with a slight abuse of notations, the  $\partial/\partial z_i$  are regarded as tangent vector fields to  $\tilde{X}$  (so,  $\tilde{\eta}$  actually means a lifting of  $\eta$  from  $\mathcal{O}_{\tilde{X}}(-1) \subset \pi^*V \subset \pi^*T_X$  to  $T_{\tilde{X}}$ , which admits  $\pi^*T_X$  as a quotient). For all sufficiently small  $\varepsilon > 0$  we get an Hermitian metric on  $\tilde{V}$  by restricting  $\tilde{\omega}_\varepsilon = \pi^*\omega + \varepsilon^2\Theta(\mathcal{O}_{\tilde{X}}(1))$  to  $\tilde{V}$ ; at the point  $(x_0, [v_0]) = (0, 0)$  with this choice of local coordinates, we have

$$\begin{aligned} \tilde{\omega}_\varepsilon \left( \frac{\partial}{\partial \xi_\lambda}, \frac{\partial}{\partial \xi_\mu} \right) &= \underbrace{\pi^* \omega \left( \frac{\partial}{\partial \xi_\lambda}, \frac{\partial}{\partial \xi_\mu} \right)}_{=0} + \varepsilon^2 \Theta(\mathcal{O}_{\tilde{X}}(1)) \left( \frac{\partial}{\partial \xi_\lambda}, \frac{\partial}{\partial \xi_\mu} \right) \\ &= \delta_{\lambda,\mu} \varepsilon^2, \\ \tilde{\omega}_\varepsilon \left( \frac{\partial}{\partial \xi_\lambda}, \tilde{\eta} \right) &= \underbrace{\pi^* \omega \left( \frac{\partial}{\partial \xi_\lambda}, \tilde{\eta} \right)}_{=0} + \varepsilon^2 \Theta(\mathcal{O}_{\tilde{X}}(1)) \left( \frac{\partial}{\partial \xi_\lambda}, \tilde{\eta} \right) \\ &= 0, \quad \text{since } \Theta(\mathcal{O}_{\tilde{X}}(1))_{(x_0,[v_0])}(\partial/\partial \xi_\lambda, \partial/\partial z_i) = 0, \end{aligned}$$

and

$$\begin{aligned} \tilde{\omega}_\varepsilon(\tilde{\eta}, \tilde{\eta}) &= \pi^* \omega(\tilde{\eta}, \tilde{\eta}) + \varepsilon^2 \Theta(\mathcal{O}_{\tilde{X}}(1))(\tilde{\eta}, \tilde{\eta}) \\ &= |\eta(0, 0)|_\omega^2 + O(\varepsilon^2) = 1 + O(\varepsilon^2). \end{aligned}$$

We now renormalize this local frame of  $\tilde{V}$  by setting

$$f_1 = \frac{1}{\varepsilon} \frac{\partial}{\partial \xi_1}, \dots, f_{r-1} = \frac{1}{\varepsilon} \frac{\partial}{\partial \xi_{r-1}}, f_r = C_\varepsilon \tilde{\eta},$$

where

$$C_\varepsilon = \frac{1}{\sqrt{\tilde{\omega}_\varepsilon(\tilde{\eta}, \tilde{\eta})}} = 1 + O(\varepsilon).$$

Then  $(f_\lambda)$  is unitary at  $(x_0, [v_0])$  with respect to  $\tilde{\omega}_\varepsilon$  and we have

$$\tilde{\omega}_\varepsilon(f_\lambda, f_\mu) = \begin{cases} -\xi_\mu \bar{\xi}_\lambda - \sum_{j,k} c_{jk\lambda\mu} z_j \bar{z}_k + \delta_{\lambda,\mu} \left( 1 - |\xi|^2 + \sum_{j,k} c_{jkr r} z_j \bar{z}_k \right), & \text{if } 1 \leq \lambda, \mu \leq r-1, \\ 0, & \text{if } 1 \leq \lambda \leq r-1 \text{ and } \mu = r \text{ or } 1 \leq \mu \leq r-1 \text{ and } \lambda = r, \\ |\eta|_\omega^2, & \text{if } \lambda = \mu = r, \end{cases}$$

modulo  $\varepsilon$  and terms of order three in  $z$  and  $\xi$ .

Next, we compute the curvature

$$\Theta(\tilde{V})_{(x_0, [v_0])} = \sum_{j,k=1}^{n+r-1} \sum_{\lambda,\mu=1}^r \gamma_{jk\lambda\mu} dz_j \wedge d\bar{z}_k \otimes f_\lambda^* \otimes f_\mu$$

for  $\varepsilon \rightarrow 0$ , where we have set  $z_{n+\lambda} = \xi_\lambda$ . Recall that for a Hermitian vector bundle  $E \rightarrow Y$ , given a holomorphic trivialization, the curvature operator at a point  $0 \in Y$  is given by

$$\Theta(E)_0 = \bar{\partial}(\bar{H}^{-1} \partial \bar{H})(0) = (\bar{\partial} \bar{H}^{-1})(0) \wedge (\partial \bar{H})(0) + \bar{H}^{-1}(0) (\bar{\partial} \partial \bar{H})(0),$$

where  $H$  is the Hermitian matrix of Hermitian products between the elements of the local frame. If the local holomorphic frame is unitary in  $0$ , so that  $H(0) = \text{Id}$ , observing that  $0 = \bar{\partial}(\bar{H}^{-1} \bar{H}) = (\bar{\partial} \bar{H}^{-1})(0) \bar{H}(0) + \bar{H}^{-1}(0) (\bar{\partial} \bar{H})(0)$ , we obtain

$$(3) \quad \Theta(E)_0 = -\bar{\partial} \bar{H}(0) \wedge \partial \bar{H}(0) + \bar{\partial} \partial \bar{H}(0).$$

Thus, in our case, it suffices to compute the part with second derivatives in (3) to get the following proposition.

**Proposition 3.1.** *Notations as given, the Chern curvature of  $\tilde{V}$  has the following expression:*

$$(4) \quad \begin{aligned} \Theta(\tilde{V}) = & \sum_{\lambda,\mu=1}^{r-1} \left( d\xi_\mu \wedge d\bar{\xi}_\lambda + \sum_{j,k=1}^n c_{jk\lambda\mu} dz_j \wedge d\bar{z}_k \right. \\ & \left. + \delta_{\lambda,\mu} \left( \sum_{v=1}^{r-1} d\xi_v \wedge d\bar{\xi}_v - \sum_{j,k=1}^n c_{jkr r} dz_j \wedge d\bar{z}_k \right) \right) \otimes f_\lambda^* \otimes f_\mu \\ & + \left( \sum_{j,k=1}^n c_{jkr r} dz_j \wedge d\bar{z}_k - \sum_{v=1}^{r-1} d\xi_v \wedge d\bar{\xi}_v \right) \otimes f_r^* \otimes f_r + O(\varepsilon). \end{aligned}$$

In particular, we get the following identities modulo  $\varepsilon$ :

$$\gamma_{jk\lambda\mu} = \begin{cases} c_{jk\lambda\mu} - \delta_{\lambda\mu}c_{jkrr}, & \text{if } 1 \leq j, k \leq n \text{ and } 1 \leq \lambda, \mu \leq r - 1, \\ \delta_{\lambda\mu}\delta_{jk} + \delta_{(j-n)\mu}\delta_{(k-n)\lambda}, & \text{if } n + 1 \leq j, k \leq n + r - 1 \text{ and } 1 \leq \lambda, \mu \leq r - 1, \end{cases}$$

$$\gamma_{jkrr} = \begin{cases} c_{jkrr}, & \text{if } 1 \leq j, k \leq n, \\ -1, & \text{if } n < j = k \leq n + r - 1, \end{cases}$$

the remaining coefficients being zero.

#### 4. A special choice of coordinates and local frames

We now pass to the tower of projective bundles  $(X_k, V_k)$  over  $(X, V)$ . We recall that we simply set  $(X, V) = (X_0, V_0)$  and, for all integer  $k > 0$ ,  $(X_k, V_k) = (\tilde{X}_{k-1}, \tilde{V}_{k-1})$  together with the projection  $\pi_{k-1,k}: X_k \rightarrow X_{k-1}$  so that the total fibration is given by  $\pi_{0,k} = \pi_{0,1} \circ \pi_{1,2} \circ \dots \circ \pi_{k-1,k}: X_k \rightarrow X$ .

For all  $k$ , we also have a tautological line bundle  $\mathcal{O}_{X_k}(-1)$  and a metric  $\omega^{(k)} = \omega^{(k)}(\varepsilon_1, \dots, \varepsilon_k)$  on  $V_k$ , with the  $\varepsilon_l$ 's positive and small enough, obtained recursively by setting  $\omega^{(k)} = (\pi_{k-1,k}^* \omega^{(k-1)} + \varepsilon_k^2 \Theta(\mathcal{O}_{X_k}(1)))|_{V_k}$ ,  $\omega^{(0)} = \omega$ .

To start with, fix a point  $x_0 \in X$ , a  $\omega$ -unitary vector  $v_0 \in V$  and a holomorphic local normal frame  $(e_\lambda^{(0)})$  for  $(V, \omega)$  such that  $e_r(x_0) = v_0$ .

**First step.** On  $X_1$ , we have local holomorphic coordinates centered at  $(x_0, [v_0])$  given by  $(z, \xi^{(1)})$  where,  $(z, \xi) \mapsto [\xi_1^{(1)}e_1^{(0)}(z) + \dots + \xi_{r-1}^{(1)}e_{r-1}^{(0)}(z) + e_r^{(0)}(z)] \in P(V_z)$ . Recall that we have, as before, a ‘‘natural’’ local section  $\eta_1$  of  $\mathcal{O}_{X_1}(-1)$  given by

$$\eta_1(z, \xi^{(1)}) = \xi_1^{(1)}e_1(z) + \dots + \xi_{r-1}^{(1)}e_{r-1}(z) + e_r(z)$$

and for all  $\varepsilon_1 > 0$  small enough, a holomorphic local frame  $(f_\lambda^{(1)})$  for  $V_1$  near  $(x_0, [v_0])$  which is a  $\omega^{(1)}$ -unitary basis for  $V_{1(x_0, [v_0])}$ .

Now, choose a  $\omega^{(1)}$ -unitary vector  $v_1 \in V_{1(x_0, [v_0])}$  and a holomorphic local normal frame  $(e_\lambda^{(1)})$  for  $V_1$  such that  $e_r^{(1)}(x_0, [v_0]) = v_1$ . Then there exist a unitary  $r \times r$  matrix  $U_1 = (a_{\lambda\mu}^{(1)})$  such that at  $(x_0, [v_0])$  we have

$$f_\mu^{(1)} = \sum_{\lambda=1}^r a_{\lambda\mu}^{(1)} e_\lambda^{(1)}.$$

So, if we call respectively  $\gamma_{ij\lambda\mu}^{(1)}$  and  $c_{ij\lambda\mu}^{(1)}$  the coefficients of curvature of  $V_1$  at  $(x_0, [v_0])$  with respect to the basis  $(f_\lambda^{(1)})$  and  $(e_\lambda^{(1)})$  we have

$$c_{ij\lambda\mu}^{(1)} = \sum_{\alpha, \beta=1}^r \gamma_{ij\alpha\beta}^{(1)} \bar{a}_{\lambda\alpha}^{(1)} a_{\mu\beta}^{(1)},$$

with  $i, j = 1, \dots, n + (r - 1)$  and  $\lambda, \mu = 1, \dots, r$ .

**General step.** For the general case, suppose for all  $\varepsilon_1, \dots, \varepsilon_{k-1} > 0$  small enough we have built a system of holomorphic coordinates  $(z, \xi^{(1)}, \dots, \xi^{(k-1)})$  for  $X_{k-1}$  and a holomorphic local normal frame  $(e_\lambda^{(k-1)})$  for  $(V_{k-1}, \omega^{(k-1)})$ ,  $k \geq 2$ , such that  $e_r^{(k-1)}(x_0, [v_0], \dots, [v_{k-2}]) = v_{k-1}$  where  $v_{k-1}$  is a  $\omega^{(k-1)}$ -unitary vector in  $V_{k-1}(x_0, [v_0], \dots, [v_{k-2}])$ . Our procedure gives us also a holomorphic local frame  $(f_\lambda^{(k-1)})$  for  $V_{k-1}$  near  $(x_0, [v_0], \dots, [v_{k-2}])$  which is a  $\omega^{(k-1)}$ -unitary basis for  $V_{k-1}(x_0, [v_0], \dots, [v_{k-2}])$  and a unitary  $r \times r$  matrix  $U_{k-1} = (a_{\lambda\mu}^{(k-1)})$  such that

$$f_\mu^{(k-1)} = \sum_{\lambda=1}^r a_{\lambda\mu}^{(k-1)} e_\lambda^{(k-1)}.$$

Then, we put holomorphic local coordinates  $(z, \xi^{(1)}, \dots, \xi^{(k)})$  on  $X_k$  centered at the point  $(x_0, [v_0], \dots, [v_{k-1}])$  where

$$(z, \xi^{(1)}, \dots, \xi^{(k)}) \mapsto [\xi_1^{(k)} e_1^{(k-1)}(z, \dots, \xi^{(k-1)}) + \dots + \xi_{r-1}^{(k)} e_{r-1}^{(k-1)}(z, \dots, \xi^{(k-1)}) + e_r^{(k-1)}(z, \dots, \xi^{(k-1)})] \in P(V_{k-1}(z, \dots, \xi^{(k-1)}))$$

and also

$$\eta_k(z, \xi^{(1)}, \dots, \xi^{(k)}) = \xi_1^{(k)} e_1^{(k-1)}(z, \dots, \xi^{(k-1)}) + \dots + \xi_{r-1}^{(k)} e_{r-1}^{(k-1)}(z, \dots, \xi^{(k-1)}) + e_r^{(k-1)}(z, \dots, \xi^{(k-1)})$$

is a local nonzero section of  $\mathcal{O}_{X_k}(-1)$ .

As we have already done, if we call

$$f_\lambda^{(k)} = \frac{1}{\varepsilon_k} \frac{\partial}{\partial \xi_\lambda^{(k)}}, \quad \lambda = 1, \dots, r - 1,$$

$$f_r^{(k)} = C_{\varepsilon_k}^{(k)} \tilde{\eta}_k,$$

where  $C_{\varepsilon_k}^{(k)} = 1/\sqrt{\omega^{(k)}(\tilde{\eta}_k, \tilde{\eta}_k)} = 1 + \mathcal{O}(\varepsilon_k)$ , then  $(f_\lambda^{(k)})$  is a local holomorphic frame for  $V_k$ , unitary at  $(x_0, [v_0], \dots, [v_{k-1}])$ . We now fix a  $\omega^{(k)}$ -unitary vector  $v_k \in V_k(x_0, [v_0], \dots, [v_{k-1}])$  and choose a holomorphic local normal frame  $(e_\lambda^{(k)})$  for  $(V_k, \omega^{(k)})$  such that  $e_\lambda^{(k)}(x_0, [v_0], \dots, [v_{k-1}]) = v_k$  and a  $r \times r$  unitary matrix  $U_k = (a_{\lambda\mu}^{(k)})$  such that  $f_\mu^{(k)} = \sum_{\lambda=1}^r a_{\lambda\mu}^{(k)} e_\lambda^{(k)}$ .

So, if we call respectively  $\gamma_{jk\lambda\mu}^{(k)}$  and  $c_{jk\lambda\mu}^{(k)}$  the coefficients of curvature of  $V_k$  at  $(x_0, [v_0], \dots, [v_{k-1}])$  with respect to the basis  $(f_\lambda^{(k)})$  and  $(e_\lambda^{(k)})$  we have

$$(5) \quad c_{ij\lambda\mu}^{(k)} = \sum_{\alpha, \beta=1}^r \gamma_{ij\alpha\beta}^{(k)} \bar{a}_{\lambda\alpha}^{(k)} a_{\mu\beta}^{(k)},$$

with  $i, j = 1, \dots, n + k(r - 1)$  and  $\lambda, \mu = 1, \dots, r$ .

**5. Curvature of  $\mathcal{O}_{X_k}(1)$  and proof of Theorem 1.1**

We now use (4) and (5) to get the induction formulae to derive an expression for the curvature of  $\mathcal{O}_{X_k}(1)$ , when  $\varepsilon^{(k)} = (\varepsilon_1, \dots, \varepsilon_{k-1})$  tends to zero.

We start by observing that (4) shows how  $\gamma_{ij\lambda\mu}^{(s)}$  depends on  $c_{lm\alpha\beta}^{(s-1)}$ ; we rewrite here the dependence modulo  $\varepsilon_s$ :

$$(6) \quad \begin{aligned} \gamma_{ij\lambda\mu}^{(s)} &= \begin{cases} c_{ij\lambda\mu}^{(s-1)} - \delta_{\lambda\mu} c_{ijrr}^{(s-1)}, & \text{if } 1 \leq i, j \leq n+(s-1)(r-1) \text{ and } 1 \leq \lambda, \mu \leq r-1, \\ \delta_{jk} \delta_{\lambda\mu} + \delta_{(i-n-(s-1)(r-1))\mu} \delta_{(j-n-(s-1)(r-1))\lambda}, & \text{if } i, j \geq n+(s-1)(r-1)+1, \ i, j \leq n+s(r-1) \text{ and } 1 \leq \lambda, \mu \leq r-1, \end{cases} \\ \gamma_{ijrr}^{(s)} &= \begin{cases} c_{ijrr}^{(s-1)}, & \text{if } 1 \leq i, j \leq n+(s-1)(r-1), \\ -1, & \text{if } n+(s-1)(r-1)+1 \leq i = j \leq n+s(r-1), \end{cases} \end{aligned}$$

the remaining coefficients being zero. Recall also that, by (5),

$$c_{ij\lambda\mu}^{(s)} = \sum_{\alpha, \beta=1}^r \gamma_{ij\alpha\beta}^{(s)} \bar{a}_{\lambda\alpha}^{(s)} a_{\mu\beta}^{(s)}.$$

Now, we have

$$(7) \quad \Theta(\mathcal{O}_{X_k}(1))_{(x_0, [v_0], \dots, [v_{k-1}])} = \sum_{\lambda=1}^{r-1} d\xi_{\lambda}^{(k)} \wedge d\bar{\xi}_{\lambda}^{(k)} - \sum_{i,j=1}^{n+(k-1)(r-1)} c_{ijrr}^{(k-1)} dz_i \wedge d\bar{z}_j,$$

where we have set  $z_{n+(s-1)(r-1)+\lambda} = \xi_{\lambda}^{(s)}$ ,  $\lambda = 1, \dots, r-1$ , and to get the expression of this curvature with respect to the coefficients of curvature of  $V$  it suffices to perform the recursive substitutions (5) and (6) and to stop with  $c_{ij\lambda\mu}^{(0)} = c_{ij\lambda\mu}$ .

Thus, Theorem 1.1 is proved.

**5.1. The case of surfaces.** In the case  $\text{rank } V = \dim X = 2$ , we have a nice matrix representation of these formulae. First of all, note that in this case the identities (6) become much simpler:

$$\begin{aligned} \gamma_{ij11}^{(s)} &= \begin{cases} c_{ij11}^{(s-1)} - c_{ij22}^{(s-1)}, & \text{if } 1 \leq i, j \leq s+1, \\ 2, & \text{if } i = j = s+2, \end{cases} \\ \gamma_{ij22}^{(s)} &= \begin{cases} c_{ij22}^{(s-1)}, & \text{if } 1 \leq i, j \leq s+1, \\ -1, & \text{if } i = j = s+2. \end{cases} \end{aligned}$$

Now, for each  $s \geq 1$ , let  $v_s = v_s^1 f_1^{(s)} + v_s^2 f_2^{(s)}$ , with  $|v_s^1|^2 + |v_s^2|^2 = 1$ . Then we have  $a_{21}^{(s)} = \bar{v}_s^1$  and  $a_{22}^{(s)} = \bar{v}_s^2$  and so, for instance  $a_{11}^{(s)} = -v_s^2$  and  $a_{12}^{(s)} = v_s^1$  would work. It follows that

$$c_{ij11}^{(s)} = \gamma_{ij11}^{(s)} |v_s^2|^2 + \gamma_{ij22}^{(s)} |v_s^1|^2, \quad c_{ij22}^{(s)} = \gamma_{ij11}^{(s)} |v_s^1|^2 + \gamma_{ij22}^{(s)} |v_s^2|^2.$$

So, if we set

$$R_s = \begin{pmatrix} |v_s^2|^2 & |v_s^1|^2 \\ |v_s^1|^2 & |v_s^2|^2 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, \quad C_{ij}^{(s)} = \begin{pmatrix} c_{ij11}^{(s)} \\ c_{ij22}^{(s)} \end{pmatrix}$$

we have that

$$C_{ij}^{(s)} = \begin{cases} R_s \cdot T \cdot R_{s-1} \cdot T \cdot \dots \cdot R_1 \cdot T \cdot C_{ij}^{(0)}, & \text{if } 1 \leq i, j \leq 2, \\ R_s \cdot T \cdot R_{s-1} \cdot T \cdot \dots \cdot R_{i-2} \cdot T \cdot \begin{pmatrix} 1 \\ -1 \end{pmatrix}, & \text{if } 3 \leq i = j \leq s + 2 \end{cases}$$

and we are interested in the second element of the vector  $C_{ij}^{(k-1)}$ : in fact, in the surface absolute case, formula (7) can be rewritten in the form

$$(8) \quad \Theta(\mathcal{O}_{X_k}(1)) = d\xi^{(k)} \wedge d\bar{\xi}^{(k)} - \sum_{s=3}^{k+1} c_{ss22}^{(k-1)} d\xi^{(s-2)} \wedge d\bar{\xi}^{(s-2)} - \sum_{i,j=1}^2 c_{ij22}^{(k-1)} dz_i \wedge d\bar{z}_j.$$

We shall see in the next sections how this explicit formulae can be use to compute Morse-type integrals, in order to obtain the existence of nonzero global section of the bundle of invariant jet differentials.

### 6. Holomorphic Morse inequalities for jets

Let  $X$  be a smooth surface and  $V = T_X$ . From now on we will suppose that  $K_X$  is ample, so that we can take as a metric on  $X$  the Kähler-Einstein one, and we will work always modulo  $\varepsilon^k$  (this will be possible thanks to Lebesgue’s dominated convergence theorem).

**6.1. The Kähler-Einstein assumption.** So, let  $K_X$  be ample. Then we have a unique Hermitian metric  $\omega$  on  $T_X$ , such that  $\text{Ricci}(\omega) = -\omega$  and, for this metric,

$$\text{Vol}_\omega(X) = \frac{\pi^2}{2} c_1^2(X) > 0,$$

where

$$\text{Vol}_\omega(X) \stackrel{\text{def}}{=} \int_X \frac{\omega^2}{2!}.$$

Now, consider the two Hermitian matrices  $(c_{ij11})$  and  $(c_{ij22})$ . The Kähler-Einstein assumption implies that

$$(c_{ij11}) + (c_{ij22}) = (-\delta_{ij})$$

and so they are simultaneously diagonalizable. Let

$$\begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix}$$

be a diagonal form for  $(c_{ij11})$ . Then

$$\lambda + \mu = c_{1111} + c_{2211} = c_{1111} + c_{1122} = -1$$

thanks to the Kähler symmetries. If the eigenvalues of  $(c_{ij22})$  are  $\lambda', \mu'$  then  $\lambda' + \lambda = \mu + \mu' = -1$ , thus a diagonal form for  $(c_{ij22})$  is

$$\begin{pmatrix} \mu & 0 \\ 0 & \lambda \end{pmatrix}.$$

As a consequence, for  $\alpha, \beta \in \mathbb{C}$ , the eigenvalues of the matrix  $\alpha(c_{ij11}) + \beta(c_{ij22})$  are  $\alpha\lambda + \beta\mu$  and  $\alpha\mu + \beta\lambda$  and so

$$\begin{aligned} \det(\alpha(c_{ij11}) + \beta(c_{ij22})) &= (\alpha\lambda + \beta\mu)(\alpha\mu + \beta\lambda) \\ &= \alpha\beta(\lambda^2 + \mu^2) + \lambda\mu(\alpha^2 + \beta^2) \\ (9) \quad &= \alpha\beta[(\lambda + \mu)^2 - 2\lambda\mu] + \lambda\mu(\alpha^2 + \beta^2) \\ &= \alpha\beta + \lambda\mu(\alpha - \beta)^2 \end{aligned}$$

and

$$\begin{aligned} \text{tr}(\alpha(c_{ij11}) + \beta(c_{ij22})) &= (\alpha\lambda + \beta\mu) + (\alpha\mu + \beta\lambda) \\ (10) \quad &= (\alpha + \beta)(\lambda + \mu) \\ &= -(\alpha + \beta). \end{aligned}$$

For  $k = 1$ , the curvature of  $\mathcal{O}_{X_1}(1)$  is simply

$$\Theta(\mathcal{O}_{X_1}(1)) = d\xi^{(1)} \wedge d\bar{\xi}^{(1)} - \sum_{i,j=1}^2 c_{ij22} dz_i \wedge d\bar{z}_j$$

and so

$$\left(\frac{i}{2\pi}\Theta(\mathcal{O}_{X_1}(1))\right)^3 = 3!\left(\frac{i}{2\pi}\right)^3 D dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge d\xi^{(1)} \wedge d\bar{\xi}^{(1)},$$

where we have set  $D \stackrel{\text{def}}{=} \lambda\mu$ , which is, of course, a function  $X_1 \rightarrow \mathbb{R}$ . In particular,

$$\begin{aligned} &\int_{X_1} \left(\frac{i}{2\pi}\right)^3 D dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge d\xi^{(1)} \wedge d\bar{\xi}^{(1)} \\ &= \frac{1}{6} \int_{X_1} \left(\frac{i}{2\pi}\Theta(\mathcal{O}_{X_1}(1))\right)^3 \\ &= \frac{1}{6}(c_1^2(X) - c_2(X)), \end{aligned}$$

in fact, this integral over  $X_1$  is just the top self-intersection of  $c_1(\mathcal{O}_{X_1}(1))$ , and this is easily seen to be  $c_1^2(X) - c_2(X)$  by means of exact sequences (1) and (2).

Moreover, we have that

$$\left(\frac{i}{2\pi}\right)^3 dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge d\xi^{(1)} \wedge d\bar{\xi}^{(1)} = \pi_{0,1}^* \left(\frac{1}{\pi^2} dV_\omega\right) \wedge \left(\frac{i}{2\pi}\Theta(\mathcal{O}_{X_1}(1))\right),$$

so that

$$\begin{aligned} &\int_{X_1} \left(\frac{i}{2\pi}\right)^3 dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge d\xi^{(1)} \wedge d\bar{\xi}^{(1)} \\ &= \int_{X_1} \pi_{0,1}^* \left(\frac{1}{\pi^2} dV_\omega\right) \wedge \left(\frac{i}{2\pi}\Theta(\mathcal{O}_{X_1}(1))\right) \\ &= \frac{1}{2}c_1^2(X) \end{aligned}$$

by Fubini.

**6.2. A “negative” example: quotients of the ball.** Here, we wish to make an example to clarify why, if we deal with smooth metrics, we have to use the relatively nef weighted line bundles introduced above.

Suppose you want to show, using just  $\mathcal{O}_{X_k}(1)$ , the existence of global  $k$ -jet differentials on a surface  $X$ . From our point of view, a good possible “test” case is when  $X$  is a compact unramified quotient of the unit ball  $\mathbb{B}_2 \subset \mathbb{C}^2$ ; surfaces which arise in this way are Kähler-Einstein, hyperbolic and with ample cotangent bundle: the best one can hope (these surfaces have even lots of symmetric differentials).



So, let  $\mathbb{B}_2 = \{z \in \mathbb{C}^2 \mid |z| < 1\}$  endowed with the Poincaré metric

$$\begin{aligned} \omega_P &= -\frac{i}{2} \partial \bar{\partial} \log(1 - |z|^2) \\ &= \frac{i}{2} \left( \frac{dz \otimes d\bar{z}}{1 - |z|^2} + \frac{|\langle dz, z \rangle|^2}{(1 - |z|^2)^2} \right). \end{aligned}$$

Consider a compact unramified quotient  $X = \mathbb{B}_2/\Gamma$  with the quotient metric, say  $\omega$ . Then,  $\omega$  has constant curvature; in particular, the function  $D: X_1 \rightarrow \mathbb{R}$  we defined in §6.1 is constant.

This constant can be quite easily directly computed by hands. Here, we shall compute it as a very simple application of the celebrated Bogomolov-Miyaoka-Yau inequality  $c_1^2 \leq 3c_2$  for surfaces of general type with ample canonical bundle, which says moreover that the equality holds if and only if the surface is a quotient of the ball  $\mathbb{B}_2$ .

Using computations made in §6.1, we have

$$\begin{aligned} \frac{1}{6}(c_1(X)^2 - c_2(X)) &= \int_{X_1} \left( \frac{i}{2\pi} \right)^3 D dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge d\xi^{(1)} \wedge d\bar{\xi}^{(1)} \\ &= D \int_{X_1} \left( \frac{i}{2\pi} \right)^3 dz_1 \wedge d\bar{z}_1 \wedge \cdots \wedge d\xi^{(1)} \wedge d\bar{\xi}^{(1)} \\ &= \frac{1}{2}c_1(X)^2 D, \end{aligned}$$

so that, making the substitution  $c_1(X)^2 = 3c_2(X)$ , we find  $D \equiv 2/9$ .

Now, a somewhat tedious computation of the 1-index set of our curvatures leads to the following result for the “Morse” integrals for  $\mathcal{O}_{X_k}(1)$  and low values of  $k$ , using the new information about  $D$ .

$k = 1$  We have already done this integral in §6.1: in this special case it gives  $(2/3)c_1(X)^2 > 0$  and so the existence of 1-jet differentials.

$k = 2$  In this case (the line bundle is no longer relatively nef) we don’t have the equality  $(X_2, \leq 1) = X_2$  and so we have to determine the open set  $(X_2, \leq 1)$ . This is an easy matter: using notations of §5.1 and setting moreover  $|v_1^1|^2 = x$ ,  $0 \leq x \leq 1$ , one sees from the expression of the curvature that

$$(X_2, \leq 1) = \left\{ 0 < x < \frac{2}{3} \right\},$$

since the trace of the “horizontal” part is always positive for  $k = 2$ . Then we have

$$\begin{aligned} &\int_{(X_2, \leq 1)} \left( \frac{i}{2\pi} \Theta(\mathcal{O}_{X_2}(1)) \right)^4 \\ &= 4! \left( \frac{i}{2\pi} \right)^4 \int_{(X_2, \leq 1)} (1 - 3x) \left( -\frac{1}{3}x + \frac{2}{9} \right) dz_1 \wedge \cdots \wedge d\bar{\xi}^{(2)} \end{aligned}$$

$$\begin{aligned}
 &= 4! \left(\frac{i}{2\pi}\right)^3 \int_{X_1} dz_1 \wedge \cdots \wedge d\bar{\xi}^{(1)} \int_0^{2/3} (1-3x) \left(-\frac{1}{3}x + \frac{2}{9}\right) dx \\
 &= 4! \left(\frac{i}{2\pi}\right)^3 \int_{X_1} \frac{2}{81} dz_1 \wedge \cdots \wedge d\bar{\xi}^{(1)} \\
 &= \frac{8}{27} c_1(X)^2 > 0,
 \end{aligned}$$

where we make the substitution in the  $\xi^{(2)}$ -complex plane

$$\frac{i}{2\pi} d\xi^{(2)} \wedge d\bar{\xi}^{(2)} \mapsto \frac{dx d\vartheta}{2\pi}.$$

Hence the existence of 2-jet differentials (the optimal attended result should be  $(10/27)c_1(X)^2$ , by replacing  $c_2(X) = (1/3)c_1(X)^2$  in the expression of the leading term of the Euler characteristic  $\chi(E_{2,m}T_X^*)$  of the bundle of invariant 2-jet differentials on  $X$ , see [3]).

$k = 3$  Here the situation becomes much more involved. Several computations (which can be found in our PhD thesis [6]) give

$$\int_{X_3(\leq 1, \mathcal{O}_{X_3}(1))} \left(\frac{i}{2\pi} \Theta(\mathcal{O}_{X_3}(1))\right)^5 = - \underbrace{\frac{715933}{1944000}}_{\approx 0.37} c_1(X)^2 < 0,$$

and so we are not able to check the existence of 3-jet differentials by this method. Here are some considerations.

First, the value of the above integrals is, at least in these first cases, decreasing while morally one should expect an increasing sequence (the existence of  $k$ -jet differentials implies obviously the existence of  $(k + 1)$ -jet differentials).

Second, we suspect that, in fact, this sequence continues to be non-increasing in general, since going up with  $k$ , adds more and more regions of negativity along the fiber direction ( $\mathcal{O}_{X_k}(1)$  is not relatively positive over  $X$ , for  $k \geq 2$ ). Moreover, recall that we are working here on a quotient of the ball, so that we had the most favorable “horizontal” contribution in terms of positivity: thus, the problem really relies in the fibers direction.

From these considerations, we deduce that to get a Green-Griffiths type result about asymptotic (on  $k$ ) existence of section, we are naturally led to study either the smooth relatively nef line case (weighted line bundles  $\mathcal{O}_{X_k}(\mathbf{a})$ ), or to leave the “smooth world” and to study singular Hermitian metrics on  $\mathcal{O}_{X_k}(1)$  which reflects the relative base locus of this bundle.

The rest of this paper will be devoted to the first of these two different approaches.

**6.3. Minimal surfaces of general type.** If we relax the hypothesis on the canonical bundle of the surface  $X$ , and we just take it to be big and nef, then our previous

computation gives the same results.

To see this, it suffices to select an ample class  $A$  on  $X$  and, for every  $\varepsilon > 0$ , to solve the “approximate” Kähler-Einstein equation  $\text{Ricci}(\omega) = -\omega + \delta\Theta(A)$  (the existence of such a metric  $\omega$  on  $T_X$  is a well-known consequence of the theory of Monge-Ampère equations).

Once we have such a metric we just observe that, with the notations of this section, we have  $\lambda + \mu = -1 + O(\delta)$ , so that

$$\det(\alpha(c_{ij11}) + \beta(c_{ij22})) = \alpha\beta(1 + O(\delta)) + \lambda\mu(\alpha^2 - \beta^2)$$

and

$$\text{tr}(\alpha(c_{ij11}) + \beta(c_{ij22})) = -(\alpha + \beta)(1 - O(\delta)).$$

It is then clear that our integral computation will now have a final error term which is in fact a  $O(\delta)$ , and thus we obtain the same results, by letting  $\delta$  tend to zero.

### 7. Proof of Theorem 1.2

In this section we compute explicitly the Chern curvature of the weighted line bundles  $\mathcal{O}_{X_k}(\mathbf{a})$  on a surface  $X$  and we find conditions for them to be relatively positive. Next, thanks to holomorphic Morse inequalities, we study the consequences of positive self-intersection and finally we prove Theorem 1.2.

**7.1. Curvature of weighted line bundles.** We recall some notations and formulae. Let  $v_s = v_s^1 f_1^{(s)} + v_s^2 f_2^{(s)} \in V_s$ , with  $|v_s^1|^2 + |v_s^2|^2 = 1$  and set  $x_s = |v_s^1|^2$ ,  $0 \leq x_s \leq 1$ . Then, if

$$R_s = \begin{pmatrix} 1 - x_s & x_s \\ x_s & 1 - x_s \end{pmatrix}, \quad T = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$$

and

$$R_p \cdot T \cdot \dots \cdot R_q \cdot T = \begin{pmatrix} \delta_{p,q} & \gamma_{p,q} \\ \beta_{p,q} & \alpha_{p,q} \end{pmatrix}, \quad p \geq q \geq 1,$$

where  $\alpha_{p,q}$ ,  $\beta_{p,q}$ ,  $\gamma_{p,q}$  and  $\delta_{p,q}$  are functions of  $(x_q, \dots, x_p)$ , we have that, for  $k \geq 2$ ,

$$\begin{aligned} \Theta(\mathcal{O}_{X_k}(1)) &= d\xi^{(k)} \wedge d\bar{\xi}^{(k)} + \sum_{s=1}^{k-1} (\alpha_{k-1,s} - \beta_{k-1,s}) d\xi^{(s)} \wedge d\bar{\xi}^{(s)} \\ &\quad + \sum_{i,j=1}^2 (-\beta_{k-1,1} c_{ij11} - \alpha_{k-1,1} c_{ij22}) dz_i \wedge d\bar{z}_j. \end{aligned}$$

More generally, for  $\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}^k$  (or possibly  $\in \mathbb{R}^k$ ), we have

$$\begin{aligned}
 \Theta(\mathcal{O}_{X_k}(a_1, \dots, a_k)) &= a_k d\xi^{(k)} \wedge d\bar{\xi}^{(k)} + \sum_{s=1}^{k-1} \left( \sum_{j=s}^{k-1} a_{j+1} y_{j,s} + a_s \right) d\xi^{(s)} \wedge d\bar{\xi}^{(s)} \\
 (11) \quad &+ \underbrace{\sum_{i,j=1}^2 \left( \sum_{l=0}^{k-1} -a_{l+1} \beta_{l,1} c_{ij11} - a_{l+1} \alpha_{l,1} c_{ij22} \right)}_{\stackrel{\text{def}}{=} A_{ij}(a_1, \dots, a_k)} dz_i \wedge d\bar{z}_j,
 \end{aligned}$$

where  $y_{p,q}(x_q, \dots, x_p) \stackrel{\text{def}}{=} \alpha_{p,q} - \beta_{p,q}$  (we also set formally  $\alpha_{0,1} = \beta_{0,1} = 1$ ). Observe that, for the  $(2 \times 2)$ -matrix  $(A_{ij})$ , we have

$$\text{tr}(A_{ij}) = \sum_{s=0}^{k-1} a_{s+1} w_{s,1}, \quad w_{p,q}(x_q, \dots, x_p) \stackrel{\text{def}}{=} \alpha_{p,q} + \beta_{p,q},$$

thanks to the Kähler-Einstein assumption and formula (10).

Now, define  $\theta_s^k = \theta_s^k(x_s, \dots, x_{k-1})$  to be the function given by

$$(x_s, \dots, x_{k-1}) \mapsto \sum_{j=s}^{k-1} a_{j+1} y_{j,s} + a_s.$$

This is the  $s$ -th “vertical” eigenvalue of the weighted curvature.

REMARK 7.1. As the  $\theta_s^k$ 's are linear combinations of the  $y_{j,s}$ 's, we have that they all are of degree one in each variable. Hence they and their restriction to each edge of the cube  $[0, 1]^{k-s}$  are harmonic. In particular they attain their minimum on some vertex of this cube.

In  $\mathbb{R}^k$ , define the closed convex cone

$$\mathfrak{N} = \left\{ \mathbf{a} \in \mathbb{R}^k \mid a_j \geq 2 \sum_{l=j+1}^k a_l, \forall j = 1, \dots, k-1 \text{ and } a_k \geq 0 \right\}.$$

We have the following three lemmas.

**Lemma 7.1.** *The functions  $\theta_s^k$  are positive if (and only if)  $\mathbf{a} \in \mathring{\mathfrak{N}}$ .*

Proof. First of all, observe that the structure of the four functions  $\alpha_{p,q}$ ,  $\beta_{p,q}$ ,  $\gamma_{p,q}$  and  $\delta_{p,q}$  (and hence  $y_{p,q}$ ) depends only on  $p - q$ . Now, it is immediate to check by induction that we have the following expression for the  $\gamma_{p,q}$ 's and the  $\delta_{p,q}$ 's:

$$\gamma_{p,q} = - \sum_{h=q}^p \alpha_{h,q}$$

and

$$\delta_{p,q} = 1 - \sum_{h=q}^p \beta_{h,q}.$$

Next, observe that for all  $s \geq 1$ ,

$$R_s(0) \cdot T = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$$

and

$$R_s(1) \cdot T = \begin{pmatrix} 0 & 1 \\ 1 & -1 \end{pmatrix},$$

so that, if  $j \geq 1$  then  $y_{j+1,1}(\bullet, 0) = y_{j,1}(\bullet)$  and  $y_{j+1,1}(\bullet, 1) = -1 - 2y_{j,1}(\bullet) - \sum_{h=1}^{j-1} y_{h,1}(\bullet)$ ; moreover,  $y_{1,1}(0) = 1$  and  $y_{1,1}(1) = -2$ .

The lemma is clearly true for  $k = 1$ , so we proceed by induction on  $k$ . We have, for  $s \geq 2$ ,

$$\begin{aligned} \theta_s^k &= \theta_s^k(x_s, \dots, x_{k-1}; \mathbf{a}) = a_s + \sum_{j=s}^{k-1} a_{j+1} y_{j,s} \\ &= a_s + \sum_{j=s}^{k-1} a_{j+1} y_{j-s+1,1} = \theta_1^{k-s+1}(x_s, \dots, x_{k-1}; \mathbf{b}), \end{aligned}$$

where  $\mathbf{b} = (a_s, \dots, a_k) \in \mathbb{R}^{k-s+1}$  is again in the corresponding  $\mathfrak{H}$ : it remains then to show that, for a general  $k \geq 2$ , the lemma is true for  $\theta_1^k$ . Recall that, by Remark 7.1, it suffices to check positivity on the vertices of the cube  $[0, 1]^{k-1}$ . Let  $\star$  denote an arbitrary sequence of 0 and 1 of length  $k - 2$ : we shall treat the two cases  $(\star, 0)$  and  $(\star, 1)$  separately. For the first one, we have

$$\begin{aligned} \theta_1^k(\star, 0; \mathbf{a}) &= a_1 + \sum_{j=1}^{k-1} a_{j+1} y_{j,1}(\star, 0) \\ &= a_1 + \sum_{j=1}^{k-2} a_{j+1} y_{j,1}(\star) + a_k y_{k-1,1}(\star, 0) \end{aligned}$$

$$\begin{aligned}
 &= a_1 + \sum_{j=1}^{k-2} a_{j+1}y_{j,1}(\star) + a_k y_{k-2,1}(\star) \\
 &= a_1 + \sum_{j=1}^{k-3} a_{j+1}y_{j,1}(\star) + (a_{k-1} + a_k)y_{k-2,1}(\star) \\
 &= \theta_1^{k-1}(\star; \mathbf{b}')
 \end{aligned}$$

for a new  $\mathbf{b}' \in \mathbb{R}^{k-1}$  which is easily seen to be in the corresponding  $\mathfrak{N}$ . Similarly, for the second case, we have

$$\begin{aligned}
 \theta_1^k(\star, 1; \mathbf{a}) &= a_1 + \sum_{j=1}^{k-1} a_{j+1}y_{j,1}(\star, 1) \\
 &= a_1 + \sum_{j=1}^{k-2} a_{j+1}y_{j,1}(\star) + a_k y_{k-1,1}(\star, 1) \\
 &= a_1 + \sum_{j=1}^{k-2} a_{j+1}y_{j,1}(\star) + a_k \left( - \sum_{h=1}^{k-3} y_{h,1}(\star) - 2y_{k-2,1}(\star) - 1 \right) \\
 &= (a_1 - a_k) + \sum_{j=1}^{k-3} (a_{j+1} - a_k)y_{j,1}(\star) + (a_{k-1} - 2a_k)y_{k-2,1}(\star) \\
 &= \theta_1^{k-1}(\star; \mathbf{b}''),
 \end{aligned}$$

where again  $\mathbf{b}'' \in \mathbb{R}^{k-1}$  is a new weight which satisfies the (strict) inequalities defining  $\mathfrak{N}$ . The lemma is proved. □

The reason why we choose  $\mathbf{a}$  in the interior of the cone  $\mathfrak{N}$ , is that with such a choice the vertical eigenvalues of the curvature  $\Theta(\mathcal{O}_{X_k}(\mathbf{a}))$  are positive for all sufficiently small  $\varepsilon^{(k)}$ .

**REMARK 7.2.** The above lemma says in particular that if  $\mathbf{a} \in \mathfrak{N}$ , then for all  $\varepsilon > 0$  we can endow  $\mathcal{O}_{X_k}(\mathbf{a})$  with a smooth Hermitian metric  $h_k$  (namely, the one we are working with) such that  $\Theta_{h_k}(\mathcal{O}_{X_k}(\mathbf{a})) \geq -\varepsilon\omega$  along the fiber of  $X_k \rightarrow X$ , for some Hermitian metric  $\omega$  on  $T_{X_k}$  (recall that we are always working modulo  $\varepsilon^{(k)}$ ). In particular, the cone  $\mathfrak{N}$  is contained in the cone of relatively nef (over  $X$ ) line bundles.

**Lemma 7.2.** *If  $\theta_s^k \geq 0$  for all  $s = 1, \dots, k - 1$ , and  $\mathbf{a} \in \mathbb{N}^k$  with at least one of the  $a_j$ 's strictly positive, then  $\text{tr}(A_{ij}) > 0$  in the cube  $[0, 1]^{k-1}$ .*

Proof. First of all, we recover the expression of the  $w_{p,q}$ 's in terms of the  $y_{r,s}$ 's. We have  $w_{p,p} = (2 + y_{p,p})/3$  and

$$\begin{aligned} w_{p,j-1} &= \alpha_{p,j-1} + \beta_{p,j-1} = x_{j-1}(\beta_{p,j} - \alpha_{p,j}) + \alpha_{p,j} \\ &= \alpha_{p,j} - y_{p,j}x_{j-1} = \alpha_{p,j} + \frac{y_{p,j-1} + 2\beta_{p,j} - \alpha_{p,j}}{3y_{p,j}}y_{p,j} \\ &= \frac{y_{p,j-1} + 2w_{p,j}}{3}, \end{aligned}$$

as,  $x_{j-1} = -(y_{p,j-1} + 2\beta_{p,j} - \alpha_{p,j})/3y_{p,j}$  (this is easily seen from the very definitions). Then, by induction, we obtain

$$w_{p,q} = \left(\frac{2}{3}\right)^{p-q+1} + \frac{1}{3} \sum_{l=q}^p \left(\frac{2}{3}\right)^{l-q} y_{p,l}.$$

Now,  $\text{tr}(A_{ij}) = \sum_{s=0}^{k-1} a_{s+1}w_{s,1}$  and so

$$\begin{aligned} \sum_{s=0}^{k-1} a_{s+1}w_{s,1} &= a_1 + \sum_{s=1}^{k-1} \left( \left(\frac{2}{3}\right)^s + \frac{1}{3} \sum_{l=1}^s \left(\frac{2}{3}\right)^{l-1} y_{s,l} \right) \\ &= a_1 + \sum_{s=1}^{k-1} \left(\frac{2}{3}\right)^s a_{s+1} + \frac{1}{3} \sum_{l,s=1}^{k-1} \left(\frac{2}{3}\right)^{l-1} a_{s+1}y_{s,l} \\ &= a_1 + \sum_{s=1}^{k-1} \left(\frac{2}{3}\right)^s a_{s+1} + \frac{1}{3} \sum_{l=1}^{k-1} \left(\frac{2}{3}\right)^{l-1} (\theta_l^k - a_l) \\ &= \frac{2}{3}a_1 + \left(\frac{2}{3}\right)^{k-1} a_k + \sum_{s=1}^{k-2} \left(\frac{2}{3}\right)^{s+1} a_{s+1} + \frac{1}{3} \sum_{l=1}^{k-1} \left(\frac{2}{3}\right)^{l-1} \theta_l^k. \quad \square \end{aligned}$$

**Lemma 7.3.** *Let  $D: X_1 \rightarrow \mathbb{R}$  be as in §6.1. If  $\theta_s^k \geq 0$  for all  $s = 1, \dots, k - 1$ ,  $\mathbf{a} \in \mathbb{N}^k$  with at least one of the  $a_j$ 's strictly positive and  $D \equiv 2/9$ , then  $\det(A_{ij}) > 0$  in the cube  $[0, 1]^{k-1}$ .*

Proof. Set

$$\alpha(\mathbf{a}) \stackrel{\text{def}}{=} a_1 + \sum_{l=1}^{k-1} a_{l+1}\alpha_{l,1}$$

and

$$\beta(\mathbf{a}) \stackrel{\text{def}}{=} a_1 + \sum_{l=1}^{k-1} a_{l+1}\beta_{l,1}.$$

Then, formula (9) yields

$$\begin{aligned} \det(A_{ij}) &= \alpha(\mathbf{a})\beta(\mathbf{a}) + (\alpha(\mathbf{a}) - \beta(\mathbf{a}))^2 D \\ &= \frac{1}{4}(\alpha(\mathbf{a}) + \beta(\mathbf{a}))^2 - \frac{1}{36}(\alpha(\mathbf{a}) - \beta(\mathbf{a}))^2. \end{aligned}$$

But now, we observe that  $\alpha(\mathbf{a}) + \beta(\mathbf{a}) = \text{tr}(A_{ij})$  and that  $\alpha(\mathbf{a}) - \beta(\mathbf{a}) = \theta_1^k$ ; the end of the proof of Lemma 7.2 shows that  $\text{tr}(A_{ij}) > (1/3)\theta_1^k$ , so that  $\det(A_{ij}) > 0$ .  $\square$

**Proposition 7.4.** *If  $\mathbf{a} \in \mathfrak{N}$ , then the line bundle  $\mathcal{O}_{X_k}(1)$  is big as soon as the top self-intersection  $\mathcal{O}_{X_k}(\mathbf{a})^{k+2}$  is positive.*

Proof. Without loss of generality, we can suppose that  $\mathbf{a}$  is integral and that  $\mathbf{a} \in \mathring{\mathfrak{N}}$ . Then Lemma 7.1 ensures that all the “vertical” eigenvalues are positive: in this case, thanks to Lemma 7.2, we conclude that the curvature of  $\mathcal{O}_{X_k}(\mathbf{a})^{k+2}$  can have at most one negative “horizontal” eigenvalue.

Thus,  $X_k(\leq 1, \mathcal{O}_{X_k}(\mathbf{a})) = X_k$  and so

$$\begin{aligned} \int_{X_k(\leq 1, \mathcal{O}_{X_k}(\mathbf{a}))} \left( \frac{i}{2\pi} \Theta(\mathcal{O}_{X_k}(\mathbf{a})) \right)^{k+2} &= \int_{X_k} \left( \frac{i}{2\pi} \Theta(\mathcal{O}_{X_k}(\mathbf{a})) \right)^{k+2} \\ &= \mathcal{O}_{X_k}(\mathbf{a})^{k+2}. \end{aligned}$$

If  $\mathcal{O}_{X_k}(\mathbf{a})^{k+2} > 0$ , then, by Demailly’s holomorphic Morse inequalities,  $\mathcal{O}_{X_k}(\mathbf{a})$  is big and so is  $\mathcal{O}_{X_k}(1)$  (recall that if  $\mathbf{a} \in \mathbb{N}^k$ , then there is a non-trivial morphism  $\mathcal{O}_{X_k}(\mathbf{a}) \rightarrow \mathcal{O}_{X_k}(|\mathbf{a}|)$ ).  $\square$

**7.2. End of the proof.** Let  $u_j = c_1(\mathcal{O}_{X_j}(1))$  be the first Chern class of the anti-tautological line bundle on  $X_j$ . Define the (real) polynomials  $F_k, G_k: \mathbb{R}^k \rightarrow \mathbb{R}$  by

$$(a_1 u_1 + \dots + a_k u_k)^{k+2} = F_k(\mathbf{a})c_1(X)^2 - G_k(\mathbf{a})c_2(X).$$

Observe that these two polynomials do not depend on the particular surface  $X$ , but only on the relative structure of the fibration  $X_k \rightarrow X$ , which is universal.

**Lemma 7.5.** *Suppose that for each  $k \geq 1$ , there exists a minimal surface of general type  $X$  such that  $\mathcal{O}_{X_k}(1)$  is not big. Then, if  $\mathbf{a} \in \mathfrak{N}$ , we have the inequalities*

$$3F_k(\mathbf{a}) \geq G_k(\mathbf{a}) \geq 0$$

and  $G_k \not\equiv 0$ .



Proof. Since  $F_k$  and  $G_k$  are independent of the particular surface chosen, we can suppose  $X$  to be a compact unramified quotient of the ball  $\mathbb{B}_2$ . In this case,  $D \equiv 2/9$  and, for  $\mathbf{a} \in \mathfrak{N}$  rational,

$$(a_1u_1 + \dots + a_ku_k)^{k+2} = \int_{X_k} \left( \frac{i}{2\pi} \Theta(\mathcal{O}_{X_k}(\mathbf{a})) \right)^{k+2} > 0$$

by Lemmas 7.1, 7.2 and 7.3; on the other hand, by Bogomolov-Miyaoka-Yau,  $c_1(X)^2 = 3c_2(X)$  and also  $c_1(X)^2 > 0$ . Hence, by continuity,  $3F_k(\mathbf{a}) - G_k(\mathbf{a}) \geq 0$  on  $\mathfrak{N}$  (with strict inequality for  $\mathbf{a}$  rational in the interior of the cone).

Now, let us compute the intersection  $(a_1u_1 + \dots + a_ku_k)^{k+2}$  on a minimal surface of general type  $X$  as in the hypotheses: of course, such a surface cannot be a compact unramified quotient of the ball  $\mathbb{B}_2$ . In this case we have  $c_1(X)^2 < 3c_2(X)$ , and so

$$\begin{aligned} (a_1u_1 + \dots + a_ku_k)^{k+2} &= F_k(\mathbf{a})c_1(X)^2 - G_k(\mathbf{a})c_2(X) \\ &\geq \frac{1}{3}G_k(\mathbf{a})\underbrace{(c_1(X)^2 - 3c_2(X))}_{<0}. \end{aligned}$$

Thus, if there exists a point  $\mathbf{a}' \in \mathfrak{N}$  such that  $G_k(\mathbf{a}') < 0$ , we would have  $(a_1u_1 + \dots + a_ku_k)^{k+2} > 0$  and hence, by Proposition 7.4,  $\mathcal{O}_{X_k}(1)$  big, contradiction. Finally, if  $G_k \equiv 0$ , fix a rational point  $\mathbf{a}$  in the interior of the cone  $\mathfrak{N}$ : such an  $\mathbf{a}$  gives  $F_k(\mathbf{a}) > 0$ . Then, for such a point we would have  $(a_1u_1 + \dots + a_ku_k)^{k+2} = F_k(\mathbf{a})c_1(X)^2 > 0$ , again contradiction. □

REMARK 7.3. Call  $(\sharp)$  the first hypothesis of Lemma 7.5. If  $(\sharp)$  is not satisfied, then we would have that there exist a  $k \geq 1$  such that for each minimal surface of general type  $X$ , the line bundle  $\mathcal{O}_{X_k}(1)$  is big. In this case, we would already have the global sections we are looking for.

Now, let  $\Sigma_k \subset \mathbb{R}^k$  the zero locus of  $G_k$ . By the above lemma,  $\mathfrak{N} \setminus \Sigma_k$  is dense in  $\mathfrak{N}$ . Set

$$m_k \stackrel{\text{def}}{=} \sup_{\mathbf{a} \in \mathfrak{N} \setminus \Sigma_k} \frac{F_k(\mathbf{a})}{G_k(\mathbf{a})}.$$

If  $(\sharp)$  holds, then  $m_k < +\infty$ : otherwise, for each  $M > 0$  we would find an  $\mathbf{a}_M \in \mathfrak{N} \setminus \Sigma_k$  such that  $F_k(\mathbf{a}_M) > M G_k(\mathbf{a}_M)$  and so

$$\begin{aligned} F_k(\mathbf{a}_M)c_1(X)^2 - G_k(\mathbf{a}_M)c_2(X) &> M G_k(\mathbf{a}_M)c_1(X)^2 - G_k(\mathbf{a}_M)c_2(X) \\ &= G_k(\mathbf{a}_M)(M c_1(X)^2 - c_2(X)). \end{aligned}$$

We would then contradict  $(\sharp)$ , by choosing  $M > c_2(X)/c_1(X)^2$ .

On the other hand, obviously, if  $m_k > c_2(X)/c_1(X)^2$ , then  $\mathcal{O}_{X_k}(1)$  is big. Moreover, for each  $k \geq 1$ , we have  $1/3 \leq m_k \leq m_{k+1}$ . The inequality  $m_k \geq 1/3$  follows directly from Lemma 7.5. To see the monotonicity, notice that  $(a_1u_1 + \dots + a_ku_k)^{k+2}|_{a_k=0} \equiv 0$  (just for dimension reasons) and so

$$(a_1u_1 + \dots + a_ku_k)^{k+2} = a_k \cdot \underbrace{\frac{1}{a_k}(a_1u_1 + \dots + a_ku_k)^{k+2}}_{\text{well defined for } a_k = 0}.$$

But then,

$$\begin{aligned} \frac{1}{a_k}(a_1u_1 + \dots + a_ku_k)^{k+2} \Big|_{a_k=0} &= \frac{\partial}{\partial a_k}(a_1u_1 + \dots + a_ku_k)^{k+2} \Big|_{a_k=0} \\ &= (k+2)(a_1u_1 + \dots + a_{k-1}u_{k-1})^{k+1} \cdot u_k \\ &= (k+2)(a_1u_1 + \dots + a_{k-1}u_{k-1})^{k+1}, \end{aligned}$$

where the last equality is simply obtained by integrating along the fibers of  $X_k \rightarrow X_{k-1}$ . Hence, we have that

$$\frac{F_k(a_1, \dots, a_{k-1}, 0)}{G_k(a_1, \dots, a_{k-1}, 0)} = \frac{F_{k-1}(a_1, \dots, a_{k-1})}{G_{k-1}(a_1, \dots, a_{k-1})}$$

and monotonicity follows.

Finally, if we set  $m_\infty$  to be the  $\sup_{k \geq 1} m_k$ , we find that, for  $X$  a given minimal surface of general type, if  $m_\infty > c_2(X)/c_1(X)^2$ , then there exist a  $k_0 \in \mathbb{N}$  such that  $\mathcal{O}_{X_{k_0}}(1)$  is big.

**REMARK 7.4.** For the moment, we are not able to compute or even to estimate in a satisfactory way the limit term  $m_\infty$ . Of course, a divergent sequence would imply the existence of global invariant jet differentials of some order on every surface of general type. A less ambitious aim could be, for example, to encompass the case of hypersurfaces  $X$  of  $\mathbb{P}^3$  of degree greater than or equal to five (which is the minimum degree for  $X$  to be of general type). In this case, a simple Chern classes computation shows that  $m_\infty > 11$  would be sufficient.

**7.2.1. Comparison with lower bounds of [5].** For low values of  $k$ , one can compute directly the intersection product

$$(a_1u_1 + \dots + a_ku_k)^{k+2}$$

either algebraically, by means of sequences (1) and (2), or using our curvature formula and computing the corresponding integrals (for more details, see [6]), on some partic-

ular  $k$ -tuple  $\mathbf{a}$ . For instance, a natural choice is given by the sequences (2, 1), (6, 2, 1), (18, 6, 2, 1) and so on. These supply the estimates

$$m_1 \geq 1, \quad m_2 \geq \frac{13}{9} \simeq 1.44, \quad m_3 \geq \frac{1195}{742} \simeq 1.61, \quad m_4 \geq \frac{442243}{271697} \simeq 1.63,$$

which give the existence of global invariant jet differentials on a minimal surface of general type whose Chern classes satisfy the following inequalities:

order 1 jet differentials if  $c_1(X)^2 > c_2(X)$ ,

order 2 jet differentials if  $13c_1(X)^2 > 9c_2(X)$ ,

order 3 jet differentials if  $1195c_1(X)^2 > 742c_2(X)$ ,

order 4 jet differentials if  $442243c_1(X)^2 > 271697c_2(X)$ .

Unfortunately, these first terms are still very far from being close even to 11.

We would like to remark here, that, even if we are dealing with the same relatively nef bundles of [5], we get considerably better lower bounds for the degree in the case of hypersurface in  $\mathbb{P}^3$ .

The reason is quite subtle: from a Hermitian point of view, in proving Theorem 2 of [5], we tacitly used the restriction of the Fubini-Study metric of the projective space the hypersurface is embedded in, to the tangent bundle of the hypersurfaces. This is why he had to “correct” this metric by adding some positivity coming from  $\mathcal{O}(2)$  and thus loosing some effectivity.

Using instead the differential-geometric approach of the present paper, we were able to take advantage of the full strength of the Kähler-Einstein metric, which reflects directly the strong positivity properties of varieties with ample canonical bundle.

ACKNOWLEDGMENTS. We would like to thank Stefano Trapani for his suggestions, comments and remarks which have finally lead to the present version of Theorem 1.2.

---

### References

- [1] F.A. Bogomolov: *Families of curves on a surface of general type*, Dokl. Akad. Nauk SSSR **236** (1977), 1041–1044.
- [2] J.-P. Demailly: *Champs magnétiques et inégalités de Morse pour la  $d''$ -cohomologie*, Ann. Inst. Fourier (Grenoble) **35** (1985), 189–229.
- [3] J.-P. Demailly: *Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials*; in Algebraic Geometry—Santa Cruz 1995, Proc. Sympos. Pure Math. **62**, Amer. Math. Soc., Providence, RI., 1997, 285–360.
- [4] J.-P. Demailly: *Private communication*, (2007).

- [5] S. Diverio: *Differential equations on complex projective hypersurfaces of low dimension*, Compos. Math., to appear (2008).
- [6] S. Diverio: *Jet differentials, holomorphic Morse inequalities and hyperbolicity*, PhD thesis in mathematics, Sapienza Università di Roma — Université Joseph Fourier, Grenoble I (2008).
- [7] M. Green and P. Griffiths: *Two applications of algebraic geometry to entire holomorphic mappings*; in The Chern Symposium 1979 (Proc. Internat. Sympos., Berkeley, Calif., 1979), Springer, New York, 1980, 41–74.
- [8] M. McQuillan: *Diophantine approximations and foliations*, Inst. Hautes Études Sci. Publ. Math. **87** (1998), 121–174.

Istituto “Guido Castelnuovo”  
SAPIENZA Università di Roma  
Piazzale Aldo Moro 2  
00185 Roma  
Italy

Current address:  
Chargé de recherche  
Institut de Mathématiques de Jussieu  
Université Pierre et Marie Curie  
175 rue du Chevaleret  
75013 Paris  
France  
e-mail: [diverio@math.jussieu.fr](mailto:diverio@math.jussieu.fr)