



Title	On integral basis of algebraic function fields with several variables
Author(s)	Nobusawa, Nobuo
Citation	Osaka Mathematical Journal. 1959, 11(1), p. 63-69
Version Type	VoR
URL	<a href="https://doi.org/10.18910/5371">https://doi.org/10.18910/5371</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Nobusawa, Nobuo  
Osaka Math. J.  
**11** (1959), 63-69.

***On Integral Basis of Algebraic Function Fields  
with Several Variables***

By Nobuo NOBUSAWA

Let  $K$  be an algebraic function field with two variables and  $w$  a discrete valuation of rank 2 of  $K$ . Let  $L$  be a finite extension of  $K$  and  $w_1, w_2, \dots, w_g$  all the extensions of  $w$  in  $L$ . We denote the valuation rings of  $w$  in  $K$  and of  $w_i$  in  $L$  by  $\mathfrak{o}_0$  and  $\mathfrak{o}_i$ . It is clear that  $\mathfrak{o} = \bigcap_{i=1}^g \mathfrak{o}_i$  is the integral closure of  $\mathfrak{o}_0$  in  $L$ . The structure of  $\mathfrak{o}$  as an  $\mathfrak{o}_0$ -module will be determined in this paper. Let  $(e_1^{(i)}, e_2^{(i)})$  be the value of ramification of  $w_i$ :  $w_i(a) = (e_1^{(i)}, e_2^{(i)})w(a)$  for  $a \in K$ .<sup>1)</sup> The main theorem given in this paper is that  $\mathfrak{o}$  is a direct sum of  $n_0 (= \sum_i e_1^{(i)} f_i)$   $\mathfrak{o}_0$ -modules of rank 1 and of  $n - n_0$   $\mathfrak{o}_0$ -modules of infinite rank where  $n = [L : K]$ . From this we can easily conclude that  $L/K$  has integral basis with respect to  $w$  in the classical sense when and only when  $e_2^{(i)} = 1$  for every  $i$ .<sup>2)</sup> In order to get the theorem, some lemmas on the independence of valuations of rank 2 will be required, which are proved generalizing naturally the well-known proofs in case of rank 1. Then we construct concretely  $n$  linearly independent basis of  $L/K$  which are a generalization of the classical integral basis, having the following property: If  $u_1, u_2, \dots, u_n$  are those generalized integral basis and  $\sum c_i u_i \in \mathfrak{o}$  with  $c_i$  in  $K$ , then  $c_i u_i \in \mathfrak{o}$  for each  $i$ .

The above mentioned results will be inductively generalized in general case. Let  $K$  be an algebraic function field with several variables and  $w$  a discrete valuation of  $K$ . Let  $L$  be a finite extension of  $K$ . We must assume that there holds a fundamental equality with respect to the extensions of  $w$  in  $L$ :  $\sum_i e_i f_i = n$  where  $e_i$  are the ramification indices of  $w_i$  and  $f_i$  are the relative degrees of  $w_i$ . This equality holds when  $\text{rank } w + \dim w = n$ . (See Roquette [4], p. 43. Second Criterion.) In this case the main theorem is proved to be true, although we do not discuss the general case in this paper.

1) We express  $w_i(a)$  and  $w(a)$  in the normal exponential form and the order of value group will be determined by the last non-zero difference of components (contrary to the usual sense. [5]).

2) For the definition of integral basis, see [1].

### 1. Lemmas on valuations.<sup>3)</sup>

Let us denote a discrete valuation  $w$  of rank 2 of a field  $K$  in the normal exponential form :

$$w(a) = (\alpha_1, \alpha_2) \quad \text{for non zero elements } a \text{ of } K,$$

where  $\alpha_i$  range over all rational integers. The order of the value group is defined such as  $(\alpha_1, \alpha_2) \succ (\beta_1, \beta_2)$  if  $\alpha_2 \succ \beta_2$  or if  $\alpha_2 = \beta_2$  and  $\alpha_1 \succ \beta_1$ . From this valuation  $w$ , we get a valuation  $w^{(2)}$  of rank 1 putting  $w^{(2)}(a) = \alpha_2$ .

Let  $v_1, v_2, \dots, v_n$  be a set of discrete valuations of rank 1 and of rank 2 satisfying the next condition :

$$(A) \quad \begin{cases} \text{i) if both } v_i \text{ and } v_j \text{ are of rank 1, then } v_i \neq v_j, \\ \text{ii) if } v_i \text{ is of rank 1 and } v_j \text{ of rank 2, then } v_i \neq v_j^{(2)}, \\ \text{iii) if both } v_i \text{ and } v_j \text{ are of rank 2, then } v_i \neq v_j \text{ and } v_i^{(2)} = v_j^{(2)}. \end{cases}$$

**Lemma 1.** *Let  $v_1, v_2, \dots, v_n$  be a set of discrete valuations of rank 1 and of rank 2 satisfying (A) and let  $\mathfrak{o}_i$  be the valuation rings of  $v_i$ . Then  $\mathfrak{o}_i \not\subseteq \mathfrak{o}_j$  for  $i \neq j$ .*

Proof. When one of  $v_i$  and  $v_j$  is of rank 1, Lemma 1 is clear from the theory of valuations of rank 1 by (A) i) and ii). For, there exists an element  $a$  such that  $v_i^{(1)}(a) = \alpha$  and  $v_j^{(1)}(a) = \beta$  for any rational integers  $\alpha$  and  $\beta$ . Here we denote, by  $v_i^{(1)}, v_i^{(2)}$  when  $v_i$  is of rank 2 and  $v_i$  when  $v_i$  is of rank 1. Next suppose both  $v_i$  and  $v_j$  are of rank 2. Put  $\mathfrak{o}'_i = \{a \in K | v_i(a) = (\alpha, 0), \alpha \geq 0\}$  and  $\mathfrak{p}'_i = \{b \in K | v_i(b) = (\beta, 0), \beta > 0\}$ . If  $\mathfrak{o}'_i \subseteq \mathfrak{o}'_j$ , we have  $\mathfrak{o}'_i \subseteq \mathfrak{o}'_j$  by (A) iii), and  $\mathfrak{p}'_i = \mathfrak{o}'_i \cap \mathfrak{p}'_j$ , since  $\mathfrak{o}'_i \cap \mathfrak{p}'_j$  is a proper prime ideal of the semi-group  $\mathfrak{o}'_i$ <sup>4)</sup> and  $\mathfrak{p}'_i$  is the unique proper prime ideal of  $\mathfrak{o}'_i$ . If  $c$  is an element such that  $v_i(c) = (\gamma, 0)$  with  $\gamma < 0$ , then  $c^{-1} \in \mathfrak{p}'_i$  and hence  $c^{-1} \in \mathfrak{p}'_j$ . Therefore  $v_j(c) = (\delta, 0)$  with  $\delta < 0$ , which implies  $\mathfrak{o}'_i \supseteq \mathfrak{o}'_j$ , and hence  $\mathfrak{o}'_i = \mathfrak{o}'_j$  by (A) iii), that is,  $v_i = v_j$ .

**Lemma 2.** *Let  $v_1, v_2, \dots, v_n$  be a set of discrete valuations satisfying (A). Then there exists an element  $x$  such that*

$$v_1(1-x) > 0, \quad v_2(x) > 0, \dots, v_n(x) > 0.$$

Proof. We shall prove Lemma 2 by induction. First assume  $n=2$ . By Lemma 1 there exists an element  $a$  such that  $v_1(a) \geq 0$  and  $v_2(a) < 0$ .

3) The proofs in this paper are, as stated in the introduction, generalizations of the classical ones and we shall follow [3] for this purpose.

4) If we put  $\mathfrak{g}_i = \{a \in K | v_i^{(2)}(a) = 0\}$ , then  $\mathfrak{g}_i = \mathfrak{g}_j$  by (A) iii) and  $\mathfrak{g}_i = \{\mathfrak{o}'_i, \mathfrak{o}'_i^{-1}\}$ . Then  $\mathfrak{o}'_i \cap \mathfrak{p}'_j = \emptyset$  implies that  $\{\mathfrak{o}'_i, \mathfrak{o}'_i^{-1}\} \cap \mathfrak{p}'_j = \emptyset$  which is a contradiction.

Put  $x_1=1/a$  if  $v_1(a)=0$  and  $x_1=1/(1+a)$  if  $v_1(a)>0$ . Then  $v_1(x_1)=0$  and  $v_2(x_1)>0$ . Similarly there exists  $x_2$  such that  $v_1(x_2)>0$  and  $v_2(x_2)=0$ .  $x=x_1/(x_1+x_2)$  is then a required element. Assume that Lemma 2 is true for  $n-1$ , and there exist  $x_1$  and  $x_2$  such that

$$v_1(1-x_1) > 0, \quad v_3(x_1) > 0, \quad v_4(x_1) > 0, \dots, v_n(x_1) > 0,$$

and

$$v_1(1-x_2) > 0, \quad v_2(x_2) > 0, \quad v_4(x_2) > 0, \dots, v_n(x_2) > 0.$$

Then it is easy to show the following element  $x$  is a required one :

- 1)  $x = x_1 x_2$  if  $v_2(x_1) \geq 0$  and  $v_3(x_2) \geq 0$ ,
- 2)  $x = x_1/(1+x_1(1-x_1))$  if  $v_2(x_1) < 0$ ,
- 3)  $x = x_2/(1+x_2(1-x_2))$  if  $v_2(x_1) \geq 0$  and  $v_3(x_2) < 0$ .

**Lemma 3.** *Let  $w_1, w_2, \dots, w_g$  be  $g$  distinct discrete valuations of rank 2 of  $K$ . If  $w_1(a)=0$  with some element  $a$ , then there exists an element  $a'$  such that  $w_1(a-a')>0, w_2(a')>0, \dots, w_g(a')>0$ .*

Proof. We may assume that  $w_1^{(2)}=w_2^{(2)}=\dots=w_i^{(2)}$  and  $w_1^{(2)} \neq w_{i+1}^{(2)}$  for  $j \geq 1$ . All the distinct valuations in  $w_1, \dots, w_i, w_{i+1}^{(2)}, \dots, w_g^{(2)}$  will be denoted  $v_1 (= w_1), v_2, \dots, v_n$ . Then  $v_1, v_2, \dots, v_n$  satisfy (A) and hence there exists an element  $x$  by Lemma 2 such that  $v_1(1-x)>0, v_2(x)>0, \dots, v_n(x)>0$ . Put  $a'=ax^m$  with a sufficiently large number  $m$ . Then we have

$$v_1(a-a') = v_1(a) + v_1(1-x^m) \geq v_1(1-x) > 0$$

and

$$v_k(a') = v_k(a) + mv_k(x) \quad \text{for } k \neq 1.$$

When  $v_k$  is of rank 1,  $v_k(a) + mv_k(x) > 0$  with a sufficiently large number  $m$ . When  $v_k$  is of rank 2,  $v_k^{(2)}(a) = v_1^{(2)}(a) = 0$ , and hence we have also  $v_k(a') > 0$  with a sufficiently large number  $m$ .

**Lemma 4.** *When  $v_1, v_2, \dots, v_n$  are a set of discrete valuations satisfying (A), there exist  $x_i$  for every  $i$  such that*

$$v_i(x_i) = \begin{cases} 1 & \text{if } v_i \text{ is of rank 1,} \\ (1, 0) & \text{if } v_i \text{ is of rank 2,} \end{cases}$$

$$v_j(x_i) = 0 \quad \text{for } j \neq i.$$

Proof. We may prove the existence of  $x_1$ . If  $v_1$  is of rank 1, we choose an element  $a$  such that  $v_1(a)=1$  and  $v_i^{(2)}(a)=0$  for  $i \neq 1$ . This can be done by the theory of valuations of rank 1. If  $v_1$  is of rank 2, we choose an element  $a$  with  $v_1(a)=(1, 0)$ , when it is seen  $v_i^{(2)}(a)=0$  with  $v_i$  of rank 2. Let  $x$  be an element in Lemma 2 and put

$$x_1 = ax^m + (x-1)^m$$

with a sufficiently large natural number  $m$ . Then

$$v_i(x_1) = v_i(ax^m + (x-1)^m) = v_i(ax^m) = v_i(a) + mv_i(x) = v_i(a),$$

and

$$v_i(x_1) = v_i(ax^m + (x-1)^m) = v_i((x-1)^m) = 0,$$

since  $v_i(ax^m) = v_i(a) + mv_i(x) > 0$  with sufficiently large  $m$ .

**Lemma 5.** *Let  $v_1, v_2, \dots, v_n$  be a set of discrete valuations satisfying (A) and let  $\alpha_i$  be any rational integers. Then there exists an element  $x$  such that*

$$v_i(x) = \begin{cases} \alpha_i & \text{if } v_i \text{ is of rank 1,} \\ (\alpha_i, 0) & \text{if } v_i \text{ is of rank 2.} \end{cases}$$

Proof. We may put  $x = \prod x_i^{\alpha_i}$  with  $x_i$  in Lemma 4.

**Lemma 6.** *Let  $w_1, w_2, \dots, w_g$  be  $g$  distinct discrete valuations of rank 2. When  $l, m$  and  $l'$  are any rational integers, there exist such elements  $x_{i,m}^{(i)}$  for every  $i$  that  $w_i(x_{i,m}^{(i)}) = (l, m)$  and  $w_j(x_{i,m}^{(i)}) \geq (l', m)$  for  $j \neq i$ .*

Proof. We may prove Lemma 6 for  $i=1$ . We define  $v_1, v_2, \dots, v_n$  from  $w_1, w_2, \dots, w_g$  as in Lemma 3. Assume that  $v_1, v_2, \dots, v_i$  are of rank 2 and that  $v_{i+1}, \dots, v_n$  are of rank 1. Then there exists an element  $a$  such that  $v_1^{(2)}(a) = \dots = v_i^{(2)}(a) = m$  and  $v_{i+j}^{(2)}(a) > m$  for  $j \geq 1$ . When  $v_k(a) = (\alpha_k, m)$  for  $1 \leq k \leq i$ , put  $\beta_1 = l - \alpha_1$  and  $\beta_k > l' - \alpha_k$  for  $2 \leq k \leq i$ . By Lemma 5 we can choose an element  $b$  such that  $v_k(b) = (\beta_k, 0)$  for  $1 \leq k \leq i$  and  $v_{i+j}(b) \geq 0$  for  $j \geq 1$ .  $x_{i,m}^{(1)} = ab$  is a required element.

## 2. The structure of $\mathfrak{o}$ .

Let  $K$  be an algebraic function field with two variables and  $w$  a discrete valuation of rank 2 of  $K$ . We denote the valuation ring and the valuation ideal of  $w$  by  $\mathfrak{o}_0$  and  $\mathfrak{p}_0$ . Let  $L$  be a finite extension of  $K$  and  $w_1, w_2, \dots, w_g$  all the extensions of  $w$  in  $L$ . We denote the valuation rings and the valuation ideals of  $w_i$  by  $\mathfrak{o}_i$  and  $\mathfrak{p}_i$ . Let  $(e_1^{(i)}, e_2^{(i)})$  be defined such as  $w_i(a) = (e_1^{(i)}, e_2^{(i)})w(a)$  for  $a \in K$ .  $e_i = e_1^{(i)}e_2^{(i)}$  is the ramification index.  $f_i = [\mathfrak{o}_i/\mathfrak{p}_i : \mathfrak{o}_0/\mathfrak{p}_0]$  is the relative degree of  $w_i$ . Then  $\sum_{i=1}^g e_i f_i = [L : K]$  by Roquette [4]. Let  $\bar{t}_{i,j}$  ( $j = 1, 2, \dots, f_i$ ) be  $\mathfrak{o}_0/\mathfrak{p}_0$ -basis of  $\mathfrak{o}_i/\mathfrak{p}_i$  and let  $t_{i,j}$  be representatives of  $\bar{t}_{i,j}$  chosen by Lemma 3 such as

$$(1) \quad w_k(t_{i,j}) \geq 0 \quad \text{for every } k.$$

By lemma 6 we can choose  $x_{im}^{(i)}$  such that

$$(2) \quad w_i(x_{im}^{(i)}) = (l, m) \text{ and } w_j(x_{im}^{(i)}) \geq (e_1^{(j)}, m) \text{ for } j \neq i,$$

where  $l=0, 1, \dots, e_1^{(i)}-1$  and  $m=0, 1, \dots, e_2^{(i)}-1$ .

**Lemma 7.** *If  $a = \sum_{i,j,l,m} c_{ijlm} t_{ij} x_{im}^{(i)}$  with  $c_{ijlm} \in K$  such that  $w_i^{(2)}(a) \geq e_2^{(i)}$  for every  $i$  and  $w^{(2)}(c_{ijlm}) \geq 0$ , then  $w^{(2)}(c_{ijlm}) > 0$ .*

Proof. Assume that  $w^{(2)}(c_{ijlm}) = 0$  for some  $c_{ijlm}$  and put

$$(3) \quad b = \sum'_{i,j,l,m} c_{ijlm} t_{ij} x_{im}^{(i)}$$

where  $\sum'$  denotes the sum of all  $c_{ijlm}$  such that  $w^{(2)}(c_{ijlm}) = 0$ . It is clear  $w_i^{(2)}(b) \geq e_2^{(i)}$  for every  $i$ . Let  $m_0$  be the smallest of  $m$  with  $c_{ijlm} \neq 0$  in (3), and let  $c = c_{i_0 j_0 l_0 m_0}$  be one of  $c_{ijlm_0}$  having the smallest value with respect to  $w$ . Then

$$bc^{-1} = \sum_{\substack{m \geq m_0 \\ i, j, l}} b_{ijlm} t_{ij} x_{im}^{(i)} \text{ with } b_{ijlm} = c_{ijlm} c^{-1},$$

where  $w_i^{(2)}(bc^{-1}) \geq e_2^{(i)}$ ,  $b_{ijlm_0} \in \mathfrak{p}_0$  and  $b_{i_0 j_0 l_0 m_0} = 1$ . We shall show that all  $b_{i_0 j_0 l_0 m_0} \in \mathfrak{p}_0$ . We have, by (1) and (2),

$$w_{i_0}^{(2)}(b_{ijlm} t_{ij} x_{im}^{(i)}) > m_0 \quad \text{for } m > m_0,$$

and

$$w_{i_0}(b_{ijlm_0} t_{ij} x_{im_0}^{(i)}) \geq (e_1^{(i_0)}, m_0) \quad \text{for } i \neq i_0.$$

Hence

$$(4) \quad w_{i_0}(\sum_{j,l} b_{ijlm_0} t_{ij} x_{im_0}^{(i)}) \geq (e_1^{(i_0)}, m_0).$$

We have also

$$w_{i_0}(b_{i_0 j_0 l_0 m_0} t_{i_0 j_0} x_{i_0 m_0}^{(i_0)}) > (0, m_0) \quad \text{for } l > 0,$$

and hence by (4)

$$w_{i_0}(\sum_j b_{i_0 j_0 m_0} t_{i_0 j_0} x_{i_0 m_0}^{(i_0)}) > (0, m_0),$$

from which we get

$$w_{i_0}(\sum_j b_{i_0 j_0 m_0} t_{i_0 j_0}) > 0,$$

that is,  $b_{i_0 j_0 m_0} \in \mathfrak{p}_0$  for all  $j$ . Therefore

$$(5) \quad w_{i_0}(b_{i_0 j_0 m_0}) \geq (e_1^{(i_0)}, 0).$$

Next we have, by (1), (2) and (5),

$$w_{i_0}(b_{i_0 j_0 l_0 m_0} t_{i_0 j_0} x_{i_0 m_0}^{(i_0)}) > (1, m_0) \quad \text{for } l \neq 1,$$

and hence, by (4),

$$w_{i_0} \left( \sum_j b_{i_0 j 1 m} t_{i_0 j} x_{1 m_0}^{(i_0)} \right) > (1, m_0),$$

that is,

$$w_{i_0} \left( \sum_j b_{i_0 j 1 m_0} t_{i_0 j} \right) > 0.$$

Thus  $b_{i_0 j 1 m_0} \in \mathfrak{p}_0$  for all  $j$ . Continuing the same procedure, we get that  $b_{i_0 j l m_0} \in \mathfrak{p}_0$  for all  $j$  and  $l$ . But this contradicts  $b_{i_0 j_0 l_0 m_0} = 1 \notin \mathfrak{p}_0$ .

**Lemma 8.** *If  $a = \sum_{i, j, l} c_{i j l 0} t_{i j} x_{i 0}^{(i)}$  with  $c_{i j l 0} \in K$  is an element of  $\mathfrak{o}$ , then all  $c_{i j l 0} \in \mathfrak{o}_0$ .*

Proof. Assume that  $c_{i j l 0} \notin \mathfrak{o}_0$  for some  $c_{i j l 0}$ . Let  $c = c_{i_0 j_0 l_0 0}$  be one of  $c_{i j l 0}$  having the smallest value with respect to  $w$ . Then by assumption  $c \notin \mathfrak{o}_0$  and hence  $c^{-1} \in \mathfrak{p}_0$ . Then

$$ac^{-1} = \sum_{i, j, l} b_{i j l 0} t_{i j} x_{i 0}^{(i)}$$

with  $b_{i j l 0} \in \mathfrak{o}_0$  and  $b_{i_0 j_0 l_0 0} = 1$ . We have

$$w_{i_0} (ac^{-1}) \geq w_{i_0} (c^{-1}) \geq (e_1^{(i_0)}, 0).$$

As in the proof of Lemma 7, we can show  $b_{i_0 j l 0} \in \mathfrak{p}_0$  for all  $j$  and  $l$ , which is a contradiction.

**Theorem 1.** *If  $a = \sum_{i, j, l, m} c_{i j l m} t_{i j} x_{i m}^{(i)} \in \mathfrak{o}$  with  $c_{i j l m} \in K$ , then  $c_{i j l m} t_{i j} x_{i m}^{(i)} \in \mathfrak{o}$ .*

Proof. First we shall prove  $w^{(2)}(c_{i j l m}) \geq 0$ . Assume on the contrary that  $w^{(2)}(c_{i j l m}) < 0$  for some  $c_{i j l m}$ . Let  $c_{i_0 j_0 l_0 m_0} = c$  be one of  $c_{i j l m}$  having the smallest value with respect to  $w$ . By assumption  $c^{-1} \in \mathfrak{p}_0$ . Then

$$ac^{-1} = \sum_{i, j, l, m} b_{i j l m} t_{i j} x_{i m}^{(i)}$$

with  $b_{i j l m} = c_{i j l m} c^{-1} \in \mathfrak{o}_0$  and  $b_{i_0 j_0 l_0 m_0} = 1$ , and also  $w_i^{(2)}(ac^{-1}) \geq w_i^{(2)}(c^{-1}) \geq e_2^{(i)}$ . Then, by Lemma 7, we have  $w^{(2)}(b_{i j l m}) > 0$  for all  $i, j, l$  and  $m$ , which contradicts  $b_{i_0 j_0 l_0 m_0} = 1$ . Thus we have  $c_{i j l m} t_{i j} x_{i m}^{(i)} \in \mathfrak{o}$  for  $m > 0$  and for all  $i, j, l$ . We may now assume that  $a = \sum_{i, j, l} c_{i j l 0} t_{i j} x_{i 0}^{(i)}$ . In this case, by Lemma 8, we can say that  $c_{i j l 0} \in \mathfrak{o}_0$  and hence  $c_{i j l 0} t_{i j} x_{i 0}^{(i)} \in \mathfrak{o}$ , which completes the proof.

**Corollary 1.**  *$t_{i j} x_{i m}^{(i)}$  are linearly independent over  $K$ .*

Proof. If  $0 = \sum c_{i j l m} t_{i j} x_{i m}^{(i)}$ , then  $c_{i j l m} t_{i j} x_{i m}^{(i)} \in \mathfrak{o}$  by Theorem 1. Since we may suppose that  $c_{i j l m}$  take any small values with respect to  $w$ ,  $c_{i j l m} t_{i j} x_{i m}^{(i)} \in \mathfrak{o}$  imply  $c_{i j l m} = 0$ .

**Theorem 2.** *If we put  $\mathfrak{o}_{ijlm} = \mathfrak{o} \cap \{ct_{ij}x_{lm}^{(i)} \text{ with } c \in K\}$ , then  $\mathfrak{o} = \sum_{i,j,l,m} \mathfrak{o}_{ijlm}$  (direct).*

Proof. This is a direct consequence of Theorem 1 and Corollary 1.

**Corollary 1.** *If  $e_2^{(i)} = 1$  for every  $i$ , then  $\mathfrak{o}$  is a finite  $\mathfrak{o}_0$ -module and has a linearly independent basis with respect to  $\mathfrak{o}_0$ .*

Proof. Note  $\mathfrak{o}_{ijl0} = \mathfrak{o}_0 t_{ij} x_{lm}^{(i)}$ .

**Corollary 2.** *If  $e_2^{(i)} \neq 1$  for some  $i$ , then  $\mathfrak{o}$  is not a finite  $\mathfrak{o}_0$ -module.*

Proof. It is sufficient to show that  $\mathfrak{o}_{ijlm}$  is not a finite  $\mathfrak{o}_0$ -module if  $m > 0$ . Assume on the contrary that  $\mathfrak{o}_{ijlm}$  is a finite  $\mathfrak{o}_0$ -module. Then there exists a minimal value of  $w(c)$  where  $ct_{ij}x_{lm}^{(i)} \in \mathfrak{o}_{ijlm}$ . For, if  $c_1 t_{ij} x_{lm}^{(i)}, c_2 t_{ij} x_{lm}^{(i)}, \dots, c_p t_{ij} x_{lm}^{(i)}$  constitute  $\mathfrak{o}_0$ -basis of  $\mathfrak{o}_{ijlm}$ , take  $w(c) = \min_{1 \leq s \leq p} (w(c_s))$ . If it is  $w(c_0)$ , then  $w^{(2)}(c_0) \geq 0$ , since

$$w_i(c_0 t_{ij} x_{lm}^{(i)}) = w_i(c_0) + (l, m) \geq 0.$$

Therefore  $w_k(c_0 t_{ij} x_{lm}^{(i)}) \geq (\alpha, 1)$  for every  $k$  with some integer  $\alpha$ , since  $m \geq 1$ . Let  $c'$  be an element of  $K$  such that  $w(c') = (\alpha', 0)$  with  $\alpha' < 0$ . Then

$$w_k(c' c_0 t_{ij} x_{lm}^{(i)}) \geq (\alpha + \beta_k, 1) > 0 \quad \text{for all } k,$$

where  $\beta_k = e_1^{(k)} \alpha'$ , that is  $c' c_0 t_{ij} x_{lm}^{(i)} \in \mathfrak{o}_{ijlm}$  and  $w(c' c_0) < w(c_0)$ , which contradicts the minimality of  $c_0$ .

(Received March 17, 1959)

---

### References

- [1] C. Chevalley: Introduction to the theory of algebraic functions of one variable, New York (1951).
- [2] I. S. Cohen and O. Zariski: A fundamental inequality in the theory of extensions of valuations, Illinois J. of Math. **1**, (1957) 1-8.
- [3] K. Iwasawa: Theory of algebraic function fields, (in Japanese) Tokyo (1952).
- [4] P. Roquette: On the prolongation of valuations, Trans. Amer. M. S. **88** (1958) 42-56.
- [5] O. F. G. Schilling: The theory of valuations, New York (1950).

