

Title	On the hitting properties of a class of one- dimensional Markov processes
Author(s)	Manabe, Shojiro
Citation	Osaka Journal of Mathematics. 1975, 12(2), p. 217–266
Version Type	VoR
URL	https://doi.org/10.18910/5378
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Manabe, S. Osaka J. Math. 12 (1975), 217-266

ON THE HITTING PROPERTIES OF A CLASS OF ONE-DIMENSIONAL MARKOV PROCESSES

Shôjirô MANABE

(Received April 9, 1974)

1. Introduction. Let $X = (X_t, P_x, x \in R^1)$ a one-dimensional standard Markov process with generator A

(1.1)
$$Au(x) = a(x)u'(x) + \sigma(x)^{2}u''(x)/2 + \int_{-\infty}^{\infty} \left\{ u(x+y) - u(x) - \frac{y}{-+y^{2}}u'(x) \right\} n(x, dy)$$

In this article, we shall discuss how the sample paths of X approach a single point. Let σ_0 be the first hitting time of the sample path to the origin: $\sigma_0 = \inf\{t>0; X_t=0\}$. Set $\Omega_1 = \{\omega; \sigma_0(\omega) < +\infty\}$. Define $\Omega(\Omega, \Omega\Gamma)$ and Ω_1^{\pm} by

and

$$\Omega_1^{\pm} = \{ \omega \in \Omega_1; \ \mathcal{I}_t \wedge \sigma_0(\omega) \quad s.t. \ X_{t_{2n}-1}(\omega) < 0 < X_{t_{2n}}(\omega) \} .$$

Our present problem is to decide whether $P_x(\Omega_1^+/\Omega_1)$, $P_x(\Omega_1^-/\Omega_1)$ and $P_x(\Omega_1^\pm/\Omega_1)$ are positive or not. When X is spatially homogeneous, the problem was treated by T.Takada [15] in case $\sigma \pm 0$ and by N.Ikeda-S.Watanabe [3] in case $\sigma = 0$ who also applied their results to the study of two-dimensional diffusion processes. Their method is based on the estimate of the singularity of the Green function on the diagonal set. Recently P.W.Millar [10] solved a similar problem independently. Let T_x be the first exit time of the sample path of a spatially homogeneous process from $(-\infty, x]$ (x>0). Millar gave, in terms of the exponent, a necessary and sufficient condition that $P_0(X_{T_x}=x)>0$.

Here we shall consider the class of spatially inhomogeneous Markov processes determined by A under certain regularity conditions on a, σ and Lévy measure n(x, dy). We shall give some sufficient conditions that $P_x(\Omega_1^+/\Omega_1)=1$, $P_x(\Omega_1^-/\Omega_1)=1$ and $P_x(\Omega_1^\pm/\Omega_1)=1$. Our method consists in estimating the singularity of the Green function as in [3]. More precisely, under the regularity conditions that will be given in §2, put S. MANABE

(1.2)
$$\psi(x, \xi) = ia(x)\xi - \sigma(x)^2\xi^2/2 + \int_{-\infty}^{\infty} \left(e^{i\xi} - 1 - \frac{i\xi w}{1 + w^2}\right)n(x, dw).$$

Then *A* can be written in the form:

(1.3)
$$Af(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \ell^{\xi} \psi(x,\xi) \hat{f}(\xi) d\xi,$$

namely, we can regard A as a pseudo-differential operator having its symbol $\psi(x,\xi)$. On account of the theory of pseudo-differential operators, the equation $(\lambda - A)u = fadmits$ a fundamental solution $g_{\lambda}(x, y)$, which is of the form

(1.4)
$$g_{\lambda}(x, y) = \frac{1}{2} \int_{-\infty}^{\infty} e^{i(x-y)\xi} \frac{d\xi}{dx} + g_{\lambda,1}(x, y),$$

with a continuously differentiable function $g_{\lambda,1}(x, y)$. Therefore the singularity of the Green function is the same as in the spatially homogeneous case and the results of [3], [15] concerning the manners of hitting remain true in the present case.

The organization of the present paper is as follows. In section 2, we state our theorems. In section 3, we mention some related facts from the theory of pseudo-differential operators. We construct the above mentioned fundamental solution in section 4 and estimate its singularity in section 5. In section 6, 7 and 8 we prove our theorems by making use of the estimates established in section 5.

2. Theorems. Denote by $\mathcal{B}(\mathbb{R}^n)$ ($\mathcal{B}(\mathbb{R}^n)$) the space of $C^{\infty}(\mathbb{R}^n)$ -functions whose derivatives of any order are bounded (vanishing at infinity). Let a(x) and $\sigma(x)$ be the bounded C^{∞} -functions with its first derivatives a'(x) and $\sigma'(x)$ belonging to $\mathcal{B}^{1,1}$. Let $\nu(x, y)$ be a nonnegative function of $\mathcal{B}(\mathbb{R}^2)$. We assume that $\nu(x, y)$ satisfies following two conditions:

 $(\nu.1)$ there exists a positive constant c_1 such that

$$\nu(x, y) > c_1 \text{ on } R^1 \times \{y; |y| \leq 1\},$$

 $(\nu.2)$ there exists a positive constant L such that

 $\nu(x,y)$ is independent of x, if $|x| \ge L$.

Define n(x, y) by

$$n(x,y) = \begin{cases} \frac{\nu(x, y)}{y^{1+\alpha_1}}, y > 0, \\ \frac{\nu(x, y)}{|y|^{1+\alpha_2}}, y < 0, \end{cases} \quad \text{where } 0 < \alpha_i < 2, i = 1, 2.$$

1) We denote $\mathcal{B}(\text{or } \dot{\mathcal{B}})$ for $\mathcal{B}(\mathbb{R}^1)$ (or $\dot{\mathcal{B}}(\mathbb{R}^1)$) for simplicity.

Consider the following operator A on $\dot{\mathcal{B}}$:

(2.1)
$$Au(x) = a(x)u'(x) + \sigma(x)^2 u''(x)/2 + \int_{-\infty}^{\infty} \left\{ u(x+y) - u(x) - \frac{y}{1+y^2} u'(x) \right\} n(x, y) dy.$$

It is known that there exists a unique standard Markov process $X = \{X_t, P_x, x \in \mathbb{R}^t\}$ whose generator is the closure of A (in fact, X is a Hunt process, see Sato [11]). Let G_{λ} be the resolvent of X:

$$G_{\lambda}f(x) = E_{x} \left[\int_{-\infty}^{\infty} e^{-\lambda t} f(X_{t}) dt \right]^{-1} \int_{-\infty}^{\infty} G_{\lambda}(x, dy) f(y), \quad f \in C_{b}(\mathbb{R}^{1}).^{1}$$

Then it can be shown that there exists a density $g_{\lambda}(x, y)$ of $G_{\lambda}(x, dy)$ with respect to Lebesgue measure dy (see §4).

Now our results are as follows.

Theorem 1. If (i) $\sigma(x)^2 \ge \sigma^2 > 0$ or (ii) $\sigma(x) = 0$ and $\max(\alpha_1, \alpha_2) > 1$, then $P_x(\Omega_1) > 0$ for any $x \neq 0$.

- **Theorem 2.** (1) $//\sigma(x) \ge \sigma^2 > 0$, then we have
 - (i) $P_x(\Omega_1^+ \cup \Omega_1^- | \Omega_1) = 1$ for any $x \neq 0$.
 - (ii) More precisely, for any $x \neq 0$,

(2.2a)
$$E_{x}[e^{-\lambda\sigma_{0}}; \Omega_{1}^{+}] = \frac{\sigma(0)^{2}}{2g_{\lambda}(0, 0)} \left[g_{\lambda}(0, 0)\frac{\partial g_{\lambda}}{\partial y}(x, 0) - g_{\lambda}(x, 0)\frac{\partial g_{\lambda}}{\partial y}(0, 0+)\right],$$

(2.2b)
$$E_x[e^{-\lambda\sigma_0}; \Omega\Gamma] = \int [-g_{\lambda}(0,0)\frac{\partial g_{\lambda}}{\partial v}(x,0) + g_{\lambda}(x,0) (0,0-)].$$

- (2) If $\sigma(x) = 0$ and $\alpha_1 > \alpha_2, \alpha_1 > 1$ [resp. $\alpha_1 < \alpha_2, \alpha_2 > 1$], then $P_x(\Omega_1^+ | \Omega_1) = 1$ [resp. $P_x(\Omega_1^- | \Omega_1) = 1$] for any $x \neq 0$.
- (3) $//\sigma(x)=0$ and $\alpha_1=\alpha_2>1$, then $P_x(\Omega_1^{\pm}|\Omega_1)=1$ for any $x \neq 0$.

3. Pseudo-differential operators. In this section, we shall collect a number of known facts needed for later section from the theory of pseudo-differential operators. We refer to Kumano-go [6] for details. We denote by $S_{\rho,\delta}^m, 0 \le \delta , the set of <math>C^{\infty}(R \times R)$ -functions $p(x,\xi)$ such that

$$(3.1) |D_x^{\alpha}\partial_{\xi}^{\beta}p(x\xi)| \leq C_{\alpha,\beta}(1+|\xi|)^{m+\delta\alpha-\rho\beta} \text{for any } x, \xi \in \mathbb{R}^1,$$

where $D_x^{\alpha} = (-i\partial/\partial x)^{\alpha}$ and $\partial_{\xi}^{\beta} = (\partial/\partial \xi)^{\beta}$. An element of $S_{\rho,\delta}^{m}$ is called a symbol. Set $S_{\rho,\delta}^{\infty} = \bigcup_{m} S_{\rho,\delta}^{m}$ and $S_{\rho,\delta}^{-\infty} = \bigcap_{m} S_{\rho,\delta}^{m}$. For a function $p(x,\xi)$ belonging to $S_{\gamma,\delta}^{m}$, we define a pseudo-differential operator $P = p(x, D_x)$

¹⁾ $C_b(R^1)$ is the totality of bounded continuous functions on R^1 .

by

$$Pu(x) = p(x, D_x)u(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ix\xi} p(x, \xi)\hat{u}(\xi)d\xi^{1}, \quad u \in \mathcal{S} .$$

Let $S_{\rho,\delta}^{m}$ be the totality of $p(x, D_x)$, where $p(x, \xi)$ is a function of $S_{\rho,\delta}^{m}$. Set $S_{\rho,\delta}^{\infty} = \bigcup S_{\rho,\delta}^{m}$ and $S_{\rho,\delta}^{-\infty} = \bigcup S_{\rho,\delta}^{m}$. We call E belonging to $S_{\rho,\delta}^{\infty}$ a right (or left) parametrix of P, if $PE = I - K_1$ for some $K_1 \in S_{\rho,\delta}^{-\infty}$ (or $EP = I - K_2$ for some $K_2 \in S_{\rho,\delta}^{-\infty}$) A parametrix is a right and left parametrix. Following theorem gives a sufficient condition for the existence of parametrix.

Theorem 3.1. (Hürmander [2]). Let $p(x, \xi)$ be a symbol belongs to $S^{m}_{\rho,\delta}$. If $p(x,\xi)$ satisfies the following conditions (i), (ii): (i) For some $\delta_0 > 0$ and m_1 ,

$$|p(x,\xi)| \ge \delta_0 (1+|\xi|)^{m_1}$$
 for any $x, \xi \in \mathbb{R}^1$.

(ii) For any nonnegative integers α , β , there exists a constant $C_{\alpha\beta} > 0$ such that

$$I D^{\beta}_{\mathfrak{a}} \partial^{\mathfrak{a}}_{\xi} p(x\xi) / p(x\xi) I \leq C_{\mathfrak{a}\beta} (1+I\xi I)^{-\rho \mathfrak{a}+\delta\beta} \quad \text{for any } x, \xi \in \mathbb{R}^{1}$$

Then there exists a parametrix $E = e(x, D_x)$, where $e(x, \xi) \in S_{\rho,\delta}^{-m_1}$.

REMARK 3.1. $e(x, \xi)$ can be constructed as follows (Hürmander [2]). Let $e_{j} = 0$, 1, 2, \cdots be functions determined by the following relations

(3.2)
$$e_{0}(x,\xi) = \frac{1}{p(x,\xi)},$$
$$e_{j}(x,\xi) = \frac{-1}{p(x,\xi)} \sum_{\substack{|\omega|+k=j \\ k < j}} \frac{1}{\alpha_{j}} \partial_{\xi}^{\omega} e_{k}(x,\xi) D_{x}^{\omega} p(x,\xi) \quad \text{for any } j \ge 1.$$

Let ϕ be a C^{∞} -function such that $\phi(\xi)=0$ if $|\xi| \leq 1/2$, =1 if $|\xi| \geq 1$. Choose a sequence $1 < t_1 < t_2 < \cdot < t_n \uparrow \infty$ such that

$$(3.3) \qquad |\partial_{\alpha}^{\alpha}\partial_{\xi}^{\beta}\{e_{j}(x,\xi)\phi(\xi/t_{j})\}| \leq 2^{-j}(1+|\xi|)^{-m-\rho\beta+\delta\alpha-j+1},$$

for $|\xi I \ge t_j$, $|\alpha + \beta| \le j$. Then $e(x, \xi)$ can be written in the form

(3.4)
$$e(x, \xi) = e_0(x, \xi) + \sum_{j=1}^{\infty} e_j(x,\xi) \phi(\xi/t_j)$$

Let H_s , $-\infty < s < \infty$ be the Sobolev space with norm $||u||_s^2 = \int_{-\infty}^{\infty} (1+|\xi|^2)^s |\hat{u}(\xi)|^2 d\xi$. We need the following sharp form of Gårding inequality.

Theorem 3.2. (Kumano-go [7]). Let $p(x, \xi)$ be a function of $S_{1,0}^m$. If $p(x, \xi)$ satisfies the following inequality: For some 8 > 0 and m_1 such that $0 \le m$

¹⁾ We define $u(\xi) = \int_{-\infty}^{\infty} e^{-i\xi x} u(x) dx$ for $u \in \mathscr{S}$.

^{2) &}amp; is the space of rapidly decreasing functions.

 $-m_1 \equiv \theta < 1$,

$$|p(x,\xi)| \geq \delta(1+|\xi|)^{m_1}$$
 for any $x, \xi \in \mathbb{R}^1$,

then

$$\delta^2 ||u||_{s+m-\theta/2}^2 \leq ||p(x, D_x)u||_s^2 + C_s ||u||_{s+m-1/2}^2, \qquad u \in \mathscr{A}.$$

We shall give the characterization of $S_{\rho,\delta}^{-\infty}$.

Theorem 3.3. (Kumano-go [7]). Let $P=p(x,D_x)$ be an element of $S_{\rho,\delta}^{-\infty}$. If we define K(x, w) by

$$K(x, w) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{iw\xi} p(x, \xi) d\xi ,$$

then K(x, w) belongs to $\mathcal{B}(R \times R)$ and K(x, w) satisfies the following: For any α, β , N, there exists a constant $C = C_{\alpha\beta N}$ such that

$$(3.5) \qquad |\partial_x^{\alpha}\partial_w^{\beta}K(x,w)| \leq C(1+|w|^2)^{-N/2} \quad \text{for any } x, w \in \mathbb{R}^1.$$

Pu(x) can be written in the form

(3.6)
$$Pu(x) = \int_{-\infty}^{\infty} K(x, x-y)u(y)dy, \quad u \in \mathcal{S}.$$

Proof. Because $p(x, \xi) \in S_{\rho,\delta}^{-\infty}$, we have

$$(1+|w|^2)^I\partial_x^{\alpha}\partial_w^{\beta}K(x,w)=\frac{1}{2\pi}\int_{-\infty}^{\infty}e^{iw\xi}(1+D_{\xi}^2)^I\{(i\xi)^{\beta}\partial_x^{\alpha}p(x,\xi)\}d\xi.$$

From this we get (3.5). By the Fubini theorem we have (3.6).

For later use we shall quote the index theorem of Kumano-go [8]. We call a function $p(x, \xi)$ of $S^m_{\rho,\delta}$ be slowly varying, if for any α , β ,

$$|D^{\alpha}_{x}\partial^{\beta}_{\xi}p(x,\zeta)| \leq C_{\alpha\beta}(x)(1+|\xi|)^{i+\delta} \quad \text{for any } x, \xi \in \mathbb{R}^{1},$$

where $C_{\alpha\beta}(x)$ is a bounded function such that $C_{\alpha\beta}(x) \rightarrow 0$ as $|x| \rightarrow \infty$, if $\alpha \neq 0$.

Theorem 3.4. (Kumano-go [8]). Let $p(x, \xi) \in S_{p,\delta}^m$, m > 0 be slowly varying. Suppose that there exist positive constants C_0 and $0 < \tau \leq 1$ such that $(p(x, \xi) - \zeta)^{-1}$ exists on

$$\boldsymbol{\Xi}_{\boldsymbol{\xi}} = \{ \boldsymbol{\zeta} \in \boldsymbol{C}; \; \operatorname{dist}(\boldsymbol{\zeta}, (-\infty, 0]) \leq C_{\boldsymbol{0}} (1 + |\boldsymbol{\xi}|)^{\tau_{\boldsymbol{m}}} \}$$

and the estimate of the form

$$(3.7) \qquad |(D^{\alpha}_{x}\partial^{\beta}_{\xi}p(x,\xi))(p(x,\xi)-\zeta)^{-1}| \leq C_{\alpha\beta}(x) (1+|\xi|)^{\delta\alpha-\rho\beta}$$

holds uniformly on Ξ_{ξ} , where $C_{\alpha\beta}(x)$ is a bounded function such that $C_{\alpha\beta}(x) \rightarrow 0$ as $|x| \rightarrow \infty$, if $\alpha \neq 0$. Then $P = p(x, D_x)$ as the map from L^2 into **itself** with the domain $D(P) = \{u \in L^2; Pu \in L^2\}$ has index 0.

4. Construction of the fundamental solution. Throughout this section, we always assume that (a) $\alpha_1 \ge \alpha_2$ and (b) $\sigma(x)^2 \ge \sigma^2 > 0$ or $\alpha_1 > 1$.

Let A be the integro-differential operator defined by (2.1) and $\psi(x, \xi)$ be the function defined by the following equation

(4.1)
$$\psi(x,\xi) = ia(x)\xi - \sigma(x)^2\xi^2/2 + \int_{-\infty}^{\infty} \left[e^{i\xi w} - \frac{\psi_{\xi}^{i\xi\sigma^{n}}}{1 + w} \right]_{-\infty}^{n} (x,w) dw.$$

Then it is easy to see that A can be written in the form

(4.2)
$$Au(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ix\xi} \psi(x,\xi) \hat{u}(\xi) d\xi, \quad u \in \mathcal{S}.$$

In Lemma 4.1, we collect some properties of A as a pseudo-differential operator when the following condition (c) is satisfied. (c) There exists a constant $L_1 > 1$ such that

$$\nu(x, y) = 0 \quad \text{on } R \times \{y; |y| \ge L_1\}.$$

Lemma 4.1. Let the condition (c) be satisfied. Then we have (1°) $\psi(x, \xi)$ belongs to $S_{1,0}^2$. $//\sigma(x)=0$, then ψ belongs to $S_{1,0}^{\alpha_1}$. (2°) For any $\lambda > 0$, the symbol of $\lambda - A$ satisfies the following: There exists a constant $\delta_0 > 0$ such that

(4.3)
$$|\lambda - \psi(x,\xi)| \geq \delta_0 (1+|\xi|)^{\omega_1} \quad \text{for any } x, \xi \in \mathbb{R}^1.$$

(3°) For any $\lambda > 0$ and for any nonnegative integers a, β , there exists $C_{\alpha\beta} > 0$ such that for any $x, \xi \in \mathbb{R}^1$

(4.4)
$$|(D_x^{\alpha}\partial_{\xi}^{\beta}\{\lambda-\psi(x,\xi)\})(\lambda-\psi(x,\xi))^{-1}| \leq C_{\alpha\beta}(1+|\xi|)^{-\beta+\delta\alpha}$$

where $\delta=2-\alpha_1$ if $\alpha_1>1$, $=0$ if $\sigma(x)^2 \geq \sigma^2>0$.

(4°) Choose a constant γ such that $1 < \gamma < 2if \sigma(x)^2 \ge \sigma^2 > 0$, $1 < \gamma < \alpha_1$ if $\alpha_1 > 1$ and fix it. Let A_0 be the pseudo-differential perator having $\psi_0(x, \xi) = (\lambda - ty(x, \xi))$ $(1+ I \xi I^2)^{-\gamma/2}$ as to symbol. Then we have index $(A_0)=0$.

Since a! and σ' belong to \mathcal{B} , it is obvious that $ia(x)\xi$ Proof. Proof of (1°) . $-\sigma(x)^2\xi^2/2$ belongs to $S_{1,0}^2$. Set

$$\phi(x,\xi) = \int_{-L_1}^{L_1} \left(e^{i\xi w} - 1 - \frac{i\xi w}{1 + w^2} \right) n(x, w) dw .$$

Since ϕ can be written in the form

$$\phi(x, \xi) = \int_{-L_1}^0 \left(e^{i\xi w} - 1 - \frac{i\xi w}{1 + w^2} \right) \frac{\nu(x, w)}{|w|^{1 + \omega_2}} dw$$

HITTING PROPERTIES

$$+ \int_{\mathfrak{g}}^{L_1} \left(e^{i\xi w} - 1 - \frac{i\xi w}{1+w^2} \right) \frac{\nu(x,w)}{w^{1+w_1}} dw$$

it is sufficient to consider $F(x, \xi) = \int_{J_0} \left(e^{i\xi w} - 1 - \frac{i\xi w}{1 + w^2} \right) \frac{\nu(x, w)}{w^{1+\omega}} dw$ and to prove that $F(x, w) \in S_{1,0}^1$ if $0 < \alpha < 1$, $F(x, w) \in S_{1,0}^{\alpha}$ if $1 < \alpha < 2$, $|\partial_x^m \partial_\xi^n F(x, w)| \le C_{m,n}(1 + |\xi|)^{1-} (1 + \log(|\xi| \vee 1))$ if $\alpha = 1$.

First we consider the case of $0 < \alpha < 1$. In this case, F(x, f) can be written in the form

$$(4.5) F(x, \xi) = \int_{0}^{L_{1}} (e^{i\xi w} - 1) \frac{\nu(x, w)}{w^{1+\omega}} dw - i\xi \int_{0}^{L_{1}} \frac{\nu(x, w)}{w^{\omega}(1+w^{2})} dw$$

Set $F_1(x,\xi) = \int_0^{L_1} (e^{i\xi w} - 1) \frac{\nu(x,w)}{w^{1+\omega}} dw$ and $F_2(x,\xi) = -i\xi \int_0^{L_1} \frac{\nu(x,w)}{w^{\omega}(1+w^2)} dw$. It is clear that F_2 belongs to $S_{1,0}^1$. We show F_1 belongs to $S_{1,0}^{\omega}$. Set $M_I = \sup_{x,w \in \mathbb{R}^{1,m+n} \leq I} |\partial_x^m \partial_w^n \nu(x,w)|, l=0, 1, 2, \cdots$. If n=0, then we have $\partial_x^m F_1(x,\xi) = \int_0^{L_1} (e^{i\xi w} - 1) |\partial_x^m \nu(x,w)|$ rfro. Therefore, for $|\xi| \leq 1$, noting that $|e^{i\xi w} - 1| \leq |\xi w| \leq |w|$, we get

$$|\partial_x^m F_1(x,\xi)| \leq \int_0^{L_1} \frac{|\partial_x^m \nu(x,w)|}{w^{\alpha}} dw \leq M_m \frac{L_1^{1-\alpha}}{1-\alpha}.$$

For the case $|\xi| > 1$, by putting $|\xi| w = y$, we have

$$\begin{aligned} |\partial_{\alpha}^{m}F_{1}(x,\xi)| &= |\xi|^{\alpha} \left| \int_{0}^{L_{1}|\xi|} \frac{e^{i\xi/|\xi|\cdot y}}{y^{1+\alpha}} \partial_{\alpha}^{m}\nu\left(x,\frac{y}{|\xi|}\right) dy \right| \\ &\leq |\xi|^{\alpha} \left\{ \int_{0}^{1} \frac{M_{m}}{y^{\alpha}} dy + \int_{1}^{L_{1}|\xi|} \frac{2M_{m}}{y^{1+\alpha}} dy \right\} \leq \frac{2-\alpha}{\alpha(1-\alpha)} M_{m} |\xi|^{\alpha}. \end{aligned}$$

Thus we have

(4.6)
$$|\partial_x^m F_1(x,\xi)| \leq C_{m,0}(1+|\xi|)^{\sigma}.$$

If $n \ge 1$, we have

$$\partial_x^m \partial_\xi^n F_1(x,\,\xi) = (i)^n \int_0^{L_1} \frac{e^{i\xi w}}{w^{\alpha-n+1}} \, \partial_x^m \nu(x,\,w) dw \; .$$

For $|\xi| \leq 1$, we get

$$|\partial_x^m \partial_{\xi}^n F_1(x, \xi)| \leq M_m \int_0^{L_1} \frac{dw}{w^{\alpha-n+1}} \leq \frac{M_m L_1^{n-\alpha}}{n-\alpha}.$$

For |f I > 1, we have

S. MANABE

$$\partial_x^m \partial_{\xi}^n F_1(x, \xi) = (i) |\xi|^{\omega - n} \int_0^{L_1|\xi|} e^{i \xi/|\xi| \cdot y} y^{n - \omega - 1} \partial_x^m \nu \left(x, \frac{y}{|\xi|}\right) dy$$

= $(i)^n |\xi|^{\omega - n} F_3(x, \xi).$

For the estimate of $F_3(x,\xi)$, set $F_3^{(1)}(x,\xi) = \int_0^1 e^{i\xi/|\xi|y} y^{n-\alpha-1} \partial_x^m v\left(x,\frac{1}{|\xi|}\right) dy$ and $F_3^{(2)}(x,\xi) = \int_1^{L_1|\xi|} e^{i\xi/|\xi|y} y^{n-\alpha-1} \partial_x^m v\left(x,\frac{y}{|\xi|}\right) dy$. Then we obtain for $F_3^{(1)}$, $|F_3^{(1)}(x,\xi)|$ $\leq M_m \int_0^1 y^{n-\alpha-1} dy = \frac{M}{n-\alpha}$. For $F_3^{(2)}$, by integration by parts, we have

$$F_{3}^{(2)}(x,\xi) = \sum_{l=1} \left(-\frac{|\xi|}{i\xi} \right)^{l} e^{i\xi/|\xi|y} \frac{\partial^{l}}{\partial y^{l}} \left\{ y^{n-\omega-1} \partial_{x}^{m} \nu \left(x, \frac{y}{|\xi|} \right) \right\} \Big|_{y=1} \\ + \left(-\frac{|\xi|}{i\xi} \right)^{n} \sum_{l} {n \choose l} (n-\alpha-1)(n-\alpha-2)\cdots(l-\alpha) \\ \times \int_{1}^{L_{1}|\xi|} e^{i\xi/|\xi|y} y^{l-\omega-1} (\partial_{y}^{l} \partial_{x}^{m} \nu) \left(x, \frac{y}{|\xi|} \right) \frac{dy}{|\xi|^{l}}.$$

For the terms corresponding to $l \ge 1$, we have

$$\begin{aligned} &\left|\frac{1}{|\xi|^{l}}\int_{1}^{L_{1}|\xi|}e^{i\,\xi/|\xi|\,y}y^{l-\alpha-1}(\partial_{y}^{l}\partial_{x}^{m}\nu)\left(x,\frac{y}{|\xi|}\right)dy\right| \\ &=\left|\frac{1}{|\xi|}\int_{0}^{L_{1}}\int_{1/|\xi|}^{L_{1}}e^{i\xi w}w^{l-\alpha-1}(\partial_{w}^{l}\partial_{x}^{m}\nu)(x,w)dw \leq M_{l+m}\int_{1/|\xi|}^{L_{1}}w^{l-\alpha-1}dw\right| \\ &\leq M_{l+m}\frac{L_{1}^{l-\alpha}}{l-\alpha}.\end{aligned}$$

For the term corresponding to l=0,

$$\left|\int_{1}^{L_{1}|\xi|} e^{i\,\xi/|\xi|\,y} \frac{1}{y^{1+\omega}} (\partial_{x}^{m} \nu) \left(x, \frac{y}{|\xi|}\right) dy \right| = M_{m} \int_{1}^{L_{1}|\xi|} \frac{dy}{y^{1+\omega}} \leq \frac{M_{m}}{\alpha}.$$

Thus F_3 is bounded and the inequality

(4.7)
$$I\partial_x^m \partial_{\xi}^n F_1(x, \xi) \setminus \leq C_{m,n} (1+|\xi|)^{\omega-n}, \quad n \geq 1$$

holds. It follows from (4.6) and (4.7) that F_1 belongs to $S^{\alpha}_{1,0}$. Combining the fact that F_2 belongs to $S^1_{1,0}$ and $\alpha < 1$, we have $F \in S^1_{1,0}$.

Next let us consider the case of $1 < \alpha < 2$. Set

$$F_{1}(x,\,\xi) = \int_{0}^{L_{1}} (e^{i\xi w} - 1 - i\xi w) \frac{\nu(x,\,w)}{w^{1+\omega}} dw \text{ and}$$

$$1) \quad (\partial_{y} \partial_{x}^{m} \nu) \left(x, \frac{y}{|\xi|}\right) \text{ means } \frac{\partial^{i+m} \nu(x,z)}{\partial x^{m} \partial z^{i}} \Big|_{z=\frac{y}{|\xi|}}$$

$$F_2(x,w)=i\xi\int_0^{L_1}\frac{w^{2-w}}{1+w^2}\nu(x,w)dw.$$

Then F can be written in the form $F(x, \xi) = F_1(x,\xi) + F_2(x,\xi)$. It is evident that F_2 belongs to S_{10}^1 . In order to prove (3.1) for F_1 , it suffices to show (3.1) for the case $\beta = 0$. In fact, since

$$\partial_{\xi}F_{1}(x, \xi) = i \int_{0}^{L_{1}} \frac{e^{i \xi w} - 1}{w^{1+(\omega-1)}} \nu(x, w) dw$$
,

it reduces to the case $0 < \alpha < 1$. If $\beta = 0$, then we have for $|\xi| \ge 1$,

$$\begin{split} |\partial_x^m F_1(x,\,\xi)| &= |\xi|^{\sigma} \Big| \int_0^{|\xi|L_1} \Big(e^{i\xi/|\xi|w} - 1 - i\frac{\xi}{|\xi|} w \Big) \frac{\partial_x^m \nu \Big(x,\frac{w}{|\xi|}\Big)}{w^{1+\sigma}} dw \Big| \\ &\leq |\xi|^{\sigma} \Big\{ \int_0^1 \frac{M_m}{w^{\sigma-1}} dw + \int_1^\infty \frac{2+w}{w^{1+\sigma}} M_m dw \Big\} \\ &= \Big(\frac{2}{\alpha} + \frac{1}{\alpha-1} + \frac{1}{2-\alpha} \Big) M_m |\xi|^{\sigma} \,. \end{split}$$

For $|\boldsymbol{\xi}| < 1$, noting $|e^{i\boldsymbol{\xi}w} - 1 - i\boldsymbol{\xi}w| \le \frac{1}{2} |\boldsymbol{\xi}w|^2 \le \frac{1}{2} |w|^2$, we have $|\partial_{\boldsymbol{x}}^{\boldsymbol{m}} F_1(\boldsymbol{x}, \boldsymbol{\xi})| \le M_{\boldsymbol{m}} \int_{0}^{L_1} w^{1-\boldsymbol{\omega}} dw = \frac{M_{\boldsymbol{m}}}{2-\boldsymbol{\alpha}} (L_1)^{2-\boldsymbol{\omega}}.$

Thus F_1 belongs to $S_{1,0}^{\alpha}$. Combining this and the fact that $F_2 \in S_{1,0}^1$, we obtain that F belongs to $S_{1,0}^{\alpha}$.

Finally consider the case $\alpha = 1$. We have

(4.8a)
$$\partial_x^m F(x,\xi) = \int_0^{L_1} \left\{ e^{i\xi w} - 1 - \frac{i\xi w}{1+w^2} \right\} \frac{\partial_x^m v(x,w)}{w^2} dw,$$

(4.8b)
$$\partial_x^m \partial_{\xi} F(x, \xi) = i \int_0^{L_1} (e^{i\xi w} - 1) \frac{\partial_x^m \nu(x, w)}{w} dw + i \int_0^{L_1} \frac{w \partial_x^m \nu(x, w)}{1 + w^2} dw,$$

(4.8c)
$$\partial_x^m \partial_{\xi}^n F(x, \xi) = (i)^n \int_0^{L_1} e^{i\xi w} w^{n-2} \partial_x^m \nu(x, w) dw, \ n \ge 2.$$

First we consider the case n=0. If $|\xi| \leq 1$, then we have

$$\begin{aligned} |\partial_x^m F(x,\xi)| &\leq \left| \int_0^{L_1} (e^{i\xi w} - 1 - i\xi w) \frac{\partial_x^m \nu(x,w)}{w^2} dw \right| + |\xi| \left| \int_0^{L_1} \frac{\partial_x^m \nu(x,w)}{1 + w^2} dw \right| \\ &\leq 2M_m L_1 \,. \end{aligned}$$

If $I \xi I > 1$, then we have

$$\begin{split} |\partial_{x}^{m}F(x,\xi)| &= |\xi| \Big| \int_{0}^{L_{1}|\xi|} \Big\{ e^{i \cdot sgn(\xi)y} - 1 - \frac{i \cdot sgn(\xi)y}{1 + y^{2}/\xi^{2}} \Big\} \frac{\partial_{x}^{m}\nu\left(x,\frac{y}{|\xi|}\right)}{y^{2}} dy \Big| \\ &\leq |\xi| \Big| \int_{0}^{1} \Big\{ e^{i \cdot sgn(\xi)y} - 1 - \frac{i \cdot sgn(\xi)y}{1 + y^{2}/\xi^{2}} \Big\} \frac{\partial_{x}^{m}\nu\left(x,\frac{y}{|\xi|}\right)}{y^{2}} dy \Big| \\ &+ \langle \xi \rangle \Big| \int_{1}^{L_{1}|\xi|} \Big\{ e^{i \cdot sgn(\xi)y} - 1 - \frac{i \cdot sgn(\xi)y}{1 + y^{2}/\xi^{2}} \Big\} \frac{\partial_{x}^{m}\nu\left(x,\frac{y}{|\xi|}\right)}{y^{2}} dy \Big| \\ &\equiv I_{1}|\xi| + I_{2}|\xi| \,. \end{split}$$

For I_1 , we have

$$\begin{split} I_{1} &\leq \Big| \int_{0}^{1} \{ e^{i \cdot sgn(\xi)y} - 1 - i \cdot sgn(\xi)y \} \frac{\partial_{x}^{m} \nu \left(x, \frac{y}{|\xi|}\right)}{y^{2}} dy \Big| \\ &+ \Big| \int_{0}^{1} i \cdot sgn(\xi)y \left(1 - \frac{1}{1 + y^{2}/\xi^{2}}\right) \frac{\partial_{x}^{m} \nu \left(x, \frac{y}{|\xi|}\right)}{y^{2}} dy \Big| \\ &\leq \int_{0}^{1} \Big| \partial_{x}^{m} \nu \left(x, \frac{y}{|\xi|}\right) \Big| dy + \int_{0}^{1} \frac{\Big| \partial_{x}^{m} \nu \left(x, \frac{y}{|\xi|}\right) \Big|}{y^{2} + \xi^{2}} dy \leq 2M_{m} \,. \end{split}$$

For I_2 , we have

$$\begin{split} I_{2} &\leq \left| \int_{1}^{L_{1}|\xi|} \{e^{i \cdot sgn(\xi)y} - 1\} \frac{\partial_{x}^{m} \nu \left(x, \frac{y}{|\xi|}\right)}{y^{2}} dy \right| \\ &+ \left| \int_{1}^{L_{1}|\xi|} \frac{-i \cdot sgn(\xi)y}{(1 + y^{2}/\xi^{2})y^{2}} \partial_{x}^{m} \nu \left(x, \frac{y}{|\xi|}\right) dy \right| \\ &\leq 2M_{m} \int_{1}^{\infty} \frac{dy}{y^{2}} + \int_{1}^{L_{1}|\xi|} \frac{M_{m}}{y(1 + y^{2}/\xi^{2})} dy = 2M_{m} + M_{m} \int_{1/|\xi|}^{L_{1}} \frac{dw}{w(1 + w^{2})} \\ &\leq M_{m} (2 + \log L_{1}|\xi|). \end{split}$$

Therefore we have

(4.9)
$$|\partial_x^m F(x,\xi)| \leq C_{m,0}(1+|\xi|) (1+\log(|\xi|\vee 1)).^{1}$$

Next we consider the case n=1. The second term of (4.8b) is bounded. Denote by / the first term of (4.8b). For $|\xi| > 1$, we have

¹⁾ $a \lor b = \max(a, b), a \land b = \min(a, b).$

$$|/| = \left| \int_{0}^{L_{1}|\xi|} \frac{e^{\operatorname{sgn}(\xi)y}}{y} - \frac{1}{2} \partial_{x}^{m} v\left(x \frac{y}{|\xi|}\right) dy \right| \leq 2M_{m} \log L_{1}|\xi|.$$

For $|\xi| \leq 1$, we have

$$|I| \leq \int_0^{L_1} |\xi| |\partial_x^m \nu(x, w)| dw \leq M_m L_1.$$

Therefore we get

(4.10)
$$I \partial_x^m \partial_{\xi} F(x, \xi) \setminus \leq C_{m,1} (1 + \log(|\xi| \vee 1)) .$$

Finally we consider the case $n \ge 2$. If $|\xi| \le 1$, then we obtain

$$|\partial_x^m \partial_{\xi}^n F(x, \xi)| \leq M_m \int_0^{L_1} w^{n-2} dw = \frac{M_m}{n-1} L_1^{n-1}.$$

If $I \xi I > 1$, then we have

$$\partial_x^m \partial_\xi^n F(x,\,\xi) = \frac{(i)^n}{|\xi|^{n-1}} \int_0^{L_1|\xi|} e^{i \cdot sg_n(\xi)y} y^{n-2} \partial_x^m \nu\left(x,\frac{y}{|\xi|}\right) dy$$

= $\frac{(i)^n}{|\xi|^{n-1}} \left\{ \int_0^1 e^{i \cdot sg_n(\xi)y} y^{n-2} \partial_x^m \nu\left(x,\frac{y}{|\xi|}\right) dy + \int_1^{L_1|\xi|} e^{i \cdot sg_n(\xi)y} y^{n-2} \partial_x^m \nu\left(x,\frac{y}{|\xi|}\right) dy \right\}.$

Set

$$J_1 = \int_0^1 e^{i \cdot sgn(\xi)y} y^{n-2} \partial_x^m \nu\left(x, \frac{y}{|\xi|}\right) dy \text{ and } \int_1^{L_1|\xi|} e^{i \cdot sgn(\xi)y} y^{n-2} \partial_x^m \nu\left(x, \frac{y}{|\xi|}\right) dy.$$

Then we have for J_1 ,

$$|J_1| \leq M_m/(n-1)$$
.

For J_2 , by integration by parts, we get

$$J_{2} = \sum_{l=1}^{n-1} \left(-\frac{1}{i \cdot sgn(\xi)} \right)^{l} e^{i \cdot sgn(\xi) y} \frac{\partial^{l}}{\partial y^{l}} \left\{ y^{n-2} \partial_{x}^{m} \nu \left(x, \frac{y}{|\xi|} \right) \right\} \Big|_{y=1} + \left(-\frac{1}{i \cdot sgn(\xi)} \right)^{n-1} \sum_{l=1}^{n-1} \left(\frac{h-1}{k} \right)^{l} (n-2) (n-3) \cdots l \\ \times \int_{1}^{L_{1}|\xi|} e^{i \cdot sgn(\xi) y} y^{l-1} (\partial_{y}^{l} \partial_{x}^{m} \nu) \left(x, \frac{y}{|\xi|} \right) \frac{dy}{|\xi|^{l}}.$$

For the terms corresponding to $l \ge 1$, we have

$$\left|\frac{1}{|\xi|^{i}}\int_{1}^{L_{1}|\xi|}e^{i\cdot sgn(\xi)y}y^{l-1}(\partial_{y}^{l}\partial_{x}^{m}\nu)\left(x,\frac{y}{|\xi|}\right)dy\right| \leq M_{m+n-1}(L_{1})^{n-1}.$$

For the term corresponding to l=0, we have

$$\left|\int_{1}^{L_{1}|\xi|} e^{i \cdot sgn(\xi)y} \frac{1}{y} (\partial_{x}^{m} \nu) \left(x, \frac{y}{|\xi|}\right) dy \right| \leq M_{m} \log L_{1}|\xi| .$$

Therefore we get $|J_2| \leq C(1+\log|\xi|)$ for $|\xi| > 1$. Hence we have

$$(4.11) \qquad \qquad |\partial_x^m \partial_\xi^n F(x,\varsigma) \setminus \leq C_{m,n} (1+|\xi|)^{1-n} (1+\log(|\xi|\vee 1))$$

It follows from (4.9), (4.10) and (4.11) that

$$|\partial_x^m \partial_\xi^n F(x,\xi) \setminus \leq C_{m,n}(1+|\xi|)^{1-n}(1+\log(|\xi \setminus \vee 1)).$$

Thus the proof of (1°) is completed.

Proof of (2°) . Note that

$$\lambda - \psi(x, \xi) = \lambda + \frac{1}{2} \sigma(x)^2 \xi^2 + \int_{-\infty}^{\infty} (1 - \cos \xi w) n(x, w) dw$$
$$- i \left\{ a(x)\xi + \int_{-\infty}^{\infty} \left(\frac{\xi w}{1 + w^2} - \sin \xi w \right) n(x, w) dw \right\}.$$

Set
$$c_2 = \int_{J_0}^{1} \frac{1 - \cos y}{y}$$
. Then usi $\langle 1 \rangle$, we have for $|\xi| > 1$
 $|\lambda - \psi(x, \xi)| \ge \lambda + \frac{1}{2} \sigma(x)^2 \xi^2 + \int_{-\infty}^{\infty} (1 - \cos \xi w) n(x, w) dw$
 $\ge \lambda + \frac{1}{2} \sigma(x)^2 \xi^2 + |\xi|^{\sigma_1} \int_{0}^{1} (1 - \cos y) \frac{\nu(x, \frac{y}{|\xi|})}{y^{1+\sigma_1}} dy$
 $\ge \lambda + \frac{1}{2} \sigma(x)^2 \xi^2 + c_1 c_2 |\xi|^{\sigma_1}.$

Thus (2°) is proved.

Proof of (3°). Let $\alpha_1 > 1$. For $m \ge 1$, we have by (1°) and (2°)

$$\left|\frac{\partial_{\xi}^{n}\partial_{x}^{m}(\lambda-\psi(x,\xi))}{\lambda-\psi(x,\xi)}\right| \leq C_{m,n}(1+|\xi|)^{-n+(2-\omega_{1})} \leq C_{m,n}(1+|\xi|)^{-n+(2-\omega_{1})m}$$

For the case m=0 and $n \ge 2$, we have

$$\left|\frac{\partial_{\boldsymbol{\xi}}^{\boldsymbol{n}}(\lambda-\psi(\boldsymbol{x},\,\boldsymbol{\xi}))}{\lambda-\psi(\boldsymbol{x},\,\boldsymbol{\xi})}\right| \leq C_{\boldsymbol{0},\boldsymbol{n}}(1+|\boldsymbol{\xi}|)^{-\boldsymbol{n}}.$$

For m = 0 and n = 1, we have

HITTING PROPERTIES

$$\left|\frac{\partial_{\boldsymbol{\xi}}(\lambda - \psi(\boldsymbol{x}, \, \boldsymbol{\xi}))}{\lambda - \psi(\boldsymbol{x}, \, \boldsymbol{\xi})}\right| \leq \frac{\sigma(\boldsymbol{x})^2 |\boldsymbol{\xi}| + C_3(1 + |\boldsymbol{\xi}|)^{\boldsymbol{\sigma}_1^{-1}}}{\lambda + \frac{1}{2}\sigma(\boldsymbol{x})^2 \boldsymbol{\xi}^2 + C_4(1 + |\boldsymbol{\xi}|)^{\boldsymbol{\sigma}_1}} \leq C_{0,1}(1 + |\boldsymbol{\xi}|)^{-1}.$$

Therefore we get

$$\left|\frac{\partial_{\xi}^{n}\partial_{x}^{m}(\lambda-\psi(x,\xi))}{\lambda-\psi(x,\xi)}\right| \leq C_{m,n}(x) (1+|\xi|)^{-n+(2-\omega_{1})m}.$$

In case $\sigma(x)^2 \ge \sigma^2 > 0$, (4.4) is clear. The proof of (3°) is completed.

Proof of (4°). It is sufficient to verify that the conditions in Theorem 3.4 are satisfied for A_0 . First note that $\lambda - \psi(x, \xi)$ is slowly varying. This follows from the fact that a'(x) and $\sigma'(x)$ belong to \mathring{B} and $\partial_x^m \nu(x, y)$, $m \ge 1$ are zero for $|x| \ge L$. Next we show the estimate (3.7) for $\psi_0(x, \xi)$. If $\eta = \psi_0(x, \xi)$, then we have

dist
$$(\eta, (-\infty, 0]) = |\eta| \ge \operatorname{Re} \eta \ge \delta_0 (1 + |\xi|^2)^{(\alpha_1 - \gamma)/2}$$
.

Therefore $\{(\lambda - \psi(x, \xi))(1 + |\xi|^2)^{-\gamma/2} - \zeta\}^{-1}$ exists on Ξ_{ξ} if we choose $\tau < \alpha_1^{-1}(\alpha_1 - \gamma)$ Then since $\operatorname{Re}(\lambda - \psi(x, \xi)) > 0$,

$$\begin{aligned} |(\lambda - \psi(x, \xi)) (1 + \xi^2)^{-\gamma/2} - \zeta| &\geq |\operatorname{Re}(\lambda - \psi(x, \xi)) (1 + \xi^2)^{-\gamma/2} - \operatorname{Re} \zeta| \\ &\geq \operatorname{Re}(\lambda - \psi(x, \xi)) - \operatorname{Re} \zeta \quad \text{if } \operatorname{Re} \zeta \geq 0, \\ &\geq \operatorname{Re}(\lambda - \psi(x, \xi)) \quad \text{if } \operatorname{Re} \zeta \leq 0. \end{aligned}$$

In view of the proof of (2°) , we have

$$|(\lambda - \psi(x, \xi)) (1 + \xi^2)^{-\gamma/2} - \zeta| \ge C_3 |(\lambda - \psi(x, \xi)) (1 + \xi^2)^{-\gamma/2}|$$
 on Ξ_{ξ} .

It is easy to see that

$$|\partial_x^{\boldsymbol{a}}\partial_{\boldsymbol{\xi}}^{\boldsymbol{\beta}}(\lambda-\psi(x,\,\boldsymbol{\xi}))\,(1+\boldsymbol{\xi}^2)^{-\boldsymbol{\gamma}/2}| \leq C_{\boldsymbol{a},\boldsymbol{\beta}}|\partial_x^{\boldsymbol{a}}\partial_{\boldsymbol{\xi}}^{\boldsymbol{\beta}}(\lambda-\psi(x,\,\boldsymbol{\xi}))|\,(1+\boldsymbol{\xi}^2)^{-\boldsymbol{\gamma}/2}$$

Combining these two estimates and (4.4) it follows that the estimate (3.7) holds. Thus the conditions of Theorem 3.4 holds for $\psi_0(x, \xi)$. The proof of (4°) is completed.

Although the following remark is well known, we shall state here for later use.

REMARK 4.1. (Maximum principle). Let $u(\equiv 0)$ be a function of \mathcal{B} . If $u(x_0)=\sup_{x\in\mathcal{B}^+}u(x)$ for some x_0 , then $Au(x_0)\leq 0$. Moreover there exists a point x_1 such that $u(x_1)=u(x_0)$ and $Au(x_1)<0$.

Now we shall show the existence of the Green operator of A and construct its kernel.

Lemma 4.2. Let the condition (c) be satisfied and A be the operator defined by (2.1). Then we have

S. MANABE

(i) For any $f \in H_{\infty}$, there exists a unique solution $u \in \dot{\mathcal{B}}$ of the equation

$$(4.12) \qquad \qquad (\lambda - A)u = f$$

(ii) For any $f \in H_{\infty}$, define G_{λ} fas the unique solution of (4.12) Then G_{λ} is a continuous operator from H_{∞} to $\dot{\mathcal{B}}$.

Proof. First we note that A can be written in the form

(4.13)
$$Au(x) = \mathcal{F}^{-1}(\psi(x, \)\mathcal{F}u(\cdot))(x), u \in \mathcal{B},$$

where $\mathcal{F}u, \mathcal{F}^{-1}u$ denote the Fourier transform and the Fourier inverse transform in distribution sense respectively. In fact, for any $u \in \dot{\mathcal{B}}$, there exists a sequence $\{u_n\}$ of $\dot{\mathcal{S}}$ such that $u_n \rightarrow u$ in $\dot{\mathcal{B}}$. Therefore $\mathcal{F}u_n \rightarrow \mathcal{F}u$ in $\dot{\mathcal{S}}'$. Because $\psi(x, \) \in \mathcal{O}_M^{(1)}$, we have $\psi(x, \)\mathcal{F}u_n \rightarrow \psi(x, \)\mathcal{F}u$ in $\dot{\mathcal{S}}'$. Define A_0 by

$$A_{0}u(x) = \mathcal{F}^{-1}((\lambda - \psi(x, \xi)) (1 + |\xi|^{2})^{-\gamma/2} \mathcal{F}u(\xi))(x), \qquad u \in \dot{\mathcal{B}}(1 < \gamma < m^{2}).$$

Consider A_0 as a map of L^2 with domain $D(A_0) = \{u \in L^2; A_0 u \in L^2\}$. First we show that A_0 is one-to-one. Let u be a function of L^2 such that $A_0 u = 0$. Set

$$g(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{ix\xi}}{(1+|\xi|^2)^{\gamma/2}} d\xi$$
 and $v(x) = g * u(x)$.³⁾

Since the symbol ψ_0 satisfies the condition of Theorem 3.2 for ra=2 $-\gamma$, $m_1=2$ - γ if $\sigma(x)^2 \ge \sigma^2 > 0$, $\alpha_1 - \gamma$ if $\alpha_1 > 1$, we have $||u||_{s+m-\theta/2} \le C'_s ||u||_{s+m-1/2} (\forall s)$. Therefore $u \in H_{\infty}$. Combining this and $g \in L^1$, we obtain $v \in \dot{\mathcal{B}}$. So we have

$$(\lambda - A)v(x) = \mathcal{F}^{-1}\{(\lambda - \psi(x, \cdot)) \mathcal{F}v(\cdot)\}(x) = A_0v(x) = 0$$

By Remark 4.1, we have v(x)=0, hence u(x)=0. Therefore $\ker(A_0)=\{0\}$. Combining this and $\operatorname{index}(A_0)=0$, we have $\operatorname{coker}(A_0)=L^2/\operatorname{Im}(A_0)=\{0\}$. Thus A_0 maps $D(A_0)$ onto $L^2(\mathbb{R}^1)$. Let f be any function of H_∞ . Then since / belongs to L^2 , it follows that there exists a unique $v \in L^2$ such that $A_0v=f$. Set $u=g_*v$. Then by Theorem 3.2, $u \in \mathcal{B}$ and using the same argument as above, we have $(\lambda - A)u=f$. Uniqueness follows easily from Remark 4.1.

Finally we shall prove the continuity of G_{λ} . By the closed graph theorem, it is sufficient to show the following:

$$f, f_n \in H_{\infty}, f_n \to f i H_{\infty}, G_{\lambda} f_n \to v \text{ in } \dot{\mathcal{B}}, \text{ then } v = G_{\lambda} f.$$

Since it follows from (2.1) that A maps $\dot{\mathcal{B}}$ into $\dot{\mathcal{B}}$ continuously, we have $f_n = (\lambda - A)G_{\lambda}f_n \rightarrow (\lambda - A)in \dot{\mathcal{B}}$. On the other hand, $f_n \rightarrow f$ in H_{∞} , we have

3) $g^*u(x) = \int_{-\infty}^{\infty} g(x-y)u(y)dy.$

¹⁾ $\mathcal{O}_{\mathbf{M}}$ is the totality of C^{∞} -functions such that each derivative is dominated by some polynomial.

²⁾ m=2 if $\sigma(x)^2 \ge \sigma^2 > 0$, $=\alpha_1$ if $\alpha_1 > 1$.

 $f=(\lambda - A)v$ in $\dot{\mathcal{B}}$. The proof is completed.

By Lemma 4.1, there exists a parametrix $Q=q(x, D_x)$ of $\lambda - A$, where $q \in S_{1,2-\alpha_1}^{-\alpha_1}$. Define K by the following equation:

$$(4.14) K = I - (\lambda - A)Q.$$

Then K maps $H_{-\infty}$ to H_{∞} continuously. Hence $G_{\lambda}K=G_{\lambda}-Q$ maps $H_{-\infty}$ to \mathcal{B} continuously. This implies that $G_{\lambda}-Q$ maps \mathcal{E}' to 6 continuously. By Schwartz' kernel theorem (Schwartz [13]), there exists a C^{∞} -function k(x, y) such that

(4.15)
$$G_{\lambda}f(x) - Qf(x) \int_{-\infty}^{\infty} f(x,y)f(y)dy, \quad f \in \mathcal{D}^{1}.$$

Set

(4.16)
$$g_{\lambda,0}(x, y) = \int_{2\pi} \int_{-\infty} e^{i(x-y)\xi} q_0(x, \xi) d\xi,$$

(4.17)
$$g_{\lambda,1}(x,y) = \sum_{j=1}^{\infty} \frac{1}{2\pi J} \int_{-\infty}^{\infty} e^{i(x-y)\xi} q_j(x,\xi) \phi\left(\frac{\xi}{t_j}\right) d\xi + k(xy)$$

and

(4.18)
$$g_{\lambda}(x,y) = g_{\lambda,0}(x, y) + g_{\lambda,1}(x, y)$$

Since fceS $\sum_{2-\infty}^{\alpha_1-}$, $k=0, 1, 2, \cdots, \int_{-\infty}^{\infty} e^{i(x-y)\xi} q_0(x,\xi) d\xi$ and $\int_{-\infty}^{\infty} e^{i(x-y)\xi} q_j(x,\xi) \phi\left(\frac{\xi}{t_j}\right) d\xi$ are well-defined. By (3.2) of Remark 3.1, the right hand side of (4.17) converges uniformly in x, y.

Lemma 4.3. Under the same condition in Lemma 4.2, (i) G_{λ} has the kernel representation:

(4.19)
$$G_{\lambda}f(x) = \int_{-\infty}^{\infty} g_{\lambda}(x, y)f(y)dy, \quad f \in \mathcal{D},$$

where g_{λ} is the function defined by (4.18).

(ii) (a) $g_{\lambda}(x,y)$ is a nonnegative function, (b) $g_{\lambda,0}(x,y)$ is a continuous function and C^{∞} -function except on the diagonal set, (c) $g_{\lambda,1}(x, y)$ is a continuously differentiable function and C^{∞} -function except on the diagonal set.

Proof. First note that

(4.20)
$$G_{\lambda}f(x) = Qf(x) + \int_{-\infty}^{\infty} k(xy)f(y)dy, \quad f \in \mathcal{D} .$$

On the other hand, Qf(x) can be written in the form

¹⁾ \mathscr{D} is the totality of C^{∞} -functions on \mathbb{R}^1 of compact support.

S. MANABE

$$Qf(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i \mathbf{x} \cdot \mathbf{\xi}} q(x, \, \mathbf{\xi}) \hat{f}(\mathbf{\xi}) d\mathbf{\xi}, \qquad f \in \mathcal{S}$$

Using the fact that $q_k \in S_{1,2-\alpha_1}^{-\alpha_1-k}$, $k=0, 1, 2, \cdots$ and (3.2), we have

$$Qf(x) = \int_{-\infty}^{\infty} \left\{ \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(x-y)\xi} q_0(x,\xi) d\xi + \sum_{j=1}^{\infty} \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(x-y)\xi} q_j(x,\xi) \phi\left(\frac{\xi}{t_j}\right) d\xi \right\} f(y) dy.$$

From this and (4.20), we obtain (4.19). The proof of (i) is completed. Next we show (ii). It is clear that $g_{\lambda,0}$ and $g_{\lambda,1}$ are continuous in x, y. So g_{λ} is continuous in x, y. By Remark 4.1, we have $G_{\lambda}f \ge 0$ if $f \ge 0$. Therefore g_{λ} is nonnegative. Although the proof of the fact that $g_{\lambda,0}$ and $g_{\lambda,1}$ belong to $C^{\infty}(R \times R - \Delta)$ is found in [2], for completeness we present the proof. By integration by parts, we have

$$g_{\lambda,0}(x,y) = \frac{(i)^{k+l}}{2\pi(x-y)^{k+l}} \int_{-\infty}^{\infty} e^{i(x-y)\xi} \frac{\partial^{k+l}}{\partial \xi^{k+l}} q_0(x,\xi) d\xi .$$

Since $q_0 \in S_{1,2-\alpha_1}^{-\alpha_1}$, we have

$$\begin{split} \left| \frac{\partial^{k+l}}{\partial x^{k} \partial y^{l}} \left\{ e^{i(x-y)\xi} \frac{\partial^{k+l}}{\partial \xi^{k+l}} q_{0}(x, \xi) \right\} \right| \\ &= \left| (-i\xi)^{l} \sum_{r=0}^{k} \binom{k}{r} (i\xi)^{r} e^{i(x-y)\xi} \frac{\partial^{k-r+k+l} q_{0}(x, \xi)}{\partial x^{k-r} \partial \xi^{k+l}} \right| \\ &\leq C_{5} |\xi|^{l} \sum_{r=0}^{k} \binom{k}{r} |\xi|^{r} (1+|\xi|)^{-\omega_{1}-(k+l)+\delta(k-r)} \\ &\leq C_{6} (1+|\xi|)^{-\omega_{1}}. \end{split}$$

Therefore we have

$$\frac{\partial^{k+l}g_{\lambda,0}}{\partial x^k \partial y^l}(x,y) = \frac{1}{2\pi} \sum_{r_1=0}^k \sum_{r_2=0}^l \binom{k}{r_1} \binom{l}{r_2} \left\{ \frac{\partial^{r_1+r_2}}{\partial x^{r_1} \partial y^{r_2}} \frac{1}{(y-x)^{k+l}} \right\} \\ \times \int_{-\infty}^\infty \frac{\partial^{(k-r_1)+(l-r_2)}}{\partial x^{k-r_1} \partial y^{l-r_2}} \left\{ e^{i(y-x)\xi} \frac{\partial^{k+l}q_0}{\partial \xi^{k+l}}(x,\xi) \right\} d\xi \,.$$

Thus we have $g_{\lambda,0} \in C^{\infty}(R \times R - \Delta)$. By the same way we have $g_{\lambda,1} \in C^{\infty}(R \times R - \Delta)$ $\prod C^{1}(R \times R)$. The proof of (ii) is completed.

5. The estimates of the singularity of $\frac{\partial g_{\lambda}}{\partial y}$. In this section we estimate the singularity of $\partial g_{\lambda}/\partial y$. By Lemma 4.3 it is sufficient to consider $g_{\lambda,0}(x, y)$ only.

Lemma 5.1. Let $g_{\lambda,0}(x, y)$ be the function defined by (4.16). Then we have

HITTING PROPERTIES

(5.1)
$$\frac{\partial g_{\lambda,0}}{\partial y}(x, y) = -\frac{1}{\pi} \int_0^\infty \frac{\xi [\lambda - \psi_R(x, \xi)] \sin(y - x)\xi}{[\lambda - \psi_R(x, \xi)]^2 + \psi_I(x, \xi)^2} d\xi + \frac{1}{\pi} \int_0^\infty \frac{\xi \psi_I(x, \xi) \cos(y - x)\xi}{[\lambda - \psi_R(x, \xi)]^2 + \psi_I(x, \xi)^2} \xi, \quad \text{for } x \neq y,$$

where ψ_R and ψ_I are the real part and the imaginary part of ψ respectively.

Proof. Since fy_R and ψ_I are even and odd functions of ξ respectively, we have

$$g_{\lambda,0}(x,y) = \frac{1}{\pi} \int_{0}^{\infty} \frac{[\lambda - v \ln(x,\xi)] \cos(v - x)\xi}{[\lambda - \psi_R(x,\xi)]^2 + \psi_I(x,\xi)^2} d\xi + \frac{1}{\pi} \int_{0}^{\infty} \frac{v \ln(x,\xi) \sin(v - x)\xi}{[\lambda - \psi_R(x,\xi)]^2 + \psi_I(x,\xi)^2} d\xi.$$

Therefore it suffices to prove that the following integrals

$$I_{1} = \int_{J_{0}}^{\infty} \frac{\xi\{\lambda - \psi(x, \xi)\}\sin(y - x)\xi}{[\lambda - \psi_{R}(x, \xi)]^{2} + \psi_{I}(x, \xi)^{2}} d\xi$$

and

$$L_{\overline{2}} \underbrace{=}_{\operatorname{Jo}} \int_{\operatorname{O}}^{\infty} \underbrace{\xi \psi_{I}(x,\xi) \cos(y-x)\xi}_{\left[\lambda - \psi_{R}(x,\xi)\right]^{2} + \psi_{I}(x,\xi)^{2}} d\xi$$

are uniformly convergent for $|x-y| \ge \delta > 0$. First we show this for I_1 . Set

$$F_1(\xi) = \int_0^t \left\{ rac{\partial}{\partial \eta} \left(rac{\eta [\lambda - \psi_R(x, \eta)]}{[\lambda - \psi_R(x, \eta)]^2 + \psi_I(x, \eta)^2}
ight)
ight\} ee 0 \ d\eta$$

and

$$F_2(\xi) = -\int_0^{\xi} \left\{ rac{\partial}{\partial \eta} \left(rac{\eta [\lambda - \psi_R(x, \eta)]}{[\lambda - \psi_R(x, \eta)]^2 + \psi_I(x, \eta)^2}
ight)
ight\} \wedge 0 \; d\eta \; .$$

Then F_1 and F_2 are monotone increasing and

$$rac{\xi[\lambda-\psi_R(x,\xi)]}{[\lambda-\psi_R(x,\xi)]^2+\psi_I(x,\xi)^2}=F_{\scriptscriptstyle 1}(\xi)\!-\!F_{\scriptscriptstyle 2}(\xi)\,.$$

Note that the set of pseudo-differential operators forms an algebra, that is if $p_i(x, D_x)$ belongs to $S_{\rho,\delta}^{m_i}$, i=1, 2, then $p_1(x, D_x)p_2(x, D_x)$ belongs to $S_{\rho,\delta}^{m_1+m_2}$ (see [6]). Using this, we have $\frac{\eta[\lambda - \psi_R(x, \eta)]}{[\lambda - \psi_R(x, \eta)]^2 + \psi_I(x, \eta)^2}$ belongs to $S_{1.0}^{-m+1}$ for some m > 1. Hence we get

$$\left|\frac{\partial}{\partial \eta}\left\{\frac{\eta[\lambda-\gamma]}{[\lambda-\psi_R(x,\eta)]^2+\psi_I(x,\eta)^2}\right\}\right|-\frac{c_4}{\eta^m} \text{ for } |\eta| \ge M,$$

S. Manabe

where *M* is a constant independent of *x*. According to this and $\lim_{\substack{i \in I \to \infty \\ \downarrow \land - \psi_R \end{bmatrix}} - = 0 \text{ uniformly in } x, \text{ we have } \lim_{\substack{\xi \to \infty \\ \xi \to \infty }} F_1(\xi) = \lim_{\xi \to \infty} F_2(\xi) = a. \text{ Set}$ $F_3(\xi) = F_1(\xi) - a \text{ and } F_4(\xi) = F_2(\xi) - a. \text{ Then } F_3 \text{ and } F_4 \text{ converge to } 0 \text{ uniformly}$ in *x* and

$$\frac{\xi[\lambda-\psi_R(x,\xi)]}{[\lambda-\psi_R(x,\xi)]^2+\psi_I(x,\xi)^2}=F_3(\xi)-F_4(\xi)\,.$$

Therefore by the second mean value theorem, for $|x-y| \ge \delta > 0$,

$$\begin{split} & \left| \int_{N_2}^{N_2} \xi[\lambda - \psi_R(x, \xi)] \sin(y - x) \xi}_{\lambda_1} d\xi \right| \\ & \leq \left| \int_{N_1}^{N_2} F_3(\xi) \sin(y - x) \xi d\xi \right| + \left| \int_{N_1}^{N_2} F_4(\xi) \sin(y - x) \xi d\xi \right| \\ & \leq \frac{2}{\delta} \left\{ F_3(N_1) + F_3(N_2) + F_4(N_1) + F_4(N_2) \right\} \to 0, \text{ as } N_1, N_2 \to \infty . \end{split}$$

The proof for I_2 is the same as above.

Lemma 5.2. (i) If $\sigma(x)^2 \ge \sigma^2 > 0$ on R^1 or (ii) $\sigma(x)^2 \ge \sigma^2 > 0$ on a neighborhood of the origin and $\alpha_1 > 1$, then

(5.2a)
$$\lim_{x \downarrow 0} \frac{\partial g_{\lambda,0}}{\partial y}(x,0) = \lim_{y \downarrow 0} \frac{\partial g_{\lambda,0}}{\partial y}(0,y) = -\frac{1}{\sigma(0)^2} + C,$$

(5.2b)
$$\lim_{x \downarrow 0} \frac{\partial g_{\lambda,0}}{\partial y}(x, 0) = \lim_{y \downarrow 0} \frac{\partial g_{\lambda,0}}{\partial y}(0, y) = \frac{1}{\sigma(0)^2} + C,$$

$$C = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\xi \left[-a(0)\xi + \int_{-\infty}^{\infty} \left(\frac{\xi u}{1+u^2} - \xi u \right) n(0, u) du \right]}{[\lambda - \psi_R(0, \xi)]^2 + \psi_I(0, \xi)^2} d\xi + \frac{1}{2\pi} \int_{-\infty}^{f_-} \frac{\xi \left[\int_{-\infty}^{\infty} (\xi u - \sin \xi u) n(0, u) du \right]}{[\lambda - \psi_R(0, \xi)]^2 + \psi_I(0, \xi)^2} d\xi .$$

Proof. By Lemma 5.1, we have

$$\frac{\partial g_{\lambda,0}}{\partial y}(x, 0) = \frac{1}{\pi} \int_{\mathcal{O}}^{\infty} \frac{\xi[\lambda - \psi_R(x, \xi)] \sin x\xi}{[\lambda - \psi_R(x, \xi)]^2 + \psi_I(x, \xi)^2} d\xi + \frac{1}{\pi} \int_{\mathcal{O}}^{\infty} \frac{\xi \psi_I(x, \xi) \cos x\xi}{[\lambda - \psi_R(x, \xi)]^2 + \psi_I(x, \xi)^2} d\xi$$
$$\equiv I_1 + I_2.$$

First we show $\lim_{x \neq 0} I_1 = \frac{1}{\sigma(0)^2}$.

Set

where

HITTING PROPERTIES

$$G(x,\eta) = \left[\lambda x^2 + \frac{\sigma(x)^2}{2}\eta^2 + x^2 \int_{-\infty}^{\infty} \left(1 - \cos\frac{\eta u}{x}\right) n(x,u) du\right]^2$$

and

$$H(x, \eta) = \left[-a(x)x\eta + x^2\int_{-\infty}^{\infty}\left\{\frac{\eta u}{x(1+u^2)} - \sin\frac{\eta u}{x}\right\}n(x, u)du\right]^2.$$

Then we have

$$I_{1} = \frac{1}{\pi} \int_{0}^{\infty} \frac{\eta \sin \eta \left[\lambda x^{2} + \frac{\sigma(x)^{2}}{2} \eta^{2} + x^{2} \int_{-\infty}^{\infty} \left(1 - \cos \frac{\eta u}{x} \right) n(x, u) du \right] d\eta}{G(X, \eta) + H(x, \eta)}.$$

Since

$$\Big|\frac{\partial}{\partial\eta}\Big\{\frac{\eta\Big[\lambda x^2+\frac{\sigma(x)^2}{2}\eta^2+x^2\int_{-\infty}^{\infty}\Big(1-\cos\frac{\eta u}{x}\Big)n(x,u)du\Big]}{G(x,\eta)+H(x,\eta)}\Big\}\Big|\leq \frac{\gamma}{\eta^2} \qquad |\eta|\geq M,$$

by the same argument as in the proof of Lemma 5.1, we can write

$$\frac{\eta \left[\lambda x^2 + \frac{\sigma(x)^2}{2}\eta^2 + x^2 \int_{-\infty}^{\infty} \left(1 - \cos \frac{\eta u}{x}\right) n(x, u) du\right]}{G(x, \eta) + H(x, \eta)} = F_1(\eta) - F_2(\eta)$$

 $(F_1, F_2$ have the same property as F_3, F_4 in the proof of Lemma 5.1).

Using the second mean value theorem, for any $\varepsilon > 0$, there exists a constant N > 0 such that

$$\Big|\frac{1}{\pi}\int_{N}^{\infty}\frac{\eta\sin\eta\Big[\lambda x^{2}+\frac{1}{2}\sigma(x)^{2}\eta^{2}+x^{2}\int_{-\infty}^{\infty}\Big(1-\cos\frac{\eta u}{x}\Big)n(x,u)du\Big]d\eta}{G(x,\eta)+H(x,\eta)}\Big|<\varepsilon$$

for any x. Since the integrand is bounded in [0, N],

$$\lim_{x \neq 0} \frac{1}{\pi} \int_0^N \frac{\eta \sin \eta \left[\lambda x^2 + \frac{1}{2} \sigma(x)^2 \eta^2 + x^2 \int_{-\infty}^\infty \left(1 - \cos \frac{\eta u}{x} \right) n(x, u) du \right] d\eta}{G(x, \eta) + H(x, \eta)}$$
$$= \frac{1}{\pi} \int_0^N \frac{2}{\sigma(0)^2} \frac{\sin \eta}{\eta} d\eta.$$

Consequently we obtain $\lim_{x \neq 0} I_1 = \frac{1}{\sigma(0)^2}$. In case x < 0, we have

$$I_{1} = -\frac{1}{\pi}\int_{0}^{\infty}\frac{\eta\sin\eta\left[\lambda x^{2}+\frac{1}{2}\sigma(x)^{2}\eta^{2}+x^{2}\int_{-\infty}^{\infty}\left(1-\cos\frac{\eta u}{x}\right)n(x, u)du\right]d\eta}{G(x, \eta)+H(x, \eta)},$$

so $\lim_{x \neq 0} I = \frac{1}{\sigma(0)^2}$. For I_2 , we devide I_2 into two terms $I_2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\xi \left[-a(x)\xi + \int_{-\infty}^{\infty} \left(\frac{\xi u}{1+u^2} - \xi u \right) n(x, u) du \right] \cos x\xi}{[\lambda - \psi_R(x, \xi)]^2 + \psi_I(x, \xi)^2} d\xi$ $+ \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [u \sin u) n(x, u u] \cos x\xi}{[\lambda - \psi_R(x, \xi)]^2 + \psi_I(x, \xi)^2} - d\xi.$

Using Lebesgue's convergence theorem,

$$\lim_{x \to 0} f_{2} = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\xi \left[-a(0)\xi + \int_{-\infty}^{\infty} \left(\frac{\xi u}{1+u^{2}} - \xi u \right) n(0, u) du \right]}{[\lambda - \psi_{R}(0, \xi)]^{2} + \psi_{I}(0, \xi)^{2}} d\xi + \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\xi \left[\int_{-\infty}^{\infty} (\xi u - \sin \xi u) n(0, u) du \right]}{[\lambda - \psi_{R}(0, \xi)]^{2} + \psi_{I}(0, \xi)^{2}} d\xi = C.$$

The same estimate shows that

$$\lim_{y \neq 0} \frac{\partial g_{\lambda,0}(0, y)}{\partial y} = -\frac{1}{\sigma(0)^2} + C, \lim_{n \to 0} \frac{\partial g_{\lambda,0}(0, v)}{\partial y} = \frac{1}{\sigma(0)^2} + C.$$

The proof of Lemma 5.2 is completed.

Next we shall consider the case $\sigma(x) = 0$. We shall devide into the following three cases:

(I) $1 < \alpha_2 \le \alpha_1 < 2$, (II) $0 < \alpha_2 < 1 < \alpha_1 < 2$ and (III) $1 = \alpha_2 < \alpha_1 < 2$. We write ψ in the following form according to the above cases.

Case (I).
$$\psi(x,\xi) = ic_1(x)\xi + \prod_{j \to oo} (e^{i\xi u} - 1 - i\xi u)n(x,u)du$$

where $c_1(x) = a(x) + \int_{-\infty}^{\infty} \frac{u^3 n(x, u)}{1 + u^2} du$.

Case (II).
$$\psi(x, \xi) = ic_2(x)\xi + \int_{-\infty}^{0} (e^{i\xi u} - 1) \frac{\nu(x, u)}{|u|^{1+\alpha_2}} du$$

 $+ \int_{0}^{\infty} (e^{i\xi u} - 1 - i\xi u) \frac{\nu(x, u)}{u^{1+\alpha_1}} du$

where $c_2(x) = a(x) + \int_{-\infty}^{0} \frac{1}{1+u^2} \frac{1}{|u|^{-2}} + \int_{0}^{\infty} \frac{1}{1+u^2} \nu(x, u) du$

Case (III).
$$\psi(x,\xi) = ic_3(x)\xi + \int_{-\infty}^0 \left(e^{i\xi u} - 1 - \frac{i\xi u}{1+u^2}\right) \frac{\nu(x,u)}{|u|^2} du$$

$$+\int_0^\infty (e^{i\xi u}-1-i\xi u)\frac{\nu(x,u)}{u^{1+\omega_1}}du,$$

where $c_{3}(x) = a(x) + \int_{0}^{\infty} \frac{u^{3}n(x, u)}{1+u^{2}} du$.

Set
$$a_1(x,\xi) = \int_0^\infty (1-\cos v) \frac{\nu(x,v/\xi)}{v^{1+\sigma_1}} dv, \ a_2(x,\xi) = \int_{-\infty}^0 (1-\cos v) \frac{\nu(x,v/\xi)}{v^{1+\sigma_2}} dv,$$

 $b_1(x,\xi) = \int_0^\infty (\sin v - v) \frac{\nu(x,v/\xi)}{v^{1+\sigma_1}} dv$

and

$$b_{2}(x,\xi) = \begin{cases} \int_{-\infty}^{0} (\sin v - v) \frac{\nu(x, v/\xi)}{|v|^{1+\omega_{2}}} dv, & \text{if } \alpha_{2} > 1, \\ \int_{-\infty}^{0} \sin v \frac{\nu(xv/\xi)}{v^{1+\omega_{2}}} dv, & \text{if } \alpha_{2} < 1, \\ \int_{-\infty}^{0} (\sin v - \frac{v}{1+v^{2}/\xi^{2}}) \frac{\nu(x, v/\xi)}{v^{2}} dv \frac{1}{\log \xi}, & \text{if } \alpha_{2} = 1, \end{cases}$$

Then $\psi(x, \xi)$ can be written in the form

Case (I). $(1 < \alpha_2 \le \alpha_1 < 2)$.

(5.3)
$$\psi(x,\xi) = -(a_1(x,\xi)\xi^{\omega_1} + a_2(x,\xi)\xi^{\omega_2}) + i(b_1(x,\xi)\xi^{\omega_1} + b_2(x,\xi)\xi^{\omega_2} + c_1(x)\xi).$$

Case (II). $(0 < \alpha_2 < 1 < \alpha_1 < 2)$.

(5.4)
$$\psi(x,\xi) = -(a_1(x,\xi)\xi^{\omega_1} + a_2(x,\xi)\xi^{\omega_2}) + i(b_1(x,\xi)\xi^{\omega_1} + b_2(x,\xi)\xi^{\omega_2} + c_2(x)\xi).$$

Case (III). $(1 = \alpha_2 < \alpha_1 < 2)$.

(5.5)
$$\psi(x,\xi) = -(a_1(x,\xi)\xi^{\omega_1} + a_2(x,\xi)\xi^{\omega_2}) + i(b_1(x,\xi)\xi^{\omega_1} + b_2(x,\xi)\xi\log\xi + c_3(x)\xi).$$

Set

$$egin{aligned} &a_1(x) =
u(x,0) \int_0^\infty rac{1 - \cos v}{v^{1 + lpha_1}} dv, \ a_2(x) =
u(x,\mathbf{O}) \int_{-\infty}^0 rac{1 - \cos v}{|v|^{1 + lpha_2}} dv \ , \ &b_1(x) =
u(x,0) \int_0^\infty rac{\sin v - v}{v^{1 + lpha_1}} dv \end{aligned}$$

and

S. MANABE

$$b_2(x) = egin{cases}
u(x, 0) \int_{-\infty}^0 rac{\sin v - v}{|v|^{1+ arphi_2}} dv, & ext{if} \quad lpha_2 > 1\,, \
u(x, 0) \int_{-\infty}^0 rac{\sin v}{|v|^{1+ arphi_2}} dv, & ext{if} \quad lpha_2 < 1\,, \
u(x, 0), & ext{if} \quad lpha_2 = 1 \end{cases}$$

Lemma 5.3. Let $a_i(x,\xi)$, $a_i(x)$, $b_i(x,\xi)$ and $b_i(x)$, i=1,2 be as above. Then we have the following estimates

(56)
$$a_1(x,\xi) - a_1(x) = O(\xi^{-1}),$$

(5.7)
$$a_{2}(x,\xi)-a_{2}(x) = \begin{cases} O(\xi^{-1}), & \text{if } \alpha_{2} > 1, \\ O(\xi^{-\alpha_{2}}), & \text{if } \alpha_{2} < 1, \\ O(\xi^{-1}\log\xi), & \text{if } \alpha_{2} = 1, \end{cases}$$

(5.8)
$$b_1(x,\xi)-b_1(x) = O(\xi^{1-\alpha_1})$$

and

(5.9)
$$b_{2}(x, \xi)-b_{2}(x) = \begin{cases} O(\xi^{1-\omega_{2}}), & \text{if } \alpha_{2} > 1, \\ O(\xi^{-\omega_{2}}), & \text{if } \alpha_{2} < 1, \\ O((\log \xi)^{-1}), & \text{if } \alpha_{2} = 1, \end{cases}$$

uniformly in x, as $\xi \rightarrow \infty$.

Proof. In the following, we use the notation $\frac{\partial}{\partial y}(x, u/\xi)$ for $\frac{\partial}{\partial y}(x, y)\Big|_{y=u/\xi}$. Throughout the proof we assume $\xi > 1$. For $a_1(x, \xi) - a_1(x)$, we have

$$\begin{aligned} |a_{1}(x,\xi)-a_{1}(x)| &\leq \left|\int_{0}^{1} \frac{1-\cos u}{u^{1+a_{1}}}(\nu(x,u/\xi)-\nu(x,0))du\right| \\ &+\left|\int_{1}^{\infty} \frac{1-\cos u}{u^{1+a_{1}}}(\nu(x,u/\xi)-\nu(x,0))du\right| \\ &\leq M_{1}\xi^{-1}\left\{\int_{0}^{1} \frac{1-\cos u}{u^{a_{1}}}du+\int_{1}^{\infty} \frac{du}{u^{a_{1}}}\right\} &\leq M_{1}\xi^{-1}\left(\frac{1}{2(3-\alpha_{1})}+\frac{1}{\alpha_{1}-1}\right),\end{aligned}$$

where M_1 is the same constant as in the proof of Lemma 4.1. Hence we have (5.6). For $b_1(x, \xi) - b_1(x)$, we have

$$\begin{aligned} |b_1(x,\xi)-b_1(x)| &\leq \left| \int_0^1 \frac{\sin v - v}{v^{1+\alpha_1}} (\nu(x,v|\xi) - \nu(x,0)) dv \right| \\ &+ \left| \int_1^{\xi L_1} \frac{\sin v - v}{v^{1+\alpha_1}} (\nu(x,v|\xi) - \nu(x,0)) dv \right| \\ &+ \left| \int_{\xi L_1}^\infty \frac{\sin v - v}{v^{1+\alpha_1}} \nu(x,0) dv \right| \end{aligned}$$

HITTING PROPERTIES

$$\leq M_{1} \Big\{ \xi^{-1} \int_{0}^{1} \frac{v - \sin v}{v^{\alpha_{1}}} dv + \xi^{1 - \alpha_{1}} \int_{0}^{L_{1}} \frac{dv}{v^{\alpha_{1} - 1}} \Big\} + M_{1} \int_{\xi L_{1}}^{\infty} \frac{1 + v}{v^{1 + \alpha_{1}}} dv \\ \leq M_{1} \Big\{ \frac{1}{2(3 - \alpha_{1})\xi} + \frac{L_{1}^{2 - \alpha_{1}}}{2 - \alpha_{1}} \frac{1}{\xi^{\alpha_{1} - 1}} + \frac{2}{L_{1}^{\alpha_{1} - 1}(\alpha_{1} - 2)} \frac{1}{\xi^{\alpha_{1} - 1}} \Big\} .$$

Thus we have (5.8). By the same way, we have (5.7) for $\alpha_2 > 1$. If $\alpha_2 < 1$, then we have

$$\begin{aligned} a_2(x,\xi) - a_2(x) &= \int_{-1}^0 \frac{1 - \cos u}{|u|^{1+\sigma_2}} [\nu(x,u|\xi) - \nu(x,0)] du \\ &+ \int_{-\xi L_1}^{-1} \frac{1 - \cos u}{|u|^{1+\sigma_2}} [\nu(x,u|\xi) - \nu(x,0)] du - \int_{-\infty}^{-\xi L_1} \frac{1 - \cos u}{|u|^{1+\sigma_2}} \nu(x,0) du \\ &= I_1 + I_2 + I_3 . \end{aligned}$$

For I_1 , we have

$$|I_1| = \left| \int_{-1}^{0} \frac{1 - \cos u}{|u|^{\alpha_2}} \frac{1}{\xi} \frac{\partial \nu}{\partial y}(x, u/\xi) du \right| = M_1/\xi \int_{-1}^{0} \frac{1 - \cos u}{|u|^{\alpha_2}} du = \frac{M_1}{2(3 - \alpha_2)} \frac{1}{\xi}.$$

For I_2 ,

$$\begin{split} |I_{2}| &= \left| \frac{1}{\xi^{\alpha_{2}}} \int_{-L_{1}}^{-1/\xi} \frac{1 - \cos \xi v}{|v|^{1+\alpha_{2}}} \left[\cdots (\infty - v) - \cdots (x, 0) \right] dv \right| \\ &= \left| \frac{1}{\xi^{\alpha_{2}}} \int_{-L_{1}}^{-1/\xi} \frac{1 - \cos \xi v}{|v|^{\alpha_{2}}} \frac{\partial v}{\partial y} (\cdots, v') dv \right| \leq \frac{2M_{1}}{\xi^{\alpha_{2}}} \int_{-L_{1}}^{0} \frac{dv}{|v|^{\alpha_{2}}} = \frac{2M_{1}L_{1}^{1-\alpha_{2}}}{\alpha_{2} - 1} \frac{1}{\xi^{\alpha_{2}}}, \end{split}$$

where v' is a mean value such that v < v' < 0.

For I_3 , we get

$$|I_3| \leq \frac{2M_0}{\alpha_2 L_1^{\alpha_2}} \frac{1}{\xi^{\alpha_2}}.$$

Therefore $| a_2(x, \xi) - a_2(x) | \leq c_5 \frac{1}{\xi^{a_2}},$ where c_5 depends only on L_1, α_2 and $M_i, i=1, 2$.

In case $\alpha_2 = 1$,

$$|a_{2}(x,\xi)-a_{2}(x)| = \left|\int_{0}^{\xi L_{1}} \frac{1-\cos y}{y^{2}} \nu(x,-y/\xi) dy - \nu(x,0) \int_{0}^{\infty} \frac{1-\cos y}{y^{2}} dy\right|$$
$$= \left|\int_{0}^{\xi L_{1}} \frac{1-\cos y}{y^{2}} \{\nu(x,-y/\xi) - \nu(x,0)\} dy - \nu(x,0) \int_{\xi L_{1}}^{\infty} \frac{1-\cos y}{y^{2}} dy\right|$$

S. MANABE

$$\leq \int_{0}^{1} \frac{1 - \cos y}{y^{2}} |v(x, -y|\xi) - v(x, 0)| \, dy + \int_{1}^{\xi L_{1}} \frac{1 - \cos y}{y^{2}} |v(x, -y|\xi) - v(x, 0)| \, dy$$

+ $v(x, 0) \int_{\xi L_{1}}^{\infty} \frac{1 - \cos y}{y^{2}} \, dy$
 $\equiv I_{1} + I_{2} + I_{3}.$

For I_1 ,

$$I_1 \leq M_1 / \xi \int_0^1 \frac{1 - \cos y}{y^2} dy \leq M_1 / 2\xi$$
.

For I_2 ,

$$I_{2} \leq \frac{M_{1} \log L_{1} \xi}{\xi} \frac{\xi^{\xi L_{1}} 1 - \cos y}{J_{1} y \log \xi L_{1}} \leq \frac{2M_{1} \log \xi L_{1}}{\xi}$$

For I_3 , we have

$$I_3 \leq 2M_0/L_1 \xi$$
.

Therefore

(5.10)
$$|a_2(x,\xi)-a_2(x)| \leq c_6 \frac{\log \xi}{\xi},$$

where c_6 depends only on L_1 , M_0 and M_1 . Thus (5.7) is proved. Finally for $b_2(x,\xi)-b_2(x)$, if $\alpha_2 > 1$ then the same estimate as that of $b_1(x,\xi)-b_1(x)$ holds. If $\alpha_2 < 1$, then we have

$$b_{2}(x,\xi)-b_{2}(x) = \int_{-1}^{0} \frac{\sin v}{|v|^{1+\alpha_{2}}} [\nu(x,v|\xi)-\nu(x,0)]dv$$

+
$$\int_{-\xi L_{1}}^{-1} \frac{\sin v}{|v|^{1+\alpha_{2}}} [\nu(x,v|\xi)-\nu(x,0)]dv$$

+
$$\int_{-\infty}^{-L_{1}\xi} \frac{\sin v}{|v|^{1+\alpha_{2}}} [\nu(x,v|\xi)-\nu(x,0)]dv$$

$$\equiv I_{1}+I_{2}+I_{3}.$$

For I_1 , we have

$$|I_1| = \frac{M_1}{\xi} \int_{-1}^0 \frac{dv}{|v|^{a_2}} = \frac{M_1}{1-\alpha_2} \frac{1}{\xi}.$$

For I_2 ,

$$|I_2| = \left| \frac{1}{\xi^{a_2}} \int_{-L_1}^{-1/\xi} \frac{\sin \xi w}{|w|^{1+a_2}} [\nu(x,w) - \nu(x,0)] dw \right| \le \frac{M_1}{\xi^{a_2}} \int_{-L_1}^{0} \frac{dw}{|w|^{a_2}} \le \frac{M_1 L_1^{1-a_2}}{1-\alpha_2} \frac{1}{\xi^{a_2}}.$$

For I_3 ,

$$|I_3| \leq 2M_0 \int_{-\infty}^{-\xi L_1} \frac{dv}{|v|^{1+\alpha_2}} - \frac{2M_0}{\alpha_2 L_1^{\alpha_2}} \frac{1}{\xi^{\alpha_2}}.$$

Thus we obtain

(5.11)
$$|b_2(x,\xi)-b_2(x)| \leq c_7 \xi^{-\alpha_2},$$

where c_7 does not depend on $x, \xi \in \mathbb{R}^1$.

Finally consider the case $\alpha_2 = 1$.

$$b_{2}(x, \xi) - b_{2}(x) = \frac{1}{\log \xi} \left\{ \int_{0}^{1} \frac{y - \sin y}{y^{2}} \nu(x, -y/\xi) dy + \int_{0}^{1} \left(\frac{y}{1 + y^{2}/\xi^{2}} - y \right) \frac{\nu(x, -y/\xi)}{y^{2}} dy - \int_{1}^{\xi L_{1}} \frac{\sin y}{y^{2}} \nu(x, -y/\xi) dy \right\} + \left\{ \frac{1}{\log \xi} \int_{1}^{\xi L_{1}} \frac{\nu(x, -y/\xi)}{y(1 + y^{2}/\xi^{2})} dy - \nu(x, 0) \right\} \equiv I_{1} + I_{2}.$$

For I_1 , we obtain

$$|I_1| \leq \frac{1}{\log \xi} \left(\frac{M_0}{6} + \frac{M_0}{2\xi^2} + M_0 \right).$$

For I_2 , we get

$$\begin{split} |I_{2}| &= \left| \frac{1}{\log \xi L_{1}} \int_{1}^{\xi L_{1}} \left\{ \frac{\nu(x, -y/\xi)}{y(1+y^{2}/\xi^{2})} - \frac{\nu(x, 0)}{y} \right\}_{J}^{J_{0}} \times \frac{\log \xi L_{1}}{\log \xi} \right| \\ &\leq \frac{1}{\log \xi L_{1}} \int_{1}^{\xi L_{1}} \frac{|\nu(x, -y/\xi) - \nu(x, 0)|}{y(1+y/\xi^{2})} \quad \forall n \in \mathbb{N}, 0 \leq \frac{1}{\log \xi L_{1}} \int_{1}^{\xi L_{1}} \frac{ydy}{\xi^{2}+y^{2}} \\ &\leq \frac{M_{1}}{\xi \log \xi L_{1}} \int_{1}^{\xi L_{1}} dy + \frac{M_{0}}{\log \xi L_{1}} \log \frac{\xi^{2}(1+L_{1}^{2})}{\xi^{2}+1} \leq \frac{L_{1}M_{1}+1}{\log \xi L_{1}} \,. \end{split}$$

So we have

(5.12)
$$|b_2(x,\xi)-b_2(x)| \leq c_8(\log \xi)^{-1}$$
,

where c_8 does not depend on x, $\xi \in \mathbb{R}^1$. By (5.11) and (5.12), we obtain (5.9), which completes the proof of Lemma 5.3.

Now we can estimate the singularity of $\int \frac{g_{\lambda,0}}{dy}$. Set

S. Manabe

(5.13)
$$I(x,y) = -\frac{1}{\pi} \int_0^\infty \frac{\xi[\lambda - \psi_R(x,\xi)]\sin(y-x)\xi}{[\lambda - \psi_R(x,\xi)]^2 + \psi_I(x,\xi)^2} d\xi$$

and

(5.14)
$$J(x,y) = \frac{1}{\pi} \int_0^\infty \frac{\xi \psi_I(x,\xi) \cos(y-x)\xi}{[\lambda - \psi_R(x,\xi)]^2 + \psi_I(x,\xi)^2} d\xi.$$

In the rest of this paper, we set

$$\Phi_R(x,\xi) = -(a_1(x)\xi^{\omega_1} + a_2(x)\xi^{\omega_2})$$

and

$$\Phi_{I}(x,\xi) = \begin{pmatrix} (b_{1}(x)\xi^{\alpha_{1}}+b_{2}(x)\xi^{\alpha_{2}}+c_{1}(x)\xi), & \text{if } \alpha_{2} > 1, \\ (b_{1}(x)\xi^{\alpha_{1}}+b_{2}(x)\xi^{\alpha_{2}}+c_{2}(x)\xi), & \text{if } \alpha_{2} < 1, \\ (b_{1}(x)\xi^{\alpha_{1}}+b_{2}(x)\xi\log\xi+c_{3}(x)\xi), & \text{if } \alpha_{2} = 1 \end{pmatrix}$$

Define $h_1(x)$ by

$$h_1(x) = rac{\pi}{2\Gamma(lpha_1+1)[a_1(x)^2+b_1(x)^2]}$$
.

Then we can show the following lemma.

Lemma 5.4. Assume that $\alpha_1 > \alpha_2$ Let *n* be the largest positive integer such that $\alpha_1 - 2 + n(\alpha_1 - \alpha_2) \leq 0$ Then we have (i) $//\alpha_2 \neq 1$, then

(5.15)
$$I(x, y) = -\frac{h_{1}(x)}{\pi} \Gamma(2-\alpha_{1}) \operatorname{sgn}(y-x) | y-x|^{\omega_{1}-2} + \sum_{k=1}^{k} \sum_{j=0}^{k} \left\{ D_{\omega_{2},k+1}^{(j)}(x) \int_{0}^{\infty} \xi^{j(1-\omega_{2})+1-\omega_{1}-k(\omega_{1}-\omega_{2})} \operatorname{sin}\xi d\xi \right\} X \operatorname{sgn}(y-k) y-x | \omega_{1}-2+k(\omega_{1}-\omega_{2})+j(\omega_{2}-1) + v_{5}(x, y) | y-x|^{2\omega_{1}-3} + v_{6}(x, y) ,$$

(5.16)
$$J(x, y) = -\frac{h_{1}(x)}{\pi} \Gamma(2-\alpha_{1}) | y-x|^{\omega_{1}-2} + \sum_{k=1}^{n} \sum_{j=0}^{k} \left\{ E_{\omega_{2},k+1}^{(j)}(x) \int_{0}^{\infty} \xi^{j(1-\omega_{2})+1-\omega_{1}-k(\omega_{1}-\omega_{2})} \cos \xi d\xi \right\} X | y-x|^{\omega_{1}-2+k(\omega_{1}-\omega_{2})+j(\omega_{2}-1)} + v_{7}(x, y) | y-x|^{2\omega_{1}-3} + v_{8}(x, y) ,$$

where $\{D_{a_2,k+1}^{(j)}\}_{0 \le j \le k}$, are bounded continuous functions. (ii) If $tf_2 = l$, then

HITTING PROPERTIES

(5.17)
$$I(x, y) = -\frac{h_{1}(x)}{\pi} \Gamma(2-\alpha_{1}) \operatorname{sgn}(y-x) |y-x|^{\omega_{1}-2} + \sum_{k=1}^{n} \sum_{r=0}^{k} \left\{ \sum_{j=r}^{k} D_{1,k+1}^{(j)}(x) \int_{0}^{\infty} (\log \xi)^{j-r} \xi^{1-\omega_{1}-k(\omega_{1}-\omega_{2})} \sin \xi d\xi \right\} \\ \times \left(\log \frac{1}{|y-x|} \right)^{r} \operatorname{sgn}(y-x) \operatorname{I}_{y-x} \setminus^{\omega_{1}-2+k(-1-\omega_{2})} + v_{5}(x, y) |y-x|^{2\omega_{1}-3} + v_{6}(x, y),$$

(5.18)
$$J(x,y) = -\frac{h_{1}(x)}{\pi} \Gamma(2-\alpha_{1}) | y-x|^{\omega_{1}-2} + \sum_{k=1}^{k} \sum_{r=0}^{k} \left\{ \sum_{j=r}^{k} E_{1,k+1}^{(j)}(x) \int_{0}^{\infty} (\log \xi)^{j-r} \xi^{1-\omega_{1}-k(\omega_{1}-\omega_{2})} \cos \xi d\xi \right\} \times \log \left(\frac{1}{|y-x|} \right)^{r} | y-x|^{\omega_{1}-2+k(\omega_{1}-\omega_{2})} + v_{7}(x,y) | y-x|^{2\omega_{1}-3} + v_{8}(x,y) ,$$

where $\{D_{1,k+1}^{(j)}(x)\}_{0 \leq j \leq k, 1 \leq k \leq n}$, $\{E_{1,k+1}^{(j)}(x)\}_{0 \leq j \leq k, 1 \leq k \leq n}$ and v_j , $5 \leq j \leq 8$ are bounded continuous functions. In case $\alpha_1 - 2 + n(\alpha_1 - \alpha_2) = 0$, $|y - x|^{\alpha_1 - 2 + n(\alpha_1 - \alpha_2)}$ should be replaced by $\log\left(\frac{1}{|y - x|} \lor 1\right)$.

Proof. (i) First we show (5.15). Denote by c(x) for $c_1(x)$ or $c_2(x)$. Then I(x, y) can be written as follows:

$$I(x, y) = -\frac{1}{\pi} \int_{0}^{\infty} \frac{\lambda \xi \sin((y-x)\xi d\xi}{[\lambda + a_{1}(x, \xi)\xi^{\omega_{1}} + a_{2}(x, \xi)\xi^{\omega_{2}}]^{2} + [b_{1}(x, \xi)\xi^{\omega_{1}} + b_{2}(x, \xi)\xi^{\omega_{2}} + c(x)\xi]^{2}} \\ -\frac{1}{\pi} \int_{0}^{\infty} \frac{\sin((y-x)\xi d\xi}{[\lambda + a_{1}(x, \xi)\xi^{\omega_{1}} + a_{2}(x, \xi)\xi^{\omega_{2}}]^{2} + [b_{1}(x, \xi)\xi^{\omega_{1}} + b_{2}(x, \xi)\xi^{\omega_{2}} + c(x)\xi]^{2}}}{\sin((y-x)\xi d\xi}.$$

The first term is a bounded continuous function. Denote by $I_1(x,y)$ the second term. Then we have

$$\begin{split} I_1(x,y) &= -\frac{1}{\pi} \int_0^\infty \frac{[a_1(x)\xi^{1+\pmb{\sigma}_1} + a_2(x)\xi^{1+\pmb{\sigma}_2}]\sin{(y-x)\xi}d\xi}{[\lambda - \Phi_R(x,\,\xi)]^2 + \Phi_I(x,\,\xi)^2} \\ &- \frac{1}{\pi} \int_0^\infty \left\{ \frac{a_1(x,\,\xi)\xi^{1+\pmb{\sigma}_1} + a_2(x,\,\xi)\xi^{1+\pmb{\sigma}_2}}{(\lambda - \psi_R(x,\,\xi))^2 + \psi_I(x,\,\xi)^2} - \frac{a_1(x)\xi^{1+\pmb{\sigma}_1} + a_2(x)\xi^{1+\pmb{\sigma}_2}}{(\lambda - \Phi_R(x,\,\xi))^2 + \Phi_I(x,\,\xi)^2} \right\} \\ &\times \sin{(y-x)\xi}d\xi \\ &\equiv I_2(x,\,y) + I_3(\chi,\,y) \end{split}$$

For the estimate of $I_2(x, y)$, we employ the method of Ikeda-S.Watanabe [3 p. 165]. Set $\tilde{b}_2(x, \xi) = b_2(x) + c(x)\xi^{1-\omega_2}$. Define

S. Manabe

$$\begin{split} A_1(x,\xi) &= a_1(x), B_1(x,\xi) = a_2(x), D_{a_2,k}(x,\xi) = \frac{A_k(x,\xi)}{a_1(x)^2 + o_1(x)^2}, \\ A_{k+1}(x,\xi) &= B_k(x,\xi) - \frac{2(a_1(x)a_2(x) + b_2(x)\tilde{b}_2(x,\xi))}{a_1(x)^2 + b_1(x)^2} A_k(x,\xi) \end{split}$$

and

$$\mathcal{B}_{k+1}(x,\,\xi) = \frac{\gamma_2(x)^2 + \tilde{b}_2(x,\,\xi)^2}{(u_1(x))^2 + b_1(x)^2} A_k(x,\,\xi), \, 1 \le k \le n+1 \, .$$

Then we have the following formula:

(519)
$$\int_{0}^{\infty} \sin(y \ x) \xi^{*} \frac{A_{k}(x, \xi) \xi^{1+\omega_{1}-(k-1)(\omega_{1}-\omega_{2})} + B_{k}(x, \xi) \xi^{1+\omega_{1}-k(\omega_{1}-\omega_{2})}}{(\lambda+a_{1}(x)\xi^{\omega_{1}}+a_{2}(x)\xi^{\omega_{2}})^{z} + (b_{1}(x)\xi^{\omega_{1}}+b_{2}(x)\xi^{\omega_{2}}+c(x)\xi)^{z}} d\xi$$
$$= \int_{0}^{\infty} D_{\omega_{2},k}(x, \xi) \sin(y-x)\xi \cdot \xi^{1-\omega_{1}-(k-1)(\omega_{1}-\omega_{2})} d\xi$$
$$+ \int_{0}^{\infty} \sin(y-x)\xi \frac{A_{k+1}(x, \xi)\xi^{1+\omega_{1}-k(\omega_{1}-\omega_{2})} + B_{k+1}(x, \xi)\xi^{1+\omega_{1}-(k+1)(\omega_{1}-\omega_{2})}}{(\lambda+a_{1}(x)\xi^{\omega_{1}}+a_{2}(x)\xi^{\omega_{2}})^{2} + (b_{1}(x)\xi^{\omega_{1}}+b_{2}(x)\xi^{\omega_{2}}+c(x)\xi)^{z}} d\xi$$
$$+ e_{k}(x, y),$$

where $e_k(x, y)$ is a bounded continuous function. From this formula, it is easy to see that $D_{\alpha_2,k}(x, \xi)$ is a polynomial of degree k in $\xi^{1-\alpha_2}$. Therefore $D_{\alpha_2,k}(x,\xi)$ can be written in the form

$$D_{\boldsymbol{\sigma}_{2,1}}(x, \boldsymbol{\xi}) - \frac{a_1(x)}{a_1(x)^2 + b_1(x)^2},$$

$$D_{\boldsymbol{\sigma}_{2,k+1}}(x, \boldsymbol{\xi}) = D_{\boldsymbol{\sigma}_{2,k+1}}^{(0)}(x)(\boldsymbol{\xi}^{1-\boldsymbol{\sigma}_2})^k + \dots + D_{\boldsymbol{\sigma}_{2,k+1}}^{(k)}(x), \qquad 1 \leq k \leq n,$$

where the coefficients $D_{a_2,k+1}^{(j)}(x)$, $0 \le j \le k$, are bounded continuous functions. Hence we have

where v_8 is a bounded continuous function. Noting that $D_{\alpha_{2,1}}^{(1)}(x) \equiv 0$ and $D_{\alpha_{2,1}}^{(0)}(x) = \frac{a_1(x)}{a_1(x)^2 + b_1(x)^2}$, we obtain

$$I_{2}(x, y) = -\frac{h_{1}(x)}{\pi} \Gamma(2-\alpha_{1}) \operatorname{sgn}(y-x) |y-x|^{\omega_{1}-2} + \sum_{k=2}^{n+1} \sum_{j=0}^{k} D_{\omega_{2},k}^{(j)}(x) |y-x|^{\omega_{1}-2+(k-1)(\omega_{1}-\omega_{2})+j(\omega_{2}-1)}$$

HITTING PROPERTIES

$$X \operatorname{sgn}(y-x) \int_{J_0}^{\infty} \xi^{1-\frac{1}{2}-1} (k^{-1}) (1-2)^{j+j(1-2)} \sin \xi d\xi + v_{g}(x,y) .$$

For $I_{s}(x, y)$, we have

(5.20)
$$I_{3}(x, y) = -\frac{1}{\pi} \int_{0}^{\infty} (a_{1}(x, \xi) - a_{1}(x))\xi^{1+a_{1}} + (a_{2}(x, \xi) - a_{2}(x))\xi^{1+a_{2}}}{(\lambda - \psi_{R}(x, \xi))^{2} + \psi_{I}(x, \xi)^{2}} \dot{\varsigma} \dot{\eta}_{2}(x, -x)\xi d\xi$$
$$+ \frac{1}{\pi} \int_{0}^{\infty} \left\{ \frac{1}{(\lambda - \psi_{R}(x, \xi))^{2} + \psi_{I}(x, \xi)^{2}} - \frac{1}{(\lambda - \Phi_{R}(x, \xi))^{2} + \Phi_{I}(x, \xi)^{2}} \right\}$$
$$\times \xi \Phi_{R}(x, \xi) \sin(y - x)\xi d\xi$$
$$\equiv I_{4}(x, y) + I_{5}(x, y).$$

Using Lemma 5.3, it is easy to see that $I_4(x, y)$ is a bounded continuous function. For I_5 , set

$$H_{1}(x,\xi) = (a_{1}(x,\xi)-a_{1}(x))\xi^{\bullet_{1}}+(a_{2}(x,\xi)-a_{2}(x))\xi^{\bullet_{2}},$$
(5.21) flX*. ξ = $2\lambda - (a_{1}(x,\xi)+a_{1}(x))\xi^{\bullet_{1}}-(a_{2}(x,\xi)+a_{2}(x))\xi^{\bullet_{2}},$
 $K_{1}(x,\xi) = (b_{1}(x,\xi)-b_{1}(x))\xi^{\bullet_{1}}+(b_{2}(x,\xi)-b_{2}(x))\xi^{\bullet_{2}},$
 $K_{2}(x,\xi) = (b_{1}(x,\xi)+b_{1}(x))\xi^{\bullet_{1}}+(b_{2}(x,\xi)+b_{2}(x))\xi^{\bullet_{2}}+2c(x)\xi$

and

$$\Xi(x, y; \xi) = \frac{\xi \Phi_R(x, \xi) \sin(y-x)\xi}{\{(\lambda - \psi_R(x, \xi))^2 + \psi_I(\chi, \xi)^2\} \{(\lambda - \Phi_R(x, \xi))^2 + \Phi_I(x, \xi)^2\}}.$$

Then we have

$$I_{\mathfrak{s}}(x, y) = \frac{1}{\pi} \int_{0}^{\infty} \frac{H_{\mathfrak{s}}(x, \xi)H_{\mathfrak{s}}(x, \xi)}{\Xi(x, y; \xi)} d\xi + \frac{1}{\pi} \int_{0}^{\infty} \frac{K_{\mathfrak{s}}(x, \xi)K_{\mathfrak{s}}(x, \xi)}{\Xi(x, y; \xi)} d\xi$$
$$\equiv I_{\mathfrak{s}} + I_{\mathfrak{r}}.$$

By Lemma 5.3, it is easy to see that I_6 is bounded continuous. For the estimate of I_7 , set

$$a_{1}\left(x,\frac{\eta}{y-x}\right) - a_{1}(x) = \bar{a}_{1}(x,y;\eta) \frac{|y-x|}{\eta},$$

$$b_{1}\left(x,\frac{\eta}{y-x}\right) - b_{1}(x) = \bar{b}_{1}(x,y;\eta) \frac{|y-x|^{\alpha_{1}-1}}{\eta^{\alpha_{1}-1}},$$

$$a_{2}\left(x,\frac{\eta}{y-x}\right) - a_{2}(x) = \begin{cases} a_{2}(x,y;\eta) \frac{|y-x|}{\eta}, & \text{if } \alpha_{2} > 1, \\ a_{2}(x,y;\eta) \frac{|y-x|}{\eta^{\alpha_{2}}}, & \text{if } \alpha_{2} < 1, \end{cases}$$

and

S. Manabe

$$b_{2}\left(x,\frac{\eta}{y-x}\right)-b_{2}(x) = \begin{cases} \bar{b}_{2}(x, y; \eta)\frac{|y-x|^{\omega_{2}-1}}{\eta^{\omega_{2}-1}}, & \text{if } \alpha_{2} > 1, \\ \bar{b}_{2}(x, y; \eta)\frac{|y-x|^{\omega_{2}}}{\eta^{\omega_{2}}}, & \text{if } \alpha_{2} < 1. \end{cases}$$

Then we have

$$I_{\tau}(x, y) = -\frac{\operatorname{sgn}(y-x)}{\pi} |y-x|^{2\mathfrak{\omega}_{1}-3} \int_{0}^{\infty} R(x, y \quad \eta) \sin \eta d\eta ,$$

where

$$\begin{split} R\left(x, y; \eta\right) &= \frac{R_{1}(x, y; \eta)R_{2}(x, y, \eta)R_{3}(x, y, \eta)}{(S_{1}(x, y; \eta)^{2} + S_{2}(x, y; \eta)^{2})(T_{1}(x, y, \eta)^{2} + T_{2}(x, y; \eta)^{2})},\\ R_{1}(x, y; \eta) &= a_{1}(x)\eta^{1+\alpha_{1}} + a_{2}(x)\eta^{1+\alpha_{2}}(y-x)^{\alpha_{1}-\alpha_{2}},\\ R_{2}(x, y; \eta) &= \overline{b}_{1}(x, y; \eta)\eta + \overline{b}_{2}(x, y; \eta)\eta^{\beta}(y-x)^{\gamma},\\ R_{3}(x, y; \eta) &= \left(\frac{bx}{2}, \frac{\eta}{y-x}\right) + b_{1}(x)\right)\eta^{\alpha_{1}} + \left(b_{2}\left(x, \frac{\eta}{y-x}\right) + b_{2}(x)\right)\eta^{\alpha_{2}}(y-x)^{\alpha_{1}-\alpha_{2}} \\ &\quad + 2c(x)\eta(y-x)^{\alpha_{1}-1},\\ S_{1}(x, y; \eta) &= \lambda(y-x^{\alpha_{1}} + a_{1}\left(x, \frac{\eta}{y-x}\right)\eta^{\alpha_{1}} + a_{2}\left(x, \frac{\eta}{y-x}\right)\eta^{\alpha_{2}}(y-x)^{\alpha_{1}-\alpha_{2}},\\ S_{2}(x, y; \eta) &= b_{1}\left(x, \frac{\eta}{y-x}\right)\eta^{\alpha_{1}} + b_{2}\left(x, \frac{\eta}{y-x}\right)\eta^{\alpha_{2}}(y-x)^{\alpha_{1}-\alpha_{2}} + c(x)\eta(y-x)^{\alpha_{1}-1},\\ T_{1}(x, y; \eta) &= \lambda(y-x^{\alpha_{1}} + a_{1}(x)\eta^{\alpha_{1}} + a_{2}(x)\eta^{\alpha_{2}}(x-x)^{\alpha_{1}-\alpha_{2}}) \end{split}$$

and

$$T_{2}(x, y; \eta) = b_{1}(x)\eta^{\omega_{1}} + b_{2}(x)\eta^{\omega_{2}}(y-x)^{\omega_{1}-\omega_{2}} + c(x)\eta(y-x)^{\omega_{1}-1}.$$

In the above expressions, we set $\beta = \alpha_1 - 1 + (\alpha_1 - \alpha_2)$ if $\alpha_2 > 1$, $= 2(\alpha_1 - 1)$ if $\alpha_2 < 1$ and $\gamma = \alpha_2 \lor 1$. Applying the second mean value theorem, we obtain the integral $\int_0^\infty R(x, y; \eta) d\eta$ is uniformly convergent in x, y. Hence $\int_0^\infty R(x, y; \eta) d\eta$ is bounded continuous. Thus we have (5.15). By the same argument as that of I(x,y), we obtain (5.16).

Next we show (ii). I(x, y) can be written as follows:

$$I(x, y) = -\frac{1}{\pi} \int_0^\infty \frac{(a_1(x, \xi)\xi^{1+\alpha_1} + a_2(x, \xi)\xi^{1+\alpha_2})}{(\lambda - \psi_R(x, \xi))^2 + \psi_I(x, \xi)^2} \sin(y-x)\xi d\xi + v_{\theta}(x, y),$$

where v_{g} is bounded continuous. Denote by $I_{1}(x, y)$ the first term of the right hand side of the above expression. Then we get

$$I_1(x, y) = -rac{1}{\pi} \int_0^\infty rac{(a_1(x) \xi^{1+lpha_1} + a_2(x) \xi^{1+lpha_2}) \sin{(y-x)} \xi}{(\lambda - \Phi_R(x, \, \xi))^2 + \Phi_I(x, \, \xi)^2} d\xi$$

$$-\frac{1}{\pi}\int_{0}^{\infty} \int a_{1}(x,\xi)\xi^{1+\alpha_{1}}+a_{2}(x,\xi)\xi^{1+\alpha_{2}}-a_{1}(x)\xi^{1+\alpha_{1}}+a_{2}(x)\xi^{1+\alpha_{2}}-\frac{1}{\pi}\int_{0}^{\infty} \int ((\lambda-\psi_{R}(x,\xi))^{2}+\psi_{I}(x,\xi)^{2}(\lambda-\Phi_{R}(x,\xi))^{2}+\Phi_{I}(x,\xi)^{2})$$

$$\times \sin(y-x)\xi d\xi$$

$$\equiv I_{2}(x,y)+I_{3}(x,y).$$

For the estimate of $I_2(x, y)$, we set $\tilde{b}_2(x, \xi) = b_2(x)\log \xi + c(x)$, $A(x, \xi) = a_1(x)$, $B_1(x, \xi) = a_2(x)$, $D_{1,k}(x, \xi) = \frac{A_k(x, \xi)}{a_1(x)^2 + b_1(x)^2}$, $A_{k+1}(x, \xi) = B_k(x, \xi) - \frac{2(a_1(x)a_2(x) + b_1(x)\tilde{b}_2(x, \xi))}{a_1(x)^2 + b_1(x)^2} A_k(x, \xi)$

and

$$B_{k+1}(x, \xi) = -\frac{a_2(x)^2 + \tilde{b}_2(x,\xi)^2}{a_1(x)^2 + b_1(x)^2} A_k(x,\xi) , \qquad 1 \leq k \leq n+1 .$$

Then by (5.19), we get

$$\int_{0}^{\infty} \sin(y-x)\xi \frac{A_{k}(x,\xi)\xi^{1+\sigma_{1}-(k-1)(\sigma_{1}-\sigma_{2})}+B_{k}(x,\xi)\xi^{1+\sigma_{1}-k(\sigma_{1}-\sigma_{2})}}{(\lambda+a_{1}(x)\xi^{\sigma_{1}}+a_{2}(x)\xi^{\sigma_{2}})^{2}+(b_{1}(x)\xi^{\sigma_{1}}+b_{2}(x)\xi\log\xi+c(x)\xi)^{2}}d\xi$$

$$=\int_{0}^{\infty} D_{1,k}(x,\xi) \sin(y-x)\xi \cdot\xi^{1-\sigma_{1}-(k-1)(\sigma_{1}-\sigma_{2})}d\xi$$

$$+\int_{0}^{\infty} \sin((y-x)\xi \frac{A_{k+1}(x,\xi)\xi^{1+\sigma_{1}-k(\sigma_{1}-\sigma_{2})}+B_{k+1}(x,\xi)\xi^{1+\sigma_{1}-(k+1)(\sigma_{1}-\sigma_{2})}}{(\lambda+a_{1}(x)\xi^{\sigma_{1}}+a_{2}(x)\xi^{\sigma_{2}})^{2}+(b_{1}(x)\xi^{\sigma_{1}}+b_{2}(x)\xi_{0g}\xi+c(x)\xi)^{2}}d\xi$$

$$+e_{k}(x,y),$$

where e_k is bounded continuous. From this formula, it is clear that $D_{1,k}(x,\xi)$ is a polynomial of degree k in log ξ . Hence $D_{1,k}(x,\xi)$ can be written in the form

$$\begin{split} D_{1,1}(x,\,\xi) &= \frac{a_1(x)}{a_1(x)^2 + \tilde{b}_1(x)^2}, \\ D_{1,k+1}(x,\,\xi) &= D_{1,k+1}^{(0)}(x)(\log\xi)^k + \dots + D_{1,k+1}^{(k)}(x), \qquad 1 \leq k \leq n \end{split}$$

where the coefficients $D_{1,k+1}^{(j)}$, $0 \le j \le k$ are bounded continuous functions. Therefore we have

$$\begin{split} I_2(x, y) &= \sum_{k=1}^{n+1} \sum_{j=0}^k D_{1,k}^{(j)}(x) \int_0^\infty \xi^{1-\sigma_1 - (k-1)(\sigma_1 - \sigma_2)} (\log \xi)^j \, \dot{\mathrm{str}}_{\mathcal{L}}(x) - x) \xi d\xi + v_8(x, y) \\ &= \sum_{k=1}^{n+1} \sum_{j=0}^k D_{1,k}^{(j)}(x) \, \mathrm{sgn}(y-x) \|y-x\|^{\sigma_1 - 2 + (k-1)(\sigma_1 - \sigma_2)} \left(\log \frac{1}{|y-x|}\right)^r \\ &\quad \times \sum_{r=0}^k \binom{j}{p^r} \int_0^\infty (\log \xi)^{j-r} \, \xi^{1-\sigma_1 - (k-1)(\sigma_1 - \sigma_2)} \mathrm{sin} \, \xi d\xi + v_8(xy) \,, \end{split}$$

where $v_{s}(x,y)$ is a bounded continuous function.

S. MANABE

Noting that
$$D_{1,1}^{(1)}(x) = 0$$
, $D_{1,1}^{(0)}(x) = \frac{\pi}{a_1(x)^r} + \frac{\pi}{a_1(x)^r}$, we obtain
 $I_2(x, y) = -\frac{h_1(x)}{\pi} \Gamma(2 - \alpha_1) \operatorname{sgn}(y - x) |y - x|^{-\sigma_1 - 2}$
 $+ \sum_{n=2} \sum_{r=0}^{\infty} \left\{ \sum_{j=r}^{k} {j \choose r} D_{1,k}^{(j)}(x) \int_{0}^{\infty} (\log \xi)^{j-r} \xi^{1-\sigma_1 - (k-1)(1-\sigma_2)} \operatorname{Sin} \xi d\xi \right\}$
 $\times (\log \frac{1}{\sqrt{1-\alpha_1}})^r \operatorname{sgn}(y - x) |y - x|^{-\sigma_1 - 2 + (k-1)(\sigma_1 - \sigma_2)} + v_8(x, y).$

For I_3 , we devide $I_3(x, y)$ into two terms: $I_3(x, y) = I_4(x, y) + I_5(x, y)$, where I_4 , I_5 are defined by (5.20). By Lemma 5.3, it is easy to see that $I_4(x, y)$ is a bounded continuous function. For I_5 , define $H_1(x, \xi)$, $H_2(x, \xi)$, $K_1(x, \xi)$ and $\Xi(x, y; \xi)$ by (5.21). We define $K_2(x, \xi)$ by

$$K_2(x, \xi) = (b_1(x, \xi) + b_1(x))\xi^{\omega_1} + (b_2(x, \xi) + b_2(x))\xi \log \xi + 2c(x)\xi.$$

Then we get

$$I_{\mathfrak{s}}(x, y) = \frac{1}{\pi} \int_{\mathrm{Jo}}^{\infty} \frac{H_{\mathfrak{l}}(x, \xi)H_{\mathfrak{l}}(x, \xi)}{\Xi(x, y, \xi)} d\xi + \frac{1}{\pi} \int_{Jo}^{\infty} \frac{K_{\mathfrak{l}}(x, \xi)K_{\mathfrak{l}}(x, \xi)}{\Xi(x, y; \xi)} d\xi$$
$$\equiv I_{\mathfrak{s}} + I_{\mathfrak{r}}.$$

Using Lemma 5.3, we can show that I_6 is bounded continuous. For I_7 , set

$$a_{1}\left(x,\frac{\eta}{y-x}\right) - a_{1}(x) \equiv \bar{a}_{1}(x,y;\eta) \frac{|y-x|}{\eta},$$

$$b_{1}\left(x,\frac{\eta}{y-x}\right) - b_{1}(x) \equiv b_{1}(x,y-\eta) \frac{|y-x|^{\sigma_{1}-1}}{\eta^{\sigma_{1}-1}},$$

$$a_{2}\left(x,\frac{\eta}{y-x}\right) - a_{2}(x) \equiv \bar{a}_{2}(x,y;\eta) \frac{|y-x|}{\eta} \log \frac{\eta}{|y-x|}$$

and

$$b_2\left(x, \frac{\eta}{y-x}\right) - b_2(x) = b_2(x, y \eta) \frac{-1}{\log \frac{\eta}{|y-x|}}$$

Substituting this into I_7 and applying the second mean value theorem to I_7 , we obtain

$$I_{7}(x, y) = -\frac{\operatorname{sgn}(y-x)}{\pi} |y-x|^{2\sigma_{1}-3} v_{10}(x, y),$$

where v_{10} is bounded continuous. Thus we have (5.17). By the same way, we obtain (5.18). This completes the proof.

Lemma 5.5. If $\sigma(x) \equiv 0$ and $\alpha_1 > \alpha_2$, $\alpha_1 > 1$, then $g_{\lambda}(x, y)$ satisfies one of the following conditions:

(A)
$$\lim_{y \neq x} \frac{\partial g_{\lambda}}{\partial y}(x,y) = \frac{\partial g_{\lambda}}{\partial y}(x,x-) \text{ exists finitely and } \lim_{y \neq x} \frac{\partial g_{\lambda}}{\partial y}(x,y) = -\infty.$$
$$\lim_{x \neq y} \frac{\partial g_{\lambda}}{\partial y}(x,y) = \frac{\partial g_{\lambda}}{\partial y}(y+,y) \text{ exists finitely and } \lim_{x \neq y} \frac{\partial g_{\lambda}}{\partial y}(x,y) = -\infty.$$

(B) There exist nonnegative constants c_1, c_2, a, β, r and \mathcal{E}_0 such that $c_1 > 1, c_2 > 1, 0 < \alpha < \beta < 1$ and

$$\begin{aligned} \varepsilon^{\beta}/c_{1} < g_{\lambda}(x, x) - g_{\lambda}(x, x-\varepsilon) < c_{1}\varepsilon^{\beta}, \\ \varepsilon^{\sigma} \left(\log \frac{1}{\varepsilon} \right)^{r} / c_{2} < g_{\lambda}(x, x) - g_{\lambda}(x, x+\varepsilon) < c_{2}\varepsilon^{\sigma} \left(\log \frac{1}{\varepsilon} \right)^{r}, \\ for any \, \varepsilon \in (0, \, \varepsilon_{0}] \text{ and } x. \end{aligned}$$

Proof. By virtue of Lemma 5.4, we have

$$\frac{\partial g_{\lambda}}{\partial y}(x, y) = \begin{cases} O(|y-x|^{a_1-2}), & \text{as } y \downarrow x, \\ o(|y-x|^{a_1-2}), & \text{as } y \uparrow x. \end{cases}$$

Therefore only two cases occur: (i) $\lim_{y \neq x} \frac{\partial g_{\lambda}}{\partial y}(x, y)$ and $\lim_{x \neq y} \frac{\partial g_{\lambda}}{\partial y}(x, y)$ exist finitely or (ii) $\lim_{y \neq x} \frac{\partial g_{\lambda}}{\partial y}(x, y)$ and $\lim_{x \neq y} \frac{\partial g_{\lambda}}{\partial y}(x, y)$ do not exist. If the inequality $g_{\lambda,0}(x, x) - g_{\lambda,0}(x, y) \ge 0$ holds (we show this soon later), then since $\alpha_1 < 2$ we have $\lim_{x \to y} \frac{\partial g_{\lambda}}{\partial y}(x, y) = \lim_{x \to y} \frac{\partial g_{\lambda}}{\partial y}(x, y) = \infty$. Therefore if (i) occurs, it is nothing but the case (A). Let (ii) occur. Then $\frac{\partial g_{\lambda}}{\partial y}(x, y) = O(|x - y|)$ for some $\alpha_3: \alpha_1 - 2 < \alpha_3 < 0$ and $r \ge 0$. Therefore we have

$$g_{\lambda}(x,x)-g_{\lambda}(x,x+\varepsilon)=\int_{x+\varepsilon}^{x}\frac{\partial g_{\lambda}}{\partial y}(x,y)dy=O(\varepsilon^{\omega_{1}-1})$$

and

$$g_{\lambda}(x, x) - g_{\lambda}(x, x-\varepsilon) = \int_{x-\varepsilon}^{x} \frac{\partial g_{\lambda}}{\partial y}(x, y) dy = O\left(\varepsilon^{\omega_{3}+1}\left(\log\frac{1}{\varepsilon}\right)^{r}\right).$$

Since $0 < \alpha_1 - 1 < \alpha_3 + 1 < 1$, it follows that the case (B) holds. Therefore it sufficies to prove that $g_{\Lambda,0}(x, x) - g_{\Lambda,0}(x, y) \ge 0$ for all x, y. Let x be any point in \mathbb{R}^1 and fix it. Set

$$g_{\lambda}^{(x)}(z) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-iz\xi} \frac{d\xi}{\lambda - \psi(x, \xi)}.$$

Let $\{X_t^{(x)}, P_y^{(x)}, y \in R^1\}$ be a Lévy process whose exponent $\psi^{(x)}(\xi)$ is $\psi(x, \xi)$. Then $g_{\lambda}^{(x)}(z)$ is the density of the Green measure of Lévy process $X^{(x)}$ with respect to Lebesgue measure. Let $G_{\lambda}^{(x)}$ be the corresponding Green operator and \tilde{x} be any point such that $\tilde{x} \neq 0$. Let *B* be an interval containing the origin and $B \cap \{\tilde{x}\} = \phi$. Then for any Borel set A(cB),

$$\int_{A} g_{\lambda}^{(x)}(y-\tilde{x}) dy = \int_{A} E_{\tilde{x}}^{(x)} [e^{-\lambda \sigma_B} g_{\lambda}^{(x)}(y-X_{\sigma_B}^{(x)})] dy \,.$$

Since $g_{\lambda}^{(\infty)}$ is continuous, we have

$$g_{\lambda}^{(x)}(y-\tilde{x}) = E_{\tilde{x}}^{(x)}[e^{-\lambda\sigma_B}g_{\lambda}^{(x)}(y-X_{\sigma_B}^{(x)})] \quad \text{for } y \in B.$$

Since the exponent $\psi^{(x)}$ of the Levy process $X^{(x)}$ satisfies the following inequality

$$\int_{-\infty}^{\infty} \left\{ \operatorname{Re}(\lambda - \psi^{(x)}(\xi)) \right\} |\lambda - \psi^{(x)}(\xi)|^{-2} d\xi < +\infty \qquad (\lambda > 0) ,$$

by virtue of Theorem 2 of [5] we have $P_{\tilde{L}}^{(x)}(\sigma_0 < \infty) > 0$. Letting $B \downarrow \{0\}$, by quasi-left continuity of Lévy process and by the above fact, we obtain

$$g_{\lambda}^{(x)}(-\tilde{x}) = E_{\tilde{x}}^{(x)}[e^{-\lambda\sigma_0}]g_{\lambda}^{(x)}(0) < g_{\lambda}^{(x)}(0)$$
.

Noting that $g_{\lambda}^{(x)}(z) = g_{\lambda,0}(x,x+z)$, we have

$$g_{\lambda,0}(x, x-\tilde{x}) \leq g_{\lambda,0}(x, x)$$
 for any x and $\tilde{x} \neq 0$.

Thus Lemma 5.5 is proved.

Lemma 5.6. If $\sigma(x)=0$ and $\alpha_1=\alpha_2>1$, then $g_{\lambda}(x, y)$ satisfies the following:

(5.22)
$$\lim_{\varepsilon \neq 0} \frac{g_{\lambda}(x, y) - g_{\lambda}(x, y \pm \varepsilon)}{g_{\lambda}(x, x) - g_{\lambda}(x, x \pm \varepsilon)} = 0, \quad \text{for any } y \neq x,$$

(5.23)
$$-\infty < \lim_{\overline{\mathfrak{e}} \neq 0} \frac{g_{\lambda}(-\varepsilon, -\varepsilon) - g_{\lambda}(-\varepsilon, 0)}{g_{\lambda}(0, 0) - g_{\lambda}(0, -\varepsilon)} \leq \lim_{\overline{\mathfrak{e}} \neq 0} \frac{g_{\lambda}(-\varepsilon, -\varepsilon) - g_{\lambda}(-\varepsilon, 0)}{g_{\lambda}(0, 0) - g_{\lambda}(0, -\varepsilon)} < +\infty.$$

Proof. Set $\tilde{a}_1(x, \xi) = a_1(x, \xi) + a_2(x, \xi)$, $\tilde{a}_2(x, \xi) = b_1(x,\xi) + b_2(x, \xi)$, $\tilde{a}_1(x) = a_1(x) + a_2(x)$ and $\tilde{a}_2(x) = b_1(x) + b_2(x)$. Then \hat{g}_{λ} can be expressed in the form ∂y

$$\frac{\partial g_{\lambda,0}(x,y)}{\partial y} = \frac{1}{\pi} \int_{0}^{\infty} \frac{\lambda\xi \sin(y-x)\xi d\xi}{[\lambda-\psi_{R}(x,\xi)]^{2}+\psi_{I}(x,\xi)^{2}} - \frac{1}{\pi} \int_{0}^{\infty} \frac{\tilde{a}_{1}(x)\xi^{1+s_{1}}\sin(y-x)\xi}{[\lambda-\psi_{R}(x,\xi)]^{2}+\Phi_{I}(x,\xi)^{2}} \xi \\ -\frac{1}{\pi} \int_{0}^{\infty} \left\{ \frac{\tilde{a}_{1}(x,\xi)\xi^{1+s_{1}}}{\lambda-\gamma_{R}(x,\xi)]^{2}+\psi_{I}(x,\xi)^{2}} - \frac{\tilde{a}_{1}(x)\xi^{1+s_{1}}}{[\lambda-\Phi_{R}(x,\xi)]^{2}+\Phi_{I}(x,\xi)^{2}} \right\} \\ \times \sin(y-x)\xi d\xi + \frac{1}{\pi} \int_{0}^{\infty} \frac{c_{1}(x)\xi^{2}+\tilde{a}_{2}(x,\xi)\xi^{1+s_{1}}}{[\lambda-\psi_{R}(x,\xi)]^{2}+\psi_{I}(x,\xi)^{2}} \cos(y-x)\xi d\xi$$

$$\equiv I_1 + I_2 + I_3 + I_4.$$

It is easy to see that I_1 is bounded continuous. Using Lemma 5.3, we have $I_3=O(|y-x|^{2\omega_1-3})$. For I_4 , noting that $\tilde{a}_2(x)=0$, we have $I_4=O(|y-x|^{2\omega_1-3})$. Next we consider I_2 . Let *n* be the largest positive integer such that $\alpha_1-2+(n-1)(2-2\alpha_1 \leq 0)$. Then we have

$$\int_{0}^{\infty} \frac{A_{k}(x)\xi^{1+\omega_{1}+(k-1)(2-2\omega_{1})}\sin(y-x)\xi}{(\lambda+\tilde{a}_{1}(x)\xi^{\omega_{1}})^{2}+c_{1}(x)^{2}\xi^{2}}d\xi = \int_{0}^{\infty}B_{k}(x)\xi^{1-\omega_{1}+(k-1)(2-2\omega_{1})}\sin(y-y)\xi d\xi + \int_{0}^{\infty}\frac{A_{k+1}(x)\xi^{1+\omega_{1}+k(2-2\omega_{1})}\sin(y-x)\xi}{(\lambda+\tilde{a}_{1}(x)\xi^{\omega_{1}})^{2}+c_{1}(x)^{2}\xi^{2}}d\xi + v_{11}(x,y),$$

where $A_1(x) = \tilde{a}_1(x)$, $B_k(x) = A_k(x)/\tilde{a}_1(x)^2$, $A_{k+1}(x) = -B_k(x)c_1(x)^2$, $1 \le k \le n+1$, and $v_{11}(x,y)$ is a bounded continuous function.

From this formula we have

$$I_{2}(x, y) = -\frac{h_{1}(x)}{\pi} \operatorname{sgn}(y-x) |y-x|^{\omega_{1}-2} + \sum_{j=2}^{n+1} h_{j}(x) \operatorname{sgn}(y-x) |y-x|^{-2+(j-1)(2-2\omega_{1})} + v(x, y),$$

where $h_1(x) = \frac{\tilde{h_j}}{2\Gamma(\alpha_1 - 1)\cos\frac{n \,\tilde{\alpha_1}}{2} \cdot \tilde{\alpha_1}(x)} x_2 \leq j \leq n+1$, and v are bounded con-

tinuous functions. (5.23) follows from this estimate. Since $g_{\lambda} \in C^{\infty}$ $(R \times R - \Delta)$ and $g_{\lambda}(x, x) - g_{\lambda}(x, x \pm \varepsilon) = O(\varepsilon^{\omega_1 - 2})$, we have (5.22). This completes the proof.

Now we shall show that Lemma 5.2, 5.5 and 5.6 hold without the condition (c). Let A be the operator defined by (2.1) with the conditions (a) and (b). Let $\chi(y)$ be a C^{∞} -f unction of compact support such that $0 \le \chi \le 1$ and $\chi(y)=1$ for $|y| \le L_1$, =0 for $|y| \ge L_1+1$. Set

$$n_1(x,y) = n(x,y)\chi(y), \quad n_2(x,y) = n(x,y)(1-\chi(y)) \text{ and}$$
$$a_1(x) = a(x) - \int_{-\infty}^{\infty} \frac{yn_2(x,y)}{1+y^2} dy.$$

A can be written as follows.

$$Au(x) = \left\{a_{1}(x)u'(x) + \sigma(x)^{2}u''(x)/2 + \int_{-\infty}^{\infty} \left[u(x+y) - u(x) - \frac{yu'(x)}{1+y^{2}}\right]n_{1}(x, y)dy\right\}$$

$$(5.24) \qquad \qquad + \int_{-\infty}^{\infty} \left[u(x+y) - u(x)\right]n_{2}(x, y)dy \equiv A^{(1)}u(x) + A^{(2)}u(x).$$

Since $A^{(1)}$ satisfies the condition (c), there exists the density $g_{\lambda}^{(1)}$ of the Green operator $G_{\lambda}^{(1)}$ of $A^{(1)}$. The next lemma is needed to show the regularity of g_{λ_1}

in the Theorem 5.1.

Lemma 5.7. Let k(x,y) be the function defined by (4.15) and K be the operator defined by (4.14). Then k(x, y) can be written in the form.

(5.25)
$$\mathbf{k}(\mathbf{x},\mathbf{y}) = \prod_{\mathbf{J}=\infty} g_{\boldsymbol{\lambda}}^{(1)}(\mathbf{x},\mathbf{z})\mathbf{h}(\mathbf{z},\mathbf{y})dz ,$$

where **h** is the kernel of K(h(x,y)) corresponds to k(x, x-y) in Theorem 3.3).

Proof. Note that k(x, y) is the kernel of $G_{\lambda}K$. On the other hand, we have for $f \in \mathcal{D}$,

$$G_{\lambda}Kf(x) = \int_{-\infty}^{\infty} g_{\lambda}^{(1)}(x, z)(Kf)(z)dz = \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} g_{\lambda}^{(1)}(x, z)h(z, y)dz \right\} f(y) dy.$$

Therefore we obtain (5.25), which completes the proof.

Theorem 5.1. Let A be the operator defined by (2.1) with the conditions (a) and (b). Then we have

(i) For any $f \in H_{\infty}$, there exists a unique $u \in \dot{B}$ such that

$$(\lambda - A)u = f.$$

(ii) Let G_{λ} be the inverse of $(\lambda - A)$. Then G_{λ} has the kernel representation

$$G_{\lambda}f(x)=\int_{-\infty}^{\infty}g_{\lambda}(x, y)f(y)dy,$$

where $g_{\lambda}(x, y)$ is of the form

$$g_{\lambda}(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{i(x-y)\xi}}{\lambda - \psi(x, \xi)} d\xi + g_{\lambda,1}(x, y), \qquad g_{\lambda,1} \in C^{1}(R \times R).$$

Proof. Let χ , n_1 , n_2 and a_1 be as in the paragraph before Lemma 5.7. Write A as in (5.24). It is easy to see that the proof of (i) is the same as the proof of [12, p. 537]. From (5.24), we have the following integral equation for the density $g_{\lambda}(x, y)$ of Green measure:

(5.26)
$$g_{\lambda}(x,y) = g_{\lambda}^{(1)}(x,y) + \prod_{\mathbf{J} \to \infty} g_{\lambda}^{(1)}(x,z) A_{z}^{(2)} g_{\lambda}(z,y) dz.$$

The equation (5.26) can be solved by successive approximation and $g_{\lambda}(x,y)$ is given in the form

$$g_{\lambda}(x, y) = g_{\lambda}^{(1)}(x, y) + \sum_{k=2}^{\infty} g_{\lambda}^{(k)}(x, y) ,$$

where $g_{\lambda}^{(k)}(x, y) = \int_{-\infty}^{\infty} g_{\lambda}^{(1)}(x, z) A_{z}^{(2)} g_{\lambda}^{(k-1)}(z, y) dz = 2, 3, \cdots$. Using the estimates

for $g_{\lambda}^{(1)}$ in §5, Lemma 5.4, 5.6 and 5.7, we obtain $\sum_{k=2}^{\infty} g_{\lambda}^{(k)}(x,y)$ is a continuously differentiable function. Hence setting $g_{\lambda,1}(x,y) = g_{\lambda,1}^{(1)}(x,y) + \sum_{k=2}^{\infty} g_{\lambda}^{(k)}(x,y)$, we prove the theorem.

6. **Proof of Theorem 1.** Now we can prove the Theorem 1. Let x and y be any point in \mathbb{R}^1 such that $x \neq y$. Choose an open interval **B** containing y such that $B \cap \{x\} = \phi$. Then for any Borel set $A \subset B$, we have

$$\int_A g_{\lambda}(x, z) dz = \int_A E_x[e^{-\lambda \sigma_B} g_{\lambda}(X_{\sigma_B}, z)] dz.$$

Since g_{λ} is continuous, we have

(6.1)
$$g_{\lambda}(x, z) = E_{z}[e^{-\lambda} g_{\lambda}(X_{\sigma_{B}}z)], \quad z \in B.$$

By quasi-left continuity of X, letting $B \downarrow \{y\}$, we obtain

(6.2)
$$g_{\lambda}(x,y) = E_{x}[e^{-\lambda\sigma_{y}}]g_{\lambda}(yy), \qquad x \neq y.$$

Set $g_{\lambda,0}(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{i(x-y)\xi}}{\sqrt{\psi(x,\xi)}} d\xi$. Then as is seen in the proof of Lemma 5.5, we have $g_{\lambda,0}(x,y) > 0$ for any x, y. Next we show $g_{\lambda}(x,x) > 0$. First note that $g_{\lambda} = g_{\lambda,0} + g_{\lambda,1} \ge 0$, $g_{\lambda,1} \in C^1$ and $g_{\lambda,0}(x, x) \ge g_{\lambda,0}(x, x)$. If $g_{\lambda}(x, x) = 0$ then we have $g_{\lambda,0}(x, y) - g_{\lambda,0}(x, x) \ge -g_{\lambda,1}(x, y) + g_{\lambda,1}(x, x)$ Therefore

$$\overline{\lim_{y \neq x}} \frac{g_{\lambda,0}(x,y) - g_{\lambda,0}(x,x)}{y - x} \leq \overline{\lim_{y \neq x}} \frac{-g_{\lambda,1}(x,y) + g_{\lambda,1}(x,x)}{y - x} \\
= \underbrace{\lim_{y \neq x}}_{y - x} \frac{-g_{\lambda,1}(x,y) + g_{\lambda,1}(x,x)}{y - x} \leq \underbrace{\lim_{y \neq x}}_{y - x} \frac{g_{\lambda,0}(x,y) - g_{\lambda,0}(x,x)}{y - x}.$$

Since $g_{\lambda,0}(x, \cdot)$ is not differentiable at x and $g_{\lambda,0}(x, \cdot) \leq g_{\lambda,0}(x, x)$, we have

$$\underline{\lim_{y \neq x}} \frac{g_{\lambda,0}(x,y) - g_{\lambda,0}(x,x)}{y - x} < \overline{\lim_{y \neq x}} \frac{g_{\lambda,0}(x,y) - g_{\lambda,0}(x,x)}{y - x}$$

This is a contradiction. Thus we have $g_{\lambda}(x, x) > 0$. Therefore we can choose finite number of open sets U_1, \dots, U_l such that $g_{\lambda}(x', y') > \varepsilon > 0$ if $x', y' \in U_i$, $i=1, \dots, l$ and $U_i \cap U_{i+1} \neq \phi$, $i=1, \dots, l-1$, $\bigcup_{i=1}^{l} U_i \supset [x, y]$. Let $x = x_1 < x_2 \cdots < x_l < x_{l+1} = y, x_{i+1} \in U_i \sqcap U_{i+1}, i=1, \dots, l-1$. By (6.2), we have

$$g_{\lambda}(x_i, x_{i+1}) = g_{\lambda}(x_{i+1}, x_{i+1}) E_{x_i}[e^{-\lambda \sigma_{x_{i+1}}}] \qquad i = 1, \dots, l.$$

Hence for some $\delta > 0$,

$$E_{x_i}[e^{-\lambda\sigma_{x_{i+1}}}] = \frac{x_{i+1}}{y_{\lambda}(x_{i+1}, x_{i+1})} > \delta, \qquad i = 1, ..., l.$$

S. MANABE

This implies $P_x(\sigma_x < \infty) > \delta$, $i=1, \dots, /$. By the strong Markov property, we have $P_x(\sigma_y < +\infty) > 0$ for any $x \neq y$, which completes the proof of Theorem 1.

7. Martin boundary theory. In this section, we shall prepare some lemmas from the theory of Martin boundary for the proof of the part (3) of Theorem 2. For the theory of Martin boundary, we refer to Kunita-T.Watanabe [9]. Although almost parallel arguments to [9] hold, we present this section for completeness, since we do not assume the existence of dual process. Let $X = (X_t, P_x, x \in \mathbb{R}^1)$ be the process defined in §2. Define $X^0 = (X_t^0, P_x^0, x \in \mathbb{R}^1 \setminus \{0\})$ by $X_t^0 = X_t$ if $t < \sigma_0$, =0 if $t \ge \sigma_0$ and P_x^0 ()= P_x (, $\sigma_0 < \infty$). Let G_λ^0 be the resolvent of X_t^0 . By the strong Markov property, the density $g_\lambda^0(x, y)$ of $G_\lambda^0(x, dy)$ with respect to dy is given by

(7.1)
$$g_{\lambda}^{0}(x,y) = g_{\lambda}(x,y) - E_{x}[e^{-\lambda\sigma_{0}}]g_{\lambda}(0,y)$$
$$= g_{\lambda}(x,y) - \frac{g_{\lambda}(x,0)}{g_{\lambda}(0,0)}g_{\lambda}(0,y), \qquad x \neq 0.$$

For any Borel set A of $R^1 \setminus \{0\}$, define $\tau_A = \inf\{t \ge 0; X_t^0 \in A\}$. For any bounded measurable function f and for any Borel set A of $R^1 \setminus \{0\}$, define

$$H^{\lambda}_{A}f(x) = E^{0}_{x}[e^{-\lambda\tau_{A}}f(X^{0}_{\tau_{A}}); \tau_{A} < \sigma_{0}].$$

Set $u_1(x) = E_x[e^{-\lambda\sigma_0}; \Omega_1^+]$, $u_2(x) = E_x[e^{-\lambda\sigma_0}; \Omega\Gamma]$ and $u_3(x) = E_x[e^{-\lambda\sigma_0}; \Omega_1^+]$ for $x \neq 0$. We call f a λ -harmonic function (relative to X^0) in $R^1 \setminus \{0\}$ if for any open set A with A compact in $R^1 \setminus \{0\}$,

(7.2)
$$f(x) = E_x^0 [e^{-\lambda \tau_A c} f(X^0_{\tau_A c} \tau_A c < \sigma_0].$$

REMARK 7.1. It follows from the existence of the continuous density of $G_{\lambda}(x, dy)$ that every λ -excessive function is lower semi-continuous. Indeed by Fatou's lemma, $G_{\lambda}f(x)$ lower semi-continuous (/ is nonnegative bounded measurable). Since every λ -excessive function f is an increasing limit of $G_{\lambda}f_n$ (f_n is nonnegative bounded measurable), / is lower semi-continuous. In view of [1, p. 197], we see that there exists a reference measure.

Lemma 7.1. u_i , i=1, 2, 3 are λ -harmonic (relative to X^0) in $\mathbb{R}^1 \setminus \{0\}$.

Proof. First consider the case i=1. Note that $\tau_{A^c} < \sigma_0$ on Ω_1 . Indeed by Remark 7.1, we can apply Theorem 4.2 of [16]. Therefore we have

$$E_{\mathbf{x}}[e^{-\lambda\tau_{\mathbf{A}^{c}}};X_{\tau_{\mathbf{A}^{c}}}\in \mathcal{A} X_{\tau_{\mathbf{A}^{c}}}=0]=0$$

Thus we get $P_x(\tau_{a^c} = \sigma_0, \Omega_1) = 0$. By the strong Markov property, we have

$$u_1(x) = E^0_x[e^{-\lambda \tau_{A^c}}u_1(X^0_{\tau_{A^c}}); \tau_{A^c} < \sigma_0].$$

By the same way, we obtain (7.2) for u = 2, 3, which proves the lemma.

Lemma 7.2. (1) Let A be a compact set of $\mathbb{R}^{1} \setminus \{0\}$ and $\mathbb{B}(\subset \mathbb{R}^{1} \setminus \{0\})$ be a neighborhood of A. Then we have

(7.3)
$$\inf_{x \in \mathcal{A}} \int_{\mathcal{B}} g_{\lambda}^{0}(y, x) dy > 0$$

(2) (i) For any open set A in $\mathbb{R}^n \setminus \{0\}$ and for any $y \in A$, we have

$$H^{\lambda}_{A}g^{0}_{\lambda}(x, y) = g^{0}_{\lambda}(x, y) \, .$$

(ii) For any $y \in \mathbb{R}^1 \setminus \{0\}$, $g_{\lambda}^0(, y)$ is λ -harmonic (relative to X^0) in $\mathbb{R}^1 \setminus \{0, y\}$. (3) Let $x_0 \in \mathbb{R}^1 \setminus \{0\}$ be fixed. Let V be an open interval of $\mathbb{R}^1 \setminus \{0\}$ containing x_0 with $\overline{V} \subset \mathbb{R}^1 \setminus \{0\}$. Then there exists an open interval $U(x_0)(\subset V)$ containing x_0 such that

$$g^{0}_{\lambda}(x, y) > E^{0}_{x}[e^{-\lambda \tau_{Vc}}g^{0}_{\lambda}(X^{0}_{\tau_{Vc}}, y); \tau_{vc} < \sigma_{0}] \quad \text{for any } x, y \in U(x_{0}).$$

Proof. (1) By (7.1), $g_{\lambda}^{0}(x, y)$ is continuous in $(R^{1} \times R^{1}) \setminus \{0\}$ and

$$g_{\lambda}^{0}(x, x) = g_{\lambda}(x, x) - \frac{g_{\lambda}(x, 0)}{g_{\lambda}(0, 0)} g_{\lambda}(0, x) > g_{\lambda}(x, x) - g_{\lambda}(0, x) > \mathbf{0}$$

Therefore $\inf_{x \in \mathcal{A}} \int_{\mathcal{B}} g^{0}_{\lambda}(y, x) dy > 0$. Thus (1) is proved.

(2) (i). Let A be any open set in $\mathbb{R}^1 \setminus \{0\}$ and / be any continuous function of compact support such that f=0 on \mathbb{A}^c . Then by the strong Markov property we have

$$\begin{split} &\int_{R^{1}\setminus\{0\}} E^{0}_{x} \left[e^{-\lambda\tau_{A}} g^{0}_{\lambda} (X^{0}_{\tau_{A}}, y); \tau_{A} < \sigma_{0} \right] f(y) dy = E^{0}_{x} \left[\int_{0}^{\infty} e^{-\lambda t} f(X^{0}_{t}) dt; \tau_{A} < \sigma_{0} \right] \\ &= \int_{R^{1}\setminus\{0\}} g^{0}_{\lambda} (x, y) f(y) dy \,. \end{split}$$

Since $g_{\lambda}^{0}(x, y)$ is continuous in $(R^{1}X R^{1}) \setminus \{0\}$, we get

$$E^0_x[e^{-\lambda\tau_A}g^0_\lambda(X^0_{\tau_A}, y); \tau_A < \sigma_0] = g^0_\lambda(x, y) \quad \text{for} y \in A \text{ and } x \neq 0.$$

For the proof (ii), let A be an open set in $\mathbb{R}^1 \setminus \{0, y\}$ with \overline{A} compact in $\mathbb{R}^1 \setminus \{0, y\}$. Then by (i), for $y' \in (\overline{A})^c$ we have

$$H^{\lambda}_{(\overline{A})^{c}}g^{0}_{\lambda}(x, y') = g^{0}_{\lambda}(x, y').$$

Thus (2) is proved.

(3) For any $x, y \neq 0$ ($x \neq y$), we have

S. MANABE

$$g_{\lambda}^{0}(x,y) = g_{\lambda}^{0}(y,y) E_{x}^{0}[e^{-\lambda \tau_{y}}; \tau_{y} < \sigma_{0}].$$

Since by the right continuity of paths, $P_x \{\tau_y > 0\} = 1$, we have

(7.4) $g_{\lambda}^{0}(x, y) < g_{\lambda}^{0}(y, y)$

Applying the Riemann-Lebesgue theorem to $g_{\lambda}^{(1)}(x, y)$, we have $\lim_{|x|\to\infty} g_{\lambda}^{(1)}(x,y)=0$, uniformly on any compact set of y. Therefore by virtue of (7.1), (5.26) and Lemma 5.7, we have

(7.5)
$$\lim_{|z|\to\infty} g_{\lambda}^0(z,y) = 0 \quad \text{uniformly in } y: |x_0-y| < 1.$$

Therefore it follows from (7.4), (7.5) and the continuity of g_{λ}^{0} that there exist constants δ_{0} and $\mathcal{E}_{0} > 0$ such that

$$\inf_{x \to \pi} [g^{0}_{\lambda}(y, y) - g^{0}_{\lambda}(z, y)] \geq \varepsilon_{0} \quad \text{for } y \colon |x_{0} - y| < \delta_{0}.$$

By the continuity of g_{λ}^{0} , for any $\varepsilon_{1}: 0 < \varepsilon_{1} < \varepsilon_{0}$, there exists a constant $\delta > 0$ ($\delta < \delta_{0}$) such that for $|x - x_{0}| < \delta$, $|y - x_{0}| < \delta$,

$$g_{\lambda}^{0}(y, y) - g_{\lambda}^{0}(x, y) < \varepsilon_{1}$$
.

Therefore we have

$$E^{0}_{\boldsymbol{y}}[e^{-\lambda\tau_{\boldsymbol{Y}c}}g^{0}_{\lambda}(X^{0}_{\tau_{\boldsymbol{Y}c}},y);\tau_{\boldsymbol{Y}c} < \sigma_{0}] \leq g^{0}_{\lambda}(y,y) - \mathcal{E}_{0} < g^{0}_{\lambda}(x,y) .$$

So it is sufficient to put $U(x_0) = \{x; |x-x_0| < \delta\} \prod V$, which proves the Lemma 7.2.

Lemma 7.3. Let $\{\mu_n\}$ be a sequence of measures on $R^1 \setminus \{0\}$. Define $G^0_{\lambda} \mu_n by$

$$G^{0}_{\lambda}\mu_{n}(x) = \int_{R^{1}\setminus\{0\}} g^{0}_{\lambda}(x, y)\mu_{n}(dy) \, .$$

If there exists a locally integrable function v such that $G^{0}_{\lambda}\mu_{n} \leq v$, then we have

(i) There exists a subsequence $\{\mu_{n_k}\}$ of $\{\mu_n\}$ such that $\{\mu_{n_k}\}$ converges to some μ weakly. (ii) If for the above $\{\mu_{n_k}\}, G^0_{\lambda} \mu_{n_k}$ converges to some u almost everywhere with respect to Lebesgue measure, then

$$\lim_{\alpha \neq \infty} \alpha G^0_{\alpha} u \geq G^0_{\lambda} \mu .$$

Especially if u is λ -excessive, then $u \ge G_{\lambda}^{0} \mu$.

(iii) If for any nonnegative bounded measurable function f with compact support and for any $\pounds > 0$, there exists a compact set A in $\mathbb{R}^n \setminus \{0\}$ such that

(7.6)
$$(\tilde{G}_{\lambda}^{0}f(y)\mu_{n}(dy) < \varepsilon, \quad n = 1, 2, ...,$$

HITTING PROPERTIES

then $\lim_{\sigma_1 \to \infty} \alpha G^0_{\omega} u = G^0_{\lambda} \mu$, where $\hat{G}^0_{\lambda} f(y) = \int_{\mathcal{A}^1 \setminus \{0\}} g^0_{\lambda}(x,y) f(x) dx$. Note that if there exists a compact set A_1 such that $\operatorname{supp}(\mu_n) \subset A_1$, $n = 1, 2, \cdots$, then the condition in (iii) is fulfilled.

Proof. For the assertion (i) it is enough to show that, for each compact set, $\mu_n(A)$ is bounded. Let *B* be a compact neighborhood of *A*. By Lemma 7.2, $c = \inf_{x \in A} \int_B^{0} g_{(y, x)}^0 dy > 0$. Hence

$$\infty > \int_{B} v(x) dx \ge \int_{B} G_{\lambda}^{0} \mu_{n}(x) dx \ge \int_{A} \left\{ \int_{B} g_{\lambda}^{0}(x, y) dx \right\} \mu_{n}(dy) \ge c \mu_{n}(A) .$$

For (ii), let μ_{n_k} converge weakly to μ and let f be a nonnegative bounded measurable function of compact support. Using the facts that $v \ge G_{\lambda}^{0} \mu_{n_k}$ and $\mathring{G}_{\lambda}^{0} f$ is nonnegative continuous, we have

$$\int_{R^{1}\setminus\{0\}} f(x)u(x)dx = \lim_{k\to\infty} \int_{R^{1}\setminus\{0\}} f(x)G_{\lambda}^{0}\mu_{n_{k}}(x)dx = \lim_{k\to\infty} \int_{R^{1}\setminus\{0\}} \hat{G}_{\lambda}^{0}f(y)\mu_{n_{k}}(dy)$$
$$\geq \int_{R^{1}\setminus\{0\}} \hat{G}_{\lambda}^{0}f(y)\mu(dy) = \int_{R^{1}\setminus\{0\}} f(x)G_{\lambda}^{0}\mu(x)dx.$$

Therefore $u \ge G_{\lambda}^{0} \mu$ a.e. rf^{*}. On the other hand, for any $\alpha > 0$,

$$u \equiv \lim_{k \to \infty} G^0_{\lambda} \mu_{n_k} \ge \lim_{k \to \infty} \alpha G^0_{\alpha} (G^0_{\lambda} \mu_{n_k}) \ge \alpha G^0_{\alpha} (\lim_{k \to \infty} G^0_{\lambda} \mu_{n_k}) = \alpha G^0_{\alpha} u \quad \text{a.e. } dx.$$

So we have

$$\int_{R^{1}\setminus\{0\}} f(y) \lim_{\substack{\omega \neq \infty \\ \sigma \neq \infty}} \alpha G^{0}_{\omega} u(x) dx$$

=
$$\lim_{\substack{\omega \neq \infty \\ \sigma \neq \infty}} \int_{R^{1}\setminus\{0\}} \alpha \hat{G}^{0}_{\omega} f(y) u(y) dy \ge \lim_{\substack{\sigma \neq \infty \\ \sigma \neq \infty}} \int_{R^{1}\setminus\{0\}} \alpha \hat{G}^{0}_{\omega} f(y) G^{0}_{\lambda} \mu(y) dy$$

$$\ge \int_{R^{1}\setminus\{0\}} f(y) \lim_{\substack{\omega \neq \infty \\ \sigma \neq \infty}} \alpha G^{0}_{\omega} (G^{0}_{\lambda} \mu)(y) dy = \int_{R^{1}\setminus\{0\}} f(y) G^{0}_{\lambda} \mu(y) dy .$$

Hence we have

(7.7)
$$\lim_{\omega \to \infty} \alpha G^0_{\omega} u(x) \ge G^0_{\lambda} \mu(x) \quad \text{a.e. } dx .$$

Since the both sides of (7.7) are λ -excessive, we have $\lim_{\omega_{\uparrow} \to \infty} \alpha G^0_{\omega} u \ge G^0_{\lambda} \mu$. Finally we shall prove (iii). Let *f* be a nonnegative bounded measurable function with compact support. Then by the assumption, for any £>0, there exists a compact set *A* in $R^i \setminus \{0\}$ such that (7.6) holds. So we have

$$\lim_{k\to\infty}\int_{R^{1}\setminus\{0\}} \hat{G}^{0}_{\lambda}f(y)\mu_{n_{k}}(dy) \leq \lim_{k\to\infty}\int_{A} \hat{G}^{0}_{\lambda}f(y)\mu_{n_{k}}(dy) + \lim_{k\to\infty}\int_{A^{c}} \hat{G}^{0}_{\lambda}f(y)\mu_{n_{k}}(dy)$$

S. Manabe

$$\leq \int_{R^{1}\setminus\{0\}} \hat{G}^{0}_{\lambda}f(y)\mu(dy) + \lim_{k \to \infty} \int_{A^{c}} \hat{G}^{0}_{\lambda}f(y)\mu_{n_{k}}(dy) \leq \int_{R^{1}\setminus\{0\}} \hat{G}^{0}_{\lambda}f(y)\mu(dy) + \varepsilon.$$

Therefore we have

$$\int_{\mathbb{R}^1\setminus\{0\}} f(x)u(x)dx \leq \int_{\mathbb{R}^1\setminus\{0\}} \hat{G}^0_{\lambda}f(y)\mu(dy) + \varepsilon.$$

Hence $u \leq G_{\lambda}^{0} \mu$ a.e. dx. This implies that $\lim_{\alpha \uparrow \infty} \alpha G_{\alpha}^{0} u \leq G_{\lambda}^{0} \mu$. Thus the proof is completed.

Lemma 7.4. Let A be an open set in $\mathbb{R}^1 \setminus \{0\}$ with \overline{A} compact in $\mathbb{R}^1 \setminus \{0\}$. Then there exists a measure μ_i (i=1, 2, 3) concentrated in A such that

$$H^{\lambda}_{A}u_{i}(x) = \int_{A} g^{0}_{\lambda}(x, y) \mu_{i}(dy), \quad i = 1, 2, 3$$

Proof. First we show for the case i=1. Since u_1 is λ -excessive, setting $f_n(x) = n[u_1(x) - nG^0_{\lambda+n}u_1(x)]$ we have $u_1(x) = \lim_{n \to \infty} G^0_{\lambda} f_n(x)$. We shall first prove Lemma 7.4 for $v_n(x) = G^0_{\lambda} f_n(x)$. Let B be a compact set in $R^1 \setminus \{0\}$, then by the strong Markov property

$$H^{\lambda}_{B^{c}}v_{n}(x) = E^{0}_{\pi} \left[\int_{\tau_{B^{c}}}^{\sigma_{0}} e^{-\lambda t} f_{n}(X^{0}_{t}) d\tau_{B^{c}} < \sigma_{0} \right] \to 0 \quad \text{as} \quad B \uparrow R^{1} \setminus \{0\} \; .$$

So we have $H_{B^c}^{\lambda}H_A^{\lambda}v_n(x) \leq H_{B^c}^{\lambda}v_n(x) \to 0$ as $B \uparrow R^1 \setminus \{0\}$. Since $H_A^{\lambda}v_n$ is λ excessive, by setting $f_{n,m}(x) = m[H_A^{\lambda}v_n(x) - mG_{\lambda+m}^0 H_A^{\lambda}v_n(x)]$ we have $H_A^{\lambda}v_n(x) = \lim_{m \to \infty} G_{\lambda}^0 f_{n,m}(x)$. Set $v_{n,m} = G_{\lambda}^0 f_{n,m}$ and $\mu_{n,m}(dy) = f_{n,m}(y) dy$. Then using Lemma 7.2, we have

$$\begin{split} H^{\lambda}_{B^c} v_{n,m}(x) &= H^{\lambda}_{B^c} G^0_{\lambda} \mu_{n,m}(x) = \int_B H^{\lambda}_{B^c} g^0_{\lambda}(x, y) \mu_{n,m}(dy) + \int_{B^{cc}} H^{\lambda}_{B^c} g^0_{\lambda}(x, y) \mu_{n,m}(dy) \\ &\geq \int_{B^c} H^{\lambda}_{B^c} g^0_{\lambda}(x, y) \mu_{n,m}(dy) = \int_{B^c} (x, y) \mu_{n,m}(dy). \end{split}$$

For any nonnegative bounded measurable function / with compact support,

$$\begin{split} & \int_{B} \hat{G}_{\lambda}^{0} f(y) \mu_{n,m}(dy) = \int_{\beta} \left\{ \int_{X \setminus \{0\}} g_{\lambda}^{0}(x, y) f(x) dx \right\} \mu_{n,m}(dy) \\ & \leq \int_{R^{1} \setminus \{0\}} H_{B^{c}}^{\lambda} v_{n,m}(x) f(x) dx \leq \int_{R^{1} \setminus \{0\}} H_{B^{c}}^{\lambda} H_{A}^{\lambda} v_{n}(x) f(x) dx \\ & \leq \int_{R^{1} \setminus \{0\}} H_{B^{c}}^{\lambda} v_{n}(x) f(x) dx \rightarrow 0 \quad \text{as} \quad B \uparrow R^{1} \setminus \{0\} \,. \end{split}$$

Therefore by Lemma 7.3, there exists a measure μ_n on $\mathbb{R}^1 \setminus \{0\}$ such that

 $H^{\lambda}_{A}v_{n}(x) = G^{0}_{\lambda}\mu_{n}(x)$. Next we shall show $\mu_{n}((\bar{A})^{c}) = 0$. Let us suppose that there exists a compact subset D of $(\bar{A})^{c}$ such that $\mu_{n}(D) > 0$. Then there exists a point $x_{0} \in D$ such that for any neighborhood Q containing x_{0} , $\mu_{n}(Q) > 0$. Let $V(\subset (\bar{A})^{c})$ be an open interval containing x_{0} with $\bar{V} \subset R^{1} \setminus \{0\}$. Then by Lemma 7.2, there exists an open interval $U(x_{0})(\subset V)$ containing x_{0} such that for any $x, y \in U(x_{0})$,

$$E_x^0[e^{-\lambda_{\tau_{V^c}}}g^0_\lambda(X^0_{\tau_{V^c}},y); \ au_{V^c} < \sigma_0] < g^0_\lambda(x, y)$$

Therefore

$$\int_{U(x_0) \cap D} g^0_{\lambda}(x_0, y) \mu_n(dy) > E^0_{x_0} \left[\int_{U(x_0) \cap D} e^{-\lambda_{\tau_{V_c}}} g^0_{\lambda}(X^0_{\tau_{V_c}}, y) \mu_n(dy) \right].$$

On the other hand $g_{\lambda}^{0}(, y)$ is λ -excessive (relative to X^{0}), we have

$$\int_{(\mathbb{R}^1\setminus\{0\})\setminus(U(x_0)\cap D)} g^0_{\lambda}(x_0, y)\mu_n(dy) \geq \int_{(\mathbb{R}^1\setminus\{0\})\setminus(U(x_0)\cap D)} E^0_x[e^{-\lambda_{\tau_{F_c}}}g^0_{\lambda}(X^0_{\tau_{F_c}}, y)]\mu_n(dy) .$$

Therefore we have

$$v_n(x_0) > E^0_{x_0}[e^{-\lambda \tau_{V^c}} v_n(X^0_{\tau_{V^c}}); \tau_{V^c} < \sigma_0].$$

This contradicts Lemma 7.1. So $\mu_n((\bar{A})^c)=0$. Thus Lemma 7.4 is proved for $v_n(x)=G^0_{\lambda}f_n(x)$. Again applying Lemma 7.3 to $G^0_{\lambda}\mu_n=H^{\lambda}_A v_n$, we have

$$H_A^{\lambda}u_1(x) = G_{\lambda}^0\mu(x)$$
 for some μ with supp $(\mu) \subset \overline{A}$.

The proof for $u_i(x)$, i=2, 3 is similar to that of u_1 , which completes the proof.

Using the estimates in Lemma 5.6, we can construct the Martin boundary Δ of the process X_t^0 . Set

$$\kappa(x, y) = \frac{g_{\lambda}^{0}(x, y)}{g_{\lambda}^{0}(c, y)}.$$

Let A be the set of infinite sequences $\{y_n\}$ such that $\{y_n\}$ does not converge to any point in $\mathbb{R}^1 \setminus \{0\}$ and for which $\kappa(x, y_n)$ converges. We call $\{y_n\}$, $\{z_n\}$ ($\in \tilde{\Delta}$) equivalent if $\lim_{n \to \infty} \kappa(x, y_n) = \lim_{n \to \infty} \kappa(x, z_n)$. Define Δ as the set of all equivalence class of A. Since $\lim_{y \to 0} \kappa(x, y) = \frac{g_{\lambda}(x, \gamma)}{g_{\lambda}(c, 0)}$ is by Lemma 5.6, any $\{y_n\}$ ($\in \tilde{\Delta}$) which converges to 0 are equivalent. So the origin belongs to Δ and there exists no points of Δ except the origin in [-N, N] (N > 0).

Lemma 7.5. Let $\{\mu_n\}$ be a sequence of (Radon) measures on $(R^1 \setminus \{0\}) \cup \Delta$ such that $\operatorname{supp}(\mu_n)$ are contained in [-M, M] (M>0) and $\{\mu_n[-M, M]\}$ is S. Manabe

bounded. Let μ be a weak limit of $\{\mu_{n_b}\}$. If

$$v_n = \int_{(R^1 \setminus \{0\}) \cup \Delta} \kappa(\cdot, \eta) \mu_n(d\eta)$$

is dominated by a locally integrable function and if v_{n_k} converges a.e. dx to a function u, then

$$\lim_{\alpha \uparrow \infty} \alpha G^0_{\alpha} u = \int_{(R^1 \setminus \{0\}) \cup \Delta} \kappa(, \eta) \mu(d\eta) \, .$$

Proof. Let / be a continuous function with compact support, then

$$\int_{R^{1}\setminus\{0\}} f(x)u(x)dx = \int \int_{R^{1}\setminus\{0\}} f(x) \left\{ \int_{(R^{1}\setminus\{0\})\cup\Delta} \kappa(x,\eta)\mu_{n_{k}}(d\eta) \right\} dx$$
$$= \int_{(R^{1}\setminus\{0\})\cup\Delta} \left\{ \int_{R^{1}\setminus\{0\}} f(x)\kappa(x,\eta) \int_{R^{1}\setminus\{0\}} \int_{R^{1}\setminus\{0\}} \int_{R^{1}\setminus\{0\}} \int_{R^{1}\setminus\{0\}\cup\Delta} f(x)\mu(d\eta) dx \right\} dx.$$
Thus $u(x) = \int_{R^{1}\setminus\{0\}} \int_$

Thus $u(x) = \int_{\langle R^1 \setminus \{0\} \rangle \cup \Delta} \kappa(x, \eta) \mu(d\eta)$ a.e. dx. The rest of the proof is the same as that of Lemma 7.3 (iii). This completes the proof.

After above preparation we can prove the following lemma.

Lemma 7.6. There exist nonnegative constants c_1, c_2, c_3 such that

$$\begin{split} E_{\mathbf{x}}[e^{-\lambda\sigma_{0}} \ \Omega_{1}^{+}] &= c_{1}\frac{g}{g_{\lambda}(\mathbf{U}, \mathbf{U})} & \text{for any } \mathbf{x} \neq 0, \\ E_{\mathbf{x}}[e^{-\lambda\sigma_{0}}; \ \Omega_{1}^{-}] &= c_{2}\frac{g_{\lambda}(\mathbf{x}, 0)}{g_{\lambda}(0, 0)} & \text{for any } \mathbf{x} \neq 0, \\ E_{\mathbf{x}}[e^{-\lambda\sigma_{0}}; \ \Omega_{1}^{\pm}] &= c_{3}\frac{g_{\lambda}(\mathbf{x}, 0)}{g_{\lambda}(0, 0)} & \text{for any } \mathbf{x} \neq 0. \end{split}$$

Proof. First we show this lemma for $u_1(x) = E_x[e^{-\lambda\sigma_0} \Omega l]$. For any open set A in $(\mathbb{R}^1 \setminus \{0\}) \cup \Delta$ with \overline{A} compact, we can choose a sequence $\{A_n\}$ of open sets in $\mathbb{R}^1 \setminus \{0\}$ with A_n compact in $\mathbb{R}^1 \setminus \{0\}$ such that $A_n \uparrow A \cap (\mathbb{R}^1 \setminus \{0\})$. We denote [A] for $A \cap (\mathbb{R}^1 \setminus \{0\})$ in the rest of the proof. By Lemma 7.4, there exists a measure ν_n such that

$$H_{A_n}^{\lambda}u_1(x)=\int_{\overline{A}_n}g_{\lambda}^0(x, y)\nu_n(dy)\,.$$

Define μ_n by $\mu_n(dy) = g_{\lambda}^0(c.y)\nu_n(dy)$, then $\operatorname{supp}(\mu_n) \subset A_n^-$ and $\{\mu_n((R^1 \setminus \{0\}) \cup \Delta)\}$ is bounded. Therefore there exists a subsequence $\{\mu_{n_k}\}$ of $\{\mu_n\}$ which converges to some μ_A . On the other hand since

$$H_{A_n}^{\lambda}u_1(x) = \int_{\overline{A}_n} \kappa(x, y) \mu_n(dy) \to H_{[A]}^{\lambda}u_1(x) \quad \text{as} \quad n \to \infty ,$$

and $H_{A_n}^{\lambda} u_1(x) \leq 1$ by Lemma 7.5 we have

$$\lim_{\alpha \uparrow \infty} \alpha G^{0}_{\alpha} H^{\lambda}_{[A]} u_{1} = \int_{\overline{[A]}} \kappa(\cdot, \eta) \mu_{A}(d\eta) \, .$$

Because $H_{[A]}^{\lambda} u_1(x)$ is λ -excessive (relative to X^0), we get

$$H_{[A]}^{\lambda}u_{1}(x)=\int_{\overline{[A]}}\kappa(x, \eta)\mu_{A}(d\eta).$$

Next let A be $\{0\} \in \Delta$. Choose a sequence A_n of open sets in $(\mathbb{R}^1 \setminus \{0\}) \cup \Delta$ such that $A_n \downarrow A$. Then since $H_{[A_n]}^{\lambda} u_i(x)$ is a monotone decreasing sequence and $\{\mu_{A_n}((\mathbb{R}^1 \setminus \{0\}) \cup \Delta)\}$ is bounded, it follows from Lemma 7.5 that there exists a measure μ_0 such that

$$\lim_{\alpha \downarrow \infty} \alpha G^{0}_{\alpha} (\lim_{n \to \infty} H^{\lambda}_{[A_{n}]} u_{1})(x) = \int_{A} \kappa(x, \eta) \mu_{0}(d\eta) \, .$$

On the other hand, in view of the proof of Lemma 7.1, we have $\lim_{n \to \infty} H^{\lambda}_{[A_n]} u_1(x) = u_1(x)$. **Bo we get**

$$\lim_{\alpha \uparrow \infty} \alpha G^{0}_{\alpha} u_{1}(x) = \int_{\mathcal{A}} \kappa(x, \eta) \mu_{0}(d\eta).$$

Since u_1 is λ -excessive, we have

$$u_1(x) = \int_A \kappa(x, \eta) \mu_0(d\eta) = \mu_0(\{0\}) \kappa(x, 0)$$
 .

Thus we proved the lemma for u_1 . By the same way, we can prove the lemma for u_i , i=2, 3. The proof of lemma is completed.

8. Proof of Theorem 2. In this section we shall prove Theorem 2 by using the estimates established in §5. Before entering the proof, we note that σ_0 is an accessible stopping time on Ω_1 . Indeed let R_n be the hitting time of $A_n = \left\{x; E_x[e^{-\lambda\sigma_0}] > 1 - \frac{1}{n}\right\}$ and let $\Lambda = \{R_n < \sigma_0, \forall n\}$. Since $E_x[e^{-\lambda\sigma_0}]$ is λ -excessive (relative to X) and every λ -excessive function is lower semi-continuous (see Remark 7.1), A_n contains an open set which contains the origin. Therefore in view of the proof of Lemma 7.1, $\Lambda \supset \Omega_1$. On the other hand, by Proposition (4.12) of [1. Chapter IV] lim $R_n = T$ on Λ a.s. Thus σ_0 is accessible on Ω_1 .

Proof of (1). Let X_t^0 , G_λ^0 and g_λ^0 be as in the beginning of §7. For $x \neq 0$, put

$$F_{+}(x) = \lim_{y \neq 0} \frac{a^{0}(x-x)}{g^{v}_{\lambda}(a, y)} \quad \text{and} \quad F(x) = \lim_{y \neq 0} \frac{g^{0}_{\lambda}(x, y)}{g^{0}_{\lambda}(b, y)}.$$

Then by Lemma 5.2 we have

(8.1)

$$F_{+}(x) = \lim_{y \neq 0} \frac{g_{\lambda}(x, y)g_{\lambda}(0, 0) - g_{\lambda}(x, 0)g_{\lambda}(0, y)}{g_{\lambda}(a, y)g_{\lambda}(0, 0) - g_{\lambda}(a, 0)g_{\lambda}(0, y)}$$

$$= \frac{a \begin{pmatrix} 0 & 0 \\ \lambda & 0 \end{pmatrix} \frac{\partial g_{\lambda}(x, 0) - a_{\lambda}(x, 0) \frac{\partial g_{\lambda}}{\partial y}(0, 0+)}{g_{\lambda}(0, 0) \frac{\partial g_{\lambda}}{\partial y}(a, \dots - x, -\frac{\partial y}{\partial y}, 0+)}.$$

$$F_{-}(x) = \frac{g_{\lambda}(0, 0) \frac{\partial g_{\lambda}}{\partial y}(x, 0) - g_{\lambda}(x, 0) \frac{\partial g_{\lambda}}{\partial y}(0, 0-)}{g_{\lambda}(0, 0) \frac{\partial g_{\lambda}}{\partial y}(b, 0) - g_{\lambda}(b, 0) \frac{\partial g_{\lambda}}{\partial y}(0, 0-)}.$$

We can choose constants *a* and *b* such that the denominators do not vanish by virtue of Lemma 5.2. In fact, if for all $fI\Phi\theta$

$$g_{\lambda}(0, 0)\frac{\partial g_{\lambda}}{\partial y}(a, 0) - g_{\lambda}(a, 0)\frac{\partial g_{\lambda}}{\partial y}(0, 0+) = 0,$$

then we have $g_{\lambda}(0, 0) \left(\frac{\partial g_{\lambda}}{\partial dy} (0+, 0) - \frac{\partial g_{\lambda}}{\partial y} (0, 0+) \right) = 0$. Hence by (S.2a) and (5.2b), we have $\sigma(0)^{-2} g_{\lambda}(0, 0) = 0$. This is a contradiction. By (8.1), (5.2a) and (5.2b), we get

Dy (0.1), (0.2a) and (0.2b),

$$F_{+}(0+) = \frac{2g_{\lambda}(0,0)}{\sigma(0)^{2} \left\{ g_{\lambda}(0,0) \frac{\partial g_{\lambda}}{\partial y}(a,0) - g_{\lambda}(a,0) \frac{\partial g_{\lambda}}{\partial y}(0,0+) \right\}} \pm 0, \ F_{+}(0-) = 0,$$

$$F_{-}(0-) = \frac{-2g_{\lambda}(0,0)}{\sigma(0)^{2} \left\{ g_{\lambda}(0,0) \frac{\partial g_{\lambda}}{\partial y}(b,0) - g_{\lambda}(b,0) \frac{\partial g_{\lambda}}{\partial y}(0,0-) \right\}} \pm 0 \ F_{-}(0+) = 0.$$

Since $F_+(X_t^0)$ is a λ -excessive function relative to X^0 , $e^{-\lambda t}F_+(X_t^0)$ is a supermartingale¹ having left limit with probability one. Therefore

(8.3)
$$P^{0}_{x} \{\lim_{t \uparrow \sigma_{0}} e^{-\lambda t} F_{+}(X^{0}_{t}) \text{ exists} / \Omega_{1} \} = 1 \quad \text{for } x \neq 0 .$$

Combining (8.2) and (8.3), we have $P_x(\Omega_1^{\pm} | \Omega_1) = 0$. Hence we have $P_x(\Omega_1^{\pm} \cup \Omega^- | \Omega_1) = 1$ for any $x \neq 0$, which completes the proof of (i).

Next we show (ii). Let $\{\tau_n\}$ be the sequence of stopping times such that $\tau_0=0, \ \tau_n=\inf\left\{t \ X_t^0\in\left[-\frac{1}{n}, \frac{1}{n}\right]\right\}$. Then as noted in the paragraph before the

¹⁾ $(e^{\lambda t}F_{+}(X_{t}^{0}))_{0 \le t \le \sigma_{0}}$ is a supermartingale' means $\{e^{-\lambda t}F_{+}(X_{t})I_{(0,\sigma_{0})}(t), \mathcal{G}_{t}, P_{x}\}$ is a supermartingale.

HITTING PROPERTIES

proof of (1), $\tau_n < \sigma_0$, Vn and $\lim \tau_n = \sigma_0$ on Ω_1 a.s. Set $f_{\tau_n}(\omega) = e^{-\lambda \tau_n} F_+(X_n^0(\omega))$. Then $\{f_{\tau_n}(\omega)\}_{n\geq 0}$ is a nonnegative bounded martingale. So there exists $f_{\infty}(\omega) = \lim_{n \to \infty} f_{\tau_n}(\omega)$ such that $E_x^0[f_{\infty}(\omega)/\mathfrak{B}_{\tau}^0 = f_{\tau_n}(\omega)^2)$, (a.s. P_x^0) $n \geq 0$. Hence we have

$$E^{\mathbf{0}}_{\mathbf{x}}[f_{\infty}(\omega)] = E^{\mathbf{0}}_{\mathbf{x}}[f_{\tau_{n}}(\omega)].$$

By (8.2), we have

$$f_{\infty}(\omega) = \begin{cases} e^{-\lambda\sigma_0}F_+(0+) & \text{on } \Omega_1^+, \\ 0 & \text{on } \Omega_1^-. \end{cases}$$

Hence we obtain

$$E_{x}^{0}[e^{-\lambda\sigma_{0}}F_{+}(0+); \quad \Omega_{1}^{+}] = \begin{bmatrix} g_{\lambda}(0, 0) \frac{\partial g_{\lambda}}{\partial A_{\lambda}} (X_{\tau_{n}}^{0}, 0) - g_{\lambda}(X_{\tau_{n}}^{0}, 0) \frac{\partial g_{\lambda}}{\partial A_{\lambda}} (0, 0+) \\ g_{\lambda}(0, 0) \frac{\sigma_{S\lambda}}{\partial y} (a, 0) - g_{\lambda}(a, 0) \frac{\sigma_{S\lambda}}{\partial y} (0, 0+) \end{bmatrix}$$
for $n \ge 0$.

Therefore we get

$$E_{x}^{0}[e^{-\lambda\sigma_{0}};\Omega_{1}^{+}] = \frac{\sigma(0)^{2}}{2g_{\lambda}(0>0)} \left[g_{\lambda}(0,0) \frac{\partial g_{\lambda}}{\partial y}(x,0) - g_{\lambda}(x,0) \frac{\partial g_{\lambda}}{\partial y}(0,0+) \right].$$

By the same way, we have

$$E_x^0[e^{-\lambda\sigma_0}\,\Omega_1^-] = \frac{\sigma(0)^2}{2g_\lambda(0,\,0)} \Big[-g_\lambda(0,\,0)\frac{\partial g_\lambda}{\partial \nu}(x,\,0) + g_\lambda(x,\,0)\frac{\partial g_\lambda}{\partial y}(0,\,0-) \Big] \,.$$

This completes the proof of (1).

Proof of (2). Let us suppose the condition (A) of Lemma 5.5 satisfied. We can choose a constant $c \in \mathbb{R}^n \setminus \{0\}$ such that

$$-g_{\lambda}(0, \frac{\partial}{\partial y})\frac{\partial g_{\lambda}}{\partial y}(c, \frac{\partial}{\partial y})+ \mathcal{J}_{\lambda}(c, 0)\frac{\partial g_{\lambda}}{\partial y}(0, 0-) \neq 0.$$

Put $K(x) = \lim_{y \downarrow 0} \frac{g_{\lambda}^{0}(x, y)}{g_{\lambda}^{*}(c, y)} x \neq 0$. Then we have

$$K(x) = \frac{-g_{\lambda}(0, 0) \frac{\partial g_{\lambda}}{\partial y}(x, 0) + g_{\lambda}(x, 0) \frac{\partial g_{\lambda}}{\partial y}(0, 0-)}{-g_{\lambda}(0, 0) \frac{\partial g_{\lambda}}{\partial y}(c, 0) + g_{\lambda}(c, 0) \frac{\partial g_{\lambda}}{\partial y}(0, 0-)},$$

$$\lim_{x \neq 0} K(x) = \infty \quad \text{and} \quad \lim_{x \neq 0} K(x) = 0.$$

As before, $e^{-\lambda t}K(X_t^0)$ is a nonnegative supermartingale having a left limit.

²⁾ \mathcal{F}_t^0 is the σ -field generated by X_s^0 , $0 \leq s \leq t$.

Therefore we have $P_x^0(\lim_{t \neq \sigma_0} e^{-\lambda t} K(X_t^0) \operatorname{exists}/\Omega_1) = 1$ for $x \neq 0$. It follows that $P_x(\mathfrak{A}t_n \uparrow \sigma_0, X_{t_n} < 0/\Omega_1) = 0$ for $x \neq 0$. This implies that $P_x(\Omega_1^+/\Omega_1) = 1$ for $x \neq 0$.

Next suppose the condition (B) of Lemma 5.5 is fulfilled. We construct a finite measure μ on $(-\infty, 0)$ such that

$$U(x) = \int_{-\infty}^{0} \frac{g_{\lambda}^{0}(x, y)}{g_{\lambda}^{0}(c, y)} \mu(dy)$$

has the property

$$U(x) < \infty$$
 for $x \in \mathbb{R}^{1} \setminus \{0\}$ and $\lim_{x \neq 0} U(x) = \infty$.

Fix $c \in \mathbb{R}^n \setminus \{0\}$. Note that

$$(8.4) \quad \frac{f^{\alpha}(x, -\varepsilon)}{g^{\alpha}_{\lambda}(c, -\varepsilon)} = \frac{g_{\lambda}(0, 0)[g_{\lambda}(x, -\varepsilon) - g_{\lambda}(x, 0)] + g_{\lambda}(x, 0)[g_{\lambda}(0, 0) - g_{\lambda}(0, -\varepsilon)]}{g_{\lambda}(0, 0)[g_{\lambda}(c, -\varepsilon) - g_{\lambda}(c, 0)] + g_{\lambda}(c, 0)[g_{\lambda}(0, 0) - g_{\lambda}(0, -\varepsilon)]}.$$

Putting $y = x + \varepsilon$ and using the condition (B), we have

$$egin{aligned} g_{\lambda}(x, \ -arepsilon) &= g_{\lambda}(x, \ x) - g_{\lambda}(x, x + (arepsilon - y)) - [g_{\lambda}(x, x) - g_{\lambda}(x, x - y)] \ &\geq K arepsilon^{lpha} \Big(\log rac{1}{arepsilon} \Big)^{m{r}} & ext{for } 0 \leq y \leq a arepsilon, \ 0 \leq arepsilon \leq arepsilon_{0} \ , \end{aligned}$$

that is $g_{\lambda}(x, -\varepsilon) - g_{\lambda}(x, 0) \ge K\varepsilon^{a} \left(\log \frac{1}{c^{c}} \right)^{r}$ for $-\varepsilon \le x \le -(1-a)\varepsilon$, $0 \le \varepsilon \le \varepsilon_{0}$. Hence $g_{\lambda}^{0}(x, -\varepsilon) \ge K_{1}\varepsilon^{a} \left(\log \frac{1}{c^{c}} \right)^{r}$. By the same way, we have $g_{\lambda}^{0}(c, -\varepsilon) < K_{2}\varepsilon^{\beta}$. So we have

$$\frac{g_{\lambda}^{0}(x, -\varepsilon)}{g_{\lambda}^{0}(c, -\varepsilon)} > K'\varepsilon^{\alpha - \beta} \left(\log \frac{1}{\varepsilon}\right)^{r} \quad \text{for } -\varepsilon \leq x \leq -(1-a)\varepsilon \quad \text{and} \quad 0 \leq \varepsilon \leq \varepsilon_{0} \,.$$

We conclude from this that there exists a constant K''>0 such that

$$\frac{g_{\lambda}^{0}(x, x_{n})}{g_{\lambda}^{0}(c, x_{n})} > K''(1-a)^{n(\omega-\beta)} \left(\log \frac{1}{(1-a)^{n}}\right)^{r} \quad \text{for } x_{n} \leq x \leq x_{n+1}, n = 1, 2, \dots,$$

where $x_n = -\mathcal{E}_0(1-a)^n$. Choose a constant b such that 0 < b < 1 and $(1-a)^{\alpha-\beta}b > 1$. Define a finite measure μ on $(-\infty, 0)$ by $\mu = \sum_{n=1}^{\infty} b^n \delta_{x_n}$. Because

$$\lim_{\varepsilon\to 0}\frac{g_{\lambda}^0(x,\,-\varepsilon)}{g_{\lambda}^0(c,\,-\varepsilon)}=\frac{g_{\lambda}(x,\,0)}{g_{\lambda}(c,\,\mathbf{o})},$$

we get

$$U(x) = \int_{-\infty}^{0} \frac{g_{\lambda}^{0}(x, y)}{g_{\lambda}^{0}(c, y)} \mu(dy) < +\infty \quad \text{for any } x \neq 0.$$

HITTING PROPERTIES

Since $U(x) \ge \frac{g_{\lambda}^{0}(xx_{n})}{g_{\lambda}^{0}(c, x_{n})} b^{n} = K''(1-a)^{n(o-\beta)} \left(\log - \frac{1}{(1-a)^{n}} \right)^{r} b^{n}$ for $x \in [x_{n}, x_{n+1}]$, we have $\lim_{x \neq 0} U(x) = \infty$. Since $e^{-\lambda t} U(X_{t}^{0})$ is a nonnegative supermartingale having a left limit. Therefore as before, we have $P_{x}(\exists t_{n} \uparrow \sigma_{0}, X_{t_{n}} < 0/\Omega_{1}) = 0$ for $x \neq 0$. Thus the proof of (2) is completed.

Proof of (3). First we show $P_x(\Omega_1^+/\Omega_1)=0$ or 1. Define $f_t(\omega)$ by $f_t(\omega)=e^{-\lambda t}E^0_{X_1^0(\omega)}[e^{-\lambda\sigma_0};\Omega_1^+]$. Let τ_n be as in the proof of (1). Since $\{f_{\tau_n}\}_{n\geq 0}$ is a bounded nonnegative $\mathcal{F}^0_{\tau_n}$ -martingale, it follows that there exists $af_{\infty}(\omega)=\lim_{n\to\infty}f_{\tau_n}(\omega)$ such that

$$f_{\tau_n}(\omega) = E^{\mathbf{0}}_{\mathbf{x}}[f_{\infty}(\omega)/\mathcal{F}^{\mathbf{0}}_{\tau_n}], \qquad n \ge 0.$$

On the other hand, by the strong Markov property,

$$E^0_{X^0_{ au_{\pi}}}[e^{-\lambda\sigma_0};\Omega^+_1]=e^{+\lambda au_{\pi}}E^0_x[e^{-\lambda\sigma_0}I_{\Omega^+_1}(\omega)/\mathcal{F}_{\pi}] \quad ext{on} \quad t\!<\!\sigma_0.$$

So we have

$$\lim_{n\to\infty} E^{\,0}_{X^0_{\tau_n}}[e^{-\lambda\sigma_0};\,\Omega^+_1] = I_{\Omega^+_1}(\omega)$$

Hence in view of Lemma 7.6, we have

$$f_{\infty}(\omega) = \begin{cases} e^{-\lambda_{\sigma_0}}, & \text{if } \omega \in \Omega_1^+, \\ 0, & \text{if } \omega \in \Omega_1^+, \end{cases}$$

and in view of Lemma 7.6, we have $f_{\infty}(\omega) = e^{-\lambda\sigma_0}c_1$. This implies that $P_x(\Omega_1^+/\Omega_1) = 1$ or $P_x(\Omega_1^+/\Omega_1) = 0$. By the same way, we have $P_x(\Omega_1^-/\Omega_1) = 1$ or 0 and $P_x(\Omega_1^+/\Omega_1) = 1$ or 0. Finally we show that $P_x(\Omega_1^+/\Omega_1) = 0$ and $P_x(\Omega_1^-/\Omega_1) = 0$. Let $\sigma_{-z} = \inf \{t; X_t = -\varepsilon\}$. For any $x \in \mathbb{R}^n \setminus \{0\}$,

$$P_x(\sigma_{-\varepsilon} < \sigma_0) \ge E_x[e^{-\lambda \sigma_{-\varepsilon}}; \sigma_{-\varepsilon} < \sigma_0] = \frac{g_{\lambda}^0(x, -\varepsilon)}{g_{\lambda}^0(-\varepsilon, -\varepsilon)}.$$

It follows from Lemma 5.6 that $\lim_{\epsilon \downarrow 0} P_x(\sigma_{-\epsilon} < \sigma_0) > 0$. Hence there exists a sequence $\{\mathcal{E}_n\}$ with $\mathcal{E}_n \downarrow 0$ such that

$$P_x(\overline{\lim_{n\to\infty}} \sigma_{-\varepsilon_n} < \sigma_0) > 0$$
.

Set $T_n = \sigma_{-\mathfrak{e}_n} \wedge \sigma_{-\mathfrak{e}_{n+1}} \wedge \cdots, n = 1, 2, \cdots$. Then T_n is increasing in n and $\lim_{n \to \infty} \sigma_0 \wedge T_n = \sigma_0 a.s.$ on $\{\sigma_0 < \infty\}$. So we get

$$\overline{\lim_{n \to \infty}} \{ \sigma_{-\mathfrak{r}_n} < \sigma_0 \} - \{ T_n < \sigma_0 \text{ for any } n \}.$$

Therefore

$$P_{\mathbf{x}}(T_{\mathbf{n}} < \sigma_{0}, T_{\mathbf{n}} t \sigma_{0}) = P_{\mathbf{x}}(\lim_{n \to \infty} \{\sigma_{-\mathfrak{e}_{n}} < \sigma_{0}\}, T_{\mathbf{n}} \uparrow \sigma_{0}) > 0.$$

It follows that $P_x(\Omega_1^+/\Omega_1) \neq 1$. So $P_x(\Omega_1^+/\Omega_1) = 0$. Similarly we can show that $P_x(\Omega^-/\Omega_1) = 0$. Therefore we have $P_x(\Omega_1^\pm/\Omega_1) = 1$. The proof of Theorem 2 is completed.

OSAKA UNIVERSITY

References

- [1] R.M. Blumenthal and R.K. Getoor: Markov Processes and Potential Theory, New York, 1968.
- [2] L. Hörmander: Pseudo-differentialoperators and hypoellipitc equations, Proc. Symposia in Pure Math. 10, A.M.S. 1968, 138-183.
- [3] N. Ikeda and S. Watanabe: *The local structure of a class of diffusions and related problems,* Proc. of the 2nd Japan-USSR Symp. on Probability Theory, 1972 in Lecture Notes in Math. Vol. 330.
- [4] M. Kanda: Comparison theorems on regular points for multidimensional Markov processes of transient type, Nagoya Math. J. 44 (1971), 165-214.
- [5] H. Kesten: *Hitting probabilities of single points for processes with stationary independent increments*, Mem. Amer. Math. Soc. 93 (1969).
- [6] H. Kumano-go: Algebra of pseudo-differential operators, J. Fac. Sci. Univ. Tokyo 1A, 17 (1970), 31-50.
- [7] H. Kumano-go: Pseudo-differential Operators, Seminar Note No. 25, Tokyo Univ. (in Japanese), 1970.
- [8] H. Kumano-go: On the index of hypoelliptic pseudo-differential operators on Rⁿ, Proc. Japan Acad. 48 (1972), 402-407.
- [9] H. Kunita and T. Watanabe: Markov processes and Martin boundaries, I, Illinois J. Math. 9 (1965), 458-526.
- [10] P.W. Millar: Exit properties of stochastic processes with stationary independent increments, Trans. Amer. Math. Soc. 178 (1973), 459–479.
- [11] K. Sato: Integration of Kolmogorov-Feller backward equations, J. Fac. Univ. Tokyo 1A, 9 (1961-63), 13-27.
- [12] K. Sato and T. Ueno: Multi-dimensional diffusion and the Markov process on the boundary, J. Math. Kyoto Univ. 4 (1965), 529-605.
- [13] L. Schwartz: Theorie des noyaux, Proc. of International Congress (1950), vol. 1, 220-230.
- [14] L. Schwartz: Theorie des Distributions, 3rd ed., Paris, 1966.
- [15] T. Takada: On potential densities of one-dimensional Levy processes, J. Math. Kyoto Univ. 14 (1974), 371-390.
- [16] S. Watanabe: On discontinuous additive functionals and Levy measures of a Markov process, Japan. J. Math. 34 (1964), 53-70.