

Title	Λ -L Λ -L-Bis[(μ -L-cysteinato)-1:2 κ^3 S:N,S;1:3 κ^3 S:N,S-tetrakis(ethylenediamine)-2 κ^4 N,N';3 κ^4 N,N'-dicobalt(III)silver(I) Tris(perchlorate) Pentahydrate
Author(s)	Tamura, Motoshi; Yoshinari, Nobuto; Igashira, Asako et al.
Citation	Acta Crystallographica Section E. 2007, E63(6), p. m1641-m1642
Version Type	VoR
URL	https://hdl.handle.net/11094/54378
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University



$\Lambda_L\Lambda_L$ -Bis(μ -L-cysteinato)-1:2 κ^3 S:N,S;1:3 κ^3 S:N,S-tetra-
kis(ethylenediamine)-2 κ^4 N,N';3 κ^4 N,N'-dicobalt(III)silver(I)
tris(perchlorate) pentahydrate

Motoshi Tamura, Nobuto Yoshinari, Asako Igashira-Kamiyama and Takumi Konno

Acta Cryst. (2007). E63, m1641–m1642



IUCr Journals

CRYSTALLOGRAPHY JOURNALS ONLINE

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Reproduction of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>

$\Lambda_L\Lambda_L$ -Bis(μ -L-cysteinato)-1:2 κ^3 S:N,S;1:-3 κ^3 S:N,S-tetrakis(ethylenediamine)-2 κ^4 N,N';3 κ^4 N,N'-dicobalt(III)silver(I) tris(perchlorate) pentahydrate

Motoshi Tamura, Nobuto Yoshinari, Asako Igashira-Kamiyama* and Takumi Konno

Department of Chemistry, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

Correspondence e-mail: igashira@ch.wani.osaka-u.ac.jp


Received 20 April 2007; accepted 24 April 2007

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; disorder in solvent or counterion; R factor = 0.023; wR factor = 0.060; data-to-parameter ratio = 7.3.

In the crystal structure of the title compound, $\Lambda_L\Lambda_L$ -[AgCo₂-(C₃H₅NO₂S)₂(C₂H₈N₂)₄](ClO₄)₃·5H₂O, the Ag^I atom, which lies on a twofold rotation axis, is linearly coordinated by two thiolate S atoms from two Λ_L -[Co(C₃H₅NO₂S)(C₂H₈N₂)₂]⁺ octahedral units, forming an S-bridged Co^{III}-Ag^I-Co^{III} trinuclear unit. The compound has two uncoordinated carboxylate groups.

Related literature

For related literature, see: Konno (2004); Konno *et al.* (2001).



Experimental

Crystal data

[AgCo₂(C₃H₅NO₂S)₂(C₂H₈N₂)₄]-
(ClO₄)₃·5H₂O
 $M_r = 1092.86$
Monoclinic, $C2$
 $a = 16.542$ (2) Å
 $b = 9.050$ (2) Å
 $c = 13.728$ (2) Å

$\beta = 111.507$ (10)^o
 $V = 1912.0$ (6) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 1.77$ mm⁻¹
 $T = 296$ (2) K
 $0.40 \times 0.18 \times 0.18$ mm

Data collection

Rigaku AFC-7S diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.537$, $T_{\max} = 0.741$
2426 measured reflections
2346 independent reflections

2219 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$
3 standard reflections
every 150 reflections
intensity decay: 14.4%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.060$
 $S = 1.05$
2346 reflections
322 parameters
13 restraints

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.37$ e Å⁻³
 $\Delta\rho_{\min} = -0.25$ e Å⁻³
Absolute structure: Flack (1983)
Flack parameter: 0.001 (18)

Table 1

Selected geometric parameters (Å, °).

Ag1—S1	2.3938 (9)	Co1—N5	1.975 (3)
Co1—N3	1.960 (3)	Co1—N4	1.980 (3)
Co1—N2	1.964 (3)	Co1—S1	2.2642 (10)
Co1—N1	1.972 (3)		
S1 ⁱ —Ag1—S1	172.38 (5)	N1—Co1—N5	175.53 (14)
N3—Co1—N2	175.89 (16)	N4—Co1—S1	177.46 (10)

Symmetry code: (i) $-x, y, -z$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O12 ⁱⁱ	0.90	2.30	3.14 (2)	155
N1—H2 \cdots O11	0.90	2.19	3.034 (19)	157
N1—H2 \cdots O7	0.90	2.23	3.066 (18)	155
N2—H3 \cdots O15	0.90	2.04	2.899 (5)	160
N2—H4 \cdots O13 ⁱⁱⁱ	0.90	2.28	3.057 (19)	144
N3—H5 \cdots O4	0.90	2.25	3.146 (13)	174
N3—H6 \cdots O10 ^{iv}	0.90	2.21	3.083 (12)	162
N4—H7 \cdots O2 ^v	0.90	2.03	2.868 (4)	155
N4—H8 \cdots O7	0.90	2.49	3.117 (16)	128
N4—H8 \cdots O17 ^v	0.90	2.50	3.138 (13)	128
N5—H9 \cdots O15	0.90	2.36	3.201 (5)	155
N5—H10 \cdots O16	0.90	2.25	3.079 (6)	152
O15—H22 \cdots O1 ^v	0.85 (5)	1.89 (3)	2.711 (5)	162 (7)
O16—H26 \cdots O2 ^v	0.86 (10)	2.12 (4)	2.891 (7)	149 (7)
O17—H27 \cdots O1 ^{vi}	0.85 (2)	2.2 (2)	2.822 (12)	126 (19)
O17—H28 \cdots O1	0.85 (2)	2.12 (12)	2.884 (12)	149 (19)

Symmetry codes: (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z$; (iii) $x, y - 1, z$; (iv) $x - \frac{1}{2}, y - \frac{1}{2}, z$; (v) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + 1$; (vi) $-x, y, -z + 1$.

Data collection: *MSC/AFC Diffractometer Control Software* (Molecular Structure Corporation, 1992); cell refinement: *MSC/AFC Diffractometer Control Software*; data reduction: *CrystalStructure* (Rigaku/MSC, 2004); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2256).

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Konno, T. (2004). *Bull. Chem. Soc. Jpn.*, **77**, 627–649.
- Konno, T., Yoshimura, T., Aoki, K., Okamoto, K. & Hirotsu, M. (2001). *Angew. Chem. Int. Ed.* **40**, 1765–1768.
- Molecular Structure Corporation (1992). *MSC/AFC Diffractometer Control Software*. MSC, The Woodlands, Texas, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Rigaku/MS (2004). *CrystalStructure*. Version 3.6.0. Rigaku/MS, The Woodlands, Texas, USA.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.

supporting information

Acta Cryst. (2007). E63, m1641–m1642 [doi:10.1107/S1600536807020442]

$\Lambda_L\Lambda_L$ -Bis(μ -L-cysteinato)-1:2 κ^3 S:N,S;1:3 κ^3 S:N,S-tetrakis(ethylenediamine)-2 κ^4 N,N';3 κ^4 N,N'-dicobalt(III)silver(I) tris(perchlorate) pentahydrate

Motoshi Tamura, Nobuto Yoshinari, Asako Igashira-Kamiyama and Takumi Konno

S1. Comment

Thiolato groups coordinated to a metal center possess relatively strong Lewis basicity, which allows them to bind with a second metal center (Konno, 2004). Previously, Konno *et al.* (2001) reported Λ_L -[Co(*L*-cys-*N,S*)(en)₂]⁺ (*L*-cys = *L*-cysteinate, en = ethylenediamine) reacts with AgNO₃ to give an S-bridged (Co^{III}Ag^I)_n coordination polymer, { Λ_L -[Ag{Co(*L*-cys-*N,S*)(en)₂}(NO₃)₂]_n·H₂O (II)}. In this compound, the Λ_L -[Co(*L*-cys-*N,S*)(en)₂]⁺ unit binds with two Ag^I atoms through sulfur and with another Ag^I atom through a carboxylate group, to give a sheet-like structure. In this paper, we report on the structure of $\Lambda_L\Lambda_L$ -[Ag{Co(*L*-cys-*N,S*)(en)₂]₂(ClO₄)₃·5H₂O (I), which was obtained by the 2:1 reaction of Λ_L -[Co(*L*-cys-*N,S*)(en)₂]⁺ with AgClO₄.

The cation is composed of two octahedral [Co(*L*-cys-*N,S*)(en)₂]⁺ units that are linked by an Ag^I atom through the S atoms to form a linear-type S-bridged Co^{III}Ag^ICo^{III} trinuclear structure in [Ag{Co(*L*-cys-*N,S*)(en)₂]₂³⁺ (Fig. 1). The Ag^I atom, which is located on a twofold axis, adopts an almost linear coordination geometry, unlike an angular geometry found in (II) (S—Ag—S = 149.62 (7) °). Furthermore, the Ag—S bonds in (I) are appreciably shorter than those in (II) (Ag—S = 2.501 (3), 2.511 (3) Å). These differences are ascribed to the fact that the Ag^I center in (II) is coordinated by a carboxylate group (Ag—O = 2.598 (1), 2.490 (8) Å), besides two thiolato groups. Other bond distances and angles in (I) are similar to those in (II) (Table 1). The two [Co(*L*-cys-*N,S*)(en)₂]⁺ units in (I) have an Λ configuration because of the configuration of the mononuclear Λ_L -[Co(*L*-cys-*N,S*)(en)₂]⁺ reactant. The *L*-cys *N,S*-chelate ring adopts a λ conformation; the two en *N,N*-chelate rings adopt δ and λ conformations.

The cation is connected with to four adjacent cations through N—H···O hydrogen bonds between coordinated amine groups and non-coordinated carboxylate groups to give a sheet-like structure (Fig. 2). The sheets are further linked through the perchlorate anions and water molecules through hydrogen bonds.

S2. Experimental

Treatment of Λ_L -[Co(*L*-Hcys-*N,S*)(en)₂](ClO₄)₂ with a mixture of 0.5 molar equiv. of AgClO₄ and 0.5 equiv. of NaOH in water at room temperature gave a red solution, from which red crystals (I) were isolated by adding an aqueous solution of NaClO₄.

S3. Refinement

H atoms bonded to C and N atoms were placed at calculated positions [C—H = 0.97 (methylene) and 0.98 (methine) Å, and N—H = 0.90 Å] and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. H atoms of water molecules were found in a difference Fourier map and were refined with restrained geometrical parameters [O—H = 0.85 (2) Å, H···H = 1.38 (2) Å, and $U_{\text{iso}} = 1.5U_{\text{eq}}(\text{O})$]. One H atom of one water molecule is disordered over two positions (H23/H24), which were refined with site occupancies of 0.5. Atom H26 of a water molecule was refined with a restrained geometrical parameter




Figure 2

A view of the two-dimensional sheet like structure in (I). Dashed lines indicate N–H···O hydrogen bonds. Perchlorate anions and water molecules have been omitted.

$\Lambda_1\Lambda_1$ —Bis(μ -L-cysteinato)- 1:2 κ^3 S:N,S;1:3 κ^3 S:N,S- tetrakis(ethylenediamine)-2 κ^4 N,N';3 κ^4 N,N'-dicobalt(III)silver(I) tris(perchlorate) pentahydrate

Crystal data

[AgCo₂(C₃H₅NO₂S)₂(C₂H₈N₂)₄](ClO₄)₃·5H₂O

$M_r = 1092.86$

Monoclinic, $C2$

Hall symbol: $C\ 2y$

$a = 16.542\ (2)\ \text{\AA}$

$b = 9.050\ (2)\ \text{\AA}$

$c = 13.728\ (2)\ \text{\AA}$

$\beta = 111.507\ (10)^\circ$

$V = 1912.0\ (6)\ \text{\AA}^3$

$Z = 2$

$F(000) = 1116$

$D_x = 1.898\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71069\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 14.6\text{--}15.0^\circ$

$\mu = 1.77\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Rod, dark red

$0.40 \times 0.18 \times 0.18\ \text{mm}$

Data collection

Rigaku AFC-7S

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω - 2θ scans

Absorption correction: ψ scan

(North *et al.*, 1968)

$T_{\min} = 0.537$, $T_{\max} = 0.741$

2426 measured reflections

2346 independent reflections

2219 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.015$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.5^\circ$

$h = 0 \rightarrow 21$

$k = 0 \rightarrow 11$

$l = -17 \rightarrow 16$

3 standard reflections every 150 reflections

intensity decay: 14.4%

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.060$
 $S = 1.05$
 2346 reflections
 322 parameters
 13 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0348P)^2 + 0.6577P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.37 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$
 Extinction correction: *SHELXL97*,
 $\text{Fc}^* = k\text{Fc}[1 + 0.001x\text{Fc}^2\lambda^3/\sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0150 (5)
 Absolute structure: Flack (1983)
 Absolute structure parameter: 0.001 (18)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ag1	0.0000	0.31917 (5)	0.0000	0.04457 (14)	
Co1	0.21020 (3)	0.48714 (5)	0.22914 (3)	0.02666 (12)	
S1	0.14873 (5)	0.30161 (10)	0.11542 (6)	0.03258 (19)	
O1	0.0811 (2)	0.0540 (4)	0.3873 (3)	0.0581 (9)	
O2	0.22302 (18)	0.1037 (4)	0.4561 (2)	0.0458 (7)	
N1	0.2271 (2)	0.6004 (4)	0.1154 (2)	0.0395 (7)	
H1	0.1798	0.5912	0.0564	0.047*	
H2	0.2340	0.6967	0.1327	0.047*	
N2	0.3263 (2)	0.4071 (4)	0.2530 (3)	0.0396 (7)	
H3	0.3576	0.4039	0.3222	0.048*	
H4	0.3216	0.3145	0.2277	0.048*	
N3	0.09835 (19)	0.5808 (4)	0.2088 (3)	0.0383 (7)	
H5	0.0783	0.6268	0.1463	0.046*	
H6	0.0595	0.5112	0.2089	0.046*	
N4	0.2594 (2)	0.6494 (4)	0.3304 (2)	0.0364 (6)	
H7	0.2765	0.6141	0.3961	0.044*	
H8	0.3061	0.6878	0.3207	0.044*	
N5	0.1989 (2)	0.3614 (3)	0.3410 (2)	0.0322 (6)	
H9	0.2515	0.3238	0.3787	0.039*	
H10	0.1825	0.4186	0.3842	0.039*	
C1	0.1504 (3)	0.1644 (4)	0.2126 (3)	0.0372 (8)	
H11	0.1051	0.0919	0.1813	0.045*	

H12	0.2059	0.1135	0.2369	0.045*	
C2	0.1364 (2)	0.2377 (4)	0.3044 (3)	0.0317 (7)	
H13	0.0771	0.2771	0.2810	0.038*	
C3	0.1481 (2)	0.1233 (4)	0.3914 (3)	0.0361 (8)	
C4	0.3050 (3)	0.5450 (7)	0.0968 (4)	0.0573 (13)	
H14	0.2897	0.4603	0.0502	0.069*	
H15	0.3280	0.6216	0.0648	0.069*	
C5	0.3707 (3)	0.5023 (7)	0.2000 (4)	0.0551 (12)	
H16	0.4181	0.4487	0.1904	0.066*	
H17	0.3943	0.5896	0.2417	0.066*	
C6	0.1082 (3)	0.6900 (5)	0.2938 (4)	0.0477 (10)	
H18	0.0610	0.7610	0.2716	0.057*	
H19	0.1078	0.6402	0.3562	0.057*	
C7	0.1931 (3)	0.7657 (4)	0.3154 (4)	0.0456 (10)	
H20	0.1900	0.8290	0.2571	0.055*	
H21	0.2078	0.8261	0.3779	0.055*	
C11	0.0000	0.86644 (15)	0.0000	0.0397 (3)	
O3	0.0194 (6)	0.8851 (10)	0.1120 (5)	0.070 (2)	0.50
O4	0.0409 (7)	0.7347 (13)	-0.0117 (9)	0.112 (4)	0.50
O5	-0.0919 (4)	0.8481 (10)	-0.0499 (6)	0.067 (2)	0.50
O6	0.0282 (5)	0.9912 (12)	-0.0383 (7)	0.079 (2)	0.50
C12	0.37527 (6)	0.98034 (14)	0.19897 (9)	0.0495 (2)	
O7	0.3094 (11)	0.898 (2)	0.2056 (16)	0.115 (6)	0.50
O8	0.3534 (12)	1.1056 (13)	0.1333 (12)	0.121 (6)	0.50
O9	0.4166 (17)	1.056 (3)	0.2960 (10)	0.168 (10)	0.50
O10	0.4352 (8)	0.8937 (16)	0.1788 (16)	0.097 (6)	0.50
O11	0.2889 (9)	0.919 (2)	0.1446 (11)	0.084 (4)	0.50
O12	0.3949 (13)	1.028 (5)	0.1154 (12)	0.218 (17)	0.50
O13	0.3750 (11)	1.080 (2)	0.2633 (16)	0.159 (10)	0.50
O14	0.4421 (13)	0.887 (2)	0.248 (2)	0.193 (13)	0.50
O15	0.4028 (2)	0.3330 (6)	0.4736 (3)	0.0604 (9)	
H22	0.403 (4)	0.389 (6)	0.524 (4)	0.091*	
H23	0.419 (11)	0.247 (6)	0.497 (7)	0.091*	0.50
H24	0.454 (3)	0.309 (17)	0.482 (8)	0.091*	0.50
O16	0.1116 (4)	0.4563 (10)	0.4944 (4)	0.115 (2)	
H25	0.083 (6)	0.438 (14)	0.534 (6)	0.173*	
H26	0.156 (4)	0.510 (12)	0.530 (7)	0.173*	
O17	0.0413 (7)	0.2037 (15)	0.5497 (11)	0.096 (3)	0.50
H27	-0.013 (3)	0.20 (2)	0.538 (15)	0.144*	0.50
H28	0.053 (12)	0.131 (15)	0.517 (16)	0.144*	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.0396 (2)	0.0497 (3)	0.0313 (2)	0.000	-0.00251 (16)	0.000
Co1	0.0272 (2)	0.0267 (2)	0.0234 (2)	0.00172 (18)	0.00598 (16)	-0.00014 (18)
S1	0.0351 (4)	0.0334 (4)	0.0248 (4)	-0.0001 (4)	0.0058 (3)	-0.0024 (4)
O1	0.0459 (17)	0.067 (2)	0.064 (2)	-0.0068 (15)	0.0227 (15)	0.0198 (17)

O2	0.0473 (15)	0.0469 (16)	0.0363 (13)	0.0033 (14)	0.0072 (11)	0.0092 (13)
N1	0.0432 (16)	0.0415 (17)	0.0322 (14)	-0.0023 (15)	0.0119 (13)	0.0047 (14)
N2	0.0345 (15)	0.0474 (19)	0.0366 (16)	0.0048 (14)	0.0124 (13)	0.0015 (15)
N3	0.0332 (15)	0.0337 (15)	0.0421 (17)	0.0054 (13)	0.0068 (13)	-0.0008 (14)
N4	0.0412 (16)	0.0320 (15)	0.0321 (14)	-0.0028 (13)	0.0086 (12)	-0.0044 (13)
N5	0.0379 (15)	0.0315 (15)	0.0264 (13)	0.0027 (12)	0.0109 (12)	-0.0027 (11)
C1	0.044 (2)	0.0312 (18)	0.0295 (17)	0.0022 (16)	0.0056 (15)	0.0003 (15)
C2	0.0305 (17)	0.0318 (17)	0.0316 (17)	0.0029 (14)	0.0100 (14)	0.0057 (14)
C3	0.0417 (18)	0.031 (2)	0.0365 (17)	0.0003 (16)	0.0159 (15)	0.0012 (15)
C4	0.062 (3)	0.072 (3)	0.050 (2)	0.008 (2)	0.035 (2)	0.016 (2)
C5	0.038 (2)	0.074 (3)	0.060 (3)	0.001 (2)	0.0259 (19)	0.007 (3)
C6	0.048 (2)	0.042 (2)	0.054 (3)	0.0119 (19)	0.019 (2)	-0.0067 (19)
C7	0.059 (3)	0.0250 (17)	0.050 (2)	0.0016 (16)	0.016 (2)	-0.0096 (16)
C11	0.0393 (6)	0.0405 (7)	0.0334 (6)	0.000	0.0064 (5)	0.000
O3	0.093 (5)	0.078 (5)	0.034 (3)	-0.023 (5)	0.017 (3)	-0.007 (3)
O4	0.117 (9)	0.099 (7)	0.088 (7)	0.074 (6)	-0.002 (6)	-0.025 (6)
O5	0.038 (3)	0.080 (6)	0.069 (4)	-0.008 (4)	0.002 (3)	-0.008 (4)
O6	0.073 (5)	0.083 (6)	0.090 (6)	-0.017 (5)	0.039 (4)	0.026 (6)
C12	0.0404 (5)	0.0508 (6)	0.0594 (6)	-0.0013 (5)	0.0209 (4)	-0.0041 (5)
O7	0.107 (13)	0.077 (8)	0.215 (19)	-0.032 (9)	0.120 (14)	-0.020 (13)
O8	0.185 (17)	0.056 (6)	0.096 (10)	0.015 (7)	0.021 (9)	0.039 (6)
O9	0.25 (2)	0.159 (18)	0.050 (6)	-0.088 (18)	0.008 (11)	-0.023 (8)
O10	0.060 (7)	0.075 (7)	0.174 (16)	-0.003 (6)	0.063 (10)	-0.041 (10)
O11	0.054 (5)	0.083 (8)	0.108 (9)	-0.021 (5)	0.021 (6)	-0.006 (8)
O12	0.135 (15)	0.45 (5)	0.067 (7)	-0.13 (2)	0.039 (9)	0.004 (18)
O13	0.097 (9)	0.140 (13)	0.173 (19)	0.042 (9)	-0.028 (10)	-0.103 (14)
O14	0.112 (12)	0.132 (15)	0.26 (3)	0.089 (11)	-0.015 (15)	-0.002 (18)
O15	0.062 (2)	0.076 (3)	0.0410 (16)	-0.007 (2)	0.0156 (15)	-0.0118 (19)
O16	0.104 (4)	0.169 (7)	0.089 (3)	-0.027 (4)	0.054 (3)	-0.029 (4)
O17	0.083 (6)	0.113 (9)	0.109 (9)	-0.020 (6)	0.054 (6)	-0.028 (7)

Geometric parameters (Å, °)

Ag1—S1 ⁱ	2.3938 (9)	C2—H13	0.9800
Ag1—S1	2.3938 (9)	C4—C5	1.486 (7)
Co1—N3	1.960 (3)	C4—H14	0.9700
Co1—N2	1.964 (3)	C4—H15	0.9700
Co1—N1	1.972 (3)	C5—H16	0.9700
Co1—N5	1.975 (3)	C5—H17	0.9700
Co1—N4	1.980 (3)	C6—C7	1.492 (6)
Co1—S1	2.2642 (10)	C6—H18	0.9700
S1—C1	1.815 (4)	C6—H19	0.9700
O1—C3	1.256 (5)	C7—H20	0.9700
O2—C3	1.245 (5)	C7—H21	0.9700
N1—C4	1.490 (5)	C11—O6	1.396 (8)
N1—H1	0.9000	C11—O4	1.409 (8)
N1—H2	0.9000	C11—O5	1.430 (6)
N2—C5	1.486 (6)	C11—O3	1.462 (6)

N2—H3	0.9000	C12—O13	1.264 (15)
N2—H4	0.9000	C12—O7	1.351 (14)
N3—C6	1.492 (5)	C12—O14	1.359 (16)
N3—H5	0.9000	C12—O10	1.370 (11)
N3—H6	0.9000	C12—O12	1.372 (15)
N4—C7	1.479 (5)	C12—O8	1.410 (10)
N4—H7	0.9000	C12—O9	1.430 (15)
N4—H8	0.9000	C12—O11	1.459 (14)
N5—C2	1.482 (5)	O15—H22	0.85 (5)
N5—H9	0.9000	O15—H23	0.84 (2)
N5—H10	0.9000	O15—H24	0.84 (2)
C1—C2	1.516 (5)	O16—H25	0.85 (9)
C1—H11	0.9700	O16—H26	0.86 (10)
C1—H12	0.9700	O17—H27	0.85 (2)
C2—C3	1.538 (5)	O17—H28	0.85 (2)
S1 ⁱ —Ag1—S1	172.38 (5)	N5—C2—C1	108.2 (3)
N3—Co1—N2	175.89 (16)	N5—C2—C3	111.8 (3)
N3—Co1—N1	93.59 (14)	C1—C2—C3	109.7 (3)
N2—Co1—N1	84.87 (14)	N5—C2—H13	109.0
N3—Co1—N5	90.38 (13)	C1—C2—H13	109.0
N2—Co1—N5	91.31 (14)	C3—C2—H13	109.0
N1—Co1—N5	175.53 (14)	O2—C3—O1	126.4 (4)
N3—Co1—N4	84.43 (14)	O2—C3—C2	117.3 (3)
N2—Co1—N4	91.80 (14)	O1—C3—C2	116.3 (3)
N1—Co1—N4	91.68 (15)	C5—C4—N1	107.5 (4)
N5—Co1—N4	90.76 (13)	C5—C4—H14	110.2
N3—Co1—S1	93.46 (10)	N1—C4—H14	110.2
N2—Co1—S1	90.35 (11)	C5—C4—H15	110.2
N1—Co1—S1	89.86 (11)	N1—C4—H15	110.2
N5—Co1—S1	87.84 (9)	H14—C4—H15	108.5
N4—Co1—S1	177.46 (10)	N2—C5—C4	107.3 (4)
C1—S1—Co1	96.81 (12)	N2—C5—H16	110.3
C1—S1—Ag1	105.49 (13)	C4—C5—H16	110.3
Co1—S1—Ag1	120.00 (4)	N2—C5—H17	110.3
C4—N1—Co1	110.1 (3)	C4—C5—H17	110.3
C4—N1—H1	109.6	H16—C5—H17	108.5
Co1—N1—H1	109.6	C7—C6—N3	106.6 (3)
C4—N1—H2	109.6	C7—C6—H18	110.4
Co1—N1—H2	109.6	N3—C6—H18	110.4
H1—N1—H2	108.2	C7—C6—H19	110.4
C5—N2—Co1	109.5 (3)	N3—C6—H19	110.4
C5—N2—H3	109.8	H18—C6—H19	108.6
Co1—N2—H3	109.8	N4—C7—C6	107.2 (3)
C5—N2—H4	109.8	N4—C7—H20	110.3
Co1—N2—H4	109.8	C6—C7—H20	110.3
H3—N2—H4	108.2	N4—C7—H21	110.3
C6—N3—Co1	110.4 (2)	C6—C7—H21	110.3

C6—N3—H5	109.6	H20—C7—H21	108.5
Co1—N3—H5	109.6	O6—C11—O4	113.6 (7)
C6—N3—H6	109.6	O6—C11—O5	110.9 (5)
Co1—N3—H6	109.6	O4—C11—O5	108.5 (6)
H5—N3—H6	108.1	O6—C11—O3	109.1 (5)
C7—N4—Co1	109.7 (2)	O4—C11—O3	106.8 (6)
C7—N4—H7	109.7	O5—C11—O3	107.5 (5)
Co1—N4—H7	109.7	O13—C12—O14	107.9 (13)
C7—N4—H8	109.7	O7—C12—O10	111.0 (10)
Co1—N4—H8	109.7	O13—C12—O12	115 (2)
H7—N4—H8	108.2	O14—C12—O12	102.4 (17)
C2—N5—Co1	115.1 (2)	O7—C12—O8	117.5 (11)
C2—N5—H9	108.5	O10—C12—O8	111.5 (10)
Co1—N5—H9	108.5	O7—C12—O9	108.9 (12)
C2—N5—H10	108.5	O10—C12—O9	109.2 (13)
Co1—N5—H10	108.5	O8—C12—O9	97.6 (11)
H9—N5—H10	107.5	O13—C12—O11	112.1 (11)
C2—C1—S1	110.2 (3)	O14—C12—O11	118.9 (13)
C2—C1—H11	109.6	O12—C12—O11	100.4 (9)
S1—C1—H11	109.6	H22—O15—H23	109 (4)
C2—C1—H12	109.6	H22—O15—H24	109 (4)
S1—C1—H12	109.6	H25—O16—H26	108 (9)
H11—C1—H12	108.1	H27—O17—H28	108 (4)
N3—Co1—S1—C1	-98.88 (17)	N1—Co1—N4—C7	79.2 (3)
N2—Co1—S1—C1	82.67 (17)	N5—Co1—N4—C7	-104.6 (3)
N1—Co1—S1—C1	167.53 (16)	N3—Co1—N5—C2	75.3 (3)
N5—Co1—S1—C1	-8.63 (16)	N2—Co1—N5—C2	-108.4 (3)
N3—Co1—S1—Ag1	13.43 (11)	N4—Co1—N5—C2	159.7 (2)
N2—Co1—S1—Ag1	-165.03 (11)	S1—Co1—N5—C2	-18.1 (2)
N1—Co1—S1—Ag1	-80.16 (11)	Co1—S1—C1—C2	34.0 (3)
N5—Co1—S1—Ag1	103.68 (10)	Ag1—S1—C1—C2	-89.7 (3)
N3—Co1—N1—C4	-173.1 (3)	Co1—N5—C2—C1	44.4 (3)
N2—Co1—N1—C4	10.7 (3)	Co1—N5—C2—C3	165.4 (2)
N4—Co1—N1—C4	102.4 (3)	S1—C1—C2—N5	-51.5 (3)
S1—Co1—N1—C4	-79.7 (3)	S1—C1—C2—C3	-173.7 (3)
N1—Co1—N2—C5	16.7 (3)	N5—C2—C3—O2	-32.8 (5)
N5—Co1—N2—C5	-165.6 (3)	C1—C2—C3—O2	87.2 (4)
N4—Co1—N2—C5	-74.8 (3)	N5—C2—C3—O1	150.2 (4)
S1—Co1—N2—C5	106.5 (3)	C1—C2—C3—O1	-89.7 (4)
N1—Co1—N3—C6	-105.3 (3)	Co1—N1—C4—C5	-35.7 (5)
N5—Co1—N3—C6	76.8 (3)	Co1—N2—C5—C4	-40.6 (5)
N4—Co1—N3—C6	-14.0 (3)	N1—C4—C5—N2	49.3 (6)
S1—Co1—N3—C6	164.6 (3)	Co1—N3—C6—C7	38.9 (4)
N3—Co1—N4—C7	-14.3 (3)	Co1—N4—C7—C6	39.3 (4)
N2—Co1—N4—C7	164.1 (3)	N3—C6—C7—N4	-50.3 (5)

Symmetry code: (i) $-x, y, -z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1···O12 ⁱⁱ	0.90	2.30	3.14 (2)	155
N1—H2···O11	0.90	2.19	3.034 (19)	157
N1—H2···O7	0.90	2.23	3.066 (18)	155
N2—H3···O15	0.90	2.04	2.899 (5)	160
N2—H4···O13 ⁱⁱⁱ	0.90	2.28	3.057 (19)	144
N3—H5···O4	0.90	2.25	3.146 (13)	174
N3—H6···O10 ^{iv}	0.90	2.21	3.083 (12)	162
N4—H7···O2 ^v	0.90	2.03	2.868 (4)	155
N4—H8···O7	0.90	2.49	3.117 (16)	128
N4—H8···O17 ^v	0.90	2.50	3.138 (13)	128
N5—H9···O15	0.90	2.36	3.201 (5)	155
N5—H10···O16	0.90	2.25	3.079 (6)	152
O15—H22···O1 ^v	0.85 (5)	1.89 (3)	2.711 (5)	162 (7)
O15—H24···O15 ^{vi}	0.84 (2)	2.23 (5)	3.027 (8)	158 (12)
O16—H25···O17	0.85 (9)	2.27 (11)	2.794 (15)	120 (10)
O16—H26···O2 ^v	0.86 (10)	2.12 (4)	2.891 (7)	149 (7)
O17—H27···O1 ^{vii}	0.85 (2)	2.2 (2)	2.822 (12)	126 (19)
O17—H28···O1	0.85 (2)	2.12 (12)	2.884 (12)	149 (19)

Symmetry codes: (ii) $-x+1/2, y-1/2, -z$; (iii) $x, y-1, z$; (iv) $x-1/2, y-1/2, z$; (v) $-x+1/2, y+1/2, -z+1$; (vi) $-x+1, y, -z+1$; (vii) $-x, y, -z+1$.