

Title	A note on direct sums of cyclic modules over commutative regular rings
Author(s)	Oshiro, Kiyochi
Citation	Osaka Journal of Mathematics. 12(3) P.715-P.718
Issue Date	1975
Text Version	publisher
URL	<a href="https://doi.org/10.18910/5466">https://doi.org/10.18910/5466</a>
DOI	10.18910/5466
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## A NOTE ON DIRECT SUMS OF CYCLIC MODULES OVER COMMUTATIVE REGULAR RINGS

KIYOICHI OSHIRO

(Received December 18, 1974)

Throughout this note  $R$  is a commutative ring with identity and all modules are unital.

We denote the maximal ring of quotients of  $R$  by  $Q(R)$  and the ring generated by the set of all idempotents of  $Q(R)$  over  $R$  by  $C(R)$ . In case  $R$  is semi-prime,  $C(R)$  coincides with the Baer hull of  $R$  in the sense of A. C. Mewborn ([3, Proposition 2.5]). For an  $R$ -module  $M$ , we denote its injective hull by  $E_R(M)$ . It is well known (e.g. [1]) that if  $R$  is semi-prime, then  $Q(R) = E_R(R)$ .

Let  $M$  be an  $R$ -module. We put

$$\begin{aligned} T(M) &= \{x \in M \mid \text{Hom}_R(Rx, E_R(R)) = 0\} \\ &= \{x \in M \mid (0:x) \text{ is a dense}^\dagger \text{ ideal of } R\} \end{aligned}$$

where  $(0:x) = \{r \in R \mid rx = 0\}$  (see [7]).  $M$  is said to be torsion if  $T(M) = M$  and torsion free if  $T(M) = 0$ .

Now, for an  $R$ -module  $M$ , we shall consider the following condition studied in [4]:

(\*)  $M$  is embedded in a direct sum of cyclic  $R$ -modules as an essential  $R$ -submodule.

In [4] the author proved the following

**Theorem 1.** *Let  $R$  be a regular ring. Then the following conditions are equivalent:*

- (a)  $Q(R) = C(R)$ .
- (b) *Every finitely generated torsion free  $R$ -module  $M$  satisfies the condition (\*)*

The purpose of this note is to prove the following two theorems.

**Theorem 2.** *Let  $R$  be a regular ring. Then the following conditions are equivalent:*

- (a)  $Q(R/I) = C(R/I)$  for every dense ideal  $I$  of  $R$ .

---

<sup>†</sup> An ideal  $I$  of  $R$  is said to be dense provided that for any  $r$  and  $r'$  in  $R$  with  $r' \neq 0$ , there exists  $s$  in  $R$  such that  $sr \in I$  and  $sr' \neq 0$ .

(b) Every finitely generated torsion  $R$ -module  $M$  satisfies the condition (\*).

**Theorem 3.** Let  $R$  be a regular ring. Then the following conditions are equivalent:

- (a)  $Q(R/I)=C(R/I)$  for every ideal  $I$  of  $R$ .  
 (b) Every finitely generated  $R$ -module  $M$  satisfies the condition (\*).

To prove the theorems above, we use the following lemmas.

**Lemma 4.** For any  $R$ -module  $M$  with  $n$  generators, there exist  $n$  elements  $x_1, \dots, x_n$  in  $E_R(M)$  such that  $E_R(M)$  is a direct sum of  $E_R(Rx_1), \dots, E_R(Rx_n)$ . Moreover we can take  $Rx_1 + \dots + Rx_n$  to be torsion if  $M$  is torsion.

*Proof.* We proceed the proof by induction on the number  $n$  of the generators of  $M$ . If  $n=1$ , our assertion is obvious. We assume that the lemma is true for every  $R$ -module with  $n(<m)$  generators, and let us suppose that  $M$  has  $m$  generators, say  $M=Ra_1 + \dots + Ra_m$ .

By a usual property of injectivity of  $E_R(Ra_m)$ , we have that

$$E_R(Ra_m) + M = E_R(Ra_m) \oplus S \subseteq E_R(M)$$

for some submodule  $S$  of  $E_R(Ra_m) + M$ . Putting  $a_i = b_i + s_i$ ,  $b_i \in E_R(Ra_m)$ ,  $s_i \in S$ ,  $i=1, 2, \dots, m-1$ , we obtain

$$M + E_R(Ra_m) = (\sum_{i=1}^{m-1} Rs_i) \oplus E_R(Ra_m).$$

Hence by making use of our induction hypothesis on  $\sum_{i=1}^{m-1} Rs_i$ , there are  $m-1$  elements  $x_1, \dots, x_{m-1}$  in  $E_R(\sum_{i=1}^{m-1} Rs_i) \subseteq E_R(M)$  such that

$$E_R(\sum_{i=1}^{m-1} Rs_i) = E_R(Rx_1) \oplus \dots \oplus E_R(Rx_{m-1}).$$

Hence it follows that

$$E_R(M) = E_R(Rx_1) \oplus \dots \oplus E_R(Rx_{m-1}) \oplus E_R(Ra_m).$$

Moreover, if  $M$  is torsion, so is each  $Ra_i$  and hence each  $Rs_i$  is also torsion. This implies that  $(\sum_{i=1}^{m-1} Rx_i) + Ra_m$  is torsion.

The following lemma is due to R. S. Pierce [6, Corollary 23.7].

**Lemma 5.** Let  $R$  be a regular ring,  $I$  an ideal of  $R$  and  $M$  an  $R$ -module with  $IM=0$ . Then  $M$  is injective as an  $R$ -module if and only if  $M$  is injective as an  $R/I$ -module.

**REMARK.** Let  $R$  be a regular ring and  $I$  an ideal of  $R$ . If  $M$  is an  $R$ -module and  $IM=0$ , then it is easily seen that  $IE_R(M)=0$ . In particular we have that  $IE_R(R/I)=0$  and hence, by the lemma above,  $E_R(R/I)=E_{R/I}(R/I)$  ( $=Q(R/I)$ ).

Proof of Theorem 3. (a) $\Rightarrow$ (b). Let  $M=Ra_1+\dots+Ra_n$  be a finitely generated  $R$ -module. By Lemma 4 and the above remark, there are ideals  $I_1, \dots, I_n$  of  $R$  such that  $M$  is embedded in the external direct sum of  $Q(R/I_1), \dots, Q(R/I_n)$  as an essential submodule. Let us write

$$\begin{aligned} a_1 &= x_{11} \times \dots \times x_{1n}, \\ a_2 &= x_{21} \times \dots \times x_{2n}, \\ &\dots\dots\dots \\ a_n &= x_{n1} \times \dots \times x_{nn}, \end{aligned}$$

where  $x_{ij} \in Q(R/I_j)$ ,  $i, j=1, 2, \dots, n$ , and denote  $\sum_{i=1}^n Rx_{ij}$  by  $A_j, j=1, 2, \dots, n$ . Then  $A_j$  is a finitely generated  $R/I_j$ -submodule of  $Q(R/I_j), j=1, 2, \dots, n$  and  $M$  is embedded in  $A_1 \oplus \dots \oplus A_n$  as an essential  $R$ -submodule. Here, applying Theorem 1, each  $A_j$  satisfies the condition (\*) as an  $R/I_j$ -module and so does as an  $R$ -module. Hence  $M$  satisfies the condition (\*) as an  $R$ -module.

In the proof above, we can take each  $I_i$  to be dense ideal of  $R$  if  $M$  is torsion. Therefore this yields the proof of (a) $\Rightarrow$ (b) in Theorem 2 at the same time.

(b) $\Rightarrow$ (a). Let  $I$  be an ideal of  $R$ . By Theorem 1, to prove that  $Q(R/I)=C(R/I)$ , we may show that every finitely generated torsion free  $R/I$ -module satisfies the condition (\*) as an  $R/I$ -module. But this is evident, since every finitely generated torsion free  $R/I$ -module satisfies the condition (\*) as an  $R$ -module and so does as an  $R/I$ -module.

Note that if  $I$  is a dense ideal and  $M$  is an  $R/I$ -module, then  $M$  is torsion as an  $R$ -module since  $IM=0$ . Hence we also obtain the proof of (b) $\Rightarrow$ (a) in Theorem 2.

**Corollary 6.** *Let  $R$  be a regular ring such that  $Q(R/I)=C(R/I)$  for every dense ideal  $I$  of  $R$ . Then every finitely generated torsion injective  $R$ -module is a direct sum of cyclic  $R$ -modules.*

**Corollary 7.** *Let  $R$  be a regular ring such that  $Q(R/I)=C(R/I)$  for every ideal  $I$  of  $R$ . Then every finitely generated injective  $R$ -module is a direct sum of cyclic  $R$ -modules.*

Corollary 7 was shown by R. S. Pierce [6, Theorem 23.5] for the ring of all global sections of the simple  $F$ -sheaf over a Boolean space where  $F$  is a finite field. Let us note that a regular ring  $R$  is isomorphic to such a regular ring if and only if there exist finite elements, say  $r_1, \dots, r_n$ , in  $R$  with the property that all  $R/\mathfrak{m}, \mathfrak{m} \in \text{Spec}(R)$  are fields with just  $n$  elements  $r_1 + \mathfrak{m}, \dots, r_n + \mathfrak{m}$  and  $R/\mathfrak{n} \cong R/\mathfrak{n}'$  for any  $\mathfrak{n}, \mathfrak{n}'$  in  $\text{Spec}(R)$  by the canonical mapping:  $r_i + \mathfrak{n} \rightarrow r_i + \mathfrak{n}'$  (cf. [5, Proposition 2.1]). This class of regular rings contains Boolean rings and more generally  $p$ -rings in the sense of McCoy and Montgomery [2] ([6, p. 53]).

On the other hand, since this class of regular rings is clearly closed under homomorphic images, it is contained, by [5, Theorem 2.4], in the class of those regular rings  $R$  with  $Q(R/I)=C(R/I)$  for every ideal  $I$  of  $R$ . So, Corollary 7 can be seen as a generalization of the result due to R. S. Pierce.

YAMAGUCHI UNIVERSITY

---

### References

- [1] J. Lambek: *Lectures on Rings and Modules*. Blaisdell Publishing Co., Waltham, Mass., 1966.
- [2] N.H. McCoy and D. Montgomery: *A representation of generalized Boolean rings*, *Duke Math. J.* **3** (1937), 455–459.
- [3] A.C. Mewborn: *Regular rings and Baer rings*, *Math. Z.* **12** (1971), 211–219.
- [4] K. Oshiro: *On torsion free modules over regular rings*, *Math. J. Okayama Univ.* **16** (1973), 107–114.
- [5] K. Oshiro: *On torsion free modules over regular rings II*, *Math. J. Okayama Univ.* **16** (1974), 199–205.
- [6] R.S. Pierce: *Modules over Commutative Regular Rings*, *Memoirs Amer. Math. Soc.* **70**, 1967.
- [7] J. Zelmanowitz: *Injective hulls of torsion free modules*, *Canad. J. Math.* **23** (1971), 1094–1101.