



Title	On the loop-order of a fibre space
Author(s)	Furukawa, Yasukuni; Nomura, Yasutoshi
Citation	Osaka Journal of Mathematics. 1975, 12(3), p. 607-622
Version Type	VoR
URL	https://doi.org/10.18910/5538
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

ON THE LOOP-ORDER OF A FIBRE SPACE

Dedicated to Professor Ryoji Shizuma on his 60-th birthday

YASUKUNI FURUKAWA AND YASUTOSHI NOMURA

(Received January 20, 1975)

Introduction

Let ΩX denote the space of loops on a based topological space X . M. Sugawara [8] called the order of the identity class $1_{\Omega X}$ of ΩX in the group $[\Omega X, \Omega X]$ the loop-order of X , denoted by $l(X)$, and proved ([8], Theorem 3) that, for a Hurewicz fibration $F \rightarrow E \rightarrow B$, $l(E)$ is a divisor of the multiple $l(B) \cdot l(F)$.

The aim in this note is to determine, using a technique of Larmore and Thomas [2], the loop-order of a total space obtained as a 2-stage Postnikov tower and to discuss that of a space obtained as a 3-stage Postnikov tower.

In this note, let p denote a fixed prime. Let $\mathcal{A}(p)$ denote the mod p Steenrod algebra, and let $\varepsilon: \mathcal{A}(p) \rightarrow \mathcal{A}(p)$ denote the Kristensen map of degree -1 , which is a derivation and is given by

$$\begin{aligned} \varepsilon(Sq^n) &= Sq^{n-1} \quad (n \geq 1) & \text{if } p = 2, \\ \varepsilon(\Delta) &= 1, \quad \varepsilon(P^k) = 0 \quad (k \geq 0) & \text{if } p > 2, \end{aligned}$$

(cf. [2], Proposition 3.5; [5]). We shall write $\varepsilon(\alpha) = \bar{\alpha}$.

Also denote by $K_n = K(Z_p, n)$ the Eilenberg-MacLane complex of type (Z_p, n) . Let E_1 and E_2 be principal fibre spaces with classifying classes

$$\{\theta_1, \theta_2, \dots, \theta_m\}: K_n \rightarrow \bigotimes_{j=1}^m K_{n+r_j}, \quad 0 < r_1 \leq r_2 \leq \dots \leq r_m \leq n-3$$

and

$$\sum_{i=1}^k \pi_i^* \gamma_i: \bigotimes_{i=1}^k K_{n+s_i} \rightarrow K_{n+r}, \quad s_1 = 0 \leq s_2 \leq \dots \leq s_k < r \leq n-3$$

respectively, where θ_j and γ_i are cohomology operations of degree r_j and $r - s_i$, regarded as elements of $\mathcal{A}(p)$, and $\pi_i: \bigotimes_{t=1}^k K_{n+s_t} \rightarrow K_{n+s_i}$ is the projection on the i -th factor. We then obtain

Theorem A. $l(E_1) = p^2$ if, and only if, there exists j , $1 \leq j \leq m$, such that $\bar{\theta}_j$ does not belong to the left $\mathcal{A}(p)$ -module, $\sum_{i=1}^{j-1} \mathcal{A}(p)\theta_i$, of $\mathcal{A}(p)$ generated by

$\theta_1, \dots, \theta_{j-1}$.

Theorem B. $l(E_2)=p^2$ if, and only if, there exists i , $1 \leq i \leq k$, such that $\tilde{\gamma}_i$ does not belong to the right $\mathcal{A}(p)$ -module, $\sum_{i=i+1}^k \gamma_i \mathcal{A}(p)$, of $\mathcal{A}(p)$ generated by $\gamma_{i+1}, \dots, \gamma_k$.

The following corollary is a restatement of Theorem 1.3 of L. Smith [5].

Corollary 1. Let E be a fibre space induced from the path-fibration on K_{n+r} by $\theta = \theta \iota_n: K_n \rightarrow K_{n+r}$, where $0 < r \leq n-3$ and ι_n denotes the fundamental class. Then $l(E)$ is p^2 if, and only if, $\tilde{\theta} \neq 0$.

We next consider the situation shown in the diagram below:

$$\begin{array}{ccccc}
 \Omega L & \xrightarrow{j} & E & & \\
 & & \downarrow \pi & & \\
 \Omega B & \xrightarrow{l} & K & \xrightarrow{\theta} & L \\
 & & \downarrow \rho & & \\
 & & A & \xrightarrow{\alpha} & B
 \end{array}
 \quad (*)$$

where we set

$$\begin{aligned}
 A &= K_n, \quad B = \bigtimes_{i=1}^m K_{n+r_i}, \quad L = K_{n+s}, \quad 0 < r_1 \leq r_2 \leq \dots \leq r_m \leq s \leq n-3, \\
 \alpha &= \{\alpha_1, \dots, \alpha_m\}, \quad \alpha_i \in \mathcal{A}(p), \quad \deg \alpha_i = r_i, \\
 \beta &= \theta l = \sum_{i=1}^m (\Omega \pi_i)^* \beta_i, \quad \beta_i \in \mathcal{A}(p), \quad \deg \beta_i = s - r_i + 1,
 \end{aligned}$$

and where K and E are principal fibre spaces with classifying classes α and θ . Let

$$\psi: \bigcap_{i=1}^m (\text{Ker } \alpha_i \cap \text{Ker } \tilde{\alpha}_i) \rightarrow \text{Coker } \sum_{i=1}^m (\beta_i + \tilde{\beta}_i)$$

denote a secondary operation associated with the relation $\sum_{i=1}^m [\tilde{\beta}_i \alpha_i + (-1)^{s-r_i+1} \beta_i \tilde{\alpha}_i] = 0$, which is deduced from $\sum_{i=1}^m \beta_i \alpha_i = 0$ by taking the map ε .

Theorem C. Suppose that, for all $i=1, \dots, m$, $\tilde{\alpha}_i \in \sum_{k=1}^{i-1} \mathcal{A}(p) \alpha_k$.

- 1) If there exists j such that $\tilde{\beta}_j \notin \sum_{k=j+1}^m \beta_k \mathcal{A}(p)$, then $l(E) = p^2$.
- 2) If $\deg \beta_m > 1$ (i.e., $s > r_m$) and if

$$\psi(\Omega\rho) \equiv 0 \bmod \sum_{i=1}^m [\beta_i H^{n+r_i-3}(\Omega K; Z_p) + \tilde{\beta}_i H^{n+r_i-2}(\Omega K; Z_p)] \\ + (\Omega\rho)^* H^{n+s-2}(\Omega A; Z_p),$$

then $l(E)=p^2$.

3) If for all $i=1, \dots, m$, $\tilde{\beta}_i \in \sum_{k=j+1}^m \beta_k \mathcal{A}(p)$, and if $\deg \beta_m > 1$ and

$$(\Omega\rho)^* H^{n+s-2}(\Omega A; Z_p) \subset \sum_{i=1}^m \beta_i H^{n+r_i-3}(\Omega K; Z_p),$$

$$\psi(\Omega\rho) \equiv 0 \bmod \sum_{i=1}^m \beta_i H^{n+r_i-3}(\Omega K; Z_p),$$

then $l(E)=p$.

Corollary 2. Suppose that, for all i , $\tilde{\alpha}_i \in \sum_{k=1}^{i-1} \mathcal{A}(p)\alpha_k$ and $\tilde{\beta}_i \in \sum_{k=j+1}^m \beta_k \mathcal{A}(p)$ and that the homogeneous part $\mathcal{A}(p)$ of degree $s-1$ is contained in $\sum_{k=1}^m \beta_k \mathcal{A}(p) + \sum_{k=1}^m \mathcal{A}(p)\alpha_k$. If $\deg \beta_m > 1$ and the homogeneous part of $\mathcal{A}(p)$ of degree $s-r_i$ is trivial for all i , then $l(E)=p$.

Theorem D. Suppose that there exists i such that $\tilde{\alpha}_i \in \sum_{k=1}^{i-1} \mathcal{A}(p)\alpha_k$. If $(\Omega\rho)^* [\sum_{i=1}^m (-1)^{r_i} \tilde{\beta}_i \tilde{\alpha}_i] \equiv 0 \bmod \sum_{i=1}^m \beta_i H^{n+r_i-3}(\Omega K; Z_p)$, then $l(E)=p^3$; otherwise $l(E)=p^2$.

Corollary 3. Suppose that there exists i such that $\tilde{\alpha}_i \in \sum_{k=1}^{i-1} \mathcal{A}(p)\alpha_k$.

1) If $\sum_{i=1}^m (-1)^{r_i} \tilde{\beta}_i \tilde{\alpha}_i \in \sum_{k=1}^m \{\beta_k \mathcal{A}(p) + \mathcal{A}(p)\alpha_k\}$ and if

$$\sum_{i=1}^m \beta_i : \bigoplus_{i=1}^m H^{n+r_i-3}(\Omega^2 B) \rightarrow H^{n+s-2}(\Omega^2 B)$$

is monic, then $l(E)=p^3$.

2) If $\sum_{i=1}^m (-1)^{r_i} \tilde{\beta}_i \tilde{\alpha}_i \in \sum_{i=1}^m \{\beta_k \mathcal{A}(p) + \mathcal{A}(p)\alpha_k\}$, then $l(E)=p^2$.

REMARK. $\sum_{i=1}^m \beta_i$ is monic in each of the following cases:

- i) $\beta_i = Sq^{a_i}$, $a_1 > a_2 > \dots > a_m$, $a_i \geq 2(r_i - r_1 - 1)$ for $p=2$;
- ii) β_i are of the form P^{a_i} or ΔP^{a_i} and are all distinct, and $(2p-2)a_i \geq p(r_i - r_1 - 1)$ for $p > 2$.

1. A basic theorem

In this note we work in the category of based spaces having the homotopy types of CW complexes and based continuous maps, and we don't distinguish

between a map and the homotopy class it represents. Let $\pi: E \rightarrow K$ be the principal fibre space with $\theta: K \rightarrow L$ as classifying map and let $j: \Omega L \rightarrow E$ denote the fibre inclusion. Let p denote a fixed prime. A map of degree $p^k (k > 0)$ of $S = S^1$ yields the Puppe sequence

$$S \xrightarrow{p^k} S \xrightarrow{i} P \xrightarrow{q} S^2 \xrightarrow{p^k} S^2 \longrightarrow \dots$$

Form the commutative diagram

$$\begin{array}{ccccccc}
 & & & & & L^{S^2} & \\
 & & & & & \downarrow p^{\sharp\sharp} & \\
 & & & & K^{S^2} & \xrightarrow{\theta^{S^2}} & L^{S^2} \\
 & & & & \downarrow q^{\sharp} & & \downarrow q^{\sharp} \\
 & & & & K^P & \xrightarrow{\theta^P} & L^P \\
 & & & & \downarrow i^{\sharp} & & \downarrow i^{\sharp} \\
 \Omega L^S & \xrightarrow{j^S} & E^S & \xrightarrow{\pi^S} & K^S & \xrightarrow{\theta^S} & L^S \\
 \downarrow p^{\sharp\sharp} & & \downarrow p^{\sharp\sharp} & & \downarrow p^{\sharp\sharp} & & \\
 \Omega K^S & \xrightarrow{(\Omega\theta)^S} & \Omega L^S & \xrightarrow{j^S} & E^S & \xrightarrow{\pi^S} & K^S
 \end{array}$$

where rows and columns are fibration sequences and \sharp indicates induced maps of function spaces.

We now assume that K and L are loop spaces. Larmore and Thomas [2] have defined a sort of functional operation

$$\Phi_{\sharp}: [X, K^S] \cap \text{Ker}(p^{\sharp\sharp})_* \cap \text{Ker} \theta_*^S \rightarrow [X, L^{S^2}] / \theta_*^{S^2} [X, K^{S^2}] + (p^{\sharp\sharp})_* [X, L^{S^2}]$$

by setting $\Phi_{\sharp} = (q^{\sharp})_*^{-1} \theta_*^P (i^{\sharp})_*^{-1}$, with the property that, for $x \in [X, E^S]$ such that $(p^{\sharp\sharp})_* \pi_*^S x = 0$,

$$(1.1) \quad p^{\sharp} x \equiv -j_*^S \Phi_{\sharp}(\pi_*^S x) \pmod{j_*^S p^{\sharp} [X, \Omega L^S]},$$

where we have made the adjoint identification $[X, L^{S^2}] = [X, \Omega L^S]$ (cf. Theorem 3.2 of [3]).

In what follows we assume that

$$(1.2) \quad l(K) \text{ and } l(L) \text{ are divisors of } p^k;$$

$$(1.3) \quad [\Omega^2 L, \Omega^2 K] = 0;$$

$$(1.4) \quad [\Omega^2 L, Y] \xleftarrow{(\Omega j)^*} [\Omega E, Y] \xleftarrow{(\Omega \pi)^*} [\Omega K, Y] \xleftarrow{(\Omega \theta)^*} [\Omega L, Y]$$

is exact for $Y = \Omega^2 L$ and $\Omega^2 K$, (this condition may be verified using Theorem 6.5 of Sugawara [7]).

Taking $X = \Omega E$, $x = 1_{\Omega E}$ in (1.1), we then have

Theorem 1.5. *With the hypotheses (1.2), (1.3) and (1.4), we have*

- 1) $p^*1_{\Omega E} = -(\Omega j)_*\Phi_k(\Omega\pi).$
- 2) Write $\Psi_k(E)$ for the subset $(\Omega\pi)^{*^{-1}}\Phi_k(\Omega\pi)$ of $[\Omega K, \Omega^2 L]$.

Then $\Psi_k(E)$ is non-empty and is a coset of $(\Omega\theta)^[\Omega L, \Omega^2 L] + (\Omega^2\theta)_*[\Omega K, \Omega^2 K]$ such that $p^*1_{\Omega E} = 0$ if, and only if,*

$$\Psi_k(E) \equiv 0 \pmod{(\Omega\theta)^*[\Omega L, \Omega^2 L] + (\Omega^2\theta)_*[\Omega K, \Omega^2 K]}.$$

Proof. 1) is obvious by (1.1) and (1.2). Consider the commutative diagram

$$\begin{array}{ccccccc}
 & & [\Omega E, \Omega E] & & & & \\
 & & (\Omega j)_* & \swarrow & & & \\
 [\Omega^2 L, L^{S^2}] & \xleftarrow{(p^{k*})_*} & [\Omega^2 L, L^{S^2}] & \xleftarrow{(\Omega j)^*} & [\Omega E, L^{S^2}] & \xleftarrow{(\Omega\pi)^*} & [\Omega K, L^{S^2}] \xleftarrow{(\Omega\theta)^*} [\Omega L, \Omega^2 L] \\
 & & \downarrow q_*^* & & \downarrow q_*^* & & \downarrow (\Omega^2\theta)_* \\
 [\Omega^2 L, K^{S^2}] & & [\Omega^2 L, L^P] & \xleftarrow{(\Omega j)^*} & [\Omega E, L^P] & & [\Omega E, \Omega^2 K] \xleftarrow{(\Omega\pi)^*} [\Omega K, \Omega^2 K] \\
 & & \downarrow \theta_*^P & & \downarrow \theta_*^P & & \downarrow (\Omega j)_* \\
 & & [\Omega^2 L, K^P] & \xleftarrow{(\Omega j)^*} & [\Omega E, K^P] & & [\Omega^2 L, \Omega^2 K] \\
 & & \downarrow i_*^* & & \downarrow i_*^* & & \\
 & & [\Omega^2 L, K^S] & \xleftarrow{(\Omega j)^*} & [\Omega E, K^S] & &
 \end{array}$$

Since $(\Omega j)^*\Omega\pi = 0 = i_*^*(\Omega j)^*(i_*^*)^{-1}\Omega\pi$ and q_*^* and the left i_*^* are monic by virtue of (1.2) and (1.3), we see that $(\Omega j)^*\Phi_k(\Omega\pi) = 0$, and hence there exists $y \in [\Omega K, \Omega^2 L]$ with $(\Omega\pi)^*y \in \Phi_k(\Omega\pi)$, which shows that $\Psi_k(E)$ is non-empty. By diagram-chasing we may easily verify that $(\Omega\pi)^{*^{-1}}\text{Ker}(\Omega j)_* = (\Omega\pi)^{*^{-1}}(\Omega^2\theta)_*[\Omega E, \Omega^2 K]$ coincides with $(\Omega\theta)^*[\Omega L, \Omega^2 L] + (\Omega^2\theta)_*[\Omega K, \Omega^2 K]$. The last assertion follows from 1), since $p^*1_{\Omega E} = 0$ iff $\Phi_k(\Omega\pi) = \text{Ker}(\Omega j)_*$.

We note that the assignment $\theta \rightarrow \Psi_1(E)$ is dual to Toda's derivative θ ([9], p. 209).

2. Proofs of Theorems A and B

We may prove Corollary 1 in the introduction as follows. Let $\theta: K_n \rightarrow K_{n+r}$. Then, by Corollary 3.7 of [2], $\Psi_1(E) = (-1)^{n+r+1}\tilde{\theta}_{\ell_{n-1}}$. Hence our assertion follows from 2) of Theorem 1.5.

We now consider more general situation. Let

$$\begin{aligned} K &= \bigotimes_{i=1}^k K_{n+s_i}, \quad L = \bigotimes_{j=1}^m K_{n+r_j}, \\ 0 &= s_1 \leq s_2 \leq \cdots \leq s_k < r_1 \leq r_2 \leq \cdots \leq r_m \leq n-3, \\ \theta &= \{\theta_1, \dots, \theta_m\}, \\ \theta_j &= \sum_{i=1}^k \pi_i^* \theta_{ji}, \quad \theta_{ji} \in \mathcal{A}(p), \quad \deg \theta_{ji} = r_j - s_i, \end{aligned}$$

where $\pi_i: K \rightarrow K_{n+s_i}$ is the projection on the i -th factor. Then Theorems A and B are consequences of the following

Theorem 2.1. *Let E be the principal fibre space with the above θ as classifying class. Then $l(E) = p^2$ if, and only if, there exist j and i , $1 \leq j \leq m$, $1 \leq i \leq k$, such that*

$$\tilde{\theta}_{ji} \in \sum_{t=1}^{j-1} \mathcal{A}(p) \theta_{ti} + \sum_{t=i+1}^k \theta_{jt} \mathcal{A}(p).$$

Proof. Introduce the diagram

$$\begin{array}{ccccccc} K_{n+r_j-2} & \xrightarrow{l_j} & K_{n+r_j-2} \times K_{n+r_j-1} & \xleftarrow{\varphi} & \bigotimes_{i=1}^k (K_{n+s_i-2} \times K_{n+s_i-1}) & \xrightarrow{\bigotimes_{i=1}^k p_i} & \bigotimes_{i=1}^k K_{n+s_i-1} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ K_{n+r_j}^{S^2} & \xrightarrow{q^*} & K_{n+r_j}^P & \xleftarrow{\theta_j^P} & \bigotimes_{i=1}^k K_{n+s_i}^P & \xrightarrow{i^*} & K^S = \bigotimes_{i=1}^k K_{n+s_i}^S \end{array}$$

where p_i denotes the projection on the second factor, l_j the injection and vertical maps are homotopy equivalences as given in Proposition 3.3 of [2]. Here we take the cofibre of $p: S \rightarrow S$ for P . φ is defined by

$$\begin{aligned} \varphi^*(\iota_{n+r_j-2} \times 1) &= \sum_{i=1}^k \pi_i^* [\theta_{ji} \iota_{n+s_i-2} \times 1 + (-1)^{n+r_j-1} \times \tilde{\theta}_{ji} \iota_{n+s_i-1}], \\ \varphi^*(1 \times \iota_{n+r_j-1}) &= \sum_{i=1}^k \pi_i^* (1 \times \theta_{ji} \iota_{n+s_i-1}). \end{aligned}$$

We see from Theorem 3.6 of [2] that the above diagram homotopy-commutes.

Apply $[\Omega E, \]$ to the above diagram. Since $\theta_j^P = \sum_{i=1}^k \theta_{ji}^P \pi_i^P$, $\theta^P = \{\theta_1^P, \dots, \theta_m^P\}$ and since

$$\begin{aligned} & l_j^* \left(\sum_{i=1}^k (-1)^{n+r_j} (\Omega \pi)^* (\Omega \pi_i)^* \tilde{\theta}_{ji} \iota_{n+s_i-1} \right) \\ &= \left(\sum_{i=1}^k (-1)^{n+r_j} (\Omega \pi)^* (\Omega \pi_i)^* \tilde{\theta}_{ji} \iota_{n+s_i-1}, 0 \right), \end{aligned}$$

$$(\bigotimes_{i=1}^k p_i)_*(0, \Omega(\pi_1\pi); \dots; 0, \Omega(\pi_k\pi)) = \Omega\pi,$$

$$\varphi_*(0, \Omega(\pi_1\pi); \dots; 0, \Omega(\pi_k\pi)) = \left(\sum_{i=1}^k (-1)^{n+r_j} (\Omega\pi_i)^* (\Omega\pi_i)^* \tilde{\theta}_{ji} \iota_{n+s_i-1}, 0 \right)$$

by $\theta_j(\Omega\pi)=0$, it follows that the j -th component of $\Phi_1(\Omega\pi)$ has a representative

$$(\Omega\pi)^* \sum_{i=1}^k (-1)^{n+r_j} (\Omega\pi_i)^* \tilde{\theta}_{ji} \iota_{n+s_i-1}. \quad \text{Hence}$$

$$\sum_{i=1}^k (-1)^{n+r_j} (\Omega\pi_i)^* \tilde{\theta}_{ji} \iota_{n+s_i-1}$$

represents the j -th component of $\Psi_1(E)$.

Now, by the Künneth theorem, we compute $(\Omega^2\theta_j)_*[\Omega K, \Omega^2 K] + (\Omega^2\pi_j)_*(\Omega\theta)^*[\Omega L, \Omega^2 L]$ as follows:

$$\begin{aligned} (\Omega^2\theta_j)_*[\Omega K, \Omega^2 K] &= \sum_{i=1}^k (\Omega^2\theta_{jt})_* H^{n+s_i-2}(\Omega K; Z_p) \\ &= \left\{ \sum_{t=1}^k \theta_{jt} \sum_{i=1}^k (\Omega\pi_i)^* \alpha_{ti} \iota_{n+s_i-1}; \alpha_{ti} \in \mathcal{A}(p), \right. \\ &\quad \left. \deg \alpha_{ti} = s_t - s_i - 1 \right\}, \\ (\Omega^2\pi_j)_*(\Omega\theta)^*[\Omega L, \Omega^2 L] &= H^{n+r_j-2}(\Omega L; Z_p)(\Omega\theta_1, \dots, \Omega\theta_m) \\ &= \sum_{t=1}^m H^{n+r_j-2}(Z_p, n+r_t-1; Z_p)(\Omega\theta_t) \\ &= \sum_{t=1}^m \sum_{i=1}^k H^{n+r_j-2}(Z_p, n+r_t-1; Z_p)(\theta_{ti})(\Omega\pi_i). \end{aligned}$$

These complete the proof of Theorem 2.1.

In connection with Corollary 1 we examine some elements in the kernel of the Kristensen map $\varepsilon: \mathcal{A}(2) \rightarrow \mathcal{A}(2)$. Let $Sq(i_1, \dots, i_k)$ denote $Sq^{i_1} \dots Sq^{i_k}$. Then, using the Adem relation $Sq(2m-1, m)=0$ ($m \geq 1$), we may easily verify

Proposition 2.2. *The following elements are in the kernel of ε :*

$$Sq(3k) + \sum_{i=1}^k Sq(3k-i, i), \quad k \geq 1;$$

$$Sq(6k+1) + Sq(6k, 1) + \sum_{i=1}^k Sq(6k+1-2i, 2i) + \sum_{j=2}^{2k} Sq(6k-j, j, 1), \quad k \geq 1;$$

$$\sum_{i=1}^k Sq(6k+3-2i, 2i+1) + \sum_{j=2}^{2k+1} Sq(6k+3-j, j, 1), \quad k \geq 1;$$

$$Q + Sq(6k-1, 2, 1) + Sq(6k-2, 3, 1) + \sum_{j=2}^k Sq(6k-2j+1, 2j, 1) + \sum_{r=4}^{2k} Sq(6k-r, r, 2),$$

where

$$Q = \begin{cases} Sq(6k+2) + Sq(6k+1, 1) + Sq(6k, 2) + Sq(6k-2, 4) \\ \quad + \sum_{i=2}^{k/2} [Sq(6k-4i+3, 4i-1) + Sq(6k-4i+2, 4i)] & \text{for } k \text{ even,} \\ Sq(6k-1, 3) + \sum_{i=1}^{(k-1)/2} [Sq(6k-4i+1, 4i+1) + Sq(6k-4i, 4i+2)] & \text{for } k \text{ odd;} \end{cases}$$

$$R + \sum_{j=2}^k Sq(6k-2j+3, 2j+1, 1) + \sum_{r=4}^{2k+1} Sq(6k-r+3, r, 2),$$

where

$$R = \begin{cases} Sq(6k+5) + \sum_{i=1}^3 Sq(6k+5-i, i) + \sum_{i=1}^{k/2} [Sq(6k+5-4i, 4i) \\ \quad + Sq(6k+4-4i, 4i+1)] & \text{for } k \text{ even,} \\ \sum_{i=1}^{(k-1)/2} [Sq(6k-4i+3, 4i+2) + Sq(6k-4i+2, 4i+3)] & \text{for } k \text{ odd.} \end{cases}$$

We mention some examples. The loop-order of the fibre space with classifying class $\{Sq^{i_1}, \dots, Sq^{i_k}\}$, $0 < i_1 \leq i_2 \leq \dots \leq i_k$, is 4, but those of fibre spaces with classifying classes $Sq^3 + Sq^2Sq^1$, $Sq^4Sq^2 + Sq^2Sq^4$, $Sq^7 + Sq^6Sq^1 + Sq^5Sq^2 + Sq^4Sq^2Sq^1$ are 2. The loop-order of the fibre space with classifying class $\{P^k, \Delta P^k\}$ ($k \geq 1$) is p .

3. Proof of Theorem C

First we prove 1). Introduce the commutative diagram

$$\begin{array}{ccccc} E_0 & \xrightarrow{l_0} & E & & \\ \pi_0 \downarrow & & \downarrow \pi & \searrow \theta & \\ \Omega B & \xrightarrow{l} & K & \longrightarrow & L \end{array}$$

where the square is a pull-back. Observe that the fibre of l_0 is homotopy-equivalent to that of l , i.e., ΩA . Since $\pi_0: E_0 \rightarrow \Omega B$ is a principal fibration with $\beta = \theta l$ as classifying map, we have $l(E_0) = p^2$ by Theorem B, and hence it follows from the exact sequence

$$[\Omega E_0, \Omega^2 A] \longrightarrow [\Omega E_0, \Omega E_0] \xrightarrow{(\Omega l_0)_*} [\Omega E_0, \Omega E]$$

and from the $(n+r_1-2)$ -connectedness of E_0 that the order of Ωl_0 is p^2 and $l(E)$ is a multiple of p^2 . Also, since $l(K) = p$ by Theorem A, we see that $l(E) = p^2$.

We now proceed to prove 2) and 3). Note that, in the situation (*), θ determines a secondary operation $\varphi: \bigcap_{i=1}^m \text{Ker } \alpha_i \rightarrow \text{Coker } \sum_{i=1}^m \beta_i$ associated with the relation $\sum_{i=1}^m \beta_i \alpha_i = 0$ (cf. Adams [1], Spanier [6]). Take the cofibre P of $p^k: S \rightarrow S$

($k=1, 2$). Applying the functor $(\)^P$ to the diagram $(*)$, we see similarly that θ^P determines a secondary operation

$$\bar{\varphi}: [X, A^P] \cap \text{Ker } \alpha^P \rightarrow [X, L^P]/\text{Im } \beta^P$$

associated with $\beta^P(\Omega\alpha)^P=0$, where

$$\begin{aligned} (\Omega\alpha)^P &= \bigotimes_{i=1}^m (\alpha_i \times 1 + (-1)^{n+r_i-1} \lambda_k (1 \times \tilde{\alpha}_i), 1 \times \alpha_i), \\ \beta^P &= \sum_{i=1}^m (\Omega\pi_i^P)^* \{ \beta_i \times 1 + (-1)^{n+s} \lambda_k (1 \times \tilde{\beta}_i), 1 \times \beta_i \}, \quad (\lambda_1 = 1, \lambda_2 = 0) \end{aligned}$$

Let $t: L^P \rightarrow \Omega^2 L$ denote a projection with $tq^* \simeq 1$ and let $e: \Omega A \rightarrow A^P$, $e: \Omega B \rightarrow B^P$ denote injections with $i^* e \simeq 1$. Then

$$\begin{aligned} \alpha^P e &= \{ (-1)^{n+r_1} \lambda_k \tilde{\alpha}_1, \alpha_1; \dots; (-1)^{n+r_m} \lambda_k \tilde{\alpha}_m, \alpha_m \}, \\ t\beta^P &= \sum_{i=1}^m (\Omega\pi_i^P)^* (\beta_i \times 1 + (-1)^{n+s} \lambda_k (1 \times \tilde{\beta}_i)). \end{aligned}$$

Consider the following commutative diagram

$$(3.1) \quad \begin{array}{ccccccc} \Omega^2 B & \xrightarrow{\Omega e} & \Omega B^P & \xlongequal{\quad} & \Omega B^P & & \\ \Omega l \downarrow & & \downarrow \bar{l} & & \downarrow l^P & & \\ \Omega K & \xrightarrow{f} & \bar{K} & \xrightarrow{\varepsilon} & K^P & \xrightarrow{\theta^P} & L^P \xrightarrow{t} \Omega^2 L \\ \Omega \rho \downarrow & & \downarrow \bar{\rho} & & \downarrow \rho^P & & \\ \Omega A & \xlongequal{\quad} & \Omega A & \xrightarrow{e} & A^P & \xrightarrow{\alpha^P} & B^P \end{array}$$

where $\bar{\rho}$ is the pull-back of ρ^P by e , hence the principal fibration with classifying map $\alpha^P e$. We denote by $\psi_k(\theta)$ the secondary operation determined by $t\theta^P \varepsilon$, which is associated with $(t\beta^P)\Omega(\alpha^P e)=0$. Since $\alpha_i(\Omega\rho)=0$ yields $\tilde{\alpha}_i(\Omega\rho)=0$ for $k=1$ with $\tilde{\alpha}_i \in \sum_{j=1}^{i-1} \mathcal{A}(p)\alpha_j$ and since $\lambda_2=0$, we may define $\bar{\varphi}(0, \Omega\rho)$ and $\psi_k(\theta)(\Omega\rho)$. Note that $\psi_k(\theta)(\Omega\rho)$ is the first component of $\bar{\varphi}(0, \Omega\rho)$.

Lemma 3.2. *Let $k=1$ or 2 . Suppose $\deg \beta_m > 1$ for $k=1$. Then there exists $f: \Omega K \rightarrow \bar{K}$ such that $\bar{\rho} f = \Omega\rho$ and $t\theta^P \varepsilon f$ represents both $\psi_k(\theta)(\Omega\rho)$ and $\Psi_k(E)$. Moreover, if $k=2$, $i^* \varepsilon f \simeq 1$ and $f(\Omega l) \simeq \bar{l}(\Omega e)$.*

Proof. Assume first $k=1$ and $\deg \beta_m > 1$. Take $x: \Omega E \rightarrow K^P$ with $i_*^* x = \pi^S$. Since $[\Omega^2 L, K^P]=0$ by $s > r_m$, we have $(\Omega j)^* x = 0$, and hence we may pick $y \in [K^S, K^P]$ with $x = (\Omega\pi)^* y$. Further, since $[\Omega K, A^{S^2}]=0$, we may set $\rho^P y = (0, z)$ for $z = i_*^* \rho^P y = (\Omega\rho) i_*^* y$. We have

$$(0, z(\Omega\pi)) = (0, z)(\Omega\pi) = \rho^P y(\Omega\pi) = \rho^P x = (0, (\rho\pi)^S)$$

by $i_*^* \rho^P x = (\rho\pi)^S$ and $[\Omega E, \Omega^2 A] = 0$. Therefore,

$$z - \Omega\rho \in \text{Ker}(\Omega\pi)^* = (\Omega\theta)^*[\Omega L, \Omega A] = 0.$$

This gives rise to $\rho^P y = (0, \Omega\rho) = e(\Omega\rho)$, which yields $f: \Omega K \rightarrow \bar{K}$ with $\bar{\rho}f = \Omega\rho$, $\varepsilon f = y$. Now $\Phi_1(\Omega\pi)$ has, by definition, a representative $(q_*^*)^{-1}\theta^P(x)$. Thus

$$\Phi_1(\Omega\pi) = t_* q_*^* \Phi_1(\Omega\pi) \ni t_* \theta^P(x) = t_* \theta^P y(\Omega\pi).$$

This shows that $t\theta^P y = t\theta^P \varepsilon f$ represents $\Psi_1(E)$ and $\psi_1(\theta)(\Omega\rho)$.

Next let $k=2$; then, $\alpha^P e \simeq e(\Omega\alpha)$ by virtue of the expression of $\alpha^P e$, and hence one gets an induced map $\bar{e}: \Omega K \rightarrow K^P$ which makes the following diagram homotopy-commute:

$$\begin{array}{ccccccc} \Omega^2 B & \xrightarrow{\Omega l} & \Omega K & \xrightarrow{\Omega \rho} & \Omega A & \xrightarrow{\Omega \alpha} & \Omega B \\ \Omega e \downarrow & & \downarrow \bar{e} & & \downarrow e & & \downarrow e \\ \Omega B^P & \xrightarrow{l^P} & K^P & \xrightarrow{\rho^P} & A^P & \xrightarrow{\alpha^P} & B^P \\ i^* \downarrow & & \downarrow i^* & & \downarrow i^* & & \downarrow i^* \\ \Omega^2 B & \xrightarrow{\Omega l} & \Omega K & \xrightarrow{\Omega \rho} & \Omega A & \xrightarrow{\Omega \alpha} & \Omega B \end{array}$$

Since $i^* e \simeq 1$, it follows from the five lemma that $i^* \bar{e}$ is a homotopy equivalence with a homotopy inverse $\xi: \Omega K \rightarrow \Omega K$. Thus, by factoring \bar{e} , we may find $f: \Omega K \rightarrow \bar{K}$ such that $\bar{e}\xi = \varepsilon f$, $\bar{\rho}f = \Omega\rho$, $i^* \varepsilon f \simeq 1$ and $\varepsilon f(\Omega l) \simeq \varepsilon \bar{l}(\Omega e)$. Since the fibre of $e: \Omega A \rightarrow A^P$ is homotopy-equivalent to the loop space of that of i^* by inspection of the relative mapping sequence for $i^* e \simeq 1$ (cf. [4], Lemma 2.1 (ii)), and since the fibre of i^* is $\Omega^2 A$, we see from $[\Omega^2 B, \Omega^3 A] = 0$ that $\varepsilon_*: [\Omega^2 B, \bar{K}] \rightarrow [\Omega^2 B, K^P]$ is monic. This implies that $f(\Omega l) \simeq \bar{l}(\Omega e)$. $i^* \varepsilon f \simeq 1$ implies $i^*(\varepsilon f(\Omega\pi)) \simeq \Omega\pi$, hence $tq_*^*(q_*^*)^{-1}\theta^P \varepsilon f(\Omega\pi)$ represents $\Phi_2(\Omega\pi)$. q.e.d.

Now let $k=1$. We observe that

$$\begin{aligned} t_* \beta^P[\Omega K, \Omega B^P] &\supset t_* \beta^P q_*^*[\Omega K, \Omega^3 B] \\ &= (\Omega^2 \beta)_*[\Omega K, \Omega^3 B] \\ &= (\Omega^2 \theta)_*[\Omega K, \Omega^2 K] \quad \text{by } [\Omega K, \Omega^2 A] = 0, \end{aligned}$$

and that, if $\tilde{\beta}_i \in \sum_{j=i+1}^m \beta_j \mathcal{A}(p)$ then

$$t_* \beta^P[\Omega K, \Omega B^P] = (\Omega^2 \beta)_*[\Omega K, \Omega^3 B].$$

Thus we may infer from Theorem 1.5, 2) that $\psi_1(\theta)(\Omega\rho) \equiv 0 \pmod{t_* \beta^P[\Omega K, \Omega B^P]}$

implies $p1_{\Omega E} \neq 0$. Since $\psi(\Omega\rho)$ differs from $\psi_1(\theta)(\Omega\rho)$ by an element of $(\bar{p}f)^*[\Omega A, \Omega^2 L] = (\Omega\rho)^*[\Omega A, \Omega^2 L]$, the assertions 2) and 3) of Theorem C are obtained.

Corollary 2 is obtained from 3) of Theorem C, by noting that the sequence $H^{n+s-2}(\Omega^2 B) = H^{n+s-2}(\bigotimes_{i=1}^m K_{n+r_i-2}) \leftarrow H^{n+s-2}(\Omega K) \xleftarrow{(\Omega\rho)^*} H^{n+s-2}(\Omega A)$ is exact and $H^{n+s-2}(\Omega A)$ is contained in $\sum_{j=i+1}^m \beta_j \mathcal{A}(p) + \text{Ker}(\Omega\rho)^*$.

By the way, we examine the extent to which $\psi_k(\theta)(\Omega\rho)$ may be altered with θ being a universal example of a secondary operation associated with $\beta(\Omega\alpha) = 0$.

Proposition 3.3. $\psi_1(\theta + \rho^* \gamma)(\Omega\rho) = \psi_1(\theta)(\Omega\rho) \pm (\Omega\rho)^* \widetilde{\Omega\gamma}$,
 $\psi_2(\theta + \rho^* \gamma)(\Omega\rho) = \psi_2(\theta)(\Omega\rho)$ for $\gamma \in [A, L]$.

Proof. Since t can be delooped, we have

$$\begin{aligned} t(\theta^P + \gamma^P \rho^P) \varepsilon f &= t\theta^P \varepsilon f + t\gamma^P \rho^P \varepsilon f \\ &= t\theta^P \varepsilon f + t\gamma^P e(\Omega\rho) \\ &= t\theta^P \varepsilon f + (\Omega^2 \gamma \times 1)e(\Omega\rho) \pm \lambda_k(1 \times \widetilde{\Omega\gamma})e(\Omega\rho) \\ &= t\theta^P \varepsilon f \pm \lambda_k(\widetilde{\Omega\gamma})(\Omega\rho). \end{aligned}$$

4. Proof of Theorem D

In this section let P and P' be cofibres of $p^2: S \rightarrow S$ and of $p: S \rightarrow S$ respectively. Given a generalized Eilenberg-MacLane space Z , let

$$Z^{S^2} \xrightleftharpoons[t]{q^*} Z^P \xrightleftharpoons[e]{i^*} Z^S$$

and

$$Z^{S^2} \xrightleftharpoons[t']{q'^*} Z^{P'} \xrightleftharpoons[e']{i'^*} Z^S$$

denote product representations.

Introduce the following commutative diagram

$$(4.1) \quad \begin{array}{ccccccc} S & \xrightarrow{p} & S & \xrightarrow{i'} & P' & \xrightarrow{q'} & S^2 \\ \parallel & & \downarrow p & & \downarrow (1, p) & & \parallel \\ S & \xrightarrow{p^2} & S & \xrightarrow{i} & P & \xrightarrow{q} & S^2 \\ \downarrow & & \downarrow i' & & \downarrow (0, i') & & \downarrow \\ * & \longrightarrow & P' & \xlongequal{\quad} & P' & \longrightarrow & * \\ & & \downarrow q' & & \downarrow (0, q') & & \\ & & S^2 & \longrightarrow & SP' & & \end{array}$$

in which rows and columns are Puppe sequences by the 3×3 lemma (cf. Nomura [4], Lemma 1.2) and $(1, p)$ and $(0, i')$ are induced maps.

Lemma 4.2. (4.1) induces a fibration sequence

$$K_n^{P'} \xleftarrow{(1, p)^*} K_n^P \xleftarrow{(0, i')^*} K_n^{P'} \xleftarrow{(0, q')^*} K_n^{SP'}$$

which is homotopically equivalent to

$$K_{n-2} \times K_{n-1} \xleftarrow{1 \times 0} K_{n-2} \times K_{n-1} \xleftarrow{0 \times 1} K_{n-2} \times K_{n-1} \xleftarrow{T(0 \times 1)} K_{n-3} \times K_{n-2}$$

where $T: K_{n-1} \times K_{n-2} \rightarrow K_{n-2} \times K_{n-1}$ denotes the switching map.

Proof. From the diagram (4.1) one can form the homotopy-commutative diagram

$$\begin{array}{ccccccc} K_n^{S^2} & \xlongequal{\quad} & K_n^{S^2} & & & & \\ t' \downarrow q'^* & & t \downarrow q^* & & & & \\ K_n^{P'} & \xleftarrow{(1, p)^*} & K_n^P & \xleftarrow{(0, i')^*} & K_n^{P'} & \xleftarrow{(0, q')^*} & K_n^{SP'} \\ \downarrow i'^* & & \downarrow i^* & & \parallel & & \downarrow (Si')^* \\ K_n^S & \xleftarrow{p^*=0} & K_n^S & \xleftarrow{i'^*} & K_n^{P'} & \xleftarrow{q'^*} & K_n^{S^2} \end{array}$$

Then $t'(1, p)^* \in H^{n-2}(K_n^P; Z_p) \cong H^{n-2}(K_{n-2} \times K_{n-1}; Z_p)$ is a multiple of the projection $t: K_{n-2} \times K_{n-1} \rightarrow K_{n-2}$. Since $t'(1, p)^* q^* \simeq t' q'^* \simeq 1$, it follows that $t'(1, p)^* \simeq t$. This shows that $(1, p)^*$ is essentially 1×0 and that $t(0, i')^* \simeq t'(1, p)^*(0, i')^* \simeq 0$. Hence $(0, i')^*$ is essentially 0×1 and, by $i'^*(0, q')^* \simeq 0$ and $t'(0, q')^* \simeq (Si')^*$, we see that $(0, q')^*$ is homotopy-equivalent to $T(0 \times 1)$.


Consider now the homotopy-commutative diagram

$$\begin{array}{ccccccc} & & & & B^{SP'} & & \\ & & & & \downarrow T(0 \times 1) & \searrow (Si')^* & \\ & & A^{P'} & \xrightarrow{\alpha^{P'}} & B^{P'} & \xrightarrow{t'} & B^{S^2} \\ & & \downarrow 0 \times 1 & & \downarrow 0 \times 1 & & \\ \bar{K} & \xrightarrow{\bar{p}} & \Omega A & \xrightarrow{e} & A^P & \xrightarrow{\alpha^P} & B^P \\ \chi \downarrow & & \parallel & & \downarrow 1 \times 0 & & \downarrow 1 \times 0 \\ K' & \xrightarrow{\rho'} & \Omega A & \xrightarrow{(1 \times 0)e} & A^{P'} & \xrightarrow{\alpha^{P'}} & B^{P'} \end{array}$$

where \bar{K} is, as in (3.1), the fibre of $\alpha^P e$ and K' is the fibre of $\alpha^{P'}(1 \times 0)e$. Note that K' is homotopy-equivalent to $\Omega A \times \Omega B^{P'}$ because of $(1 \times 0)e \simeq 0$. The maps

1×0 induce a map $\chi: \bar{K} \rightarrow K'$.

Let $f: \Omega K \rightarrow \bar{K}$ be a map constructed in Lemma 3.2 for $k=2$. Then one gets the homotopy-commutative diagram



Since $\chi f(\Omega l) \simeq \chi \bar{l}(\Omega e) \simeq l' \Omega(1 \times 0)(\Omega e) \simeq 0$ by $\Omega(1 \times 0)(\Omega e) \simeq 0$, we may find $f': \Omega A \rightarrow K'$ such that

$$(4.3) \quad f'(\Omega \rho) \simeq \chi f$$

and so

$$(4.4) \quad t \theta^P \varepsilon f \simeq t' \theta^{P'} \varepsilon' f'(\Omega \rho).$$

Further, since $\rho^{P'} \varepsilon' f' \simeq 0$, there exists $g: \Omega A \rightarrow \Omega B^{P'}$ such that

$$(4.5) \quad l^{P'} g \simeq \varepsilon' f'.$$

Therefore, by (4.4) and $\theta l = \beta$,

$$(4.6) \quad t \theta^P \varepsilon f \simeq t' \beta^{P'} g(\Omega \rho).$$

We next show that

$$(4.7) \quad T(0 \times 1)g(\Omega \rho) \simeq -\alpha^{P'} e'(\Omega \rho).$$

For this purpose, introduce the commutative diagram

$$\begin{array}{ccccccc}
 & & & & & B^{SP} & \\
 & & & & & \downarrow 1 \times 0 & \\
 & & & & A^{SP'} & \xrightarrow{\alpha^{SP'}} & B^{SP'} \\
 & & & & \downarrow & & \downarrow T(0 \times 1) \\
 & & & & A^{P'} & \xrightarrow{\alpha^{P'}} & B^{P'} \\
 & & & & \downarrow 0 \times 1 & & \downarrow 0 \times 1 \\
 \Omega B^P & \xrightarrow{l^P} & K^P & \xrightarrow{\rho^P} & A^P & \xrightarrow{\alpha^P} & B^P \\
 & & \downarrow \Omega(1 \times 0) & \downarrow (1, p)^* & \downarrow 1 \times 0 & & \downarrow 1 \times 0 \\
 \Omega A^{P'} & \xrightarrow{\Omega \alpha^{P'}} & \Omega B^{P'} & \xrightarrow{l^{P'}} & K^{P'} & \xrightarrow{\alpha^{P'}} & B^{P'}
 \end{array}$$

Apply the functor $[\Omega K, \]$ to the above diagram and observe that

$$\alpha^P e(\Omega \rho) \simeq \alpha^P \rho^P f \simeq \alpha^P \rho^P \varepsilon f \simeq 0, \quad (1 \times 0)e(\Omega \rho) \simeq 0.$$

Since, by Lemma 3.2, (4.3) and (4.5),

$$\rho^P \varepsilon f = e(\Omega \rho), \quad (1, p)^* \varepsilon f \simeq l^{P'} g(\Omega \rho), \quad (0 \times 1)e'(\Omega \rho) \simeq e(\Omega \rho),$$

we can apply two kinds of functional operations to $e(\Omega \rho) \in [\Omega K, A^P]$ to yield $g(\Omega \rho) \in [\Omega K, \Omega B^{P'}]$ and $[T_*(0 \times 1)_*]^{-1} \alpha^{P'} e'(\Omega \rho) \in [\Omega K, B^{SP'}]$. Thus, according to Spanier [6],

$$-g(\Omega \rho) \equiv [T_*(0 \times 1)_*]^{-1} \alpha^{P'} e'(\Omega \rho) \bmod \alpha_*^{SP'}[\Omega K, A^{SP'}] + (1 \times 0)_*[\Omega K, B^{SP}]$$

under the adjoint isomorphism. Hence (4.7) follows from the fact that $[\Omega K, \Omega A^{P'}] = 0$ and $[T(0 \times 1)_*](1 \times 0)_* = 0$.

We now compute, by the expression for $t' \beta^{P'}$ and $\alpha^{P'} e'$ in §3,

$$\begin{aligned}
 t' \beta^{P'} g(\Omega \rho) &= (\Omega \rho)^* g^* \sum_{j=1}^m \pi_j^{P'} * (\beta_j \times 1 + (-1)^{n+s} 1 \times \tilde{\beta}_j) \\
 &\equiv (\Omega \rho)^* g^* \sum_{j=1}^m (-1)^{n+s} \pi_j^{P'} * (1 \times \tilde{\beta}_j) \bmod (\Omega \rho)^*(\Omega^2 \beta)_*[\Omega A, \Omega^3 B] \\
 &= (-1)^{n+s} (\Omega \rho)^* g^* \sum_{j=1}^m \pi_j^{P'} * (\Omega i'^*)^* \tilde{\beta}_j \\
 &= (-1)^{n+s} (\Omega \rho)^* g^* (\Omega i'^*)^* \sum_{j=1}^m \pi_j^* \tilde{\beta}_j \\
 &= (-1)^{n+s} (\Omega \rho)^* g^* (t' T(0 \times 1))^* \sum_{j=1}^m \pi_j^* \tilde{\beta}_j \\
 &= (-1)^{n+s+1} (\Omega \rho)^* (t' \alpha^{P'} e')^* \sum_{j=1}^m \pi_j^* \tilde{\beta}_j \\
 &= (-1)^{s+1} (\Omega \rho)^* \sum_{j=1}^m (-1)^r i_j \tilde{\beta}_j \tilde{\alpha}_j.
 \end{aligned}$$

This reveals that $(-1)^{s+1} \sum_{j=1}^m (-1)^r i \beta_j \tilde{\alpha}_j$ represents $\Psi_2(E)$ by Lemma 3.2, since $(\Omega\rho)^*(\Omega^2\beta)_*[\Omega A, \Omega^2 B]$ is contained in the indeterminacy, $(\Omega^2\theta)_*[\Omega K, \Omega^2 K] = (\Omega^2\beta)_*[\Omega K, \Omega^3 B]$, of $\Psi_2(E)$. Therefore, Theorem *D* follows from Theorem 1.5 and from the fact $p^2 | l(E)$ is a consequence of the exact sequence

$$[\Omega E, \Omega K] \xleftarrow{(\Omega\pi)^*} [\Omega K, \Omega K] \xleftarrow{(\Omega\theta)^*} [\Omega L, \Omega K] = 0.$$

Corollary 3, 1) follows from Theorem *D* by inspecting the exact ladder

$$\begin{array}{ccccccc} [\Omega^2 B, \Omega^3 B] & \longleftarrow & [\Omega K, \Omega^3 B] & \xleftarrow{(\Omega\rho)^*} & [\Omega A, \Omega^3 B] & & \\ \downarrow (\Omega^2\beta)_* & & \downarrow (\Omega^2\beta)_* & & \downarrow (\Omega^2\beta)_* & & \\ H^{n+s-2}(\Omega^2 B) & \longleftarrow & H^{n+s-2}(\Omega K) & \xleftarrow{(\Omega\rho)^*} & H^{n+s-2}(\Omega A) & \xleftarrow{(\Omega\alpha)^*} & H^{n+s-2}(\Omega B) \end{array}$$

and by observing that the left hand $(\Omega^2\beta)_*$ may be identified with

$$\sum_{i=1}^m \beta_i: \oplus H^{n+r_i-3}(\Omega^2 B) \rightarrow H^{n+s-2}(\Omega^2 B).$$

5. Some examples

As an illustration of Theorems C and D in the introduction, we list some relations in $\mathcal{A}(p)$ to which the theorems are applicable:

i) Relations to which Theorem C, 1), is applicable:

$$\begin{aligned} (P^k \Delta) P^{p-1} &= 0 \quad (2 \leq k < p), \\ (P^p \Delta) P^k + (k-1) \Delta P^{p+k} - (\Delta P^{p+k-1}) P^1 &= 0 \quad (1 < k < p). \end{aligned}$$

ii) Relations to which Theorem C, 2) is applicable:

$$\begin{aligned} (\Delta P^k) P^{k-1} - P^k (\Delta P^{k-1}) - P^{k-1} (\Delta P^k) &= 0 \quad (k \geq 2, k \not\equiv 0 \pmod{p}, \\ p > 3, k < (p^{2p-4} + 2p - 3)(p^2 - 1)^{-1}). \end{aligned}$$

iii) Relations to which Corollary 2 is applicable:

$$\begin{aligned} P^{p-1} P^1 &= 0 \quad (p > 3), \\ P^p P^{p+2} - P^{2p+1} P^1 &= 0. \end{aligned}$$

iv) Relations to which Corollary 3, 2) is applicable:

$$\begin{aligned} Sq^{2k-1} Sq^{k-1} + Sq^{2k-2} Sq^k &= 0 \quad (k \geq 2), \\ Sq^{2k-1} Sq^{k-3} + Sq^{2k-2} Sq^{k-2} + Sq^{2k-4} Sq^k &= 0 \quad (k \geq 4), \\ Sq^{2k-1} Sq^{k-5} + Sq^{2k-2} Sq^{k-4} + Sq^{2k-3} Sq^{k-3} + Sq^{2k-6} Sq^k &= 0 \quad (k \geq 6). \end{aligned}$$

v) Relations to which Corollary 3, 1) is applicable:

$$\begin{aligned}
 l(E) = 8 \quad \text{iff} \quad Sq^{2k-2}Sq^{k-1} \notin \mathcal{A}(2)Sq^k + Sq^{2k-1}\mathcal{A}(2) \\
 \text{for } Sq^{2k-1}Sq^k = 0 \quad (k \geq 1), \\
 l(E) = 8 \quad \text{iff} \quad Sq^{2k-2}Sq^{k-7} \notin \mathcal{A}(2)Sq^{k-6} + \mathcal{A}(2)Sq^{k-4} + \mathcal{A}(2)Sq^k \\
 + Sq^{2k-1}\mathcal{A}(2) + Sq^{2k-3}\mathcal{A}(2) + Sq^{2k-7}\mathcal{A}(2) \\
 \text{for } Sq^{2k-1}Sq^{k-6} + Sq^{2k-3}Sq^{k-4} + Sq^{2k-7}Sq^k = 0 \quad (k \geq 9).
 \end{aligned}$$

AICHI UNIVERSITY OF EDUCATION
 OSAKA UNIVERSITY

References

- [1] J.F. Adams: *On the non-existence of elements of Hopf invariant 1*, Ann. of Math. **72** (1960), 20–104.
- [2] L.L. Larmore and E. Thomas: *Group extensions and principal fibrations*, Math. Scand. **30** (1972), 227–248.
- [3] L.L. Larmore and E. Thomas: *Group extensions and twisted cohomology theories*, Illinois J. Math. **17** (1973), 397–410.
- [4] Y. Nomura: *The Whitney join and its dual*, Osaka J. Math. **7** (1970), 353–373.
- [5] L. Smith: *Secondary cohomology theories*, Indiana Univ. Math. J. **23** (1974), 899–923.
- [6] E. Spanier: *Secondary operations on mappings and cohomology*, Ann. of Math. **75** (1962), 260–282.
- [7] M. Sugawara: *On the homotopy-commutativity of groups and loop spaces*, Mem. Coll. Sci., Univ. of Kyoto, Ser A, **33** (1960), 257–269.
- [8] M. Sugawara: *Order of the identity class of a loop space*, J. Sci. Hiroshima Univ. Ser A-I, **30** (1966), 131–136.
- [9] H. Toda: *Algebra of stable homotopy of Z_p -spaces and applications*, J. Math. Kyoto Univ. **11** (1971), 197–251.