

Title	Theory of multi-fans
Author(s)	Hattori, Akio; Masuda, Mikiya
Citation	Osaka Journal of Mathematics. 2003, 40(1), p. 1– 68
Version Type	VoR
URL	https://doi.org/10.18910/5581
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Hattori, A. and Masuda, M. Osaka J. Math. **40** (2003), 1–68

# THEORY OF MULTI-FANS

AKIO HATTORI and MIKIYA MASUDA

(Received May 1, 2001)

# 1. Introduction

The purpose of the present paper is to develop a theory of multi-fans which is an outgrowth of our study initiated in the work [27] on the topology of torus manifolds (the precise definition will be given later). A multi-fan is a combinatorial object generalizing the notion of a fan in algebraic geometry. Our theory is combinatorial by nature but it is built so as to keep a close connection with the topology of torus manifolds.

It is known that there is a one-to-one correspondence between toric varieties and fans. A toric variety is a normal complex algebraic variety of dimension n with a  $(\mathbb{C}^*)^n$ -action having a dense orbit. The dense orbit is unique and isomorphic to  $(\mathbb{C}^*)^n$ , and other orbits have smaller dimensions. The fan associated with the toric variety is a collection of cones in  $\mathbb{R}^n$  with apex at the origin. To each orbit there corresponds a cone of dimension equal to the codimension of the orbit. Thus the origin is the cone corresponding to the dense orbit, one-dimensional cones correspond to maximal singular orbits and so on. The important point is the fact that the original toric variety can be reconstructed from the associated fan, and algebro-geometric properties of the toric variety can be described in terms of combinatorial data of the associated fan.

If one restricts the action of  $(\mathbb{C}^*)^n$  to the usual torus  $T = (S^1)^n$ , one can still find the fan, because the orbit types of the action of the total group  $(\mathbb{C}^*)^n$  can be detected by the isotropy types of the action of the subgroup T. Take a circle subgroup S of T which appears as an isotropy subgroup of the action. Then each connected component of the closure of the set of those points whose isotropy subgroup equals S is a T-invariant submanifold of real codimension 2, and contains a unique  $(\mathbb{C}^*)^n$  orbit of complex codimension 1. We shall call such a submanifold a characteristic submanifold. If  $M_1, \ldots, M_k$  are characteristic submanifolds such that  $M_1 \cap \cdots \cap M_k$  is nonempty, then the submanifold  $M_1 \cap \cdots \cap M_k$  contains a unique  $(\mathbb{C}^*)^n$ -orbit of complex codimension k. This suggests the following definition of torus manifolds and associated multi-fans.

Let M be an oriented closed manifold of dimension 2n with an effective action of an n dimensional torus T with non-empty fixed point set  $M^T$ . A closed, connected, codimension two submanifold of M will be called characteristic if it is a connected component of the fixed point set of a certain circle subgroup S of T, and if it contains at least one *T*-fixed point. The manifold *M* together with a prefered orientation of each characteristic submanifold will be called a torus manifold. The multi-fan associated with the torus manifold *M* involves cones in the Lie algebra L(T) of *T*, with apex at the origin. If  $M_i$  is a characteristic submanifold and  $S_i$  is the circle subgroup of *T* which pointwise fixes  $M_i$ , then  $S_i$  together with the orientation of  $M_i$  determines an element  $v_i$  of Hom( $S^1, T$ ), and hence a one dimensional cone in the vector space Hom( $S^1, T$ )  $\otimes \mathbb{R}$  canonically identified with L(T). If  $M_{i_1}, \ldots, M_{i_k}$  are characteristic submanifolds such that their intersection contains at least one *T*-fixed point, and if  $v_{i_1}, \ldots, v_{i_k}$  are the corresponding elements in Hom( $S^1, T$ ), then the *k*-dimensional cone spanned by  $v_{i_1}, \ldots, v_{i_k}$  lies in the multi-fan associated with *M*. It should be noted that the intersection of characteristic submanifolds may not be connected in contrast with the case of toric manifolds where the intersection is always connected. For example, the intersection of a family of *n* characteristic submanifolds is a finite set consisting of *T*-fixed points. These data are also incorporated in the definition of the associated multi-fan in Section 2.

One of the differences between a fan and a multi-fan is that, while cones in a fan intersect only at their faces and their union covers the space L(T) just once without overlap for complete toric varieties, it happens that the union of cones in a multi-fan covers L(T) with overlap for torus manifolds. Also the same multi-fan corresponds to different torus manifolds. Nevertheless it turns out that important topological invariants of a torus manifold can be described in terms of the associated multi-fan. In fact it is furthermore possible to develop an abstract theory of multi-fans and to define various "topological" invariants of a multi-fan in such a way that, when the multi-fan is associated with a torus manifold, they coincide with the ordinary topological invariants of the manifold. For example, the "multiplicity of overlap", which we call the degree of the multi-fan, equals the Todd genus for a unitary torus manifold (unitary toric manifold in the terminology in [27]; the precise definition will be given in Section 9).

Another feature of the theory of toric varieties is the correspondence between ample line bundles over a complete toric variety and convex polytopes. From a topological point of view this can be explained in the following way. Let  $(M, \omega)$  be a compact symplectic manifold with a Hamiltonian *T*-action, and let  $\Psi: M \to L(T)^*$  be an associated moment map. Then it is well-known ([1], [16]) that the image *P* of  $\Psi$  is a convex polytope. Moreover, if the de Rham cohomology class of  $\omega$  is an integral class, then the polytope *P* is a lattice polytope up to translations in  $L(T)^*$  identified with  $\mathbb{R}^n$ . Delzant [9] showed that the original symplectic manifold  $(M, \omega)$  is equivariantly symplectomorphic to a complete non-singular toric variety and the form  $\omega$  is transformed into the first Chern form of an ample line bundle *L* over the toric variety. It is known that the number of lattice points in *P* is equal to the Riemann-Roch number

$$\int_M e^{c_1(L)} \mathcal{T}(M)$$

#### THEORY OF MULTI-FANS

where  $\mathcal{T}$  is the Todd class of M, see e.g. [11]. This sort of phenomenon was generalized to "presymplectic" toric manifolds by Karshon and Tolman [22], then to Spin<sup>c</sup> toric manifolds by Grossberg and Karshon [12] and also to unitary toric manifolds by the second-named author [27] in the form which relates the equivariant index of the line bundle L regarded as an element of  $K_T(M)$  to the Duistermaat-Heckman measure of the moment map associated with L. In these extended cases the form  $\omega$  may be degenerate or the line bundle may not be ample, and consequently the image of the moment map may not be convex any longer. This leads us to consider more general figures which we call multi-polytopes. A multi-polytope is a pair of a multi-fan and an arrangement of affine hyperplanes in  $L(T)^*$ . A similar notion was introduced by Karshon and Tolman [22] and also by Khovanskii and Pukhlikov [25] for ordinary fans under the name twisted polytope and virtual polytope respectively. We shall develop a combinatorial theory of multi-polytopes as well; we define the Duistermaat-Heckman measure and the equivariant index in a purely combinatorial fashion for multi-fans and multi-polytopes, and generalize the above results in the combinatorial context. Also we shall introduce a combinatorial counterpart of a moment map which can be used to interpret the combinatorial Duistermaat-Heckman measure.

In carrying out the above program, the use of equivariant homology and cohomology plays an important role. First note that the group  $\text{Hom}(S^1, T)$  can be canonically identified with the equivariant integral homology group  $H_2(BT)$ , and hence the vector space L(T) with  $H_2(BT, \mathbb{R})$ . In this way we regard vectors  $v_i$  in a multi-fan as lying in  $H_2(BT, \mathbb{R})$ . On the other hand a characteristic submanifold  $M_i$  with a fixed orientation determines a cohomology class  $\xi_i$  in  $H_T^2(M)$ , the equivariant Poincaré dual of  $M_i$ . These cohomology classes are fundamental for describing the first Chern class of an equivariant line bundle over M. This fact enables us to associate a multi-polytope and a generalization of the Duistermaat-Heckman measure with an equivariant line bundle. To a T-line bundle L whose equivariant first Chern class has the form  $c_1^T(L) = \sum c_i \xi_i$ , we associate an arrangement of affine hyperplanes  $F_i$  in  $H^2(BT; \mathbb{R}) = L(T)^*$  defined by

$$F_i = \{ u \in H^2(BT; \mathbb{R}) \mid \langle u, v_i \rangle = c_i \}.$$

This arrangement defines the multi-polytope associated with the line bundle L. Moreover it is possible to define the equivariant cohomology of a complete simplicial multifan and extend the results to such abstract multi-fans and multi-polytopes.

If  $v_{i_1}, \ldots, v_{i_n}$  are primitive vectors generating an *n*-dimensional cone in the multifan associated with a torus manifold, then they form a basis of Hom( $S^1, T$ ). However, in the definition of abstract multi-fans, this condition is not postulated. From this point of view, it is natural to deal with torus orbifolds besides torus manifolds. This can be achieved without much change technically. More importantly every complete simplicial multi-fan (the precise definition will be given later) can be realized as a multi-fan associated with a torus orbifold in dimensions greater than 2. In dimensions 1 and 2, realizable multi-fans are characterized.

Concerning the realization problem we are not sure at this moment whether every non-singular complete simplicial multi-fan is realized as the multi-fan associated with a torus manifold. In any case it should be noticed that a multi-fan may correspond to more than one torus manifolds unlike the case of toric varieties.

We now explain the contents of each section. In Section 2 we give a definition of a multi-fan and introduce certain related notions. The completeness of multi-fans is most important. It is a generalization of the notion of completeness of fans. But the definition takes a somewhat sophisticated form. Section 3 is devoted to the  $T_y$ -genus of a complete multi-fan. It is defined in such a way that, when the multi-fan is associated with a unitary torus manifold M, it coincides with the  $T_y$ -genus of M. In Lemma 3.1 we exhibit an equality which is an analogue of the relation between h-vectors and fvectors in combinatorics (see e.g. [32]), and which, we hope, sheds more insight on that relation.

In Sections 4 and 5 the notion of a multi-polytope and the associated Duistermaat-Heckman function are defined. As explained above, a multi-polytope is a pair  $\mathcal{P}$  =  $(\Delta, \mathcal{F})$  of an *n*-dimensional complete multi-fan  $\Delta$  and an arrangement of hyperplanes  $\mathcal{F} = \{F_i\}$  in  $H^2(BT;\mathbb{R})$  with the same index set as the set of 1-dimensional cones in  $\Delta$ . It is called simple if the multi-fan  $\Delta$  is simplicial. The Duistermaat-Heckman function  $DH_{\mathcal{P}}$  associated with a simple multi-polytope  $\mathcal{P}$  is a locally constant integervalued function with bounded support defined on the complement of the hyperplanes  $\{F_i\}$ . The wall crossing formula (Lemma 5.3) which describes the difference of the values of the function on adjacent components is important for later use. In Section 6 another locally constant function on the complement of the hyperplanes  $\{F_i\}$  in a multi-polytope  $\mathcal{P}$ , called the winding number, is introduced. It satisfies a wall crossing formula entirely similar to the Duistermaat-Heckman function. When the multi-fan  $\Delta$ is associated with a torus manifold or a torus orbifold M and if there is an equivariant complex line bundle L over M, then there is a simple multi-polytope  $\mathcal{P}$  naturally associated with L, and the winding number  $WN_{\mathcal{P}}$  is closely related to the moment map of L. In fact it can be regarded as the density function of the Duistermaat-Heckman measure associated with the moment map. Theorem 6.6, the main theorem in Section 6, states that the Duistermaat-Heckman function and the winding number coincide for any simple multi-polytope.

Section 7 is devoted to a generalization of the Ehrhart polynomial to multipolytopes. If P is a convex lattice polytope and if  $\nu P$  denotes the multiplied polytope by a positive integer  $\nu$ , then the number of lattice points  $\sharp(\nu P)$  contained in  $\nu P$  is developed as a polynomial in  $\nu$ . It is called the Ehrhart polynomial of P. The generalization to multi-polytopes is straightforward and properties similar to that of the ordinary Ehrhart polynomial hold (Theorem 7.2). If  $\mathcal{P}$  is a simple lattice multi-polytope, then the associated Ehrhart polynomial  $\sharp(\nu \mathcal{P})$  is defined by

$$\sharp(\nu\mathcal{P}) = \sum_{u \in H^2(BT;\mathbb{Z})} \mathrm{DH}_{\nu\mathcal{P}_+}(u),$$

where  $\mathcal{P}_+$  denotes a multi-polytope obtained from  $\mathcal{P}$  by a small enlargement. Lemma 7.3 is crucial for the proof of Theorem 7.2 and for the later development of the theory. Its corollary, Corollary 7.4, gives a localization formula for the Laurent polynomial  $\sum_{u \in H^2(BT;\mathbb{Z})} DH_{\mathcal{P}_+}(u)t^u$  regarded as a character of T. It can be considered as a combinatorial generalization of Theorem 11.1. It reduces to  $\#\mathcal{P}$  when evaluated at the identity. Using this fact, in Section 8, a cohomological formula expressing  $\#\mathcal{P}$  in terms of the "Todd class" of the multi-fan and the first "Chern class" of the multipolytope is given in Theorem 8.5. The formula can be thought of as a generalization of the formula expressing the number of lattice points in a convex lattice polytope by the Riemann-Roch number of the corresponding ample line bundle. The argument is completely combinatorial. We define the equivariant cohomology  $H_T^*(\Delta)$  of a multi-fan  $\Delta$  which is a module over  $H^*(BT)$ , the index map (Gysin homomorphism)  $\pi_1 \colon H_T^*(\Delta) \to H^{*-2n}(BT)$ , the cohomology  $H^*(\Delta)$  of  $\Delta$  and finally the evaluation on the "fundamental class". As a corollary a generalization of Khovanskii-Pukhlikov formula ([25]) for simple lattice multi-polytopes is given in Theorem 8.7.

In Section 9 it is shown how to associate a multi-fan with a torus manifold. It is also shown that the associated multi-fan is complete. Then, in Section 10, the  $T_y$ -genus of a general torus manifold is defined and is proved to coincide with the  $T_y$ -genus of the associated multi-fan in Theorem 10.1. As a corollary a formula for the signature of a torus manifold is given. In the same spirit the definition of the equivariant index of a line bundle over a general torus manifold is given in Section 11 using a localization formula which holds in the case of unitary torus manifolds. The main theorem of this section, Theorem 11.1, gives a formula describing that equivariant index using the winding number. It generalizes the results of [22], [12] and [27] as indicated before. Results of Section 5 and 6 are crucially used here.

In Section 12 necessary changes to deal with torus orbifolds are explained briefly. One of the remarkable points is that the torus action and the orbifold structure are closely related to each other for a torus orbifold as is explained in Lemma 12.3. In the last section the realization problem is dealt with. Main results of the section are Theorems 13.1, 13.2 and 13.3.

# 2. Multi-fans

In [27], we introduced the notion of a unitary toric manifold, which contains a compact non-singular toric variety as an example, and associated with it a combinatorial object called a multi-fan, which is a more general notion than a complete non-singular fan. In this section, we define a multi-fan in a combinatorial way and in full generality. The reader will find that our notion of a multi-fan is a complete general-

ization of a fan. We also define the completeness and non-singularity of a multi-fan, which generalize the corresponding notion of a fan. To do this, we begin with reviewing the definition of a fan.

Let N be a lattice of rank n, which is isomorphic to  $\mathbb{Z}^n$ . We denote the real vector space  $N \otimes \mathbb{R}$  by  $N_{\mathbb{R}}$ . A subset  $\sigma$  of  $N_{\mathbb{R}}$  is called a *strongly convex rational polyhedral cone* (with apex at the origin) if there exits a finite number of vectors  $v_1, \ldots, v_m$  in N such that

$$\sigma = \{r_1v_1 + \dots + r_mv_m \mid r_i \in \mathbb{R} \text{ and } r_i \geq 0 \text{ for all } i\}$$

and  $\sigma \cap (-\sigma) = \{0\}$ . Here "rational" means that it is generated by vectors in the lattice N, and "strong" convexity means that it contains no line through the origin. We will often call a strongly convex rational polyhedral cone in  $N_{\mathbb{R}}$  simply a *cone* in N. The dimension dim  $\sigma$  of a cone  $\sigma$  is the dimension of the linear space spanned by vectors in  $\sigma$ . A subset  $\tau$  of  $\sigma$  is called a *face* of  $\sigma$  if there is a linear function  $l: N_{\mathbb{R}} \to \mathbb{R}$  such that l takes nonnegative values on  $\sigma$  and  $\tau = l^{-1}(0) \cap \sigma$ . A cone is regarded as a face of itself, while others are called *proper* faces.

DEFINITION. A fan  $\Delta$  in N is a set of a finite number of strongly convex rational polyhedral cones in  $N_{\mathbb{R}}$  such that

- (1) Each face of a cone in  $\Delta$  is also a cone in  $\Delta$ ;
- (2) The intersection of two cones in  $\Delta$  is a face of each.

DEFINITION. A fan  $\Delta$  is said to be *complete* if the union of cones in  $\Delta$  covers the entire space  $N_{\mathbb{R}}$ .

A cone is called *simplicial* if it is generated by linearly independent vectors. If the generating vectors can be taken as a part of a basis of N, then the cone is called *non-singular*.

DEFINITION. A fan  $\Delta$  is said to be *simplicial* (resp. *non-singular*) if every cone in  $\Delta$  is simplicial (resp. non-singular).

The basic theory of toric varieties tells us that a fan is complete (resp. simplicial or non-singular) if and only if the corresponding toric variety is compact (resp. an orbifold or non-singular).

For each  $\tau \in \Delta$ , we define  $N^{\tau}$  to be the quotient lattice of N by the sublattice generated (as a group) by  $\tau \cap N$ ; so the rank of  $N^{\tau}$  is  $n - \dim \tau$ . We consider cones in  $\Delta$  that contain  $\tau$  as a face, and project them on  $(N^{\tau})_{\mathbb{R}}$ . These projected cones form a fan in  $N^{\tau}$ , which we denote by  $\Delta_{\tau}$  and call the *projected fan* with respect to  $\tau$ . The dimensions of the projected cones decrease by dim  $\tau$ . The completeness, simpliciality and non-singularity of  $\Delta$  are inherited to  $\Delta_{\tau}$  for any  $\tau$ . We now generalize these notions of a fan. Let N be as before. Denote by Cone(N) the set of all cones in N. An ordinary fan is a subset of Cone(N). The set Cone(N) has a (strict) partial ordering  $\prec$  defined by:  $\tau \prec \sigma$  if and only if  $\tau$  is a proper face of  $\sigma$ . The cone  $\{0\}$  consisting of the origin is the unique minimum element in Cone(N). On the other hand, let  $\Sigma$  be a partially ordered finite set with a unique minimum element. We denote the (strict) partial ordering by < and the minimum element by \*. An example of  $\Sigma$  used later is an abstract simplicial set with an empty set added as a member, which we call an *augmented simplicial set*. In this case the partial ordering is defined by the inclusion relation and the empty set is the unique minimum element which may be considered as a (-1)-simplex. Suppose that there is a map

$$C: \Sigma \to \operatorname{Cone}(N)$$

such that

(1)  $C(*) = \{0\};$ 

(2) If I < J for  $I, J \in \Sigma$ , then  $C(I) \prec C(J)$ ;

(3) For any  $J \in \Sigma$  the map *C* restricted on  $\{I \in \Sigma \mid I \leq J\}$  is an isomorphism of ordered sets onto  $\{\kappa \in \text{Cone}(N) \mid \kappa \preceq C(J)\}$ .

For an integer m such that  $0 \le m \le n$ , we set

$$\Sigma^{(m)} := \{ I \in \Sigma \mid \dim C(I) = m \}.$$

One can easily check that  $\Sigma^{(m)}$  does not depend on *C*. When  $\Sigma$  is an augmented simplicial set,  $I \in \Sigma$  belongs to  $\Sigma^{(m)}$  if and only if the cardinality |I| of *I* is *m*, namely *I* is an (m-1)-simplex. Therefore, even if  $\Sigma$  is not an augmented simplicial set, we use the notation |I| for *m* when  $I \in \Sigma^{(m)}$ .

The image  $C(\Sigma)$  is a finite set of cones in N. We may think of a pair  $(\Sigma, C)$  as a set of cones in N labeled by the ordered set  $\Sigma$ . Cones in an ordinary fan intersect only at their faces, but cones in  $C(\Sigma)$  may overlap, even the same cone may appear repeatedly with different labels. The pair  $(\Sigma, C)$  is almost what we call a multi-fan, but we incorporate a pair of weight functions on cones in  $C(\Sigma)$  of the highest dimension  $n = \operatorname{rank} N$ . More precisely, we consider two functions

$$w^{\pm} \colon \Sigma^{(n)} \to \mathbb{Z}_{>0}.$$

We assume that  $w^+(I) > 0$  or  $w^-(I) > 0$  for every  $I \in \Sigma^{(n)}$ . These two functions have its origin from geometry. In fact if M is a torus manifold of dimension 2n and if  $M_{i_1}, \ldots, M_{i_n}$  are characteristic submanifolds such that their intersection contains at least one T-fixed point, then the intersection  $M_I = \bigcap_{\nu} M_{i_{\nu}}$  consists of a finite number of T-fixed points. At each fixed point  $p \in M_I$  the tangent space  $\tau_p$  has two orientations; one is endowed by the orientation of M and the other comes from the intersection of the oriented submanifolds  $M_{i_{\nu}}$ . Denoting the ratio of the above two orientations by  $\epsilon_p$  we define the number  $w^+(I)$  to be the number of points  $p \in M_I$  with  $\epsilon_p = +1$ and similarly for  $w^-(I)$ . More detailed explanation will be given in Section 9.

DEFINITION. We call a triple  $\Delta := (\Sigma, C, w^{\pm})$  a *multi-fan* in N. We define the dimension of  $\Delta$  to be the rank of N (or the dimension of  $N_{\mathbb{R}}$ ).

Since an ordinary fan  $\Delta$  in N is a subset of Cone(N), one can view it as a multifan by taking  $\Sigma = \Delta$ , C = the inclusion map,  $w^+ = 1$ , and  $w^- = 0$ . In a similar way as in the case of ordinary fans, we say that a multi-fan  $\Delta = (\Sigma, C, w^{\pm})$  is *simplicial* (resp. *non-singular*) if every cone in  $C(\Sigma)$  is simplicial (resp. non-singular). The following lemma is easy.

**Lemma 2.1.** A multi-fan  $\Delta = (\Sigma, C, w^{\pm})$  is simplicial if and only if  $\Sigma$  is isomorphic to an augmented simplicial set as partially ordered sets.

The definition of completeness of a multi-fan  $\Delta$  is rather complicated. A naive definition of the completeness would be that the union of cones in  $C(\Sigma)$  covers the entire space  $N_{\mathbb{R}}$ . But it turns out that this is not a right definition if we look at multi-fans associated with unitary torus manifolds, see Section 9. Although the two weighted functions  $w^{\pm}$  are incorporated in the definition of a multi-fan, only the difference

$$w := w^{+} - w^{-}$$

matters in this paper except Section 13. We shall introduce the following intermediate notion of pre-completeness at first. A vector  $v \in N_{\mathbb{R}}$  will be called generic if v does not lie on any linear subspace spanned by a cone in  $C(\Sigma)$  of dimension less than n. For a generic vector v we set  $d_v = \sum_{v \in C(I)} w(I)$ , where the sum is understood to be zero if there is no such I.

DEFINITION. We call a multi-fan  $\Delta = (\Sigma, C, w^{\pm})$  of dimension *n* pre-complete if  $\Sigma^{(n)} \neq \emptyset$  and the integer  $d_v$  is independent of the choice of generic vectors v. We call this integer the *degree* of  $\Delta$  and denote it by deg( $\Delta$ ).

REMARK. For an ordinary fan, pre-completeness is the same as completeness.

To define the completeness for a multi-fan  $\Delta$ , we need to define a projected multifan with respect to an element in  $\Sigma$ . We do it as follows. For each  $K \in \Sigma$ , we set

$$\Sigma_K \coloneqq \{J \in \Sigma \mid K \leq J\}.$$

It inherits the partial ordering from  $\Sigma$ , and K is the unique minimum element in  $\Sigma_K$ .



Fig. 1.

A map

$$C_K \colon \Sigma_K \to \operatorname{Cone}(N^{C(K)})$$

sending  $J \in \Sigma_K$  to the cone C(J) projected on  $(N^{C(K)})_{\mathbb{R}}$  satisfies the three properties above required for *C*. We define two functions

$$w_K^{\pm} \colon \Sigma_K^{(n-|K|)} \subset \Sigma^{(n)} \to \mathbb{Z}_{\geq 0}$$

to be the restrictions of  $w^{\pm}$  to  $\Sigma_K^{(n-|K|)}$ . The triple  $\Delta_K := (\Sigma_K, C_K, w_K^{\pm})$  is a multifan in  $N^{C(K)}$ , and this is the desired *projected multi-fan* with respect to  $K \in \Sigma$ . When  $\Delta$  is an ordinary fan, this definition agrees with the previous one.

DEFINITION. A pre-complete multi-fan  $\Delta = (\Sigma, C, w^{\pm})$  is said to be *complete* if the projected multi-fan  $\Delta_K$  is pre-complete for any  $K \in \Sigma$ .

REMARK. A multi-fan  $\Delta$  is complete if and only if the projected multi-fan  $\Delta_J$  is pre-complete for any  $J \in \Sigma^{(n-1)}$ . The argument is as follows. The pre-completeness of  $\Delta_J$  for  $J \in \Sigma^{(n-1)}$  implies that  $d_v = \sum_{v \in C(I)} w(I)$  remains unchanged when v gets across the codimension one cone C(J), which means the pre-completeness of  $\Delta$ . Since  $\Sigma_K^{(n-|K|-1)}$  is contained in  $\Sigma^{(n-1)}$  for any  $K \in \Sigma$ , the pre-completeness of  $\Delta_J$  for any  $J \in \Sigma^{(n-1)}$  also implies the pre-completeness of  $\Delta_K$  for any  $K \in \Sigma$ .

EXAMPLE 2.2. Here is an example of a complete non-singular multi-fan of degree two. Let  $v_1, \ldots, v_5$  be integral vectors shown in Fig. 1, where the dots denote lattice points.

The vectors are rotating around the origin twice in counterclockwise. We take

 $\Sigma = \{\phi, \{1\}, \dots, \{5\}, \{1, 2\}, \{2, 3\}, \{3, 4\}, \{4, 5\}, \{5, 1\}\},\$ 



Fig. 2.

define  $C: \Sigma \to \operatorname{Cone}(N)$  by

$$C(\{i\}) =$$
 the cone spanned by  $v_i$ ,  
 $C(\{i, i+1\}) =$  the cone spanned by  $v_i$  and  $v_{i+1}$ 

where i = 1, ..., 5 and 6 is understood to be 1, and take  $w^{\pm}$  such that w = 1 on every two dimensional cone. Then  $\Delta = (\Sigma, C, w^{\pm})$  is a complete non-singular two-dimensional multi-fan with deg $(\Delta) = 2$ .

EXAMPLE 2.3. Here is an example of a complete multi-fan "with folds". Let  $v_1, \ldots, v_5$  be vectors shown in Fig. 2.

We take the same  $\Sigma$  and C as in Example 2.2 and take  $w^\pm$  such that

$$w(\{3,4\}) = -1$$
 and  $w(\{i,i+1\}) = 1$  for  $i \neq 3$ .

Then  $\Delta = (\Sigma, C, w^{\pm})$  is a complete two-dimensional multi-fan with deg( $\Delta$ ) = 1.

A similar example can be constructed for a number of vectors  $v_1, \ldots, v_d$   $(d \ge 3)$  by defining

 $w(\{i, i+1\}) = 1$  if  $v_i$  and  $v_{i+1}$  are rotating in counterclockwise,  $w(\{i, i+1\}) = -1$  if  $v_i$  and  $v_{i+1}$  are rotating in clockwise,

where d + 1 is understood to be 1. The degree deg( $\Delta$ ) is the rotation number of the vectors  $v_1, \ldots, v_d$  around the origin in counterclockwise and may not be one.

EXAMPLE 2.4. Here is an example of a multi-fan which is pre-complete but not complete. Let  $v_1, \ldots, v_5$  be vectors shown in Fig. 3.

We take

$$\Sigma = \{\phi, \{1\}, \dots, \{5\}, \{1, 2\}, \{2, 3\}, \{3, 1\}, \{4, 5\}\},\$$



Fig. 3.

define  $C: \Sigma \to \operatorname{Cone}(N)$  as in Example 2.2, and take  $w^{\pm}$  such that

$$w(\{1,2\}) = 2, w(\{2,3\}) = 1, w(\{3,1\}) = 1, w(\{4,5\}) = -1.$$

Then  $\Delta = (\Sigma, C, w^{\pm})$  is a two-dimensional multi-fan which is pre-complete (in fact, deg( $\Delta$ ) = 1) but not complete because the projected multi-fan  $\Delta_{\{i\}}$  for  $i \neq 3$  is not pre-complete.

So far, we treated *rational* cones that are generated by vectors in the lattice N. But, most of the notions introduced above make sense even if we allow cones generated by vectors in  $N_{\mathbb{R}}$  which may not be in N. In fact, the notion of non-singularity requires the lattice N, but others do not. Therefore, one can define a multi-fan and its completeness and simpliciality in this extended category as well. The reader will find that the arguments developed in Sections 3 through 6 work in this extended category.

### 3. $T_y$ -genus of a multi-fan

A unitary torus manifold M determines a complete non-singular multi-fan. (This will be discussed and extended to torus manifolds in Section 9.) On the other hand, the  $T_y$ -genus (also called  $\chi_y$ -genus) for unitary manifolds introduced by Hirzebruch in his famous book [20] is defined for M. Its characteristic power series is given by  $x(1 + ye^{-(1+y)x})/(1 - e^{-(1+y)x})$ . It is a polynomial in one variable y of degree (at most)  $(1/2) \dim M$ . The Kosniowski formula about the  $T_y$ -genus for unitary  $S^1$ -manifolds (see [18], [23]) and the results in [27] imply that the  $T_y$ -genus of M should be described in terms of the multi-fan associated with M. In this section (and in Section 10) we give the explicit description. In fact, our argument is rather more general. We think of the  $T_y$ -genus of M as a polynomial invariant of the associated multi-fan which is complete and non-singular. It turns out that the polynomial invariant can be defined not only for the multi-fans associated with unitary torus manifolds but also for all com-

plete simplicial multi-fans.

Since the lattice N is unnecessary from now until the end of Section 6, we shall denote the vector space, in which cones sit, by V instead of  $N_{\mathbb{R}}$ . Let  $\Delta = (\Sigma, C, w^{\pm})$ be a complete simplicial multi-fan defined on V. By Lemma 2.1 we may assume that  $\Sigma$  is an augmented simplicial set, say, consisting of subsets of  $\{1, \ldots, d\}$  and  $\Sigma^{(1)} =$  $\{\{1\}, \ldots, \{d\}\}$  where d is the number of elements in  $\Sigma^{(1)}$ . For each  $i = 1, \ldots, d$ , let  $v_i$  denote a nonzero vector in the one-dimensional cone  $C(\{i\})$ . Choose a generic vector  $v \in V$ . Let  $I \in \Sigma^{(n)}$ . Since  $v_i$ 's  $(i \in I)$  are linearly independent, v has a unique expression  $\sum_{i \in I} a_i v_i$  with real numbers  $a_i$ 's. The coefficients  $a_i$ 's are all nonzero because v is generic. We set

$$\mu(I) := \sharp \{ i \in I \mid a_i > 0 \}.$$

This depends on v although v is not recorded in the notation  $\mu(I)$ .

DEFINITION. For an integer q with  $0 \le q \le n$ , we define

$$h_q(\Delta) \coloneqq \sum_{\mu(I)=q} w(I) \text{ and } e_q(\Delta) \coloneqq \sum_{K \in \Sigma^{(q)}} \deg(\Delta_K).$$

Note that  $h_n(\Delta) = \deg(\Delta) = e_0(\Delta)$ , and  $e_q(\Delta)$ 's are independent of v. If  $\Delta$  is a complete simplicial multi-fan such that  $\deg(\Delta) = 1$  and w(I) = 1 for all  $I \in \Sigma^{(n)}$  (e.g. this is the case if  $\Delta$  is a complete simplicial ordinary fan), then  $\deg(\Delta_K)$  equals 1 for all  $K \in \Sigma$  and hence  $e_q(\Delta)$  agrees with the number of cones of dimension q in the multi-fan.

The following lemma reminds us of the relation between the h-vectors and the f-vectors for simplicial sets studied in combinatorics (see [32]).

**Lemma 3.1.** 
$$\sum_{q=0}^{n} h_q(\Delta)(s+1)^q = \sum_{m=0}^{n} e_{n-m}(\Delta)s^m \text{ where } s \text{ is an indeterminate.}$$

Proof. The lemma is equivalent to the following equality:

(3.1) 
$$\sum_{q=m}^{n} h_q(\Delta) \begin{pmatrix} q \\ m \end{pmatrix} = e_{n-m}(\Delta).$$

It follows from the definition of  $h_q(\Delta)$  that

On the other hand, we shall rewrite  $e_{n-m}(\Delta)$ . It follows from the definition of

 $deg(\Delta_K)$  that

$$\deg(\Delta_K) = \sum_{J \in \Sigma_K^{(n-|K|)} s.t. \ v_K \in C_K(J)} w_K(J)$$

where  $v_K$  denotes the projection image of v on the quotient vector space of V by the subspace  $V_K$  spanned by the cone C(K). Note that  $v_K$  lies in  $C_K(J)$  if and only if v lies in  $C(J \cup K)$  modulo  $V_K$ , and that  $w_K(J) = w(J \cup K)$  by definition. Therefore, writing  $J \cup K$  as I, the equality above turns into

$$\deg(\Delta_K) = \sum_I w(I),$$

where *I* runs over elements in  $\Sigma^{(n)}$  such that  $K \subset I$  and  $v \in C(I)$  modulo  $V_K$ . Putting this in the defining equation of  $e_{n-m}(\Delta)$ , we have

(3.3) 
$$e_{n-m}(\Delta) = \sum_{K,I} w(I),$$

where the sum is taken over elements  $K \in \Sigma^{(n-m)}$  and  $I \in \Sigma^{(n)}$  such that  $K \subset I$  and  $v \in C(I)$  modulo  $V_K$ . Fix  $I \in \Sigma^{(n)}$  with  $\mu(I) = q$ , and observe how many times I appears in the above sum. It is equal to the number of  $K \in \Sigma^{(n-m)}$  such that  $K \subset I$  and  $v \in C(I)$  modulo  $V_K$ . But the number of such K is  $\binom{q}{m}$ . To see this, we note that  $\mu(I) = q$  means that  $\sharp\{i \in I \mid a_i > 0\} = q$  by definition, where  $v = \sum_{i \in I} a_i v_i$ , and that the condition that  $v \in C(I)$  modulo  $V_K$  is equivalent to saying that K contains the complement of the set  $\{i \in I \mid a_i > 0\}$  in I. Therefore, any such K is obtained as the complement of a subset of  $\{i \in I \mid a_i > 0\}$  with cardinality m, so that the number of such K is  $\binom{q}{m}$ . This together with (3.2) and (3.3) proves the equality (3.1).

#### Corollary 3.2.

- (1)  $h_q(\Delta)$ 's are independent of the choice of the generic vector v.
- (2)  $h_q(\Delta) = h_{n-q}(\Delta)$  for any q.

Proof. (1) This immediately follows from Lemma 3.1 because  $e_q(\Delta)$ 's are independent of v.

(2) If we take -v instead of v, then  $\mu(I)$  turns into  $n - \mu(I)$ , so that  $h_q(\Delta)$  turns into  $h_{n-q}(\Delta)$ . Since  $h_q(\Delta)$ 's are independent of v as shown in (1) above, this proves  $h_q(\Delta) = h_{n-q}(\Delta)$ .

When  $\Delta$  is associated with a unitary torus manifold M, the  $T_y$ -genus of M turns out to be given by  $\sum_{q=0}^{n} h_q(\Delta)(-y)^q$ . (This will be discussed in Section 10 later.) Motivated by this observation,

DEFINITION. For a complete simplicial multi-fan  $\Delta$ , we define

$$T_{y}[\Delta] \coloneqq \sum_{q=0}^{n} h_{q}(\Delta)(-y)^{q}$$

and call it the  $T_y$ -genus of  $\Delta$ . Note that  $T_0[\Delta] = h_0(\Delta) = h_n(\Delta) = \deg(\Delta)$ .

Lemma 3.1 can be restated as

**Corollary 3.3.** Let  $\Delta$  be a complete simplicial multi-fan. Then

$$T_{y}[\Delta] = \sum_{m=0}^{n} e_{n-m}(\Delta)(-1-y)^{m}.$$

#### 4. Multi-polytopes

A convex polytope P in  $V^* = \text{Hom}(V, \mathbb{R})$  is the convex hull of a finite set of points in  $V^*$ . It is the intersection of a finite number of half spaces in  $V^*$  separated by affine hyperplanes, so there are a finite number of nonzero vectors  $v_1, \ldots, v_d$  in V and real numbers  $c_1, \ldots, c_d$  such that

$$P = \{ u \in V^* \mid \langle u, v_i \rangle \le c_i \text{ for all } i \},\$$

where  $\langle , \rangle$  denotes the natural pairing between  $V^*$  and V. (Warning: In this paper, we take  $v_i$  to be "outward normal" to the corresponding face of P contrary to the usual convention in algebraic geometry, cf. e.g. [29].) The convex polytope P can be recovered from the data  $\{(v_i, c_i) \mid i = 1, ..., d\}$ . But, a more general figure like Qshaded in Fig. 4 cannot be determined by the data  $\{(v_i, c_i) \mid i = 1, ..., d\}$ . We need to prescribe the vertices of Q, in other words, which pairs of lines  $l_i$ 's are presumed to intersect. For instance, if four points  $l_1 \cap l_2$ ,  $l_2 \cap l_3$ ,  $l_3 \cap l_4$  and  $l_4 \cap l_1$  are presumed to be vertices (and the others such as  $l_2 \cap l_4$  are not), then we can find the figure Qin Fig. 4. But, if different four points  $l_1 \cap l_4$ ,  $l_4 \cap l_2$ ,  $l_2 \cap l_3$  and  $l_3 \cap l_1$  are presumed to be vertices, then we obtain a figure Q' shaded in Fig. 4.

The data of whether two lines  $l_i$  and  $l_j$  are presumed to intersect is equivalent to the data of whether the corresponding vectors  $v_i$  and  $v_j$  span a cone. In the former (resp. latter) example above, resulting cones are four two-dimensional ones shown in Fig. 5 (1) (resp. (2)). Needless to say,  $l_i$  is 'perpendicular' to the half line spanned by  $v_i$ .

A polytope gives rise to a multi-fan in this way. One notes that a convex polytope gives rise to a complete fan. Taking this observation into account, we reverse a gear. We start with a complete multi-fan  $\Delta = (\Sigma, C, w^{\pm})$ . Let HP(V<sup>\*</sup>) be the set of all affine hyperplanes in V<sup>\*</sup>.

THEORY OF MULTI-FANS







Fig. 5.

DEFINITION. Let  $\Delta = (\Sigma, C, w^{\pm})$  be a complete multi-fan and let  $\mathcal{F}: \Sigma^{(1)} \to$ HP( $V^*$ ) be a map such that the affine hyperplane  $\mathcal{F}(I)$  is 'perpendicular' to the half line C(I) for each  $I \in \Sigma^{(1)}$ , i.e., an element in C(I) takes a constant on  $\mathcal{F}(I)$ . We call a pair  $(\Delta, \mathcal{F})$  a *multi-polytope* and denote it by  $\mathcal{P}$ . The dimension of a multi-polytope  $\mathcal{P}$  is defined to be the dimension of the multi-fan  $\Delta$ . We say that a multi-polytope  $\mathcal{P}$ is *simple* if  $\Delta$  is simplicial.

REMARK. The completeness assumption for  $\Delta$  is not needed for the definition of multi-polytopes. We incorporated it because most of our results depend on that assumption. Similar notions were introduced by Karshon-Tolman [22] and Khovanskii-

Pukhlikov [25] when  $\Delta$  is an ordinary fan. They use the terminology *twisted polytope* and *virtual polytope* respectively. The notion of multi-polytopes is a direct generalization of that of twisted polytopes. The relation between virtual polytopes and multi-polytopes is clarified by [28].

EXAMPLE 4.1. A convex polytope determines a complete fan together with an arrangement of affine hyperplanes containing the facets of the polytope (as explained above), so it uniquely determines a multi-polytope.

EXAMPLE 4.2. Associated with the multi-fan in Example 2.2, one obtains the arrangement of lines drawn in Fig. 6 with a suitable choice of the map  $\mathcal{F}$ . The pentagon shown up in Fig. 6 produces the same arrangement of lines and can be viewed as a multi-polytope as explained in Example 4.1 above, but these two multi-polytopes are different because the underlying multi-fans are different; one is a multi-fan of degree two while the other is an ordinary fan. The reader will find a star-shaped figure in the former multi-polytope.



# 5. Duistermaat-Heckman functions

A multi-polytope  $\mathcal{P} = (\Delta, \mathcal{F})$  defines an arrangement of affine hyperplanes in  $V^*$ . In this section, we associate with  $\mathcal{P}$  a function on  $V^*$  minus the affine hyperplanes when  $\mathcal{P}$  is simple. This function is locally constant and Guillemin-Lerman-Sternberg formula ([14], [15]) tells us that it agrees with the density function of a Duistermaat-Heckman measure when  $\mathcal{P}$  arises from a moment map.

Hereafter our multi-polytope  $\mathcal{P}$  is assumed to be simple, so that the multi-fan  $\Delta = (\Sigma, C, w^{\pm})$  is complete and simplicial unless otherwise stated. As before, we may assume that  $\Sigma$  consists of subsets of  $\{1, \ldots, d\}$  and  $\Sigma^{(1)} = \{\{1\}, \ldots, \{d\}\}$ , and denote by  $v_i$  a nonzero vector in the one-dimensional cone  $C(\{i\})$ . To simplify notation, we denote  $\mathcal{F}(\{i\})$  by  $F_i$  and set

$$F_I \coloneqq \bigcap_{i \in I} F_i \quad \text{for } I \in \Sigma.$$

 $F_I$  is an affine space of dimension n - |I|. In particular, if |I| = n (i.e.,  $I \in \Sigma^{(n)}$ ), then  $F_I$  is a point, denoted by  $u_I$ .

Suppose  $I \in \Sigma^{(n)}$ . Then the set  $\{v_i \mid i \in I\}$  forms a basis of V. Denote its dual basis of  $V^*$  by  $\{u_i^I \mid i \in I\}$ , i.e.,  $\langle u_i^I, v_j \rangle = \delta_{ij}$  where  $\delta_{ij}$  denotes the Kronecker delta. Take a generic vector  $v \in V$ . Then  $\langle u_i^I, v \rangle \neq 0$  for all  $I \in \Sigma^{(n)}$  and  $i \in I$ . Set

$$(-1)^{I} := (-1)^{\sharp\{i \in I \mid \langle u_{i}^{I}, v \rangle > 0\}} \quad \text{and} \quad (u_{i}^{I})^{+} := \begin{cases} u_{i}^{I} & \text{if } \langle u_{i}^{I}, v \rangle > 0\\ -u_{i}^{I} & \text{if } \langle u_{i}^{I}, v \rangle < 0 \end{cases}$$

We denote by  $C^*(I)^+$  the cone in  $V^*$  spanned by  $(u_i^I)^+$ 's  $(i \in I)$  with apex at  $u_I$ , and by  $\phi_I$  its characteristic function.

DEFINITION. We define a function  $DH_{\mathcal{P}}$  on  $V^* \setminus \bigcup_{i=1}^d F_i$  by

$$\mathsf{DH}_{\mathcal{P}} \coloneqq \sum_{I \in \Sigma^{(n)}} (-1)^{I} w(I) \phi_{I}$$

and call it the Duistermaat-Heckman function associated with  $\mathcal{P}$ .

REMARK. Apparently, the function  $DH_{\mathcal{P}}$  is defined on the whole space  $V^*$  and depends on the choice of the generic vector  $v \in V$ , but we will see in Lemma 5.4 below that it is independent of v on  $V^* \setminus \bigcup F_i$ . This is the reason why we restricted the domain of the function to  $V^* \setminus \bigcup F_i$ .

For the moment, we shall see the independence of v when dim  $\mathcal{P} = 1$ .

EXAMPLE 5.1. Suppose dim  $\mathcal{P} = 1$ . We identify V with  $\mathbb{R}$ , so that  $V^*$  is also identified with  $\mathbb{R}$ . Let E be the subset of  $\{1, \ldots, d\}$  such that  $i \in E$  if and only if  $C(\{i\})$  is the half line consisting of nonnegative real numbers. Then the completeness of  $\Delta$  means that

(5.1) 
$$\sum_{i \in E} w(\{i\}) = \sum_{i \notin E} w(\{i\}) = \deg(\Delta).$$

Take a nonzero vector v. Since  $V^*$  is identified with  $\mathbb{R}$ , each affine hyperplane  $F_i$  is nothing but a real number. Suppose that v is toward the positive direction. Then

(5.2) 
$$(-1)^{\{i\}} = \begin{cases} -1 & \text{if } i \in E \\ 1 & \text{if } i \notin E \end{cases}$$

and the support of the characteristic function  $\phi_{\{i\}}$  is the half line given by

$$\{u \in \mathbb{R} \mid F_i \leq u\}.$$

Therefore

(5.3) 
$$DH_{\mathcal{P}}(u) = \sum_{i \in E \ s.t. \ F_i < u} (-w(\{i\})) + \sum_{i \notin E \ s.t. \ F_i < u} w(\{i\})$$

for  $u \in \mathbb{R} \setminus \bigcup F_i$ . If *u* is sufficiently small, then the sum above is empty; so it is zero. If *u* is sufficiently large, then the sum is also zero by (5.1). Hence the support of the function  $DH_{\mathcal{P}}$  is bounded.

Now, suppose that v is toward the negative direction. Then  $(-1)^{\{i\}}$  above is multiplied by -1 and the inequality  $\leq$  above turns into  $\geq$ . Therefore

(5.4) 
$$DH_{\mathcal{P}}(u) = \sum_{i \in E \ s.t. \ u < F_i} w(\{i\}) + \sum_{i \notin E \ s.t. \ u < F_i} (-w(\{i\})).$$

It follows that

r.h.s. of (5.3) – r.h.s. of (5.4) = 
$$-\sum_{i \in E} w(\{i\}) + \sum_{i \notin E} w(\{i\}),$$

which is zero by (5.1). This shows that the function  $DH_{\mathcal{P}}$  is independent of v when dim  $\mathcal{P} = 1$ .

EXAMPLE 5.2. For the star-shaped multi-polytope in Example 4.2,  $DH_{\mathcal{P}}$  takes 2 on the pentagon, 1 on the five triangles adjacent to the pentagon and 0 on other (unbounded) regions. The check is left to the reader.

Assume  $n = \dim \Delta > 1$ . For each  $\{i\} \in \Sigma^{(1)}$ , the projected multi-fan  $\Delta_{\{i\}} = (\Sigma_{\{i\}}, C_{\{i\}}, w_{\{i\}}^{\pm})$ , which we abbreviate as  $\Delta_i = (\Sigma_i, C_i, w_i^{\pm})$ , is defined on the quotient vector space  $V/V_i$  of V by the one-dimensional subspace  $V_i$  spanned by  $v_i$ . Since  $\Delta$  is complete and simplicial, so is  $\Delta_i$ . We identify the dual space  $(V/V_i)^*$  with

$$(V^*)_i := \{ u \in V^* \mid \langle u, v_i \rangle = 0 \}$$

in a natural way. We choose an element  $f_i \in F_i$  arbitrarily and translate  $F_i$  onto  $(V^*)_i$ by  $-f_i$ . If  $\{i, j\} \in \Sigma^{(2)}$ , then  $F_j$  intersects  $F_i$  and their intersection will be translated



Fig. 7.

into  $(V^*)_i$  by  $-f_i$ . This observation leads us to consider the map

$$\mathcal{F}_i: \Sigma_i \to \operatorname{HP}((V^*)_i)$$

sending  $\{j\} \in \Sigma_i^{(1)}$  to  $F_i \cap F_j$  translated by  $-f_i$ . The pair  $\mathcal{P}_i = (\Delta_i, \mathcal{F}_i)$  is a multipolytope in  $(V/V_i)^* \cong (V^*)_i$ .

Let  $I \in \Sigma^{(n)}$  such that  $i \in I$ . Since  $\langle u_j^I, v_i \rangle = \delta_{ij}, u_j^I$  for  $j \neq i$  is an element of  $(V^*)_i$ , which we also regard as an element of  $(V/V_i)^*$  through the isomorphism  $(V/V_i)^* \cong (V^*)_i$ . We denote the projection image of the generic element  $v \in V$  on  $V/V_i$  by  $\bar{v}$ . Then we have  $\langle \bar{v}, u_j^I \rangle = \langle v, u_j^I \rangle$  for  $j \neq i$ , where  $u_j^I$  at the left-hand side is viewed as an element of  $(V/V_i)^*$  while the one at the right-hand side is viewed as an element of  $(V^*)_i$ . Since  $\langle \bar{v}, u_j^I \rangle = \langle v, u_j^I \rangle \neq 0$  for  $j \neq i$ , we use  $\bar{v}$  to define  $DH_{\mathcal{P}_i}$ .

**Lemma 5.3** (Wall crossing formula.). Let F be one of  $F_i$ 's. Let  $u_\alpha$  and  $u_\beta$  be elements in  $V^* \setminus \bigcup_{i=1}^d F_i$  such that the segment from  $u_\alpha$  to  $u_\beta$  intersects the wall F transversely at  $\mu$ , and does not intersect any other  $F_j \neq F$ . Then

$$\mathrm{DH}_{\mathcal{P}}(u_{\alpha}) - \mathrm{DH}_{\mathcal{P}}(u_{\beta}) = \sum_{i:F_i=F} \mathrm{sign} \langle u_{\beta} - u_{\alpha}, v_i \rangle \mathrm{DH}_{\mathcal{P}_i}(\mu - f_i).$$

Proof. For simplicity we assume that there is only one *i* such that  $F_i = F$ . We may assume that  $\langle u_\beta - u_\alpha, v_i \rangle$  is positive without loss of generality. The situation is as in Fig. 7.

It follows from the definition of  $DH_{\mathcal{P}}$  that the difference between  $DH_{\mathcal{P}}(u_{\alpha})$  and  $DH_{\mathcal{P}}(u_{\beta})$  arises from the cones  $C^*(I)^+$ 's for  $I \in \Sigma^{(n)}$  such that  $i \in I$  and  $\langle u_I, v \rangle < \langle \mu, v \rangle$ . In fact, one sees that

$$\mathrm{DH}_{\mathcal{P}}(u_{\alpha}) + \sum_{I} \operatorname{sign} \langle u_{i}^{I}, v \rangle (-1)^{I} w(I) \phi_{I}(\mu) = \mathrm{DH}_{\mathcal{P}}(u_{\beta})$$

where *I* runs over the elements as above. Since  $\operatorname{sign}\langle u_i^I, v \rangle (-1)^I = -(-1)^{I \setminus \{i\}}$  and  $w(I) = w_i(I \setminus \{i\})$ , the equality above turns into

$$\mathrm{DH}_{\mathcal{P}}(u_{\alpha}) - \mathrm{DH}_{\mathcal{P}}(u_{\beta}) = \sum_{I} (-1)^{I \setminus \{i\}} w_{i}(I \setminus \{i\}) \phi_{I}(\mu).$$

Here  $\phi_I(\mu)$  may be viewed as the value at  $\mu$  of the characteristic function of the cones in  $F_i$  with apex  $u_I$  spanned by  $(u_j^I)^{+,s}$   $(j \in I, j \neq i)$ . This shows that the right-hand side at the equality above agrees with  $DH_{\mathcal{P}_i}(\mu - f_i)$ , proving the lemma.

**Lemma 5.4.** The support of the function  $DH_{\mathcal{P}}$  is bounded, and the function is independent of the choice of the generic element  $v \in V$ .

Proof. Induction on the dimension of simple multi-polytopes  $\mathcal{P}$ . We have observed the lemma in Example 5.1 when dim  $\mathcal{P} = 1$ . Suppose dim  $\mathcal{P} = n > 1$  and suppose that the lemma is true for simple multi-polytopes of dimension n - 1. Then the support of  $DH_{\mathcal{P}_i}$  is bounded by the induction assumption. This together with Lemma 5.3 implies that  $DH_{\mathcal{P}}$  takes the same constant on unbounded regions in  $V^* \setminus \bigcup F_i$ . On the other hand, it follows from the definition of  $DH_{\mathcal{P}}$  that  $DH_{\mathcal{P}}$  vanishes on a half space  $H_r := \{u \in V^* \mid \langle u, v \rangle < r\}$  for a sufficiently small real number r, because for each  $I \in \Sigma^{(n)}$  the cone  $C^*(I)^+$  is contained in the complement of  $H_r$  if r is sufficiently small. Therefore the constant which  $DH_{\mathcal{P}}$  takes on the unbounded regions in  $V^* \setminus \bigcup F_i$  is zero, proving the former assertion in the lemma.

As for the latter assertion in the lemma, it follows from the induction assumption that the right-hand side of the wall crossing formula in Lemma 5.3 is independent of v, and we have seen above that  $DH_{\mathcal{P}}$  vanishes on unbounded regions regardless of the choice of v. Thus, it follows from Lemma 5.3 that  $DH_{\mathcal{P}}$  is independent of v on any regions of  $V^* \setminus \bigcup F_i$ .

### 6. Winding numbers

We continue to assume that our multi-polytope  $\mathcal{P} = (\Delta, \mathcal{F})$  is simple and that  $\Sigma$  is an augmented simplicial set consisting of subsets of  $\{1, \ldots, d\}$ . In this section, we associate another locally constant function on  $V^* \setminus \bigcup F_i$  with  $\mathcal{P}$  from a topological viewpoint, and show that it agrees with the Duistermaat-Heckman function defined in Section 5.

Choose an orientation on V and fix it. We define an orientation on  $I = \{i_1, \ldots, i_n\} \in \Sigma^{(n)}$  as follows. If an ordered basis  $(v_{i_1}, \ldots, v_{i_n})$  gives the chosen orientation on V, then we say that the oriented simplex  $\langle i_1, \ldots, i_n \rangle$  has a positive orientation, and otherwise a negative orientation. We define

$$\langle I \rangle := \begin{cases} \langle i_1, \dots, i_n \rangle & \text{if } \langle i_1, \dots, i_n \rangle \text{ has a positive orientation,} \\ -\langle i_1, \dots, i_n \rangle & \text{if } \langle i_1, \dots, i_n \rangle \text{ has a negative orientation.} \end{cases}$$

The completeness of  $\Delta$  (equivalently, the pre-completeness of the projected multi-fan  $\Delta_J$  for any  $J \in \Sigma^{(n-1)}$ ) implies that

$$\sum_{I\in\Sigma^{(n)}}w(I)\langle I
angle$$

is a cycle in the chain complex of the simplicial set  $\Sigma$ . In fact, the converse also holds, i.e., the completeness of  $\Delta$  is equivalent to  $\sum_{I \in \Sigma^{(n)}} w(I) \langle I \rangle$  being a cycle. We denote by  $[\Delta]$  the homology class that the cycle defines in  $H_{n-1}(\Sigma)$ . Actually  $[\Delta]$  lies in the reduced homology  $\tilde{H}_{n-1}(\Sigma)$ , see Example 6.3 discussed later.

Let *S* be the realization of the first barycentric subdivision of  $\Sigma$ . For each  $i \in \{1, ..., d\}$ , we denote by  $S_i$  the union of simplicies in *S* which contain the vertex  $\{i\}$ , and by  $S_I$  the intersection  $\bigcap_{i \in I} S_i$  for  $I \in \Sigma$ . Note that  $\partial S_i$  can be identified with the realization of the first barycentric subdivision of  $\Sigma_i$ , where  $\Sigma_i$  is the augmented simplicial set of the projected multi-fan  $\Delta_i = (\Sigma_i, C_i, w_i^{\pm})$ .

The projected multi-fan  $\Delta_i$  is defined on  $V/V_i$  where  $V_i$  is the one-dimensional subspace spanned by  $v_i$ . We orient  $V/V_i$  as follows: if an ordered basis  $(v_i, v_{j_1}, \ldots, v_{j_{n-1}})$  defines the given orientation on V, then we give  $V/V_i$  the orientation determined by  $(v_{j_1}, \ldots, v_{j_{n-1}})$ , and otherwise give the opposite orientation. Then  $[\Delta_i]$  is defined in  $\tilde{H}_{n-2}(\Sigma_i) = \tilde{H}_{n-2}(\partial S_i)$ .

**Lemma 6.1.**  $[\Delta]$  maps to  $[\Delta_i]$  through the composition of maps

$$\tilde{H}_{n-1}(\Sigma) = \tilde{H}_{n-1}(S) \xrightarrow{\iota_*} H_{n-1}(S, S \setminus \operatorname{Int} S_i) \xleftarrow{excision}_{\cong} H_{n-1}(S_i, \partial S_i) \xrightarrow{\partial}_{\cong} \tilde{H}_{n-2}(\partial S_i),$$

where  $\iota$  is the inclusion.

Proof. Through  $\iota_*$  and the inverse of the excision isomorphism, the cycle  $\sum_{I \in \Sigma^{(n)}} w(I) \langle I \rangle$  maps to  $\sum_{i \in I \in \Sigma^{(n)}} w(I) \langle I \rangle$ . We express  $\langle I \rangle$  as  $\epsilon \langle i, j_1, \ldots, j_{n-1} \rangle$  where  $\epsilon = +1$  or -1 and define an oriented (n-2)-simplex  $\langle I \setminus \{i\} \rangle$  in  $\Sigma_i^{(n-1)}$  by  $\epsilon \langle j_1, \ldots, j_{n-1} \rangle$ . It follows that

$$\partial\left(\sum_{i\in I\in\Sigma^{(n)}}w(I)\langle I\rangle\right)=\sum_{i\in I\in\Sigma^{(n)}}w(I)\langle I\backslash\{i\}\rangle.$$

Here  $w(I) = w_i(I \setminus \{i\})$  by the definition of  $w_i$ , and  $i \in I \in \Sigma^{(n)}$  if and only of  $I \setminus \{i\} \in \Sigma_i^{(n-1)}$ . Therefore, the right-hand side above reduces to  $\sum_{J \in \Sigma_i^{(n-1)}} w_i(J) \langle J \rangle$ , that is  $[\Delta_i]$  in  $\tilde{H}_{n-2}(\partial S_i)$ .

The following lemma will be used later several times.

**Lemma 6.2.** Let X and Y be topological spaces with subspaces  $X_i \subset X$  and  $Y_i \subset Y$  for each  $i \in \Sigma^{(1)}$ . For  $I \in \Sigma$ , we set  $X_I := \bigcap_{i \in I} X_i$  and  $Y_I := \bigcap_{i \in I} Y_i$ . If

- (1)  $X = \bigcup_{i=1}^d X_i$ ,
- (2)  $X_I$ 's for  $I \in \Sigma^{(n)}$  are disjoint, and
- (3)  $Y_I$  is nonempty and contractible for any non-empty set  $I \in \Sigma$ ,

then there is a continuous map  $\psi: X \to Y$  sending the stratum  $X_I$  to  $Y_I$  for each  $I \in \Sigma$ , and such a map is unique up to homotopy preserving the stratifications.

Proof. *Existence*. We will construct  $\psi$  inductively using decending induction on |I|. If |I| = n, then we map  $X_I$  to any point in  $Y_I$ . Thus  $\psi$  is defined on  $\bigcup_{|I|=n} X_I$  with the image in  $\bigcup_{|I|=n} Y_I$ . Let k be a nonnegative integer less than n and |I| = k. Suppose that  $\psi$  is defined on  $\bigcup_{|J|>k+1} X_J$  with the image in  $\bigcup_{|J|>k+1} Y_J$ . Then

$$\psi \colon X_I \cap \left( \bigcup_{|J| \ge k+1} X_J \right) \to Y_I \cap \left( \bigcup_{|J| \ge k+1} Y_J \right) \subset Y_I$$

extends to a continuous map from  $X_I$  to  $Y_I$  because  $Y_I$  is contractible. Thus  $\psi$  is defined on  $\bigcup_{|I| \ge k} X_I$  with the image in  $\bigcup_{|I| \ge k} Y_I$ . This completes the induction step, so that we obtain the desired map  $\psi$  defined on X.

Uniqueness. We construct a homotopy  $H: X \times [0, 1] \to Y$  of given two maps  $\psi_0$ and  $\psi_1$  in the lemma. The argument is almost same as above. Since  $Y_I$  is contractible, H can be defined on  $\bigcup_{|I|=n} X_I \times [0, 1]$  with  $\bigcup_{|I|=n} Y_I$  as the image. Let k be as above and |I| = k. Suppose that H is defined on  $(\bigcup_{|J| \ge k+1} X_J) \times [0, 1]$  with the image in  $\bigcup_{|J| \ge k+1} Y_J$  and that H agrees with  $\psi_t$  on  $(\bigcup_{|J| \ge k+1} X_J) \times \{t\}$  for t = 0, 1. Then a map

$$H \cup \psi_0 \cup \psi_1 \colon (X_I \cap (\bigcup_{|J| \ge k+1} X_J)) \times [0, 1] \cup X_I \times \{0\} \cup X_I \times \{1\}$$
$$\rightarrow (Y_I \cap (\bigcup_{|J| \ge k+1} Y_J)) \cup Y_I \cup Y_I = Y_I$$

extends to a continuous map from  $X_I \times [0, 1]$  to  $Y_I$  because  $Y_I$  is contractible. Thus H is defined on  $(\bigcup_{|I| \ge k} X_I) \times [0, 1]$  with the image in  $\bigcup_{|I| \ge k} Y_I$ . This completes the induction step, so that we obtain the desired homotopy H defined on  $X \times [0, 1]$ .

Lemma 6.2 can be applied with X = S,  $X_i = S_i$ ,  $Y = V^*$  and  $Y_i = F_i$ . It follows that the multi-polytope  $\mathcal{P}$  associates a continuous map

$$\Psi\colon S\to \bigcup_{i=1}^d F_i\subset V^*$$

sending  $S_I$  to  $F_I$  for each  $I \in \Sigma$  by Lemma 6.2, and  $\Psi$  induces a homomorphism

$$\Psi_* \colon \tilde{H}_{n-1}(S) = \tilde{H}_{n-1}(\Sigma) \to \tilde{H}_{n-1}(V^* \setminus \{u\})$$

for each  $u \in V^* \setminus \bigcup F_i$ . Such  $\Psi$  was first introduced in [17] and plays the role of a moment map. The orientation on V chosen at the beginning of this section induces an orientation on  $V^*$  in a natural way. This determines a fundamental class

22

in  $H_n(V^*, V^* \setminus \{u\})$  and hence in  $\tilde{H}_{n-1}(V^* \setminus \{u\})$  through  $\partial : H_n(V^*, V^* \setminus \{u\}) \cong \tilde{H}_{n-1}(V^* \setminus \{u\})$ . We denote the fundamental class in  $\tilde{H}_{n-1}(V^* \setminus \{u\})$  by  $[V^* \setminus \{u\}]$ .

DEFINITION. For each  $u \in V^* \setminus \bigcup F_i$ , we define an integer  $WN_{\mathcal{P}}(u)$  by

$$\Psi_*([\Delta]) = WN_{\mathcal{P}}(u)[V^* \setminus \{u\}]$$

and call it the *winding number* of the multi-polytope  $\mathcal{P} = (\Delta, \mathcal{F})$  around u.

REMARK. The function  $WN_{\mathcal{P}}$  is independent of the choice of an orientation on V because if the orientation on V is reversed, then  $[\Delta]$  and  $[V^* \setminus \{u\}]$  are multiplied by -1 simultaneously. Moreover, it is locally constant and vanishes on unbounded regions separated by  $F_i$ 's, which immediately follows from the definition of  $WN_{\mathcal{P}}$ .

We will see in Theorem 6.6 below that  $WN_{\mathcal{P}} = DH_{\mathcal{P}}$ . For the moment, we shall check this coincidence when dim  $\mathcal{P} = 1$ .

EXAMPLE 6.3. We use the notation in Example 5.1. We identify V with  $\mathbb{R}$ , so that  $V^*$  is also identified with  $\mathbb{R}$ . Then V and  $V^*$  have standard orientations, and since  $v_i$  gives the orientation on V if and only if  $i \in E$ , the cycle which defines  $[\Delta]$  is given by

$$\sum_{i \in E} w(\{i\})\langle i \rangle + \sum_{i \notin E} w(\{i\})(-\langle i \rangle) = -\sum_{i=1}^d (-1)^{\{i\}} w(\{i\})\langle i \rangle$$

where  $(-1)^{\{i\}}$  is the same as in (5.2). Since  $\Delta$  is complete,  $\sum_{i=1}^{d} (-1)^{\{i\}} w(\{i\}) = 0$ ; so  $[\Delta]$  actually lies in  $\tilde{H}_0(\Sigma) = \tilde{H}_0(S)$  and one can rewrite the cycle above as

$$\sum_{i=1}^d (-1)^{\{i\}} w(\{i\}) (\langle j \rangle - \langle i \rangle)$$

for any  $j \in \{1, ..., d\}$ . Since  $S_i = \{i\}$  and  $\Psi(\{i\}) = F_i$ ,  $WN_{\mathcal{P}}(u) = 0$  unless u is between the minimum value and the maximum value of  $\{F_1, ..., F_d\}$ . Suppose u is between them and take j such that  $F_j$  is the maximum. Then one easily sees that

WN<sub>P</sub>(u) = 
$$\sum_{F_i < u} (-1)^{\{i\}} w(\{i\}).$$

This together with (5.3) shows that  $WN_{\mathcal{P}} = DH_{\mathcal{P}}$  when dim  $\mathcal{P} = 1$ .

We will show that WN satisfies the same wall crossing formula as in Lemma 5.3. For that, we first state a lemma which expresses the winding number as a sum of local winding numbers so to speak. Assume dim  $\mathcal{P} > 1$ . We orient  $F_i$  in such a way

that the juxtaposition of a normal vector to  $F_i$ , whose evaluation on  $v_i$  is positive, and the orientation on  $F_i$  agrees with the prescribed orientation on  $V^*$ . By Lemma 6.2,  $\Psi$ maps a pair  $(S_i, \partial S_i)$  into a pair  $(F_i, F_i \setminus \{\mu\})$  for any  $\mu \in F_i \setminus (F_i \cap (\bigcup_{j \in \Sigma_i^{(1)}} F_j))$ . If we identify  $F_i$  with  $(V^*)_i$  through the translation by  $-f_i$  as before, then the map  $\Psi$ restricted to  $\partial S_i$  agrees with the map (up to homotopy) constructed from the multipolytope  $\mathcal{P}_i = (\Delta_i, \mathcal{F}_i)$ . It follows that

(6.1) 
$$\Psi_*([\Delta_i]) = WN_{\mathcal{P}_i}(\mu - f_i)[F_i \setminus \{\mu\}].$$

Let  $u \in V^* \setminus \bigcup F_i$ . We choose a generic ray R starting from u with direction  $\gamma \in V^*$ , so that the intersection  $F_i \cap R$  is one point for each i if it is nonempty. We denote the point  $F_i \cap R$  by  $R_i$ .

**Lemma 6.4.** WN<sub>$$\mathcal{P}$$</sub>( $u$ ) =  $\sum_{i:F_i \cap R \neq \phi} \operatorname{sign} \langle \gamma, v_i \rangle WN_{\mathcal{P}_i}(R_i - f_i).$ 

Proof. Consider the following commutative diagram:

$$\begin{split} \tilde{H}_{n-1}(S) & \to H_{n-1}(S, S \setminus \bigcup_{i} \operatorname{Int} S_{i}) \xleftarrow{\operatorname{excision}}_{\cong} \bigoplus_{i} H_{n-1}(S_{i}, \partial S_{i}) \xrightarrow{\partial}_{\cong} \bigoplus_{i} \tilde{H}_{n-2}(\partial S_{i}) \\ \psi_{*} \downarrow & \psi_{*} \downarrow & \psi_{*} \downarrow \\ \tilde{H}_{n-1}(V^{*} \setminus \{u\}) \xrightarrow{\cong} H_{n-1}(V^{*} \setminus \{u\}, V^{*} \setminus R) \leftarrow \bigoplus_{i} H_{n-1}(F_{i}, F_{i} \setminus \{R_{i}\}) \xrightarrow{\partial}_{\cong} \bigoplus_{i} \tilde{H}_{n-2}(F_{i} \setminus \{R_{i}\}) \end{split}$$

where *i* runs over the indices of  $F_i$ 's which intersect *R*. The element  $[\Delta] \in \tilde{H}_{n-1}(S)$  maps to  $\bigoplus_i [\Delta_i] \in \bigoplus_i \tilde{H}_{n-2}(\partial S_i)$  through the upper horizontal sequence by Lemma 6.1 and down to  $\bigoplus_i WN_{\mathcal{P}_i}(R_i - f_i)[F_i \setminus \{R_i\}]$  by (6.1).

Now we trace the lower horizontal sequence from the right to the left. Through the inverse of  $\partial$ ,  $[F_i \setminus \{R_i\}]$  maps to the fundamental class  $[F_i, F_i \setminus \{R_i\}]$ , and further maps to sign $\langle \gamma, v_i \rangle [V^* \setminus \{u\}] \in \tilde{H}_{n-1}(V^* \setminus \{u\})$ , where the sign arises from the choice of the orientation on  $F_i$ . These together with the commutativity of the diagram above show that

$$\Psi_*([\Delta]) = \sum_{i:F_i \cap R \neq \phi} \operatorname{sign}\langle \gamma, v_i \rangle \operatorname{WN}_{\mathcal{P}_i}(R_i - f_i)[V^* \setminus \{u\}].$$

On the other hand,  $\Psi_*([\Delta]) = WN_{\mathcal{P}}(u)[V^* \setminus \{u\}]$  by definition. The lemma follows by comparing these two equalities.

**Lemma 6.5.** The wall crossing formula as in Lemma 5.3 holds for WN instead of DH.

Proof. Subtract the equality in Lemma 6.4 for  $u = u_{\beta}$  from that for  $u = u_{\alpha}$ . Since one can take  $\gamma$  to be  $u_{\beta} - u_{\alpha}$ , the lemma follows. **Theorem 6.6.**  $DH_{\mathcal{P}} = WN_{\mathcal{P}}$  for any simple multi-polytope  $\mathcal{P}$ .

Proof. The equality is established in Example 6.3 when dim  $\mathcal{P} = 1$ . Suppose dim  $\mathcal{P} = n > 1$  and suppose that the equality holds for simple multi-polytopes of dimension n-1. Both DH<sub> $\mathcal{P}$ </sub> and WN<sub> $\mathcal{P}$ </sub> are locally constant, satisfy the same wall crossing formula (Lemma 5.3, Lemma 6.5) and DH<sub> $\mathcal{P}_i$ </sub> = WN<sub> $\mathcal{P}_i$ </sub> by induction assumption. Therefore, it suffices to see that DH<sub> $\mathcal{P}$ </sub> and WN<sub> $\mathcal{P}$ </sub> agree on one region. But we know that they vanish on unbounded regions (Lemma 5.4 and the remark after the definition of WN<sub> $\mathcal{P}$ </sub>), hence they agree on the whole domain. This completes the induction step, proving the theorem.

#### 7. Ehrhart polynomials

Let *P* be a convex lattice polytope of dimension *n* in *V*<sup>\*</sup>, where "lattice polytope" means that each vertex of *P* lies in the lattice  $N^* = \text{Hom}(N, \mathbb{Z})$  of  $V^* = \text{Hom}(V, \mathbb{R})$ . For a positive integer  $\nu$ , let  $\nu P := \{\nu u \mid u \in P\}$ . It is again a convex lattice polytope in *V*<sup>\*</sup>. We denote by  $\sharp(\nu P)$  (resp.  $\sharp(\nu P^\circ)$ ) the number of lattice points in  $\nu P$  (resp. in the interior of  $\nu P$ ). The lattice  $N^*$  determines a volume element on  $V^*$  by requiring that the volume of the unit cube determined by a basis of  $N^*$  is 1. Thus the volume of *P*, denoted by vol(*P*), is defined. The following theorem is well known.

**Theorem 7.1** (see [11], [29] for example.). Let P be an n-dimensional convex lattice polytope.

(1)  $\sharp(\nu P)$  and  $\sharp(\nu P^{\circ})$  are polynomials in  $\nu$  of degree n.

(2)  $\sharp(\nu P^{\circ}) = (-1)^n \sharp(-\nu P)$ , where  $\sharp(-\nu P)$  denotes the polynomial  $\sharp(\nu P)$  with  $\nu$  replaced by  $-\nu$ .

(3) The coefficient of  $\nu^n$  in  $\sharp(\nu P)$  is vol(P) and the constant term in  $\sharp(\nu P)$  is 1.

The fan  $\Delta$  associated with P may not be simplicial, but if we subdivide  $\Delta$ , then we can always take a simplicial fan that is compatible with P. In this section, we show that the theorem above holds for a *simple* lattice multi-polytope  $\mathcal{P} = (\Delta, \mathcal{F})$ . For that, we need to define  $\sharp(\mathcal{P})$  and  $\sharp(\mathcal{P}^\circ)$ . This is done as follows. Let  $v_i$  (i = 1, ..., d)be a primitive integral vector in the half line  $C(\{i\})$ . In our convention,  $v_i$  is chosen "outward "normal" to the face  $\mathcal{F}(\{i\})$  when  $\mathcal{P}$  arises from a convex polytope. We slightly move  $\mathcal{F}(\{i\})$  in the direction of  $v_i$  (resp.  $-v_i$ ) for each i, so that we obtain a map  $\mathcal{F}_+$  (resp.  $\mathcal{F}_-$ ):  $\Sigma^{(1)} \to \operatorname{HP}(V^*)$ . We denote the multi-polytopes  $(\Delta, \mathcal{F}_+)$  and  $(\Delta, \mathcal{F}_-)$  by  $\mathcal{P}_+$  and  $\mathcal{P}_-$  respectively. Since the affine hyperplanes  $\mathcal{F}_{\pm}(\{i\})$ 's miss the lattice  $N^*$ , the functions  $\operatorname{DH}_{\mathcal{P}_{\pm}}$  and  $\operatorname{WN}_{\mathcal{P}_{\pm}}$  are defined on  $N^*$ . DEFINITION. We define

$$\sharp(\mathcal{P}) := \sum_{u \in N^*} \mathrm{DH}_{\mathcal{P}_+}(u) = \sum_{u \in N^*} \mathrm{WN}_{\mathcal{P}_+}(u),$$
$$\sharp(\mathcal{P}^\circ) := \sum_{u \in N^*} \mathrm{DH}_{\mathcal{P}_-}(u) = \sum_{u \in N^*} \mathrm{WN}_{\mathcal{P}_-}(u).$$

When  $\mathcal{P}$  arises from a convex polytope P,  $DH_{\mathcal{P}_+} = WN_{\mathcal{P}_+}$  (resp.  $DH_{\mathcal{P}_-} = WN_{\mathcal{P}_-}$ ) takes 1 on u in P (resp. in the interior of P) and 0 otherwise. Therefore,  $\sharp(\mathcal{P})$  (resp.  $\sharp(\mathcal{P}^\circ)$ ) agrees with the number of lattice points in P (resp. in the interior of P) in this case.

Denote the volume element on  $V^*$  by  $dV^*$ , and define the volume  $vol(\mathcal{P})$  of  $\mathcal{P}$  by

$$\operatorname{vol}(\mathcal{P}) := \int_{V^*} \mathrm{DH}_{\mathcal{P}} \, dV^* = \int_{V^*} \mathrm{WN}_{\mathcal{P}} \, dV^*.$$

When  $\mathcal{P}$  arises from a (convex) polytope P,  $vol(\mathcal{P})$  agrees with the actual volume of P, but otherwise it can be zero or negative.

For a (not necessarily positive) integer  $\nu$ , we denote  $(\Delta, \nu \mathcal{F})$  by  $\nu \mathcal{P}$ , where

$$(\nu \mathcal{F})(\{i\}) \coloneqq \{u \in V^* \mid \langle u, v_i \rangle = \nu c_i\}$$

when  $\mathcal{F}(\{i\}) = \{u \in V^* \mid \langle u, v_i \rangle = c_i\}$  for a constant  $c_i$ .

**Theorem 7.2.** Let  $\mathcal{P} = (\Delta, \mathcal{F})$  be a simple lattice multi-polytope of dimension *n*. (1)  $\sharp(\nu \mathcal{P})$  and  $\sharp(\nu \mathcal{P}^{\circ})$  are polynomials in  $\nu$  of degree (at most) *n*.

(2)  $\sharp(\nu \mathcal{P}^{\circ}) = (-1)^n \sharp(-\nu \mathcal{P})$  for any integer  $\nu$ .

(3) The coefficient of  $\nu^n$  in  $\sharp(\nu \mathcal{P})$  is  $\operatorname{vol}(\mathcal{P})$  and the constant term in  $\sharp(\nu \mathcal{P})$  is  $\operatorname{deg}(\Delta)$ . (see Section 2 for  $\operatorname{deg}(\Delta)$ .)

In order to prove this theorem, we need some notations and a lemma. Basic ideas in the following arguments are in [4] and [5]. Let  $I \in \Sigma^{(n)}$ . Although the integral vectors  $\{v_i \mid i \in I\}$  are not necessarily a basis of the lattice N, they are linearly independent. Therefore, the sublattice  $N_I$  of N generated by  $v_i$ 's  $(i \in I)$  is of the same rank as N, hence  $N/N_I$  is a finite group. Needless to say,  $N/N_I$  is trivial for any  $I \in \Sigma^{(n)}$  if  $\Delta$  is non-singular. For  $u \in N_I^* = \text{Hom}(N_I, \mathbb{Z}) \supset N^*$  and  $g \in N/N_I$ , we define

(7.1) 
$$\chi_I(u,g) := \exp(2\pi\sqrt{-1}\langle u, v_g \rangle)$$

where  $v_g \in N$  is a representative of g. The right-hand side does not depend on the choice of the representative  $v_g$ , and  $\chi_I(u, \cdot)$  (resp.  $\chi(\cdot, g)$ ) is a homomorphism from

 $N/N_I$  (resp.  $N_I^*$ ) to  $\mathbb{C}^*$ . Note that  $\chi_I(u, : N/N_I \to \mathbb{C}^*$  is trivial if and only if  $u \in N^*$ . It follows that

(7.2) 
$$\sum_{g \in N/N_I} \chi_I(u,g) = \begin{cases} |N/N_I| & \text{if } u \in N^*, \\ 0 & \text{otherwise.} \end{cases}$$

**Lemma 7.3.** For each  $I \in \Sigma^{(n)}$  let  $u_I$  be the corresponding vertex of  $\mathcal{P}$  and let  $\{u_i^I \mid i \in I\}$  be the dual basis of  $\{v_i \mid i \in I\}$  as in Section 5. Then, for  $v \in N$  such that  $\langle u_i^I, v \rangle$  is a nonzero integer for any  $I \in \Sigma^{(n)}$  and  $i \in I$ , we have

$$\sum_{I\in\Sigma^{(n)}}\frac{w(I)z^{\langle u_I,v\rangle}}{|N/N_I|}\sum_{g\in N/N_I}\frac{1}{\prod_{i\in I}(1-\chi_I(u_i^I,g)z^{-\langle u_i^I,v\rangle})}=\sum_{u\in N^*}\mathrm{DH}_{\mathcal{P}_+}(u)z^{\langle u,v\rangle}$$

as functions of  $z \in \mathbb{C}$ .

Proof. The Maclaurin expansion of  $1/(1 - az^{-m})$   $(a \in \mathbb{C}^*, m \in \mathbb{Z})$  is given by

$$\begin{cases} -a^{-1}z^m - a^{-2}z^{2m} - \cdots & \text{if } m > 0\\ 1 + az^{-m} + a^2 z^{-2m} + \cdots & \text{if } m < 0. \end{cases}$$

Taking this into account, we expand the sum

$$S_I := \sum_{g \in N/N_I} \frac{1}{\prod_{i \in I} (1 - \chi_I(u_i^I, g) z^{-\langle u_i^I, v \rangle})}$$

into Maclaurin series and get

$$S_{I} = \sum_{g \in N/N_{I}} (-1)^{I} \prod_{i \in I} \sum_{\{b_{i}\}} (\chi_{I}(u_{i}^{I}, g)^{-b_{i}} z^{b_{i} \langle u_{i}^{I}, v \rangle})$$
  
= 
$$\sum_{g \in N/N_{I}} (-1)^{I} \sum_{\{b_{i}\}} \chi_{I}(-\sum_{i \in I} b_{i} u_{i}^{I}, g) z^{\langle \sum_{i \in I} b_{i} u_{i}^{I}, v \rangle},$$

where the summation  $\sum_{\{b_i\}}$  runs over the collection of such  $\{b_i \mid i \in I, \ b_i \in \mathbb{Z}\}$  that

(7.3) 
$$b_i \ge 1$$
 for  $i$  with  $\langle u_i^I, v \rangle > 0$  and  $b_i \le 0$  for  $i$  with  $\langle u_i^I, v \rangle < 0$ ,

(see Section 5 for  $(-1)^I$ ). Since

$$\sum_{g \in N/N_I} \chi_I \left( -\sum_{i \in I} b_i u_i^I, g \right) = \begin{cases} |N/N_I| & \text{if } \sum_{i \in I} b_i u_i^I \in N^*, \\ 0 & \text{otherwise,} \end{cases}$$

by (7.2), the Maclaurin expansion of the left-hand side of the equality in Lemma 7.3

has the form

$$\sum_{u \in N^*} \left( \sum_{I \in \Sigma^{(n)}} (-1)^I w(I) \phi_I'(u) \right) z^{\langle u, v \rangle}$$

where

$$\phi_I'(u) = \begin{cases} 1 & \text{if } u = u_I + \sum_{i \in I} b_i u_i^I, \ b_i \text{'s are as in (7.3) and } \sum_{i \in I} b_i u_i^I \in N^*, \\ 0 & \text{otherwise.} \end{cases}$$

One easily checks that  $\sum_{I \in \Sigma^{(n)}} (-1)^I w(I) \phi'_I(u)$  agrees with  $DH_{\mathcal{P}_+}(u)$ , proving the lemma.

Proof of Theorem 7.2. We shall prove (2) first. It suffices to prove  $\sharp(\mathcal{P}^\circ) = (-1)^n \sharp(-\mathcal{P})$ . Since  $\sharp(\mathcal{P}^\circ) = \sum_{u \in N^*} WN_{\mathcal{P}_-}(u)$  by definition, it suffices to prove that

(7.4) 
$$WN_{\mathcal{P}_{-}}(u) = (-1)^{n}WN_{(-\mathcal{P})_{+}}(u) \text{ for any } u \in N^{*}.$$

Let  $\Psi_{\mathcal{P}_{-}}$  and  $\Psi_{(-\mathcal{P})_{+}}$  be the maps introduced in Section 6 which are associated with multi-polytopes  $\mathcal{P}_{-}$  and  $(-\mathcal{P})_{+}$  respectively. We note that  $\Psi_{\mathcal{P}_{-}}$  and  $-\Psi_{(-\mathcal{P})_{+}}$  considered as maps from S to  $V^* \setminus \{u\}$  for  $u \in N^*$  are homotopic. Since the multiplication by -1 on  $V^*$  sends the fundamental class  $[V^* \setminus \{-u\}]$  to  $(-1)^n [V^* \setminus \{u\}]$ , we obtain (7.4).

We shall prove (1). Because of (2), it suffices to prove (1) for  $\sharp(\nu \mathcal{P})$ . We apply Lemma 7.3 to  $\nu \mathcal{P}$  in place of  $\mathcal{P}$  (so that  $u_I$  is replaced by  $\nu u_I$ ), and approach z to 1 in the equality. Since the right-hand side approaches  $\sharp(\nu \mathcal{P})$ , it suffices to show that the left-hand side approaches a polynomial in  $\nu$  of degree at most n. When  $g \in N/N_I$ is the identity element,  $\chi_I(u_i^I, g) = 1$ . Therefore, the term in the summand  $\sum_{g \in N/N_I}$ in the left-hand side has a pole at z = 1 of degree exactly n when g is the identity element, and of degree at most n otherwise. Thus the left-hand side of the equality in Lemma 7.3 applied to  $\nu \mathcal{P}$  can be written as

$$\frac{\sum_{I\in\Sigma^{(n)}} z^{\nu\langle u_I,v\rangle} h_I(z)}{(1-z)^n f(z)}$$

where  $h_I(z)$  and f(z) are polynomials in z and  $f(1) \neq 0$ . Then the repeated use of L'Hospital's Theorem implies that when z approaches 1, the limit of the above rational function is a polynomial in  $\nu$  of degree at most n.

Finally we prove (3). Since

$$\sharp(\nu\mathcal{P}) = \sum_{u \in H^2(BT)} \mathrm{DH}_{(\nu\mathcal{P})_+}(u) = \sum_{u \in H^2(BT)/\nu} \mathrm{DH}_{\mathcal{P}_+}(u),$$

28

it follows from the definition of definite integral that

$$\lim_{\nu\to\infty}\frac{1}{\nu^n}\sharp(\nu\mathcal{P})=\lim_{n\to\infty}\frac{1}{\nu^n}\sum_{u\in H^2(BT)/\nu}\mathrm{DH}_{\mathcal{P}_+}(u)=\int_{V^*}\mathrm{DH}_{\mathcal{P}}\ dV^*=\mathrm{vol}(\mathcal{P}),$$

proving that the coefficient of  $\nu^n$  in  $\sharp(\nu \mathcal{P})$  is vol $(\mathcal{P})$ .

We apply Lemma 7.3 to  $0\mathcal{P}$ , that is  $\nu\mathcal{P}$  with  $\nu = 0$ . Then the  $u_1$  in the lemma is zero, and  $DH_{(0\mathcal{P})_+}(u) = WN_{(0\mathcal{P})_+}(u) = 0$  unless u = 0 because the origin is the only vertex of  $0\mathcal{P}$  so that the vertices of  $(0\mathcal{P})_+$  are very close to the origin. Thus the righthand side of the equality in the lemma applied to  $0\mathcal{P}$  is a constant, say c, which is nothing but the constant term in  $\sharp(\nu\mathcal{P})$ . Now we approach z to  $\infty$ . Then the equality reduces to

$$\sum_{v \in C(I)} w(I) = c$$

because  $\langle u_i^I, v \rangle > 0$  for all  $i \in I$  if and only if  $v = \sum_{i \in I} a_i v_i$  with  $a_i > 0$  for all  $i \in I$ , and the latter is equivalent to saying that v belongs to the cone C(I) spanned by  $v_i$ 's  $(i \in I)$ . Since  $\sum_{v \in C(I)} w(I) = \deg(\Delta)$  by definition, the constant term in  $\sharp(\nu \mathcal{P})$ , that is c, agrees with  $\deg(\Delta)$ .

Let  $N_{\Delta}^*$  be the lattice of  $N_{\mathbb{R}}^*$  generated by all  $u_i^I$ 's for  $I \in \Sigma^{(n)}$  and  $i \in I$ . If  $\Delta$  is non-singular, then  $N_{\Delta}^* = N^*$ . The group ring  $\mathbb{C}[N_{\Delta}^*]$  is an integral domain, and it has a basis  $t^u$  ( $u \in N_{\Delta}^*$ ) as a complex vector space with multiplication determined by the addition in  $N_{\Delta}^*$ :

$$t^{u} \cdot t^{u'} := t^{u+u'}.$$

The quotient field of  $\mathbb{C}[N_{\Delta}^*]$  will be denoted by  $\mathbb{C}(N_{\Delta}^*)$ . It contains  $\mathbb{C}[N_{\Delta}^*]$ . Each  $v \in N$  such that  $\langle u_i^I, v \rangle$  is an integer for any  $I \in \Sigma^{(n)}$  and  $i \in I$  determines a map from  $\mathbb{C}[N_{\Delta}^*]$  to a Laurent polynomial ring  $\mathbb{C}[z, z^{-1}]$  sending  $t^u$  to  $z^{\langle u, v \rangle}$ . This map extends to a map from  $\mathbb{C}(N_{\Delta}^*)$  to  $\mathbb{C}(z)$ , the field of rational functions in z. Since Lemma 7.3 holds for any such v that  $\langle u_i^I, v \rangle \neq 0$ , we obtain

Corollary 7.4. Let the notation be the same as in Lemma 7.3. Then

$$\sum_{I\in\Sigma^{(n)}}\frac{w(I)t^{u_I}}{|N/N_I|}\sum_{g\in N/N_I}\frac{1}{\prod_{i\in I}(1-\chi_I(u_i^I,g)t^{-u_i^I})}=\sum_{u\in N^*}\mathrm{DH}_{\mathcal{P}_+}(u)t^u\in\mathbb{C}[N^*]$$

as elements in  $\mathbb{C}(N^*_{\Delta})$ . In particular, if the multi-fan  $\Delta$  is non-singular, then  $N^*_{\Delta} = N^*$ and

$$\sum_{I\in\Sigma^{(n)}}\frac{w(I)t^{u_I}}{\prod_{i\in I}(1-t^{-u_i^I})}=\sum_{u\in N^*}\mathrm{DH}_{\mathcal{P}_+}(u)t^u.$$

For a later use, we shall rewrite  $\chi_I(u_i^I, g)$ . Consider a homomorphism  $\eta \colon \mathbb{R}^d \to N_{\mathbb{R}}$  mapping  $\mathbf{a} = (a_1, \ldots, a_d) \in \mathbb{R}^d$  to  $\sum_{i=1}^d a_i v_i \in N_{\mathbb{R}}$ . For  $I \in \Sigma^{(n)}$ , we define

$$G'_I := \{ \mathbf{a} \in \mathbb{R}^d \mid \eta(\mathbf{a}) \in N \text{ and } a_j = 0 \text{ for } j \notin I \}$$

and define  $G_I$  to be the projection image of  $G'_I$  on  $\mathbb{R}^d/\mathbb{Z}^d$ . Since  $v_i$ 's  $(i \in I)$  are linearly independent and belong to N,  $G_I$  is a finite subgroup of  $\mathbb{R}^d/\mathbb{Z}^d$  and  $\eta$  restricted to  $G'_I$  induces an isomorphism

$$\eta_I \colon G_I \cong N/N_I$$
.

Note that  $\eta_I([\mathbf{a}]) = [\sum_{i \in I} a_i v_i]$  where [ ] denotes the equivalence class.

On the other hand, for  $i = 1, \ldots, d$ , let

$$\rho_i \colon \mathbb{R}^d / \mathbb{Z}^d \to \mathbb{C}^*$$

be a homomorphism defined by  $\rho_i([\mathbf{a}]) = \exp(2\pi\sqrt{-1}a_i)$ .

**Lemma 7.5.** For  $[\mathbf{a}] \in G_I \subset \mathbb{R}^d / \mathbb{Z}^d$  and  $i \in I$ , we have  $\rho_i([\mathbf{a}]) = \chi_I(u_i^I, \eta_I([\mathbf{a}]))$ .

Proof. Since  $\eta_I([\mathbf{a}]) = [\sum_{i \in I} a_i v_i]$  and  $\langle u_i^I, \sum_{i \in I} a_i v_i \rangle = a_i$ , it follows from the definition (7.1) of  $\chi_I$  that  $\chi_I(u_i^I, \eta_I([\mathbf{a}])) = \exp(2\pi\sqrt{-1}a_i)$ , which is equal to  $\rho_i([\mathbf{a}])$  by definition.

Since  $G_I$  is isomorphic to  $N/N_I$ , Corollary 7.4 can be restated as follows.

Corollary 7.6. Let the notation be as above. Then

$$\sum_{I\in\Sigma^{(n)}}\frac{w(I)t^{u_I}}{|G_I|}\sum_{g\in G_I}\frac{1}{\prod_{i\in I}(1-\rho_i(g)t^{-u_i^I})}=\sum_{u\in N^*}\mathrm{DH}_{\mathcal{P}_+}(u)t^u\in\mathbb{C}[N^*]$$

as elements in  $\mathbb{C}(N^*_{\Delta})$ .

# 8. Cohomological formula for $\#(\mathcal{P})$

Motivated by the geometrical observation which will be explained in subsequent sections 9 and 11, we define the "(equivariant) cohomology" of a complete simplicial multi-fan and the "(equivariant) first Chern class" of a multi-polytope. We then define an index map "in cohomology" and establish a "cohomological formula" describing  $\sharp(\mathcal{P})$  for a lattice multi-polytope. This cohomological formula is a counterpart in combinatorics to the Hirzebruch-Riemann-Roch formula applied to a complex *T*-line bundle over a torus manifold. As an application of the cohomological formula, we show that the Khovanskii-Pukhlikov formula for a simple lattice convex polytope ([21], [25], [6], [7], [13], [4], [5]) can be generalized to a simple lattice multi-polytope.

30

Let *T* be a compact torus of dimension  $n = \operatorname{rank}_{\mathbb{Z}} N$  and let *BT* be the classifying space of *T*. Then  $H_2(BT)$  is canonically isomorphic to  $\operatorname{Hom}(S^1, T)$ , the group consisting of homomorphisms from  $S^1$  to *T*. In fact, a homomorphism  $f: S^1 \to T$  induces a continuous map  $Bf: BS^1 \to BT$  and once we fix a generator  $\alpha$  of  $H_2(BS^1) \cong \mathbb{Z}$ ,  $(Bf)_*\alpha$  defines an element of  $H_2(BT)$ . The correspondence  $f \mapsto (Bf)_*\alpha$  is known to be an isomorphism from  $\operatorname{Hom}(S^1, T)$  to  $H_2(BT)$ . In the following we assume  $N = H_2(BT)$  and identify it with  $\operatorname{Hom}(S^1, T)$ . Then  $N^* = H^2(BT)$  is identified with  $\operatorname{Hom}(T, S^1)$  and the group ring  $\mathbb{C}[N^*]$  can be identified with the representation ring R(T) of *T*.

Let  $\Delta = (\Sigma, C, w^{\pm})$  be a complete simplicial multi-fan in *N*. Let  $v_i \in H_2(BT)$  be a unique primitive vector in  $C(\{i\})$  for each  $i = 1, \ldots, d$  as before. Motivated by the description of the equivariant cohomology of a compact non-singular toric variety (see Proposition 9.2 in the next section), we define  $H_T^*(\Delta)$  to be the face ring of the augmented simplicial set  $\Sigma$ , i.e.,

$$H_T^*(\Delta) := \mathbb{Z}[x_1, \ldots, x_d]/(x_I \mid I \notin \Sigma),$$

where  $x_I = \prod_{i \in I} x_i$  and the degree of  $x_i$  is two, and call  $H_T^*(\Delta)$  the *equivariant co-homology* of  $\Delta$ . We also define a homomorphism  $\pi^* \colon H^2(BT) \to H_T^2(\Delta)$  by

(8.1) 
$$\pi^*(u) = \sum_{i=1}^d \langle u, v_i \rangle x_i,$$

where  $\langle , \rangle$  denotes the natural pairing between cohomology and homology. It extends to an algebra homomorphism  $H^*(BT) \to H^*_T(\Delta)$ , which we also denote by  $\pi^*$ . One can think of  $H^*_T(\Delta)$  as a module (or more generally an algebra) over  $H^*(BT)$  through  $\pi^*$ .

In the following we will mainly work with  $\mathbb{Q}$  coefficients but the argument will work with  $\mathbb{Z}$  coefficients when the multi-fan  $\Delta$  is non-singular. Any homomorphism  $f: A \to B$  between additive groups induces a homomorphism :  $A \otimes \mathbb{Q} \to B \otimes \mathbb{Q}$  (or  $A \otimes \mathbb{R} \to B \otimes \mathbb{R}$ ), which we also denote by f.

**Lemma 8.1.** Any element in  $H_T^*(\Delta) \otimes \mathbb{Q}$  can be written in the form  $\sum_{J \in \Sigma} \pi^*(a_J) x_J$ with  $a_J \in H^*(BT; \mathbb{Q})$  (not necessarily uniquely), in other words,  $H_T^*(\Delta) \otimes \mathbb{Q}$  is generated by  $x_J$ 's  $(J \in \Sigma)$  as an  $H^*(BT; \mathbb{Q})$ -module.

Proof. Let  $\mathcal{I}$  denote a finite set which consists of elements in  $\{1, \ldots, d\}$  taken with multiplicity, i.e., elements in  $\{1, \ldots, d\}$  may appear in  $\mathcal{I}$  repeatedly. Set  $x_{\mathcal{I}} :=$  $\prod_{i \in \mathcal{I}} x_i$  and denote by  $\overline{\mathcal{I}}$  the subset of  $\{1, \ldots, d\}$  consisting of elements appearing in  $\mathcal{I}$ . It follows from the definition that  $H_T^*(\Delta)$  is additively generated by  $x_{\mathcal{I}}$ 's such that  $\overline{\mathcal{I}} \in \Sigma$ , so it suffices to prove the lemma for such  $x_{\mathcal{I}}$ . We shall prove it by induction on  $[\mathcal{I}] := |\mathcal{I}| - |\overline{\mathcal{I}}|$ . If  $[\mathcal{I}] = 0$ , then  $\mathcal{I} = \overline{\mathcal{I}} \in \Sigma$ ; so  $x_{\mathcal{I}}$  is obviously of the form in the lemma in this case. Suppose  $[\mathcal{I}] \ge 1$ . Then there is an  $i \in \mathcal{I}$  which appears in  $\mathcal{I}$  at least twice. Set  $\mathcal{J} := \mathcal{I} \setminus \{i\}$ . Then  $\overline{\mathcal{J}} = \overline{\mathcal{I}} \in \Sigma$  and  $[\mathcal{J}] = [\mathcal{I}] - 1$ . Multiplying the both sides at (8.1) by  $x_{\mathcal{J}}$ , we obtain

$$\pi^*(u)x_{\mathcal{J}} = \langle u, v_i \rangle x_{\mathcal{I}} + \sum_{k \neq i} \langle u, v_k \rangle x_{\mathcal{J} \cup \{k\}}$$

for any  $u \in H^2(BT; \mathbb{Q})$ . We choose u such that  $\langle u, v_i \rangle = 1$  and  $\langle u, v_j \rangle = 0$  for all  $j \in \mathcal{J}$  different from i. (Such u exists because  $\{v_j \mid j \in \overline{\mathcal{J}}\}$  is a subset of a basis of  $N_{\mathbb{Q}}$ .) Then the equality above reduces to

$$x_{\mathcal{I}} = \pi^*(u) x_{\mathcal{J}} - \sum_{k \neq i, k \notin \mathcal{J}} \langle u, v_k \rangle x_{\mathcal{J} \cup \{k\}}.$$

Here  $[\mathcal{J} \cup \{k\}] = [\mathcal{J}](= [\mathcal{I}] - 1)$  for  $k \notin \mathcal{J}$ , so the right-hand side above are of the form in the lemma by the induction assumption, showing that so is  $x_{\mathcal{I}}$ . This completes the induction step and proves the lemma.

For  $I \in \Sigma^{(n)}$ , let  $\{u_i^I \mid i \in I\}$  be the dual basis of  $\{v_i \mid i \in I\}$  as before. We define a ring homomorphism  $\iota_I^* \colon H_T^*(\Delta) \otimes \mathbb{Q} \to H^*(BT; \mathbb{Q})$  by

$$\iota_I^*(x_i) = \begin{cases} u_i^I & \text{if } i \in I, \\ 0 & \text{otherwise.} \end{cases}$$

This map is well-defined because  $x_J$  for  $J \notin \Sigma$ , which is zero in  $H^*_T(\Delta) \otimes \mathbb{Q}$ , maps to zero through  $\iota^*_I$ .

**Lemma 8.2.** The composition  $\iota_I^* \circ \pi^*$  is the identity map. In particular  $\iota_I^*$  is an  $H^*(BT; \mathbb{Q})$ -module map.

Proof. Both  $\pi^*$  and  $\iota_I^*$  are ring homomorphisms and  $H^*(BT)$  is a polynomial ring generated by elements in  $H^2(BT)$ , so it suffices to check the lemma on  $H^2(BT)$ . Let  $u \in H^2(BT)$ . It follows from the definitions of  $\pi^*$  and  $\iota_I^*$  that

$$(\iota_I^* \circ \pi^*)(u) = \iota_I^* \left( \sum_{i=1}^d \langle u, v_i \rangle x_i \right) = \sum_{i=1}^d \langle u, v_i \rangle u_i^I,$$

which agrees with *u* because  $\{u_i^I \mid i \in I\}$  is the dual basis of  $\{v_i \mid i \in I\}$ . Since *u* is arbitrary, this proves that  $\iota_I^* \circ \pi^*$  is the identity on  $H^2(BT)$ .

A multi-polytope  $\mathcal{P} = (\Delta, \mathcal{F})$  is associated with real numbers  $c_i$ 's by

$$\mathcal{F}(\{i\}) = \{u \in H^2(BT; \mathbb{R}) \mid \langle u, v_i \rangle = c_i\},\$$

#### THEORY OF MULTI-FANS

and these numbers determine an element  $c_1^T(\mathcal{P}) := \sum_{i=1}^d c_i x_i$  of  $H_T^2(\Delta) \otimes \mathbb{R}$ , which we call the *equivariant first Chern class* of  $\mathcal{P}$ . This gives a bijective correspondence between the set of multi-polytopes defined on  $\Delta$  and  $H_T^2(\Delta) \otimes \mathbb{R}$ . Note that  $\iota_I^*(c_1^T(\mathcal{P}))$  agrees with the vertex  $u_I = \bigcap_{i \in I} \mathcal{F}(\{i\})$ , see Section 5. When  $\Delta$  is non-singular,  $\mathcal{P}$  is a lattice multi-polytope if and only if the  $c_i$ 's are all integers, but otherwise the "if" part does not hold, in other words, an element of  $H_T^2(\Delta)$  is not necessarily realized as the equivariant first Chern class of a lattice multi-polytope. However, there is a nonzero integer *m* such that mx for any  $x \in H_T^2(\Delta)$  is realized as the equivariant first Chern class of a lattice multi-polytope. However, there is

We set  $H^{**}(BT; \mathbb{Q}) = \prod_{a=0}^{\infty} H^q(BT; \mathbb{Q})$ . It is a formal power series ring.

**Lemma 8.3.** For any  $J \in \Sigma$ , the element

$$\sum_{I \in \Sigma^{(n)}} \frac{w(I)\iota_I^*(\prod_{j \in J} (e^{mx_j} - 1))}{|G_I|} \sum_{g \in G_I} \frac{1}{\prod_{i \in I} (1 - \rho_i(g)e^{-u_i^I})}$$

in the quotient field of  $H^{**}(BT; \mathbb{Q})$  actually belongs to  $H^{**}(BT; \mathbb{Q})$ .

Proof. Since  $\prod_{j \in J} (e^{mx_j} - 1)$  is a linear combination of  $\prod_{k \in K} e^{mx_k} = e^{m\sum_{k \in K} x_k}$  for  $K \in \Sigma$ , it suffices to show that

(8.2) 
$$\sum_{I \in \Sigma^{(n)}} \frac{w(I)\iota_I^*(e^{m\sum_{k \in K} x_k})}{|G_I|} \sum_{g \in G_I} \frac{1}{\prod_{i \in I} (1 - \rho_i(g)e^{-u_i^I})} \in H^{**}(BT; \mathbb{Q}).$$

As remarked above,  $m \sum_{k \in K} x_k$  is realized as the equivariant first Chern class of a lattice multi-polytope, so it follows from Corollary 7.6 that

$$\sum_{I \in \Sigma^{(n)}} \frac{w(I)t^{\iota_{I}^{*}(m\sum x_{k})}}{|G_{I}|} \sum_{g \in G_{I}} \frac{1}{\prod_{i \in I} (1 - \rho_{i}(g)t^{-u_{i}'})} \in \mathbb{C}[N^{*}] = R(T).$$

The Chern character  $\mathbb{C}[N^*] = R(T) \to H^{**}(BT; \mathbb{Q})$  mapping  $t^u$  to  $e^u$  extends to a map from  $\mathbb{C}[N^*_{\Delta}]$  and it further extends to a map between the quotient fields. Sending the element above by this extended Chern character, we obtain (8.2).

Let S be the multiplicative set consisting of nonzero homogeneous elements of positive degree in  $H^*(BT; \mathbb{Q})$ . Since  $H^*(BT; \mathbb{Q})$  is a polynomial ring, it can be thought of as a subring of the localized ring  $S^{-1}H^*(BT; \mathbb{Q})$ . We define the index map

$$\pi_! \colon H^*_T(\Delta) \otimes \mathbb{Q} \to S^{-1}H^*(BT;\mathbb{Q})$$

"in cohomology" by

$$\pi_{!}(x) \coloneqq \sum_{I \in \Sigma^{(n)}} \frac{w(I)\iota_{I}^{*}(x)}{|G_{I}| \prod_{i \in I} u_{i}^{I}}$$

(cf. [2, (3.8)]). This map decreases degrees by 2n, and is an  $H^*(BT; \mathbb{Q})$ -module map by Lemma 8.2.

**Lemma 8.4.** The image of  $\pi_1$  lies in  $H^*(BT; \mathbb{Q})$ .

Proof. Since  $\pi_1$  is an  $H^*(BT; \mathbb{Q})$ -module map, it suffices to check the lemma for elements  $x_J$ 's  $(J \in \Sigma)$  by Lemma 8.1. We distinguish two cases.

CASE 1. The case where |J| = n, i.e.,  $J \in \Sigma^{(n)}$ . In this case

$$\iota_I^*(x_J) = \begin{cases} \prod_{i \in J} u_i^I & \text{if } I = J, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore

$$\pi_!(x_J) = \sum_{I \in \Sigma^{(n)}} \frac{w(I)\iota_I^*(x_J)}{|G_I| \prod_{i \in I} u_i^I} = \frac{w(J)}{|G_J|} \in H^0(BT; \mathbb{Q}).$$

CASE 2. The case where |J| < n. In this case we will show that  $\pi_1(x_J) = 0$ . Since  $\rho_i(g) = 1$  for any  $i \in I$  if and only if g is the identity, and

$$\prod_{i \in I} \left( 1 - e^{-u_i^I} \right) = \left( \prod_{i \in I} u_i^I \right) (1 + \text{higher degree term})$$
$$\prod_{j \in J} \left( e^{mx_j} - 1 \right) = m^{|J|} x_J (1 + \text{higher degree term}),$$

the term of lowest degree in Lemma 8.3 (up to a nonzero constant multiple) is

$$\sum_{I\in\Sigma^{(n)}}\frac{w(I)\iota_I^*(x_J)}{|G_I|\prod_{i\in I}u_i^I},$$

that is,  $\pi_!(x_J)$ , and Lemma 8.3 tells us that it is an element of  $H^*(BT; \mathbb{Q})$ . This means that  $\pi_!(x_J) = 0$  because the degree of  $\pi_!(x_J)$  is equal to 2|J| - 2n < 0.

Now, motivated by the description of the cohomology ring of a compact nonsingular toric variety (see p. 106 in [11]), we define  $H^*(\Delta)$  to be the quotient ring of  $H^*_T(\Delta)$  by the ideal generated by  $\pi^*(H^2(BT))$ , in other words,

$$H^*(\Delta) := \mathbb{Z}[x_1, \ldots, x_d]/\mathfrak{A},$$

where  $\mathfrak{A}$  is the ideal generated by all

- (1)  $x_I$  for  $I \notin \Sigma$ ,
- (2)  $\sum_{i=1}^{d} \langle u, v_i \rangle x_i$  for  $u \in N$ .

Since  $\pi_1$  is an  $H^*(BT; \mathbb{Q})$ -module map and  $H^*(BT; \mathbb{Q})/(H^2(BT; \mathbb{Q}))$  is isomorphic to  $H^0(BT; \mathbb{Q}) = \mathbb{Q}, \pi_1$  induces a homomorphism

$$\int_{\Delta} \colon H^*(\Delta) \otimes \mathbb{Q} \to \mathbb{Q},$$

where only elements of degree 2n in  $H^*(\Delta) \otimes \mathbb{Q}$  survive through the map  $\int_{\Lambda}$ .

Remember that  $G_I$  is a finite subgroup of  $\mathbb{R}^d/\mathbb{Z}^d$ . We denote by  $G_\Delta$  the union of  $G_I$  over all  $I \in \Sigma^{(n)}$ . Since  $\rho_i$  is defined on  $\mathbb{R}^d/\mathbb{Z}^d$ ,  $\rho_i(g)$  makes sense for  $g \in G_\Delta$ . It follows from the definition of  $G_I$  and  $\rho_i$  that if  $g \in G_I$ , then  $\rho_i(g) = 1$  for  $i \notin I$ .

We define the *equivariant Todd class*  $\mathcal{T}^{T}(\Delta)$  of the complete simplicial multi-fan  $\Delta$  by

$$\mathcal{T}^{T}(\Delta) \coloneqq \sum_{g \in G_{\Delta}} \prod_{i=1}^{d} \frac{x_{i}}{1 - \rho_{i}(g)e^{-x_{i}}} \in H_{T}^{**}(\Delta) \otimes \mathbb{Q},$$

and the *Todd class*  $\mathcal{T}(\Delta)$  of  $\Delta$  by

$$\mathcal{T}(\Delta) \coloneqq \sum_{g \in G_{\Delta}} \prod_{i=1}^{d} \frac{\bar{x}_{i}}{1 - \rho_{i}(g)e^{-\bar{x}_{i}}} \in H^{**}(\Delta) \otimes \mathbb{Q},$$

where  $\bar{x}_i$  denotes the image of  $x_i \in H_T^*(\Delta)$  in  $H^*(\Delta)$  (cf. [5]). We also define the *first* Chern class  $c_1(\mathcal{P})$  of a multi-polytope  $\mathcal{P}$  defined on  $\Delta$  to be the image of  $c_1^T(\mathcal{P}) \in H_T^2(\Delta) \otimes \mathbb{R}$  in  $H^2(\Delta) \otimes \mathbb{R}$ .

**Theorem 8.5.** If  $\mathcal{P}$  is a simple lattice multi-polytope, then  $\int_{\Lambda} e^{c_1(\mathcal{P})} \mathcal{T}(\Delta) = \sharp(\mathcal{P})$ .

Proof. We shall compute  $\pi_!(e^{c_1^T(\mathcal{P})}\mathcal{T}^T(\Delta))$ . For that, we need to see  $\iota_I^*(\mathcal{T}^T(\Delta))$ . Let  $g \in G_{\Delta}$ . If  $g \notin G_I$ , then there is an  $i \notin I$  such that  $\rho_i(g) \neq 1$ ; so

$$\iota_I^*\left(\frac{x_i}{1-\rho_i(g)e^{-x_i}}\right)=0$$

for such *i* because the Maclaurin expansion of  $x_i/(1 - \rho_i(g)e^{-x_i})$  has no constant term and  $\iota_I^*(x_i) = 0$ . Therefore, only elements *g* in  $G_I$  contribute to  $\iota_I^*(\mathcal{T}^T(\Delta))$ . Now suppose  $g \in G_I$ . Then  $\rho_i(g) = 1$  for  $i \notin I$ , so

$$\iota_I^*\left(\frac{x_i}{1-\rho_i(g)e^{-x_i}}\right)=1$$

for such *i* because the Maclaurin expansion of  $x_i/(1-\rho_i(g)e^{-x_i})$  has the constant term 1 and  $\iota_I^*(x_i) = 0$ . Finally, since  $\iota_I^*(x_i) = u_i^I$  for  $i \in I$ , we thus have

$$\iota_I^*(\mathcal{T}^T(\Delta)) = \sum_{g \in G_I} \prod_{i \in I} \frac{u_i^I}{1 - \rho_i(g)e^{-u_i^I}}.$$

This together with the definition of  $\pi_1$  and Corollary 7.6 shows that

$$\pi_{!}(e^{c_{1}^{T}(\mathcal{P})}\mathcal{T}^{T}(\Delta)) = \pi_{!}\left(e^{c_{1}^{T}(\mathcal{P})}\sum_{g\in G_{\Delta}}\prod_{i=1}^{d}\frac{x_{i}}{1-\rho_{i}(g)e^{-x_{i}}}\right)$$
$$= \sum_{I\in\Sigma^{(n)}}\frac{w(I)e^{\iota_{I}^{*}(c_{1}^{T}(\mathcal{P}))}}{|G_{I}|}\sum_{g\in G_{I}}\frac{1}{\prod_{i\in I}(1-\rho_{i}(g)e^{-u_{i}^{I}})}$$
$$= \sum_{u\in H^{2}(BT)}\mathrm{DH}_{\mathcal{P}_{+}}(u)e^{u}.$$

This implies that

$$\int_{\Delta} e^{c_1(\mathcal{P})} \mathcal{T}(\Delta) = \sum_{u \in H^2(BT)} \mathrm{DH}_{\mathcal{P}_+}(u) = \sharp(\mathcal{P}).$$

REMARK. The argument developed above in this section is purely combinatorial, but it is possible to take a topological approach. Namely, associated with a complete simplicial multi-fan  $\Delta$ , one can construct a torus space  $M_{\Delta}$  with  $H_T^*(M_{\Delta}; \mathbb{Q}) =$  $H_T^*(\Delta) \otimes \mathbb{Q}$  (see [8]). It is not necessarily a manifold but has a fundamental class so that the equivariant Gysin homomorphism  $\pi_1: H_T^*(M_{\Delta}; \mathbb{Q}) = H_T^*(\Delta) \otimes \mathbb{Q} \to$  $H_T^{*-2n}(pt; \mathbb{Q}) = H^{*-2n}(BT; \mathbb{Q})$ , that is, the index map, can be defined.

As an application of the theorem above, we shall show that Khovanskii-Pukhlikov formula, which relates a certain variation of the volume of a simple convex lattice polytope to the number of lattice points in it, can be generalized to simple multipolytopes. We begin with

Lemma 8.6.

$$\operatorname{vol}(\mathcal{P}) = \frac{1}{n!} \int_{\Delta} c_1(\mathcal{P})^n = \int_{\Delta} e^{c_1(\mathcal{P})}$$

for a simple multi-polytope  $\mathcal{P}$ .

Proof. The latter equality is obvious because only elements of degree 2n in  $H^*(\Delta) \otimes \mathbb{R}$  survive through the map  $\int_{\Delta}$ . We shall prove the former equality.

STEP 1. If  $\mathcal{P}$  is a lattice multi-polytope, then Theorem 8.5 applied to  $\nu \mathcal{P}$  for any integer  $\nu$  implies

$$\int_{\Delta} e^{c_1(\nu \mathcal{P})} \mathcal{T}(\Delta) = \sharp(\nu \mathcal{P}).$$

We compare the coefficients of  $\nu^n$  at the both sides above. Since  $c_1(\nu \mathcal{P}) = \nu c_1(\mathcal{P})$ , the coefficient of  $\nu^n$  at the left-hand side is  $(1/n!) \int_{\Lambda} c_1(\mathcal{P})^n$ , while the one at the right-

hand side is  $vol(\mathcal{P})$  by Theorem 7.2 (3). Therefore the lemma is proven for a lattice multi-polytope  $\mathcal{P}$ .

STEP 2. If  $\mathcal{P}$  is *rational*, by which we mean that there is a nonzero integer m such that  $m\mathcal{P}$  is a lattice multi-polytope, then  $\operatorname{vol}(m\mathcal{P}) = (1/n!) \int_{\Delta} c_1(m\mathcal{P})^n$  by Step 1. Since  $\operatorname{vol}(m\mathcal{P}) = m^n \operatorname{vol}(\mathcal{P})$  and  $c_1(m\mathcal{P}) = mc_1(\mathcal{P})$ , the lemma is proven for a rational multi-polytope  $\mathcal{P}$ .

STEP 3. The functions  $vol(\cdot)$  and  $\int_{\Delta} c_1(\cdot)^n$  are defined on the vector space  $H_T^2(\Delta) \otimes \mathbb{R}$  through the equivariant first Chern class, and they are obviously continuous. By Step 2 they agree on all rational multi-polytopes which form a dense subset of the vector space, so they must agree on the entire vector space by continuity. This completes the proof of the lemma.

Multi-polytopes defined on  $\Delta$  form a vector space isomorphic to  $H_T^2(\Delta) \otimes \mathbb{R}$ through the equivariant first Chern class, and Lemma 8.6 implies that the volume function is a homogeneous polynomial function of degree *n*. In fact, if one writes  $c_1^T(\mathcal{P}) = \sum_{i=1}^d c_i x_i$ , then  $\operatorname{vol}(\mathcal{P})$  is a homogeneous polynomial in  $c_1, \ldots, c_d$  of degree *n*.

For  $h = (h_1, \ldots, h_d) \in \mathbb{R}^d$ , we denote by  $\mathcal{P}_h$  a multi-polytope with  $c_1^T(\mathcal{P}_h) = \sum_{i=1}^d (c_i + h_i)x_i$ . Since  $c_1(\mathcal{P}_h) = \sum_{i=1}^d (c_i + h_i)\bar{x}_i$ , Lemma 8.6 applied to  $\mathcal{P}_h$  implies that  $\operatorname{vol}(\mathcal{P}_h)$  is a polynomial in  $h_1, \ldots, h_d$  (of total degree *n*). We define the *Todd operator* as follows:

$$\mathcal{T}\left(\frac{\partial}{\partial h}\right) \coloneqq \sum_{g \in G_{\Delta}} \prod_{i=1}^{d} \frac{\partial/\partial h_{i}}{1 - \rho_{i}(g)e^{-\partial/\partial h_{i}}}.$$

Although the Todd operator is of infinite order, its operation on  $vol(\mathcal{P}_h)$  converges because  $vol(\mathcal{P}_h)$  is a polynomial in  $h_1, \ldots, h_d$ . The following theorem extends the Khovanskii-Pukhlikov formula to simple lattice multi-polytopes.

**Theorem 8.7.** If  $\mathcal{P}$  is a simple lattice multi-polytope, then

$$\mathcal{T}\left(\frac{\partial}{\partial h}\right) \operatorname{vol}(\mathcal{P}_h)|_{h=0} = \sharp(\mathcal{P}).$$

Proof. An elementary computation shows that

$$\frac{\partial/\partial h_i}{1-\rho_i(g)e^{-\partial/\partial h_i}}e^{(c_i+h_i)\bar{x}_i}|_{h_i=0}=e^{c_i\bar{x}_i}\frac{\bar{x}_i}{1-\rho_i(g)e^{-\bar{x}_i}}.$$

Therefore, it follows from Lemma 8.6 and Theorem 8.5 that

$$\mathcal{T}\left(\frac{\partial}{\partial h}\right)\operatorname{vol}(\mathcal{P}_{h})|_{h=0} = \mathcal{T}\left(\frac{\partial}{\partial h}\right)\int_{\Delta}e^{c_{1}(\mathcal{P}_{h})}|_{h=0} = \int_{\Delta}\mathcal{T}\left(\frac{\partial}{\partial h}\right)e^{c_{1}(\mathcal{P}_{h})}|_{h=0}$$

A. HATTORI AND M. MASUDA

$$\begin{split} &= \int_{\Delta} \sum_{g \in G_{\Delta}} \prod_{i=1}^{d} \frac{\partial/\partial h_{i}}{1 - \rho_{i}(g)e^{-\partial/\partial h_{i}}} e^{(c_{i}+h_{i})\bar{x}_{i}}|_{h_{i}=0} \\ &= \int_{\Delta} \sum_{g \in G_{\Delta}} \prod_{i=1}^{d} e^{c_{i}\bar{x}_{i}} \frac{\bar{x}_{i}}{1 - \rho_{i}(g)e^{-\bar{x}_{i}}} \\ &= \int_{\Delta} e^{c_{1}(\mathcal{P})} \mathcal{T}(\Delta) = \sharp(\mathcal{P}), \end{split}$$

proving the theorem.

REMARK. One can reformulate the Khovanskii-Pukhlikov formula as follows. As remarked above, the volume function vol is a polynomial in  $c_1, \ldots, c_d$ , so one can apply the Todd operator  $\mathcal{T}(\partial/\partial c)$  (with the variables  $c = (c_1, \ldots, c_d)$  instead of  $h = (h_1, \ldots, h_d)$ ) to the volume function vol and evaluate at a simple lattice multi-polytope  $\mathcal{P}$ . The same argument as in the proof of Theorem 8.7 shows that the evaluated value agrees with  $\sharp(\mathcal{P})$ .

### 9. Multi-fan of a torus manifold

In this section we introduce the notion of a torus manifold and associate a complete non-singular multi-fan with it. A compact non-singular toric variety provides an example of a torus manifold, but the class of torus manifolds is much wider than that of compact non-singular toric varieties, (apparently, even wider than that of unitary toric manifolds introduced in [27]). The basic theory of toric varieties says that there is a one-to-one correspondence between compact non-singular toric varieties and complete non-singular fans. This correspondence is extended in one direction, namely from torus manifolds to complete non-singular multi-fans. But the usual way to associate a fan with a toric variety (see [11, Section 2.3]) does not work in our extended category. However, when a toric variety is compact and non-singular, the corresponding (complete and non-singular) fan can be reproduced using equivariant cohomology and this argument works even for torus manifolds. The idea is essentially same as in [27].

We begin with the definition of a torus manifold. An elementary representation theory of a torus group tells us that if an *m*-dimensional torus  $(S^1)^m$  acts effectively and smoothly on a connected smooth manifold of dimension 2n with non-empty fixed point set, then  $m \leq n$  and the dimension of the fixed point set is at most 2(n - m). We are interested in an extreme case m = n. Let M be a closed, connected, smooth manifold of dimension 2n with an effective smooth action of an *n*-dimensional torus group  $T = (S^1)^n$  such that the fixed point set  $M^T$  is non-empty. Then  $M^T$  is necessarily isolated. A closed, connected, codimension two submanifold of M is called *characteristic* if it is a connected component of the set fixed pointwise by a certain circle subgroup of T and contains at least one T-fixed point. Since M is compact, there are only finitely many characteristic submanifolds. We denote them by  $M_i$  (i = 1, ..., d).

They are orientable if M is orientable.

DEFINITION. Let M be a closed, connected, oriented, smooth manifold M of dimension 2n with an effective smooth action of an *n*-dimensional torus group T with non-empty fixed point set  $M^T$ . M will be called a *torus manifold* if a prefered orientation is given for each characteristic submanifold  $M_i$ .

A toric variety X (of dimension n) is a normal complex algebraic variety of complex dimension n with an effective algebraic action of  $(\mathbb{C}^*)^n$  having a dense orbit. If X is compact and non-singular, then X with the restricted action of  $T (\subset (\mathbb{C}^*)^n)$  provides an example of a torus manifold of dimension 2n. In this case, characteristic submanifolds are  $(\mathbb{C}^*)^n$ -invariant divisors. They have canonical orientations since they are complex manifolds. Similarly, when a torus manifold is equipped with a T-invariant unitary structure, characteristic submanifolds have canonical orientations. With these orientations of characteristic submanifolds, the torus manifold will be called a *unitary torus manifold* (also called a unitary toric manifold in [27]).

EXAMPLE 9.1. A complex projective space  $\mathbb{C}P^n$  with an action of  $(\mathbb{C}^*)^n$  given by

$$[z_0, z_1, \ldots, z_n] \rightarrow [z_0, g_1 z_1, \ldots, g_n z_n],$$

where  $[z_0, z_1, \ldots, z_n] \in \mathbb{C}P^n$  and  $(g_1, \ldots, g_n) \in (\mathbb{C}^*)^n$ , is a compact non-singular toric variety. This with the restricted *T*-action is a torus manifold and there are n + 1 characteristic submanifolds, that are respectively defined by  $z_i = 0$  for  $i = 0, 1, \ldots, n$ .

There are many torus manifolds which do not arise from compact non-singular toric varieties, see [8], [27], [30].

Henceforth M will denote a torus manifold of dimension 2n. Let  $p \in M^T$ . Since  $M^T$  is isolated, the tangential T-representation  $\tau_p M$  of M at p has no trivial factor, so it decomposes into the direct sum of n irreducible real two-dimensional T-representations. This implies that there are exactly n characteristic submanifolds which contain p. In fact, an irreducible factor in  $\tau_p M$  corresponds to the normal direction to a characteristic submanifold at p. We set

$$\Sigma(M) \coloneqq \left\{ I \subset \{1, \ldots, d\} \mid \left(\bigcap_{i \in I} M_i\right)^T \neq \phi \right\}.$$

We add an empty set to  $\Sigma(M)$  as a member, so that  $\Sigma(M)$  becomes an augmented simplicial set. The observation above implies that the cardinality of an element in  $\Sigma(M)$  is at most *n* and there is an element in  $\Sigma(M)$  with cardinality *n*.

The augmented simplicial set  $\Sigma(M)$  is closely related to the ring structure of the equivariant cohomology  $H_T^*(M)$  of M with integer coefficients. Let us explain this

briefly. Since  $M_i$  and M are oriented closed T-manifolds and the codimension of  $M_i$ is two, the inclusion map from  $M_i$  to M induces a Gysin homomorphism  $H_T^*(M_i) \rightarrow H_T^{*+2}(M)$  in equivariant cohomology which raises dgrees by two (see [23] for example). Denote by  $\xi_i \in H_T^2(M)$  the image of the identity element in  $H_T^0(M_i)$ . We may think of  $\xi_i$  as the Poincaré dual of  $M_i$  (considered as a cycle in M) in equivariant cohomology. If the orientation on M or  $M_i$  is reversed, then  $\xi_i$  turns into  $-\xi_i$ .

We take a polynomial ring  $\mathbb{Z}[x_1, \ldots, x_d]$  in *d*-variables and consider a map

$$\varphi \colon \mathbb{Z}[x_1,\ldots,x_d] \to H^*_T(M)$$

which sends  $x_i$  to  $\xi_i$ . This map is often surjective. Here is a case.

**Proposition 9.2** ([27], Proposition 3.4.). Suppose that  $H^*(M)$  is generated by elements in  $H^2(M)$  as a ring (this is the case when M is a compact non-singular toric variety). Then the map  $\varphi$  is surjective and the kernel is the ideal generated by monomials  $\prod_{i \in I} x_i$  for all subsets  $I \subset \{1, \ldots, d\}$  such that  $I \notin \Sigma(M)$ . In other words,  $H_T^*(M)$  is isomorphic to the face ring (or Stanley-Reisner ring) of  $\Sigma(M)$ .

The equivariant cohomology  $H_T^*(M)$  has a finer structure than the ring structure. The map  $\pi$  collapsing M to a point induces a homomorphism  $\pi^* \colon H_T^*(pt) = H^*(BT) \to H_T^*(M)$ , so that  $H_T^*(M)$  can be viewed as an algebra over  $H^*(BT)$ through  $\pi^*$ . This algebra structure over  $H^*(BT)$  cannot be determined by  $\Sigma(M)$  and contains more information on the torus manifold M. To see the algebra structure, it is enough to see the image of  $H^2(BT)$  by  $\pi^*$  because  $H^*(BT)$  is a polynomial ring generated by elements in  $H^2(BT)$ .

**Lemma 9.3** ([27], Lemma 1.5.). For each  $i \in \{1, ..., d\}$  there exists a unique element  $v_i \in H_2(BT)$  such that

$$\pi^*(u) = \sum_{i=1}^d \langle u, v_i \rangle \xi_i \qquad modulo \ H^*(BT) \text{-torsions}$$

for any  $u \in H^2(BT)$ , where  $\langle , \rangle$  denotes the usual pairing between cohomology and homology.

Proof. The proof is given in [27], but we shall give a simple proof for the reader's convenience when M is as in Proposition 9.2. Since  $H_T^2(M)$  is additively generated by  $\xi_i$ 's, one can express

$$\pi^*(u) = \sum_{i=1}^d v_i(u)\xi_i$$

with a unique integer  $v_i(u)$  depending on u for each i. We view  $v_i(u)$  as a function of

 $u \in H^2(BT)$ . Since it is linear, it defines an element  $v_i$  of  $Hom(H^2(BT), \mathbb{Z}) = H_2(BT)$ such that  $v_i(u) = \langle u, v_i \rangle$ .

Note. A geometrical interpretation of the vectors  $v_i$  will be given in Section 12.

In order to introduce a multi-fan, we adopt  $H_2(BT)$  as the lattice N and identify  $H_2(BT; \mathbb{R})$  with the vector space  $N_{\mathbb{R}} = N \otimes \mathbb{R}$ . Then we define a map

$$C(M): \Sigma(M) \rightarrow \operatorname{Cone}(N)$$

by sending  $I \in \Sigma(M)$  to the cone in  $H_2(BT; \mathbb{R})$  spanned by  $v_i$ 's  $(i \in I)$  (and the empty set to  $\{0\}$ ).

Finally we shall define a pair of weight functions on maximal cones of dimension *n*. Remember that a characteristic submanifold  $M_i$  is a connected component of the set fixed pointwise by a certain circle subgroup, say  $T_i$ , of *T*. It turns out that  $T_i$ agrees with the circle subgroup determined by  $v_i \in H_2(BT)$  through the natural identification  $H_2(BT) \cong \text{Hom}(S^1, T)$  ([27], Lemma 1.10). Therefore  $M_I := \bigcap_{i \in I} M_i$  is fixed pointwise by a subtorus  $T_I$  generated by  $T_i$ 's for  $i \in I$ .

**Lemma 9.4** ([27], Lemma 1.7.). Suppose  $I \in \Sigma(M)^{(n)}$ . Then the set  $\{v_i \mid i \in I\}$  forms a basis of  $H_2(BT)$ , so that  $M_I$  is a subset of  $M^T$  and the cone C(M)(I) is of dimension n.

A fixed point  $p \in M^T$  belongs to  $M_I$  for some  $I \in \Sigma^{(n)}$ , and the tangent space  $\tau_p M$  at  $p \in M_I$  naturally decomposes into

$$au_p M \cong igoplus_{i \in I} ( au_p M / au_p M_i).$$

The orientations on M and  $M_i$  determine an orientation on  $\tau_p M / \tau_p M_i$  for each i, and then an orientation on  $\tau_p M$  through the above isomorphism. On the other hand,  $\tau_p M$ has a given orientation since M is oriented. These two orientations on  $\tau_p M$  may disagree. We define the sign  $\epsilon_p$  at p to be +1 or -1 according as the two orientations agree or disagree, and set

$$w(M)^+(I) :=$$
 the number of  $\{p \in M_I \mid \epsilon_p = +1\},$   
 $w(M)^-(I) :=$  the number of  $\{p \in M_I \mid \epsilon_p = -1\}.$ 

Note that  $w(M)^+(I) = 1$  and  $w(M)^-(I) = 0$  for all  $I \in \Sigma^{(n)}$  if M is a compact non-singular toric variety.

DEFINITION. We call the triple  $\Delta(M) := (\Sigma(M), C(M), w(M)^{\pm})$  the multi-fan of M.

A. HATTORI AND M. MASUDA

A characteristic submanifold of  $M_i$  is a connected component of  $M_i \cap M_j$  for some j containing a T-fixed point. We give it the orientation induced from those on  $M_i$  and  $M_j$ . With these orientations equipped,  $M_i$ , on which  $T/T_i$  acts effectively, is considered as a torus manifold. If  $M_i \cap M_j$  is connected for any  $j \in \Sigma(M)_i^{(1)}$  (this is the case when M is a compact non-singular toric variety), then the multi-fan  $\Delta(M_i)$  of  $M_i$  agrees with the projected multi-fan  $\Delta(M)_i$  with respect to  $\{i\} \in \Sigma(M)^{(1)}$ . They are different otherwise but there is a natural surjective map from  $\Sigma(M_i)$  to  $\Sigma(M)_i$ .

Similarly, a connected component of  $M_K$  for  $K \in \Sigma(M)$  containing a *T*-fixed point is considered as a torus manifold, and  $\Delta(M_K)$  agrees with  $\Delta(M)_K$  if  $M_K$  and  $M_K \cap M_j$  are connected for all  $j \in \Sigma(M)_K^{(1)}$ , but otherwise they are different although there is a natural surjective map from  $\Sigma(M_K)$  to  $\Sigma(M)_K$ , where  $\Sigma(M_K)$  is an augmented simplicial set obtained from the union of the simplicial sets associated with the connected components of  $M_K$ .

The multi-fan  $\Delta(M)$  is non-singular by Lemma 9.4. We shall show that it is complete.

#### **Lemma 9.5.** $\Delta(M)$ is complete.

Proof. As we remarked in Section 2 after the definition of the completeness of a multi-fan, it suffices to prove the pre-completeness of  $\Delta(M)_J$  for any  $J \in \Sigma(M)^{(n-1)}$ . Choose a generic vector v from  $N = H_2(BT)$ . The sign  $(-1)^{\{i\}}$  for  $i \in \Sigma(M)_J^{(1)}$  is defined as in Section 5 with respect to the projection image of v on the quotient lattice of N by the sublattice generated by  $C(M)(J) \cap N$ . The pre-completeness of  $\Delta(M)_J$  is equivalent to the equality:

$$\sum_{\{i\}\in\Sigma(M)_{I}^{(i)}} (-1)^{\{i\}} w(M)_{J}(\{i\}) = 0,$$

which we will verify in the following. Since |J| = n - 1, a connected component of  $M_J$  containing a *T*-fixed point is a 2-dimensional sphere on which  $T^J := T/T_J$  acts effectively. We denote those connected components by  $S_{\alpha}^2$ 's. They are torus manifolds equipped with the orientations discussed before this lemma. Since  $S_{\alpha}^2$  has two  $T^J$ -fixed points,  $\Sigma(S_{\alpha}^2)^{(1)}$  consists of two elements, denoted by  $\alpha_{\pm}$ , corresponding to the  $T^J$ -fixed points. One easily checks that the multi-fan  $\Delta(S_{\alpha}^2)$  of  $S_{\alpha}^2$  is complete, which is equivalent to the equality:

(9.1) 
$$(-1)^{\alpha_+} w(S_{\alpha}^2)(\alpha_+) + (-1)^{\alpha_-} w(S_{\alpha}^2)(\alpha_-) = 0.$$

As discussed before this lemma, we have a natural map  $\pi_J: \Sigma(M_J) \to \Sigma(M)_J$ . Note that if  $\pi_J(\alpha_{\epsilon}) = \{i\}$  where  $\epsilon$  stands for + or -, then  $(-1)^{\alpha_{\epsilon}} = (-1)^{\{i\}}$ . On the other hand, we have

$$w(M)_J(\{i\}) = \sum_{\pi_J(\alpha_\epsilon) = \{i\}} w(S_\alpha^2)(\alpha_\epsilon).$$

Therefore

$$\sum_{[i]\in\Sigma(M)_J^{(l)}} (-1)^{\{i\}} w(M)_J(\{i\}) = \sum_{\alpha_{\epsilon}} (-1)^{\alpha_{\epsilon}} w(S_{\alpha}^2)(\alpha_{\epsilon}),$$

which vanishes by (9.1), proving the lemma.

We make a remark on orientations at this point. Choose an orientation on T and fix it. It induces an orientation on  $H_2(BT; \mathbb{R})$ , so that  $[\Delta(M)] \in H_{n-1}(\Sigma(M))$  is defined. If the orientation on T or M is reversed, then  $[\Delta(M)]$  turns into  $-[\Delta(M)]$ . But we have

**Lemma 9.6.**  $[\Delta(M)]$  does not depend on the orientations on  $M_i$ 's.

Proof. Recall that the cycle which defines  $[\Delta(M)]$  is  $\sum_{I \in \Sigma(M)^{(n)}} w(M)(I) \langle I \rangle$ . We reverse the orientation on  $M_i$ . Obviously, w(M)(I) and  $\langle I \rangle$  remain unchanged unless  $i \in I$ . Suppose  $i \in I$ . Then, since the orientation on  $\tau_p M / \tau_p M_i$  is reversed,  $w(M)^+(I)$  and  $w(M)^-(I)$  will be interchanged, so that w(M)(I) turns into -w(M)(I). As for  $\langle I \rangle$ ,  $\xi_i$  turns into  $-\xi_i$  as remarked before and hence so does  $v_i$  by Lemma 9.3. Thus,  $\langle I \rangle$  turns into  $-\langle I \rangle$  if  $i \in I$ . After all,  $w(M)(I) \langle I \rangle$  does not depend on the orientations on  $M_i$ 's for any  $I \in \Sigma(M)^{(n)}$ .

Remember that there is a canonical isomorphism  $\text{Hom}(T, S^1) \cong H^2(BT)$ . We denote by  $t^u$  the element in  $\text{Hom}(T, S^1)$  corresponding to  $u \in H^2(BT)$ . Elements of  $\text{Hom}(T, S^1)$  are complex one-dimensional representations of T and they generate the representation ring R(T) of T which is identified with the group ring of  $H^2(BT)$ . Since  $\xi_i$  is the image of  $1 \in H^0_T(M_i)$  by the equivariant Gysin map from  $M_i$  to M, its restriction to a T-fixed point p in  $M_i$ , denoted by  $\xi_i|_p$ , gives the equivariant Euler class of the T-representation  $\tau_p M/\tau_p M_i$ ; so  $\tau_p M/\tau_p M_i = t^{\xi_i|_p}$ . On the other hand, the equality in Lemma 9.3 restricted to p shows that  $\{\xi_i|_p \mid i \in I\}$  is the dual basis of  $\{v_i \mid i \in I\}$ , so  $\xi_i|_p$  is independent of the choice of  $p \in M_I$  and  $\xi_i|_p = u_i^I$  in the notation of Section 7. Therefore we have

$$\tau_p M = \bigoplus_{i \in I} t^{u_i^I}$$

as a *T*-representation whenever  $p \in M_I$ .

The elements  $\xi_i$ 's (i = 1, ..., d) generate  $H_T^2(M)$  additively modulo  $H^*(BT)$ -torsions ([27, Lemma 3.2]) and the torsion elements vanish when restricted to the fixed

point set  $M^T$  because  $H^*_T(M^T)$  is a free  $H^*(BT)$ -module. Since the restriction  $\xi_i|_p$  $(p \in M_I)$  depends on only I, we shall denote the restriction of an element  $\xi \in H^2_T(M)$  to a point in  $M_I$  by  $\xi|_I$ . Note that

(9.2) 
$$\xi_i|_I = \begin{cases} u_i^I & \text{if } i \in I, \\ 0 & \text{otherwise.} \end{cases}$$

**Lemma 9.7.** For any  $\xi \in H^2_T(M)$ ,

$$\sum_{I\in\Sigma(M)^{(n)}}\frac{w(M)(I)t^{\xi|I}}{\prod_{i\in I}(1-t^{-u_i^I})}$$

is an element of R(T) when M is a torus manifold.

Proof. Since  $\xi_i$ 's generate  $H_T^2(M)$  additively modulo  $H^*(BT)$ -torsions,  $\xi = \sum_{i=1}^d c_i \xi_i$  modulo  $H^*(BT)$ -torsions with some integers  $c_i$ 's. We define a map  $\mathcal{F}_{\xi}$ :  $\Sigma(M)^{(1)} \to \operatorname{HP}(H^2(BT;\mathbb{R}))$  by

$$\mathcal{F}_{\xi}(\{i\}) := \{ u \in H^2(BT; \mathbb{R}) \mid \langle u, v_i \rangle = c_i \}.$$

The pair  $(\Delta(M), \mathcal{F}_{\xi})$  is a lattice multi-polytope, and  $\bigcap_{i \in I} \mathcal{F}_{\xi}(\{i\}) = \xi|_{I}$  for  $I \in \Sigma(M)^{(n)}$  which follows from (9.2). Since  $\Delta(M)$  is non-singular by Lemma 9.4 and complete by Lemma 9.5, the lemma follows from Corollary 7.4 applied to the multi-polytope  $(\Delta(M), \mathcal{F}_{\xi})$ .

#### 10. $T_{y}$ -genus of a torus manifold

When M is a unitary torus manifold, the localization formula of the  $T_y$ -genus  $T_y[M]$  of M tells us that

(10.1) 
$$T_{y}[M] = \sum_{I \in \Sigma^{(n)}} w(M)(I) \frac{\prod_{i \in I} (1 + yt^{-u_{i}^{l}})}{\prod_{i \in I} (1 - t^{-u_{i}^{l}})}$$

and this is actually a polynomial in y with constant coefficients. As is well known,  $T_0[M]$  agrees with the Todd genus of M and  $T_1[M]$  agrees with the signature of M, see [20]. The  $T_y$ -genus is a genus for unitary manifolds and it is not defined for general torus manifolds. But the right-hand side of (10.1) makes sense even for a torus manifold, and we take it as the definition of the  $T_y$ -genus  $T_y[M]$  of M and define the Todd genus of M to be  $T_0[M]$ . Note that the signature of M is already defined for a torus manifold M because M is an oriented closed manifold, and that it agrees with  $T_1[M]$  which follows from the Atiyah-Singer G-signature theorem.

**Theorem 10.1.** Let M be a torus manifold of dimension 2n. Then

$$T_{y}[M] = T_{y}[\Delta(M)] = \sum_{m=0}^{n} e_{n-m}(\Delta(M))(-1-y)^{m}.$$

(see Section 3 for  $e_q(\Delta(M))$ .) In particular, the Todd genus  $T_0[M]$  of M equals  $deg(\Delta)$ .

Proof. Look at the expansion of the right-hand side of (10.1) with respect to y. It follows from (9.2) and Lemma 9.7 that all coefficients of powers of y in (10.1) are elements of R(T). Take a generic vector  $v \in H_2(BT)$  and evaluate the right-hand side of (10.1) on v. Then we get the following polynomial in y whose coefficients are Laurent polynomials in z:

(10.2) 
$$\sum_{I \in \Sigma^{(n)}} w(M)(I) \frac{\prod_{i \in I} (1 + yz^{-\langle u_i^I, v \rangle})}{\prod_{i \in I} (1 - z^{-\langle u_i^I, v \rangle})}$$

It is easily seen that (10.2) approaches to a polynomial in y with constant coefficients if z tends either to 0 or to  $\infty$ . This means that (10.2) itself is a polynomial with constant coefficients. Since v is generic, this implies that (10.1), that is  $T_y[M]$ , is actually a polynomial with constant coefficients equal to (10.2). Then, by letting z tend to 0, we obtain

$$T_{y}[M] = \sum_{I \in \Sigma^{(n)}} w(M)(I)(-y)^{\mu(I)},$$

where  $\mu(I) = \sharp\{i \in I \mid \langle u_i^I, v \rangle > 0\}$ . This  $\mu(I)$  agrees with the  $\mu(I)$  in Section 3 because  $\{u_i^I \mid i \in I\}$  is the dual basis of  $\{v_i \mid i \in I\}$ . Hence  $T_y[M] = T_y[\Delta(M)]$ , proving the former equality in the theorem. The latter follows from Corollary 3.3.

As noted in the definition of  $T_y[\Delta]$  in Section 3,  $T_0[\Delta(M)] = \deg(\Delta(M))$ . Since  $T_0[M] = T_0[\Delta(M)]$ , the last statement in the theorem follows.

**Corollary 10.2.** The signature Sign(M) of a torus manifold M is given by

$$\operatorname{Sign}(M) = \sum_{m=0}^{n} (-2)^{m} e_{n-m}(\Delta(M)).$$

If T[M] = 1 and w(M)(I) = 1 for all  $I \in \Sigma(M)^{(n)}$ , then  $e_q(\Delta(M))$  agrees with the number of cones of dimension q in  $\Delta(M)$ .

Proof. Since Sign(M) equals  $T_1[M]$ , the former statement follows from Theorem 10.1. The latter statement is noted in the definition of  $e_q(\Delta)$  in Section 3. REMARK. If M is a compact non-singular toric variety, then T[M] = 1 and w(M)(I) = 1 for all  $I \in \Sigma(M)^{(n)}$ , and the formula above is already known in that case ([29, Theorem 3.12 (3)]).

### 11. Equivariant index of a torus manifold

If M is a unitary torus manifold, then the map  $\pi$  collapsing M to a point induces, in equivariant K-theory, the equivariant Gysin homomorphism

$$\pi_!$$
:  $K_T(M) \to K_T(pt) = R(T)$ .

If E is a complex T-vector bundle over M, then  $\pi_1(E)$  equals the index of a Dirac operator twisted by E. It is sometimes called the equivariant Riemann-Roch number. The Todd genus of M is equal to  $\pi_1(1)$ .

Let L be a complex T-line bundle over a unitary torus manifold M. Since  $\pi_!(L)$  is an element of R(T), one can express

(11.1) 
$$\pi_!(L) = \sum_{u \in H^2(BT)} m_L(u) t^u$$

with integers  $m_L(u)$  which are zero for all but finitely many elements u. In this section we describe the multiplicity  $m_L(u)$  of  $t^u$  in terms of the (shifted) moment map associated with L when M is a torus manifold. For that, we need to define  $\pi_1(L)$  when Mis a torus manifold. This is done as follows. When M is a unitary torus manifold, the localization formula applied to  $\pi_1(L)$  tells us that

(11.2) 
$$\pi_!(L) = \sum_{I \in \Sigma(M)^{(n)}} \frac{w(M)(I)t^{c_1^*(L)|_I}}{\prod_{i \in I} (1 - t^{-u_i^I})}$$

where  $c_1^T(L) \in H_T^2(M)$  denotes the equivariant first Chern class of L. (Note that  $t^{c_1^T(L)|_I}$  is nothing but the complex one-dimensional T-representation obtained by restricting L to a point in  $M_I$ .) The right-hand side of (11.2) is an element of R(T) by Lemma 9.7 whenever M is a torus manifold although  $\pi_1$  may not be defined. Thus we define  $\pi_1(L)$  as the right-hand side of (11.2) when M is a torus manifold, and then define  $m_L(u)$  as before using (11.1).

In the following, we will make the following assumption on a torus manifold M, which is satisfied for compact non-singular toric varieties with restricted T-actions: all isotropy subgroups of M are subtori of T and each connected component fixed pointwise by a subtorus contains at least one T-fixed point. Then the union  $\bigcup_{i=1}^{d} M_i$  is the set of points with nontrivial isotropy subgroups, and it follows from the slice theorem that the orbit space M/T is a compact connected smooth manifold of dimension n with  $\bigcup_{i=1}^{d} M_i/T$  as boundary (after the corners are rounded).

We make a further remark on orientations. The orbit space M/T is orientable (see

THEORY OF MULTI-FANS

[27], Lemma 6.7) and we orient it in such a way that the orientation on T followed by that of M/T agrees with that of M times  $(-1)^{n(n-1)/2}$ . This determines a fundamental class in  $H_n(M/T, \partial(M/T))$  and hence in  $H_{n-1}(\partial(M/T))$ , denoted by  $[\partial(M/T)]$ , through the boundary homomorphism from  $H_n(M/T, \partial(M/T))$  to  $H_{n-1}(\partial(M/T))$ .

Since  $H_T^2(M)$  is additively generated by  $\xi_i$ 's (i = 1, ..., d) modulo  $H^*(BT)$ torsions,  $c_1^T(L) = \sum_i c_i \xi_i$  modulo  $H^*(BT)$ -torsions with some integers  $c_i$ 's. Associated
with L, there is defined the moment map  $\Phi_L \colon M \to H^2(BT; \mathbb{R}) = L(T)^*$ . It maps  $M_i$  into an affine hyperplane  $\{u \in H^2(BT; \mathbb{R}) \mid \langle u, v_i \rangle = c_i\}$  for each *i* (see [27],
Lemma 6.5). We slightly shift  $\Phi_L$  so that the shifted map  $\Phi'_L$  maps  $M_i$  into

$$\mathcal{F}_L'(\{i\}) \coloneqq \left\{ u \in H^2(BT; \mathbb{R}) \mid \langle u, v_i \rangle = c_i + \frac{1}{2} \right\}$$

for each *i*. In fact,  $\Phi'_L$  is defined as follows. Let *K* be a complex *T*-line bundle over *M* with  $c_1^T(K) = -\sum_{i=1}^d \xi_i$ . Such *K* exists ([19]). When *M* is a compact non-singular toric variety, *K* is the canonical line bundle of *M*. Using the moment map  $\Phi_K \colon M \to H^2(BT; \mathbb{R})$  associated with *K*, we define

$$\Phi'_L \coloneqq \Phi_L - \frac{1}{2} \Phi_K.$$

The moment maps  $\Phi_L$  and  $\Phi_K$  are equivariant, the *T*-action on the target  $H^2(BT;\mathbb{R})$  being trivial; so  $\Phi'_L$  induces a map

$$\bar{\Phi}'_I \colon M/T \to H^2(BT;\mathbb{R}).$$

The shifted affine hyperplanes  $\mathcal{F}'_L(\{i\})$ 's miss the lattice  $H^2(BT)$ . Since  $\partial(M/T) = \bigcup_i (M_i/T)$  and  $\bar{\Phi}'_L$  maps  $M_i/T$  to  $\mathcal{F}'_L(\{i\})$  for each i,  $\bar{\Phi}'_L$  induces a homomorphism

$$(\bar{\Phi}'_L)_* \colon H_{n-1}(\partial(M/T)) \to H_{n-1}(H^2(BT;\mathbb{R}) \setminus \{u\})$$

for each lattice point  $u \in H^2(BT)$ . We define

$$d'_L(u) :=$$
 the mapping degree of  $(\Phi'_L)_*$ 

where the orientation on  $H^2(BT; \mathbb{R})$  is determined by that on T. Our main theorem in this section is the following.

**Theorem 11.1.** Let M be a torus manifold. Then  $m_L = d'_L$  on  $H^2(BT)$ .

REMARK. This theorem was first established by Karshon-Tolman [22] when M is a compact non-singular toric variety, and then extended to Spin<sup>*c*</sup> manifolds with torus actions by Grossberg-Karshon [12] and to a unitary torus manifold by the second named author [27]. The family of torus manifolds contains these manifolds.

Let S(M) be the realization of the first barycentric subdivision of  $\Sigma(M)$  and let  $S(M)_i$  be the union of simplicies in S(M) which contain the vertex  $\{i\}$  as in Section 6. Since  $S(M)_I = \bigcap_{i \in I} S(M)_i$  is contractible for any non-empty set  $I \in \Sigma(M)$  and  $\partial(M/T) = \bigcup_{i=1}^d (M_i/T)$ , it follows from Lemma 6.2 that there is a continuous map

$$\rho_M : \partial(M/T) \to S(M)$$

sending  $\bigcap_{i \in I} (M_i/T)$  to  $S(M)_I$  for each  $I \in \Sigma(M)$ , and that such a map is unique up to homotopy preserving the stratifications, where the stratifications on  $\partial(M/T)$  and S(M) mean subspaces  $\bigcap_{i \in I} \partial(M_i/T)$  and  $S(M)_I$  indexed by elements I's in  $\Sigma(M)$ .

If the orientation on T or M is reversed, then  $[\partial(M/T)]$  and  $[\Delta(M)]$  will be multiplied by -1 simultaneously; so the following lemma makes sense.

**Lemma 11.2.**  $\rho_{M_*}([\partial(M/T)]) = [\Delta(M)].$ 

Proof. We prove the lemma by induction on the dimension  $n = \dim(M/T)$ . When n = 1, M is  $S^2$  with a nontrivial smooth  $S^1$ -action. In this case, it is not difficult to check the lemma, which we leave to the reader.

Assume that n > 1. Since a characteristic submanifold of  $M_i$  is a connected component of  $M_i \cap M_j$  for some j and such j is uniquely determined by the characteristic submanifold of  $M_i$ , there is a natural map  $\pi_i \colon \Sigma(M_i) \to \Sigma(M)_i$ . This map is an isomorphism if  $M_i \cap M_j$  is connected for any j, but otherwise it is only surjective. As we did in Lemma 6.1, we identify the realization of  $\Sigma(M)_i$  with  $\partial(S(M)_i)$ . One sees that

(11.3) 
$$\pi_{i*}([\Delta(M_i)]) = \sum_{i \in I \in \Sigma(M)^{(n)}} w(M)(I) \langle I \setminus \{i\} \rangle \in H_{n-2}(\partial(S(M)_i)) = H_{n-2}(\Sigma(M)_i).$$

Since  $M_i$  is itself a torus manifold, the spaces  $\partial(M_i/T)$  and  $S(M_i)$  have stratifications like for M, and hence we have a map  $\rho_{M_i} : \partial(M_i/T) \to S(M_i)$  preserving the stratifications. By the induction assumption

(11.4) 
$$\rho_{M_{i*}}([\partial(M_{i}/T)]) = [\Delta(M_{i})] \in H_{n-1}(S(M_{i})) = H_{n-1}(\Sigma(M_{i})).$$

On the other hand,  $\partial(S(M)_i)$  has a stratification induced from S(M) and each stratum is contractible. Since  $\rho_M$  restricted to  $\partial(M_i/T)$  is a map from  $\partial(M_i/T)$  to  $\partial(S(M)_i)$  preserving the stratifications and so is  $\pi_i \circ \rho_{M_i}$  as well, they are homotopic preserving the stratifications by Lemma 6.2. Therefore, we have the following commu-

tative diagram:

where the left horizontal maps are natural ones. Tracing the upper horizontal sequence from the left to the right,  $[\partial(M/T)] \in H_{n-1}(\partial(M/T))$  maps to  $\bigoplus_i [\partial(M_i/T)]$ , and down to  $\sum_{i \in I \in \Sigma(M)^{(n)}} w(M)(I) \langle I \setminus \{i\} \rangle \in \bigoplus_i H_{n-2}(\partial(S(M)_i))$  by (11.3) and (11.4), while  $[\Delta(M)] \in H_{n-1}(S(M))$  maps through the lower horizontal sequence to the same element as observed in Lemma 6.1. Since the horizontal sequences above are injective, the lemma follows.

Proof of Theorem 11.1. By Lemma 6.2 we have a map  $S(M) \to H^2(BT; \mathbb{R})$ associated with the multi-polytope  $\mathcal{P}'_L := (\Delta(M), \mathcal{F}'_L)$ . We denote the map by  $\Psi'_L$ . The composition  $\Psi'_L \circ \rho_M$  is a map from  $\partial(M/T)$  to  $H^2(BT; \mathbb{R})$  sending  $\bigcap_{i \in I} (M_i/T)$  to  $\bigcap_{i \in I} \mathcal{F}'_L(\{i\})$  for any  $I \in \Sigma(M)$ , and so is  $\overline{\Phi}'_L$  as well. Therefore,  $\Psi'_L \circ \rho_M$  and  $\overline{\Phi}'_L$  are homotopic preserving the stratifications by Lemma 6.2. It follows from Lemma 11.2 that

 $\begin{aligned} d'_{L}(u) &= \text{ the mapping degree of } (\bar{\Phi}'_{L})_{*} : H_{n-1}(\partial(M/T)) \to H_{n-1}(H^{2}(BT;\mathbb{R})\backslash\{u\}) \\ &= \text{ the mapping degree of } (\Psi'_{L} \circ \rho_{M})_{*} : H_{n-1}(\partial(M/T)) \to H_{n-1}(H^{2}(BT;\mathbb{R})\backslash\{u\}) \\ &= \text{ the mapping degree of } (\Psi'_{L})_{*} : H_{n-1}(S(M)) \to H_{n-1}(H^{2}(BT;\mathbb{R})\backslash\{u\}) \\ &= \text{WN}_{\mathcal{P}'_{L}}(u) = \text{DH}_{\mathcal{P}'_{L}}(u) = \text{DH}_{(\mathcal{P}_{L})_{*}}(u). \end{aligned}$ 

This together with Corollary 7.4 and the definition of  $m_L$  (i.e., (11.1) and (11.2)) proves the theorem.

#### 12. Torus orbifolds

The aim of this section is to give the definition of a torus orbifold and provide its basic properties for generalizing the results of Sections 10 and 11. We first recall basic definitions concerning orbifolds. We refer to [31], [24] or [10] for details. The reference [26] will be also useful; it deals with torus actions on symplectic orbifolds. If Mis an orbifold of dimension n, then there is a family  $\{(U_{\alpha}, V_{\alpha}, H_{\alpha}, p_{\alpha})\}$  of orbifold charts, where  $\{U_{\alpha}\}$  is an open covering of M,  $V_{\alpha}$  is an n-dimensional manifold,  $H_{\alpha}$ is a finite group acting on  $V_{\alpha}$  and  $p_{\alpha} : V_{\alpha} \to U_{\alpha}$  is a map which induces a homeomorphism from  $V_{\alpha}/H_{\alpha}$  onto  $U_{\alpha}$ . If  $U_{\alpha}$  and  $U_{\beta}$  intersect each other, then the charts  $(U_{\alpha}, V_{\alpha}, H_{\alpha}, p_{\alpha})$  and  $(U_{\beta}, V_{\beta}, H_{\beta}, p_{\beta})$  satisfy suitable compatibility conditions. Such a family  $\{(U_{\alpha}, V_{\alpha}, H_{\alpha}, p_{\alpha})\}$  is called an orbifold atlas. For any point x in M, there exists a special type of orbifold chart  $(U_x, V_x, H_x, p_x)$  with the property that  $p_x^{-1}(x)$  is a single point  $\tilde{x} \in V_x$ . The isomorphism class of the group  $H_x$  is uniquely determined by x and is called the isotropy group of x. The order of  $H_x$ , denoted by  $d_x$ , is called the multiplicity of the point x. Such an orbifold chart will be called a reduced orbifold chart. When M is connected, the minimum of the multiplicities is called the multiplicity of the orbifold M and is denoted by d(M). The set  $\{x \in M \mid d_x = d(M)\}$ is open and dense in M. It is a manifold. This set is called the principal stratum of the orbifold M. We have d(M) = 1 if and only if the actions of all the isotropy groups are effective.

A map  $f: M \to M'$  from an orbifold M into another orbifold M' is called smooth if, near every point x in M, there is a homomorphism  $\rho_{\alpha}: H_{\alpha} \to H'_{\alpha}$ and a  $\rho_{\alpha}$ -equivariant smooth map  $f_{\alpha}: V_{\alpha} \to V'_{\alpha}$  for suitable orbifold charts  $(U_{\alpha}, V_{\alpha}, H_{\alpha}, p_{\alpha})$  for M around x and  $(U'_{\alpha}, V'_{\alpha}, H'_{\alpha}, p'_{\alpha})$  for M' around f(x) satisfying the commutativity relation  $p'_{\alpha} \circ f_{\alpha} = f \circ p_{\alpha}$ . A subset M of an orbifold M' is called a suborbifold if, for each orbifold chart  $(U'_{\alpha}, V'_{\alpha}, H'_{\alpha}, p'_{\alpha})$  of  $M', V_{\alpha} = p'_{\alpha}^{-1}(M \cap U'_{\alpha})$ is an  $H'_{\alpha}$ -invariant submanifold of  $V'_{\alpha}$ . If this is the case, M becomes an orbifold with orbifold charts  $(U_{\alpha}, V_{\alpha}, H'_{\alpha}, p'_{\alpha})$  where  $U_{\alpha} = M \cap U'_{\alpha}$ , and the inclusion  $M \to M'$ becomes a smooth map. It may happen that d(M) > d(M') (M and M' are assumed connected). The integer  $d(M \mid M') = d(M)/d(M')$  will be called the relative multiplicity of the pair (M, M').

Orbifold vector bundles are also defined. Typical examples are the tangent bundle of an orbifold and the normal bundle of a suborbifold. An orbifold is orientable if its tangent bundle is orientable. If  $E \to M$  is an orbifold vector bundle over a connected orbifold, then the relative multiplicity of the orbifold vector bundle E is defined to be  $d(M \mid E)$  where M is identified with the zero-section and is considered as a suborbifold of E. If M is a suborbifold of M' and  $\nu$  is the normal bundle of M in M', then  $d(M \mid \nu)$  equals  $d(M \mid M')$ .

Let G be a Lie group. An action of G on an orbifold M is a smooth map  $\psi: G \times M \to M$  satisfying the usual rule of group action. Suppose that G is connected. If  $x \in M$  is a fixed point of the action, and  $(U_x, V_x, H_x, p_x)$  is a reduced orbifold chart around x such that  $U_x$  is invariant under the action of G, then there is a finite covering group  $\tilde{G}_x$  of G and an action of  $\tilde{G}_x$  on  $V_x$  which covers the action of G on  $U_x$ . If G is compact, the fixed point set of the action is a suborbifold.

Now let M be an oriented, closed orbifold of dimension 2n with an effective action of an *n*-dimensional torus T. A connected component of the fixed point set by a circle subgroup is a suborbifold. A suborbifold of this type which has codimension two and contains at least one fixed point of the T-action will be called a *characteristic* suborbifold. Let  $M_i$  be a characteristic suborbifold and  $x \in M_i$ . We take, as we may, a reduced orbifold chart  $(U_x, V_x, H_x, p_x)$  around x such that  $V_x$  is an open disk in  $\mathbb{R}^{2n}$ and the action of  $H_x$  on  $V_x$  is linear. We denote by the same symbol  $V_x$  the tangent space to  $V_x$  at the point  $\tilde{x} = p_x^{-1}(x)$ . Then the vector space  $V_x$  decomposes into a direct sum  $V_{ix} \oplus V_{ix}^{\perp}$  where  $V_{ix}^{\perp}$  is tangent to  $p_x^{-1}(U_x \cap M_i)$ , and the vector space  $V_{ix}$  represents the fiber direction of the normal bundle of  $M_i$  in M. The isotropy group  $H_x$  acts on  $V_{ix}$ .

**Lemma 12.1.** Let M be an oriented closed orbifold as above and  $M_i$  a characteristic suborbifold. Let  $S_i$  denote the circle subgroup which fixes the points of  $M_i$ . Then there exists a finite covering group  $\tilde{S}_i$  of  $S_i$  and a lifting of the action of  $S_i$  to the action of  $\tilde{S}_i$  on  $V_x$  for any point  $x \in M_i$ . The lifted action of  $\tilde{S}_i$  preserves  $V_{ix}$ .

Proof. To  $x \in M_i$  we correspond the degree of the minimal finite covering  $\tilde{S}_{ix}$  of  $S_i$  such that there is a lifting of the action to  $\tilde{S}_{ix}$ . The lifted action necessarily preserves  $V_{ix}$ . It is not difficult to see that the correspondence is locally constant. Since  $M_i$  is connected the correspondence must be constant.

Hereafter we denote by  $\rho_i : \tilde{S}_i \to S_i$  the minimal finite covering of  $S_i$  with the above property.  $\tilde{S}_i$  acts effectively on  $V_x$ .

An oriented, closed orbifold M of dimension 2n with an effective action of a torus T of dimension n with non-empty fixed point set  $M^T$  equipped with a preferred orientation of the normal bundle of each characteristic suborbifold will be called a *torus orbifold* if, for each  $M_i$  and at each point  $x \in M_i$ , the action of  $H_x$  preserves the orientation of each  $V_{ix}$ . Note that choosing an orientation of a characteristic submanifold is equivalent to choosing an orientation of its normal bundle. Thus a torus manifold is a torus orbifold in the above sense. Another example is a unitary torus orbifold is a torus orbifold such that  $V_{\alpha}$  is a unitary manifold, the action of  $H_{\alpha}$  preserves the unitary structure of  $V_{\alpha}$  for each orbifold chart  $(U_{\alpha}, V_{\alpha}, H_{\alpha}, p_{\alpha})$  and the action of T on M also preserves the unitary structure of  $V'_{\alpha}$ s.

Let M be a torus orbifold. The preferred orientation of the normal bundle  $\nu_i$  of  $M_i$  makes it a complex orbifold line bundle. Then there is a unique isomorphism  $\varphi_i \colon S^1 \to \tilde{S}_i$  such that  $\varphi(z)$  acts by the complex multiplication of z on each  $V_{ix}$ . We identify  $\tilde{S}_i$  with  $S^1$  via  $\varphi_i$ . The homomorphism  $\rho_i \colon S^1 = \tilde{S}_i \to T$  defines an element  $v_i \in \text{Hom}(S^1, T) = H_2(BT; \mathbb{Z})$ . We are now ready to define the multi-fan  $\Delta(M) = (\Sigma(M), C(M), w(M)^{\pm})$  associated with a torus orbifold M in an entirely similar way to the case of torus manifolds. Specifically

$$\Sigma(M) = \left\{ I \mid \left(\bigcap_{i \in I} M_i\right)^T \neq \emptyset \right\},\$$

and C(M)(I) is the cone in  $H_2(BT; \mathbb{R})$  with apex at 0 and spanned by  $\{v_i \mid i \in I\}$ . Furthermore  $w(M)^{\pm}(I) = \#\{x \in M_I \mid \epsilon_x = \pm 1\}$  for  $I \in \Sigma(M)^{(n)}$ , where  $\epsilon_x$  is defined to be the ratio of two orientations at x, one which is given by the orientation of M and the other by that of the oriented vector space  $V_x = \bigoplus_{i \in I} V_{ix}$ . We set  $\tilde{T}_I = \prod_{i \in I} \tilde{S}_i$  for  $I \in \Sigma(M)^{(k)}$  and  $\rho_I = \prod_{i \in I} \rho_i : \tilde{T}_I \to T$ . The image of  $\rho_I$  is denoted by  $T_I$ .  $\rho_I : \tilde{T}_I \to T_I$  is a finite covering.  $T_I$  fixes the points of  $M_I = \bigcap_{i \in I} M_i$ . If  $I \in \Sigma(M)^{(n)}$ , then  $T_I = T$ . Let x be a fixed point of the action of T on M. Then there is a unique  $I \in \Sigma(M)^{(n)}$  such that x belongs to  $M_I$ . The inclusion  $S^1 = \tilde{S}_i \to \tilde{T}_I$  defines an element  $\tilde{v}_i \in \text{Hom}(S^1, \tilde{T}_I) = H_2(B\tilde{T}_I; \mathbb{Z})$ , and we have  $\rho_{I*}(\tilde{v}_i) = v_i$ .  $V_x$  and  $V_{ix}$ ,  $i \in I$ , are complex  $\tilde{T}_I$ -modules, and the decomposition  $V_x = \bigoplus_{i \in I} V_{ix}$  is compatible with the action of  $\tilde{T}_I$ . The effectiveness of the T-action on M implies that  $\tilde{T}_I$  effectively acts on  $V_x$ ; equivalently, it implies that  $\{\tilde{v}_i \mid i \in I\}$  is a basis of  $H_2(B\tilde{T}_I;\mathbb{Z})$ . Since  $\rho_{I*}: H_2(B\tilde{T}_I;\mathbb{Z}) \to H_2(BT;\mathbb{Z})$  is injective, the  $v_i$ ,  $i \in I$ , are linearly independent in  $H_2(BT;\mathbb{R})$ .

**Lemma 12.2.**  $\Delta(M)$  is a complete multi-fan.

Proof. The argument is almost similar to the case of torus manifolds. One has only to observe that the characteristic suborbifolds and their intersections are torus orbifolds and a 2-dimensional torus orbifold is topologically a 2-sphere acted on by a circle group with exactly two fixed points.  $\Box$ 

**Lemma 12.3.** Suppose d(M) = 1. Let  $I \in \Sigma(M)^{(k)}$ , and let x be a point in the principal stratum (as an orbifold) of  $M_I$ . Then the isotropy group  $H_x$  of x is isomorphic to the kernel of  $\rho_I : \tilde{T}_I \to T$ .

Proof. Let  $(U_x, V_x, H_x, p_x)$  be an orbifold chart around x. We may regard  $V_x$  as an n dimensional  $\tilde{T}_I$ -module as before. As such,  $V_x$  is decomposed as a direct sum of  $\tilde{T}_I$ -modules

$$V_x = \left(\bigoplus_{i \in I} V_{ix}\right) \oplus V'$$

where V' is projected into  $M_I$  by  $p_x$ .  $\tilde{T}_I = \prod_{i \in I} \tilde{S}_i$  can be regarded as embedded in the general linear group of  $\bigoplus_{i \in I} V_{ix}$ . Since  $H_x$  acts on each  $V_{ix}$  preserving its orientation, there is a homomorphism  $H_x \to \tilde{T}_I$ . The action of  $H_x$  on V' is trivial. Moreover the action of  $H_x$  on  $V_x$  is effective because d(M) = 1. It follows that the homomorphism above embeds  $H_x$  into  $\tilde{T}_I$ . Since the kernel of  $\rho_I$  is equal to the intersection of  $\tilde{T}_I$  with the image of  $H_x$ , it is isomorphic to  $H_x$ .

It is known that a closed oriented orbifold M of dimension n has the fundamental class  $[M] \in H_n(M;\mathbb{Z})$ , and that the Poincaré duality holds, i.e., the operation  $\vartheta = [M] \cap : H^q(M;\mathbb{Q}) \to H_{n-q}(M;\mathbb{Q})$  is an isomorphism. If  $f: M \to M'$ is a smooth map from an oriented close orbifold M to another such M', then the Gysin homomorphism  $f_!: H^q(M;\mathbb{Q}) \to H^{q+n-n'}(M';\mathbb{Q})$  is defined to be the compostion  $\vartheta^{-1} \circ f_* \circ \vartheta$ , where n' is the dimension of M'. If a compact Lie group G acts on M and M', and f is equivariant, then the equivariant Gysin homomorphism  $f_!: H^q_G(M; \mathbb{Q}) \to H^{q+n-n'}_G(M'; \mathbb{Q})$  is also defined.

Henceforth M will be a torus orbifold. For each  $i \in \Sigma(M)^{(1)}$ , we set

$$\xi_i = (f_i)_!(1) \in H^2_T(M; \mathbb{Q}),$$

where  $f_i: M_i \to M$  is the inclusion.

**Lemma 12.4.** Let  $c_1^T(\nu_i)$  be the equivariant first Chern class of the normal bundle  $\nu_i$ . Then we have

$$c_1^T(\nu_i) = f_i^*(\xi_i).$$

Proof. We may assume that d(M) = 1. Take an equivariant Thom form  $\phi$  for the equivariant orbifold bundle  $\nu_i$  (we refer to [3] for Thom form and Chern form, cf. also [10]). Let x be a point in the principal stratum of  $M_i$ , and  $(U_x, V_x, H_x, p_x)$  a reduced orbifold chart around x. The restriction of  $\phi$  to  $V_x$  is invariant under the action of  $H_x$  and its support is contained in a tubular neighborhood  $W_i$  of  $V_i = p_x^{-1}(U_i)$ , where  $U_i = U_x \cap M_i$ . Moreover, with respect to the fibering  $\tilde{\pi}_i : W_i \to V_i$ , we have  $|H_x|^{-1}(\tilde{\pi}_i)_*(\phi) = 1$ , where  $(\tilde{\pi}_i)_*$  is the integration along the fiber of  $\tilde{\pi}_i$ . Note that the fiber is  $V_{ix}$ , and that the action of  $H_x$  preserves the orientation of  $V_{ix}$ . The equivariant Chern class  $c_1^T(\nu_i)$  is the restriction to  $M_i$  of the cohomology class  $[\phi]$  of  $\phi$ . Here  $[\phi]$ is considered as a relative class in  $H_T^2(W, W \setminus M_i; \mathbb{R})$  where W is a tubular neighborhood of  $M_i$ .

On the other hand,  $\xi_i$  is the restriction of a cohomology class  $\psi \in H^2_T(W, W \setminus M_i; \mathbb{R})$  such that

$$\pi_*(\psi) = 1 \in H^0_T(W; \mathbb{R}) = H^0_T(M_i; \mathbb{R}),$$

where  $\pi: W \to M_i$  denotes the projection of the fibration. Note that the fiber of  $\pi$  is  $U_{ix} = V_{ix}/H_x$ , where  $H_x$  acts effectively on  $V_{ix}$ . We have

$$\pi_*([\phi]) = |H_{\chi}|^{-1}(\tilde{\pi}_i)_*([\phi]) = 1 = \pi_*(\psi).$$

But  $\pi_*$  is an isomorphism (Thom isomorphism). Hence we have  $[\phi] = \psi$ , and consequently

$$c_1^T(\nu_i) = [\phi] \mid M_i = \psi \mid M_i = f_i^*(\xi_i).$$

We noticed that, for  $I \in \Sigma(M)^{(n)}$ ,  $\{v_i \mid i \in I\}$  was a basis of  $H_2(BT; \mathbb{R})$ . Let  $\{u_i^I\}$  be the dual basis in  $H^2(BT; \mathbb{R})$ . This can be interpreted in the following way. Let  $\{\tilde{u}_i \mid i \in I\}$  be the basis of  $H^2(B\tilde{T}_I; \mathbb{Z})$  dual to  $\{\tilde{v}_i \mid i \in I\}$ . We have  $\rho_I^*(u_i^I) = \tilde{u}_i$ , since  $\rho_{I*}(\tilde{v}_i) = v_i$ . We identify  $H^2(B\tilde{T}_I; \mathbb{R})$  with  $H^2(BT; \mathbb{R})$  by the isomorphism  $\rho_I^*$ . Then  $H^2(B\tilde{T}_I;\mathbb{Z})$  can be considered as embedded in  $H^2(BT;\mathbb{R})$ . With this convention we have  $u_i^I = \tilde{u}_i$ .

Let  $x \in M^T$  be a fixed point of the *T*-action. In the sequel we identify  $H^2_T(x;\mathbb{R})$  with  $H^2(BT;\mathbb{R})$ .

**Lemma 12.5.** Let  $I \in \Sigma(M)^{(n)}$  and  $x \in M_I$ . Then  $\xi_i \mid x = u_i^I \in H^2(BT; \mathbb{R})$  for  $i \in I$ . If  $j \notin I$ , then  $\xi_j \mid x = 0$ .

Proof. By Lemma 12.4 we have

$$\xi_i \mid x = c_1^T(\nu_i \mid x).$$

But  $\nu_i \mid x$  viewed as  $\tilde{T}_I$ -module is  $V_{ix}$ . It follows that  $c_1^{\tilde{T}_I}(\nu_i \mid x) = \tilde{u}_i$ . Hence

$$c_1^T(\nu_i \mid x) = u_i^I$$
.

If  $j \notin I$ , then  $x \notin M_j$ . Therefore  $\xi_j | x = 0$ .

If we consider  $u_i^I = \tilde{u}_i$  as an element of  $\operatorname{Hom}(\tilde{T}_I, S^1) = H^2(B\tilde{T}_I; \mathbb{Z})$ , then Lemmas 12.5 and 12.6 imply that  $u_i^I$  is nothing but the  $\tilde{T}_I$ -module  $V_{ix}$ . The following Lemma describes the algebra structure of  $H_T^*(M; \mathbb{R})$  over  $H^*(BT; \mathbb{R})$  modulo  $H^*(BT; \mathbb{R})$ -torsion as in the case of torus manifolds (Lemma 9.3).

**Lemma 12.6.** The following equality holds for any  $u \in H^2(BT; \mathbb{R})$ :

$$\pi^*(u) = \sum_{i \in \Sigma(M)^{(1)}} \langle u, v_i \rangle \xi_i \quad modulo \ H^*(BT; \mathbb{R}) \text{-torsion.}$$

Proof. Let  $x \in M_I \subset M^T$  be a fixed point of the *T*-action. We restrict both sides of the equality in Lemma 12.6 to *x*. On the left hand side we get *u*. On the right hand side the result is

$$\sum_{i\in I} \langle u, v_i \rangle u_i^I$$

by virtue of Lemma 12.5. But this is equal to u by the definition of the  $u_i^I$ . Thus both sides coincide after the restriction to each  $x \in M^T$ . Since the restriction homomorphism  $\pi^* \colon H_T^*(M;\mathbb{R}) \to H_T^*(M^T;\mathbb{R})$  is injective modulo  $H^*(BT;\mathbb{R})$ -torsion, the equality is confirmed.

REMARK. The equality in Lemma 12.6 characterizes the vectors  $v_i$  in terms of the  $\xi_i$  as in Lemma 9.3.

We set  $N = H_2(BT; \mathbb{Z})$  and define  $N_I$  for  $I \in \Sigma(M)^{(n)}$  to be the lattice generated

54

by the  $v_i$ ,  $i \in I$ .

**Lemma 12.7.** Assume that d(M) = 1. Let  $x \in M_I$  with  $I \in \Sigma(M)^{(n)}$ . Then  $H_x$  is isomorphic to Ker  $\rho_I$ . Moreover Ker  $\rho_I$  is isomorphic to  $N/N_I$ .

Proof. We have already shown that  $H_x$  is isomorphic to the kernel of  $\rho_I$  in Lemma 12.3. For the second part it suffices to note that N and  $N_I$  can be identified with the fundamental group of T and  $\tilde{T}_I$ . Therefore the kernel of  $\rho_I$  is isomorphic to  $N/N_I$ .

REMARK. Hereafter we identify  $H_x$  and  $N/N_I$  with  $\operatorname{Ker} \rho_I \subset \tilde{T}_I$  through the isomorphisms given in Lemma 12.7. We put  $\chi_I(u, v) = \exp(2\pi\sqrt{-1}\langle u, v\rangle)$  for  $u \in H^2(B\tilde{T}_I;\mathbb{Z})$  and  $v \in H_2(BT;\mathbb{R})$ . If u is fixed, then the value  $\chi_I(u, v)$  depends only on the equivalence class of v modulo  $N_I$ . Hence, if we identify  $\tilde{S}_i$  with  $S^1$  via  $\varphi_i$  as before and  $\tilde{T}_I$  with  $\prod_{i \in I} S^1$  via  $\prod_{i \in I} \varphi_i$ , then the map  $\exp: H_2(BT;\mathbb{R}) \to \tilde{T}_I$  defined by  $\exp(v) = \prod_{i \in I} \exp(2\pi\sqrt{-1}\langle u_i^I, v \rangle)$  is a universal covering map and its kernel is  $N_I$ . It induces an isomorphism from  $H_x = N/N_I$  onto  $\operatorname{Ker} \rho_I$ . We shall write  $\chi_I(u, g)$  instead of  $\chi_I(u, v)$  for  $g = \exp(v) \in \tilde{T}_I$  as in Section 7. Let V be a one dimensional  $\tilde{T}_I$ -module. It defines an element  $u \in \operatorname{Hom}(\tilde{T}_I, S^1) = H^2(B\tilde{T}_I;\mathbb{Z})$ . Then the action of  $g \in \tilde{T}_I$  on V is given by the complex multiplication by  $\chi_I(u, g)$ .

Suppose that M is a unitary torus orbifold such that d(M) = 1. Let L be a T-invariant complex line bundle over M. By using a hermitian connection of M and a hermitian connection of L, a Dirac operator twisted by L is defined as in the case of torus manifolds. Its index is a T-module. It is called the equivariant Riemann-Roch number with coefficient in L, and is denoted by  $RR^T(M, L) \in R(T)$ . It can be expressed by the fixed point formula due to Vergne [33]; cf. also [10]. The formula is particularly simple when all the fixed points are isolated. It is convenient to write down the image of  $RR^T(M, L)$  by ch:  $R(T) \to H^{**}(BT; \mathbb{R})$ ; the result is

**Lemma 12.8.** Let  $\xi = c_1^T(L)$  be the equivariant Chern class of L. Then

$$\operatorname{ch}\left(RR^{T}(M,L)\right) = \sum_{x \in M^{T}} \frac{\epsilon_{x} e^{\xi \mid x}}{\mid H_{x} \mid} \sum_{g \in H_{x}} \frac{\chi_{I_{x}}(\xi \mid x,g)}{\prod_{i \in I_{x}} (1 - \chi_{I_{x}}(u_{i}^{I_{x}},g)^{-1}e^{-u_{i}^{I_{x}}})}$$

where  $I_x \in \Sigma(M)^{(n)}$  is such that  $x \in M_{I_x}$ .

It can be shown that, if x and y both lie in the same  $M_I$ , then  $\xi \mid x = \xi \mid y$  for  $\xi = c_1^T(L)$ . The proof is same as in the case of torus manifolds as was given in [27]. We shall write  $u_I = c_1^T(L) \mid x$  for  $x \in M_I$ . Taking Remark below Lemma 12.7 in account, we get

Proposition 12.9.

$$\operatorname{ch}\left(RR^{T}(M,L)\right) = \sum_{I \in \Sigma(M)^{(n)}} \frac{w(M)(I)e^{u_{I}}}{|N/N_{I}|} \sum_{g \in N/N_{I}} \frac{\chi_{I}(u_{I},g)}{\prod_{i \in I}(1 - \chi_{I}(u_{i}^{I},g)^{-1}e^{-u_{i}^{I}})} \ .$$

Since ch:  $R(T) \to H^{**}(BT; \mathbb{R})$  is injective, the formula in Proposition 12.9 characterizes  $RR^{T}(M, L)$ . Using the notation in Section 7, we obtain

# Corollary 12.10.

$$RR^{T}(M,L) = \sum_{I \in \Sigma(M)^{(n)}} \frac{w(M)(I)t^{u_{I}}}{|N/N_{I}|} \sum_{g \in N/N_{I}} \frac{\chi_{I}(u_{I},g)}{\prod_{i \in I}(1 - \chi_{I}(u_{i}^{I},g)^{-1}t^{-u_{i}^{I}})}$$

When  $u_I = c_1^T(L) \mid x, x \in M_I$ , lies in  $N^* = H^2(BT; \mathbb{Z})$ , then  $\chi_I(u_I, g) = 1$  for all  $g \in N/N_I$ . Therefore, if  $u_I \in N^*$  for all  $I \in \Sigma(M)^{(n)}$ , then

$$RR^{T}(M,L) = \sum_{I \in \Sigma(M)^{(n)}} \frac{w(M)(I)t^{u_{I}}}{|N/N_{I}|} \sum_{g \in N/N_{I}} \frac{1}{\prod_{i \in I} (1 - \chi_{I}(u_{i}^{I},g)^{-1}t^{-u_{i}^{I}})}$$

By observing that  $g \mapsto \chi_I(u, g)$  is a character of  $N/N_I$  for any  $u \in H^2(B\tilde{T}_I, \mathbb{Z}) = \text{Hom}(\tilde{T}_I, S^1)$ , the formula above can be rewritten in the following form:

(12.1) 
$$RR^{T}(M,L) = \sum_{I \in \Sigma(M)^{(n)}} \frac{w(M)(I)t^{u_{I}}}{|N/N_{I}|} \sum_{g \in N/N_{I}} \frac{1}{\prod_{i \in I} (1 - \chi_{I}(u_{i}^{I},g)t^{-u_{i}^{I}})}$$

The right hand side of this formula (12.1) appeared in Corollary 7.4. There, it was related to a lattice multi-polytope  $\mathcal{P}$ , in which  $u_I$  is contained in  $N^*$  for all  $I \in \Sigma^{(n)}$ , and the Duistermaat-Heckman function  $DH_{\mathcal{P}_+}$ . Suppose that  $c_1^T(L)$  is of the form  $c_1^T(L) = \sum_{i \in \Sigma(M)^{(1)}} c_i \xi_i \in H^2_T(M; \mathbb{R})$ . Then the above multi-polytope  $\mathcal{P}$  is nothing but the one whose first Chern class is  $c_1(\mathcal{P}) = \sum c_i x_i$  as in Section 8. Note that  $\mathcal{P}$  is not always a lattice multi-polytope in this case.

REMARK. Corollary 7.4 shows that the right hand side of the formula (12.1) depends only on  $\Delta(M)$  and  $\mathcal{P}$ ; namely, it does not depend on the choice of generating vectors  $v_i \in H^2(BT; \mathbb{R})$  in so far as they lie in  $N = H^2(BT; \mathbb{Z})$  and  $\{u_i^I \mid i \in I\}$  is interpreted as the dual basis of  $\{v_i \mid i \in I\}$ .

When M is a torus manifold, the Duistermaat-Heckman function has a geometric meaning coming from the moment map of the line bundle L as was explained in Section 11. There the role of the winding number was also explained. These notions are generalized to the case of torus orbifolds and similar results hold in this case too. The details can be worked out without much alteration and are left to the reader.

56

#### THEORY OF MULTI-FANS

The  $T_y$ -genus of a torus orbifold is also defined by using the fixed point formula due to Vergne in a similar way as in Section 10, and the analogue of Theorem 10.1 holds. the details are left to the reader.

#### 13. Realizing multi-fans by torus orbifolds

In the previous section, we associated a complete simplicial multi-fan of dimension *n* with a torus orbifold of dimension 2*n*. In this section, we consider the converse problem. If a multi-fan  $\Delta$  is associated with a torus orbifold *M*, then we say that  $\Delta$ is (geometrically) realized by *M*, or *M* realizes  $\Delta$ .

We recall how the multi-fan of M changes when the orientations on M or  $M_i$ are reversed. If the orientation of M is unchanged but that of  $M_i$  is reversed, then the orientation of the normal bundle of  $M_i$  is reversed and, hence, 1-dimensional cone C(i) turns into the cone -C(i), and the pair  $(w(M)^+(I), w(M)^-(I))$  turns into  $(w(M)^-(I), w(M)^+(I))$  for  $I \in \Sigma(M)^{(n)}$  containing i while others remain unchanged. If the orientations of M and of all the  $M_i$ 's are reversed, then all the cones C(i)'s remain unchanged but  $(w(M)^+(I), w(M)^-(I))$  turns into  $(w(M)^-(I), w(M)^+(I))$  for any  $I \in \Sigma(M)^{(n)}$  so that w(M)(I) turns into -w(M)(I) for any  $I \in \Sigma(M)^{(n)}$ . The torus orbifold M with the reversed orientations of M and all the  $M_i$ 's will be denoted by -M.

The underlying space of a torus orbifold of dimension 2 is  $S^2$  with the standard  $S^1$ -action. In this case, there are two characteristic submanifolds. They are  $S^1$ -fixed points. Taking orientations on  $S^2$  and its characteristic submanifolds into account, we easily have the following theorem.

**Theorem 13.1.** A complete simplicial multi-fan  $\Delta = (\Sigma, C, w^{\pm})$  of dimension 1 is geometrically realized if and only if  $\Sigma$  is isomorphic to the argumented simplicial set obtained from the boundary of a 1-simplex and  $\{w^+(I), w^-(I)\} = \{1, 0\}$  as a set for  $I \in \Sigma^{(1)}$ .

The analysis of a torus orbifold of dimension 4 is more complicated. In this case, each characteristic suborbifold is homeomorphic to  $S^2$  and has two fixed points. Therefore, if two of the characteristic suborbifolds intersect, then they intersect at one point or two points, and if they intersect at two points, then they do not intersect at any other characteristic suborbifolds. We also note that a *T*-fixed point is an intersection of two characteristic suborbifolds. These facts imply the "only if" part in the following theorem. We will prove the "if" part later.

**Theorem 13.2.** A complete simplicial multi-fan  $\Delta = (\Sigma, C, w^{\pm})$  of dimension 2 is geometrically realized if and only if the following two conditions are satisfied for each  $I \in \Sigma^{(2)}$ : (1)  $\{w^+(I), w^-(I)\} = \{1, 0\}$  or  $\{1, 1\}$ , (2) when  $\{w^+(I), w^-(I)\} = \{1, 0\}$ , there are exactly two elements, say I' and I'', in  $\Sigma^{(2)}$  such that  $I \cap I'$  and  $I \cap I''$  are in  $\Sigma^{(1)}$  and  $I \cap I' \cap I'' = \emptyset$ , and when  $\{w^+(I), w^-(I)\} = \{1, 1\}$ , there is no element  $I' \in \Sigma^{(2)}$  such that  $I \cap I' \in \Sigma^{(1)}$ .

In contrast to the low dimensional cases above, we have

**Theorem 13.3.** Any complete simplicial multi-fan of dimension  $\geq 3$  is geometrically realized.

In the following  $\Delta = (\Sigma, C, w^{\pm})$  will be a complete simplicial multi-fan of dimension  $n \ge 2$  unless otherwise stated. Here is an outline of how to realize  $\Delta$  by a torus orbifold. We choose and fix a generic (rational) 1-dimensional cone in  $N_{\mathbb{R}}$ , and decompose  $\Delta$  using it into a number of what we call *minimal* multi-fans. Minimal multi-fans can essentially be realized by weighted projective spaces. We paste them together by performing equivariant connected sum along characteristic suborbifolds and at *T*-fixed points to obtain a desired torus orbifold realizing the given  $\Delta$ .

Equivariant connected sum is performed through two isomorphic orbifold charts. In this way attention should be paid to orbifold structures. So we make a remark on orbifold structures at this point. There are many choices of an orbifold structure on M(e.g.  $S^2$  with the standard  $S^1$ -action admits infinitely many orbifold structures), but the associated multi-fan does not depend on the choice of an orbifold structure. In fact, the circle subgroup  $S_i$  determined by the vector  $v_i$  in the previous section is the one which fixes points in the characteristic suborbifold  $M_i$ , so the line generated by  $v_i$  is independent of the orbifold structure. Moreover the direction of  $v_i$  is determined by the choice of orientations on M and  $M_i$ , so the cone spanned by  $v_i$  is independent of the orbifold structure. What depends on the orbifold structure is the length of  $v_i$  which is equal to the degree of the covering map  $\tilde{S}_i \to S_i$ . In this way the vectors  $v_i$  reflect the orbifold structure related to the torus action. We shall call the vector  $v_i$  the edge vector of the 1-dimensional cone C(i).

We shall use two types of equivariant connected sum; one is at *T*-fixed points and the other is along characteristic suborbifolds. Let us explain the former first. Suppose that torus orbifolds *M* and *M'* with d(M) = d(M') have *T*-fixed points *q* and *q'* respectively such that the *n*-dimensional cones and the edge vectors corresponding to them are the same and the signs  $\epsilon_q$  and  $\epsilon_{q'}$  at *q* and *q'* are opposites. Then there are a finite covering  $\tilde{T}$  of *T*, a finite subgroup *H* of  $\tilde{T}$  and orbifold charts (U, V, H, p)and (U', V, H, p') around *q* and *q'* respectively such that *V* is an invariant open disk centered at the origin in a  $\tilde{T}$ -module. In particular a diffeomorphism (in the sense of orbifold) *f* from the closure of *U* onto that of *U'* is induced. Moreover *f* sends characteristic suborbifolds that contain *q* onto characteristic suborbifolds that contain *q'*. It should be noticed that *f* is orientation reversing on *U* and on all the characteristic suborbifolds. We remove *U* and *U'* from *M* and *M'* respectively and glue their boundaries through the diffeomorphism f restricted to the boundaries. The resulting space is a torus orbifold with the orientations compatible with the torus orbifolds M and M'.

Let us explain the equivariant connected sum along characteristic suborbifolds. For the sake of simplicity we assume that d(M) = 1. Let  $M_i$  be a characteristic suborbifold, p a point in the principal stratum of the orbifold  $M_i$ . We may assume that the isotropy subgroup at p of the T-action is the circle group  $S_i$ . Let  $\tilde{S}_i$  be the covering group of  $S_i$  corresponding to the edge vector  $v_i$  as introduced in the previous section. Denote by  $V_i$  the standard complex 1-dimensional  $\tilde{S}_i$ -module and by  $D(V_i)$  the unit disk of  $V_i$ . Then it follows from the Slice Theorem and Lemma 12.3 that the T-orbit of p has a closed invariant tubular neighborhood  $\bar{U}_i$  in M equivariantly diffeomorphic to

(13.1) 
$$(T \times_{\tilde{S}_i} D(V_i)) \times D^{n-1}$$

where  $T \times_{\tilde{S}_i} D(V_i)$  denotes the orbit space of  $T \times D(V_i)$  by the  $\tilde{S}_i$ -action defined by  $s(t, x) = (t\rho_i(s)^{-1}, sx)$  for  $s \in \tilde{S}_i$ ,  $t \in T$  and  $x \in D(V_i)$ .

Suppose that there are characteristic suborbifolds  $M_i$  and  $M'_{i'}$  of torus orbifolds M and M' with d(M) = d(M') = 1 respectively such that the corresponding edge vectors coincide. Then the corresponding circle subgroups  $\tilde{S}_i$  and  $\tilde{S}'_{i'}$  agree and there is an equivariant diffeomorpism between  $\bar{U}_i$  and  $\bar{U}'_{i'}$  reversing the orientations induced from M,  $M_i$ , M' and  $M'_{i'}$  because both  $\bar{U}_i$  and  $\bar{U}'_{i'}$  are equivariantly diffeomorphic to the space in (13.1) and  $D^{n-1}$  ( $n \ge 2$ ) has an orientation reversing self-diffeomorphism. We remove the interior of  $\bar{U}_i$  and  $\bar{U}'_{i'}$  from M and M' and paste them together along the boundaries of  $\bar{U}_i$  and  $\bar{U}'_{i'}$  through the orientation reversing equivariant diffeomorphism restricted to the boundaries, producing a new torus orbifold, say M''. We call this procedure the equivariant connected sum of M and M' along  $M_i$  and  $M'_{i'}$ . The codimension of the principal orbits in  $M_i$  and  $M'_{i'}$  is n-1, so when  $n \ge 3$ ,  $M_i$  and  $M'_{i'}$  are pasted together to become one characteristic suborbifold in M'' and  $\Delta(M'')$ is obtained from  $\Delta(M)$  and  $\Delta(M')$  by identifying *i* with *i'*. However, when n = 2, the characteristic suborbifolds  $M_i$  and  $M'_{i'}$  are  $S^2$  and the principal orbits in them are circles; so the orbits separate  $M_i$  and  $M'_{i'}$  into two connected components respectively and hence two characteristic suborbifolds of M'' are produced.

Let  $I \in \Sigma(M)^{(n)}$  and  $I' \in \Sigma(M')^{(n)}$  be such that C(M)(I) = C(M)(I'). Suppose that the corresponding edge vectors are the same for I and I'. Then one can make equivariant connected sum of M and M' along each pair of characteristic suborbifolds  $M_i$  and  $M'_{i'}$  such that C(M)(i) = C(M')(i') for  $i \in I$  and  $i' \in I'$ , and then elements in I and I' will be identified in pairs in the multi-fan of the resulting torus orbifold and the weights  $w^{\pm}$  on the identified n-dimensional cone is the sum of those at I and I'.

We say that  $\Delta$  is *connected* if  $\Sigma$  is connected. According to the decomposition of  $\Sigma$  into connected components, the multi-fan  $\Delta$  decomposes into connected multi-fans which are again complete simplicial and of dimension *n*.

**Lemma 13.4.** Suppose  $n \ge 2$ . Then the multi-fan  $\Delta$  is geometrically realized if all connected components of  $\Delta$  are geometrically realized.

Proof. Let M be a torus orbifold of dimension 2n and let p be a point in the principal stratum of M. We may suppose that d(M) = 1. A closed tubular neighborhood  $\overline{U}$  of the orbit of p is equivariantly diffeomorphic to  $T \times D^n$  and the complement of  $\overline{U}$  is connected because M is connected and the orbit has codimension  $n \ge 2$ .

Let M' be another torus orbifold of dimension 2n with d(M') = 1, and let  $\overline{U'}$ be a closed subset in M' corresponding to  $\overline{U}$  in M. Since both  $\overline{U}$  and  $\overline{U'}$  are equivariantly diffeomorphic to  $T \times D^n$  and  $D^n$  has an orientation reversing diffeomorphim, there is an orientation reversing equivariant diffeomorphism between  $\overline{U}$  and  $\overline{U'}$ . We remove the interior of  $\overline{U}$  and  $\overline{U'}$  from M and M' respectively and glue their boundaries through the diffeomorphism restricted to the boundaries and obtain a new torus orbifold M''. The multi-fan  $\Delta(M'')$  is the disjoint union of  $\Delta(M)$  and  $\Delta(M')$ . (Precisely speaking,  $\Sigma(M'')$  is the disjoint union of  $\Sigma(M)$  and  $\Sigma(M')$  with the empty sets in them identified.)

If all connected components of  $\Delta$  are geometrically realized, then we connect torus orbifolds that realize the connected components of  $\Delta$  by the above method. Then the resulting torus orbifold realizes  $\Delta$ .

As is shown in the proof of Lemma 13.4, whenever we have more than two torus orbifolds of dimension  $n \ge 2$ , we can connect them and the multi-fan of the resulting torus orbifold is the disjoint union of the multi-fans of the torus orbifolds we had.

DEFINITION. We say that a complete simplicial multi-fan  $\Delta = (\Sigma, C, w^{\pm})$  of dimension *n* is *minimal* if

(1)  $\Sigma$  is isomorphic to the argumented simplicial set obtained from the boundary of an *n*-simplex, and

(2) the set  $\{w^+(I), w^-(I)\}$  is independent of  $I \in \Sigma^{(n)}$ .

Although the set  $\{w^+(I), w^-(I)\}$  is independent of I for a minimal multi-fan  $\Delta$ , the pair  $(w^+(I), w^-(I))$  may not be independent of  $I \in \Sigma^{(n)}$ . But one can convert  $\Delta$ into another minimal multi-fan  $\overline{\Delta} = (\Sigma, \overline{C}, \overline{w}^{\pm})$  such that the pair  $(\overline{w}^+(I), \overline{w}^-(I))$  is independent of I. The definition of  $\overline{\Delta}$  is as follows. Since  $\Delta$  is of dimension n and the cardinality of  $\Sigma^{(1)}$  is n + 1, there is a relation  $\sum_{i \in \Sigma^{(1)}} b_i v_i = 0$  among the edge vectors  $v_i$  with non-zero real numbers  $b_i$ . We then define

$$ar{C}(i) := egin{cases} C(i) & ext{if } b_i > 0, \\ -C(i) & ext{if } b_i < 0, \end{cases}$$

and define  $\bar{C}(K)$  for  $K \in \Sigma^{(m)}$  with  $m \ge 2$  to be the cone spanned by  $\bar{C}(k)$ 's for

 $k \in K$ . We also define

$$(\bar{w}^{+}(I), \bar{w}^{-}(I)) \coloneqq \begin{cases} (w^{+}(I), w^{-}(I)) & \text{if } \sharp\{i \in I \mid b_{i} < 0\} \text{ is even,} \\ (w^{-}(I), w^{+}(I)) & \text{if } \sharp\{i \in I \mid b_{i} < 0\} \text{ is odd,} \end{cases}$$

for  $I \in \Sigma^{(n)}$ .

**Lemma 13.5.**  $\overline{\Delta}$  is minimal and satisfies the following two conditions: (1) the n-dimensional cones  $\overline{C}(I)$  ( $I \in \Sigma^{(n)}$ ) do not overlap and their union covers the entire space  $N_{\mathbb{R}}$ , and (2)  $d = \sum_{i=1}^{n} \overline{C}(I)$  is in the space  $L \in \Sigma^{(n)}$ 

(2) the pair  $(\bar{w}^+(I), \bar{w}^-(I))$  is independent of  $I \in \Sigma^{(n)}$ . Moreover  $\Delta$  is geometrically realized if and only if so is  $\bar{\Delta}$ .

Proof. Let  $\bar{v}_i$  be a non-zero vector in the cone  $\bar{C}(i)$ . One may choose it to be  $v_i$  if  $b_i > 0$  and  $-v_i$  if  $b_i < 0$ . Then one has a relation  $\sum_{i \in \Sigma^{(1)}} \bar{b}_i \bar{v}_i = 0$  with positive numbers  $\bar{b}_i$ . This implies the statement (1) in the lemma.

We shall prove the statement (2) in the lemma. Let  $J \in \Sigma^{(n-1)}$ . Since the cardinality of  $\Sigma^{(1)}$  is n + 1, there are exactly two elements  $i, i' \in \Sigma^{(1)}$  not contained in J, and  $J \cup \{i\}$  and  $J \cup \{i'\}$  are in  $\Sigma^{(n)}$ , in other words, the (n-1)-dimensional cone C(J)is a facet of only two *n*-dimensional cones  $C(J \cup \{i\})$  and  $C(J \cup \{i'\})$ . We project them on  $N_{\mathbb{R}}^{C(J)}$  (the quotient space of  $N_{\mathbb{R}}$  by the subspace generated by C(J)). Then the vectors projected from  $v_i$  and  $v_{i'}$  are toward opposite directions if and only if  $b_i b_{i'} >$ 0. It follows from the completeness of  $\Delta$  that  $w(J \cup \{i\}) = \operatorname{sign}(b_i b_{i'})w(J \cup \{i'\})$ . This together with the definition of  $\bar{w}^{\pm}$  shows that  $\bar{w}(J \cup \{i\}) = \bar{w}(J \cup \{i'\})$ . Since  $J \in \Sigma^{(n-1)}$  is arbitrary, this proves the statement (2). It also proves the completeness of  $\bar{\Delta}$ , so that  $\bar{\Delta}$  is minimal.

The procedure from  $\Delta$  to  $\overline{\Delta}$  corresponds to reversing orientations on characteristic suborbifolds  $M_i$  with  $b_i < 0$ , so the latter statement in the lemma is obvious.

**Lemma 13.6.** Let  $\Delta$  be a minimal multi-fan of dimension  $n \geq 2$ . If  $n \geq 3$ , then  $\Delta$  is geometrically realized. If n = 2, then  $\Delta$  is geometrically realized if (and only if)  $\{w^+(I), w^-(I)\} = \{1, 0\}$  for any  $I \in \Sigma^{(2)}$ . In any case we can take an orbifold structure on the realizing torus orbifold such that the corresponding edge vectors  $\{v_i\}$  are all primitive; that is, if  $v_i = a_i v'_i$  for some  $v'_i \in N$  and  $a_i \in \mathbb{Z}$ , then  $a_i = \pm 1$ .

Proof. By Lemma 13.5, we may assume that the union of cones C(I) over  $I \in \Sigma^{(n)}$  covers the entire space  $N_{\mathbb{R}}$  and the pair  $(w^+(I), w^-(I))$ , which we denote by (p, q), is independent of I. When (p, q) = (1, 0),  $\Delta$  can be realized by a weighted projective space, say X. There is an orbifold structure on a weighted projective space such that the edge vectors are all primitive. We admit these facts for a moment; the proof will be give in the appendix at the end of this section. Then -X realizes the case when (p, q) = (0, 1). This completes the proof when n = 2.

Suppose  $n \ge 3$ . For a general value of (p, q), we prepare p copies of X and q copies of -X and do equivariant connected sum along all  $X_i$ 's and  $-X_i$ 's for each  $i \in \Sigma(X)$ . Then the resulting torus orbifold realizes  $\Delta$ . The edge vectors are all primitive in this construction since it is so for X.

Now let  $\Delta$  be an arbitrary complete simplicial multi-fan of dimension  $n \geq 2$ . We decompose  $\Delta$  into a number of minimal multi-fans as follows. We choose and fix a generic (rational) 1-dimensional cone in  $N_{\mathbb{R}}$ , say l, which is not contained in any subspaces spanned by cones of dimension  $\leq n - 1$  in  $\Delta$ . We label l as  $\star$ . To each *n*-dimensional cone C(I) for  $I \in \Sigma^{(n)}$ , we form *n* cones which are respectively spanned by l and facets of C(I). These *n* cones together with C(I) determine a simplicial multi-fan  $\Delta[I] = (\Sigma[I], C[I], w[I]^{\pm})$ , where  $\Sigma[I]$  consists of all proper subsets of  $I \cup \{\star\}$ . The weight functions  $w[I]^{\pm}$  are defined as follows. Let  $v_i$  be a non-zero vector in C(i) for each  $i \in I$  and  $v_{\star}$  a non-zero vector in l. Then there is a relation

$$v_{\star} + \sum_{i \in I} a_i v_i = 0$$

with non-zero real numbers  $a_i$ 's. Let  $\mathcal{I} \in \Sigma[I]^{(n)}$ . Then  $\mathcal{I} = I$  or  $(I \setminus \{i\}) \cup \{\star\}$  for  $i \in I$ . We define

(13.3)

$$(w[I]^{+}(\mathcal{I}), w[I]^{-}(\mathcal{I})) \coloneqq \begin{cases} (w^{+}(I), w^{-}(I)) & \text{if } \mathcal{I} = I \text{ or} \\ & \mathcal{I} = (I \setminus \{i\}) \cup \{\star\} \text{ and } a_i > 0, \\ (w^{-}(I), w^{+}(I)) & \text{if } \mathcal{I} = (I \setminus \{i\}) \cup \{\star\} \text{ and } a_i < 0. \end{cases}$$

**Lemma 13.7.**  $\Delta[I]$  is complete and hence minimal.

Proof. The proof is essentially the same as that of lemma 13.5. As remarked in Section 2, it suffices to show that, when a generic vector v gets across an (n-1)-dimensional cone, the integer  $d_v$  in Section 2 remains unchanged. Let  $\mathcal{J}$  be an element of  $\Sigma[I]^{(n-1)}$  and let i and i' be the two elements in  $(I \cup \{\star\}) \setminus \mathcal{J}$ . Then  $\mathcal{I} := \mathcal{J} \cup \{i\}$  and  $\mathcal{I}' := \mathcal{J} \cup \{i'\}$  are the elements in  $\Sigma[I]^{(n)}$  which contain  $\mathcal{J}$ . We project cones  $C[I](\mathcal{I})$  and  $C[I](\mathcal{I}')$  on  $N_{\mathbb{R}}^{C[I](\mathcal{J})}$ . Then it follows from (13.2) that the vectors projected from  $v_i$  and  $v_{i'}$  are toward opposite directions if and only if  $a_i a_{i'} > 0$ , where  $a_{\star}$  is understood to be 1. This together with the definition (13.3) of  $w[I]^{\pm}$  implies that  $d_v$  remains unchanged regardless of the sign of  $a_i a_{i'}$  when v gets across the (n-1)-dimensional cone  $C[I](\mathcal{J})$ .

Let  $J \in \Sigma^{(n-1)}$  and let  $I_1, \ldots, I_r$  be the elements in  $\Sigma^{(n)}$  containing J. The *n*-dimensional cone spanned by C(J) and l appears in  $\Delta[I_k]$  for  $k = 1, 2, \ldots, r$  with the form  $C[I_k](J \cup \{\star\})$ .

**Lemma 13.8.**  $\sum_{k=1}^{r} w[I_k](J \cup \{\star\}) = 0.$ 

Proof. Consider the projection of the cones  $C(I_k)$ 's on  $N_{\mathbb{R}}^{C(J)}$ We define sign $(I_k) = 1$  or -1 according as the projection image of  $C(I_k)$  disagrees or agrees with that of I. Applying (13.3) with  $I = I_k$  and  $I \setminus \{i\} = J$ , one sees that

$$w[I_k](J \cup \{\star\}) = \operatorname{sign}(I_k)w(I_k).$$

On the other hand, it follows from the completeness of  $\Delta$  that

$$\sum_{\operatorname{sign}(I_s)=1} w(I_s) = \sum_{\operatorname{sign}(I_t)=-1} w(I_t).$$

These two equalities imply the lemma.

Proof of Theorem 13.3. By lemma 13.4 we may assume that  $\Delta$  is connected. We choose a generic (rational) 1-dimensional cone l and decompose  $\Delta$  using l into minimal multi-fans  $\Delta[I]$ 's  $(I \in \Sigma^{(n)})$ . By Lemma 13.6  $\Delta[I]$  is realized by a torus orbifold, say M[I], such that all its edge vectors are primitive. We consider the disjoint union of M[I] over  $I \in \Sigma^{(n)}$  and piece them together using equivariant connected sum in the following way. For each  $i \in \Sigma^{(1)}$  we do equivariant connected sum of  $\{M[I] \mid i \in I\}$  successively along  $M[I]_i$ 's, and similarly do equivariant connected sum of all M[I]'s along  $M[I]_{\star}$  as well. The resulting space is connected because  $\Delta$ is connected, and becomes a torus orbifold. Its multi-fan is close to  $\Delta$  but contains extra cones which are the cones spanned by l and C(J) for  $J \in \Sigma^{(m)}$  with  $m \le n-1$ . For a fixed  $J \in \Sigma^{(n-1)}$ , it follows from Lemma 13.8 that there are the same number of T-fixed points p with  $\epsilon_p = 1$  and q with  $\epsilon_q = -1$  contained in the union of  $M[I_k]$  with  $J \subset I_k$  and corresponding to the cone spanned by l and C(J). Hence one can do equivariant connected sum at pairs of T-fixed points p and q so that those T-fixed points will be eliminated. Doing this for each  $J \in \Sigma^{(n-1)}$ , we obtain a torus orbifold, say M, realizing  $\Delta$ . In fact, the characteristic suborbifolds  $M[I]_{\star}$  turn into a codimension two suborbifold of M, which is fixed by the circle subgroup determined by l but has no T-fixed point, so it is not a characteristic suborbifold of M by definition. This means that all the cones in  $\Delta[I]$ 's containing l as an edge do not show up in the mulit-fan of M.

Proof of Theorem 13.2. We already observed the "only if" part, so we prove the "if" part. By Lemma 13.4 we may assume that our  $\Delta$ , which satisfies the conditions (1) and (2) in Theorem 13.2, is connected. Then (the realization of)  $\Sigma$  is either

CASE 1. a 1-simplex, or

CASE 2. the boundary of a *d*-gon where  $d \ge 3$ ,

63

and that

$$\{w^+(I), w^-(I)\} = \begin{cases} \{1, 1\} & \text{in Case 1,} \\ \{1, 0\} & \text{in Case 2.} \end{cases}$$

Using the latter statement in Lemma 13.6, the same argument as in the proof of Theorem 13.3 shows that  $\Delta$  in Case 2 is geometrically realized. As for Case 1, let  $I \in \Sigma^{(2)}$ be the unique simplex. There exist a finite covering  $\tilde{T} \to T$  whose kernel H is isomorphic to  $N/N_I$  where  $N_I$  is the sublattice generated by the primitive vectors  $v_i$ 's for  $i \in I$ , and a 2-dimensional  $\tilde{T}$ -module V corresponding to the cone C(I), as was explained in Section 12. Then the one point compactification of V/H, i.e., the orbit space of  $S^4$  by an action of  $N/N_I$ , realizes our  $\Delta$  in Case 1.

### Appendix. Realization of minimal multi-fans by weighted projective spaces

We identify the (n + 1)-dimensional torus  $T^{n+1} = S^1 \times \cdots \times S^1$  with the standard maximal torus of  $GL(n+1,\mathbb{C})$  consisting of diagonal matrices. We set  $\tilde{T} = T^{n+1}/D$ where D denotes the subgroup of diagonal elements  $(z, \ldots, z)$ . It is a maximal torus in  $PGL(n+1,\mathbb{C})$  and acts effectively on the projective space  $\mathbb{P}^n$ . Let  $\tilde{S}_i$  denote the *i*th factor of  $T^{n+1}$ . It is mapped injectively into  $\tilde{T}$ . We shall denote by the same letter  $\tilde{S}_i$  its image in  $\tilde{T}$ . We set  $\tilde{M}_i = \{[z_0, \ldots, z_n] \mid z_i = 0\}$ , for  $i = 0, \ldots, n$ . They are the characteristic submanifolds of  $\mathbb{P}^n$  regarded as a torus manifold with the orientations induced from the complex structure. If H is a finite subgroup of  $\tilde{T}$ , then the quotient  $M_H = \mathbb{P}^n/H$  is a torus orbifold acted on by  $T = \tilde{T}/H$  for which  $(M_H, \mathbb{P}^n, H, p)$  is an orbifold chart, where  $p: \mathbb{P}^n \to M_H$  is the projection. It is called a weighted projective space. Its characteristic suborbifolds are  $M_i = p(\tilde{M}_i), i = 0, ..., n$ , and the corresponding circle subgroups are  $S_i = \pi(\tilde{S}_i)$ , where  $\pi \colon \tilde{T} \to T$  is the projection. The symmetric group  $S_{n+1}$  of degree n+1 acts on  $T^{n+1}$  and also induces an action on  $\tilde{T}$ . It also acts on  $\mathbb{P}^n$ . If  $H^{\sigma}$  denotes the transform of H by an element  $\sigma \in \mathcal{S}_{n+1}$ , then the transformation  $\sigma \colon \mathbb{P}^n \to \mathbb{P}^n$  induces an isomorphism of torus manifolds  $M_H \to M_{H^{\sigma}}$ . We set

$$WP = \{H \mid \text{finite subgroup of } \tilde{T}\}/S_{n+1}.$$

Every element in WP represents an isomorphism class of weighted projective spaces.

In order to describe the multi-fan  $\Delta_H$  associated with the torus orbifold  $M_H$  we introduce the following notations:

$$\tilde{N} = \mathbb{Z}^{n+1}$$
/diagonal submodule,  $\tilde{v}_i$  = image of  $e_i$  in  $\tilde{N}$ ,  $N = \mathbb{Z}^n$ ,

where  $e_i$  is the *i*-th fundamental unit vector in  $\mathbb{Z}^{n+1}$ .  $\tilde{N}$  is canonically identified with Hom $(S^1, \tilde{T})$ . If one chooses an identification of Hom $(S^1, T) = H_2(BT; \mathbb{Z})$  with N, then the finite covering map  $\pi: \tilde{T} \to T$  induces an injective homomorphism  $\varphi: \tilde{N} \to N$ .

The vectors  $v_i = \varphi(\tilde{v}_i)$  are the edge vectors of the 1-dimensional cones of  $\Delta_H$ . Note that they satisfy the equality

(13.4) 
$$\sum_{i} v_i = 0,$$

since the  $\tilde{v}_i$ 's satisfy a similar equality. This implies that  $\Delta_H$  is a minimal multi-fan satisfying the conditon (1) in Lemma 13.5. It is also clear that  $(w^+(I), w^-(I)) = (1, 0)$ . We shall denote by  $\mathcal{MF}$  the set of minimal multi-fans satisfying the above two conditions. If one chooses another identification of  $\operatorname{Hom}(S^1, T)$  with N, then  $\varphi$  is transformed to  $\psi \circ \varphi$  where  $\psi \in GL(n, \mathbb{Z})$ .  $GL(n, \mathbb{Z})$  acts on  $\mathcal{MF}$  from left by transforming the cones simultaneously by its elements. Let  $d_H \in \mathbb{Z}$  be the maximal common divisor of the edge vectors  $v_i$  of  $\Delta_H$ . We get a correspondence

$$\alpha \colon \mathcal{WP}/\mathcal{S}_{n+1} \to GL(n,\mathbb{Z}) \backslash \mathcal{MF} \times \mathbb{Z}_{>0}$$

which sends H to  $(\Delta_H, d_H)$ .

**Lemma 13.9.** The correspondence  $\alpha$  is a bijection. In particular, every minimal multi-fan  $\Delta$  in  $\mathcal{MF}$  is realizable.

Proof. We shall define a correspondence  $\beta: GL(n, \mathbb{Z}) \setminus \mathcal{MF} \times \mathbb{Z}_{>0} \to \mathcal{WP}/S_{n+1}$ which is to be the inverse of  $\alpha$ . Take a multi-fan  $\Delta$  in  $\mathcal{MF}$  and  $d \in \mathbb{Z}_{>0}$ . It is easy to see there is a unique set  $\{v_i\}$  of edge vectors of  $\Delta$  such that  $\sum_i v_i = 0$ and the maximal common divisor of  $\{v_i\}$  is d. Define a homomorphism  $\varphi: \tilde{N} \to N$ by requiring  $\varphi(\tilde{v}_i) = v_i$ . Then there is a unique finite covering map  $\pi: \tilde{T} \to T$ which induces  $\varphi: \tilde{N} = \operatorname{Hom}(S^1, \tilde{T}) \to N = \operatorname{Hom}(S^1, T)$ . Let H be the kernel of  $\pi$ . The homomorphism  $\varphi$ , hence H either, does not depend on the choice of identification  $N = \operatorname{Hom}(S^1, T)$ , but it depends on the numbering of  $v_i$ 's. So if we put  $\beta(\Delta, d) =$  the class of H in  $\mathcal{WP}/S_{n+1}$ , it induces a correspondence  $\beta$  as above. It is clear that  $\beta$  is in fact the inverse of  $\alpha$ .

REMARK. Let *a* be a positive integer. The correspondence  $T^{n+1} \ni (z_0, z_1, \ldots, z_n) \mapsto (z_0^a, z_1^a, \ldots, z_n^a) \in T^{n+1}$  induces a homomorphism  $\rho: \tilde{T} \to \tilde{T}$ . For a finite group *H* of  $\tilde{T}$  define  $H' = \rho^{-1}(H)$ . The edge vectors  $\{v'_i\}$  corresponding to the torus manifold  $M_{H'}$  are of the form  $v'_i = av_i$ , where  $\{v_i\}$  correspond to  $M_H$ . Hence  $\Delta_H = \Delta_{H'}$  and  $d_{H'} = ad_H$ . Let  $g: \mathbb{P}^n \to \mathbb{P}^n$  be the map defined by  $g[z_0, z_1, \ldots, z_n] = [z_0^a, z_1^a, \ldots, z_n^a]$ . Then it induces a homeomorphism  $M_{H'} \to M_H$  which is equivariant with respect to the isomorphism of tori between  $\tilde{T}/H'$  and  $\tilde{T}/H$  induced by  $\rho$ . If  $M_H$  and  $M_{H'}$  are considered as algebraic varieties then the homeomorphism becomes an equivalence. It is a fundamental fact in the theory of toric varieties that to each fan corresponds a toric variety. The above equivalence gives an interpretation of this fact within this special case in our context. Related results are found in [26]. Related to the above remark,

for a later use, we point out the following fact. Let  $a_i, \ldots, a_n$  be positive integers, and let  $\mathbb{Z}/a_i \subset S^1$  be the subgroup of  $a_i$ -th roots of unity. Set  $G = \prod_i \mathbb{Z}/a_i$ . Then the map  $\mathbb{C}^n \ni (z_1, \ldots, z_n) \mapsto (z_1^{a_1}, \ldots, z_n^{a_n}) \in \mathbb{C}^n$  induces an equivalence of affine algebraic varieties  $\mathbb{C}^n/G \to \mathbb{C}^n$ .

Let  $M_H \in WP$  and let  $\{v_i\}$  be the edge vectors corresponding to the orbifold structure as given above. Even if  $d_H = 1$ , it may happen that some of  $v_i$ 's are not primitive. We will show that there always exists a torus orbifold structure on  $M_H$  such that the corresponding edge vectors are all primitive. More generally we have

**Lemma 13.10.** Let  $M_H$  be a weighted projective space and  $\{v_i\}$  the corresponding edge vectors satisfying  $\sum_i v_i = 0$  as given above. Suppose that  $\{v'_i\}$  are vectors in N such that  $v_i = a_i v'_i$  with  $a_i \in \mathbb{Z}_{>0}$ . Then there is an orbifold structure on  $M_H$  which admits  $\{v'_i\}$  as the corresponding edge vectors.

Proof. For each  $x \in M_H$  let  $\tilde{T}_x \subset \tilde{T}$  be the isotropy subgroup at  $\tilde{x}$  of the  $\tilde{T}$ action on  $\mathbb{P}^n$  where  $\tilde{x} \in p^{-1}(x)$ .  $\tilde{T}_x$  does not depend on the choice of  $\tilde{x}$  in  $p^{-1}(x)$ . If x lies in Int  $M_I = M_I \setminus \bigcup_{J \supseteq I} M_J$  for  $I \in \Sigma(M_H)^{(k)}$ , then  $\tilde{T}_x = \tilde{S}_I = \prod_{i \in I} \tilde{S}_i$ . We put  $H_x = H \cap \tilde{T}_x$ . We take a family  $\{V_{x,\mu} \mid \mu \in \mathbb{Z}_{>0}\}$  of small  $\tilde{T}_x$ -invariant open neighborhoods of  $\tilde{x}$  such that  $V_{x,\mu}$  converges to  $\tilde{x}$  when  $\mu$  tends to infinity. We may assume that  $V_{x,\mu}$  is equivariantly diffeomorphic to an  $\tilde{S}_I$ -invariant open disk in  $\mathbb{C}^n$ . It is possible to make  $V_{x,\mu}$ 's so small that they satisfy the following condition:

(13.5) 
$$H_x = \{h \in H \mid h \cdot V_{x,\mu} \cap V_{x,\mu} \neq \emptyset\}.$$

Then  $U_{x,\mu} = V_{x,\mu}/H_x$  is an open neighborhood of x in  $M_H$ , and  $(U_{x,\mu}, V_{x,\mu}, H_x, p \mid V_{x,\mu})$  is an orbifold chart of  $M_H$  compatible with  $(M_H, \mathbb{P}^n, H, p)$ .

On the other hand the fact that  $v_i = a_i v'_i$  implies that the kernel of  $p: \tilde{S}_i \to S_i$ contains  $\mathbb{Z}/a_i$ , which we denote by  $G_i$ . Since H is the kernel of  $p: \tilde{T} \to T$ ,  $G_i$  is contained in H. We put  $G_I = \prod_{i \in I} G_i$  for  $I \in \Sigma(M_H)^{(k)}$  and define

$$V'_{x,\mu} = V_{x,\mu}/G_I, \ H'_x = H_x/G_I \ \text{for } x \in \text{Int } M_I.$$

 $V'_{x,\mu}$  can be considered as an open disk in  $\mathbb{C}^n$  as pointed out in Remark above. The projection  $p \mid V_x \colon V_x \to U_x$  induces a map  $p'_{x,\mu} \colon V'_{x,\mu} \to U_x$  which induces a homeomorphism  $V'_{x,\mu}/H'_x \to U_x$ .

We shall prove that the family  $\{(U'_{x,\mu}, V'_{x,\mu}, H'_x, p'_{x,\mu}) \mid x \in M, \mu \in \mathbb{Z}_{>0}\}$  forms a set of orbifold charts of an orbifold structure on  $M_H$ . For that purpose it suffices to show that, if  $U'_{x,\mu} \subset U'_{y,\nu}$ , then there are an injective homomorphism  $\rho: H'_x \to H'_y$  and a  $\rho$ -equivariant open embedding  $\phi: V'_{x,\mu} \to V'_{y,\nu}$  such that

(13.6) 
$$\rho(H'_x) = \{h \in H'_y \mid h \cdot \phi(V'_{x,\mu}) \cap \phi(V'_{x,\mu}) \neq \emptyset\}.$$

The condition (13.5) implies that, if  $x \in \text{Int } M_I$  and  $y \in \text{Int } M_J$  with I and  $J \in \Sigma(M_H)$ , and if  $U'_{x,\mu} \subset U'_{y,\nu}$ , then  $I \supset J$ . Therefore

$$H_x \subset H_y$$
 and  $G_I \cap H_y = G_J$ .

It follows that the inclusion  $H_x \to H_y$  induces an injective homomorphism  $\rho: H'_x = H_x/G_I \to H_y/G_J = H'_y$ . If  $\tilde{x}$  is taken in  $V_{y,\nu}$ , then  $V_{x,\mu}$  is contained in  $V_{y,\nu}$ . The inclusion induces an embedding  $\phi: V'_{x,\mu} \to V'_{y,\nu}$ .  $\phi$  is clearly  $\rho$ -equivariant. The condition (13.6) follows from (13.5).

If x lies in  $M_i$ , then the action of  $S_i$  lifts to the action of  $\tilde{S}'_i = \tilde{S}_i/G_i$  on  $V'_{x,\mu}$  and the lifting is minimal. Hence the edge vector of C(i) corresponding to the orbifold structure defined above must be  $v'_i$ .

#### References

- [1] M. Atiyah: Convexity and commuting Hamiltonians, Bull. London Math. Soc. 14 (1981), 1–15.
- [2] M. Atiyah and R. Bott: *The moment map and equivariant cohomology*, Topology 23 (1984), 1–28.
- [3] R. Bott and L.W. Tu: Differential Forms in Algebraic Topology, GTM 82, Springer-Verlag, 1982.
- [4] M. Brion and M. Vergne: *Lattice points in simple polytopes*, J. Amer. Math. Soc. **10** (1997), 371–392.
- [5] M. Brion and M. Vergne: An equivariant Riemann-Roch theorem for complete, simplicial toric varieties, J. reine angew. Math. 482 (1997), 67–92.
- [6] S.E. Cappell and J.L. Shaneson: Genera of algebraic varieties and counting lattice points, Bull. Amer. Math. Soc., 30 (1994), 62–69.
- S.E. Cappell and J.L. Shaneson: *Euler-Maclaurin expansions for lattices above dimension one*, C. R. Acad. Sci. Paris Sr. I Math. **321** (1995), 885–890.
- [8] M. Davis and T. Januszkiewicz: Convex polytopes, Coxeter orbifolds and torus actions, Duke Math. J. 61 (1991), 417–451.
- T. Delzant: Hamiltoniens périodiques et image convexe de l'application moment, Bull. Soc. math. France 116 (1988), 315–339.
- [10] J.J. Duistermaat: The Heat Kernel Lefschetz Fixed Point formula for Spin-c Dirac Operator, Progress in Nonlinear Differential Equations and Their Applications, 8, Birkhäuser, 1996.
- [11] W. Fulton: Introduction to Toric Varieties, Ann. of Math. Studies, 131, Princeton Univ., 1993.
- [12] M. Grossberg and Y. Karshon: Equivariant index and the moment map for completely integrable torus actions, Adv. in Math. 133 (1998), 185–223.
- [13] V. Guillemin: Riemann-Roch for toric orbifolds, J. Diff. Geom. 45 (1997), 53-73.
- [14] V. Guillemin, E. Lerman and S. Sternberg: On the Konstant multiplicity formula, J. Geom. Phys. 5 (1988), 721–750.
- [15] V. Guillemin, E. Lerman and S. Sternberg: Symplectic fibrations and multiplicity diagrams, Cambridge Univ. Press, Cambridge, 1996.
- [16] V. Guillemin and S. Sternberg: Convexity properties of the momont mapping, Invent. math. 67 (1982), 491–513.
- [17] A. Hattori: Almost complex toric manifolds and positive line bundles, Homotopy and Geometry, Banach Center Publications, 45 (1998), 95–114.
- [18] A. Hattori and H. Taniguchi: Smooth S<sup>1</sup>-actions and bordism, J. Math. Soc. Japan 24 (1972),

#### A. HATTORI AND M. MASUDA

701-731.

- [19] A. Hattori and T. Yoshida: Lifting compact group actions into fiber bundles, Japan. J. Math. 2 (1976), 13–25.
- [20] F. Hirzebruch: Topological Methods in Algebraic Geometry, Springer-Verlag 1966.
- [21] J.M. Kantor and A.G. Khovanskii: Une application du théorème de Riemann-Roch combinatoire au polynome d'Ehrhart des polytopes entiers de R<sup>d</sup>, C. R. Acad. Sci. Paris Ser. I Math. 317 (1993), 501–507.
- [22] Y. Karshon and S. Tolman: The moment map and line bundles over presymplectic toric manifolds, J. Diff. Geom. 38 (1993), 465–484.
- [23] K. Kawakubo: The Theory of Transformation Groups, Oxford Univ. Press, 1991.
- [24] T. Kawasaki: The signature theorem for V-manifolds, Topology 17 (1978), 75-83.
- [25] A.G. Khovanskii and A.V. Pukhlikov: A Riemann-Roch theorem for integrals and sums of quasipolynomials over virtual polytopes, St. Petersburg Math. J. 4 (1993), 789–812.
- [26] E. Lerman and S. Tolman: Hamiltonian torus actions on symplectic orbifolds and toric varieties, Trans. Amer. Math. Soc. 349 (1997), 4201–4230.
- [27] M. Masuda: Unitary toric manifolds, multi-fans and equivariant index, Tohoku Math. J. 51 (1999), 237–265.
- [28] Y. Nishimura: Convex chains and multi-polytopes, preprint.
- [29] T. Oda: Convex Bodies and Algebraic Geometry, Springer-Verlag, 1988.
- [30] P. Orlik and F. Raymond: Actions of the torus on 4-manifolds. I, Trans. Amer. Math. Soc. 152 (1970), 531–559.
- [31] I. Satake: The Gauss-Bonnet theorem for V-manifolds, J. Math. Soc. Japan 9 (1957), 464-492.
- [32] R. Stanley: Combinatorics and Commutative Algebra (second edition), Progress in Math. 41, Birkhäuser, 1996.
- [33] M. Vergne: Equivariant index formula for orbifolds, Duke Math. J. 82 (1996), 637–652.

Mikiya Masuda Department of Mathematics Osaka City University Osaka, Japan e-mail: masuda@sci.osaka-cu.ac.jp

Akio Hattori Graduate School of Mathematical Science University of Tokyo Tokyo, Japan e-mail: hattori@ms.u-tokyo.ac.jp

68