



Title	An asymptotic behavior of the mean of some exponential functionals of a random walk
Author(s)	Hirano, Katsuhiro
Citation	Osaka Journal of Mathematics. 1997, 34(4), p. 953-968
Version Type	VoR
URL	<a href="https://doi.org/10.18910/5709">https://doi.org/10.18910/5709</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Hirano, K.  
 Osaka J. Math.  
**34** (1997), 953–968

# AN ASYMPTOTIC BEHAVIOR OF THE MEAN OF SOME EXPONENTIAL FUNCTIONALS OF A RANDOM WALK

KATSUHIRO HIRANO

(Received November 19, 1996)

## 1. Introduction

Let  $\{\xi_i\}_{i \geq 1}$  be a sequence of independent identically distributed random variables, and set  $S_0 = 0$ ,  $S_n = \sum_{i=1}^n \xi_i$ ,  $n \geq 1$ . Let  $f(x)$ ,  $h(x)$  be two functions on  $[0, \infty)$ . For  $n \geq 1$ , set

$$P_n = E \left[ f(e^{S_n}) h \left( \sum_{i=1}^n e^{S_i} \right) \right].$$

One object of this paper is to obtain an estimate of  $P_n$  under certain conditions of  $\xi_1$ ,  $f(x)$  and  $h(x)$ .

We can consider a similar problem in case of Brownian motion. Let  $\{B(t) : t \geq 0\}$  be a Brownian motion with  $B(0) = 0$ . We assume that  $f(x)$  and  $h(x)$  satisfy the following: for some  $0 < \alpha < \beta < \infty$ ,

$$\sup_{x \geq 0} x^{-\alpha} |f(x)| < \infty, \quad \sup_{x \geq 0} x^\beta |h(x)| < \infty.$$

Set

$$P(t) = E \left[ f(e^{B(t)}) h \left( \int_0^t e^{B(s)} ds \right) \right].$$

Applying a very useful formula of Yor ([9],(6.e)), we see as in [5] that

$$P(t) \sim ct^{-3/2} \quad \text{as } t \rightarrow \infty,$$

where

$$c = 2^{\frac{5}{2}} \pi^{-\frac{1}{2}} \int_0^\infty \int_0^\infty \int_0^\infty f(y^2) h(4/z) e^{-\lambda z} u \sinh u dy dz du, \quad \lambda = (1 + y^2)/2 + y \cosh u.$$

Kawazu-Tanaka [5] used this fact to obtain the rate of decay of the tail probability of the maximum of a diffusion process in a drifted Brownian environment. This

result in case of a Brownian motion suggests that an analogous result in a random walk case also holds. In fact we obtain Theorem 1 below. But to our regret, we can not find the precise value corresponding to  $c$ .

The original motivation of this problem is to clarify and simplify the proof of Afanas'ev's result [1] on a random walk in a random medium. In the last section we get Theorem 2 which is a slight extention of the theorem in [1] by using our Theorem 1.

Now we state the conditions on  $\xi = \xi_1$  and Theorem 1.

Conditions (a)-(d).

- (a)  $E\xi = 0$ ,  $0 < E\xi^2 = \sigma^2 < \infty$ .
- (b) The distribution of  $\xi$  is continuous and  $E|\xi|^3 < \infty$ .
- (c)  $Ee^{\theta\xi}$  converges for some  $\theta > 0$ .

The condition (b) can be replaced by the following condition (d).

- (d) The distribution of  $\xi$  is concentrated on the set  $\{id : i \in \mathbb{Z}\}$  with some  $d > 0$ .

**Theorem 1.** *Let functions  $f$ ,  $h$  and  $W$  satisfy the following conditions:*

- (i)  $f$  and  $h$  are continuous on  $[0, \infty)$ .
- (ii)  $W$  is continuous on  $\mathbb{R}$  and non-negative.
- (iii) There exist positive numbers  $\alpha, \beta, \varepsilon, \eta$  such that

$$\sup_{x \geq 0} x^{-\alpha} |f(x)| < \infty, \quad \sup_{x \geq 0} x^{\beta} |h(x)| < \infty,$$

$$\limsup_{x \rightarrow -\infty} W(x)e^{-\varepsilon x} < \infty, \quad \liminf_{x \rightarrow +\infty} W(x)e^{-\eta x} > 0,$$

with  $\beta\eta > \alpha > 0$ . If the conditions (a), (b) and (c) are satisfied, then

$$E \left[ f(e^{S_n}) h \left( \sum_{i=1}^n W(S_i) \right) \right] \sim cn^{-3/2},$$

as  $n \rightarrow \infty$  with a constant  $c$ . The same conclusion holds under the conditions (a), (d) and (c).

**REMARK.** A sufficient condition for the positivity of  $c$  is given in Corollary 10.

## 2. Preliminaries

We introduce the following quantities together with their corresponding Laplace transforms.

$$u_n(x) = P(S_1 > 0, \dots, S_{n-1} > 0, 0 < S_n \leq x),$$

$$\begin{aligned}
v_n(x) &= P(S_1 \leq 0, \dots, S_{n-1} \leq 0, -x \leq S_n \leq 0), \\
u_0(x) &= v_0(x) = 1_{[0, \infty)}(x), \\
\varphi_n(\theta) &= \int_0^\infty e^{-\theta x} du_n(x), \quad \psi_n(\theta) = \int_0^\infty e^{-\theta x} dv_n(x).
\end{aligned}$$

The three lemmas below will be used later. We state them without proof.

**Lemma 1** (Spitzer-Baxter identity. See [6], p. 49). *For any  $\theta > 0$ ,  $|t| < 1$ ,*

$$1 + \sum_{n=1}^\infty \varphi_n(\theta) t^n = \exp \left\{ \sum_{n=1}^\infty \frac{t^n}{n} E(e^{-\theta S_n}; S_n > 0) \right\}.$$

**Lemma 2** (See [2]). *If the conditions (a) and (b) hold, then for any  $\theta > 0$ ,*

$$E(e^{-\theta S_n}; S_n > 0) \sim (\sqrt{2\pi}\sigma\theta)^{-1} n^{-\frac{1}{2}} \quad \text{as } n \rightarrow \infty.$$

**Lemma 3** (See [4] and [8]).

- (1) *Let  $\sum_{n=0}^\infty a_n t^n = \exp(\sum_{n=1}^\infty b_n t^n)$  for  $|t| < 1$ . If  $b_n \sim bn^{-3/2}$ , then  $a_n \sim (b \exp B)n^{-3/2}$  with  $B = \sum_{n=1}^\infty b_n$ .*
- (2) *Let  $c_n \geq 0$ ,  $d_n \geq 0$ ,  $c_n \sim cn^{-3/2}$  and  $d_n \sim dn^{-3/2}$ . If  $a_n = \sum_{j=0}^n c_{n-j} d_j$ , then  $a_n \sim (cD + dC)n^{-3/2}$  with  $C = \sum_{n=0}^\infty c_n$  and  $D = \sum_{n=0}^\infty d_n$ .*

Set

$$u(x) = \sum_{n=0}^\infty u_n(x), \quad v(x) = \sum_{n=0}^\infty v_n(x),$$

and

$$\begin{aligned}
U(x) &= \frac{1}{\sqrt{2\pi}\sigma} \int_0^x u(y) dy, & \varphi(\theta) &= \int_0^\infty e^{-\theta x} dU(x), \\
V(x) &= \frac{1}{\sqrt{2\pi}\sigma} \int_0^x v(y) dy, & \psi(\theta) &= \int_0^\infty e^{-\theta x} dV(x).
\end{aligned}$$

Then we can prove the following lemma.

**Lemma 4.** *If the conditions (a) and (b) hold, then*

- (1)  $\lim_{n \rightarrow \infty} n^{3/2} \varphi_n(\theta) = \varphi(\theta)$ ,  $\lim_{n \rightarrow \infty} n^{3/2} \psi_n(\theta) = \psi(\theta)$  for each  $\theta > 0$ .
- (2)  $\lim_{n \rightarrow \infty} n^{3/2} u_n(x) = U(x)$ ,  $\lim_{n \rightarrow \infty} n^{3/2} v_n(x) = V(x)$  compact uniformly on  $[0, \infty)$ .
- (3)  $\lim_{n \rightarrow \infty} n^{3/2} P(M_n \leq x, M_n - S_n \leq y) = U(x)v(y) + u(x)V(y)$  where  $M_n = \max_{0 \leq j \leq n} S_j$ .

Proof. Combining Lemmas 1, 2 with (1) of Lemma 3, for each  $\theta > 0$ , we have

$$\lim_{n \rightarrow \infty} n^{\frac{3}{2}} \varphi_n(\theta) = \frac{1}{\sqrt{2\pi}\sigma\theta} \left( 1 + \sum_{n=1}^{\infty} \varphi_n(\theta) \right).$$

It is easy to see that the right hand side of the above is  $\varphi(\theta)$ . If we consider a reversed random walk  $\{-S_n\}_{n \geq 0}$ , we get the assertion for  $\psi$  also. By the extended continuity theorem for Laplace transform, (1) yields (2). In view of (2), (3) can be proved by applying (2) of Lemma 3 to the following identity ([6], p. 25).

$$(2.1) \quad P(M_n \leq x, M_n - S_n \leq y) = \sum_{j=0}^n u_j(x) v_{n-j}(y) \quad n \geq 0.$$

The proof of Lemma 4 is complete.  $\square$

We give some definitions. Fix a constant  $K > 0$ . For a function  $f$  on  $(-\infty, K]$  and a constant  $a \in \mathbb{R}$ , set

$$\|f\|_a = \sup_{x \leq K} e^{ax} |f(x)|,$$

and define a function space  $C(a)$ :

$$C(a) = \{f : f \text{ is continuous on } (-\infty, K] \text{ and } \|f\|_a < \infty\}.$$

$C(a)$  is a Banach space with respect to the norm  $\| \cdot \|_a$ . From now on we omit  $a$  in  $\| \cdot \|_a$ . By the condition (c), we can take  $\delta > 0$  such that  $E(e^{\delta\xi}) < \infty$ . Let this  $\delta > 0$  and arbitrary  $\mu > \delta$  be fixed, and set  $\lambda = \mu - \delta > 0$ . Let non-negative  $W \in C(-\mu)$  be fixed. For any  $f \in C(\lambda)$ , we define a family of transforms  $\{R_t\}_{t \geq 0}$  depending on  $W$  by

$$(2.2) \quad R_t f(x) = E_x \left[ f(\xi) \{1 - e^{-tW(\xi)}\}; \xi \leq K \right],$$

where  $E_x$  denotes the expectation of the random walk starting at  $x$ . For any  $f \in C(-\delta)$ , we define a sequence of transforms  $\{T_n\}_{n \geq 0}$  by

$$(2.3) \quad T_n f(x) = E_x [f(S_n); M_n \leq K], \quad n \geq 0.$$

In the sequel, the conditions (a)-(c) are assumed to be satisfied.

**Lemma 5.** *Each  $R_t$  is a bounded operator from  $C(\lambda)$  to  $C(-\delta)$ .*

Proof. Let  $f \in C(\lambda)$ . The continuity of  $R_t f$  is derived from the condition (b) and the continuity of  $f$  and  $W$ . From the assumptions of  $f$  and  $W$ ,

$$|f(x) \{1 - e^{-tW(x)}\}| \leq tW(x) |f(x)| \leq t \|W\| \|f\| e^{(\mu-\lambda)x}.$$

By the definition of  $\lambda$  and (2.2),

$$\begin{aligned}|R_t f(x)| &\leq t\|W\|\|f\|E(e^{\delta(x+\xi)}; \xi \leq K-x) \\ &\leq t\|W\|\|f\|e^{\delta x}E(e^{\delta\xi}).\end{aligned}$$

Therefore we obtain

$$\|R_t\| \leq t\|W\|E(e^{\delta\xi}) < \infty.$$

The proof of Lemma 5 is complete.  $\square$

**Lemma 6.** *Each  $T_n$  is a bounded operator from  $C(-\delta)$  to  $C(\lambda)$ , and there exists a constant  $C$  such that  $\|T_n\| \leq Cn^{-3/2}$  for  $n \geq 1$ . Here  $C$  does not depend on  $n$ .*

**Proof.** Let  $f \in C(-\delta)$ . The continuity of  $T_n f$  is derived from the condition (b) and the continuity of  $f$ . Direct calculations show  $\|T_0\| = e^{\mu K}$ . We will estimate  $\|T_n\|$  for  $n \geq 1$ . Since  $f \in C(-\delta)$ , we have  $|f(x)| \leq \|f\|e^{\delta x}$ . Applying this and (2.1) to (2.3), we see

$$\begin{aligned}|T_n f(x)| &\leq \|f\|e^{\delta x}E(e^{\delta S_n}; M_n \leq K-x) \\ &\leq \|f\|e^{\delta x} \sum_{j=0}^n \psi_{n-j}(\delta) \int_0^{K-x} e^{\delta y} du_j(y).\end{aligned}$$

Using Chebyshev's inequality, we see

$$\int_0^{K-x} e^{\delta y} du_j(y) \leq e^{\delta(K-x)} u_j(K-x) \leq e^{(\delta+\lambda)(K-x)} \varphi_j(\lambda).$$

Thus we have

$$|T_n f(x)| \leq \|f\|e^{\mu K - \lambda x} \sum_{j=0}^n \psi_{n-j}(\delta) \varphi_j(\lambda),$$

which shows

$$\|T_n\| \leq e^{\mu K} \sum_{j=0}^n \psi_{n-j}(\delta) \varphi_j(\lambda).$$

Applying (2) of Lemma 3 and (1) of Lemma 4 to the right hand side of the above, we have

$$C := e^{\mu K} \sup_{n \geq 1} \left\{ n^{\frac{3}{2}} \sum_{j=0}^n \psi_{n-j}(\delta) \varphi_j(\lambda) \right\} < \infty.$$

This completes the proof of Lemma 6.  $\square$

From Lemma 6 we get the following corollary.

**Corollary 7.** *Let  $S = \sum_{n=0}^{\infty} T_n$ . Then  $S$  is a bounded operator from  $C(-\delta)$  to  $C(\lambda)$ .*

Now we consider the limit of  $n^{3/2}T_n$ . Let  $f \in C(-\delta)$ ,  $x \leq K$  and  $L > 0$  be fixed. Then

$$\begin{aligned} T_n f(x) &= E_x [f(S_n); M_n \leq K, M_n - S_n \leq L] \\ &\quad + E_x [f(S_n); M_n \leq K, M_n - S_n > L] = a_n + b_n. \end{aligned}$$

Applying (3) of Lemma 4 to  $a_n$ , we have

$$\lim_{n \rightarrow \infty} n^{3/2} a_n = \int_0^L \int_0^{K-x} f(x+a-b) d(U(a)v(b) + u(a)V(b)).$$

Using the method employed in the proof of Lemma 6, we get

$$|b_n| \leq \|f\| e^{\mu K - \lambda x} \sum_{j=0}^n \varphi_{n-j}(\lambda) \int_L^\infty e^{-\delta b} dv_j(b).$$

In view of (1) and (2) of Lemma 4, we apply (2) of Lemma 3 to the right hand side

$$\limsup_{n \rightarrow \infty} n^{3/2} |b_n| \leq \|f\| e^{\mu K - \lambda x} \left\{ \sum_{n=0}^{\infty} \varphi_n(\lambda) \int_L^\infty e^{-\delta b} dV(b) + \varphi(\lambda) \int_L^\infty e^{-\delta b} dv(b) \right\}.$$

The last term goes to 0 as  $L \rightarrow \infty$ . Without loss of generality, we assume  $f \geq 0$ . Then we have

$$n^{3/2} a_n \leq n^{3/2} T_n f(x) \leq n^{3/2} a_n + \sup_{m \geq n} (m^{3/2} b_m).$$

Going to the limit, first with respect to  $n$ , and then with respect to  $L$ , we obtain

$$(2.4) \quad \lim_{n \rightarrow \infty} n^{3/2} T_n f(x) = \int_0^\infty \int_0^{K-x} f(x+a-b) d(U(a)v(b) + u(a)V(b)).$$

Denote by  $Qf(x)$  the right hand side of (2.4). It is clear that  $Qf$  is continuous by (2) and (3) of Lemma 4. The definition of  $Q$  and Lemma 6 show that

$$|Qf(x)| \leq \limsup_{n \rightarrow \infty} n^{3/2} |T_n f(x)| \leq \left( \limsup_{n \rightarrow \infty} n^{3/2} \|T_n\| \right) \|f\| e^{-\lambda x}.$$

Hence we have

$$\|Q\| \leq \limsup_{n \rightarrow \infty} \left( n^{\frac{3}{2}} \|T_n\| \right) \leq C.$$

Collecting the above results, we get the following lemma.

**Lemma 8.** *Let  $f \in C(-\delta)$ . Then we can define  $Qf(x) = \lim_{n \rightarrow \infty} n^{3/2} T_n f(x)$  and  $Q$  is a bounded operator from  $C(-\delta)$  to  $C(\lambda)$ .*

The following corollary will be used hereafter.

**Corollary 9.** *If  $f \in C(-\delta)$  is positive, then  $Qf(0) > 0$ .*

Proof. From (2.4) and the assumption of  $f$ , we have

$$Qf(0) \geq \int_0^\infty \int_0^K f(a-b) dU(a) dv(b).$$

By the definition of  $U(x)$ , we see  $U(K) \geq K/\sqrt{2\pi}\sigma > 0$ . The renewal theory ([3], Chap. XI) applied to  $v(x)$  yields  $\lim_{x \rightarrow \infty} v(x) = \infty$ , which shows the corollary.  $\square$

### 3. Main Lemma

The lemmas established in the previous section enable us to show Theorem 1 if  $h(x) = e^{-tx}$  ( $t \geq 0$ ) and  $W(x) = \infty$  for  $x > K$ . That is, we have the following lemma.

**Main Lemma.** *For any  $f \in C(-\delta)$ , non-negative  $W \in C(-\mu)$  and  $t \geq 0$ , the finite limit*

$$\lim_{n \rightarrow \infty} n^{3/2} E \left[ f(S_n) \exp \left\{ -t \sum_{i=1}^n W(S_i) \right\}; M_n \leq K \right]$$

*exists.*

To prove Main Lemma we need some preparations. Let non-negative  $W \in C(-\mu)$  be fixed. For each  $f \in C(-\delta)$  and  $t \geq 0$ , we put

$$A_0^{(t)} f(x) = f(x), \quad A_n^{(t)} f(x) = E_x \left[ f(S_n) \exp \left\{ -t \sum_{i=1}^n W(S_i) \right\}; M_n \leq K \right], \quad n \geq 1.$$

Then  $A_n^{(t)} = \{A_1^{(t)}\}^n$  as a transform, for  $\{\xi_i\}_{i \geq 1}$  is i.i.d. (2.2) and (2.3) yield  $T_1 - A_1^{(t)} = R_t$ . Put  $f_n^{(t)} = A_n^{(t)}f$ . Then  $f_n^{(t)} = A_1^{(t)}f_{n-1}^{(t)} = (T_1 - R_t)f_{n-1}^{(t)}$ . By induction we get the following equations in  $C(\lambda)$ .

$$(3.1) \quad f_0^{(t)} = f, \quad f_n^{(t)} = T_n f - \sum_{j=1}^n T_{n-j} R_t f_{j-1}^{(t)} \quad n \geq 1.$$

Taking account of the definitions of  $A_n^{(t)}$ ,  $T_n$  and Lemma 6, we have

$$(3.2) \quad \|f_n^{(t)}\| \leq \|T_n\| \|f\| \leq C \|f\| n^{-\frac{3}{2}}.$$

The operator  $SR_t$  maps  $C(\lambda)$  to  $C(\lambda)$ . Hence we can define

$$c_1 = \sup\{t > 0 : \|SR_t\| < 1\}.$$

For a while we assume  $0 \leq t < c_1$  and omit  $t$  in  $R_t$  and  $f_n^{(t)}$ . Then there exists  $(1 + SR)^{-1}$ . By (3.1) and (3.2),  $F \equiv \sum_{n=0}^{\infty} f_n = (1 + SR)^{-1} S f$ . Let  $g \in C(\lambda)$  satisfy the equation

$$(3.3) \quad g = Qf - QRF - SRg.$$

After some tedious calculations, we rewrite  $g$  as follows.

$$g = (1 + SR)^{-1} Q(1 + RS)^{-1} f.$$

We set

$$\bar{g}(x) = \limsup_{n \rightarrow \infty} |n^{\frac{3}{2}} f_n(x) - g(x)|.$$

By (3.2),  $\|\bar{g}\| \leq C \|f\| + \|g\| < \infty$ . Now we can start the proof of Main Lemma.

**Proof.** For fixed  $N \in \mathbb{N}$ , we have

$$n^{\frac{3}{2}} \sum_{j=1}^n T_{n-j} R f_{j-1} = n^{\frac{3}{2}} \left[ \sum_{j=1}^N + \sum_{j=N+1}^{n-N} + \sum_{j=n-N+1}^n \right] T_{n-j} R f_{j-1} = I_n + J_n + K_n.$$

Combining (3.1) and (3.3) with the above, we see

$$(3.4) \quad |n^{\frac{3}{2}} f_n - g| \leq |n^{\frac{3}{2}} T_n f - Qf| + |I_n - QRF| + |K_n - SRg| + |J_n|.$$

We estimate each term of the right hand side of (3.4). Using Lemmas 5, 6 and 8, we have

$$I_n = \sum_{j=1}^N \left( \frac{n}{n-j} \right)^{\frac{3}{2}} (n-j)^{\frac{3}{2}} T_{n-j} R f_{j-1} \longrightarrow \sum_{j=1}^N QRF_{j-1} \quad \text{as } n \rightarrow \infty.$$

Therefore we get

$$(3.5) \quad \lim_{n \rightarrow \infty} (QRF - I_n) = \sum_{j=N}^{\infty} QRf_j.$$

Rewriting  $K_n$ , we have

$$K_n = n^{\frac{3}{2}} \sum_{j=1}^N T_{j-1} R f_{n-j} = \sum_{j=1}^N \left( \frac{n}{n-j} \right)^{\frac{3}{2}} T_{j-1} R \left\{ (n-j)^{\frac{3}{2}} f_{n-j} \right\}.$$

By Corollary 7,

$$|K_n - SRg| \leq \left| K_n - \sum_{j=1}^N T_{j-1} R g \right| + \sum_{j=N+1}^{\infty} T_{j-1} R |g|.$$

The first term of the right hand side is bounded from above by the quantity

$$\sum_{j=1}^N \left( \frac{n}{n-j} \right)^{\frac{3}{2}} T_{j-1} R \left| (n-j)^{\frac{3}{2}} f_{n-j} - g \right| + \sum_{j=1}^N \left\{ \left( \frac{n}{n-j} \right)^{\frac{3}{2}} - 1 \right\} T_{j-1} R |g|.$$

Using Fatou's lemma, we see

$$(3.6) \quad \limsup_{n \rightarrow \infty} |K_n - SRg| \leq \sum_{j=0}^{N-1} T_j R \bar{g} + \sum_{j=N}^{\infty} T_j R |g|.$$

Here we note that  $T_j R \bar{g}$  is well-defined. Applying Lemmas 5, 6 and (3.2) to  $J_n$ ,

$$\|J_n\| \leq C^2 \|R\| \|f\| n^{\frac{3}{2}} \sum_{j=N+1}^{n-N} (n-j)^{-\frac{3}{2}} (j-1)^{-\frac{3}{2}}.$$

Using (2) of Lemma 3, we have

$$(3.7) \quad \limsup_{n \rightarrow \infty} \|J_n\| \leq 2C^2 \|R\| \|f\| \sum_{j=N}^{\infty} j^{-\frac{3}{2}}.$$

Collecting (3.4)-(3.7) and letting  $N \rightarrow \infty$ , we have  $0 \leq \bar{g} \leq SR\bar{g}$ . Thus  $\|\bar{g}\| \leq \|SR_t\| \|\bar{g}\|$ . Since  $t \in [0, c_1)$ , this estimate implies  $\|\bar{g}\| = 0$ , i.e.,  $\bar{g} \equiv 0$ , which shows the following: If  $0 \leq t < c_1$ , then for each  $f \in C(-\delta)$  and  $x \leq K$ ,

$$(3.8) \quad \lim_{n \rightarrow \infty} n^{\frac{3}{2}} A_n^{(t)} f(x) = (1 + SR_t)^{-1} Q (1 + R_t S)^{-1} f(x).$$

If  $c_1 = \infty$ , this completes the proof of Main Lemma. We consider the case of  $c_1 < \infty$ . Let arbitrary  $t_0 \in (0, c_1)$  be fixed. For  $f \in C(-\delta)$  and  $x \leq K$ , set

$$\bar{T}_n f(x) = A_n^{(t_0)} f(x) \quad n \geq 0 \text{ and} \quad \bar{S} = \sum_{n=0}^{\infty} \bar{T}_n.$$

Since  $0 \leq \bar{T}_n f(x) \leq T_n f(x)$  if  $f \geq 0$ , everything of the above is well-defined. For  $f \in C(\lambda)$  and  $t \geq 0$ , set

$$\bar{R}_t f(x) = E_x \left[ f(\xi) e^{-t_0 W(\xi)} \{1 - e^{-t W(\xi)}\}; \xi \leq K \right].$$

Let  $f \in C(-\delta)$ . In view of (3.8), we define  $\bar{Q}f(x)$  as follows.

$$\bar{Q}f(x) = \lim_{n \rightarrow \infty} n^{\frac{3}{2}} \bar{T}_n f(x) = (1 + SR_{t_0})^{-1} Q (1 + R_{t_0} S)^{-1} f(x).$$

Now exactly the same argument as the above is possible if we replace  $T_n$ ,  $R_t$ ,  $Q$  by  $\bar{T}_n$ ,  $\bar{R}_t$ ,  $\bar{Q}$  respectively. Therefore setting  $c_2 = \sup\{t > 0 : \|\bar{S}\bar{R}_t\| < 1\}$  where  $\bar{S}\bar{R}_t = \bar{S}(\bar{R}_t)$ , we get the following: If  $0 < t \leq c_2$ , then

$$(3.9) \quad \lim_{n \rightarrow \infty} n^{\frac{3}{2}} A_n^{(t+t_0)} f(x) = (1 + \bar{S}\bar{R}_t)^{-1} \bar{Q} (1 + \bar{R}_t \bar{S})^{-1} f(x).$$

On the other hand, we have the following relations from the definitions.

$$\bar{R}_t = R_{t+t_0} - R_{t_0}, \quad \bar{S} = S - SR_{t_0} \bar{S}. \quad (\text{cf. (3.1)})$$

Using the above relations, we get

$$\begin{aligned} (1 + SR_{t_0})(1 + \bar{S}\bar{R}_t) &= (1 + SR_{t_0+t}), \\ (1 + \bar{R}_t \bar{S})(1 + R_{t_0} S) &= (1 + R_{t_0+t} S). \end{aligned}$$

From these identities and the definition of  $\bar{Q}$ , we see that the right hand side of (3.9) is equivalent to  $(1 + SR_{t_0+t})^{-1} Q (1 + R_{t+t_0} S)^{-1} f(x)$ . Hence we can rewrite (3.9) in the following way: If  $0 \leq t < c_1 + c_2$ , then

$$(3.10) \quad \lim_{n \rightarrow \infty} n^{\frac{3}{2}} A_n^{(t)} f(x) = (1 + SR_t)^{-1} Q (1 + R_t S)^{-1} f(x).$$

Comparing  $\bar{S}$ ,  $\bar{R}_t$  with  $S$ ,  $R_t$ , we easily see  $\|\bar{S}\bar{R}_t\| \leq \|SR_t\|$ . Thus  $0 < c_1 \leq c_2$ . From this fact, the method on the above can be iterated infinitely often. Therefore (3.10) holds for all  $t \geq 0$ . This finishes the proof of Main Lemma.  $\square$

#### 4. Proof of Theorem 1

In this section functions  $f$ ,  $h$  and  $W$  are assumed to satisfy the conditions (i), (ii) and (iii) in Theorem 1. We devide the proof into three parts. For any  $K > 0$ , set

$$\begin{aligned} L_n &= E \left[ f(e^{S_n}) h \left( \sum_{i=1}^n W(S_i) \right) \right] \\ &= E \left[ f(e^{S_n}) h \left( \sum_{i=1}^n W(S_i) \right); M_n \leq K \right] + E \left[ f(e^{S_n}) h \left( \sum_{i=1}^n W(S_i) \right); M_n > K \right] \\ &= H_n(K) + J_n(K). \end{aligned}$$

STEP 1. For any  $K > 0$ , the finite limit  $\lim_{n \rightarrow \infty} n^{3/2} H_n(K)$  exists.

Proof. Let  $K > 0$  be fixed. From the conditions (c) and (iii), we can choose two numbers  $\mu$  and  $\delta$  such that  $0 < \delta < \mu < \min(\alpha, \varepsilon)$  and  $E(e^{\delta \xi}) < \infty$ . We easily see  $f(e^x) \in C(-\delta)$  and  $W \in C(-\mu)$ . From the conditions of  $h$ , there exists a sequence of functions  $\{h_m\}_{m \geq 1}$  satisfying  $\sup_{x \geq 0} |h(x) - h_m(x)| \leq m^{-1}$  and  $h_m(x)$  is a finite linear combinations of  $\{e^{-nx}\}_{n \geq 0}$ . Hence by Main Lemma we can define  $C_m$  as follows.

$$C_m = \lim_{n \rightarrow \infty} n^{3/2} E \left[ f(e^{S_n}) h_m \left( \sum_{i=1}^n W(S_i) \right); M_n \leq K \right].$$

It is obvious that  $\{C_m\}_{m \geq 1}$  is a Cauchy sequence. Set  $C_\infty = \lim_{m \rightarrow \infty} C_m$ . Then

$$\begin{aligned} |n^{3/2} H_n(K) - C_\infty| &\leq \frac{1}{m} n^{3/2} E [|f(e^{S_n})|; M_n \leq K] \\ &\quad + \left| n^{3/2} E \left[ f(e^{S_n}) h_m \left( \sum_{i=1}^n W(S_i) \right); M_n \leq K \right] - C_\infty \right|. \end{aligned}$$

Therefore we have

$$\limsup_{n \rightarrow \infty} |n^{3/2} H_n(K) - C_\infty| \leq \frac{a}{m} + |C_m - C_\infty| \rightarrow 0 \quad \text{as } m \rightarrow \infty,$$

where  $a = \lim_{n \rightarrow \infty} n^{3/2} E [|f(e^{S_n})|; M_n \leq K]$ . Thus we obtain the assertion of Step 1.  $\square$

STEP 2.  $\limsup_{K \rightarrow \infty} \limsup_{n \rightarrow \infty} n^{3/2} |J_n(K)| = 0$ .

Proof. From the condition (iii), there exist positive constants  $A$ ,  $B$  and  $K$  which satisfy the following:  $|f(x)| \leq Ax^\alpha$ ,  $|h(x)| \leq Ax^{-\beta}$  for  $x \geq 0$ , and

$W(x) \geq Be^{\eta x}$  for  $x \geq K$ . Hence we have  $|f(e^{S_n})| \leq Ae^{\alpha S_n}$  and the inequality,  $|h(\sum_{i=1}^n W(S_i))| \leq AB^{-\beta}e^{-\beta\eta M_n}$ , holds on  $(M_n > K)$ . Applying these estimates to  $J_n(K)$  and using (2.1), we have

$$\begin{aligned} |J_n(K)| &\leq \text{const} E(e^{\alpha S_n - \beta\eta M_n}; M_n > K) \\ &\leq \text{const} \sum_{j=0}^n \psi_{n-j}(\alpha) \int_K^\infty e^{-bx} du_j(x), \end{aligned}$$

where  $\text{const} = A^2 B^{-\beta}$ ,  $b = \beta\eta - \alpha > 0$ . In view of (1) (2) of Lemma 4, we apply (2) of Lemma 3 to the last term of the above.

$$\limsup_{n \rightarrow \infty} n^{\frac{3}{2}} |J_n(K)| \leq \text{const} \left\{ \sum_{n=0}^{\infty} \psi_n(\alpha) \int_K^\infty e^{-bx} dU(x) + \psi(\alpha) \int_K^\infty e^{-bx} du(x) \right\}.$$

Since the last term goes to 0 as  $K \rightarrow \infty$ , Step 2 is proved.  $\square$

STEP 3. The finite limit  $\lim_{n \rightarrow \infty} n^{3/2} L_n$  exists.

Proof. Without loss of generality, we assume that  $f$  and  $h$  are non-negative. By Step 1 we can define  $H(K) = \lim_{n \rightarrow \infty} n^{3/2} H_n(K)$ . Then  $H(K)$  is a non-decreasing and bounded. Set  $H = \lim_{K \rightarrow \infty} H(K)$ . Since

$$n^{\frac{3}{2}} H_n(K) \leq n^{\frac{3}{2}} L_n \leq n^{\frac{3}{2}} H_n(K) + \sup_{m \geq n} \left( m^{\frac{3}{2}} J_m(K) \right),$$

taking account of Step 2 we easily see that  $n^{3/2} L_n$  goes to  $H$  as  $n \rightarrow \infty$ . Finally Theorem 1 is established under the conditions (a), (b) and (c).  $\square$

REMARK. To prove Theorem 1 under the conditions (a), (c) and (d), we enumerate the modified points. Under the conditions (a) and (d), the local limit theorem

$$\lim_{n \rightarrow \infty} \left[ \sup_{k \in \mathbb{Z}} \left| \sqrt{n} P(S_n = dk) - \frac{d}{\sqrt{2\pi\sigma}} \exp \left\{ -\frac{(dk)^2}{2n\sigma^2} \right\} \right| \right] = 0$$

holds (See e.g. [3]), which implies, for  $\theta > 0$

$$E(e^{-\theta S_n}; S_n > 0) \sim \frac{d}{\sqrt{2\pi\sigma}} \frac{e^{-d\theta}}{1 - e^{-d\theta}} n^{-\frac{1}{2}} \quad \text{as } n \rightarrow \infty.$$

This corresponds to Lemma 2. If  $U$  and  $V$  are defined by

$$\begin{aligned} U(x) &= \frac{d}{\sqrt{2\pi\sigma}} \sum_{0 \leq j \leq i-1} u(dj) \quad (id \leq x < (i+1)d), \quad = 0 \quad (0 \leq x < d), \\ V(x) &= \frac{d}{\sqrt{2\pi\sigma}} \sum_{0 \leq j \leq i} v(dj) \quad (id \leq x < (i+1)d), \end{aligned}$$

then (1) of Lemma 4 holds, and (2) (3) of Lemma 4 hold at every continuous point. Fix  $K \in \mathbb{N}$ . For a function  $f$  on  $D \equiv \{id : i \leq K\}$  and  $a \in \mathbb{R}$ , set  $\|f\|_a = \sup_{x \in D} e^{ax} |f(x)|$  and  $C(a) = \{f : \|f\|_a < \infty\}$ . Then  $x$  in the expectation symbol  $E_x$  takes value on  $D$ . Having these modifications, the proof of Theorem 1 under the conditions (a), (c) and (d) are carried out analogously. We give the following corollary used in the next section.

**Corollary 10.** *If  $f$  is positive on  $(0, \infty)$  and  $h$  is positive on  $[0, \infty)$ , then the finite limit in Theorem 1 is positive.*

Proof. Set  $\nu_n(x) = E[f(e^{S_n}); \sum_{i=1}^n W(S_i) \leq x]$ . Then for any  $t > 0$ ,

$$\int_0^\infty e^{-tx} d\nu_n(x) = E \left[ f(e^{S_n}) \exp \left\{ -t \sum_{i=1}^n W(S_i) \right\} \right].$$

Apply Theorem 1 and the extended continuity theorem for Laplace transform to the above. Then there exists a measure  $\nu$  satisfying the following: For each  $t > 0$ ,

$$\lim_{n \rightarrow \infty} n^{\frac{3}{2}} \int_0^\infty e^{-tx} d\nu_n(x) = \int_0^\infty e^{-tx} d\nu(x) < \infty.$$

On the other hand, (3.10) implies

$$\int_0^\infty e^{-tx} d\nu(x) \geq (1 + SR_t)^{-1} Q(1 + R_t S)^{-1} \hat{f}(0),$$

where  $\hat{f}(x) = f(e^x)$ . Letting  $t \rightarrow 0$ , we see  $\nu(\infty) \geq Q\hat{f}(0) > 0$ . The last inequality follows from Corollary 9. Hence we can choose a positive constant  $x$  such that  $0 < \nu(x) < \infty$  and  $\lim_{n \rightarrow \infty} n^{3/2} \nu_n(x) = \nu(x)$ . By the assumptions of  $h$ , we have

$$E \left[ f(e^{S_n}) h \left( \sum_{i=1}^n W(S_i) \right) \right] = \int_0^\infty h(s) d\nu_n(s) \geq \min_{0 \leq s \leq x} h(s) \nu_n(x),$$

which proves Corollary 10.  $\square$

## 5. Application to a random walk in a random medium

Let  $\{p_i : i \in \mathbb{Z}\}$  be a sequence of i.i.d. random variables with values in  $[0, 1]$ , and let  $\mathcal{F}$  be the  $\sigma$ -field generated by the sequence  $\{p_i\}$ . A random walk in a random medium  $\{p_i\}$  is a sequence of random variables  $\{X_t : t = 0, 1, 2, \dots\}$  which satisfy the following:

$$\begin{aligned} X_0 &= 0, \quad P(X_{t+1} = X_t + 1 | \mathcal{F}, X_0, X_1, \dots, X_t = i) = p_i, \\ P(X_{t+1} &= X_t - 1 | \mathcal{F}, X_0, X_1, \dots, X_t = i) = 1 - p_i. \end{aligned}$$

Set  $\sigma_i = p_i/(1-p_i)$ . In [7] it was shown that the condition  $-\infty \leq E \log \sigma_0 < 0$  implies  $\lim_{t \rightarrow \infty} X_t = -\infty$  a.s. Hence  $\max_{t \geq 0} X_t$  is finite a.s. In [1] Afanas'ev showed the asymptotic behavior of the probability  $P_n \equiv P(\max_{t \geq 0} X_t \geq n)$  under the condition  $E\sigma_0 < 1$  which implies  $-\infty \leq E \log \sigma_0 < 0$  by Jensen's inequality. We consider the same problem under the conditions

$$(5.1) \quad -\infty \leq E \log \sigma_0 < 0, \quad E\sigma_0 < \infty.$$

**Theorem 2.** *Let  $\sigma_0$  satisfy (5.1).*

(1) *If  $E\sigma_0 \log \sigma_0 < 0$ , then*

$$P_n \sim c_1 (E\sigma_0)^n \quad \text{as } n \rightarrow \infty.$$

(2) *If  $E\sigma_0 \log \sigma_0 = 0$  and  $E\sigma_0 \log^2 \sigma_0 < \infty$ , then*

$$P_n \sim c_2 n^{-1/2} (E\sigma_0)^n \quad \text{as } n \rightarrow \infty.$$

(3) *If  $E\sigma_0 \log \sigma_0 > 0$  and the distribution of  $\log \sigma_0$  is continuous or supported on a lattice, then*

$$P_n \sim c_3 n^{-3/2} \gamma^n \quad \text{as } n \rightarrow \infty,$$

where  $\gamma = \min_{0 \leq t \leq 1} E\sigma_0^t$  and  $c_1, c_2, c_3$  are positive numbers independent of  $n$ . In case of (1) and (2),  $E\sigma_0 < 1$  holds automatically.

Proof. Set  $\zeta_i = \log \sigma_{i-1}$ ,  $\widehat{S}_n = \sum_{i=1}^n \zeta_i$ ,  $n \geq 1$ . Then we have the following identity (See [1]).

$$P_n = E\rho \left( \rho + \sum_{i=1}^n e^{-\widehat{S}_i} \right)^{-1} \quad \text{where} \quad \rho = 1 + \sum_{i=1}^{\infty} \sigma_{-1}\sigma_{-2}\cdots\sigma_{-i} < \infty \quad \text{a.s.}$$

Since  $\{p_i\}$  are i.i.d.,  $\{\zeta_i\}_{i \geq 1}$  are i.i.d. and independent of  $\rho$ . Set  $h(x) = E\rho(\rho+x)^{-1}$ . Then we can rewrite  $P_n$  using  $h(x)$ .

$$P_n = Eh \left( \sum_{i=1}^n e^{-\widehat{S}_i} \right).$$

Set  $\phi(t) = E \exp t\zeta_1$ . The condition (5.1) ensures that  $\phi(t)$  is well defined on  $[0, 1]$ ,  $\phi''(t) > 0$  on  $(0, 1)$  and  $\lim_{t \searrow 0} \phi'(t) = E \log \sigma_0 < 0$ . From the assumptions of (1) and (2),  $\lim_{t \nearrow 1} \phi'(t) = E\sigma_0 \log \sigma_0 \leq 0$ . Since  $\phi'(t)$  is monotone increasing on  $(0, 1)$ , these estimates imply  $\phi(1) = E\sigma_0 < 1$ . Thus (1) and (2) are reduced to Afanas'ev's theorem in [1].

In case of (3), we have  $\lim_{t \searrow 0} \phi'(t) < 0$  and  $\lim_{t \nearrow 1} \phi'(t) > 0$ . Therefore there exists a unique point  $\tau \in (0, 1)$  where  $\phi(t)$  attains its minimum  $\gamma < 1$ , i.e.,  $\phi'(\tau) = 0$  and  $\phi(\tau) = \gamma < 1$ . Set a proper distribution function

$$G(x) = \frac{1}{\gamma} \int_{-\infty}^x e^{\tau y} dP(\zeta_1 \leq y).$$

Let  $\{\xi_i\}_{i \geq 1}$  be i.i.d. and  $-\xi_1$  has the distribution function  $G(x)$ . By the assumption of (3) and the definition of  $G(x)$ , we see that  $\xi_1$  satisfies the conditions in Section 1. In fact we obtain

$$Ee^{\theta \xi_1} = Ee^{(\tau-\theta)\xi_1} = \frac{\phi(\tau-\theta)}{\gamma} \quad \text{and} \quad E\xi_1 = -\frac{\phi'(\tau)}{\gamma} = 0.$$

The former of the above shows that the Laplace transform of  $\xi_1$  exists in some neighborhood of the origin and  $E|\xi_1|^k < \infty$  for all  $k \in \mathbb{N}$ . Put  $S_0 = 0$  and  $S_n = \sum_{i=1}^n \xi_i$ ,  $n \geq 1$ . We express  $P_n$  by  $S_i$ .

$$(5.2) \quad P_n = Eh \left( \sum_{i=1}^n e^{-\widehat{S}_i} \right) = \gamma^n E \left[ e^{\tau S_n} h \left( \sum_{i=1}^n e^{S_i} \right) \right].$$

Let a number  $\beta$  satisfying  $\tau < \beta < 1$  and  $\phi(\beta) < 1$  be fixed. By the definitions of  $\rho$  and  $\phi(t)$ ,  $B \equiv E\rho^\beta \leq \sum_{n=0}^{\infty} \phi(\beta)^n < \infty$ . Using Chebyshev's inequality,  $P(\rho > y) \leq By^{-\beta}$  for  $y > 0$ . Applying integration by parts to  $h(x)$  and using the above result, we have

$$\begin{aligned} h(x) &= \int_0^\infty \frac{x}{(y+x)^2} P(\rho > y) dy \\ &\leq \int_0^\infty \frac{x}{(y+x)^2} By^{-\beta} dy \\ &= Bx^{-\beta} \int_0^\infty \frac{t^{-\beta}}{(1+t)^2} dt. \end{aligned}$$

Hence for  $x > 0$ ,

$$(5.3) \quad 0 < h(x) \leq \frac{\pi\beta}{\sin \pi\beta} Bx^{-\beta}.$$

Taking account of (5.2) and (5.3), we can apply Theorem 1 and Corollary 10 to  $P_n \gamma^{-n}$ . Consequently we have the following positive finite limit

$$c_3 \equiv \lim_{n \rightarrow \infty} n^{3/2} \gamma^{-n} P_n = \lim_{n \rightarrow \infty} n^{3/2} E \left[ e^{\tau S_n} h \left( \sum_{i=1}^n e^{S_i} \right) \right].$$

This finishes the proof of (3). □

ACKNOWLEDGEMENT. I would like to express my appreciation to Professor S. Kotani for his advice and encouragement.

---

### References

- [1] V.I. Afanas'ev: *On a maximum of a transient random walk in random environment*, Theory Probab. Appl. **35** (1990), 205–215.
- [2] R.R. Bahadur and R.R. Rao: *On deviation of the sample mean*, Ann. Math. Statist. **31** (1960), 1015–1027.
- [3] W. Feller: *An Introduction to Probability Theory and Its Applications*, volume II, second edition. New York: John Wiley & Sons, (1971).
- [4] D.L. Iglehart: *Random walks with negative drift conditioned to stay positive*, J. Appl. Probab. **11** (1974), 742–751.
- [5] K. Kawazu and H. Tanaka: *On the maximum of a diffusion process in a drifted Brownian environment*, Seminaire de probabilités XXVII (Lect. Notes Math. **1557**, 78–85), Springer-Verlag Berlin Heidelberg (1993).
- [6] N.U. Prabhu: *Stochastic Storage Processes*. New York, Springer-Verlag, (1980).
- [7] F. Solomon: *Random walks in a random environment*, Ann. Probab. **3** (1975), 1–31.
- [8] N. Veraverbeke and J.L. Teugels: *The exponential rate of convergence of the distribution of the maximum of a random walk*, part II. J. Appl. Probab. **13** (1976), 733–740.
- [9] M. Yor: *On some exponential functionals of Brownian motion*, Adv. Appl. Probab. **24** (1992), 509–531.

Department of Mathematics  
 Osaka University  
 Toyonaka, Osaka 560  
 Japan