

Title	GLOBAL QUOTIENTS AMONG TORIC DELIGNE-MUMFORD STACKS
Author(s)	Harada, Megumi; Krepski, Derek
Citation	Osaka Journal of Mathematics. 2015, 52(1), p. 237-269
Version Type	VoR
URL	https://doi.org/10.18910/57679
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

GLOBAL QUOTIENTS AMONG TORIC DELIGNE–MUMFORD STACKS

MEGUMI HARADA and DEREK KREPSKI

(Received February 12, 2013, revised September 25, 2013)

Abstract

This work characterizes global quotient stacks—smooth stacks associated to a finite group acting on a manifold—among smooth quotient stacks $[M/G]$, where M is a smooth manifold equipped with a smooth proper action by a Lie group G . The characterization is described in terms of the action of the connected component G_0 on M and is related to (stacky) fundamental group and covering theory. This characterization is then applied to smooth toric Deligne–Mumford stacks, and global quotients among toric DM stacks are then characterized in terms of their associated combinatorial data of stacky fans.

1. Introduction

This note has two parts: first, we consider topological properties of smooth orbifolds that describe obstructions to being a *global (resp. discrete) quotient*¹—i.e. equivalent (in a sense made more precise below) to an orbifold associated to a finite (resp. discrete) group action on a smooth manifold, and second, we apply our general results on global quotients to toric Deligne–Mumford stacks. Recall that an orbifold structure, considered from the classical viewpoint (originating in the work of Satake [25]), can be described via local charts, each of which are quotients U/Γ of a linear action of a finite group Γ on an open subset U of Euclidean space. The point is that for a general orbifold, these local charts do not necessarily ‘patch together’ into a global finite group action on a smooth manifold, but for a global quotient, they do.

It is worth emphasizing the following few points at the outset. Firstly, in this paper, by a ‘smooth (resp. topological) orbifold’ we mean a *smooth (resp. topological) stack*, or more precisely, a *stack in the category Diff (resp. Top)*. (In our exposition we have attempted to make the language of stacks accessible to a non-expert audience (more on this below), although in no way do we aim to be a textbook. We suggest [4, 17, 19] for more about stacks from a topologist’s point of view; a complete beginner may wish to start with [6] or [7].) In particular, the notion of ‘equivalence’ in the previous paragraph is a (weak) equivalence of the underlying categories

2010 Mathematics Subject Classification. Primary 57R18, 53D20; Secondary 14M25, 14D23.

MH is partially supported by an NSERC Discovery Grant, an NSERC University Faculty Award, and an Ontario Ministry of Research and Innovation Early Researcher Award.

¹In the literature, global quotients are also called *good* or *developable* orbifolds.

of the stacks. It is important to note that such an equivalence is more restrictive than a homeomorphism of underlying topological spaces; this is because such a homeomorphism does not retain any ‘stacky’ information relating to the orbifold singularities. A basic class of examples illustrating this distinction are the so-called *weighted projective stacks*, i.e. $\mathbb{P}(a_0, \dots, a_n) = [(\mathbb{C}^{n+1} \setminus \{0\})/\mathbb{C}^*]$, where \mathbb{C}^* acts diagonally on $\mathbb{C}^{n+1} \setminus \{0\}$ with weights $a_0, \dots, a_n \in \mathbb{Z}_+$. (Here we follow the convention in the literature and denote by $[X/G]$ the *stack* associated to a G -action on a space X ; the topological quotient (orbit) *space* is denoted X/G .) Although the underlying orbit space $(\mathbb{C}^{n+1} \setminus \{0\})/\mathbb{C}^*$ is homeomorphic to the orbit space of a finite group action on complex projective space $\mathbb{C}P^n$ (see [15] for details), the stack $\mathbb{P}(a_0, \dots, a_n)$ is not equivalent *as a stack* to a global quotient (except in the trivial case when all of the weights are equal to 1 and $\mathbb{P}(1, \dots, 1) = \mathbb{C}P^n$ is a smooth manifold). This follows from Theorem 4.10 below, but is also well-known—see e.g. [1].

Secondly, we emphasize that we restrict ourselves throughout this paper to orbifolds arising as quotient stacks $[X/G]$, where G is a Lie group acting smoothly and properly on a smooth manifold X . It is worth noting that all *reduced*, or *effective* orbifolds—orbifolds whose local isotropy groups act effectively—are known to be of this type by a frame-bundle construction (see, for example, [1]), so this is not a very restrictive condition in practice. Moreover, as we already mentioned, the main application we have in mind of our Theorem 4.10 is to the beautiful class of quotient stacks known as toric Deligne–Mumford stacks, which are stack analogues of smooth toric varieties.

Thirdly, we recall that in his foundational work [21], Noohi deals with topological stacks and the theory of (stacky) fundamental groups and coverings in a very general framework. We owe much to [21] for both the mathematical content and the exposition in Sections 2, 3, and 4. Indeed, our Theorem 4.10, quoted below, can be viewed as an extension of [21, Theorem 18.24] in the special case of quotient stacks. By restricting to quotient stacks, we are able to connect aspects of stacky algebraic topology with a distinctly geometric (and classical) condition on a Lie group acting on a manifold; in particular, our proofs are our own. In our exposition we have attempted to explicitly preserve the classical perspective and language as much as possible; we hope this serves to illustrate to a broad audience the appeal of the stack perspective, and to further elucidate the insights of [21] in a setting that is common in differential geometry, namely Lie groups acting on manifolds.

With these points in mind we now state our main general result on stacks arising as global quotients (Theorem 4.10 in Section 4). We refer to Section 2.2 for the definition of the inertia homomorphism in (3), and Section 4 for a discussion of (stacky) covering projections appearing in (4). For now, the reader may keep in mind that the *inertia groups* I_x mentioned below can be identified with isotropy groups $\text{Stab}(p) \subset G$ of certain points $p \in X$.

Theorem 4.10. *Let X be a simply connected manifold, equipped with a smooth proper action of a Lie group G . Let $G_0 \subset G$ denote the connected component of the*

identity element of G , and I_x the inertia group of $x \in [X/G]$. The following statements are equivalent.

- (1) $[X/G]$ is equivalent to a discrete quotient.
- (2) G_0 acts freely on X .
- (3) For all x in X , the inertia homomorphism $\omega_x: I_x \rightarrow \pi_1([X/G], x)$ is injective.
- (4) The (stacky) universal cover of $[X/G]$ is equivalent to a smooth manifold.

Though conditions (1), (3), and (4) are known to be equivalent by [21, Theorem 18.24], we provide a different proof of these equivalencies for the case of quotient stacks by showing each of these conditions is in turn equivalent to (2). The distinctly geometric appeal of condition (2), in comparison to the topological nature of conditions (3) and (4), also carries over in our interpretation of this result in the setting of smooth toric DM stacks.

We now take a moment to briefly recall the context of our discussion of *toric Deligne–Mumford (DM) stacks* in Section 5. In their foundational paper [5], Borisov, Chen, and Smith introduce the notion of a *stacky fan*, the combinatorial data from which one constructs a toric Deligne–Mumford (DM) stack using an analogue of the Cox quotient construction in algebraic geometry. In [9], an abstract definition of a smooth toric DM stack was given, which was shown to be compatible with the construction of Borisov, Chen, and Smith [5] (see also [14], [22], [10, 11] for related approaches). From the symplectic geometry perspective, Lerman and Malkin [18] gave a definition of symplectic toric DM stacks (in the smooth category Diff), offering a modern perspective on symplectic toric orbifolds. In fact, parallel to the classical theory of toric varieties, there is a subfamily of toric DM stacks—namely those toric DM stacks whose underlying fan is polytopal—that admit a construction from symplectic geometry via *stacky polytopes*, using an adaptation of the construction of Borisov, Chen, and Smith (see [24]). In particular, from this construction, this subfamily can be seen to give examples of symplectic toric DM stacks. (In earlier work, Lerman and Tolman extended the Delzant classification of (compact) symplectic toric manifolds to symplectic toric orbifolds; in the orbifold case, the classification is by ‘labelled polytopes’—i.e. polytopes with positive integer labels attached to each facet [16].) In this manuscript, we interpret our analysis of orbifolds as global quotients in this class of examples; our results are explicit and combinatorial, stated in terms of the stacky fan/polytope.

In order to state our main results for smooth toric DM stacks, recall that a stacky fan is a triple (N, Σ, β) consisting of a finitely generated \mathbb{Z} -module N , a simplicial fan Σ in $N \otimes \mathbb{R}$ with n rays ρ_1, \dots, ρ_n , and a homomorphism $\beta: \mathbb{Z}^n \rightarrow N$ satisfying certain conditions (see Definition 5.1). By interpreting Theorem 4.10 in this case, we can characterize global quotients among toric DM stacks in terms of their underlying stacky fans. Let N' denote the image of β , and for a cone σ in Σ , let N_σ denote $\text{span}\{\beta(e_i) \mid \rho_i \text{ is a ray in } \sigma\}$, where e_i denotes the i -th standard basis vector in \mathbb{Z}^n . In this context, condition (2) of Theorem 4.10 results in the following corollary, which

characterizes the stacky fans yielding global quotients. We note that the equivalence of condition (2) of Theorem 4.10 and the combinatorial condition stated in Corollary 5.7, in the context of toric DM stacks, was first proved via a different method—a combinatorial analysis of the local isotropy groups—in joint work of the authors with Goldin and Johanssen; this approach is described in [12]. We also note that in [23], the authors study quasi-toric orbifolds using techniques from toric topology, and obtain similar results describing universal covers of quasi-toric orbifolds in that framework.

Corollary 5.7. *Let (N, Σ, β) be a stacky fan, and let \mathcal{X} denote the corresponding toric DM stack. Then \mathcal{X} is equivalent to a (finite) global quotient if and only if $N' = N_\sigma$ for all maximal cones σ in Σ .*

In addition to Corollary 5.7, we interpret the other obstructions appearing in conditions (3) and (4) of Theorem 4.10 for toric DM stacks in terms of stacky fans. Specifically, using results in [12], we see that the inertia homomorphisms of condition (3) can be identified with very natural homomorphisms defined in terms of the stacky fan (Proposition 5.3). Additionally, we describe the universal cover of a toric DM stack in terms of its stacky fan in Proposition 5.5. It is interesting to note that the universal cover of a toric DM stack is given in terms of its stacky fan data, and is hence also a toric DM stack arising from a stacky fan. In particular, global quotients among toric DM stacks are quotients of toric *manifolds* by finite group actions. We expect such a nice description to be useful in computations of orbifold/stacky invariants, particularly for global quotients (cf. [2], [8], for example).

The contents of this paper are as follows. After a brief discussion of stacks and fundamental groups of stacks in Section 2, we specialize to quotient stacks in Section 3, where we describe in detail the fundamental group, and inertia homomorphism of quotient stacks. In Section 4, we describe the universal cover of a quotient stack and prove Theorem 4.10 stated above; analogous results for connected but non-simply connected X are also explored in this section. In Section 5, we turn our attention to toric DM stacks, where we describe the inertia homomorphism (Section 5.3) and universal cover (Section 5.4) of toric DM stacks. We also verify an expected relation between the symplectic volumes (in the stacky polytope case) of a symplectic toric DM stack and its universal cover, and the corresponding volumes of the underlying polytopes. We conclude with some examples in Section 5.5.

2. Preliminaries

This section establishes notation and collects some facts about stacks. We mainly follow the notation of [21] and [4].

2.1. Stacks. We will mainly work over the base category Diff (smooth manifolds and smooth maps), although occasionally we shall work over Top (topological spaces

and continuous maps). To streamline some of the discussion below, we use \mathbf{Sp} to denote either \mathbf{Diff} or \mathbf{Top} , and we refer to the corresponding objects simply as spaces and the morphisms as maps.

For a space M , let \underline{M} denote its associated stack, with objects $\{f: E \rightarrow M\}$, i.e. the collection of maps in \mathbf{Sp} with target M , and with morphisms $\{g: E \rightarrow E' \mid f' \circ g = f\}$. Given a map $F: M \rightarrow M'$, we write $\underline{F}: \underline{M} \rightarrow \underline{M}'$ for the induced morphism of stacks. Fix a terminal object \star in \mathbf{Sp} . For a choice of point p in a space M , let \underline{p} be the corresponding point in \underline{M} (i.e. by abuse of notation, p also denotes the map $\star \mapsto p \in M$ and \underline{p} the induced morphism of stacks). More generally, recall that a point x in a stack \mathcal{X} is a morphism $x: \star \rightarrow \mathcal{X}$.

A morphism $F: \mathcal{Y} \rightarrow \mathcal{X}$ of stacks is *representable* if for every morphism $\underline{W} \rightarrow \mathcal{X}$ whose source is (the stack associated to) a space, the fiber product $\mathcal{Y} \times_{\mathcal{X}} \underline{W}$ is equivalent to (the stack associated to) a space. In this case, we shall call the induced map $\mathcal{Y} \times_{\mathcal{X}} \underline{W} \rightarrow \underline{W}$ a representative of F . Notice that the representative of a representable map is (induced by) a map of spaces.

DEFINITION 2.1 ([19, Definition 66]). A stack \mathcal{X} over \mathbf{Sp} is *locally representable* if there exists a representable epimorphism of stacks $p: \underline{X} \rightarrow \mathcal{X}$. In this case, p is called a *presentation* of \mathcal{X} .

REMARK 2.2. When $\mathbf{Sp} = \mathbf{Top}$, a locally representable stack over \mathbf{Top} is called a *pretopological stack* [21].

Many properties of representable morphisms of stacks are defined by the corresponding properties of their representatives, which are maps of spaces. For example we have the following (cf. [21]).

DEFINITION 2.3. A morphism $\mathcal{Y} \rightarrow \mathcal{X}$ of pretopological stacks in \mathbf{Top} is a *covering projection* if it is representable and if every representative is a covering projection.

Several other properties of maps in \mathbf{Sp} can be defined similarly as properties of representable maps of stacks (e.g. proper, closed, submersion, etc. see [19, Section 3.3]). For now, we simply recall that an important necessary condition for a property of maps to be defined in this manner (i.e. analogous to Definition 2.3) is that this property be stable under taking pullbacks. (If $\mathbf{Sp} = \mathbf{Diff}$, it is additionally required that the property be stable under pullbacks via submersions.) We record the following definition from [19], which connects our point of view with that of orbifolds and orbifold charts.

DEFINITION 2.4. A locally representable stack \mathcal{X} over \mathbf{Diff} is called an *orbifold* or a *Deligne–Mumford stack* if it admits a presentation $p: \underline{X} \rightarrow \mathcal{X}$ that is étale and if the diagonal map $\Delta: \mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ is proper.

In the literature, the term orbifold is at times reserved for smooth Deligne–Mumford stacks with trivial global (generic) stabilizer (inertia group), what is elsewhere called an *effective* or *reduced* orbifold. The above definition allows for non-trivial global stabilizer.

An important instance of the above definition is the following. Let X be a manifold equipped with a smooth action by a compact Lie group G . If the action is locally free (i.e. with finite isotropy groups), then the associated quotient stack $[X/G]$ (see Section 3) is an orbifold. This will be our main source of examples.

We shall at times wish to view a *geometric* stack over Diff —a locally representable stack whose presentation is a surjective submersion—as a stack over Top . To say this more precisely, recall that given a geometric stack with presentation $\overline{X}_0 \rightarrow \mathcal{X}$, there is a Lie groupoid $\mathcal{G} = (X_1 \rightrightarrows X_0)$ whose associated stack $B\mathcal{G}$ (see [17, Section 4]) is equivalent to \mathcal{X} . (In this case, we say that \mathcal{G} is a *groupoid presentation* for \mathcal{X} .) Considering the Lie groupoid \mathcal{G} as a groupoid object in Top , we view $B\mathcal{G}$, and hence \mathcal{X} , as a stack over Top .

2.2. Fundamental group of topological stacks. Following the work of Noohi [21], we now recall some of the main definitions surrounding the fundamental group of a topological stack that we later interpret more explicitly for quotient stacks $\mathcal{X} = [X/G]$ as above.

In this subsection, we work over Top . Let \mathcal{X} and \mathcal{Y} be stacks and x, y points in \mathcal{X}, \mathcal{Y} respectively. Recall that a pointed map $(f, \alpha): (\mathcal{Y}, y) \rightarrow (\mathcal{X}, x)$ of stacks consists of a morphism $f: \mathcal{Y} \rightarrow \mathcal{X}$ and a 2-isomorphism $\alpha: x \rightsquigarrow f(y) = f \circ y$, where as in [21] we sometimes use the symbol \rightsquigarrow for a 2-isomorphism of points. (When we do not wish to emphasize that a 2-isomorphism is between two points, we shall instead use the symbol \Rightarrow .)


More generally, one may consider the more general situation of a pair of stacks $(\mathcal{X}, \mathcal{A})$, consisting of stacks \mathcal{X} and \mathcal{A} together with a given morphism $i: \mathcal{A} \rightarrow \mathcal{X}$. A morphism $(\mathcal{Y}, \mathcal{B}) \rightarrow (\mathcal{X}, \mathcal{A})$ between such pairs consists of a pair of morphisms $f: \mathcal{Y} \rightarrow \mathcal{X}$, $f': \mathcal{B} \rightarrow \mathcal{A}$ together with a 2-isomorphism $\alpha: i \circ f' \Rightarrow f \circ j$. Such a morphism (f, f') is often represented by a 2-cell,

$$\begin{array}{ccc} \mathcal{B} & \xrightarrow{f'} & \mathcal{A} \\ j \downarrow & \swarrow \alpha & \downarrow i \\ \mathcal{Y} & \xrightarrow{f} & \mathcal{X} \end{array}$$


In this paper, we shall work only with pairs $(\mathcal{X}, \mathcal{A})$ where the second factor \mathcal{A} is a topological space (often simply a point). In this case, the notion of 2-isomorphism simplifies, as we recall next.

For maps $(f, f'), (g, g'): (\mathcal{Y}, \mathcal{B}) \rightarrow (\mathcal{X}, \mathcal{A})$ of pairs, a 2-isomorphism consists of a pair of 2-isomorphisms $\phi: f \Rightarrow g$ and $\phi': f' \Rightarrow g'$, such that the following 2-cell

(with 2-isomorphisms $\alpha: i \circ f' \Rightarrow f \circ j$ and $\beta: i \circ g' \Rightarrow g \circ j$ suppressed) commutes.



If both \mathcal{B} and \mathcal{A} are topological spaces, this forces $f' = g'$ and $\phi' = \text{id}$. Moreover, if $\mathcal{B} = \mathcal{A} = \star$, then $f' = g' = \text{id}$; therefore, for pointed maps, $(f, \alpha), (g, \beta): (\mathcal{Y}, y) \rightarrow (\mathcal{X}, x)$, a (pointed) 2-isomorphism $\epsilon: (f, \alpha) \Rightarrow (g, \beta)$ is a 2-isomorphism $\epsilon: f \Rightarrow g$ such the following 2-cell (again, with α and β suppressed) commutes:



In other words, the induced 2-isomorphism $\epsilon(y): f(y) \rightsquigarrow g(y)$ satisfies $\epsilon(y)\alpha = \beta$. (Here $\epsilon(y) = \epsilon * \text{id}_y$, the ‘horizontal’ composition of 2-morphisms, while $\epsilon(y)\alpha$ denotes the ‘vertical’ composition of 2-morphisms as in [17].)

Let M be a topological space with a chosen point p and (\underline{M}, p) its associated pointed stack. We begin with a remark regarding morphisms (resp. pointed morphisms) from \underline{M} (resp. (\underline{M}, p)) to a stack \mathcal{X} (resp. (\mathcal{X}, x)).

REMARK 2.5. Let M be an object in Top . A morphism $f: \underline{M} \rightarrow \mathcal{X}$ determines an object E_f in \mathcal{X} (over M) given by evaluation at the identity $E_f := f(\text{id}_M: M \rightarrow M)$. In fact, evaluation at the identity defines a functor ev_M from $\text{HOM}(\underline{M}, \mathcal{X})$ to \mathcal{X}_M which is surjective on objects and fully faithful (Proposition 2.20 in [4]). That is, given an object E in \mathcal{X} over M there is a morphism $g: \underline{M} \rightarrow \mathcal{X}$ such that $E_g = E$. Moreover, given an isomorphism $\varphi: E_g \rightarrow E_f$ in \mathcal{X}_M , there exists a unique 2-isomorphism, which we will also denote $\varphi: g \Rightarrow f$, whose evaluation at the identity is φ . Thus, up to canonical 2-isomorphism a map $\underline{M} \rightarrow \mathcal{X}$ is determined by the data of an object E in \mathcal{X} over M .

A similar reasoning applies to pointed maps. Fix points p in \underline{M} and x in \mathcal{X} . Let $E_x = x(\text{id}_\star: \star \rightarrow \star)$ in \mathcal{X}_\star . A pointed map $(f, \alpha): (\underline{M}, p) \rightarrow (\mathcal{X}, x)$ determines objects E_f and $E_{f(p)}$ (over M and \star , respectively), a morphism $E_{f(p)} \rightarrow E_f$ (over $p: \star \rightarrow M$) and an isomorphism $\alpha: E_x \rightarrow E_{f(p)}$. Conversely, as in the proof of Proposition 2.20 in

[4], given objects E in \mathcal{X}_M and E_0 in \mathcal{X}_\star and a morphism $E_0 \rightarrow E$ over $p: \star \rightarrow M$, there is a morphism $f: \underline{M} \rightarrow \mathcal{X}$ such that $E_f = E$, $E_{f(p)} = E_0$ and the induced morphism $E_{f(p)} \rightarrow E_f$ is the given one. Since ev_\star is fully faithful, a choice of isomorphism $\alpha: E_x \rightarrow E_0$ then determines a unique 2-isomorphism, which we will also denote $\alpha: x \rightsquigarrow f(p)$, and hence a pointed map (f, α) whose evaluation at the identity yields the data $E_0 \rightarrow E$ with the isomorphism α .

Moreover, given pointed maps (f, α) and (g, β) and isomorphisms $\varphi: E_g \rightarrow E_f$ and $\psi: E_{g(p)} \rightarrow E_{f(p)}$ such that the diagram

$$\begin{array}{ccccc}
 & & E_{g(p)} & \longrightarrow & E_g \\
 E_x & \xrightarrow{\alpha} & \downarrow \psi & & \downarrow \varphi \\
 & \xrightarrow{\beta} & E_{f(p)} & \longrightarrow & E_f
 \end{array}$$

commutes, there is a unique 2-isomorphism $\epsilon: (g, \beta) \Rightarrow (f, \alpha)$ with $\text{ev}_M(\epsilon) = \varphi$ and $\text{ev}_\star(\epsilon(p)) = \psi$. As in the unpointed case above, we conclude that a pointed map $(\underline{M}, \underline{p}) \rightarrow (\mathcal{X}, x)$ is determined up to unique 2-isomorphism by objects E in \mathcal{X}_M and E_0 in \mathcal{X}_\star , a morphism $E_0 \rightarrow E$ over $p: \star \rightarrow \underline{M}$, and a choice of 2-isomorphism $\alpha: E_0 \rightarrow E_x$.

Let I denote the unit interval $[0, 1]$. Recall that a *homotopy* of pointed maps $(f, \alpha), (g, \beta): (\underline{M}, \underline{p}) \rightarrow (\mathcal{X}, x)$ is triple $(H, \epsilon_0, \epsilon_1)$, where H is a map $(I \times \underline{M}, I \times \star) \rightarrow (\mathcal{X}, x)$, which in this case is a morphism $H: I \times \underline{M} \cong \underline{I} \times \underline{M} \rightarrow \mathcal{X}$ with a 2-isomorphism for the diagram,

$$\begin{array}{ccc}
 \underline{I} \times \star \cong \underline{I} & \longrightarrow & \star \\
 i_1 \downarrow & \swarrow & \downarrow x \\
 \underline{I} \times \underline{M} & \xrightarrow{H} & \mathcal{X}
 \end{array}$$

together with a pair of pointed 2-isomorphisms $\epsilon_0: f \Rightarrow H_0$ and $\epsilon_1: H_1 \Rightarrow g$. Here $i_1 = (\text{id}, p): I \rightarrow I \times M$ is the inclusion into the first factor, and H_0 and H_1 denote the pointed maps induced by restriction to $\{0\} \times \underline{M}$ and $\{1\} \times \underline{M}$ respectively. When such a homotopy exists, we shall sometimes say that (f, α) is homotopic to (g, β) or that $(H, \epsilon_0, \epsilon_1)$ is a homotopy from (f, α) to (g, β) . This induces an equivalence relation on the set of pointed maps $(\underline{M}, \underline{p}) \rightarrow (\mathcal{X}, x)$ and we denote by $[(\underline{M}, \underline{p}), (\mathcal{X}, x)]$ the resulting set of equivalence classes (i.e. the set of homotopy classes of pointed maps).

That the relation above is reflexive and symmetric is easily verified. To show it is also transitive, we must be able to ‘compose’ homotopies—that is, given a homotopy $(H, \epsilon_0, \epsilon_1)$ from (f, α) to (g, β) and a homotopy $(H', \epsilon'_0, \epsilon'_1)$ from (g, β) to (h, γ) , construct a homotopy $(H'', \epsilon''_0, \epsilon''_1)$ from (f, α) to (h, γ) . This is proved in Lemma 17.4 in [21] in greater generality. (In the case of pointed maps whose domain is the associated stack of a space M , we may observe the composition of homotopies more readily in light of

Remark 2.5. Indeed, the objects E_H and $E_{H'}$ in $\mathcal{X}_{I \times M}$ can be glued together to form the required object $E_{H''}$ that yields the desired composition of homotopies.)

As in [21, Remark 17.3]), observe that a 2-isomorphism $\epsilon: (f, \alpha) \Rightarrow (g, \beta)$ of pointed maps induces a homotopy. Indeed, let H be the composition $\underline{I} \times \underline{M} \xrightarrow{\text{pr}_2} \underline{M} \xrightarrow{f} \mathcal{X}$, with 2-isomorphism as indicated by the diagram

$$\begin{array}{ccccc}
 \underline{I} & \xrightarrow{\quad} & \star & & \\
 \downarrow \text{id}_1 & \swarrow \text{id} & \downarrow p & \searrow x & \\
 \underline{I} \times \underline{M} & \xrightarrow{\text{pr}_2} & \underline{M} & \xrightarrow{f} & \mathcal{X} \\
 & & \swarrow \alpha & & \\
 & & \mathcal{X} & &
 \end{array}$$

(i.e. the 2-isomorphism $(\text{id}_f * \text{id})(\alpha * \text{id}_\star)$), $\epsilon_0 = \text{id}$ and $\epsilon_1 = \epsilon$. This observation proves the following lemma.

Lemma 2.6. *Let p be a point in a space M and let (\mathcal{X}, x) be a pointed stack. Let (f, α) and (g, β) be pointed maps $(\underline{M}, p) \rightarrow (\mathcal{X}, x)$. If there exists a 2-isomorphism of pointed maps $(f, \alpha) \Rightarrow (g, \beta)$, then the homotopy classes of (f, α) and (g, β) are equal.*

REMARK 2.7. It follows from Remark 2.5 that the homotopy class of a pointed map $(\underline{M}, p) \rightarrow (\mathcal{X}, x)$ is determined by specifying objects E in \mathcal{X}_M and E_0 in \mathcal{X}_\star , along with a morphism $E_0 \rightarrow E$ over $p: \star \rightarrow M$ and an isomorphism $\alpha: E_x \rightarrow E_0$.

Let $1 \in S^1$, the unit circle in \mathbb{C} .

DEFINITION 2.8. Let x be a point in the stack \mathcal{X} . Define the *fundamental group of \mathcal{X}* to be the set of homotopy classes of pointed maps $(\underline{S}^1, 1) \rightarrow (\mathcal{X}, x)$, denoted $\pi_1(\mathcal{X}, x) := [(\underline{S}^1, 1), (\mathcal{X}, x)]$.

The reader may wish to consult [21] for details concerning the group structure of $\pi_1(\mathcal{X}, x)$.

In Section 4, we shall consider covering projections of (connected) quotient stacks. Making use of [21, Corollary 18.20], we say that a covering projection $\mathcal{Y} \rightarrow \mathcal{X}$ is the *universal covering projection* if $\pi_1(\mathcal{Y}, y)$ is trivial.


An interesting feature of the fundamental group of a stack is the following natural homomorphism $\omega_x: I_x \rightarrow \pi_1(\mathcal{X}, x)$, where $I_x := \{\alpha: x \rightsquigarrow x\}$ is the *inertia group of \mathcal{X} at x* . The homomorphism ω_x is defined as follows (cf. [21, Remark 17.3]). Given $\alpha \in I_x$, let $H_\alpha: \underline{I} \rightarrow \mathcal{X}$ be defined by the composition $\underline{I} \rightarrow \star \xrightarrow{x} \mathcal{X}$ and consider the pair of 2-isomorphisms $\epsilon_0 = \text{id}$ and $\epsilon_1 = \alpha$ in the following diagrams.

$$\begin{array}{ccc}
 \begin{array}{ccc}
 & \star & \\
 \swarrow \iota_0 & & \searrow x \\
 \underline{I} & \xrightarrow{H_\alpha} & \mathcal{X} \\
 & \swarrow \epsilon_0 & \\
 & \mathcal{X} &
 \end{array}
 &
 &
 \begin{array}{ccc}
 & \star & \\
 \swarrow \iota_1 & & \searrow x \\
 \underline{I} & \xrightarrow{H_\alpha} & \mathcal{X} \\
 & \swarrow \epsilon_1 & \\
 & \mathcal{X} &
 \end{array}
 \end{array}$$

This data glues together to give a pointed map $(\underline{S}^1, \underline{1}) \rightarrow (\mathcal{X}, x)$ whose homotopy class is denoted $\omega_x(\alpha)$.


3. The fundamental group of $\mathcal{X} = [X/G]$

We now specialize to our case of interest, stacks $\mathcal{X} = [X/G]$, where X is a smooth manifold equipped with a (right) action of a Lie group G . In this case, recall that the objects of \mathcal{X} are pairs of maps $(B \leftarrow E \rightarrow X)$ consisting of a (right) principal G -bundle $E \rightarrow B$ and a G -equivariant map $E \rightarrow X$ and that the morphisms of \mathcal{X} are pairs $(E \rightarrow E', B \rightarrow B')$, with $E \rightarrow E'$ a G -equivariant map, such that in the diagram,



the square is Cartesian and the triangle commutes. Fix a point x in \mathcal{X} , and write $(\star \leftarrow E_x \rightarrow X)$ to denote the object $x(\text{id}_\star)$ in \mathcal{X}_\star . By choosing a base point $e_0 \in E_x$ we shall further identify $E_x \cong G$, in which case the G -equivariant map $x: E_x \rightarrow X$ is realized by the map $x_p: G \rightarrow X, g \mapsto p \cdot g$ where $p := x(e_0)$. Thus x_p parameterizes the G -orbit through p .

An element $\alpha \in I_x$ induces a gauge transformation $\alpha: E_x \rightarrow E_x$ such that the diagram below commutes.



Moreover, any such gauge transformation determines an element in I_x . Under the identification $E_x \cong G$ determined by $e_0 \leftrightarrow 1$, any $\alpha \in I_x$ is determined by $g_\alpha = \alpha(1)$, which must lie in $\text{Stab}(p)$ since the above diagram commutes. That is, $\alpha: G \rightarrow G$ is simply left multiplication by g_α . In what follows, we will implicitly use the isomorphism $\alpha \mapsto g_\alpha$ to identify $I_x \cong \text{Stab}(p)$.

In this setting, the homomorphisms ω_x discussed at the end of Section 2.2 may be described explicitly. The data determined by $H_\alpha: \underline{1} \rightarrow \mathcal{X}$ yields the trivial bundle $I \times G \rightarrow I$ with G equivariant map $I \times G \rightarrow X$ given by $(t, g) \mapsto p \cdot g$. The 2-isomorphisms $\epsilon_0 = \text{id}$ and $\epsilon_1 = \alpha$ yield the identifications of fibers

$$\begin{aligned}
 \{0\} \times G &\rightarrow G, & (0, g) &\mapsto g, \\
 G &\rightarrow \{1\} \times G, & g &\mapsto (1, g_\alpha g).
 \end{aligned}$$

This assembles to give a bundle over S^1 and an equivariant map representing $\omega_x(\alpha)$ (cf. Remark 2.7). Namely, we take

$$(3.1) \quad \begin{aligned} E(g_\alpha) &:= (I \times G)/\sim, \\ &\text{where the equivalence relation } \sim \text{ is } (0, g) \sim (1, g_\alpha g) \text{ for } g \in G, \end{aligned}$$

and the equivariant map is given by $E(g_\alpha) \rightarrow X, [(t, g)] \mapsto p \cdot g$.

Let $BG := [\star/G]$ and let $\star: \star \rightarrow BG$ be the choice of point whose value at $S \rightarrow \star$ is the trivial bundle $S \times G \rightarrow S$.

Lemma 3.1 (Cf. [20, Example 4.2]). *Let G be a Lie group and let G_0 denote the component of the identity element in G . Then $\pi_1(BG, \star) \cong G/G_0$.*

Proof. We define a map $\sigma: G/G_0 \rightarrow \pi_1(BG, \star)$ and show that σ is an isomorphism. For $g \in G$ let $E(g) \rightarrow S^1$ be the G -bundle $E(g) = (I \times G)/(0, h) \sim (1, gh)$. Let $\sigma(gG_0) = [(f, \alpha)]$ where (f, α) is a morphism determined by the bundles $E(g) \rightarrow S^1$ and $E(g)|_{\{1\}} \rightarrow \star$, the inclusion of the fiber $E(g)|_{\{1\}} \rightarrow E(g)$ over $1: \star \rightarrow S^1$, and the 2-isomorphism α determined by the identification $G \rightarrow E(g)|_{\{1\}}, h \mapsto [(0, h)]$. Here the brackets $[]$ denotes equivalence classes with respect to the relation in (3.1). (Recall that by Remark 2.7, any other pointed map determined by this data differs from (f, α) by a unique 2-isomorphism of pointed maps, which by Lemma 2.6 lies in the same homotopy class.) To see that σ is well-defined, observe that if $\gamma: I \rightarrow G$ is any path from g to g' , we may construct a bundle isomorphism $\varphi_\gamma: E(g) \rightarrow E(g')$ by the formula $[(t, h)] \mapsto [(t, \gamma(t)g^{-1}h)]$ that fits in the diagram

$$(3.2) \quad \begin{array}{ccc} & E(g)|_{\{1\}} & \longrightarrow & E(g) \\ & \nearrow \alpha & & \downarrow \varphi \\ G & & & \\ & \searrow \alpha' & & \downarrow \varphi_{\{1\}} \\ & E(g')|_{\{1\}} & \longrightarrow & E(g') \end{array}$$

where α and α' are the resulting isomorphisms over \star . Let us call such a bundle isomorphism (i.e. one which respects the given trivializations over \star) a *pointed isomorphism*. By Lemma 2.6, we see that the map σ is well-defined.

Since every G -bundle over S^1 is isomorphic to $E(g)$ for some g , it is easy to see that σ is surjective. It remains to show that σ is injective. To that end, suppose that $[(f, \alpha)] = \sigma(gG_0) = \sigma(g'G_0) = [(f', \alpha')]$. Let $(H, \epsilon_0, \epsilon_1)$ be a homotopy from (f, α) to (f', α') . We claim that this results in a pointed isomorphism of bundles as in the diagram (3.2) above, which will complete the proof. Indeed, write the resulting bundle isomorphism $E(g) \rightarrow E(g')$ as $[(t, h)] \mapsto [(t, \phi(t)h)]$, where $\phi: I \rightarrow G$ is continuous and satisfies $g'\phi(0) = \phi(1)g$. That the isomorphism is pointed (i.e. must fit in the diagram (3.2)) forces $\phi(0) = 1$ and we may construct a path $\gamma(t) = \phi(t)g$ joining g and g' , whence $gG_0 = g'G_0$.

We shall now verify the claim that the homotopy produces the required pointed isomorphism of bundles $E(g) \rightarrow E(g')$. Let $E \rightarrow I \times S^1$ be the bundle given by $H(\text{id}: I \times S^1 \rightarrow I \times S^1)$ and let $E_j \rightarrow S^1$ ($j = 0, 1$) denote the bundles given by $H(\varepsilon_j: S^1 = \{j\} \times S^1 \hookrightarrow I \times S^1)$. Note that the 2-isomorphisms of pointed maps ε_j yield pointed isomorphisms of bundles $E(g) \rightarrow E_0$ and $E_1 \rightarrow E(g')$. Therefore, it suffices to find a pointed isomorphism $E_0 \rightarrow E_1$.

Let η denote the 2-isomorphism for the diagram

$$\begin{array}{ccc} \underline{I} & \longrightarrow & \star \\ \downarrow i_1 & \swarrow \eta & \downarrow \\ \underline{I} \times \underline{S}^1 & \xrightarrow{H} & BG \end{array}$$

that in turn induces a trivialization $\eta: I \times G \rightarrow E_I$, where E_I denotes the bundle over I corresponding to $H(i_1: I \rightarrow I \times S^1)$. We therefore seek a pointed isomorphism

$$\begin{array}{ccccc} & & E_0|_{\{1\}} & \longrightarrow & E_0 \\ G & \nearrow \eta_0 & \downarrow \varphi|_{\{1\}} & & \downarrow \varphi \\ & \searrow \eta_1 & E_1|_{\{1\}} & \longrightarrow & E_1 \end{array}$$

where the identifications η_j of the fibers over $1 \in S^1$ are induced from the trivialization η . By Theorem 9.8 in [13, Chapter 4], there is a bundle isomorphism $\psi: E \rightarrow I \times E'$ where E' is a G -bundle over S^1 , inducing an isomorphism $\psi': E_I \rightarrow I \times E'|_{\{1\}}$. Notice that the composition $\psi' \circ \eta: I \times G \rightarrow I \times E'|_{\{1\}}$ defines an identification $\eta': G \rightarrow E'|_{\{1\}}$. Consider the composition of bundle maps given by

$$\rho: E_0 \rightarrow E \xrightarrow{\psi} I \times E' \rightarrow \{1\} \times E' \rightarrow I \times E' \xrightarrow{\psi^{-1}} E$$

and observe that it fits in the commutative square below.

$$\begin{array}{ccc} E_0 & \xrightarrow{\rho} & E \\ \downarrow & & \downarrow \\ S^1 & \xrightarrow{\varpi_1} & I \times S^1 \end{array}$$

By the universal property of Cartesian squares, this induces a unique isomorphism $\varphi: E_0 \rightarrow E_1$, which is the desired pointed isomorphism. \square

REMARK 3.2. It is well-known that for a compact Lie group G , (unpointed) isomorphism classes of principal G -bundles over the circle are in bijective correspondence with the set of conjugacy classes of G/G_0 .

Lemma 3.3. *Let X be a topological space and p a point in X . Then*

$$\pi_1(\underline{X}, \underline{p}) \cong \pi_1(X, p)$$

where the right hand side is the classical fundamental group of a topological space with chosen basepoint.

Proof. A pointed map of stacks $(S^1, 1) \rightarrow (\underline{X}, \underline{p})$ is determined uniquely by a pointed map $(S^1, 1) \rightarrow (X, p)$ of topological spaces. Since there are no non-trivial 2-morphisms in stacks of the form \underline{M} , classical homotopies are in one-to-one correspondence with stack homotopies, and the claim follows. \square

There is a natural morphism $q: \underline{X} \rightarrow [X/G]$, defined on objects as

$$S \rightarrow X \mapsto (S \leftarrow S \times G \rightarrow X \times G \xrightarrow{\text{act}} X).$$

Similarly there is a natural morphism from $[X/G]$ to BG which simply forgets the equivariant map to X . By choosing identifications of the trivial G -bundle over a point, these may each be considered as pointed maps. Finally, recall the pointed map $\iota: G_0 \rightarrow X$ given by parametrizing the orbit of the base point $g \mapsto p \cdot g$. Applying the fundamental group functor to each of these maps and using the isomorphism in the previous Lemma results in the following.

Proposition 3.4. *Let G be a Lie group acting smoothly on a connected manifold X . The following sequence is exact:*


$$(3.3) \quad \pi_1(G_0, 1) \rightarrow \pi_1(X, p) \rightarrow \pi_1([X/G], x) \rightarrow G/G_0 \rightarrow 1.$$

Proof. The verification is straightforward. We prove exactness at $\pi_1(X, p)$ and exactness at G/G_0 , leaving the rest for the reader. Recall first that the trivial element in $\pi_1([X/G], x)$ may be represented by the pair (as in Remark 2.7) $(S^1 \leftarrow S^1 \times G \xrightarrow{\sigma} X)$, where $\sigma(z, g) = p \cdot g$.

We show that the composition of the first two maps in the sequence is trivial. Let $\alpha: S^1 \rightarrow G_0$ represent an element of $\pi_1(G_0, 1)$, whence its image via the composition of the first two maps in the sequence is represented by the pair $(S^1 \leftarrow S^1 \times G \xrightarrow{a} X)$ where $a(z, g) = p \cdot (\alpha(z)g)$. The map α determines a gauge transformation $\phi: S^1 \times G \rightarrow S^1 \times G$ defined by $\phi(z, g) = (z, \alpha(z)g)$. Since $\sigma \circ \phi = a$ we see that ϕ determines a 2-isomorphism of the maps $q \circ \iota \circ \alpha$ and the constant map to $[X/G]$. By Lemma 2.6 this induces a homotopy, verifying that the composition of the first two maps is trivial.

Next, suppose that $f: S^1 \rightarrow X$ is a pointed map representing a class in $\pi_1(X, p)$ whose image via q_* in $\pi_1([X/G], x)$ is trivial. We will show that f is homotopic to a composition $\iota \circ \beta$ for some pointed map $\beta: S^1 \rightarrow G_0$. Suppose that $q \circ f$ is

homotopic to the constant map $S^1 \rightarrow \star \xrightarrow{x} [X/G]$. Let $(I \times S^1 \leftarrow E_H \xrightarrow{h} X)$ be a pair of maps representing the homotopy $H: I \times S^1 \rightarrow [X/G]$. That E_H represents a homotopy implies that there exists a trivialization $\epsilon_0: S^1 \times G \rightarrow E_{H_0} = E_H|_{0 \times S^1}$ fitting in the commutative diagram,




where $\varpi_0(z) = (0, z)$. Note that by Theorem 9.8 in [13, Chapter 4], the bundle $E_H \rightarrow I \times S^1$ is trivializable. Moreover, we may choose a section s so that the composition

$$S^1 \xrightarrow{s \varpi_0} E_{H_0} \xrightarrow{\epsilon_0^{-1}} S^1 \times G$$

is simply inclusion $z \mapsto (z, 1)$.

Since E_H represents a homotopy, there exists a trivialization $\epsilon_1: E_{H_1} = E_H|_{1 \times S^1} \rightarrow S^1 \times G$. Note that $\epsilon_1 \circ s \circ \varpi_1(z) = (z, \beta(z))$ for some loop $\beta: S^1 \rightarrow G$, and that $\beta(1) \in \text{Stab}(p)$ because the homotopy H is a homotopy of pointed maps. By replacing the 2-isomorphism $\epsilon_1: E_{H_1} \rightarrow S^1 \times G$ (that is part of the data of the homotopy H) with the composition of 2-isomorphisms below if necessary,



(where L_u denotes the gauge transformation $(z, g) \mapsto (z, ug)$) we may assume that $\beta(1) = 1$.

The map $\tau = h \circ s: I \times S^1 \rightarrow X$ is the desired homotopy. It is readily verified that $\tau(0, z) = f(z)$, and that $\tau(1, z) = p \cdot \beta(z) = (\iota \circ \beta)(z)$, as required.

Finally, suppose $E(g) \rightarrow S^1$ represents an element of $\pi_1(BG, *) \cong G/G_0$. It suffices to construct an equivariant map $E(g) \rightarrow X$. Choose a point $z \in X$ and a path $\gamma: I \rightarrow X$ with $\gamma(0) = z$ and $\gamma(1) = z \cdot g^{-1}$. The map $I \times G \rightarrow X$ given by $(t, h) \mapsto \gamma(t) \cdot h$ descends to $E(g)$ and is G -equivariant. \square

Corollary 3.5. *Let G be a Lie group acting smoothly on a simply connected manifold X . The fundamental group $\pi_1([X/G], x) \cong G/G_0$. In particular, if in addition G is connected, then $\pi_1([X/G], x)$ is trivial.*

Proposition 3.6. *Under the identifications $I_x \cong \text{Stab}(p)$ and $\pi_1(BG, *) \cong G/G_0$ given above, the composition $\text{Stab}(p) \cong I_x \xrightarrow{\omega_x} \pi_1([X/G], x) \rightarrow G/G_0$ is the natural homomorphism $\phi_p: \text{Stab}(p) \hookrightarrow G \rightarrow G/G_0$.*

Proof. For $g \in \text{Stab}(p)$, the image of $\omega_x(g)$ in $\pi_1(BG, *)$ is represented by the bundle $E(g)$ (in the notation above), which by the proof of Lemma 3.1 is represented by the coset gG_0 . □

4. Global quotients and universal coverings

A main purpose of this paper is to determine conditions under which a Deligne–Mumford stack is a global quotient in the sense of the following definition.

DEFINITION 4.1. A Deligne–Mumford stack \mathcal{X} is a *discrete* (resp. *global*) *quotient* if \mathcal{X} is equivalent to a quotient stack $[Y/\Gamma]$, where Γ is a discrete (resp. finite) group acting on a smooth manifold Y .

Recall that a morphism of representable stacks $\mathcal{X} \rightarrow \mathcal{Y}$ is an *equivalence* if it is an equivalence of categories. An equivalence may also be represented in terms of the corresponding representing groupoids as a principal bi-bundle (e.g. see Definition 3.25 and Remark 3.33 in [17]).

We shall deal only with the special case of quotient stacks $\mathcal{X} = [X/G]$ arising from a smooth proper action of a Lie group G on a connected manifold X . Note that an equivalence of such a pair of quotient stacks $[X/G] \rightarrow [Y/H]$ may then be represented as a bi-bundle of the action groupoids $X \times G \rightrightarrows X$ and $Y \times H \rightrightarrows Y$, which in this case amounts to a $G \times H$ -space P that is simultaneously a principal G -bundle $P \rightarrow Y$ (with H -equivariant projection) and a principal H -bundle $P \rightarrow X$ (with G -equivariant projection).

REMARK 4.2. The properness assumption does not appear in every result in this section, though our main interest remains the case of proper Lie group actions on manifolds (giving rise to Deligne–Mumford stacks). When properness is stated as a hypothesis, the reader will note that its main use is to ensure that a certain quotient (appearing in the proof of that statement) is a smooth manifold.

4.1. Global quotients and the fundamental group. The following Lemma provides a natural setting to discuss a class of examples of quotient stacks that are equivalent to global quotients.

Lemma 4.3. *Let $1 \rightarrow H \rightarrow G \rightarrow \Gamma \rightarrow 1$ be an exact sequence of topological (respectively, Lie) groups. Suppose that G acts on a topological space (resp. smooth manifold) X and that the restriction of this action to H is free (resp. free and proper). Then $[X/G]$ and $[(X/H)/\Gamma]$ are equivalent as stacks over Top (resp. Diff).*

Proof. We shall work over Diff , noting that the proof is the same over Top . Since the H -action is free and proper, the orbit space X/H is indeed a smooth manifold. Define $F: [X/G] \rightarrow [(X/H)/\Gamma]$ to be the functor defined by the assignment

$$(B \leftarrow E \rightarrow X) \mapsto (B \leftarrow E/H \rightarrow X/H)$$

on objects, and

$$\begin{array}{ccc}
 & & X \\
 & \nearrow & \nearrow \\
 E & \longrightarrow & E' \\
 \downarrow & & \downarrow \\
 B & \longrightarrow & B'
 \end{array}
 \mapsto
 \begin{array}{ccc}
 & & X/H \\
 & \nearrow & \nearrow \\
 E/H & \longrightarrow & E'/H \\
 \downarrow & & \downarrow \\
 B & \longrightarrow & B'
 \end{array}$$

on arrows. By construction, F commutes with the projections to the base category Diff . We wish to show that F is an equivalence of categories. To see this, we define a functor $K: [(X/H)/\Gamma] \rightarrow [X/G]$ as follows. Suppose given a pair $(B \leftarrow P \rightarrow X/H)$ consisting of a Γ -bundle $P \rightarrow B$ and a Γ -equivariant map $P \rightarrow X/H$. Let $E := P \times_{X/H} X$ be the fiber product, and define a G -action on E by setting $(p, z) \cdot g = (p \cdot \rho(g), z \cdot g)$, where $\rho: G \rightarrow \Gamma$ denotes the given map in the exact sequence. We claim that the composition $E \xrightarrow{\text{pr}_1} P \rightarrow B$ is a principal G -bundle. Indeed suppose $(p, z) \cdot g = (p \rho(g), z g) = (p, z)$. Then since P is a principal Γ -bundle, $\rho(g)$ is the identity element in Γ and hence $g \in H$. On the other hand, by assumption H acts freely on X so $z g = z$ implies $g = \text{id}$ in G . Hence G acts freely on E . Next suppose $(p, z) \in E, (p', z') \in E$ map to the same point b in B . Then since P is a principal Γ -bundle over B , there exists $g \in G$ such that $p' = p \rho(g)$. By definition of the fiber product $E = P \times_{X/H} X$, the equivariance of the map $P \rightarrow X/H$, and normality of H , we conclude there exists $h \in H$ such that $z' = z h g$. Since $\rho(h g) = \rho(g) \in \Gamma$ we conclude $(p, z) h g = (p', z')$ and that G acts transitively on fibers of $E \rightarrow B$. Hence $E \rightarrow B$ is a principal G -bundle, as desired. The projection map $E \rightarrow X$ is G -equivariant by construction so $(B \leftarrow E \rightarrow X)$ is an object in $[X/G]$. Given an arrow

$$\begin{array}{ccc}
 & & X/H \\
 & \nearrow & \nearrow \\
 E/H & \longrightarrow & E'/H \\
 \downarrow & & \downarrow \\
 B & \longrightarrow & B'
 \end{array}$$

in $[(X/H)/\Gamma]$, it induces a unique arrow

$$\begin{array}{ccc}
 & & X \\
 & \nearrow & \nearrow \\
 E/H \times_{X/H} X & \longrightarrow & E'/H \times_{X/H} X \\
 \downarrow & & \downarrow \\
 B & \longrightarrow & B'
 \end{array}$$

which defines the functor K on morphisms. Again by construction K commutes with projection to the base category.

Finally, we sketch the constructions of the natural transformations between $F \circ K$ (resp. $K \circ F$) and the identity functor on $[(X/H)/\Gamma]$ (resp. $[X/G]$). For $K \circ F$, observe that for any object $(B \leftarrow E \rightarrow X)$ in $[X/G]$, there is a unique isomorphism

$$\begin{array}{ccc}
 & & X \\
 & \nearrow & \nearrow \\
 E & \longrightarrow & E/H \times_{X/H} X \\
 \searrow & & \swarrow \\
 & B &
 \end{array}$$

in $[X/G]$. For $F \circ K$, observe that $(P \times_{X/H} X)/H$ is isomorphic to P via the map $[(p, z)] \mapsto p$. From here it is straightforward to check that these yields the desired natural transformations. \square

Observe that the requirement in Lemma 4.3 that the G -action restricted to H be a free H -action is necessary over both Diff and Top. (Compare with Proposition 4.6.) For example, consider the exact sequence

$$1 \rightarrow \mathbb{Z}_2 \rightarrow S^1 \xrightarrow{2} S^1 \rightarrow 1$$

where $2: S^1 \rightarrow S^1$ denotes the squaring map. Let $t \in G = S^1$ act on $X = S^3 \subset \mathbb{C}^2$ according to $t \cdot (z, w) = (t^2z, t^2w)$. The quotient stack $[X/G]$ is the weighted projective space $\mathbb{P}(2, 2)$. The restriction of the G -action to $H = \mathbb{Z}_2 = \{\pm 1\}$ is trivial, and the resulting residual action of $\Gamma = S^1$ on $X/H = X = S^3$ is the standard action of S^1 on S^3 giving the quotient $[(X/H)/\Gamma] = \mathbb{P}(1, 1)$, the complex projective plane. As a stack, $\mathbb{P}(2, 2)$ has a non-trivial inertia group isomorphic to \mathbb{Z}_2 at each point and is thus not equivalent to the smooth manifold $\mathbb{P}(1, 1)$.

If G is compact, applying the above Lemma to the case $H = G_0$, the connected component of the identity element, provides a natural class of examples of stacks equivalent to global quotients. (Since G is compact, the quotient G/G_0 is automatically a finite group.)

Corollary 4.4. *Suppose a Lie group G acts smoothly and properly on a connected smooth manifold X . Let Γ denote the (discrete) group G/G_0 of G . If the restriction of the G -action to G_0 is free, then $[X/G]$ and $[(X/G_0)/\Gamma]$ are equivalent as stacks over Diff and hence $[X/G]$ is a discrete quotient. If in addition G is compact, then $[X/G]$ is a global quotient.*

If in addition X is simply connected, then we shall see in Theorem 4.10 that the above examples characterize global quotients among quotient stacks. Proposition 4.5 below illustrates how the freeness of the G_0 -action on X relates to the fundamental group $\pi_1([X/G], x)$.

Proposition 4.5. *Suppose a Lie group G acts smoothly on a connected smooth manifold X . If the restriction of the G -action to $G_0 \subset G$ is free, then the homomorphism ω_x is injective for all points x in $[X/G]$. Moreover, if X is simply connected, the converse holds as well.*

Proof. From Proposition 3.6 we know that the composition of the homomorphism ω_x with the second arrow in (3.3) is the natural homomorphism $\phi_p: \text{Stab}(p) \rightarrow G/G_0$ obtained as the composition of the natural inclusion $\text{Stab}(p) \hookrightarrow G$ with the canonical quotient map $G \rightarrow G/G_0$. If the restriction of the G -action to G_0 is free, then ϕ_p is injective for all p and hence ω_x is injective for all x in $[X/G]$. This proves the first claim.

For the second claim, if X is simply connected then by Corollary 3.5 $\pi_1([X/G], x) \cong \pi_1(BG, *) \cong G/G_0$ and $\ker \omega_x = \text{Stab}(p) \cap G_0$. Therefore, if ω_x is injective for all x in $[X/G]$ then G_0 acts freely on X . \square

4.2. On covers of quotient stacks. Lemma 4.3 may be generalized to the context of group actions on stacks, which then fits nicely with covering theory. In preparation for the statement of Proposition 4.6, we begin with a summary of some ideas found in the work of Lerman and Malkin [18], which the reader should consult for details.

For a Lie group Λ , recall that a Λ -action on a stack \mathcal{X} can be encoded using a Λ -presentation, a groupoid presentation $\mathcal{G} = \mathcal{G}_1 \rightrightarrows \mathcal{G}_0$ of \mathcal{X} equipped with smooth and free Λ -actions on both the manifold of arrows \mathcal{G}_1 and the manifold of objects \mathcal{G}_0 that is compatible with the structure maps of the groupoid \mathcal{G} .

Towards generalizing Lemma 4.3, suppose that

$$1 \rightarrow H \hookrightarrow G \rightarrow \Gamma \rightarrow 1$$

is an exact sequence of Lie groups and that G acts smoothly on a manifold X . The exact sequence above naturally defines a G -action on $X \times \Gamma$ and the translation groupoid

$$(4.1) \quad G \times (X \times \Gamma) \rightrightarrows X \times \Gamma$$

is then a groupoid presentation for the quotient stack $[(X \times \Gamma)/G]$. As in [18, Section 4.1], since translation by Γ commutes with the above G -actions, we see that (4.1) is a Γ -presentation. By [18, Proposition 4.2] (4.1) is also a Γ -presentation for the quotient stack $[X/H]$. This gives a Γ -action on $[X/H]$, which (see [18, Section 3.3]) shows that the translation groupoid $G \times X \rightrightarrows X$ is a groupoid presentation of the stack quotient $[X/H]/\Gamma$. This verifies the following generalization of Lemma 4.3.

Proposition 4.6. *Let $1 \rightarrow H \rightarrow G \rightarrow \Gamma \rightarrow 1$ be an exact sequence of Lie groups. Suppose that G acts smoothly on a smooth manifold X . Then the quotient stack $[X/H]$ inherits a Γ -action; moreover, $[X/G]$ and $[X/H]/\Gamma$ are equivalent as stacks over Diff .*

As in previous discussions, we wish to interpret the above Proposition in the case $H = G_0$. This interpretation may be placed in the context of covering theory for stacks [21]. In particular, we shall see in Proposition 4.8 that the natural map $p: [X/G_0] \rightarrow [X/G]$ is a covering projection. (More generally, there is a natural map $[X/H] \rightarrow [X/G]$, given by the associated bundle construction, which is representable by Lemma 4.7 below.) In other words, we may view p as a quotient map, for $[X/G] \cong [X/G_0]/\Gamma$ where Γ is the discrete group G/G_0 .

Parallel to classical covering space theory, one may define universal covering projections. For simplicity, we shall define a *universal covering projection* $(\tilde{\mathcal{X}}, \tilde{x}) \rightarrow (\mathcal{X}, x)$, as a covering projection with $\pi_1(\tilde{\mathcal{X}}, \tilde{x}) = \{1\}$ (cf. [21, Corollary 18.20]).

Lemma 4.7. *Let H be a closed subgroup of a Lie group G that acts smoothly on a manifold X . The natural map $[X/H] \rightarrow [X/G]$ given by the associated bundle construction is representable.*

Proof. Let $\varphi: \underline{W} \rightarrow [X/G]$ and let $(W \leftarrow E_\varphi \rightarrow X)$ denote $\varphi(\text{id}_W)$. Recall that the fiber product $\mathcal{Z} = [X/H] \times_{[X/G]} \underline{W}$ has objects given by triples $(B \leftarrow E \rightarrow X, f: B \rightarrow W, \alpha)$ where $\alpha \in [X/G]_B$ is an isomorphism of G -bundles $\alpha: E \times_H G \rightarrow f^*E_\varphi$ (compatible with the maps to X). An arrow in \mathcal{Z} between two such objects is an arrow $(E \rightarrow E', B \rightarrow B')$ in $[X/H]$ such that $B \rightarrow B'$ is compatible with the maps to W and that the resulting (vertical) induced maps in the diagram below commute.

$$\begin{array}{ccc} E \times_H G & \xrightarrow{\alpha} & f^*E_\varphi \\ \downarrow & & \downarrow \\ E' \times_H G & \xrightarrow{\alpha'} & (f')^*E_\varphi \end{array}$$

Since $E_\varphi \rightarrow W$ is a principal G -bundle, the G -action on E_φ is free and proper; therefore, the restriction of this action to $H \subset G$ is also free and proper, whence E_φ/H is a manifold in Diff . Define $F: \mathcal{Z} \rightarrow \underline{E_\varphi/H}$ as follows. Using the section $B \rightarrow$

$(E \times_H G)/H$, which sends $b \in B$ to the H -orbit of $[(e, 1)]$ where $e \in E$ is any element in the fiber over b , and composing with the induced composition

$$(E \times_H G)/H \xrightarrow{\alpha/H} f^* E_\varphi/H \rightarrow E_\varphi/H$$

we obtain a map $B \rightarrow E_\varphi/H$. Hence, on objects, we define $F(B \leftarrow E \rightarrow X, f: B \rightarrow W, \alpha) = (B \rightarrow E_\varphi/H)$. (The effect of F on arrows is the natural one.)

To show that F is an equivalence, we next define a morphism $K: \underline{E_\varphi/H} \rightarrow \mathcal{Z}$ as follows. Given a $B \rightarrow E_\varphi/H$, let P denote the pullback H -bundle

$$\begin{array}{ccc} P & \longrightarrow & E_\varphi \\ \downarrow & & \downarrow \\ B & \longrightarrow & E_\varphi/H \end{array}$$

let f denote the composition $f: B \rightarrow E_\varphi/H \rightarrow W$. Since the map $P \times_H G \rightarrow E_\varphi$ given by $[(b, e, g)] \mapsto e \cdot g$ covers f , there is a unique isomorphism $P \times_H G \rightarrow f^* E_\varphi$ which we denote by α . Hence, on objects, we define $K(B \rightarrow E_\varphi/H)$ to be the triple $(B \leftarrow P \rightarrow E_\varphi \rightarrow X, f: B \rightarrow W, \alpha)$. (The effect of K on an arrow $B \rightarrow B'$ in $\underline{E_\varphi/H}$ is hence determined.)

It is straightforward to verify that $F \circ K$ is the identity. To realize the natural transformation between $K \circ F$ and the identity, we simply note that pulling back an H -bundle via a composition yields a canonical bundle isomorphism, so that the first factor in the triple for $K \circ F(B \leftarrow E \rightarrow X, f: B \rightarrow W, \alpha)$ is thus canonically isomorphic to $B \leftarrow E \rightarrow X$, which results in the desired natural transformation. □

Proposition 4.8. *Suppose a Lie group G acts on a smooth manifold X and let G_0 denote the identity component of G . The natural map $p: [X/G_0] \rightarrow [X/G]$ is a covering projection. Moreover, if X is simply connected, p is the universal covering projection.*

Proof. From Lemma 4.7, the natural map p is representable, and the proof of the Lemma shows that given $\underline{W} \rightarrow [X/G]$, that the induced G/G_0 -bundle $E_\varphi/G_0 \rightarrow W$ (a covering projection) is a representative for p . If X is simply connected, by Corollary 3.5, p is the universal covering projection. □

REMARK 4.9. The above proposition identifies the universal cover of the quotient stack $[X/G]$ in the setting when X is simply connected. In Proposition 4.14 below, we identify the universal cover of $[X/G]$ when X is *not* simply connected.

4.3. Characterizations of global quotients among quotient stacks of simply connected manifolds X . We now state the main result of this section, which in a sense also summarizes the previous subsections. Theorem 4.10 below characterizes discrete (resp. global) quotients among quotient stacks of simply connected manifolds. (The reader may wish to compare with [21, Theorem 18.24], which discusses a more general setting.)

Theorem 4.10. *Let X be a simply connected manifold, equipped with a smooth proper action of a Lie group G . Let $G_0 \subset G$ denote the connected component of the identity, and I_x the inertia group of $x \in [X/G]$. The following statements are equivalent.*

- (1) $[X/G]$ is equivalent to a discrete quotient.
- (2) G_0 acts freely on X .
- (3) $\omega_x : I_x \rightarrow \pi_1([X/G], x)$ is injective for all x in $[X/G]$.
- (4) The universal cover of $[X/G]$ is equivalent to a smooth manifold.

Proof. Most implications follow directly from work in previous sections: (2) \Rightarrow (1) is Corollary 4.4; (3) \Leftrightarrow (2) follows from Proposition 4.5. By Proposition 4.8, the universal cover of $[X/G]$ is $[X/G_0]$, which verifies (4) \Rightarrow (2). Conversely, if G_0 acts freely on X , the principal G_0 -bundle $X \rightarrow X/G_0$ may be viewed as a bi-bundle equivalence $[X/G_0] \cong \underline{X}/G_0$, and hence (2) \Rightarrow (4).

It remains to show (1) \Rightarrow (2). To that end, suppose given a principal bi-bundle representing an equivalence $[X/G] \cong [Z/\Lambda]$ where Λ is discrete. Recall that this yields a $G \times \Lambda$ -space P that is simultaneously a G -bundle $P \rightarrow Z$ (with Λ -equivariant projection) and a Λ -bundle $P \rightarrow X$ (with G -equivariant projection). Since X is simply connected, we have $P \cong X \times \Lambda$. And since the G and Λ -actions commute, the G -action on $X \times \Lambda$ may be written

$$g \cdot (x, \lambda) = (g \cdot x, \phi(g, x)\lambda)$$

where in the first factor \cdot signifies the original G -action on X . Since Λ is discrete, ϕ only depends on the component of g , yielding a homomorphism $\varphi : G/G_0 \rightarrow \Lambda$ with $\varphi(gG_0) = \phi(g, x)$ for any x . Finally, if $g \in G_0$ stabilizes x in X , then $g \cdot (x, \lambda) = (x, \lambda)$ and hence g is the identity element, as required. □

REMARK 4.11. In the proof of (1) \Rightarrow (2) above, one may obtain P by forming the Cartesian square

$$\begin{array}{ccc} P & \longrightarrow & Z \\ \downarrow & & \downarrow \\ X & \longrightarrow & [X/G] \cong [Z/\Lambda] \end{array}$$

The authors wish to thank the referee for this observation.

The condition that X be simply connected in the above theorem is necessary, as illustrated by the following example.

EXAMPLE 4.12. Let $G = S^1$ act on $X = S^1$ with weight 2. This action has a global stabilizer $\mathbb{Z}_2 = \{\pm 1\} \subset G_0 = G$. Nevertheless, we may readily verify that $[X/G] \cong B\mathbb{Z}_2 = [\star/\mathbb{Z}_2]$. Explicitly, consider the following functors F and K . On objects, let $F: B\mathbb{Z}_2 \rightarrow [X/G]$ be defined by taking associated bundle

$$F(E \rightarrow B) = (B \leftarrow E \times_{\mathbb{Z}_2} S^1 \xrightarrow{f} X)$$

with $f([(e, z)]) = z^2$, and let $K: [X/G] \rightarrow B\mathbb{Z}_2$ be given by $K(B \leftarrow P \xrightarrow{f} S^1) = (f^{-1}(1) \rightarrow B)$. (The effects of F and K on arrows are the natural ones.) Alternatively, see Example 4.15 and Proposition 4.14 below.

4.4. The universal cover of a quotient stack of non-simply connected manifold X . We next work towards a statement in the spirit of the equivalence (1) \Leftrightarrow (2) of Theorem 4.10 for the case of connected manifolds X that are not necessarily simply connected. Let X and G be as in Proposition 4.8, and let \tilde{X} denote the universal cover of a smooth manifold X and consider the induced action of \tilde{G}_0 on \tilde{X} . Let Λ denote the image of $\pi_1(G_0, 1) \rightarrow \pi_1(X, p)$ (cf. (3.3)). Since Λ is a subgroup of deck transformations, \tilde{X}/Λ is a smooth manifold. Moreover, $G_0 = \tilde{G}_0/\pi_1(G_0, 1)$ acts on \tilde{X}/Λ , and the covering projection $\tilde{X}/\Lambda \rightarrow X$ is G_0 equivariant.

The following technical lemma helps to identify the natural map $[(\tilde{X}/\Lambda)/G_0] \rightarrow [X/G_0]$ as a covering projection in the following proposition.

Lemma 4.13. *Suppose a Lie group G acts smoothly on connected manifolds Y and X . Assume the G -action on Y is proper and that $f: Y \rightarrow X$ is a G -equivariant submersion. Then the canonical map $[Y/G] \rightarrow [X/G]$ is representable.*

Proof. Let $\varphi: \underline{W} \rightarrow [X/G]$ be given and let $(W \leftarrow E_\varphi \rightarrow X)$ denote $\varphi(\text{id}_W)$. The fiber product $Y \times_X E_\varphi$ is a smooth G -invariant manifold since $Y \rightarrow X$ is a G -equivariant submersion. Moreover, the diagonal G -action on $Y \times_X E_\varphi$ is proper since the G -actions on Y and E_φ are both proper. Further, since the canonical map $Y \times_X E_\varphi \rightarrow E_\varphi$ is G -equivariant and G acts freely on E_φ , then G also acts freely on $Y \times_X E_\varphi$. By the slice theorem for proper G -actions, we conclude that the quotient $U = (Y \times_X E_\varphi)/G$ is a smooth manifold.

We claim that $Z = [Y/G] \times_{[X/G]} \underline{W} \cong \underline{U}$. Recall that the objects of Z are triples $(B \leftarrow E \rightarrow Y, f: B \rightarrow W, \alpha)$ where $\alpha: E \rightarrow f^*E_\varphi$ is a bundle isomorphism compatible with the equivariant maps to X . An object in Z therefore determines a canonical map $E \rightarrow Y \times_X E_\varphi$ that is G -equivariant. This map descends to a map $B \rightarrow U$, which defines a functor $F: Z \rightarrow \underline{U}$.

Towards showing that F is an equivalence, we next define $K: \underline{U} \rightarrow Z$. Given a map $B \rightarrow U$, by pulling back and composing with the natural projection, we obtain $B \leftarrow P \rightarrow Y \times_X E_\varphi \rightarrow Y$, and we may set $K(B \rightarrow U)$ to be the triple $(B \leftarrow P \rightarrow Y, B \rightarrow U \rightarrow W, \alpha)$ where α is the canonical isomorphism given by pulling back along a composition.

That $F \circ K$ is the identity is easily verified. Similarly, the composition $K \circ F$ is canonically isomorphic to the identity functor. \square

Proposition 4.14. *Suppose a compact Lie group G acts smoothly on a connected manifold X . The quotient stack $[X/G]$ is equivalent to a quotient of a discrete group action over Diff if and only if the restriction of the induced G_0 -action on \tilde{X}/Λ is free, where Λ denotes the image of $\pi_1(G_0, 1) \rightarrow \pi_1(X, p)$. Additionally, the composition of natural maps $[(\tilde{X}/\Lambda)/G_0] \rightarrow [X/G_0] \rightarrow [X/G]$ is a universal covering projection.*

Proof. By [21, Theorem 18.24] $[X/G]$ is equivalent (over Diff) to a quotient by a discrete group action if and only if its universal cover is equivalent to a manifold. By Proposition 4.8, it suffices to determine conditions under which the universal cover of $[X/G_0]$ is equivalent to a manifold, which is done next.

The representable map $p: [(\tilde{X}/\Lambda)/G_0] \rightarrow [X/G_0]$ is a covering projection. Indeed, by the proof of the previous Lemma, given $\varphi: \underline{W} \rightarrow [X/G_0]$, the natural projection $(\tilde{X}/\Lambda) \times_X E_\varphi \rightarrow E_\varphi$ is a G_0 -equivariant covering projection, which induces $((\tilde{X}/\Lambda) \times_X E_\varphi)/G_0 \rightarrow E_\varphi/G_0 = W$, a covering projection that represents p .

Applying the exact sequence (3.3) to $[(\tilde{X}/\Lambda)/G_0]$, and noting that the first map in this exact sequence is a surjection, we see that $[(\tilde{X}/\Lambda)/G_0]$ is the universal cover of $[X/G_0]$. Finally, $[(\tilde{X}/\Lambda)/G_0]$ is equivalent to a manifold if and only if (the compact group) G_0 acts freely on \tilde{X}/Λ . \square

EXAMPLE 4.15 (Example 4.12, revisited). If $G = S^1 = \mathbb{R}/\mathbb{Z}$ acts on $X = S^1 = \mathbb{R}/\mathbb{Z}$ with weight 2, then $\Lambda = 2\mathbb{Z} \subset \mathbb{Z} = \pi_1(X, 1)$ and the induced G -action on $\tilde{X}/\Lambda = \mathbb{R}/\Lambda$ may be written $e^{2\pi i\theta} \cdot e^{\pi i t} = e^{\pi i(t+2\theta)}$, which is free (and transitive). Therefore, as in the proof of the previous proposition, the universal cover of $[X/G]$ is $[(\mathbb{R}/\Lambda)/G] = \star$ and $[X/G] \cong [\star/\mathbb{Z}_2]$.

5. Toric DM stacks

We now apply the ideas of the previous section to toric Deligne–Mumford stacks arising from the combinatorial data of *stacky fans* [5] and *stacky polytopes* [24]. As we shall review below, these stacks arise as quotients $[X/G]$ of a simply connected spaces X ; therefore, we may apply Theorem 4.10.

5.1. Stacky fans and polytopes—brief review. Mainly to establish notation, we briefly recall some basic definitions of the combinatorial data appearing in the above discussion. In the following we use $(-)^*$ to denote the functor $\text{Hom}_{\mathbb{Z}}(-, \mathbb{Z})$ or $\text{Hom}_{\mathbb{R}}(-, \mathbb{R})$; it should be clear from context which one is meant. We use angled brackets $\langle -, - \rangle$ to indicate a natural pairing defined by duality. Also, $- \otimes -$ signifies $- \otimes_{\mathbb{Z}} -$.

Let $\{e_1, \dots, e_n\}$ be the standard basis vectors in $\mathbb{Z}^n \subset \mathbb{R}^n$.

DEFINITION 5.1 ([5]). A *stacky fan* is a triple (N, Σ, β) consisting of a rank d finitely generated abelian group N , a rational simplicial fan Σ in $N \otimes \mathbb{R}$ with rays ρ_1, \dots, ρ_n and a homomorphism $\beta: \mathbb{Z}^n \rightarrow N$ satisfying:

- (1) the rays ρ_1, \dots, ρ_n span $N \otimes \mathbb{R}$, and
- (2) for $1 \leq j \leq n$, $\beta(e_j) \otimes 1$ is on the ray ρ_j .

Given a polytope $\Delta \subseteq \mathbb{R}^d$, recall that the corresponding fan $\Sigma = \Sigma(\Delta)$ is obtained by setting the one dimensional cones $\Sigma^{(1)}$ to be the positive rays spanned by the inward-pointing normals to the facets of Δ ; a subset σ of these rays is a cone in Σ precisely when the corresponding facets intersect nontrivially in Δ . Observe that under this correspondence, facets intersecting in a vertex of Δ yield maximal cones (with respect to inclusion) in $\Sigma(\Delta)$.

DEFINITION 5.2 ([24]). A *stacky polytope* is a triple (N, Δ, β) consisting of a rank d finitely generated abelian group N , a simple polytope Δ in $(N \otimes \mathbb{R})^*$ with n facets F_1, \dots, F_n and a homomorphism $\beta: \mathbb{Z}^n \rightarrow N$ satisfying:

- (1) the cokernel of β is finite, and
- (2) for $1 \leq j \leq n$, $\beta(e_j) \otimes 1$ in $N \otimes \mathbb{R}$ is an inward pointing normal to the facet F_j .

Condition (2) above implies that the polytope Δ in Definition 5.2 is a rational polytope. Also, from the preceding discussion it follows immediately that the data of a stacky polytope (N, Δ, β) specifies the data of a stacky fan by the correspondence $(N, \Delta, \beta) \mapsto (N, \Sigma(\Delta), \beta)$. Indeed, Δ is simple if and only if $\Sigma(\Delta)$ is simplicial. Moreover, the fan $\Sigma(\Delta)$ is rational by condition 5.2 (2). Finally, (N, Δ, β) satisfies conditions (1) and (2) of Definition 5.2 if and only if $(N, \Sigma(\Delta), \beta)$ satisfies conditions (1) and (2) of Definition 5.1.

The extra information encoded in a stacky polytope (N, Δ, β) (compared with the stacky fan $(N, \Sigma(\Delta), \beta)$) results in a symplectic structure on the associated toric DM stack. Given a presentation of a rational polytope Δ as the intersection of half-spaces

$$(5.1) \quad \Delta = \bigcap_{i=1}^n \{x \in (N \otimes \mathbb{R})^* \mid \langle x, \beta(e_i) \otimes 1 \rangle \geq -c_i\}$$

for some $c_i \in \mathbb{R}$ and where each $\beta(e_i) \otimes 1 \in N \otimes \mathbb{R}$ is the inward pointing normal to the facet F_i , the fan $\Sigma(\Delta)$ only retains the data of the positive ray spanned by the normals,

and not the parameters c_i , which encode the symplectic structure on the resulting DM stack (see [24] for details).

Stacky polytopes can be thought of as generalizations of Lerman and Tolman’s labelled polytopes. In its original form [16], a labelled polytope is a pair $(\Delta, \{m_i\}_{i=1}^n)$ consisting of a convex simple polytope Δ in $(N \otimes \mathbb{R})^*$, where N is a lattice, with n facets F_1, \dots, F_n whose relative interiors are labelled with positive integers m_1, \dots, m_n . If we denote the primitive inward pointing normals $v_1 \otimes 1, \dots, v_n \otimes 1$, then defining $\beta: \mathbb{Z}^d \rightarrow N$ by the formula $\beta(e_i) = m_i v_i$ realizes (N, Δ, β) as a stacky polytope. Thus labelled polytopes are precisely the subset of the stacky polytopes for which the \mathbb{Z} -module N is a free module. By results of Fantechi, Mann, and Nironi [9, Lemma 7.15] this is equivalent to the geometric condition that the associated toric DM stack has no global stabilizers.

5.2. Toric DM stacks from stacky fans and polytopes. Recall (as in [5]) that given a stacky fan (N, Σ, β) , the corresponding DM stack may be constructed as a quotient stack $[Z_\Sigma/G]$ as follows. As with classical toric varieties, the fan Σ determines an ideal J_Σ generated by the monomials $\prod_{\rho_i \notin \sigma} z_i \in \mathbb{C}[z_1, \dots, z_n]$ corresponding to the cones σ in Σ . Let Z_Σ denote the complement $\mathbb{C}^n \setminus V(J_\Sigma)$ of the vanishing locus of J_Σ . Note that Z_Σ is the complement of a union of coordinate subspaces of complex codimension at least 2; therefore, Z_Σ is simply connected. Next, we recall a certain group action on Z_Σ .

Choose a free resolution

$$0 \rightarrow \mathbb{Z}^r \xrightarrow{Q} \mathbb{Z}^{d+r} \rightarrow N \rightarrow 0$$

of the \mathbb{Z} -module N , and let $B: \mathbb{Z}^n \rightarrow \mathbb{Z}^{d+r}$ be a lift of β . With these choices, define the dual group $\text{DG}(\beta) = (\mathbb{Z}^{n+r})^*/\text{im}[B \ Q]^*$ where $[B \ Q]: \mathbb{Z}^{n+r} = \mathbb{Z}^n \oplus \mathbb{Z}^r \rightarrow \mathbb{Z}^{d+r}$ denotes the map whose restrictions to the first and second summands are B and Q , respectively. Let $\beta^\vee: (\mathbb{Z}^n)^* \rightarrow \text{DG}(\beta)$ be the composition of the inclusion $(\mathbb{Z}^n)^* \rightarrow (\mathbb{Z}^{n+r})^*$ (into the first n coordinates) and the quotient map $(\mathbb{Z}^{n+r})^* \rightarrow \text{DG}(\beta)$. Applying the functor $\text{Hom}_{\mathbb{Z}}(-, \mathbb{C}^*)$ to β^\vee yields a homomorphism $G := \text{Hom}_{\mathbb{Z}}(\text{DG}(\beta), \mathbb{C}^*) \rightarrow (\mathbb{C}^*)^n$, which defines a G -action on \mathbb{C}^n , which leaves $Z_\Sigma \subset \mathbb{C}^n$ invariant. Define $\mathcal{X}(N, \Sigma, \beta) = [Z_\Sigma/G]$. By Proposition 3.2 in [5], $\mathcal{X}(N, \Sigma, \beta)$ is a DM stack.

The above construction was adapted to stacky polytopes by Sakai in [24]. As the reader may verify, the DM stack $\mathcal{X}(N, \Delta, \beta)$ obtained from a stacky polytope is a quotient stack $[Z_\Delta/H]$ where Z_Δ is a retract of Z_Σ (cf. [24, Lemma 27]) equipped with an action of the compact abelian Lie group $H = \text{Hom}_{\mathbb{Z}}(\text{DG}(\beta), S^1)$. Similar to the discussion in the preceding paragraph, H acts on \mathbb{C}^n and the invariant subset Z_Δ is a certain level set $\mu^{-1}(c)$ of the moment map $\mu: \mathbb{C}^n \rightarrow \mathfrak{h}^*$ for this H -action (where \mathfrak{h} denotes the Lie algebra of H). In particular, the regular value c is determined by the constants c_1, \dots, c_n appearing in (5.1) (see [24, Lemma 16]).

5.3. The fundamental group and inertia homomorphism of a toric DM stack associated to a stacky fan. By Corollary 3.5, the fundamental group of a toric DM stack $\mathcal{X}(N, \Sigma, \beta) = [Z_\Sigma/G]$ associated to a stacky fan (N, Σ, β) is $\pi_1(\mathcal{X}(N, \Sigma, \beta), z) \cong G/G_0$, where G_0 is the connected component of the identity element. Using Proposition 3.6, we compute the inertia homomorphisms $\omega_z: G_z \rightarrow \pi_1(\mathcal{X}(N, \Sigma, \beta), z) \cong G/G_0$, for the various isotropy groups G_z that arise.

In [12], both the isotropy groups G_z and the quotient G/G_0 are described in terms of the stacky fan data, which we summarize next. The isotropy group of a point in $\mathcal{X}(N, \Sigma, \beta)$ arises as the stabilizer $\text{Stab}(z) \subset G$ of $z \in Z_\Sigma \subset \mathbb{C}^n$. These stabilizers depend only on the cone σ in Σ satisfying $\{i: z_i = 0\} = \{i: \rho_i \subset \sigma\}$; namely, for such a cone σ , the corresponding isotropy group Γ_σ is the kernel of the composition

$$G \xrightarrow{(\beta^\vee)^*} (\mathbb{C}^*)^n \rightarrow (\mathbb{C}^*)^{|J_\sigma|},$$

where $J_\sigma = \{j: \rho_j \not\subset \sigma\}$. Hence we shall write the inertia homomorphisms as $\omega_\sigma: \Gamma_\sigma \rightarrow \pi_1(\mathcal{X}(N, \Sigma, \beta))$.

As shown in [12], we may identify Γ_σ with $\text{Tor}(N/N_\sigma)$, the torsion submodule of the quotient N/N_σ , where $N_\sigma = \text{span}\{\beta(\epsilon_i) \mid \rho_i \subset \sigma\}$. Moreover, the inclusion $\Gamma_\sigma \rightarrow G$ may be modelled by an explicit homomorphism $\gamma_\sigma: \text{Tor}(N/N_\sigma) \rightarrow G$ constructed in [12]. Additionally, the quotient $G \rightarrow G/G_0$, which is obtained by applying $\text{Hom}(-, \mathbb{C}^*)$ to the inclusion of the torsion submodule $\text{Tor}(\text{DG}(\beta)) \hookrightarrow \text{DG}(\beta)$, may also be modelled by an explicit isomorphism $\text{Hom}(\text{Tor}(\text{DG}(\beta)), \mathbb{C}^*) \xrightarrow{\cong} \text{coker } \beta$. (See [12] for details.)

It is then straightforward to verify that the diagram

$$\begin{array}{ccccc} \text{Tor}(N/N_\sigma) & \hookrightarrow & N/N_\sigma & \hookrightarrow & N/\text{im}(\beta) \\ \cong \downarrow & \searrow \gamma_\sigma & & & \uparrow \cong \\ \Gamma_\sigma & \hookrightarrow & \text{Hom}(\text{DG}(\beta), \mathbb{C}^*) & \longrightarrow & \text{Hom}(\text{Tor}(\text{DG}(\beta)), \mathbb{C}^*) \end{array}$$

commutes; therefore, the inertia homomorphism may be identified with the composition in the top row, which proves the following.

Proposition 5.3. *Let (N, Σ, β) be a stacky fan and let σ be a cone in Σ . Using the identifications above, the inertia homomorphisms $\omega_\sigma: \Gamma_\sigma \rightarrow \pi_1(\mathcal{X}(N, \Sigma, \beta), z)$ may be identified with the composition*

$$\omega_\sigma: \text{Tor}(N/N_\sigma) \hookrightarrow N/N_\sigma \rightarrow \text{coker } \beta.$$

We may apply the above Proposition to characterize global quotients among toric DM stacks in terms of their stacky fan data, giving another proof of Corollary 5.7 below. By Theorem 4.10, it follows that $\mathcal{X}(N, \Sigma, \beta)$ is a global quotient if and only if the kernels $\ker \omega_\sigma = \text{Tor}(\text{im } \beta/N_\sigma)$ are trivial for all cones σ , if and only if $\text{im } \beta/N_\sigma$ is trivial for all maximal cones σ .

5.4. The universal cover of a toric DM stack associated to a stacky fan. By Proposition 4.8, the universal cover of the DM stack $\mathcal{X}(N, \Sigma, \beta)$ is $[Z_\Sigma/G_0]$, where G_0 is the connected component of the identity element of $G = \text{Hom}_{\mathbb{Z}}(\text{DG}(\beta), \mathbb{C}^*)$. Next we describe G_0 in terms of the stacky fan (N, Σ, β) . As we shall see, this can be roughly described as replacing the abelian group N with the image of β .

Let $N' \subset N$ denote the image of β , and let Σ' be the fan in $N' \otimes \mathbb{R} \cong N \otimes \mathbb{R}$ corresponding to Σ under the natural isomorphism $N' \otimes \mathbb{R} \cong N \otimes \mathbb{R}$. Finally, let $\beta' : \mathbb{Z}^n \rightarrow N'$ be β with restricted codomain. The following lemma is easily verified.

Lemma 5.4. *Let (N, Σ, β) be a stacky fan, and let (N', Σ', β') be defined as above. Then (N', Σ', β') is a stacky fan.*

Proposition 5.5. *Let (N, Σ, β) be a stacky fan, and let (N', Σ', β') be defined as above. Then the toric DM stack $\mathcal{X}(N', \Sigma', \beta')$ is the universal cover of $\mathcal{X}(N, \Sigma, \beta)$.*

Proof. By Proposition 4.8, it suffices to verify that the toric DM stack $\mathcal{X}(N', \Sigma', \beta') = [Z_\Sigma/G_0]$, where the group $G = \text{Hom}_{\mathbb{Z}}(\text{DG}(\beta), \mathbb{C}^*)$, which is verified in [12]. □

For a stacky fan (N, Σ, β) , given a cone σ in Σ , let $N_\sigma \subset N$ denote $\text{span}\{\beta(e_i) \mid \rho_i \subset \sigma\}$. The following lemma describes well-known conditions on a stacky fan (N, Σ, β) that characterize when the toric DM stack $\mathcal{X}(N, \Sigma, \beta)$ is in fact a smooth (toric) manifold. The corollary that follows then immediately characterizes global quotients among toric DM stacks.

Lemma 5.6. *Let (N, Σ, β) be a stacky fan. Then the toric DM stack $\mathcal{X}(N, \Sigma, \beta)$ is (equivalent to) a smooth manifold if and only if $N = N_\sigma$ for all maximal cones $\sigma \in \Sigma$.*

Proof. Recall that since $\mathcal{X}(N, \Sigma, \beta)$ is Deligne–Mumford, it admits an étale presentation, and the diagonal map $\Delta : \mathcal{X}(N, \Sigma, \beta) \rightarrow \mathcal{X}(N, \Sigma, \beta) \times \mathcal{X}(N, \Sigma, \beta)$ is proper therefore a closed embedding. By Proposition 74 in [19] it suffices to check that all isotropy groups are trivial. This follows from Theorem 4.2 [12]. □

Corollary 5.7. *Let (N, Σ, β) be a stacky fan. The following are equivalent:*

- (1) *The toric DM stack $\mathcal{X}(N, \Sigma, \beta)$ is equivalent to a global quotient over Diff.*
- (2) *$N' = N'_\sigma$ for all maximal cones σ in Σ .*

REMARK 5.8. Corollary 5.7 is also obtained in joint work of the authors with Goldin and Johanssen [12] by working out the combinatorial condition (2) directly from the equivalent condition that the connected component of the identity $G_0 \subset G = \text{Hom}(\text{DG}(\beta), \mathbb{C}^*)$ act freely on Z_Σ .

REMARK 5.9. A similar result to Proposition 5.5 was obtained by Poddar and Sarkar for quasi-toric orbifolds, which are effective/reduced orbifolds studied using methods from toric topology (see Theorem 3.2 in [23]).

We may recast the above in terms of stacky polytopes. Given a stacky polytope (N, Δ, β) , let $N' = \text{im } \beta$ and $\beta': \mathbb{Z}^n \rightarrow N'$ as before and let Δ' be the polytope in $(N' \otimes \mathbb{R})^*$ described by

$$\Delta' = \bigcap_{i=1}^n \{x \in (N' \otimes \mathbb{R})^* \mid \langle x, \beta'(e_i) \otimes 1 \rangle \geq -c_i\}$$

where the numbers c_1, \dots, c_n are the same as those appearing in (5.1) for the polytope Δ . This ensures that the corresponding level sets Z_Δ and Z'_Δ coincide. Equivalently, $\Delta' \subset (N' \otimes \mathbb{R})^*$ is the polytope corresponding to Δ under the dual of the natural identification $N' \otimes \mathbb{R} \cong N \otimes \mathbb{R}$. Analogous to Proposition 5.5, $\mathcal{X}(N', \Delta', \beta')$ is the universal cover of $\mathcal{X}(N, \Delta, \beta)$. In addition, the natural covering $p: \mathcal{X}(N', \Delta', \beta') \rightarrow \mathcal{X}(N, \Delta, \beta)$ (as in Proposition 4.8) is compatible with the underlying symplectic structures.

Proposition 5.10. *Let (N, Δ, β) be a stacky polytope, (N', Δ', β') be as above, and let $p: \mathcal{X}(N', \Delta', \beta') \rightarrow \mathcal{X}(N, \Delta, \beta)$ be the universal covering projection. If ω and ω' denote the symplectic forms on $\mathcal{X}(N, \Delta, \beta)$ and $\mathcal{X}(N', \Delta', \beta')$, respectively, then $p^*\omega = \omega'$.*

Proof. Recall that the symplectic form ω on a toric DM stack arising from a stacky polytope (N, Δ, β) can be identified with the differential 2-form $\xi^*\omega$ on Z_Δ where $\xi: Z_\Delta \rightarrow [Z_\Delta/G]$ is a presentation (see Proposition 2.9 in [18] and Theorem 14 (and the discussion preceding it) in [24]). Furthermore, if $\xi': Z_\Delta \rightarrow [Z_\Delta/G_0]$ denotes a presentation for $\mathcal{X}(N', \Delta', \beta')$ we have that $(\xi')^*\omega' = \xi^*\omega$, as they are each the restriction of the same 2-form on \mathbb{C}^n . Therefore, it suffices to verify that $(\xi')^*p^*\omega = \xi^*\omega$.

The natural isomorphism of G -bundles $(B \times G_0) \times_{G_0} G \cong B \times G$ (over any base B) gives rise to the 2-commutative diagram,

$$\begin{array}{ccc} & Z_\Delta & \\ \xi' \swarrow & & \searrow \xi \\ [Z_\Delta/G_0] & \xrightarrow{p} & [Z_\Delta/G] \end{array}$$

which shows that $(\xi')^*p^*\omega = \xi^*\omega$. Indeed, a differential form ω on a stack \mathcal{Y} is an assignment of a differential form $\omega(y) \in \Omega^*(U)$ for every object y over U that is compatible with maps, in the sense that an arrow $x \rightarrow y$ over $f: V \rightarrow U$ forces $\omega(x) = f^*\omega(y)$. It follows that $\xi^*\omega$ is a the 2-form assigned to the object $Z_\Delta \leftarrow Z_\Delta \times G \rightarrow Z_\Delta$ and that $(\xi')^*p^*\omega$ is the 2-form assigned to the object $Z_\Delta \leftarrow (Z_\Delta \times G_0) \times_{G_0} G \rightarrow$

Z_Δ . But the natural isomorphism covering the identity between these objects (i.e. the 2-isomorphism in the diagram above) and the compatibility condition forces these 2-forms to coincide. \square

Notice that the polytopes Δ' and Δ are the same up to a rescaling of the underlying lattices via the natural identification $N' \otimes \mathbb{R} \cong N \otimes \mathbb{R}$; therefore, their corresponding volumes satisfy the relation $\text{vol}(\Delta') = |N/N'| \text{vol}(\Delta)$. Corollary 5.11 below verifies a similar relation among the corresponding symplectic volumes.

Corollary 5.11. *Let (N, Δ, β) be a stacky polytope, and let (N', Δ', β') be as above. The corresponding symplectic volumes satisfy $\text{Vol}(\mathcal{X}(N', \Delta', \beta')) = |\text{coker } \beta| \cdot \text{Vol}(\mathcal{X}(N, \Delta, \beta))$.*

Proof. Let $\mathcal{X} = \mathcal{X}(N, \Delta, \beta)$ and $\mathcal{X}' = \mathcal{X}(N', \Delta', \beta')$ and $p: \mathcal{X}' \rightarrow \mathcal{X}$ denote the universal covering projection. Choose an étale presentation $\varphi: \underline{X}_0 \rightarrow \mathcal{X}$ with a partition of unity so that $\text{Vol}(\mathcal{X}) = \int_{X_0} \omega^d$ (see [3] for details about integration on stacks). Since p is a covering projection, the fiber product $\mathcal{X}' \times_{\mathcal{X}} X_0 \cong \underline{W}$ for some manifold W and $W \rightarrow X_0$ is a covering projection with fiber $G/G_0 \cong \text{coker } \beta$, where G_0 is the connected component of $G = \text{Hom}_{\mathbb{Z}}(\text{DG}(\beta), S^1)$. (In fact, as in Lemma 4.7, we may take $W = E_\varphi/G_0$ where $X_0 \leftarrow E_\varphi \rightarrow Z_\Delta$ is the object representing $\varphi(\text{id}_{X_0})$.) Then we have the following 2-Cartesian diagram

$$\begin{CD} W @>p_0>> X_0 \\ @V\varphi'VV @VV\varphi V \\ \mathcal{X}' @>p>> \mathcal{X} \end{CD}$$

It follows that $\varphi': \underline{W} \rightarrow \mathcal{X}'$ is an étale presentation for \mathcal{X}' . Moreover, we may pull back the partition of unity on X_0 to W . By the previous proposition,

$$\text{Vol}(\mathcal{X}') = \int_W (\varphi')^* p^* \omega = \int_W (p_0)^* \varphi^* \omega = \text{deg}(p_0) \cdot \int_{X_0} \varphi^* \omega = |G/G_0| \cdot \text{Vol}(\mathcal{X}). \quad \square$$

5.5. Examples. We conclude with some examples illustrating the discussion above.

The following class of examples is studied in [12].

EXAMPLE 5.12 (Labelled sheared simplices). Let $\mathbf{a} = (a_1, \dots, a_d)$ be a primitive vector in $N = \mathbb{Z}^d$ with $a_i \geq 1$ and let $m_0, \dots, m_d \in \mathbb{Z}_{>0}$. Let $\Sigma(\mathbf{a})$ be the fan in $N \otimes \mathbb{R} \cong \mathbb{R}^d$ whose rays are generated by $-\mathbf{a}$ and the standard basis vectors. Note that $\Sigma(\mathbf{a})$ is the normal fan of a *sheared simplex* $\Delta(\mathbf{a})$. Letting f_0, \dots, f_d be the standard basis vectors for \mathbb{Z}^{d+1} , set $\beta: \mathbb{Z}^{d+1} \rightarrow \mathbb{Z}^d$ to be $\beta(f_0) = -m_0 \mathbf{a}$ and $\beta(f_j) = m_j e_j$ for




Fig. 5.1. A fan $\Sigma(\mathbf{a}) \subset N \otimes \mathbb{R}$ and its corresponding labelled polytope $\Delta(\mathbf{a}) \subset (N \otimes \mathbb{R})^*$.




Fig. 5.2. The fan $\Sigma(\mathbf{a}')$ and corresponding labelled polytope $\Delta(\mathbf{a}')$ of the universal cover of the symplectic toric DM stack represented in Fig. 5.1.

$j = 1, \dots, d$. It is straightforward to verify that $(N, \Sigma(\mathbf{a}), \beta)$ is the stacky fan associated to the labelled polytope $(\Delta(\mathbf{a}), \{m_j\}_{j=0}^d)$. (See Fig. 5.1 illustrating a concrete example with $\mathbf{a} = (1, 2)$ and labels $m_0 = m_1 = 1$ and $m_2 = 2$.)

In [12], it is shown that the toric DM stack $\mathcal{X}(N, \Sigma(\mathbf{a}), \beta)$ is a global quotient if and only if $m_j = m_0 a_j$ for all $j = 1, \dots, d$. In this case, $N' = \bigoplus_{j=1}^d m_j \mathbb{Z} \subset N$ and that under an identification $N' \cong \mathbb{Z}^d$ we find Σ' is the fan in \mathbb{R}^d whose rays are generated by $-\sum e_j$ and the standard basis vectors e_1, \dots, e_d and that $\beta': \mathbb{Z}^{d+1} \rightarrow N' \cong \mathbb{Z}^d$ may be expressed by the formulas $\beta'(f_0) = -\sum e_j$, $\beta'(f_j) = e_j$. That is, $\mathcal{X}(N', \Sigma', \beta') = \mathbb{C}P^d$ so that $\mathcal{X}(N, \Sigma(\mathbf{a}), \beta)$ is a quotient of complex projective space whenever it is a global quotient. (See Fig. 5.2 illustrating the (stacky) fan and polytope of the universal cover of the symplectic toric DM stack whose fan and polytope appear in Fig. 5.1.)




Fig. 5.3. The labelled polytopes $\Delta(\mathbf{a})$ and $\Delta(\mathbf{a})'$ of a of a quotient of a Hirzebruch surface and its universal cover (a Hirzebruch surface), respectively.

EXAMPLE 5.13. Let $\mathbf{a} = (a_1, a_2)$ be a primitive vector in $N = \mathbb{Z}^2$ with $a_1, a_2 \geq 1$. Consider the fan Σ in $N \otimes \mathbb{R} \cong \mathbb{R}^2$ with four rays ρ_1, \dots, ρ_4 generated by $-\mathbf{a}, e_2, e_1, -e_2$ respectively, and maximal cones $\sigma_{12}, \sigma_{23}, \sigma_{34}$, and σ_{41} , where σ_{ij} denotes the two dimensional cone generated by ρ_i and ρ_j . Let m_1, \dots, m_4 be positive integers and let $\beta: \mathbb{Z}^4 \rightarrow N$ be

$$\beta = \begin{bmatrix} -m_1 a_1 & 0 & m_3 & 0 \\ -m_1 a_2 & m_2 & 0 & -m_4 \end{bmatrix}.$$

Note that the stacky fan (N, Σ, β) corresponds to a labelled right trapezoid. (See Fig. 5.3 (a) illustrating a concrete example with $\mathbf{a} = (1, 2)$.)

By Corollary 5.7, the toric DM stack $\mathcal{X}(N, \Sigma, \beta)$ is a global quotient if and only if $N' = N'_\sigma$ for all maximal cones σ . This occurs precisely when $m_1 a_1 = m_3, m_2 = m_4$, and $m_2 | m_1 a_2$. In this case, $N' = m_3 \mathbb{Z} \oplus m_2 \mathbb{Z} \subset N$ and that under an identification $N' \cong \mathbb{Z}^2$ we see that Σ' is the fan in \mathbb{R}^2 with four rays generated by $-(1, b), e_2, e_1$, and $-e_2$, where $b = m_1 a_2 / m_2$. Moreover, under this identification

$$\beta' = \begin{bmatrix} -1 & 0 & 1 & 0 \\ -b & 1 & 0 & -1 \end{bmatrix};$$

therefore, $\mathcal{X}(N, \Sigma, \beta)$ is a quotient of a Hirzebruch surface whenever it is a global quotient. (See Fig. 5.3 illustrating the labelled polytope of a global quotient of a Hirzebruch surface.)

The remaining three examples consider toric DM stacks with non-trivial global stabilizer (due to the presence of torsion in the abelian group N). The first example exhibits a global quotient with global isotropy, while the last two illustrate how the condition in Corollary 5.7 may fail.

EXAMPLE 5.14. Let $N = \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ and Σ be the fan in $N \otimes \mathbb{R} \cong \mathbb{R}$ with rays ± 1 . Let $\beta: \mathbb{Z}^2 \rightarrow N$ be given by $(a, b) \mapsto (2a - 2b, a + b \bmod 2)$. Then N' is generated by $(2, 1 \bmod 2)$, as is N'_σ for each maximal cone σ . Therefore, $\mathcal{X}(N, \Sigma, \beta)$ is a global quotient. Under an identification $N' \cong \mathbb{Z}$, we see that $\beta'(a, b) = a - b$ and that $\mathcal{X}(N', \Sigma', \beta') = \mathbb{C}P^1$. Moreover, a direct calculation shows that $G/G_0 \cong \text{coker } \beta \cong \mathbb{Z}/4\mathbb{Z}$ so that $\mathbb{Z}/4$ acts on $\mathbb{C}P^1$ with global stabilizer isomorphic to $\mathbb{Z}/2\mathbb{Z}$ (the torsion submodule of N), and $\mathcal{X}(N, \Sigma, \beta) \cong [\mathbb{C}P^1/(\mathbb{Z}/4\mathbb{Z})]$.

EXAMPLE 5.15. Let (N, Σ, β) be any stacky fan with N containing a non-trivial torsion subgroup and β surjective (e.g. the stacky fan of a weighted projective space with non-trivial global stabilizer). Then $\mathcal{X}(N, \Sigma, \beta)$ is not equivalent to a global quotient since $N' = N$ has torsion and N'_σ is necessarily torsion free for any cone σ in the rational simplicial fan Σ (cf. the proof of Theorem 3.1 in [12]). (More generally, if $N' \subset N$ contains non-trivial torsion, then $\mathcal{X}(N, \Sigma, \beta)$ is not equivalent to a global quotient.)

EXAMPLE 5.16. Let $N = \mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$ and Σ be the fan in $N \otimes \mathbb{R} \cong \mathbb{R}$ with rays ± 1 . Let $\beta: \mathbb{Z}^2 \rightarrow N$ be given by $(a, b) \mapsto (a - 2b, a + 2b \bmod 4)$. Then N' is torsion-free, generated by $(1, 1) \in N$; however, N'_σ is generated by $(2, 2)$ for the cone σ generated by -1 . By Corollary 5.7, $\mathcal{X}(N, \Sigma, \beta)$ is not equivalent to a global quotient.

ACKNOWLEDGEMENTS. We thank Anthony Bahri for asking us the question which was the catalyst for this project, and Andrew Nicas for continuing support and many useful conversations.

References

- [1] A. Adem, J. Leida and Y. Ruan: *Orbifolds and Stringy Topology*, Cambridge Tracts in Mathematics **171**, Cambridge Univ. Press, Cambridge, 2007.
- [2] A. Adem and Y. Ruan: *Twisted orbifold K-theory*, Comm. Math. Phys. **237** (2003), 533–556.
- [3] K. Behrend: *Cohomology of stacks*; in *Intersection Theory and Moduli*, ICTP Lect. Notes **19**, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004, 249–294.
- [4] K. Behrend, B. Conrad, D. Edidin, B. Fantechi, W. Fulton, L. Göttsche and A. Kresch: *Algebraic stacks*, in progress.
- [5] L.A. Borisov, L. Chen and G.G. Smith: *The orbifold Chow ring of toric Deligne–Mumford stacks*, J. Amer. Math. Soc. **18** (2005), 193–215.
- [6] D. Edidin: *What is a stack?*, Notices, Amer. Math. Soc. **50** (2003), 458–459.
- [7] B. Fantechi: *Stacks for everybody*; in *European Congress of Mathematics, I* (Barcelona, 2000), Progr. Math. **201**, Birkhäuser, Basel, 2001, 349–359.
- [8] B. Fantechi and L. Göttsche: *Orbifold cohomology for global quotients*, Duke Math. J. **117** (2003), 197–227.
- [9] B. Fantechi, E. Mann and F. Nironi: *Smooth toric Deligne–Mumford stacks*, J. Reine Angew. Math. **648** (2010), 201–244.

- [10] A. Geraschenko and M. Satriano: *Toric stacks I: The theory of stacky fans*, (2011), arXiv:1107.1906.
- [11] A. Geraschenko and M. Satriano: *Toric stacks II: Intrinsic characterization of toric stacks*, (2011), arXiv:1107.1907.
- [12] R. Goldin, M. Harada, D. Johannsen and D. Krepski: *Inertia groups of a toric DM stack, fake weighted projective spaces, and labelled sheared simplices*, to appear in Rocky Mountain J. Math., arXiv:1306.5036.
- [13] D. Husemoller: *Fibre Bundles*, third edition, Graduate Texts in Mathematics **20**, Springer, New York, 1994.
- [14] I. Iwanari: *Logarithmic geometry, minimal free resolutions and toric algebraic stacks*, Publ. Res. Inst. Math. Sci. **45** (2009), 1095–1140.
- [15] T. Kawasaki: *Cohomology of twisted projective spaces and lens complexes*, Math. Ann. **206** (1973), 243–248.
- [16] E. Lerman and S. Tolman: *Hamiltonian torus actions on symplectic orbifolds and toric varieties*, Trans. Amer. Math. Soc. **349** (1997), 4201–4230.
- [17] E. Lerman: *Orbifolds as stacks?*, Enseign. Math. (2) **56** (2010), 315–363.
- [18] E. Lerman and A. Malkin: *Hamiltonian group actions on symplectic Deligne–Mumford stacks and toric orbifolds*, Adv. Math. **229** (2012), 984–1000.
- [19] D. Metzler: *Topological and smooth stacks*, (2003), arXiv:math/0306176.
- [20] B. Noohi: *Fundamental groups of algebraic stacks*, J. Inst. Math. Jussieu **3** (2004), 69–103.
- [21] B. Noohi: *Foundations of topological stacks I*, arXiv:math/0503247.
- [22] F. Perroni: *A note on toric Deligne–Mumford stacks*, Tohoku Math. J. (2) **60** (2008), 441–458.
- [23] M. Poddar and S. Sarkar: *On quasitoric orbifolds*, Osaka J. Math. **47** (2010), 1055–1076.
- [24] H. Sakai: *The symplectic Deligne–Mumford stack associated to a stacky polytope*, Results in Mathematics (2012).
- [25] I. Satake: *On a generalization of the notion of manifold*, Proc. Nat. Acad. Sci. U.S.A. **42** (1956), 359–363.

Megumi Harada
Department of Mathematics and Statistics
McMaster University
1280 Main Street West
Hamilton, Ontario L8S4K1
Canada
e-mail: Megumi.Harada@math.mcmaster.ca
<http://www.math.mcmaster.ca/Megumi.Harada/>

Derek Krepski
Department of Mathematics
University of Western Ontario
Middlesex College
London, Ontario N6A 5B7
Canada
Current address:
Department of Mathematics
University of Manitoba
342 Machray Hall, 186 Dysart Road
Winnipeg, Manitoba R3T2N2
Canada
e-mail: Derek.Krepski@umanitoba.ca
<http://server.math.umanitoba.ca/~dkrepski/>