

Title	Search for X-particle as a bound state of Kaon and pion
Author(s)	Fahmida, Khanam
Citation	大阪大学, 2009, 博士論文
Version Type	
URL	https://hdl.handle.net/11094/58012
rights	
Note	著者からインターネット公開の許諾が得られていないため、論文の要旨のみを公開しています。全文のご利用をご希望の場合は、 〈a href="https://www.library.osaka-u.ac.jp/thesis/#closed"〉 大阪大学の博士論文について 〈/a〉 をご参照ください。

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

[4]

氏名	ファミダ カナム Fahmida Khanam
博士の専攻分野の名称	博士(理学)
学位記番号	第 23333 号
学位授与年月日	平成21年9月25日
学位授与の要件	学位規則第4条第1項該当 理学研究科物理学専攻
学位論文名	Search for X-particle as a bound state of Kaon and pion (K中間子とpi中間子の束縛状態におけるX粒子の探索)
論文審査委員	(主査) 教授 岸本 忠史 (副査) 教授 能町 正治 教授 野海 博之 准教授 阪口 篤志 准教授 奥曾井 優

論文内容の要旨

The experiment at KEK-PS E548 was performed to search a new “Scalar particle, X” which was predicted as a bound system of “ $K\pi$ ”, that could explain the properties, mainly the narrow width of penta-quark Θ^+ , by predicting it as a $K\pi N$ bound system. We used 1.2 GeV/c K^+ beam on the proton target, to study $p(K^+, X^+)p$ reaction, where X-particle decay to $X^+ \rightarrow K^+\gamma\gamma$, in the experiment we took data with outgoing K^+ and one γ . We studied possible background and signal region, which was calculated from simulation. One of the main background was from the production of π^0 and its decay $\pi^0 \rightarrow \gamma\gamma$, which gave similar products at the final stage. As K^+ momentum from π^0 background can have maximum of 1.0 GeV/c, so we mainly studied the higher K^+ region (above 1.0 GeV/c) to find the signal corresponding to X-particle.

論文審査の結果の要旨

The experiment at KEK-PS E548 was performed to search a new “Scalar particle, X” which was predicted as a bound system of “ $K\pi$ ”, that could

explain the properties, mainly the narrow width of penta-quark Θ^+ , by predicting it as a $K\pi N$ bound system. We used 1.2 GeV/c K^+ beam on the CH_2 target, to study mainly the $p(K^+, X^+)p$ reaction, where X -particle decay to $X^+ \rightarrow K^+\gamma\gamma$. In the experiment we measured outgoing K^+ and γ rays in coincidence with the K^+ . One of the main background was from the production of π^0 and its decay $\pi^0 \rightarrow \gamma\gamma$, which gave similar products in the final stage. As K^+ momentum from π^0 background can have maximum of 1.0 GeV/c, so we mainly studied the K^+ momentum region above 1.0 GeV/c to find the signal corresponding to X -particle. We observed events in the signal region. We studied possible origin of the backgrounds and found no known physical processes that can give the events in the signal region. This study indicates observation of the X particle. The result is so important that it will open further study on this direction.

よって、本論文は博士（理学）の学位論文として十分価値あるものと認める。