



Title	SUMMATION FORMULA INEQUALITIES FOR EIGENVALUES OF THE PERTURBED HARMONIC OSCILLATOR
Author(s)	Fereitas, Pedro; Kennedy, James B
Citation	Osaka Journal of Mathematics. 2016, 53(2), p. 397-416
Version Type	VoR
URL	https://doi.org/10.18910/58874
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

SUMMATION FORMULA INEQUALITIES FOR EIGENVALUES OF THE PERTURBED HARMONIC OSCILLATOR

PEDRO FREITAS and JAMES B. KENNEDY

(Received July 25, 2014, revised January 27, 2015)

Abstract

We derive explicit inequalities for sums of eigenvalues of one-dimensional Schrödinger operators on the whole line. In the case of the perturbed harmonic oscillator, these bounds converge to the corresponding trace formula in the limit as the number of eigenvalues covers the whole spectrum.

1. Introduction

Consider the eigenvalue equation

$$(1.1) \quad -u''(x) + V(x)u(x) = \lambda u(x), \quad x \in (a, b) \subseteq \mathbb{R},$$

associated with a one-dimensional Schrödinger operator $H = -d^2/dx^2 + V$, where the potential $V: (a, b) \rightarrow \mathbb{R}$, and the boundary condition if $(a, b) \neq \mathbb{R}$, are chosen such that the spectrum consists of a discrete sequence of eigenvalues $\{\lambda_k\}$. One possible way of linking the behaviour of this sequence to properties of the potential V is via a regularized trace formula for the sum of the eigenvalues. The classical example is the formula attributed to Gelfand and Levitan, which, if we take $(a, b) = (0, \pi)$ with Dirichlet boundary conditions on the endpoints, reads

$$(1.2) \quad \sum_{k=1}^{\infty} \left[\lambda_k - k^2 - \frac{1}{\pi} \int_0^{\pi} V(x) dx \right] = \frac{1}{2\pi} \int_0^{\pi} V(x) dx - \frac{V(0) + V(\pi)}{4}$$

(see, e.g., the book [6], also for other similar formulae). Since the values k^2 are in fact the eigenvalues of the Dirichlet Laplacian, that is, the corresponding Schrödinger operator with zero potential, this is a comparison between the eigenvalues of the operators H and $H_0 := -d^2/dx^2$.

More recently it has also been shown that an analogous trace formula holds for the eigenvalues of (1.1) on the whole line $(a, b) = \mathbb{R}$ [2, 7]. The comparison case is

2010 Mathematics Subject Classification. Primary 34L15; Secondary 33C45, 34L20, 34L40, 81Q15.

J.B.K. was supported by a fellowship of the Alexander von Humboldt Foundation, Germany. Both authors were partially supported by FCT's project PEst-OE/MAT/UI0208/2011.

now provided by the quantum harmonic oscillator

$$(1.3) \quad -u''(x) + x^2 u(x) = \lambda u(x), \quad x \in \mathbb{R},$$

whose eigenvalues are given by $\lambda_k^0 = 2k + 1$ for $k \in \mathbb{N}$. Writing the potential in (1.1) as $V(x) = x^2 + q(x)$, that is, as a perturbed harmonic oscillator, if the perturbation $q: \mathbb{R} \rightarrow \mathbb{R}$ is small enough in an appropriate sense, then the eigenvalues of (1.1), which we denote by λ_k for $k \in \mathbb{N}$, satisfy the trace formula

$$(1.4) \quad \sum_{k=0}^{\infty} \left[\lambda_k - \lambda_k^0 - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx \right] = -\frac{Z_0(1/2)}{\pi} \int_{\mathbb{R}} q(x) dx,$$

where

$$(1.5) \quad Z_0(s) = (1 - 2^{-s})\zeta(s) = \sum_{k=1}^{\infty} \frac{1}{(\lambda_k^0)^s}$$

is the spectral zeta function associated with (1.3), the second equality being valid for $\operatorname{Re} s > 1$, and $\zeta(\cdot)$ is the Riemann zeta function; see [2, Theorem 2] or [7, Equation (1.12)]. We refer to [8] for a wide-ranging general survey on the theory of regularized traces.

In a separate paper [5] we show that formula (1.2) is in fact the limit as $n \rightarrow \infty$ of a sequence of inequalities for the (finite) sums of the first n eigenvalues given in terms of the Fourier coefficients of the potential, and that (1.2) can be proved by combining these inequalities with knowledge of the asymptotic behaviour of the eigenvalues and eigenfunctions [5]. In the present paper, which may be viewed as a continuation of [5], we show that a similar family of inequalities is valid for the perturbed harmonic oscillator assuming that the perturbation q is non-negative and of finite $L^1(\mathbb{R})$ -norm. More precisely, we shall prove in Theorem 3.1 below that there is a sequence of inequalities of the form

$$\sum_{k=0}^n \left[\lambda_k - \lambda_k^0 - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx \right] \leq \frac{\chi_n}{\pi} \int_{\mathbb{R}} q(x) dx$$

for all $n \in \mathbb{N}$ if $V(x) = x^2 + q(x)$ with $0 \leq q \in L^1(\mathbb{R})$, where the sequence χ_n , which is given explicitly, depends only on properties of the eigenfunctions and eigenvalues of the quantum harmonic oscillator (1.3) and converges to $-Z_0(1/2)$ like $O(1/\sqrt{n})$ as $n \rightarrow \infty$. A similar sequence of bounds will also be shown to hold for a certain class of negative or indefinite potentials (see Theorem 4.1), and although the corresponding bounding sequence we obtain is larger than χ_n , it is still explicit, and the order of convergence to the known trace formula remains $O(1/\sqrt{n})$.

These results will be established via test function methods, using for this purpose the eigenfunctions of (1.3) in a suitable Rayleigh quotient expression for the eigenvalues of

the perturbed harmonic oscillator, and then combining this with properties of Hermite polynomials to analyze the resulting expression. We believe one of these properties, namely Lemma 3.3, which provides an upper bound for the function $e^{-x^2}[H_{n+1}^2(x) - H_n(x)H_{n+2}(x)]$ to be new and interesting in its own right.

In fact, these results—and the corresponding proofs—differ from those in [5] in that for them we do not use a decomposition of the potential in terms of the eigenfunctions of the unperturbed problem. However, such an approach is also possible in this case and we carry it out to obtain a different type of bound; see Theorem 5.1. For this particular result we assume that $V \in L^2(\mathbb{R}, e^{-x^2} dx)$, that is, that the potential is no longer necessarily a perturbation of x^2 , but rather more generally merely square integrable with respect to the weighted L^2 -measure most naturally associated with the problem (1.3). The resulting bounds (which are once again explicit) are expressed in terms of the Fourier-like coefficients of V expanded as a sum of Hermite polynomials. These are actually stronger than Theorems 3.1 and 4.1, as the only inequality used now is that which arises from the substitution of test functions in the Rayleigh quotient (see Remark 5.2 (i)). However, now the finite sums converging to the the left-hand side of the trace formula (1.4) do not appear in a natural way; this will then be derived as a simple corollary by writing the potential $V(x)$ as $x^2 + q(x)$ and using the Fourier coefficients for q instead.

We also generalize Theorems 3.1 and 4.1 to obtain bounds on sums of powers of the eigenvalues in Section 6.

2. Schrödinger operators on the real line

Throughout this paper we will consider one-dimensional Schrödinger operators on the real line, that is, associated with the equation (1.1) for $x \in \mathbb{R}$, where the potential $V: \mathbb{R} \rightarrow \mathbb{R}$ is a locally measurable function on which we will impose various (and varying) assumptions. We will always assume that $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$, so that the operator associated with the problem (1.1) considered as an operator on $L^2(\mathbb{R})$ has discrete spectrum, and we will in general denote the associated eigenvalues by $\lambda_0 < \lambda_1 \leq \dots \rightarrow \infty$.

As is well known, the eigenvalues of the quantum harmonic oscillator (1.3), which will play the role of our “default” problem, are given by $\lambda_k^0 = 2k + 1$ for $k \in \mathbb{N}$, with corresponding eigenfunctions $\psi_k(x) = e^{-x^2/2}H_k(x)$, which form an orthonormal basis of $L^2(\mathbb{R})$. Here H_k denotes the k^{th} Hermite polynomial (see, e.g., [10, Chapter 5]).

Of particular interest to us will be the perturbed harmonic oscillator

$$(2.1) \quad -u''(x) + [x^2 + q(x)]u(x) = \lambda u(x), \quad x \in \mathbb{R},$$

which is easily seen to have discrete spectrum if $q \in L^p(\mathbb{R})$ for some $p \in [1, \infty]$.

For a general potential $V: \mathbb{R} \rightarrow \mathbb{R}$, we can characterize the associated eigenvalues via classical variational methods. Denoting by $\varphi \in H^1(\mathbb{R}) \cap L^2(\mathbb{R}, V(x) dx)$ an arbitrary

test function, we let

$$(2.2) \quad \mathcal{R}[V, \varphi] := \frac{\int_{\mathbb{R}} (\varphi'(x))^2 dx + \int_{\mathbb{R}} V(x) \varphi^2(x) dx}{\int_{\mathbb{R}} \varphi^2(x) dx}$$

be the Rayleigh quotient associated with (the Schrödinger operator with potential) V at φ . A standard generalization of the usual minimax formula for eigenvalues states that if $\varphi_0, \dots, \varphi_n$ is a collection of $n+1$ such functions orthogonal in $L^2(\mathbb{R})$, for any $n \in \mathbb{N}$, then

$$\sum_{k=0}^n \lambda_k \leq \sum_{k=0}^n \mathcal{R}[V, \varphi_k]$$

(see, e.g., [3]), with equality being achieved when the φ_k are the first $n+1$ eigenfunctions. For us the most natural choice of test functions will be the eigenfunctions ψ_k of the quantum harmonic operator.

3. Bounds for the perturbed harmonic oscillator with a non-negative perturbation

In this section we will state and prove our main theorem, obtaining the aforementioned finite version of the trace formula (1.4) for the general perturbed harmonic oscillator (2.1).

Theorem 3.1. *Let q be a non-negative potential defined on the real line having finite $L^1(\mathbb{R})$ norm. Then the eigenvalues of (2.1) satisfy the inequalities*

$$(3.1) \quad \sum_{k=0}^n \left[\lambda_k - \lambda_k^0 - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx \right] \leq \frac{\chi_n}{\pi} \int_{\mathbb{R}} q(x) dx, \quad n = 0, 1, \dots,$$

where

$$(3.2) \quad \chi_n = \begin{cases} \frac{2n+3}{n+1} \frac{\Gamma(n/2+1)}{\Gamma((n+1)/2)} - \sum_{k=0}^n \frac{1}{\sqrt{\lambda_k^0}}, & n \text{ odd}, \\ (n+1) \frac{\Gamma((n+1)/2)}{\Gamma(n/2+1)} - \sum_{k=0}^n \frac{1}{\sqrt{\lambda_k^0}}, & n \text{ even}. \end{cases}$$

Furthermore, $\chi_n = -Z_0(1/2) + O(1/\sqrt{n})$, where $Z_0(s) = (1 - 2^{-s})\zeta(s)$.

REMARK 3.2. (i) It is essential for our method of proof that q be non-negative. In Theorem 4.1 below, we weaken this assumption and obtain a slightly weaker set

of inequalities which nevertheless still converge in the limit to the trace formula (1.4) with the same order of convergence $O(1/\sqrt{n})$. It is not clear if the inequalities (3.1) are true for arbitrary $q \in L^1(\mathbb{R})$; the trace formula (1.4) is itself currently only known to hold under stronger assumptions on q : in [2] a certain rate of decay of q at infinity is assumed, and in [7] it is assumed q has compact support. We remark however that having convergence of order $O(1/\sqrt{n})$ is most probably optimal, since this is the rate at which we have convergence of the sequence whose limit defines $\zeta(1/2)$ (cf. (3.9) and (3.10)).

(ii) There do not exist corresponding lower bounds for *finite* sums of eigenvalues: for any fixed $n \geq 0$ is it always possible to find a function $0 \leq q \in L^1(\mathbb{R})$ for which the left-hand side of (3.1) is arbitrarily large negative; see Proposition 3.4 below. However, for a *fixed* potential it is a natural question as to whether we can recover a lower bound valid in the asymptotic limit. Indeed, it might be possible to extend our result to give a new proof of the trace formula (1.4) for a different class of (non-negative) potentials q from those considered in [2, 7], namely $q \in L^1(\mathbb{R})$. The idea would be to argue as in [4] (or [5]), to show that the degree of “error” which arises from using the eigenfunctions ψ_k of the unperturbed problem as test functions becomes asymptotically small as $k \rightarrow \infty$: denoting by φ_k the eigenfunction associated with λ_k (corresponding to the potential $V(x) = x^2 + q(x)$), we see that the trace formula holds whenever

$$(3.3) \quad \lim_{n \rightarrow \infty} \sum_{k=0}^n (\mathcal{R}[x^2 + q(x), \varphi_k] - \mathcal{R}[x^2 + q(x), \psi_k]) = 0,$$

since by definition $\mathcal{R}[x^2 + q(x), \varphi_k] = \lambda_k$. We can rewrite (3.3) as a type of “change of basis” formula

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n (\langle \varphi_k, H\varphi_k \rangle - \langle \psi_k, H\psi_k \rangle) = 0,$$

where $H: D(H) \subset L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ is the operator associated with the potential $x^2 + q(x)$. We *expect* this to hold whenever the asymptotics for λ_k and φ_k are similar enough to those of λ_k^0 and ψ_k , respectively, when $k \rightarrow \infty$. This is, however, likely to be a difficult problem, and we shall not attempt an investigation of it here.

For notational convenience, for $n \geq 0$ we define

$$(3.4) \quad \omega_n := \chi_n + \sum_{k=0}^n \frac{1}{\sqrt{\lambda_k^0}} = \begin{cases} \frac{2n+3}{n+1} \frac{\Gamma(n/2+1)}{\Gamma((n+1)/2)}, & n \text{ odd}, \\ (n+1) \frac{\Gamma((n+1)/2)}{\Gamma(n/2+1)}, & n \text{ even}, \end{cases}$$

and we also set $\omega_{-1} := 0$.

Proof of Theorem 3.1. Using the first $n + 1$ eigenfunctions of the unperturbed harmonic oscillator (1.3), given by $\psi_k(x) = e^{-x^2/2} H_k(x)$, $k = 0, \dots, n$, as test functions in the Rayleigh quotient (2.2) for $V(x) = x^2 + q(x)$ yields

$$(3.5) \quad \begin{aligned} \sum_{k=0}^n \lambda_k &\leq \sum_{k=0}^n \frac{\int_{\mathbb{R}} [(d/dx)[e^{-x^2/2} H_k(x)]]^2 + [x^2 + q(x)]e^{-x^2} H_k^2(x) dx}{\int_{\mathbb{R}} e^{-x^2} H_k^2(x) dx} \\ &= \sum_{k=0}^n \lambda_k^0 + \int_{\mathbb{R}} e^{-x^2} q(x) \sum_{k=0}^n \frac{1}{2^k k! \sqrt{\pi}} H_k^2(x) dx. \end{aligned}$$

From basic properties of Hermite polynomials we have the identity

$$(3.6) \quad \sum_{k=0}^n \frac{1}{2^k k!} H_k^2(x) = \frac{1}{2^{n+1} n!} [H_{n+1}^2(x) - H_n(x)H_{n+2}(x)].$$

This arises in the context of Turán's inequality for Hermite polynomials (cf. [9, p. 404]), and can easily be derived directly by induction in n —see also, for instance, [10, p. 106]. By using the estimate of the function

$$(3.7) \quad h_n(x) := e^{-x^2} [H_{n+1}^2(x) - H_n(x)H_{n+2}(x)]$$

given in Lemma 3.3 below in (3.6) and inserting this into (3.5), we obtain

$$(3.8) \quad \begin{aligned} \sum_{k=0}^n \lambda_k &\leq \sum_{k=0}^n \lambda_k^0 + \int_{\mathbb{R}} e^{-x^2} q(x) \sum_{k=0}^n \frac{1}{2^k k! \sqrt{\pi}} H_k^2(x) dx \\ &\leq \sum_{k=0}^n \lambda_k^0 + \frac{\omega_n}{\pi} \int_{\mathbb{R}} q(x) dx, \end{aligned}$$

which upon rearranging yields (3.1).

We now give the (routine) proof that $\chi_n = -Z_0(1/2) + O(1/\sqrt{n})$ as $n \rightarrow \infty$. We first note that

$$(3.9) \quad \zeta(s) = \sum_{k=1}^n k^{-s} + s \int_n^\infty \frac{\lfloor x \rfloor - x + 1/2}{x^{s+1}} dx + \frac{n^{1-s}}{s-1} - \frac{1}{2n^s},$$

valid for $s > 0$ (see [11], Equation (3.5.3), pp. 49–50). Setting $s = 1/2$ and passing to the limit as $n \rightarrow \infty$, this means we can write

$$(3.10) \quad -Z_0(1/2) = -\left(1 - \frac{1}{\sqrt{2}}\right) \zeta(1/2) = \left(1 - \frac{1}{\sqrt{2}}\right) \lim_{n \rightarrow \infty} a_n,$$

where for ease of notation we have set

$$(3.11) \quad a_n := 2\sqrt{n} - \sum_{k=1}^n \frac{1}{\sqrt{k}}$$

for $n \geq 1$. Now, recalling that $\lambda_k^0 = 2k + 1$ for $k \in \mathbb{N}$, we have

$$(3.12) \quad \chi_n = \omega_n - \sum_{k=0}^n \frac{1}{\sqrt{2k+1}} = \omega_n - \left(1 - \frac{1}{\sqrt{2}}\right) \sum_{k=1}^n \frac{1}{\sqrt{k}} - \sum_{k=n+1}^{2n+1} \frac{1}{\sqrt{k}};$$

we wish to show that this converges to $-Z_0(1/2)$ as $n \rightarrow \infty$. We first establish that

$$(3.13) \quad \omega_n = \sqrt{2n} + O\left(\frac{1}{\sqrt{n}}\right),$$

using the following asymptotics for the quotient of two gamma functions (see [1], formula 6.1.47, for instance):

$$(3.14) \quad \frac{\Gamma(z + 1/2)}{\Gamma(z)} = \sqrt{z} + O\left(\frac{1}{\sqrt{z}}\right),$$

for large z . For n odd we obtain

$$\omega_n = \frac{2n+3}{n+1} \left[\sqrt{\frac{n+1}{2}} + O\left(\frac{1}{\sqrt{n}}\right) \right] = \sqrt{2n} + O\left(\frac{1}{\sqrt{n}}\right).$$

A similar calculation when n is even gives

$$\omega_n = 2 \frac{\Gamma((n+3)/2)}{\Gamma(n/2+1)} = 2 \left[\sqrt{\frac{n}{2} + 1} + O\left(\frac{1}{\sqrt{n}}\right) \right],$$

proving (3.13). Next, we observe that

$$\sum_{k=n+1}^{2n+1} \frac{1}{\sqrt{k}} = 2(\sqrt{2} - 1)\sqrt{n} + O\left(\frac{1}{\sqrt{n}}\right)$$

for large n , as can be seen, for example, by noting that

$$\int_{n+1}^{2n+2} \frac{1}{\sqrt{x}} dx \leq \sum_{k=n+1}^{2n+1} \frac{1}{\sqrt{k}} \leq \int_{n+1}^{2n+2} \frac{1}{\sqrt{x-1}} dx$$

and evaluating the integrals. Substituting these two estimates into (3.12) yields

$$\begin{aligned}\chi_n &= \sqrt{2n} - \left(1 - \frac{1}{\sqrt{2}}\right) \sum_{k=1}^n \frac{1}{\sqrt{k}} - 2(\sqrt{2} - 1)\sqrt{n} + O\left(\frac{1}{\sqrt{n}}\right) \\ &= \left(1 - \frac{1}{\sqrt{2}}\right) a_n + O\left(\frac{1}{\sqrt{n}}\right).\end{aligned}$$

Letting s equal $1/2$ in (3.9) and using $-1 < \lfloor x \rfloor - x \leq 0$ we obtain

$$-\frac{1}{\sqrt{n}} < \zeta(1/2) + a_n \leq 0,$$

from which it follows that

$$\chi_n = -Z_0(1/2) + O\left(\frac{1}{\sqrt{n}}\right),$$

as desired. \square

Lemma 3.3. *The function h_n defined by (3.7) is positive and satisfies*

$$h_n(x) \leq \begin{cases} \frac{4^{n+1}}{2\pi} \frac{2n+3}{n+1} \Gamma^2\left(\frac{n}{2} + 1\right), & n \text{ odd}, \\ \frac{4^{n+1}}{2\pi} (n+1) \Gamma^2\left(\frac{n+1}{2}\right), & n \text{ even}. \end{cases}$$

Proof. Positivity of h_n is a direct consequence of (3.6). Taking derivatives in x and using the property $H'_n(x) = 2nH_{n-1}(x)$ yields

$$\begin{aligned}h'_n(x) &= e^{-x^2} \{ -2x[H_{n+1}^2(x) - H_n(x)H_{n+2}(x)] \\ &\quad + 2H_{n+1}(x)H'_{n+1}(x) - H'_n(x)H_{n+2}(x) - H_n(x)H'_{n+2}(x) \} \\ &= e^{-x^2} \{ 2H_{n+1}(x)[-xH_{n+1}(x) + 2(n+1)H_n(x)] \\ &\quad + 2xH_n(x)H_{n+2}(x) - 2nH_{n-1}(x)H_{n+2}(x) \\ &\quad - 2(n+2)H_n(x)H_{n+1}(x) \} \\ &= e^{-x^2} \{ 2H_{n+1}(x)[-xH_{n+1}(x) + nH_n(x)] \\ &\quad + 2H_{n+2}[xH_n(x) - nH_{n-1}(x)] \}.\end{aligned}$$

Using the identity $H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x)$ in the above expression yields

$$h'_n(x) = -2e^{-x^2} H_n(x)H_{n+1}(x),$$

which integrated between zero and x becomes

$$\begin{aligned}
 h_n(x) - h_n(0) &= -2 \int_0^x e^{-t^2} H_n(t) H_{n+1}(t) dt \\
 &= -\frac{1}{2(n+1)} \int_0^x e^{-t^2} \frac{d}{dt} H_{n+1}^2(t) dt \\
 &= -\frac{1}{2(n+1)} \left[e^{-x^2} H_{n+1}^2(x) - H_{n+1}^2(0) + 2 \int_0^x t e^{-t^2} H_{n+1}^2(t) dt \right].
 \end{aligned}$$

Noting that the terms which depend on x on the right-hand side above are non-positive, we obtain

$$(3.15) \quad h_n(x) - h_n(0) \leq \frac{1}{2(n+1)} H_{n+1}^2(0).$$

For odd n , $h_n(0) = H_{n+1}^2(0)$ and the above becomes

$$\begin{aligned}
 h_n(x) &\leq \frac{2n+3}{2n+2} H_{n+1}^2(0) \\
 &= \frac{2n+3}{2n+2} \Gamma^2(n+2) \Gamma^{-2}\left(\frac{n+3}{2}\right) \\
 &= \frac{4^{n+1}}{2\pi} \frac{2n+3}{n+1} \Gamma^2\left(\frac{n}{2}+1\right).
 \end{aligned}$$

For even values of n the right-hand side of (3.15) vanishes and we obtain

$$h_n(x) \leq h_n(0) = \frac{4^{n+1}}{2\pi} (n+1) \Gamma^2\left(\frac{n+1}{2}\right). \quad \square$$

We will now construct an example showing that no lower bound of the same form as in Theorem 3.1 is possible.

Proposition 3.4. *For any $n \geq 0$ and any $N > 0$, there exists $0 \leq q \in L^1(\mathbb{R})$ such that*

$$\sum_{k=0}^n \left[\lambda_k - \lambda_k^0 - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx \right] \leq -N.$$

Before giving the proof, we note two points: firstly, that there *exists* a potential for which the corresponding first n eigenvalues are arbitrarily large negative is trivial; the key point here is that q satisfies the same assumptions as in Theorem 3.1. Secondly, the sum here has to be regularized, since for any $q \geq 0$ we automatically have $\lambda_k \geq \lambda_k^0$ for all $k \geq 0$.

Proof. Fix $n \geq 0$ and $N > 0$. If we use the $n + 1$ functions $\psi_k(x) = e^{-x^2/2} H_k(x)$ for $k = 1, 3, \dots, 2n + 1$, as test functions in the Rayleigh quotient, then for any $0 \leq q \in L^1(\mathbb{R})$ we obtain after a certain amount of rearranging

$$(3.16) \quad \begin{aligned} & \sum_{k=0}^n \left[\lambda_k - \lambda_k^0 - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx \right] \\ & \leq C_n + \sum_{k=0}^n \int_{\mathbb{R}} e^{-x^2} q(x) \frac{H_{2k+1}^2(x)}{2^{k+1}(2k+1)! \sqrt{\pi}} dx - \sum_{k=0}^n \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx, \end{aligned}$$

where the constant

$$C_n := \sum_{k=0}^n \lambda_{2k+1}^0 - \sum_{k=0}^n \lambda_k^0 \geq 0$$

depends only on $n \geq 0$. We will show that we can find q for which the first sum on the right-hand side of (3.16) is arbitrarily small, while the second sum is arbitrarily large. The idea is to choose q to have support in a very small neighbourhood of 0 and use that all odd Hermite polynomials H_{2k+1} satisfy $H_{2k+1}(0) = 0$ (and hence are very small close to 0). We start by fixing $K = K(n, N) > 0$ large enough that

$$(3.17) \quad C_n + 1 - K \sum_{k=0}^n \frac{1}{\pi \sqrt{\lambda_k^0}} < -N$$

and for given $\delta > 0$, to be specified later, we choose $q_{\delta}(x) := K\delta^{-1}\chi_{\delta}(x)$, where χ_{δ} is the indicator function of the set $[-\delta/2, \delta/2]$. Then obviously $q_{\delta} \geq 0$ has L^1 -norm equal to K for any $\delta > 0$. Since, as mentioned, $H_{2k+1}^2(0) = 0$ for all $k = 0, \dots, n$, and H_{2k+1}^2 is obviously continuous, for any $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon, n) > 0$ such that

$$0 \leq \frac{e^{-x^2} H_{2k+1}^2(x)}{2^{k+1}(2k+1)! \sqrt{\pi}} < \varepsilon$$

for all $x \in [-\delta/2, \delta/2]$ and all $k = 0, \dots, n$. It follows that for this δ , we have

$$\sum_{k=0}^n \int_{\mathbb{R}} e^{-x^2} q_{\delta}(x) \frac{H_{2k+1}^2(x)}{2^{k+1}(2k+1)! \sqrt{\pi}} dx < \varepsilon(n+1) < 1,$$

if we choose $\varepsilon < 1/(n+1)$. Inserting this estimate together with (3.17) into (3.16) yields the proposition. \square

4. Bounds for the perturbed harmonic oscillator with an integrable perturbation

Here we generalize Theorem 3.1 to allow for a class of perturbations q which may now take on negative values. Although the resulting estimate is not quite as tight as in Theorem 3.1, we still have convergence to the trace formula (1.4) at the same rate as before.

Theorem 4.1. *Given the function $q \in L^1(\mathbb{R})$, suppose that there exists a non-negative constant q_m for which $q(x) + q_m e^{-x^2}$ is non-negative for almost all real values of x . Then the eigenvalues of the corresponding perturbed harmonic oscillator (2.1) satisfy the inequalities*

$$(4.1) \quad \sum_{k=0}^n \left[\lambda_k - \lambda_k^0 - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx \right] \leq \frac{\chi_n}{\pi} \int_{\mathbb{R}} q(x) dx + \varepsilon_n \frac{q_m}{\sqrt{\pi}}$$

for $n = 0, 1, \dots$, where

$$(4.2) \quad \varepsilon_n = \omega_n - \sqrt{2} \frac{\Gamma(n + 3/2)}{\Gamma(n + 1)} \geq 0$$

and χ_n and ω_n are given by (3.2) and (3.4), respectively. Moreover, $\varepsilon_n = O(1/\sqrt{n})$ as $n \rightarrow \infty$.

Proof. We suppose $q_m \geq 0$ is as in the statement of the theorem, and mimic the proof of Theorem 3.1 to obtain

$$\begin{aligned} \sum_{k=0}^n \lambda_k &\leq \sum_{k=0}^n \lambda_k^0 + \int_{\mathbb{R}} e^{-x^2} q(x) \sum_{k=0}^n \frac{1}{2^k k! \sqrt{\pi}} H_k^2(x) dx \\ &= \sum_{k=0}^n \lambda_k^0 + \int_{\mathbb{R}} e^{-x^2} [q(x) + q_m e^{-x^2}] \sum_{k=0}^n \frac{1}{2^k k! \sqrt{\pi}} H_k^2(x) dx \\ &\quad - \frac{q_m}{\sqrt{\pi}} \sum_{k=0}^n \frac{1}{2^k k!} \int_{\mathbb{R}} e^{-2x^2} H_k^2(x) dx. \end{aligned}$$

Since $q(x) + q_m e^{-x^2} \in L^1(\mathbb{R})$ is positive by assumption, we may proceed as in the proof of Theorem 3.1 to obtain

$$\begin{aligned} \int_{\mathbb{R}} e^{-x^2} [q(x) + q_m e^{-x^2}] \sum_{k=0}^n \frac{1}{2^k k! \sqrt{\pi}} H_k^2(x) dx &\leq \frac{\omega_n}{\pi} \int_{\mathbb{R}} q(x) + q_m e^{-x^2} dx \\ &= \frac{\omega_n}{\pi} \int_{\mathbb{R}} q(x) dx + \frac{q_m}{\sqrt{\pi}} \omega_n. \end{aligned}$$

Meanwhile, since

$$\int_{\mathbb{R}} e^{-2x^2} H_k^2(x) dx = 2^{k-1/2} \Gamma\left(k + \frac{1}{2}\right),$$

we have

$$\sum_{k=0}^n \frac{1}{2^k k!} \int_{\mathbb{R}} e^{-2x^2} H_k^2(x) dx = \frac{1}{\sqrt{2}} \sum_{k=0}^n \frac{\Gamma(k + 1/2)}{k!} = \sqrt{2} \frac{\Gamma(n + 3/2)}{\Gamma(n + 1)}.$$

Combining the above expressions yields (4.1). The asymptotic behaviour of ε_n is an immediate consequence of (3.13) together with the expansion (3.14).

Although ε_n can be computed explicitly, to see that it is positive we use the following easier, indirect argument: if for a given $q \in L^1(\mathbb{R})$, (4.1) holds for some $q_m \geq 0$, then the above proof shows that it also holds for all $c \geq q_m$. This is only possible if $\varepsilon_n \geq 0$ for all $n \geq 0$. \square

5. A bound for a general potential in terms of Hermite polynomials

Here we will consider the general problem (1.1), supposing only that the potential $V: \mathbb{R} \rightarrow \mathbb{R}$ admits a series expansion in terms of Hermite polynomials in the manner of an eigenfunction decomposition

$$V(x) = \sum_{j=0}^{\infty} v_j H_j(x),$$

where we now assume that $V(x) \in L^2(\mathbb{R}, e^{-x^2} dx)$, or equivalently, since the H_j form an orthonormal basis of $L^2(\mathbb{R})$ with respect to this measure, that the sequence v_j is square summable. We will prove the following explicit estimate for the $\lambda_k = \lambda_k(V)$ based on the Fourier-type coefficients v_j .

Theorem 5.1. *Under the above conditions on the potential V , for every $n \in \mathbb{N}$, the n th eigenvalue of (1.1) with $(a, b) = \mathbb{R}$ satisfies*

$$(5.1) \quad \sum_{k=0}^n \lambda_k \leq \sum_{k=0}^n \frac{2^k (2k)!}{k!} \binom{n+1}{k+1} v_{2k} + \frac{1}{2} (n+1)^2.$$

REMARK 5.2. (i) This theorem will be proved by using the eigenfunctions of the quantum harmonic oscillator as test functions in the Rayleigh quotient, as was done in Theorem 3.1. The difference is that there we used an estimate for the sum of Hermite polynomials resulting from the test functions (Lemma 3.3), whereas here we expand out the potential as a Fourier series in Hermite polynomials and multiply this against our test functions, in the spirit of the arguments used in [5]. Since the only

inequality we use here is that which results from inserting the test functions into the Rayleigh quotient, and there is no other estimate involved, it follows that the right-hand side of (5.1) must necessarily be smaller than the right-hand side of (3.8) if V is of the form $V(x) = x^2 + q(x)$ for some $0 \leq q \in L^1(\mathbb{R})$ (indeed, it must be equal to the right-hand side of (3.5), i.e. the middle expression in (3.8)). However, in practice the two estimates are fundamentally different in nature; for example, it is not easy to see any relation between the right-hand side of (5.1) and the trace formula (1.4). See also Corollary 5.3 below.

(ii) As a trivial example to show that the above theorem is sharp, if $V(x) = x^2$, then the only two nonzero coefficients in the Fourier expansion of V are $v_2 = 1/4$, $v_0 = 1/2$, and it can easily be seen that (5.1) reduces to an equality.

Proof of Theorem 5.1. As mentioned, we will use the functions $\psi_k(x) := e^{-x^2/2} H_k(x)$ as test functions in the Rayleigh quotient. In order to do so, we shall need some more fairly standard facts about integrals of Hermite polynomials H_k , which may be found in [10], for instance: for $n, m \in \mathbb{N}$,

$$(5.2) \quad \int_{\mathbb{R}} e^{-x^2} H_n(x) H_m(x) dx = \delta_{mn} \sqrt{\pi} 2^n n!$$

where δ_{jk} is the Kronecker delta; and, for $\alpha, \beta, \gamma, s \in \mathbb{N}$ with $\alpha + \beta + \gamma = 2s$ even and $s \geq \alpha, \beta, \gamma$, we have

$$(5.3) \quad \int_{\mathbb{R}} e^{-x^2} H_\alpha(x) H_\beta(x) H_\gamma(x) dx = \sqrt{\pi} \frac{2^s \alpha! \beta! \gamma!}{(s - \alpha)! (s - \beta)! (s - \gamma)!};$$

under any other conditions on α, β, γ and s , this integral is 0. We also note that, combining a standard integration by parts, (5.2) and the formula $H'_n(x) = 2nH_{n-1}(x)$, we obtain easily that

$$(5.4) \quad \begin{aligned} \int_{\mathbb{R}} e^{-x^2} x^2 H_k^2(x) dx &= \frac{1}{2} \int_{\mathbb{R}} e^{-x^2} H_k^2(x) dx + 2k^2 \int_{\mathbb{R}} e^{-x^2} H_{k-1}^2(x) dx \\ &= \sqrt{\pi} 2^{k-1} k! + \sqrt{\pi} 2^k k k!. \end{aligned}$$

So, using the ψ_k as test functions, as well the convergence of the v_j to interchange integration and summation (noting that the functions $V(x), H_k^2(x) \in L^2(\mathbb{R}, e^{-x^2} dx)$, the latter being in $\text{span}\{H_0(x), H_2(x), \dots, H_{2k}(x)\}$) together with (5.2),

$$\begin{aligned} \sum_{k=0}^n \lambda_k &\leq \sum_{k=0}^n \frac{\int_{\mathbb{R}} [(d/dx)[e^{-x^2/2} H_k(x)]]^2 + e^{-x^2} V(x) H_k^2(x) dx}{\int_{\mathbb{R}} e^{-x^2} H_k^2(x) dx} \\ &= \sum_{k=0}^n \left(\lambda_k^0 - \frac{\int_{\mathbb{R}} e^{-x^2} x^2 H_k^2(x) dx}{\int_{\mathbb{R}} e^{-x^2} H_k^2(x) dx} \right) + \sum_{k=0}^n \sum_{j=0}^{\infty} \frac{v_j \int_{\mathbb{R}} e^{-x^2} (x) H_j(x) H_k^2(x) dx}{2^k k! \sqrt{\pi}}. \end{aligned}$$

Using (5.2) and (5.4),

$$\frac{\int_{\mathbb{R}} e^{-x^2} x^2 H_k^2(x) dx}{\int_{\mathbb{R}} e^{-x^2} H_k^2(x) dx} = k + \frac{1}{2},$$

while (5.3) with $\alpha = \beta = k$ and $\gamma = j$ implies $\int_{\mathbb{R}} e^{-x^2}(x) H_j(x) H_k^2(x) dx \neq 0$ if and only if j is even and $j \leq 2k$, and under these conditions, writing $j =: 2m$ for $m = 0, \dots, k$,

$$\int_{\mathbb{R}} e^{-x^2}(x) H_{2m}(x) H_k^2(x) dx = \sqrt{\pi} \frac{2^{k+m} (k!)^2 (2m)!}{(m!)^2 (k-m)!} = \sqrt{\pi} \frac{2^{k+m} k! (2m)!}{m!} \binom{k}{m}.$$

Combining the above yields

$$\sum_{k=0}^n \lambda_k \leq \sum_{k=0}^n \left(\lambda_k^0 - k - \frac{1}{2} \right) + \sum_{k=0}^n \sum_{m=0}^k \frac{2^m (2m)!}{m!} \binom{k}{m} v_{2m}.$$

To simplify this last sum, since $\binom{a}{b} = 0$ for $b > a$, we may just as well sum m from 0 to n , giving the sum as

$$\sum_{m=0}^n \frac{2^m (2m)!}{m!} v_{2m} \left(\sum_{k=0}^n \binom{k}{m} \right) = \sum_{m=0}^n \frac{2^m (2m)!}{m!} \binom{n+1}{m+1} v_{2m},$$

using a standard formula for binomial coefficients. This establishes the theorem. \square

We shall now assume explicitly that the potential V is a perturbation of the harmonic potential and thus return to writing it as $V(x) = x^2 + q(x)$, where we will assume that q is integrable. By adding the terms which are missing in the right-hand side of (5.1) in order to obtain a sequence which converges to the right-hand side of the trace formula (1.4), and expressing the coefficients in the left-hand side in terms of the Fourier coefficients of the function q , we obtain the following result.

Corollary 5.3.

$$\begin{aligned} (5.5) \quad & \sum_{k=0}^n \left[\lambda_k - \lambda_k^0 - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx \right] \\ & \leq \sum_{k=0}^n \left[\frac{2^k (2k)!}{k!} \binom{n+1}{k+1} q_{2k} - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx \right]. \end{aligned}$$

Proof. From $V(x) = q(x) + x^2$ we obtain the relations

$$v_j = \begin{cases} q_0 + \frac{1}{2}, & j = 0, \\ q_2 + \frac{1}{4}, & j = 2, \\ q_k, & j \neq 0, 2. \end{cases}$$

Replacing this in (5.1) and adding and subtracting the term

$$-\lambda_k^0 - \frac{1}{\pi \sqrt{\lambda_k^0}} \int_{\mathbb{R}} q(x) dx$$

inside the summation on the left-hand side of (5.1), we obtain, after some manipulations, the desired result. \square

REMARK 5.4. Clearly the integral term appearing inside both sums can be cancelled. However, in this way not only do we obtain an expression where the left-hand side converges in the limit as n goes to infinity (under additional assumptions on q as in [2, 7]), but since as noted in Remark 5.1 (i) the right-hand side of (5.5) is necessarily smaller than the right-hand side of (3.1) (or (4.1), depending on q), it follows that it must converge to the right-hand side of the trace formula (1.4) and at least as fast as $O(1/\sqrt{n})$.

6. Power generalizations of Theorems 3.1 and 4.1

In this section we generalize the summation bounds obtained in Theorems 3.1 and 4.1 to allow for the summands (arranged in various ways) to be raised to a given negative power. We keep the notation and assumptions of Sections 3 and 4, and start with the case where the perturbation q is non-negative.

Theorem 6.1. *Under the assumptions and notation of Theorem 3.1, with ω_n as in (3.4), for all $n \geq 0$ and $s > 0$,*

$$(6.1) \quad \left(\frac{1}{n+1} \right) \sum_{k=0}^n (\lambda_k - \lambda_k^0)^{-s} \geq \left[\frac{\omega_n}{(n+1)\pi} \int_{\mathbb{R}} q(x) dx \right]^{-s}.$$

Under certain additional assumptions on the potential, we can rearrange the order of the terms in the above bounds somewhat.

Proposition 6.2. *If $\int_{\mathbb{R}} q(x) dx < 32\sqrt{\pi}$, then for all $n \geq 0$ and $s > 0$,*

$$(6.2) \quad \sum_{k=0}^n \lambda_k^{-s} \geq \sum_{k=0}^n \left[\lambda_k^0 + \frac{\omega_k - \omega_{k-1}}{\pi} \int_{\mathbb{R}} q(x) dx \right]^{-s}.$$

We next consider the situation covered by Theorem 4.1, where the perturbation q may take on negative values, provided its negative part decays rapidly enough at infinity. For simplicity, we consider the special case where q has zero mean.

Theorem 6.3. *Suppose in addition to the assumptions of Theorem 4.1 that $\int_{\mathbb{R}} q(x) dx = 0$. Then for all $n \geq 0$ and $s > 0$,*

$$(6.3) \quad \sum_{k=0}^n \lambda_k^{-s} \geq (s+1) \sum_{k=0}^n (\lambda_k^0)^{-s} - sq_m \sum_{k=0}^n (\lambda_k^0)^{-s-1} (\varepsilon_k - \varepsilon_{k-1}),$$

where $q_m \geq 0$ is defined in Theorem 4.1. Here $\varepsilon_n \geq 0$ is given by (4.2) for $n \geq 0$ and we set $\varepsilon_{-1} := 0$.

These results will be proved by combining generic results on arbitrary increasing or decreasing sequences of real numbers (see Lemma 6.5 and what follows it) with the following particular properties of the ω_n .

Lemma 6.4. *The sequence $\{\omega_n\}_{n \in \mathbb{N}}$ is positive and strictly increasing, while $\{\tau_n\}_{n \in \mathbb{N}}$ given by $\tau_n := \omega_{n+1} - \omega_n$ is positive and non-increasing.*

Proof. The ω_n are obviously all positive. Using the formulae

$$\Gamma\left(\frac{z+1}{2}\right) = \frac{z! \sqrt{\pi}}{2^z (z/2)!}, \quad \Gamma\left(\frac{z}{2} + 1\right) = \left(\frac{z}{2}\right)!,$$

for $z \in \mathbb{N}$ even, if we assume $n \geq 0$ is even and set

$$C_n := (n+1) \frac{\Gamma((n+1)/2)}{\Gamma(n/2 + 1)} = \frac{(n+1)! \sqrt{\pi}}{2^n [(n/2)!]^2} > 0,$$

then an elementary calculation shows that

$$\begin{aligned} \omega_{n+1} - \omega_n &= \frac{C_n}{2(n+2)}, \\ \omega_{n+2} - \omega_{n+1} &= \frac{C_n}{2(n+2)}, \\ \omega_{n+3} - \omega_{n+2} &= \frac{(n+3)C_n}{2(n+2)(n+4)}, \end{aligned}$$

from which we see that ω_n is increasing in n , while $\tau_n = \omega_{n+1} - \omega_n$ is positive and weakly decreasing. \square

The following lemma appeared in [5], but for the sake of completeness we state and prove it here as well. Here and throughout, we will use the notation $[y]_+$, $y \in \mathbb{R}$, to denote the expression taking on the value y if $y \geq 0$ and zero otherwise; $[f(x)]_{g(x) \geq y}$ will represent $f(x)$ if $g(x) \geq y$ and zero otherwise.

Lemma 6.5. *Suppose the sequences $(a_k)_{k \in \mathbb{N}}$ and $(b_k)_{k \in \mathbb{N}}$ are positive, with $(b_k)_{k \in \mathbb{N}}$ non-decreasing in $k \geq 0$. Suppose also that the sequence $(c_k)_{k \in \mathbb{N}}$ satisfies*

$$(6.4) \quad \sum_{k=0}^m a_k \leq \sum_{k=0}^m c_k$$

for all $m \geq 0$. Then for all $s > 0$ and all $n \geq 0$ we have

$$(6.5) \quad \sum_{k=0}^n (a_k)^{-s} \geq \sum_{k=0}^n [(s+1)(b_k)^{-s} - s(b_k)^{-s-1} c_k].$$

If the sequence $(c_k)_{k \in \mathbb{N}}$ is itself positive and non-decreasing in $k \geq 0$, then the right-hand side of (6.5) is maximized when $b_k = c_k$ for all $0 \leq k \leq n$, in which case (6.5) simplifies to

$$\sum_{k=0}^n (a_k)^{-s} \geq \sum_{k=0}^n (c_k)^{-s}.$$

An examination of the proof shows that if we want (6.5) to hold for some fixed $n \geq 0$, then for the proof to work we need (6.4) to hold for all $0 \leq m \leq n$.

Proof of Lemma 6.5. For $\lambda > 0$, we use the identity, valid for all $s > 0$,

$$(6.6) \quad \lambda^{-s} = s(s+1) \int_0^\infty \alpha^{-s-2} [\alpha - \lambda]_+ d\alpha.$$

Hence for $n \geq 0$, $s > 0$ arbitrary,

$$\begin{aligned} \sum_{k=0}^n (a_k^{-s} - b_k^{-s}) &= s(s+1) \int_0^\infty \alpha^{-s-2} \sum_{k=0}^n ([\alpha - a_k]_+ - [\alpha - b_k]_+) d\alpha \\ &\geq s(s+1) \int_0^\infty \alpha^{-s-2} \sum_{k=0}^n [b_k - a_k]_{\alpha \geq b_k} d\alpha \\ &\geq s(s+1) \int_0^\infty \alpha^{-s-2} \sum_{k=0}^n [b_k - c_k]_{\alpha \geq b_k} d\alpha \\ &= \sum_{k=0}^n s(s+1)(b_k - c_k) \int_{a_k}^\infty \alpha^{-s-2} d\alpha, \end{aligned}$$

which after simplification and rearrangement gives us (6.5). For the maximizing property we consider each term on the right-hand side of (6.5) as a function of b_k

$$g_k(b_k) := (s+1)(b_k)^{-s} - s(b_k)^{-s-1}c_k.$$

Differentiating in b_k shows that g_k reaches its unique maximum when $b_k = c_k$. \square

Proof of Proposition 6.2. Lemma 6.5 may be applied directly to prove Proposition 6.2 in the obvious way; for (6.2), it merely remains to be confirmed that the sequence

$$\left\{ \lambda_k^0 + \frac{\omega_k - \omega_{k-1}}{\pi} \int_{\mathbb{R}} q(x) dx \right\}_{k \in \mathbb{N}}$$

is positive and non-decreasing. Now since $\lambda_{k+1}^0 - \lambda_k^0 = 2$ for all $k \geq 0$, we need $\int_{\mathbb{R}} q(x) dx$ (which we assume to be nonzero and hence strictly positive) to be small enough that

$$\omega_{k+2} - 2\omega_{k+1} + \omega_k \geq -\frac{2\pi}{\int_{\mathbb{R}} q(x) dx}$$

for all $k \geq 0$. If k is even, then the left-hand side is identically zero, as follows from the proof of Lemma 6.4. Otherwise, for $k+1$ odd, we have

$$\omega_{k+3} - 2\omega_{k+2} + \omega_{k+1} = \frac{(k+3)C_k}{2(k+2)(k+4)} - \frac{C_k}{2(k+2)},$$

which, using the definition of C_k , may be rearranged to give

$$-\frac{\sqrt{\pi}}{2(k+4)} \cdot \frac{k+1}{k+2} \cdot \frac{k-1}{k-2} \cdots \frac{3}{4} \cdot \frac{1}{2},$$

which we see is negative and increasing in $k+1 \geq 1$ odd. Thus $\omega_{k+3} - 2\omega_{k+2} - \omega_{k+1}$ reaches its largest negative value, namely $-C_0/16 = -\sqrt{\pi}/16$, when $k=0$. The requirement on $q(x)$ is therefore that

$$-\frac{\sqrt{\pi}}{16} \geq -\frac{2\pi}{\int_{\mathbb{R}} q(x) dx},$$

that is, we have shown the required sequence is increasing when $\int_{\mathbb{R}} q(x) dx \leq 32\sqrt{\pi}$. \square

Proof of Theorem 6.1. To prove (6.1) we use a similar idea to the one in Lemma 6.5, but since the right-hand side of (3.1) is not a sequence, the method needs

to be adapted slightly to this situation. Namely, starting with the representation (6.6) of $\lambda =: \lambda_k - \lambda_k^0$,

$$(6.7) \quad \sum_{k=0}^n (\lambda_k - \lambda_k^0)^{-s} \geq s(s+1) \int_0^\infty \alpha^{-s-2} \sum_{k=0}^n [\alpha - \lambda_k + \lambda_k^0]_{\alpha \geq M} d\alpha$$

for all $M \in \mathbb{R}$; we make the choice $M := (\omega_n/((n+1)\pi)) \int_{\mathbb{R}} q(x) dx$. Using (3.1), which, when rearranged, says that

$$(6.8) \quad \sum_{k=0}^n (\lambda_k - \lambda_k^0) \leq \frac{\omega_n}{\pi} \int_{\mathbb{R}} q(x) dx,$$

we have

$$\sum_{k=0}^n [\alpha - \lambda_k + \lambda_k^0]_{\alpha \geq \omega_n/((n+1)\pi)} \int_{\mathbb{R}} q(x) dx \geq (n+1) \left[\alpha - \frac{\omega_n}{(n+1)\pi} \int_{\mathbb{R}} q(x) dx \right]_+.$$

Substituting this into (6.7) and applying (6.6) yields (6.1). \square

Proof of Theorem 6.3. This follows directly from Theorem 4.1 and Lemma 6.5, where we take $a_k = \lambda_k$, $b_k = \lambda_k^0$ and $c_k = \lambda_k^0 + (\varepsilon_k - \varepsilon_{k-1})q_m$ (with $\varepsilon_{-1} := 0$). \square

References

- [1] M. Abramowitz and I.A. Stegun: *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards Applied Mathematics Series **55**, For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC, 1964.
- [2] E.V. Aleksandrova, O.V. Bochkareva and V.E. Podol'skiĭ: *Summation of regularized traces of the singular Sturm-Liouville operator*, *Differential Equations* **33** (1997), 287–291.
- [3] C. Bande: *Isoperimetric Inequalities and Their Applications*, Monographs and Studies in Mathematics **7**, Pitman, Boston, Mass., 1980.
- [4] L.A. Dikiĭ: *On a formula of Gelfand–Levitan*, *Uspehi Matem. Nauk (N.S.)* **8** (1953), 119–123.
- [5] P. Freitas and J.B. Kennedy: *Summation formula inequalities for eigenvalues of Schrödinger operators*, preprint.
- [6] B.M. Levitan and I.S. Sargsjan: *Sturm–Liouville and Dirac Operators*, Mathematics and its Applications (Soviet Series) **59**, Kluwer Acad. Publ., Dordrecht, 1991.
- [7] A. Pushnitski and I. Sorrell: *High energy asymptotics and trace formulas for the perturbed harmonic oscillator*, *Ann. Henri Poincaré* **7** (2006), 381–396.
- [8] V.A. Sadovnichii and V.E. Podol'skiĭ: *Traces of operators*, *Uspekhi Mat. Nauk* **61** (2006), 89–156, (Russian), translation in *Russian Math. Surveys* **61** (2006), 885–953.
- [9] G. Szegő: *On an inequality of P. Turán concerning Legendre polynomials*, *Bull. Amer. Math. Soc.* **54** (1948), 401–405.
- [10] G. Szegő: *Orthogonal Polynomials*, fourth edition, Amer. Math. Soc., Providence, RI, 1975.

[11] E.C. Titchmarsh: The Theory of the Riemann Zeta-Function, second edition, Oxford Univ. Press, New York, 1986.

Pedro Freitas
Department of Mathematics
Faculty of Human Kinetics
Universidade de Lisboa
and
Group of Mathematical Physics
Faculty of Sciences
Universidade de Lisboa
Campo Grande, Edifício C6
1749-016 Lisboa
Portugal
e-mail: psfreitas@fc.ul.pt

James B. Kennedy
Institute of Applied Analysis
University of Ulm
Helmholtzstr. 18, D-89069 Ulm
Germany
and
Institute of Analysis, Dynamics and Modelling
University of Stuttgart
Pfaffenwaldring 57, D-70569 Stuttgart
Germany
e-mail: james.kennedy@mathematik.uni-stuttgart.de