

Title	p-LOCAL STABLE SPLITTING OF QUASITORIC MANIFOLDS
Author(s)	Hasui, Sho; Kishimoto, Daisuke; Sato, Takashi
Citation	Osaka Journal of Mathematics. 2016, 53(3), p. 843-854
Version Type	VoR
URL	https://doi.org/10.18910/58883
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

p -LOCAL STABLE SPLITTING OF QUASITORIC MANIFOLDS

SHO HASUI, DAISUKE KISHIMOTO and TAKASHI SATO

(Received February 12, 2015, revised August 28, 2015)

Abstract

We show a homotopy decomposition of the p -localized suspension $\Sigma M_{(p)}$ of a quasitoric manifold M by constructing power maps. As an application we investigate the p -localized suspension of the projection π from the moment-angle complex onto M , from which we deduce its triviality for $p > \dim M/2$. We also discuss non-triviality of $\pi_{(p)}$ and $\Sigma^\infty \pi$.

1. Introduction and statement of results

Manifolds which are now known as quasitoric manifolds were introduced by Davis and Januszkiewicz [4] as a topological counterpart of smooth projective toric varieties, and have been the subject of recent interest in the study of manifolds with torus action. As well as toric varieties, quasitoric manifolds have been studied in a variety of contexts where combinatorics, geometry, and topology interact in a fruitful way. We refer the reader to the exposition [2] written by Buchstaber and Panov for basics of quasitoric manifolds. This note studies a topological aspect of quasitoric manifolds involving their p -localized suspension. A quasitoric manifold M over a simple n -polytope P is by definition a $2n$ -manifold with a locally standard action of the compact n -torus T^n such that the orbit space M/T^n is identified with the simple polytope P as manifolds with corners. A fundamental example of quasitoric manifolds is the complex projective space $\mathbb{C}P^n$ which is the only quasitoric manifold over the n -simplex, whereas there are several quasitoric manifolds on the same simple polytope in general. Observe that since $\mathbb{C}P^n$ admits power maps, the p -localization of the suspension $\Sigma \mathbb{C}P^n_{(p)}$ splits into a wedge of $p - 1$ spaces as in [6]. We prove that any quasitoric manifold also admits power maps, and as a consequence the p -localization of its suspension splits into a wedge of $p - 1$ spaces.

Theorem 1.1. *For a quasitoric manifold M there is a homotopy equivalence*

$$\Sigma M_{(p)} \simeq X_1 \vee \cdots \vee X_{p-1}$$

such that for each i , $\tilde{H}_(X_i; \mathbb{Z}) = 0$ unless $* \equiv 2i + 1 \pmod{2(p-1)}$.*

As a corollary we get a kind of rigidity of quasitoric manifolds over the same polytope, which also follows from a more general result Proposition 3.3.

Corollary 1.2. *Let M, N be quasitoric manifolds over the same simple n -polytope. For $p > n$ there is a homotopy equivalence*

$$\Sigma M_{(p)} \simeq \Sigma N_{(p)}.$$

To a simplicial complex K we can assign a space \mathcal{Z}_K which is called the moment-angle complex for K (see [4, 2]). The fundamental construction involving quasitoric manifolds is that every quasitoric manifold over a simple polytope P is obtained by the quotient of a certain free torus action on the moment-angle complex $\mathcal{Z}_{K(P)}$, where $K(P)$ denotes the boundary of the dual simplicial polytope of P . Then for a quasitoric manifold M over P the projection $\pi: \mathcal{Z}_{K(P)} \rightarrow M$ is of particular importance. We investigate the p -localization of the suspension of this projection through the p -local stable splitting of Theorem 1.1. Let K be a simplicial complex on the vertex set V . Recall from [1] that there is a homotopy equivalence

$$(1.1) \quad \Sigma \mathcal{Z}_K \simeq \bigvee_{\emptyset \neq I \subset V} \Sigma^{|I|+2} |K_I|$$

where K_I denotes the full subcomplex of K on the vertex set $I \subset V$, i.e. $K_I = \{\sigma \in K \mid \sigma \subset I\}$, and $|K_I|$ means the geometric realization of K_I . We identify the map $\Sigma \pi_{(p)}: \Sigma(\mathcal{Z}_{K(P)})_{(p)} \rightarrow \Sigma M_{(p)}$ through the homotopy equivalences of Theorem 1.1 and (1.1). Note that if P has m facets, then the vertex set of $K(P)$ is $[m] := \{1, \dots, m\}$.

Theorem 1.3. *Let M be a quasitoric manifold over a simple polytope P with m facets. Then through the homotopy equivalences of Theorem 1.1 and (1.1), the map $\Sigma \pi_{(p)}: \Sigma(\mathcal{Z}_{K(P)})_{(p)} \rightarrow \Sigma M_{(p)}$ is identified with a wedge of maps*

$$\bigvee_{\substack{\emptyset \neq I \subset [m] \\ |I| \equiv i \pmod{p-1}}} (\Sigma^{|I|+2} |K(P)_I|)_{(p)} \rightarrow X_i.$$

for $i = 1, \dots, p - 1$.

Corollary 1.4. *Let M be a quasitoric manifold over a simple n -polytope P . For $p > n$, the map $\Sigma \pi_{(p)}: \Sigma(\mathcal{Z}_{K(P)})_{(p)} \rightarrow \Sigma M_{(p)}$ is null homotopic.*

We also discuss necessity of suspension and localization for triviality of the projection $\pi: \mathcal{Z}_{K(P)} \rightarrow M$ in Corollary 1.4. Consider the complex projective space $\mathbb{C}P^1$ as a quasitoric manifold. Then the projection π is the Hopf map $S^3 \rightarrow \mathbb{C}P^1$, so neither

$\Sigma^\infty \pi$ nor $\pi_{(p)}$ for any p is null homotopic. We will discuss this problem for more general quasitoric manifolds.

The authors are grateful to Kouyemon Iriye and Shuichi Tsukuda for useful comments.

2. Cohomology of quasitoric manifolds

This section collects basic properties of the cohomology of quasitoric manifolds which will be used later. Let P be a simple n -polytope, and let M be a quasitoric manifold over P . Put $f_i(P)$ to be the number of $(n - i - 1)$ -dimensional faces of P for $i = -1, 0, \dots, n - 1$. The h -vector of P is defined by $(h_0(P), \dots, h_n(P))$ such that for $k = 0, \dots, n$,

$$h_k(P) = \sum_{i=0}^k (-1)^{k-i} \binom{n-i}{n-k} f_{i-1}(P).$$

It is known that the module structure of the cohomology of M is described by the h -vector of P , implying that the module structure depends only on P .

Proposition 2.1 (Davis and Januszkiewicz [4, Theorem 3.1] (cf. [2])). *Let M be a quasitoric manifold over P . Then we have*

$$H^{\text{odd}}(M; \mathbb{Z}) = 0$$

and

$$H^{2i}(M; \mathbb{Z}) \cong \mathbb{Z}^{h_i(P)}.$$

Let K be a simplicial complex on the vertex set $[m]$. The moment-angle complex \mathcal{Z}_K is defined by

$$\mathcal{Z}_K := \bigcup_{\sigma \in K} D(\sigma) \quad (\subset (D^2)^m)$$

where $D(\sigma) = \{(x_1, \dots, x_m) \in (D^2)^m \mid |x_i| = 1 \text{ whenever } i \notin \sigma\}$ and D^2 is regarded as the unit disk of \mathbb{C} . Then the canonical action of T^m on $(D^2)^m$ restricts to the action of T^m on \mathcal{Z}_K . Let M be a quasitoric manifold over a simple n -polytope P with m facets. Then we may regard the vertex set of $K(P)$ is $[m]$. As in [4, 2], M is obtained by quotienting out the moment-angle complex $\mathcal{Z}_{K(P)}$ by a certain free T^{m-n} -action which is the restriction of the canonical T^m -action. Then there is a homotopy fibration

$$(2.1) \quad \mathcal{Z}_{K(P)} \xrightarrow{\pi} M \xrightarrow{\alpha} BT^{m-n}.$$

One easily sees that $\mathcal{Z}_{K(P)}$ is 2-connected (cf. [2]), hence the transgression $H^1(T^{m-n}; \mathbb{Z}) \rightarrow H^2(M; \mathbb{Z})$ associated with the fibration $T^{m-n} \rightarrow \mathcal{Z}_{K(P)} \xrightarrow{\pi} M$ is an

isomorphism. In particular the induced map $\alpha^*: H^2(BT^{m-n}; \mathbb{Z}) \rightarrow H^2(M; \mathbb{Z})$ is an isomorphism. It is also known as in [4, Theorem 4.14] (cf. [2]) that the cohomology ring $H^*(M; \mathbb{Z})$ is generated by 2-dimensional elements. We record these properties of the cohomology of M .

Proposition 2.2. *Let M be a quasitoric manifold over a simple n -polytope P with m facets.*

- (1) *The transgression $H^1(T^{m-n}; \mathbb{Z}) \rightarrow H^2(M; \mathbb{Z})$ associated with the fibration $T^{m-n} \rightarrow \mathcal{Z}_{K(P)} \xrightarrow{\pi} M$ is an isomorphism.*
- (2) *The map $\alpha^*: H^2(BT^{m-n}; \mathbb{Z}) \rightarrow H^2(M; \mathbb{Z})$ is an isomorphism.*
- (3) *The cohomology ring $H^*(M; \mathbb{Z})$ is generated by $H^2(M; \mathbb{Z})$.*

3. Proofs of the main results

Let P be a simple n -polytope with m facets, and let M be a quasitoric manifold over P . We construct power maps of M . Let u be an integer. By the definition of moment-angle complexes, the degree u self-map of S^1 induces a self-map $\underline{u}: \mathcal{Z}_{K(P)} \rightarrow \mathcal{Z}_{K(P)}$.

Lemma 3.1. *There is a self-map $\underline{u}: M \rightarrow M$ satisfying*

$$\underline{u}^* = u^k : H^{2k}(M; \mathbb{Z}) \rightarrow H^{2k}(M; \mathbb{Z}),$$

where the u^k means the multiplication by u^k .

Proof. Since M is the quotient of the restriction of the canonical T^m -action to a certain subtorus, the map $\underline{u}: \mathcal{Z}_{K(P)} \rightarrow \mathcal{Z}_{K(P)}$ induces a map $\underline{u}: M \rightarrow M$ satisfying the commutative diagram

$$\begin{array}{ccccc} T^{m-n} & \longrightarrow & \mathcal{Z}_{K(P)} & \xrightarrow{\pi} & M \\ \downarrow \underline{u} & & \downarrow \underline{u} & & \downarrow \underline{u} \\ T^{m-n} & \longrightarrow & \mathcal{Z}_{K(P)} & \xrightarrow{\pi} & M \end{array}$$

where $\underline{u}: T^{m-n} \rightarrow T^{m-n}$ is the product of the degree u map of S^1 . Then by Proposition 2.2 and naturality of transgression, we see that the self-map $\underline{u}: M \rightarrow M$ has the desired property. □

We now recall the result of [6], where we reproduce the proof in order to clarify naturality. Let X be a CW-complex of finite type connected satisfying

- (1) $H_{\text{odd}}(X; \mathbb{Z}) = 0$ and $H_{\text{even}}(X; \mathbb{Z})$ is free, and
- (2) there is a self-map $\varphi: X \rightarrow X$ satisfying $\varphi_* = u^k: H_{2k}(X; \mathbb{Z}) \rightarrow H_{2k}(X; \mathbb{Z})$ for any $k \geq 0$, where u is an integer whose modulo p reduction is the primitive $(p - 1)$ th root of unity of \mathbb{Z}/p .

Define a self map $\alpha_i: \Sigma X \rightarrow \Sigma X$ by $\alpha_i := (\Sigma\varphi - u^1) \circ \cdots \circ \widehat{(\Sigma\varphi - u^i)} \circ \cdots \circ (\Sigma\varphi - u^{p-1})$ for $i = 1, \dots, p - 1$. Then $(\alpha_i)_*: \tilde{H}_{2k+1}(\Sigma X; \mathbb{Z}/p) \rightarrow \tilde{H}_{2k+1}(\Sigma X; \mathbb{Z}/p)$ is trivial for $k \not\equiv i \pmod{p-1}$ and is the isomorphism for $k \equiv i \pmod{p-1}$. Put

$$X_i = \text{hocolim}\{\Sigma X_{(p)} \xrightarrow{\alpha_i} \Sigma X_{(p)} \xrightarrow{\alpha_i} \Sigma X_{(p)} \xrightarrow{\alpha_i} \cdots\}.$$

Then it is easy to check that X_i is p -locally of finite type and

$$\tilde{H}_{2k+1}(X_i; \mathbb{Z}/p) = \begin{cases} \tilde{H}_{2k+1}(\Sigma X; \mathbb{Z}/p), & k \equiv i \pmod{p-1}, \\ 0, & k \not\equiv i \pmod{p-1} \end{cases}$$

such that the canonical map $\Sigma X_{(p)} \rightarrow X_i$ induces the projection in mod p homology. Then the composite $\Sigma X_{(p)} \rightarrow \Sigma X_{(p)} \vee \cdots \vee \Sigma X_{(p)} \rightarrow X_1 \vee \cdots \vee X_{p-1}$ is an isomorphism in mod p homology, hence an isomorphism in homology with coefficient $\mathbb{Z}_{(p)}$ since spaces on both sides are p -locally of finite type, where the first arrow in the composite is defined by using the suspension comultiplication. Therefore by the J.H.C. Whitehead theorem we obtain:

Lemma 3.2 (Mimura, Nishida and Toda [6]). *Let X and X_i be as above. There is a homotopy equivalence*

$$\Sigma X_{(p)} \simeq X_1 \vee \cdots \vee X_{p-1}$$

such that $\tilde{H}_*(X_i; \mathbb{Z}/p) = 0$ unless $* \equiv 2i + 1 \pmod{2(p-1)}$ for $i = 1, \dots, p - 1$.

We now prove the main results.

Proof of Theorem 1.1. Combine Lemmas 3.1 and 3.2. □

Proof of Corollary 1.2. Recall that M is of dimension $2n$. Apply Theorem 1.1 to M , then we get $\Sigma M_{(p)} \simeq X_1 \vee \cdots \vee X_{p-1}$. If $p > n$, the space X_i is torsion free in homology over $\mathbb{Z}_{(p)}$ and satisfies $\tilde{H}_*(X_i; \mathbb{Z}/p) = 0$ unless $* = 2i + 1$. Then since X_i is simply connected, X_i is a wedge of $S_{(p)}^{2i+1}$, where the number of spheres is the $2i$ -dimensional Betti number of M which is equal to $h_i(P)$ by Proposition 2.1. So we obtain a homotopy equivalence $\Sigma M_{(p)} \simeq \bigvee_{i=1}^{p-1} \bigvee^{h_i(P)} S_{(p)}^{2i+1}$. We can get the same homotopy equivalence for N as well, and therefore the proof is completed. □

Proof of Theorem 1.3. Define a map $\beta_i: \Sigma \mathcal{Z}_{K(P)} \rightarrow \Sigma \mathcal{Z}_{K(P)}$ by $\beta_i = (\Sigma \underline{u} - u^1) \circ \cdots \circ \widehat{(\Sigma \underline{u} - u^i)} \circ \cdots \circ (\Sigma \underline{u} - u^{p-1})$ for $i = 1, \dots, p - 1$, where u is an integer whose modulo p reduction is the primitive $(p - 1)^{\text{th}}$ root of unity of \mathbb{Z}/p . Put

$$Y_i = \text{hocolim}\{\Sigma(\mathcal{Z}_{K(P)})_{(p)} \xrightarrow{\beta_i} \Sigma(\mathcal{Z}_{K(P)})_{(p)} \xrightarrow{\beta_i} \Sigma(\mathcal{Z}_{K(P)})_{(p)} \xrightarrow{\beta_i} \cdots\}.$$

By naturality of the homotopy equivalence (1.1) with respect to self-maps of S^1 [1, Theorem 2.10], the self-map $u: \Sigma \mathcal{Z}_{K(P)} \rightarrow \Sigma \mathcal{Z}_{K(P)}$ is identified with a wedge of the degree $u^{|I|}$ maps

$$u^{|I|}: \Sigma^{|I|+2}|K(P)_I| \rightarrow \Sigma^{|I|+2}|K(P)_I|$$

for $\emptyset \neq I \subset [m]$. Then we have $Y_i = \bigvee_{\substack{\emptyset \neq I \subset [m] \\ |I| \equiv i \pmod{p-1}}} \Sigma^{|I|+2}|K(P)_I|$ and the canonical map $\Sigma(\mathcal{Z}_{K(P)})_{(p)} \rightarrow Y_i$ is the projection similarly to the proof of Proposition 3.2. So the composite $\Sigma(\mathcal{Z}_{K(P)})_{(p)} \rightarrow \Sigma(\mathcal{Z}_{K(P)})_{(p)} \vee \cdots \vee \Sigma(\mathcal{Z}_{K(P)})_{(p)} \rightarrow Y_1 \vee \cdots \vee Y_{p-1}$ is a homotopy equivalence, where the first map is defined by the suspension comultiplication and the second map is a wedge of the canonical maps into the homotopy colimits. On the other hand, by Lemma 3.1 there is a commutative diagram

$$\begin{array}{ccc} \Sigma \mathcal{Z}_{K(P)} & \xrightarrow{\beta_i} & \Sigma \mathcal{Z}_{K(P)} \\ \downarrow \Sigma \pi & & \downarrow \Sigma \pi \\ \Sigma M & \xrightarrow{\alpha_i} & \Sigma M \end{array}$$

where α_i is as above. Then there are maps $\pi_i: Y_i \rightarrow X_i$ satisfying a commutative diagram

$$\begin{array}{ccc} \Sigma(\mathcal{Z}_{K(P)})_{(p)} & \longrightarrow & Y_1 \vee \cdots \vee Y_{p-1} \\ \downarrow \Sigma \pi_{(p)} & & \downarrow \pi_1 \vee \cdots \vee \pi_{p-1} \\ \Sigma M_{(p)} & \longrightarrow & X_1 \vee \cdots \vee X_{p-1} \end{array}$$

where the horizontal arrows are the prescribed homotopy equivalences. Thus the proof is completed. □

Proof of Corollary 1.4. Since $p > n$, the map $\Sigma \pi_{(p)}: \Sigma(\mathcal{Z}_{K(P)})_{(p)} \rightarrow \Sigma M_{(p)}$ is identified with a wedge of the maps $\bigvee_{I \subset [m], |I|=i} (\Sigma^{|I|+2}|K(P)_I|)_{(p)} \rightarrow \bigvee S_{(p)}^{2i+1}$ for $i = 1, \dots, p-1$. If $\dim K(P)_I = |I| - 1$, then $K(P)$ is a simplex, so $|K(P)_I|$ is contractible. Then $\bigvee_{I \subset [m], |I|=i} \Sigma^{|I|+1}|K(P)_I|$ is homotopy equivalent to a CW-complex of dimension at most $2i$, completing the proof. □

We close this section by showing a general homotopy theoretical property of finite complexes consisting only of even cells from which Corollary 1.2 also follows since there are cell decompositions of quasitoric manifolds only by even dimensional cells.

Proposition 3.3. *Let X be a finite dimensional simply connected finite complex consisting only of 0 and odd cells. If $p > n$, then $X_{(p)}$ is homotopy equivalent to a wedge of p -localized odd spheres.*

Proof. Induct on the skeleta of X . We may assume the 1-skeleton is a point since X is simply connected, so the claim is trivially true for the 1-skeleton. Suppose that $X_{(p)}^{(2k-1)} \simeq \bigvee_{i=1}^{k-1} \bigvee^{m_i} S_{(p)}^{2i+1}$. Then the attaching maps of $(2k + 1)$ -cells of $X_{(p)}$ are identified with maps $S^{2k} \rightarrow \bigvee_{i=1}^{k-1} \bigvee^{m_i} S_{(p)}^{2i+1}$. By the Hilton–Milnor theorem, $\Omega(\bigvee_{i=1}^{k-1} \bigvee^{m_i} S_{(p)}^{2i+1})$ is homotopy equivalent to a weak product of the loop spaces of p -local odd spheres of dimension ≥ 3 . Then since $p > k$ and $\pi_{2j}(S^{2l+1})_{(p)} = 0$ for $j < l + p - 1$, the attaching maps are null homotopic, hence the induction proceeds. \square

4. Non-triviality of the projection π

Let M be a quasitoric manifold over an n -polytope P and let $\pi: \mathcal{Z}_{K(P)} \rightarrow M$ denote the projection. By Corollary 1.4, $\Sigma\pi_{(p)}$ is trivial for $p > n$. So one would ask whether $\pi_{(p)}$ and $\Sigma^\infty\pi$ are trivial or not. This section shows non-triviality of $\pi_{(p)}$ and examines non-triviality of $\Sigma^\infty\pi$ for quasitoric manifolds over a product of simplices and low dimensional quasitoric manifolds. We first consider the p -localization.

Proposition 4.1. *The p -localization $\pi_{(p)}$ is not null homotopic for any prime p .*

Proof. Recall that there is a homotopy fibration (2.1). Then if $\pi_{(0)}$ were null homotopic, we would have $T_{(0)}^{m-n} \simeq (\mathcal{Z}_{K(P)})_{(0)} \times \Omega M_{(0)}$, implying that $\mathcal{Z}_{K(P)}$ is rationally contractible since it is simply connected [2, Corollary 6.19]. On the other hand, $\mathcal{Z}_{K(P)}$ is a compact simply connected $m + n$ -dimensional manifold without boundary by [2, Lemma 6.2]. Then $\mathcal{Z}_{K(P)}$ is not rationally contractible, a contradiction. Therefore $\pi_{(0)}$ is not null homotopic, completing the proof. \square

We next consider non-triviality of $\Sigma^\infty\pi$ for quasitoric manifolds over a product of simplices. We start with the easiest case. Recall that the complex projective space $\mathbb{C}P^n$ is the only quasitoric manifold over the n -simplex Δ^n , and that the projection π is the canonical map $S^{2n+1} \rightarrow \mathbb{C}P^n$. Then since the cofiber of π is $\mathbb{C}P^{n+1}$ whose top cell does not split after stabilization, one sees that $\Sigma^\infty\pi$ is not null homotopic. We here record this almost trivial fact.

Lemma 4.2. *The projection $\pi: \mathcal{Z}_{K(\Delta^n)} \rightarrow \mathbb{C}P^n$ is not null homotopic after stabilization.*

It is helpful to recall the fact on moment-angle complexes regarding products of simple polytopes. For simple polytopes P_1, P_2 the product $P_1 \times P_2$ is also a simple polytope and $K(P_1 \times P_2) = K(P_1) * K(P_2)$, the join of $K(P_1)$ and $K(P_2)$. By definition we have $\mathcal{Z}_{K(P_1 \times P_2)} = \mathcal{Z}_{K(P_1) * K(P_2)} = \mathcal{Z}_{K(P_1)} \times \mathcal{Z}_{K(P_2)}$, and in particular $\mathcal{Z}_{K(P_1)}$ is a retract of $\mathcal{Z}_{K(P_1 \times P_2)}$. We prove a simple lemma needed later.

Lemma 4.3. *Let P be a simple polytope, and let M be a quasitoric manifold over $P \times \Delta^k$. If there is a map $q: M \rightarrow \mathbb{C}P^k$ satisfying a homotopy commutative diagram*

$$\begin{array}{ccc} \mathcal{Z}_{K(P \times \Delta^k)} & \xrightarrow{\text{proj}} & \mathcal{Z}_{K(\Delta^k)} \\ \downarrow \pi & & \downarrow \pi \\ M & \xrightarrow{q} & \mathbb{C}P^k, \end{array}$$

then the projection $\pi: \mathcal{Z}_{K(P \times \Delta^k)} \rightarrow M$ is not null homotopic after stabilization.

Proof. Since $\mathcal{Z}_{K(\Delta^k)}$ is a retract of $\mathcal{Z}_{K(P \times \Delta^k)}$, it follows from Lemma 4.2 that $\Sigma^\infty(q \circ \pi)$ is not null homotopic. Therefore since $\Sigma^\infty(q \circ \pi) = \Sigma^\infty q \circ \Sigma^\infty \pi$, the proof is completed. \square

There is a class of quasitoric manifolds over a product of simplices called generalized Bott manifolds which have been intensively studied in toric topology. See [3] for details. By definition a generalized Bott manifold B over $\Delta^{n_1} \times \dots \times \Delta^{n_l}$ satisfies a commutative diagram

$$\begin{array}{ccccccc} \mathcal{Z}_{K(\Delta^{n_1} \times \dots \times \Delta^{n_l})} & \longrightarrow & \mathcal{Z}_{K(\Delta^{n_1} \times \dots \times \Delta^{n_{l-1}})} & \longrightarrow & \dots & \longrightarrow & \mathcal{Z}_{K(\Delta^{n_1} \times \Delta^{n_2})} & \longrightarrow & \mathcal{Z}_{K(\Delta^{n_1})} \\ \downarrow \pi & & \downarrow \pi & & & & \downarrow \pi & & \downarrow \pi \\ B_l & \xrightarrow{q_l} & B_{l-1} & \xrightarrow{q_{l-1}} & \dots & \xrightarrow{q_2} & B_2 & \xrightarrow{q_1} & B_1 \end{array}$$

where the upper horizontal arrows are the projections. Since $B_1 = \mathbb{C}P^{n_1}$, we get the following by Lemma 4.3

Corollary 4.4. *If B is a generalized Bott manifold over $\Delta^{n_1} \times \dots \times \Delta^{n_l}$, then the projection $\pi: \mathcal{Z}_{K(\Delta^{n_1} \times \dots \times \Delta^{n_l})} \rightarrow B$ is not null homotopic after stabilization.*

In order to examine non-triviality of $\Sigma^\infty \pi$ for quasitoric manifolds other than Bott manifolds, we give a cohomological generalization of Lemma 4.3.

Lemma 4.5. *Let X be a space such that $H^2(X; \mathbb{Z}) = \mathbb{Z}\langle x_1, \dots, x_k \rangle$ and $H^{\text{odd}}(X; \mathbb{Z}/p) = 0$, and let F be the homotopy fiber of a map $\alpha = (x_1, \dots, x_k): X \rightarrow BT^k$. Suppose the following conditions hold:*

- (1) *There are $x \in H^{2l-2i}(BT^k; \mathbb{Z}/p)$ and transgressive $a \in H^{2l-1}(F; \mathbb{Z}/p)$ such that*

$$\tau(a) = \theta(x)$$

for some degree $2i$ Steenrod operation θ .

- (2) *There is a map $f: S^{2l-1} \rightarrow F$ such that $f^*(a) \neq 0$ in mod p cohomology.*

Then the stabilization of the fiber inclusion $F \rightarrow X$ is not null homotopic.

Proof. Let $i: F \rightarrow X$ and $j: X \rightarrow C_{i \circ f}$ denote the inclusions, where C_g means the mapping cone of a map g . Then there is a commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H^{2l-1}(S^{2l-1}; \mathbb{Z}/p) & \xrightarrow{\delta} & H^{2l}(C_{i \circ f}; \mathbb{Z}/p) & \xrightarrow{j^*} & H^{2l}(X; \mathbb{Z}/p) \longrightarrow 0 \\
 & & \uparrow f^* & & \uparrow \bar{f}^* & & \uparrow \alpha^* \\
 & & H^{2l-1}(F; \mathbb{Z}/p) & \xrightarrow{\delta} & H^{2l}(X, F; \mathbb{Z}/p) & \xleftarrow{\alpha^*} & H^{2l}(BT^k; \mathbb{Z}/p)
 \end{array}$$

with exact top row, where $\bar{f}: C_{i \circ f} \rightarrow C_i$ denotes the map induced by id_X and f . Put $\bar{x} = \bar{f}^* \circ \alpha^*(x)$. Since $\tau(a) = \theta(x)$, we have $\theta(\bar{x}) = \bar{f}^* \circ \alpha^*(\theta(x)) = \delta \circ f^*(a) \neq 0$. Then we see that any splitting of the top row

$$H^*(C_{i \circ f}; \mathbb{Z}/p) \cong A \oplus \langle \theta(\bar{x}) \rangle, \quad A \cong H^*(X; \mathbb{Z}/p),$$

as modules implies that $\theta(A) \not\subset A$ by $\bar{x} \in A$. If $\Sigma^\infty i$ were null homotopic, we would have $\theta(A) \subset A$ which contradicts to the above calculation, so $\Sigma^\infty i$ is not null homotopic. □

We apply Lemma 4.5 to quasitoric manifolds over a product of two simplices which are not necessarily generalized Bott manifolds.

Proposition 4.6. *If M is a quasitoric manifold over $\Delta^k \times \Delta^{n-k}$ and neither $k + 2$ nor $n - k + 2$ is a power of 2, then $\Sigma^\infty \pi$ is not null homotopic.*

Proof. By Lemma 4.2 we may assume $0 < k < n$. It follows from Proposition 2.2 that $H^2(M; \mathbb{Z})$ is a free abelian group with a basis \bar{x}, \bar{y} . Let $\alpha: M \rightarrow BT^2$ be the classifying map of the principal bundle $T^2 \rightarrow \mathcal{Z}_{K(P)} \rightarrow M$, and put $\beta = (\bar{x}, \bar{y}): M \rightarrow BT^2$. Then by Proposition 2.2 there is a self map h of BT^2 satisfying $\beta \simeq h \circ \alpha$, so it is sufficient to show that the inclusion of the homotopy fiber of β is not null homotopic by applying Lemma 4.5. By [3] the mod 2 cohomology of M is given by

$$H^*(M; \mathbb{Z}/2) = \mathbb{Z}/2[x, y]/(x^{k'-l+1}(x+y)^l, y^{n-k'+1})$$

for some $l \geq 0$, where $k' = k$ or $k' = n - k$ and x, y are the mod 2 reduction of \bar{x}, \bar{y} respectively. Choose $t \in H^2(BT^2; \mathbb{Z}/2)$ satisfying $\beta^*(t) = y$. Let r be the largest integer satisfying $n - k' + 1 > 2^r - 1$. Then since $n - k' + 2$ is not a power of 2, we have $(n - k' + 1) - (2^r - 1) \leq 2^r - 1$, so we get $\binom{2^r - 1}{(n - k' + 1) - (2^r - 1)} \not\equiv 0 \pmod 2$ by Lucas' theorem. Thus we obtain

$$\text{Sq}^{2\{(n - k' + 1) - (2^r - 1)\}} t^{2^r - 1} = \binom{2^r - 1}{(n - k' + 1) - (2^r - 1)} t^{n - k' + 1} = t^{n - k' + 1}.$$

Since $\mathcal{Z}_{K(\Delta^k \times \Delta^{n-k})} = S^{2k+1} \times S^{2(n-k)+1}$, there is a spherical $a \in H^{2(n-k')+1}(\mathcal{Z}_{K(\Delta^k \times \Delta^{n-k})}; \mathbb{Z}/2)$ satisfying $\tau(a) = t^{n-k'+1}$ for a degree reason. Therefore the proof is done. \square

We next specialize Lemma 4.5 for applications to low dimensional quasitoric manifolds.

Proposition 4.7. *Let M be a quasitoric manifold. If there is non-zero $x \in H^2(M; \mathbb{Z}/2)$ satisfying $x^2 = 0$, then $\Sigma^\infty \pi$ is not null homotopic.*

Proof. It is sufficient to check that the conditions of Lemma 4.5 are satisfied. Let P be a polytope on which M stands. Since $\mathcal{Z}_{K(P)}$ is 2-connected, there is $a \in H^3(\mathcal{Z}_{K(P)}; \mathbb{Z}/2)$ satisfying $\tau(a) = t^2$, where $t \in H^2(BT^{m-n}; \mathbb{Z}/2)$ satisfies $\alpha^*(t) = x$. Then for $t^2 = \text{Sq}^2 t$, the condition (1) of Lemma 4.5 is satisfied. We also have that the Hurewicz map $\pi_3(\mathcal{Z}_{K(P)}) \rightarrow H_3(\mathcal{Z}_{K(P)}; \mathbb{Z})$ is an isomorphism, so any element of $H^3(\mathcal{Z}_{K(P)}; \mathbb{Z}/2)$ is spherical. Then the condition (2) of Lemma 4.5 is satisfied, and therefore the proof is done. \square

We now apply Proposition 4.7 to low dimensional quasitoric manifolds.

Corollary 4.8. *If M is a 4-dimensional quasitoric manifold, then $\Sigma^\infty \pi$ is not null homotopic.*

Proof. Suppose that the quasitoric manifold M stands over a 2-polytope P . If $P = \Delta^2$, the corollary follows from Lemma 4.2 since $\mathbb{C}P^2$ is the only quasitoric manifold over Δ^2 . If $P \neq \Delta^2$, then P is a k -gon for $k \geq 4$, hence $h_2(P) = 1 < k - 2 = h_1(P)$. Then it follows from Proposition 2.1 that $\dim H^4(M; \mathbb{Z}/2) < \dim H^2(M; \mathbb{Z}/2)$, implying that there must be non-zero $x \in H^2(M; \mathbb{Z}/2)$ satisfying $x^2 = 0$. Thus the proof is completed by Proposition 4.7. \square

REMARK 4.9. We here remark that $h_1(P) = h_2(P)$ by the Dehn–Sommerville equation for $\dim P = 3$ and $h_1(P) \leq h_2(P)$ for $\dim P > 3$ by the g -theorem (cf. [2]), so the argument in the proof of Corollary 4.8 does not work for $\dim P \geq 3$.

Corollary 4.10. *If M is a quasitoric manifold over the 3-cube, then $\Sigma^\infty \pi$ is not null homotopic.*

Proof. It is calculated in [3, 5] that the mod 2 cohomology of M is given by

$$H^*(M; \mathbb{Z}/2) = \mathbb{Z}/2[x, y, z]/(x^2 + x(ay + bz), y^2 + y(cx + dz), z^2 + z(ex + fy))$$

for $a, b, c, d, e, f \in \mathbb{Z}/2$ satisfying

$$ac = df = 0, \quad \begin{vmatrix} 1 & c & e \\ a & 1 & f \\ b & d & 1 \end{vmatrix} = 1.$$

We now suppose that $w^2 \neq 0$ for all non-zero $w \in H^2(M; \mathbb{Z}/2)$. Then for $x_1^2 \neq 0$ we have (a, b) is either $(1, 0)$, $(0, 1)$, $(1, 1)$. Consider the case $(a, b) = (1, 0)$. That $a = 1$ implies $c = 0$, so $d = 1$ since $y^2 \neq 0$. Then $f = 0$, implying $e = 1$ since $z^2 \neq 0$. Hence we obtain $\begin{vmatrix} 1 & c & e \\ a & 1 & f \\ b & d & 1 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix} = 0$, a contradiction. In the case $(a, b) = (0, 1)$, $(1, 1)$ we can similarly get $(c, d, e, f) = (0, 1, 1, 0)$, so a contradiction occurs. Thus there is non-zero $w \in H^2(M; \mathbb{Z}/2)$ with $w^2 = 0$, and therefore the proof is done by Proposition 4.7. □

For the last we dare to conjecture the following from Propositions 4.6, 4.7 and Corollaries 4.4, 4.8 and 4.10.

CONJECTURE 4.11. For any quasitoric manifold M , $\Sigma^\infty \pi$ is not null homotopic.

References

- [1] A. Bahri, M. Bendersky, F.R. Cohen and S. Gitler: *The polyhedral product functor: a method of decomposition for moment-angle complexes, arrangements and related spaces*, Adv. Math. **225** (2010), 1634–1668.
- [2] V.M. Buchstaber and T.E. Panov: *Torus Actions and Their Applications in Topology and Combinatorics*, University Lecture Series **24**, Amer. Math. Soc., Providence, RI, 2002.
- [3] S. Choi, M. Masuda and D.Y. Suh: *Quasitoric manifolds over a product of simplices*, Osaka J. Math. **47** (2010), 109–129.
- [4] M.W. Davis and T. Januszkiewicz: *Convex polytopes, Coxeter orbifolds and torus actions*, Duke Math. J. **62** (1991), 417–451.
- [5] S. Hasui: *On the cohomology equivalences between bundle-type quasitoric manifolds over a cube*, arXiv:math-AT/1409.7980.
- [6] M. Mimura, G. Nishida and H. Toda: *Localization of CW-complexes and its applications*, J. Math. Soc. Japan **23** (1971), 593–624.

Sho Hasui
Department of Mathematics
Kyoto University
Kyoto, 606-8502
Japan
e-mail: s.hasui@math.kyoto-u.ac.jp

Daisuke Kishimoto
Department of Mathematics
Kyoto University
Kyoto, 606-8502
Japan
e-mail: kishi@math.kyoto-u.ac.jp

Takashi Sato
Department of Mathematics
Kyoto University
Kyoto, 606-8502
Japan
e-mail: t-sato@math.kyoto-u.ac.jp