



Title	On multiply transitive permutation groups. I
Author(s)	Bannai, Eiichi
Citation	Osaka Journal of Mathematics. 1974, 11(2), p. 401-411
Version Type	VoR
URL	<a href="https://doi.org/10.18910/6086">https://doi.org/10.18910/6086</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## ON MULTIPLY TRANSITIVE PERMUTATION GROUPS I

EIICHI BANNAI<sup>\*)</sup>

(Received October 8, 1973)

### Introduction

In [5], M. Hall determined 4-ply transitive permutation groups whose stabilizer of 4 points is of odd order. (See also Nagao [11].) On the other hand, in Bannai [1] and Miyamoto [9],  $t$ -ply transitive finite permutation groups in which the stabilizer of  $t$  points is of order prime to an odd prime  $p$  have been determined for  $t=p^2+p$  and  $3p$  respectively. The purpose of this series of notes is to strengthen those results. In this first note, we will improve Lemma 2.1 in Miyamoto [9]. Namely, we will prove the following result.

**Theorem 1.** *Let  $p$  be an odd prime. Then there exists no permutation group  $G$  on a set  $\Omega=\{1, 2, \dots, n\}$  which satisfies the following three conditions:*

- (i)  *$G$  is  $(p+2)$ -ply transitive, and  $n \equiv 2 \pmod{p}$ ,*
- (ii) *a Sylow  $p$  subgroup  $P_0$  of  $G_{1,2,\dots,p+2}$  is semiregular on  $\Omega - \{1, 2, \dots, p+2\}$ , and*
- (iii)  *$|P_0| \geq p^2$ .*

**Corollary to Theorem 1.** *Let  $p$  be an odd prime. Let  $G$  be a  $(2p+2)$ -ply transitive permutation group on a set  $\Omega=\{1, 2, \dots, n\}$ . If the order of  $G_{1,2,\dots,2p+2}$  is not divisible by  $p$ , then  $G$  must be  $S_n$  ( $2p+2 \leq n \leq 3p+1$ ) or  $A_n$  ( $2p+4 \leq n \leq 3p+1$ ).*

This corollary is immediately proved by combining Theorem 1 with a result of Miyamoto [9]. To be more precise, if the order of  $G_{1,2,\dots,p+2}$  is not divisible by  $p^2$ , then the  $2p$ -ply transitive group  $G_{1,2}$  on  $\Omega - \{1, 2\}$  must contain  $A^{\Omega - \{1,2\}}$  by the result of Miyamoto [9, §1], and so  $G$  must be one of the groups listed in the conclusion of the corollary. If the order of  $G_{1,2,\dots,p+2}$  is divisible by  $p^2$ , then the  $(p+2)$ -ply transitive group  $G_{1,2,\dots,i}$  on  $\Omega - \{1, 2, \dots, i\}$  (if  $n \equiv i+2 \pmod{p}$  with  $0 \leq i \leq p-1$ ) satisfies the three conditions of Theorem 1, and we have a contradiction.

In our proof of Theorem 1, the following result is very important. This result is a kind of generalization of a result of Jordan [8, Chap. IV], and will be of independent interest.

---

<sup>\*</sup>) Supported in part by the Sakkokai Foundation.  
Present address: The Ohio State University.

**Theorem A.** *Let  $p$  be an odd prime. Then  $A_{p+2}$  (hence  $S_{p+2}$ ) is not involved in  $GL(p, p)$ .*

Theorem A will be proved in §1 by exploiting the theory of modular representations of the symmetric groups due to Nakayama [12] together with some other results (theory of projective representations of the symmetric groups due to Schur [13], theory of  $p$  groups and so on).

In Appendix, we will discuss some partial generalization of Theorem 1.

**Notation.** Our notation will be standard.  $S^\Delta$  and  $A^\Delta$  denote the symmetric and alternating groups on a set  $\Delta$ . If  $|\Delta|$ , the cardinality of  $\Delta$ , is  $m$ , we denote them by  $S_m$  and  $A_m$  instead of  $S^\Delta$  and  $A^\Delta$ . If  $X$  is a permutation group on a set  $\Omega$ , and if  $\Delta$  is a subset of  $\Omega$  which is fixed as a whole by  $X$ , we denote by  $X^\Delta$  the restriction of  $X$  to  $\Delta$ . For a subset  $\Delta = \{1, 2, \dots, i\}$  of  $\Omega$ , we denote by  $X_{1,2,\dots,i}$  the pointwise stabilizer of  $\Delta$  in  $X$ .  $GL(m, K)$  denotes the general linear group of dimension  $m$  over a field  $K$ .  $PGL(m, K)$  denotes the projective linear group of dimension  $m$  over  $K$ ,  $PGL(m, K) = GL(m, K)/Z(GL(m, K))$ , where  $Z(GL(m, K))$  denotes the center of  $GL(m, K)$ . When  $K$  is of cardinality  $p$ , we denote  $GL(m, K)$  by  $GL(m, p)$ . For a group  $X$ ,  $\text{Aut}(X)$  denotes the automorphism group of  $X$ .

### 1. $A_{p+2}$ is not involved in $GL(p, p)$

The purpose of this section is to prove Theorem A that  $A_{p+2}$  is not involved in  $GL(p, p)$ .

We first remark the following lemma.

**Lemma 1.** *Theorem A is true for  $p=3$  and 5.*

*Proof.*  $A_5$  is not involved in  $GL(3, 3)$ , because the order of  $GL(3, 3)$  is not divisible by 5. Similarly,  $A_7$  is not involved in  $GL(5, 5)$ , because the order of  $GL(5, 5)$  is not divisible by 7.

From now on, we always assume that  $p \geq 7$ . In case of  $p \geq 7$ , we can prove Theorem A in a little stronger form as in Lemma 4 mentioned later.

**Lemma 2.** *Let  $p \geq 7$ . Then  $S_{p+2}$  is not a subgroup of  $GL(p, K)$ , where  $K$  is an algebraically closed field of characteristic  $p$ .*

*Proof.* We have only to prove that  $S_{p+2}$  has no faithful  $p$ -modular (absolutely) irreducible representation of degree  $\leq p$  over  $K$ . Lemma 2 will be proved through the following steps (1) and (2).

(1) The degree of any not 1 dimensional ordinary irreducible representation of  $S_k$  ( $k \geq 5$ ) is  $\geq k-1$ . Therefore, the degree of any irreducible  $p$ -modular representation of  $S_{p+2}$  over  $K$  which is contained in a  $p$ -block of defect 0 is more than  $p$ .

The first assertion is immediately proved by using the Schur's recursive formula (a special case of Murnaghan-Nakayama's recursive formula) (see [13, §44]). The last assertion is obvious from an elementary properties of a  $p$ -block of defect 0.

(2) The degree of any not 1 dimensional  $p$ -modular irreducible representation of  $S_{p+2}$  over  $K$  which is contained in a  $p$ -block of defect 1 is more than  $p$ .

By Nakayama [12], we obtain that there exist just two  $p$ -blocks of defect 1 for  $S_{p+2}$ . Moreover, one block (say  $B_0$ ) with  $p$ -core of type [2] consists of  $p$  ordinary irreducible representations  $T_{0,r}$  with  $0 \leq r \leq p-1$ , where  $T_{0,r}$  is the representation associated with the Young diagram of type  $[p+2]$  (for  $r=0$ ),  $[p-r, 3, 1^{r-1}]$  (for  $1 \leq r \leq p-3$ ),  $[2, 2, 1^{p-2}]$  (for  $r=p-2$ ) and  $[2, 1^p]$  (for  $r=p-1$ ). While, the other block (say  $B_1$ ) with  $p$ -core of type  $[1^2]$  consists of  $p$  ordinary irreducible representations  $T_{1,r}$  with  $0 \leq r \leq p-1$ , where  $T_{1,r}$  is the representation associated with the Young diagram which is obtained by transposing that of  $T_{0,p-1-r}$ . Also by a result of Nakayama [12],  $T_{i,r}$  and  $T_{i,r+1}$  ( $i=0,1$ ,  $r=0, 1, \dots, p-1$ ) have just one  $p$ -modular irreducible representation (over  $K$ ) in common, say, let us denote it by  $\phi_{i,r}$  ( $0 \leq r \leq p-2$ ), and  $T_{i,r}$  and  $T_{i,s}$  with  $s > r+1$  have no  $p$ -modular irreducible representation in common. That is to say, the Brauer graphs associated with the  $p$ -blocks  $B_i$  ( $i=0, 1$ ) are trees without branches and their nodes are arranged on natural order on  $r$ . (For the definition of Brauer graphs, see, e.g., [3, §68].) Therefore, we can calculate the degree  $|\phi_{i,r}|$  of  $\phi_{i,r}$  inductively for  $r=0, 1, 2, \dots$  (and for  $r=p-2, p-3, p-4, \dots$ ), because the degree  $|T_{i,r}|$  of  $T_{i,r}$  is given explicitly by the following formula:

$$|T_{i,r}| = (p+2)! / (\text{the product of all hook lengths of the Young diagram of } T_{i,r}).$$

In the case of  $p=7$ , we can immediately calculate all the values of  $|\phi_{i,r}|$  ( $i=0, 1$ ,  $r=0, 1, \dots, 5$ ), and we obtain that they are all  $\geq 8 > 7$  except  $|\phi_{0,0}|$  and  $|\phi_{1,5}|$  which are equal to 1. Thus, in the following we may assume that  $p \geq 11$ .

$$\text{Now, we obtain that } |T_{0,0}|=1, |T_{0,1}| = \frac{(p+2)(p+1)(p-3)}{6} \geq p^2, |T_{0,p-3}| = \frac{(p+2)(p+1)(p-2)(p-3)}{12} \geq p^2, |T_{0,p-2}| = \frac{(p+1)(p-1)}{2} \text{ and } |T_{0,p-1}| = p+1.$$

Therefore, we obtain  $|\phi_{0,0}|=1, |\phi_{0,p-3}| > p, |\phi_{0,p-2}| = p+1 > p$ . Moreover, when  $1 \leq r \leq p-4$ , we obtain that  $|T_{0,r}| / |T_{0,r+1}| = \frac{r(r+3)(p-r-2)}{(r+2)(p-r)(p-3-r)}$ . Now, we

obtain that  $\frac{1}{p} < \frac{r(r+3)(p-r-2)}{(r+2)(p-r)(p-3-r)} < p$  for any  $r=1, 2, \dots, p-2$ , and

$$\frac{r(r+3)(p-r-2)}{(r+2)(p-r)(p-3-r)} < 1 \text{ when } r \leq \frac{p-1}{2}, \text{ and } > 1 \text{ when } r \geq \frac{p+1}{2}.$$

Therefore, we obtain that  $|T_{0,1}| < |T_{0,2}| < \dots < |T_{0,(p+1)/2}|$ , and  $|T_{0,(p+1)/2}| > \dots > |T_{0,p-2}| > |T_{0,p-1}|$ . Hence, we obtain that  $|\phi_{0,r}| > p$  (for  $r=1, 2, \dots, p-4$ ), because of the

fact that the Brauer graph of the block  $B_0$  is a tree without branches and of natural order on  $r$ . Therefore, we obtain that  $|\phi_{0,r}| > p$  for any  $r \neq 0$  ( $1 \leq r \leq p-2$ ). Since  $|T_{0,r}| = |T_{1,p-1-r}|$  and  $|\phi_{0,r}| = |\phi_{1,p-2-r}|$  for any  $r$ , we also obtain that  $|\phi_{1,r}| > p$  for any  $r \neq p-2$  ( $0 \leq r \leq p-3$ ) and  $|\phi_{1,p-2}| = 1$ . Thus, we have proved the assertion of (2).

Since any  $p$ -block of  $S_{p+2}$  is either of defect 0 or 1, we have completed the proof of Lemma 2 by (1) and (2).

We also have

**Lemma 2'.** *Let  $p \geq 7$ . Then  $A_{p+2}$  is not a subgroup of  $GL(p, K)$ , where  $K$  is an algebraically closed field of characteristic  $p$ .*

*Proof.* The assertion corresponding to step (1) in Lemma 2 is easily obtained similarly by using the Schur's recursive formula for the characters of the symmetric groups. Namely, we have

(1') The degree of any ( $p$ -modular) irreducible representation of  $A_{p+2}$  over  $K$  which is contained in a  $p$ -block of defect 0 is more than  $p$ .

Since  $T_{i,r}$  ( $i=0, 1, r=0, 1, \dots, p-1$ ) are all irreducible representations of  $A_{p+2}$ , we immediately obtain the following assertion.

(2') The degree of not 1 dimensional ( $p$ -modular) irreducible representation of  $A_{p+2}$  over  $K$  which is contained in a  $p$ -block of defect 1 is more than  $p$ .

Thus, we have proved Lemma 2'.

**Lemma 3.** *Let  $p$  be an odd prime  $\geq 7$ . Then  $S_{p+2}$  is not a subgroup of  $PGL(p, K)$ , where  $K$  is an algebraically closed field of characteristic  $p$ .*

*Proof.* We have only to prove that  $S_{p+2}$  has no not 1 dimensional projective irreducible representation of degree  $\leq p$  over  $K$ . Since we have already proved in Lemma 2 that  $S_{p+2}$  is not a subgroup of  $GL(p, K)$ , we have only to prove that  $S_{p+2}$  has no projective representation of degree  $\leq p$  over  $K$  which is not a linear representation. As is easily seen from a result of Schur (and a slight extension of it) (cf. Yamazaki [15, §3.3, Corollary 1]), there is a finite group (which is a central extension of  $S_{p+2}$  and is called a representation group of  $S_{p+2}$  over  $K$ ) such that any projective representation of  $S_{p+2}$  is induced by a linear representation of the representation group. Moreover, by Yamazaki [15, §3, e.g., Proposition 3.3, 2) and Proposition 3.5], we may take as a representation group of  $S_{p+2}$  over  $K$  the following group  $T_{p+2}$  defined by the generators

$$\{J, X_i \ (i = 1, 2, \dots, p+1)\}$$

with the defining relations

$$\begin{aligned} J^2 &= 1, \\ X_\alpha^2 &= J \ (\alpha = 1, 2, \dots, p+1), \end{aligned}$$

$$(X_\beta X_{\beta+1})^3 = J \quad (\beta = 1, 2, \dots, p) \quad \text{and} \\ X_\gamma X_\delta = J X_\delta X_\gamma \quad (\gamma = 1, 2, \dots, p-1, \delta = \gamma+2, \dots, p+1).$$

(Note that  $Z(T_{p+2}) = \langle J \rangle$  (which is contained in the commutator subgroup of  $T_{p+2}$ ) is a cyclic group of order 2, and  $T_{p+2}/Z(T_{p+2}) = S_{p+1}$ .  $T_{p+2}$  is the group denoted  $\mathfrak{T}_{p+2}$  in Schur [13]. Also note that  $H^2(S_{p+2}, K^*) = H^2(S_{p+2}, \mathbb{C}^*) = \mathbb{Z}_2$ .)

The ordinary irreducible representations of  $T_{p+2}$  were completely determined by Schur [13]. As in [13], let us call an ordinary irreducible representation of  $T_{p+2}$  is of the first kind (resp. of the second kind) if the kernel of the representation contains  $Z(T_{p+2})$  (resp. does not contain  $Z(T_{p+2})$ ). The proof of Lemma 3 will be done through the following steps (1), (2) and (3).

(1) The degree of any ordinary irreducible representation of  $T_{p+2}$  of the second kind is more than  $2^{\lfloor (p+1)/2 \rfloor}$ . Moreover  $2^{\lfloor (p+1)/2 \rfloor} > p$ .

The degree of any ordinary irreducible representations of  $T_{p+2}$  of the second kind is given as follows (Schur [13]):

$$f_{\nu_1, \nu_2, \dots, \nu_m} = 2^{\lfloor (p+2-m)/2 \rfloor} g_{\nu_1, \nu_2, \dots, \nu_m},$$

with

$$g_{\nu_1, \nu_2, \dots, \nu_m} = \frac{(p+2)!}{\nu_1! \nu_2! \dots \nu_m!} \prod_{\alpha < \beta} \frac{\nu_\alpha - \nu_\beta}{\nu_\alpha + \nu_\beta},$$

where  $\nu_1 + \nu_2 + \dots + \nu_m = p+2$  and  $\nu_1 > \nu_2 > \dots > \nu_m > 0$ . Moreover, by Schur [13, §44], it is proved that

$$f_{\nu_1, \nu_2, \dots, \nu_m} \geq 2^{\lfloor (p+2-1)/2 \rfloor} = 2^{\lfloor (p+1)/2 \rfloor}$$

for any  $f_{\nu_1, \nu_2, \dots, \nu_m}$ . Thus we obtain the first assertion. The last assertion is clear, because  $p \geq 7$ .

(2) The degree of any ordinary irreducible representation of  $T_{p+2}$  of the second kind which is not divisible by  $p$  is divisible by  $2^{\lfloor (p-1)/2 \rfloor}$ . Moreover,  $2^{\lfloor (p-1)/2 \rfloor} > p$ .

Since  $f_{\nu_1, \nu_2, \dots, \nu_m}$  is not divisible by  $p$ , we obtain that  $m \leq 3$ , by noticing the formula of  $f_{\nu_1, \nu_2, \dots, \nu_m}$ . Since  $f_{\nu_1, \nu_2, \dots, \nu_m} = 2^{\lfloor (p+2-m)/2 \rfloor} g_{\nu_1, \nu_2, \dots, \nu_m}$  and  $g_{\nu_1, \nu_2, \dots, \nu_m}$  is an integer (Schur [13, §40]), we obtain the first assertion. The last assertion is clear, because  $p \geq 7$ .

(3) The degree of any not 1 dimensional ( $p$ -modular) irreducible representation of  $T_{p+2}$  over  $K$  is more than  $p$ .

Let  $\phi$  be an irreducible representation of  $T_{p+2}$  over  $K$  of degree  $> 1$ . If  $\phi$  is contained in a  $p$ -block of defect 0 of  $T_{p+2}$ , then by step (1) and the step (1) in Lemma 2, we obtain that the degree of  $\phi$  is more than  $p$ . Now, let us assume that  $\phi$  is contained in a  $p$ -block of defect 1. Since any block of defect 1 contains at most  $p$  ordinary irreducible representations in general (and in this case) (cf. [3, §68]),  $B_0$  and  $B_1$  ( $p$ -blocks of  $S_{p+2}$ ) themselves also become  $p$ -blocks

of  $T_{p+2}$  of defect 1 (all representation of  $S_{p+2}$  are naturally regarded as representations of  $T_{p+2}$ ). Therefore, any ordinary irreducible representation of  $T_{p+2}$  which is contained in a  $p$ -block of defect 1 and not contained in  $B_0$  and  $B_1$  (as blocks of  $T_{p+2}$ ) must be of the second kind. Therefore, the degree of any ordinary irreducible representation of  $T_{p+2}$  contained in a  $p$ -block of defect 1 and not contained in  $B_0$  and  $B_1$  must be divisible by  $2^{\lfloor (p-1)/2 \rfloor}$  by step (2). Since  $p$  is to the first power in the order of  $T_{p+2}$ , the Brauer graph of any  $p$ -block of defect 1 of  $T_{p+2}$  must be a tree (cf. [3, §68]), and so the degree of any irreducible representation of  $T_{p+2}$  over  $K$  is divisible by  $2^{\lfloor (p-1)/2 \rfloor} > p$ . Thus, we obtain the assertion of (3).

Thus, we have completed the proof of Lemma 3.

We also have

**Lemma 3'.** *Let  $p \geq 7$ . Then  $A_{p+2}$  is not a subgroup of  $PGL(p, K)$ , where  $K$  is an algebraically closed field of characteristic  $p$ .*

Proof. The commutator subgroup  $T_{p+2}'$  of  $T_{p+2}$  with index 2 becomes a representation group of  $A_{p+2}$  over  $K$ .

(1') The degree of any ordinary irreducible representation of  $T_{p+2}'$  of the second kind is more than  $2^{\lfloor (p+1)/2 \rfloor - 1} > p$ .

Proof is clear.

(2') The degree of any ordinary irreducible representation of  $T_{p+2}'$  of the second kind which is not divisible by  $p$  is divisible by  $2^{\lfloor (p-1)/2 \rfloor - 1}$  and divisible by 8 if  $p=7$ . Moreover,  $2^{\lfloor (p-1)/2 \rfloor - 1} > p$  when  $p \geq 11$ .

Proof of the first assertion is clear. The second assertion for  $p=7$  is proved directly and easily.

(3') The degree of not 1 dimensional ( $p$ -modular) irreducible representation of  $T_{p+2}'$  over  $K$  is more than  $p$ .

The proof is quite the same as that of step (3) in Lemma 3.

Thus, we have proved Lemma 3'.

**Lemma 4.**  *$A_{p+2}$  is not involved in a finite subgroup of  $GL(p, K)$ , where  $K$  is an algebraically closed field of characteristic  $p$ .*

Proof. Let us assume that  $l$  is the smallest integer  $\leq p$  such that  $A_{p+2}$  is involved in a finite subgroup  $X$  of  $GL(l, K)$ . Moreover, let us take  $X$  being of the least order among them, then  $X$  contains a normal subgroup  $Y$  such that  $X/Y = A_{p+2}$ . Now, we will derive a contradiction. By the assumption, we may assume that  $X$  is an irreducible subgroup of  $GL(l, K)$ , and moreover that  $X$  is a primitive subgroup of  $GL(l, K)$ , because  $A_{p+2}$  is obviously not involved in  $S_l$ . (Cf. Dixon [2, §4], see also [2] for some fundamental properties of (finite) linear groups). By Lemma 2 and Lemma 3, we may assume that  $Y$  is not contained in  $Z(GL(l, K))$ . Thus, there exists a Sylow  $q$  subgroup  $Q$  (for some prime

$q$ ) of  $Y$  such that  $Q$  is not contained in  $Z(GL(l, K))$ . By the theorem of Sylow (Frattini argument), and since  $A_{p+2}$  is not involved in  $Y$  by the minimality of the order of  $X$ , we obtain that  $X$  normalizes the Sylow  $q$  subgroup  $Q$  which is not contained in  $Z(GL(l, K))$ . The proof of Lemma 4 will be completed through the following steps (1) to (6).

(1)  $p \neq q$ .

Otherwise,  $X$  becomes not irreducible as a subgroup of  $GL(l, K)$ , and this contradicts the minimality of  $l$ . (Cf. Dixon [2, §§2.2 and 2.8, or §4.2]).

(2)  $Q$  does not contain any characteristic abelian subgroup of rank  $\geq 2$ .

Otherwise,  $X$  becomes imprimitive or not irreducible as a subgroup of  $GL(l, K)$ , and this contradicts the minimality of  $l$ . (Cf. Dixon [2, §4.2].)

(3)  $Q$  is a central product of groups  $Q_1$  and  $Q_2$ , where  $Q_1$  is either 1 or extraspecial  $q$  group, say of order  $q^{2r+1}$ , and  $Q_2$  is either cyclic or  $q=2$  and isomorphic to one of dihedral, generalized quaternion and semidihedral groups of order  $\geq 2^4$ .

Since  $Q$  contains no characteristic abelian subgroup of rank  $\geq 2$ , we obtain the assertion by a result of P. Hall (cf. Gorenstein [4, Theorem 5.4.9]).

Next, we utilize the following important result of Jordan.

**Lemma of Jordan** ([8, Chap. (V, page 56, (3))]. *Let  $q$  be a prime. If  $r$  is a prime such that  $r \neq q$  and  $r \leq k-2$ , then  $A_k$  is not involved in  $GL(r-2, q)$ .*

As a special case of Lemma of Jordan, we obtain the following assertion.

(4)  $A_{p+2}$  is not involved in  $GL(p-2, q)$ , where  $q$  is a prime different from  $p$ .

(5) Let  $x$  be an element of  $GL(l, K)$  which is of order prime to  $p$  and not lying in  $Z(GL(l, K))$ . Then  $A_{p+2}$  is not involved in  $C_{GL(l, K)}(x)$ .

This assertion is well known and immediately proved, e.g., by Dixon [2, §4.2], because  $C_{GL(l, K)}(x)$  becomes either not irreducible or imprimitive as a subgroup of  $GL(l, K)$ .

(6)  $A_{p+2}$  is not involved in  $\text{Aut}(Q)$ .

We obtain that all irreducible components of the natural representation of  $Q$  in  $GL(l, K)$  are equivalent (cf. [2, §4.2]), and so it is a faithful representation of  $Q$ . Now, any faithful ordinary absolutely irreducible representation of  $Q$  (and hence any faithful absolutely irreducible representation of  $Q$  over a field of characteristic  $p \neq q$  (cf. Dixon [2, §3.8]) is) either of degree  $q^r$  (when  $Q_1$  is extraspecial of order  $q^{2r+1}$  and  $Q_2$  is cyclic) or  $q^{r+1}$  (when  $Q_1$  is extraspecial of order  $q^{2r+1}$  and  $Q_2$  is one of dihedral, generalized quaternion and semidihedral and  $q=2$ ), or  $\leq 2$  (when  $Q_1=1$ ) (cf. Gorenstein [4, Theorem 5.5.5 and Theorem 3.7.2]). If  $Q_1=1$ , then we easily have that  $A_{p+2}$  is not involved in  $\text{Aut}(Q)$ , and so in the following we assume that  $Q_1 \neq 1$ . Thus, we obtain in every case that  $q^r \leq l$  ( $\leq p$ ) or  $q^{r+1} \leq l$  ( $\leq p$ ). Now, investigating the structures of the group



$Q$  in every possible case, we obtain that  $Q$  contains a series of characteristic subgroups  $Q_{(i)}$  such that

$$Q = Q_{(0)} > Q_{(1)} > \cdots > Q_{(k)} = 1,$$

and  $Q_{(i)}/Q_{(i+1)}$  ( $i=0, 1, \dots, k-1$ ) are elementary abelian  $q$  subgroups of rank  $\leq 2r$ . Here, note that in every case  $Q/Z(Q)$  is a direct product of  $Q_1/Z(Q)$  (an elementary abelian group of order  $q^{2r}$ ) and a group  $Q_2/Z(Q)$  which is either trivial or one of cyclic subgroups of order  $\geq q^2$  (since, if of order  $q$  then  $Q$  becomes an extraspecial  $q$  group of order  $q^{2r+2}$ , and this is a contradiction) or  $q=2$  and dihedral group of order  $\geq 2^3$ . Therefore, in any way, since  $q^r \leq p$  or  $q^{r+1} \leq p$ , we obtain that  $p-2 \geq 2r$  whenever  $p \geq 7$ . Therefore, in order that  $A_{p+2}$  is involved in  $\text{Aut}(Q)$ ,  $A_{p+2}$  must be involved in  $GL(2r, q)$ , because  $\text{Aut}(Q)/(\text{the stabilizer group of the above chain of characteristic subgroups})$  is a subgroup of the direct product of  $GL(l_i, q)$ 's with  $l_i \leq 2r$ , and the stabilizer group of the chain is a  $q$  group (cf. Gorenstein [4, §5.3]). But, since  $p-2 \geq 2r$ , this contradicts the assertion of (4). Thus, we have obtained the assertion of (6).

Now, we will complete the proof of Lemma 4. Since  $A_{p+2}$  is not involved in  $C_{GL(L, K)}(Q)$  by step (5), and since  $\text{Aut}(Q)$  is a subgroup of  $N_{GL(L, K)}/C_{GL(L, K)}(Q)$ , we obtain that  $A_{p+2}$  is not involved in  $N_{GL(L, K)}(Q)$ . But this is a contradiction, and we have completed the proof of Lemma 4.

Thus, we have completed the proof of Theorem A.

REMARK 1. Theorem A improves Lemma of Jordan (stated preceding step (4) in Lemma 4) a little. That is, we can omit the assumption that  $r \neq q$  in Lemma of Jordan.

REMARK 2. Since it will be not easy for us to follow the proof of Lemma of Jordan along the original paper [8] of Jordan, because of its old fashionedness of its way of description and its terminologies (but not of its context), we give a sketch of an alternative proof.

(a) Let  $q$  be a prime  $\neq p$ . Then  $A_{p+2}$  is not a subgroup of  $GL(p-2, F)$ , where  $F$  is an algebraically closed field of characteristic  $q$ .

$A_{p+2}$  contains a Frobenius group  $H$  of order  $p(p-1)$  whose any Sylow subgroups are cyclic. Since the Schur multipliers of any cyclic subgroups are trivial,  $H^2(H, K^*)$  also becomes trivial (cf. Yamazaki [15, §3]). Therefore, we obtain the assertion by Lemma 1.4 in Harris and Hering [6].

The next assertion will be of independent interest.

(b) Let  $G$  be a finite simple group which is not involved in  $A_8 \cong GL(4, 2)^{11}$ . If the degree of any not 1 dimensional projective (including linear) irreducible

---

1) The assumption that  $G$  is not involved in  $A_8$  is unnecessary in practice, as we can easily see by the case by case considerations of such simple groups.

representation over any (algebraically closed) field of any characteristic is more than  $t$ , then  $G$  is not involved in a finite subgroup of  $GL(t, K)$ , where  $K$  is any (algebraically closed) field of any characteristic.

**Proof.** Let  $l (< t)$  be the smallest integer such that  $G$  is contained in a finite subgroup  $X$  of  $GL(l, K)$  with some algebraically closed field  $K$  of characteristic, say  $s$ . Among them, let us take  $X$  to be of the least order. Because of the assumption, we obtain by quite the same argument as used in the proof of Lemma 3, that  $X$  contains a nontrivial normal Sylow  $q (\neq s)$  subgroup  $Q$  which is not contained in  $Z(GL(l, K))$ , and that  $X$  is not involved in  $C_{GL(l, K)}(Q)$ . Moreover, since  $G$  must be involved in  $\text{Aut}(Q)$ ,  $G$  must be involved in  $GL(2r, q)$ , where  $l \geq q^r$  (or  $q^{r+1}$ ) holds. From the minimality of  $l$ ,  $q^r \leq 2r$ . This asserts that  $q=2$  and  $r=2$  and  $l=4$ . Hence  $G$  must be involved in  $GL(4, 2)$ .

Proof of Lemma of Jordan follows immediately from steps (a) and (b) together with Lemma 3' and Lemma 1.

## 2. Proof of Theorem 1

Let us assume that  $G$  satisfies the three conditions of Theorem 1. Now, we will derive a contradiction.

There is an element  $a$  of  $G$  of order  $p$  such that

$$a = (1)(2)(3, \dots, p+2)(p+3) \cdots (2p+2) \cdots,$$

i.e.,  $a$  fixes  $p+2$  points. Then there exists a Sylow  $p$  subgroup of  $G_{1,2,\dots,p+2}$  which is normalized by the element  $a$ . We may denote it by  $P_0$  without loss of generality. Now, let us set  $P$  be the subgroup generated by  $a$  and  $P_0$ . Then  $P$  is a Sylow  $p$  subgroup of  $G$ .

(1)  $P$  is of maximal class (in the sense of Blackburn). Therefore,  $|Z(P)| = p$ .

Since we obtain that  $|C_{P_0}(a)| = p$  from the semiregularity of  $P_0$  on  $\Omega - \{1, 2, \dots, p+2\}$  (cf. Lemma of Nagao [11]), we have  $|C_P(a)| = p^2$ , and so we have the first assertion (cf. [7, Kapitel III, Satz 14.23]). The last assertion is immediate from the assumption that  $|P_0| \geq p^2$ .

(2)  $N_G(P_0)^{\{1,2,\dots,p+2\}} = S^{\{1,2,\dots,p+2\}}$ .

This assertion is an immediate consequence of Lemma of Witt (cf. [14, Theorem 9.3]).

(3)  $C_G(P_0)^{\{1,2,\dots,p+2\}} \geq A^{\{1,2,\dots,p+2\}}$ .

Otherwise,  $C_G(P_0)^{\{1,2,\dots,p+2\}} = 1$  (because  $p+2 \geq 5$ ), and  $S_{p+2}$  must be involved in  $\text{Aut}(P_0)$ , because  $N_G(P_0)/C_G(P_0)$  is a subgroup of  $\text{Aut}(P_0)$ . Now,  $P_0$  has an automorphism  $\sigma$  (induced from the element  $a$ ) such that the following condition (\*) is satisfied:

$$(*) \quad \sigma \text{ is of order } p \text{ and } |C_{P_0}(\sigma)| = p.$$

If a  $p$  group  $X$  has an automorphism  $\sigma$  satisfying the condition (\*), then any  $\sigma$ -invariant subgroup of  $X$  and any factor group  $X/Y$  for a  $\sigma$ -invariant normal subgroup  $Y$  of  $X$  have the automorphism (naturally induced by  $\sigma$ ) satisfying the condition (\*) provided  $\sigma$  acts nontrivially on them (cf. Huppert [7, Kapitel III, §14], or the argument in Zassenhaus [16, pp. 18–19]), because the map  $\tau$  of  $X$  to  $X$  defined by  $\tau(x)=x^{-1}x^\sigma$  is  $p$  to 1, and if  $(xY)^\sigma=xY$  then  $\tau(x)$  is contained in  $Y$ . Moreover, by a lemma of Ito in Nagao [10], an elementary abelian  $p$  group which has an automorphism with the property (\*) is of rank  $\leq p$ . Thus, if we take a chain of Frattini subgroups  $\Phi^{(i)}(P_0)$  of  $P_0$ :

$$P_0 > \Phi^{(1)}(P_0) > \Phi^{(2)}(P_0) > \dots > \Phi^{(k)}(P_0) = 1,$$

where  $P_0 = \Phi^{(0)}(P_0)$  and  $\Phi^{(1)}(P_0)$  is the Frattini subgroup of  $P_0$  and  $\Phi^{(i+1)}(P_0) = \Phi^{(1)}(\Phi^{(i)}(P_0))$  for  $i \geq 2$ , then  $\Phi^{(i)}(P_0)/\Phi^{(i+1)}(P_0)$  is an elementary abelian  $p$  group of rank  $r_i \leq p$  ( $i=0, 1, \dots, k-1$ ). Therefore, we obtain that

$$\text{Aut}(P_0)/(\text{the stabilizer group of the above chain})$$

is a subgroup of the direct product of the groups  $GL(r_i, p)$  with  $r_i \leq p$  ( $i=0, 1, \dots, k-1$ ), and the stabilizer group of the chain is a  $p$  group. Therefore, since  $S_{p+2}$  is not involved in  $GL(p, p)$  by Theorem A, we obtain that  $S_{p+2}$  is not involved in  $\text{Aut}(P_0)$ . But, this is a contradiction.

Since  $C_G(P_0)^{(1,2,\dots,p+2)} \geq A^{(1,2,\dots,p+2)}$  we obtain that  $|Z(P)| \geq p^2$ . But, this contradicts the fact (1) that  $P$  is of maximal class.

Thus, we have completed the proof of Theorem 1.

## Appendix

In this appendix, we will prove the following result.

**Theorem 2.** *Let  $p$  be an odd prime  $\geq 11$ . Let  $G$  be a permutation group on a set  $\Omega = \{1, 2, \dots, n\}$  which satisfies the following conditions:*

- (i)  *$G$  is  $(p+1)$ -ply transitive, and  $n \equiv 1 \pmod{p}$ ,*
- (ii) *a Sylow  $p$  subgroup  $P_0$  of  $G_{1,2,\dots,p+1}$  is semiregular on  $\Omega - \{1, 2, \dots, p+1\}$ , and*
- (iii)  *$|P_0| \geq p^2$ .*

*Then we obtain that  $P_0$  is an elementary abelian  $p$  group of order  $p^p$  and that a Sylow  $p$  subgroup  $P$  of  $G$  is isomorphic to  $\mathbf{Z}_p \wr \mathbf{Z}_p$  (wreathed product).*

The next Theorem B is proved by quite the same argument as in Theorem A, and so we omit the proof.

**Theorem B.** *Let  $p$  be an odd prime  $\geq 11$ . Then  $S_{p+1}$  is not involved in  $GL(p-1, p)$ .*

**Proof of Theorem 2.** Let  $P$  be a Sylow subgroup of  $G$  which contains  $P_0$ . Then  $P$  is of maximal class. We obtain that  $|P_0/\Phi(P_0)| \leq p^p$ , because of Lemma of Ito in Nagao [10]. Since  $S_{p+1}$  must be involved in  $\text{Aut}(P_0)$  (cf. the proof of Theorem 1) and since  $S_{p+1}$  is not involved in  $GL(p-1, p)$  by Theorem B, we obtain that  $|P_0/\Phi(P_0)| = p^p$ , because of a result of Burnside (cf. Gorenstein 4, Theorem 5.1.4.) (The use of the result of Burnside simplifies the argument of the proof of Theorem 1 a little, i.e., in step (3) we have only to show that  $S_{p+1}$  is not involved in  $\text{Aut}(P_0/\Phi(P_0))$ .) Now,  $P/\Phi(P_0)$  is a homomorphic image of  $P$  and is isomorphic to  $\mathbf{Z}_p \wr \mathbf{Z}_p$ . Therefore, by a result of Blackburn (cf. Huppert [7, Kapitel III, Satz 14.20]) we obtain that  $\Phi(P_0) = 1$ , and so we obtain the assertion of Theorem 2.

UNIVERSITY OF TOKYO

### References

- [1] E. Bannai: *A note on multiply transitive permutation groups* (to appear in J. Algebra).
- [2] J.D. Dixon: *The Structure of Linear Groups*, Van Nostrand Reinhold Math. Studies 37, London, 1971.
- [3] L. Dornhoff: *Group Representation Theory*, Part B, Dekker, New York, 1972.
- [4] D. Gorenstein: *Finite Groups*, Harper and Row, New York, 1968.
- [5] M. Hall, Jr.: *On a theorem of Jordan*, Pacific. J. Math. **4** (1954), 219–226.
- [6] M.E. Harris and C. Hering: *On the smallest degrees of projective representations of the groups  $PSL(n, q)$* , Canad. J. Math. **28** (1971), 90–102.
- [7] B. Huppert: *Endliche Gruppen I*, Springer, Berlin-Heidelberg-New York, 1967.
- [8] C. Jordan: *Sur la limite de transitivité des groupes non alternés*, Bull. Soc. Math. France **1** (1875), 40–71.
- [9] I. Miyamoto: *Multiply transitive permutation groups and odd primes*, Osaka J. Math. **11** (1974), 9–13.
- [10] H. Nagao: *On multiply transitive groups*, Nagoya Math. J. **27** (1966), 15–19.
- [11] H. Nagao: *On multiply transitive groups V*, J. Algebra **9** (1968), 240–248.
- [12] T. Nakayama: *On some modular properties of irreducible representations of a symmetric group II*, Japanese J. Math. **17** (1940), 411–423.
- [13] I. Schur: *Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene lineare Substitutionen*, Crelle J. **139** (1911), 155–250.
- [14] H. Wielandt: *Finite Permutation Groups*, Academic Press, New York and London, 1964.
- [15] K. Yamazaki: *On projective representations and ring extensions of finite groups*, J. Fac. Sci. Univ. Tokyo **10** (1963–4), 147–195.
- [16] H. Zassenhaus: *Kennzeichnung endlicher linearer Gruppen als Permutationsgruppen*, Abh. Math. Sem. Univ. Hamburg, **11** (1934), 17–40.

