

Title	Studies on the Nickel- and Platinum-Catalyzed Borylation of Aromatic Carbon-Hydrogen Bonds
Author(s)	古川, 嵩幸
Citation	大阪大学, 2017, 博士論文
Version Type	VoR
URL	https://doi.org/10.18910/61784
rights	
Note	

### Osaka University Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

Osaka University

**Doctoral Dissertation** 

Studies on the Nickel- and Platinum-Catalyzed Borylation of Aromatic Carbon–Hydrogen Bonds

Takayuki Furukawa

January 2017

**Department of Applied Chemistry,** 

Graduate School of Engineering,

**Osaka University** 

#### **Preface and Acknowledgement**

The research presented in this thesis was carried out under the direction of Professor Naoto Chatani of the Department of Applied Chemistry, Faculty of Engineering, Osaka University from April 2011 to March 2017. The thesis is concerned with the nickel- and platinum-catalyzed borylation of aromatic carbon-hydrogen bonds.

This thesis could not have been finished without the heartfelt help and support from numerous people. I wish to express my sincerest appreciation to all of those people who I spent the time with during the last 6 years in the Chatani group.

I express my utmost gratitude to Professor Naoto Chatani for his guidance and support throughout this work. He always gave me fruitful advice, support and encouragement for anything I attempted to do. He has a great enthusiasm for chemistry and deserves respect from his personality.

I would like to express my deepest gratitude to Professor Mamoru Tobisu for his valuable advice and continuous support. The advice derived from his logical and creative mind always encouraged me to work out issues and provided me new points of view to consider problems. Thanks to his respectable character and personality, there was always pleasant atmosphere to study in his group. It was a great delight for me to experience exciting chemistry and satisfying days under his supervision.

I deeply appreciate Professor Yoshiya Fukumoto for his instructive and expert advice based on his extensive and profound understanding in chemistry.

I wish to express my special appreciation to the secretaries in our laboratory, including Ms. Mayuko Nakamura, Ms. Yoshimi Shinomiya and Ms. Junko Ohmagari for their generous assistance.

I would also like to express my appreciation to the past and present members of the Chatani group. The marvelous and talented senior members: Dr. Takeshi Uemura, Dr. Yusuke Ano (*Ano-san*), Dr. Isao Hyodo, and Dr. Masahiro Onoe (The first captain of the Chatani-lab horse-racing club), influenced me a lot through their enthusiastic attitude regarding chemistry and offered much to emulate. I wish to thank the excellent and humorous senior members; Dr. Katsuaki Baba, Dr. Hirotaka Kinuta, Dr. Yoshinori Aihara (Fun at drinking party) and Dr. Keisuke Nakamura (The initial member of the Chatani-lab video-game club). I could spent fruitful and happy times through discussing chemistry, chatting and drinking with them even when my research was not going well. I also appreciate my respected seniors, including Mr. Motohiro Shiratani, Ms. Nao Hasegawa, Mr. Masato Daijo, Ms. Tian Xu, Mr. Akihiro Ohmae, Mr. Junya Hasegawa and Mr. Yasunori Soga.

I also wish to express my special thanks to my classmates: Dr. Kaname Shibata (*Shibatch*), Ms. Miki Iyanaga, Mr. Hiroto Shimizu and Ms. Ayaka Yasutome. They were highly motivated and they were my helpful colleagues to be able to encourage each other.

I would also wish to express my gratitude to the kind and motivated junior members of the Chatani group, including Ms. Aya Tashiro, Mr. Masaya Hirano, Mr. Toshifumi Morioka (a former member of the 3<sup>rd</sup> floor), Ms. Ayana Yokota, Mr. Motonobu Kamiya, Mr. Tsuyoshi Takahira, Mr. Yuto Tamura, Ms. Moe Noguchi, Mr. Yoshihiro Masuya (the 2<sup>nd</sup> generation of the video game club), Mr. Takuya Igarashi (*Igaiga*), Mr. Teruhiko Kubo, Mr. Jiangning Zhao (*Cho-kun*), Ms. Natsuki Okazaki (*Zaki*), Mr. Yuta Seo, Mr. Kosuke Yasui (*Yasui-kun*), Mr. Takuma Yamaguchi (*Gussan*), Ms. Mao Yamaguchi, Mr. Sodai Yamada, Mr. Yasuaki Iyori, Mr. Atsushi Obata, Ms. Satoko Natsui (*Natzy*), Mr. Akihiro Nishizawa (*Zawa*), Mr. Kosuke Yanagisawa, Mr. Yuki Amano, Ms. Akane Sasagawa, Mr. Akira Haito, Mr. Masaya Higashino (*Ninja*) and Mr. Nao Matsubara.

Furthermore, I express my appreciation to Dr. Guy Rouquet, Mr. Ho Jordan Sun, Dr. Luis Carlos Misal Castro, Mr. Jendrik Wuelbern, Dr. Akimichi Ohtsuki, Mr. Mikhail Konev, Dr. Yadagiri Kommagalla and Mr. Daichi Koseki who worked in the Chatani group as visiting scientists or postdoctoral fellows.

Professor Chatani provided me with an opportunity to be engaged in total synthesis in the research group of Professor Dirk Trauner at Ludwig-Maximilians-Universität München (LMU) in Germany from July to December 2015. It was a great experience in my life and what I learned through this opportunity represents an important contributions to me as a researcher. I would like to express my sincere gratitude to Professor Dirk Trauner for accepting me as a visiting student. In addition, I deeply appreciate Mr. Felix Hartrampf, who was my dear colleague during the research stay and taught me a lot of techniques and extended my knowledge of total synthesis, in addition to the language and culture of Germany.

I am grateful for and wish to acknowledge a JSPS Research Fellowship for Young Scientists.

At last but not least, I would like to express my thanks to my parents Mr. Yoshikazu Furukawa and Ms. Mayumi Furukawa for their perpetual support.

Suita, Osaka January 2017

Takayuki Furukawa

#### Contents

#### **General Introduction**

#### References

# Chapter 1 Nickel-Catalyzed Borylation of Aromatic C-H Bonds 1.1 Introduction 1.2 Results and Discussion

- 1.3 Conclusion
- 1.4 Experimental Section
- 1.5 References and Notes

#### Chapter 2 Platinum-Catalyzed Borylation of Aromatic C-H Bonds

- 2.1 Introduction
- 2.2 Results and Discussion
- 2.3 Conclusion
- 2.4 Experimental Section
- 2.5 References and Notes

#### Conclusion

#### List of Publications / Supplementary List of Publications

#### **General Introduction**

Catalytic C–H bond functionalization is one of the simplest and most straightforward methods for the synthesis of complex organic molecules from readily available hydrocarbons and their derivatives.<sup>1</sup> The catalytic transformation of C-H bonds omits the pre-activation steps prior to the introduction of the targeted functionality and therefore makes the process more efficient than traditional synthetic methods. One of the most reliable methods for C–H functionalization involves the use of a substrate containing a preinstalled coordinating group, which can direct a metal catalyst to react with a targeted C–H bond in a regioselective manner (Figure 1a, i.e., directed methodology). This method is very useful for the activation of C-H bonds that are located in close proximity to a directing group, which is normally the *ortho* position to the directing group in an aromatic system. C-H functionalization methods that do not rely on a directing group (i.e., undirected C-H functionalization) are highly desired from the view point that it could be used to achieve unique patterns of regioselectivity that would be otherwise unattainable using directed methods. The regioselectivity in undirected C-H functionalization is often determined by the electronic and steric properties of the substrate (Figure 1b). **Figure 1.** Strategy for Regioselective C-H Functionalization



Among the undirected functionalization of C-H bonds, C-H borylation is currently recognized as one of the most powerful and reliable methods (eq. 1).<sup>2,3</sup> The catalytic borylation of C-H bonds allows for the direct introduction of a synthetically useful boron functionality from readily available starting materials.<sup>3</sup> Among the variety of catalysts that have been used in C-H borylation, the Ir/dtbpy-based complexes developed by Miyaura, Ishiyama and Hartwig are capable of catalyzing the borylation of aromatic C-H bonds with a high degree of efficiency and selectivity in an undirected manner (Figure 2a).<sup>2</sup> The regioselectivity of the C-H borylation reaction is normally controlled by the steric



environment surrounding the reacting C-H bonds. For example, the C-H borylation of 1,3-disubstituted benzene substrates occurs exclusively at the 5-position (Figure 2a, right). The outstanding catalytic activity and selectivity of iridium systems has resulted in their practical application in various areas in materials science and in the synthesis of fine chemicals. Since the first report on Ir/dtbpy catalysts, considerable progress has been made in the field of C-H borylation by modifying the properties of the ligands to the iridium complex. These attempts have resulted in considerable success in terms of improving catalytic efficiency<sup>4</sup> and establishing novel levels of regioselectivity that go beyond simple steric control.<sup>5</sup>





Another promising approach for the development of new C-H borylation catalysts involves the replacement of iridium with other metals. The catalytic activity of rhodium for C-H borylation reactions has long been established (Figure 2b).<sup>6</sup> Cobalt-based catalysts have been studied extensively by Chirik, who demonstrated that well-defined pincer cobalt complexes could effectively promote C-H borylation reactions (Figure 2c).<sup>7</sup> Iron complexes have also been shown to be suitable metal centers

for C-H borylation reactions when they are combined with oxidants or performed under photoirradiation conditions (Figure 2d).<sup>8</sup>

Compared to the development of catalytic reactions using group 9 metals such as Ir, Rh and Co for C-H borylation, the potential reactivity of group 10 metals for these reactions represents a less developed area. Except for the seminal work by Miyaura using a Pd/C catalyst for benzylic borylation,<sup>9</sup> there were no other examples of group 10 metal-catalyzed C-H borylations when the author started the research.<sup>10</sup> Varying the central metal could provide opportunities for the development of new types of catalysts for C–H borylation reactions with new reactivities and/or selectivity profiles.

The prime objective of this research was to investigate the catalytic activity of nickel and platinum metals in C-H borylation reactions and to explore their unique properties (eq. 2). This thesis consists of the following two chapters.



In chapter 1, nickel-catalyzed borylation of aromatic C-H bonds is discussed. This is the first demonstration of the catalytic activity of a nickel complex in the borylation of aromatic C-H bonds. This method can be applied to the borylation of indoles on a gram scale, which demonstrates that earth abundant nickel is a promising candidate as a substitute for a precious metal catalyst such as iridium and rhodium. The choice of *N*-heterocyclic carbene (NHC) as a ligand is important in achieving nickel catalysis, which represents a rare example of a nickel-catalyzed C-H bond activation process in the absence of a directing group.

In chapter 2, the platinum-catalyzed borylation of C-H bonds is discussed. NHC-ligated platinum catalysts have been found to be efficient catalysts for the borylation of aromatic  $C(sp^2)$ –H bonds. This is the first example of a catalytic C-H borylation in which a platinum catalyst is used. The most remarkable feature of these Pt catalysts is their lack of sensitivity with respect to the degree of steric hindrance around the C–H bonds that participate in the borylation reaction. These Pt catalysts allow for the synthesis of sterically congested 2,6-disubstituted phenylboronic esters, which are otherwise difficult to synthesize using existing C–H borylation methods. Furthermore, platinum catalysis allows for the regioselective borylation of the C–H bonds *ortho* to fluorine substituents in fluoroarene systems. Preliminary mechanistic studies and work directed at synthetic applications of this platinum catalyzed C–H borylation process are described.

#### References

- Selected reviews: a) T. W. Lyons, M. S. Sanford, *Chem. Rev.* 2010, *110*, 1147. b) J. Yamaguchi, A. D. Yamaguchi, K. Itami, *Angew. Chem. Int., Ed.* 2012, *51*, 8960. c) G. Rouquet, N. Chatani, *Angew. Chem., Int. Ed.* 2013, *52*, 11726. d) F. Kakiuchi, T. Kochi, S. Murai, *Synlett* 2014, *25*, 2390. e) M. Miura, T. Satoh, K. Hirano, *Bull. Chem. Soc. Jpn.* 2014, *87*, 751. f) Z. Chen, B. Wang, J. Zhang, W. Yu, Z. Liu, Y. Zhang, *Org. Chem. Front.* 2015, *2*, 1107. g) B. Su, Z.-C. Cao, Z.-J. Shi, *Acc. Chem. Res.* 2015, *48*, 886. h) T. Gensch, M. N. Hopkinson, F. Glorius, J. Wencel-Delord, *Chem. Soc. Rev.* 2016, *45*, 2900. i) R.-Y. Zhu, M. E. Farmer, Y.-Q. Chen, J.-Q. Yu, *Angew. Chem., Int. Ed.* 2016, *55*, 10578. j) L. C. M. Castro, N. Chatani, *Chem. Lett.* 2015, *44*, 410.
- Seminal works: a) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaura, N. R. Anastasi, J. F. Hartwig, J. Am. Chem. Soc. 2002, 124, 390. b) J.-Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka, M. R. Smith, Science. 2002, 295, 305.
- (3) Reviews: a) T. Ishiyama, N. Miyaura, J. Organomet. Chem. 2003, 680, 3. b) I. A. I. Mkhalid, J. H. Barnard, T. B. Marder, J. M. Murphy, J. F. Hartwig, Chem. Rev. 2010, 110, 890. c) J. F. Hartwig, Chem. Soc. Rev. 2011, 40, 1992. d) J. F. Hartwig, Acc. Chem. Res. 2012, 45, 864.
- (4) Selected examples: a) S. Kawamorita, H. Ohmiya, K. Hara, A. Fukuoka, M. Sawamura, J. Am. Chem. Soc. 2009, 131, 5058. b) S. M. Preshlock, B. Ghaffari, P. E. Maligres, S. W. Krska, R. E. Maleczka, M. R. Smith, J. Am. Chem. Soc. 2013, 135, 7572. c) G. Wang, L. Xu, P. Li, J. Am. Chem. Soc. 2015, 137, 8058. d) L. P. Press, A. J. Kosanovich, B. J. McCulloch, O. V. Ozerov, J. Am. Chem. Soc. 2016, 138, 9487.
- (5) Selected examples: a) Y. Kuninobu, H. Ida, M. Nishi, M. Kanai, *Nat. Chem.* 2015, 7, 712. b) Y. Saito, Y. Segawa, K. Itami, *J. Am. Chem. Soc.* 2015, *137*, 5193. c) R. Bisht, B. Chattopadhyay, *J. Am. Chem. Soc.* 2016, *138*, 84. d) H. J. Davis, M. T. Mihai, R. J. Phipps, *J. Am. Chem. Soc.* 2016, *138*, 12759.
- (6) Seminal works: a) H. Chen, S. Schlecht, T. C. Semple, J. F. Hartwig, *Science*. 2000, 287, 1995.
  b) J.-Y. Cho, C. N. Iverson, M. R. Smith, *J. Am. Chem. Soc.* 2000, *122*, 12868.
- (7) A seminal work: J. V. Obligacion, S. P. Semproni, P. J. Chirik, J. Am. Chem. Soc. 2014, 136, 4133.
- (8) a) K. M. Waltz, X. He, C. Muhoro, J. F. Hartwig, J. Am. Chem. Soc. 1995, 117, 11357. b) G. Yan, Y. Jiang, C. Kuang, S. Wang, H. Liu, Y. Zhang, J. Wang, Chem. Commun. 2010, 46, 3170.
  c) T. Hatanaka, Y. Ohki, K. Tatsumi, Chem. Asian J. 2010, 5, 1657. d) T. J. Mazzacano, N. P. Mankad, J. Am. Chem. Soc. 2013, 135, 17258. e) T. Dombray, C. G. Werncke, S. Jiang, M. Grellier, L. Vendier, S. Bontemps, J.-B. Sortais, S. Sabo-Etienne, C. Darcel, J. Am. Chem. Soc. 2015, 137, 4062.
- (9) T. Ishiyama, K. Ishida, J. Takagi, N. Miyaura, *Chem. Lett.* **2001**, *2*, 1082.
- (10) Several examples of Pd-catalyzed borylation using directing groups were reported. See the review: A. Ros, R. Fernández, J. M. Lassaletta, *Chem. Soc. Rev.* **2014**, *43*, 3229.

#### Chapter 1

#### Nickel-Catalyzed Borylation of Aromatic C-H Bonds

#### **1.1 Introduction**

As described in the general introduction, the catalytic borylation of arenes is arguably the method of paramount importance, since it allows the introduction of a synthetically useful boron functionality directly at the C-H bond of an arene without the need for any directing groups. One remaining challenge in the catalytic C-H borylation methodology is the development of base metal catalysts, driven by recent concerns regarding the limited availability of precious metals. A pioneering work by Hartwig in 1995 demonstrated the potential activity of base metal complexes such as Fe and Mn in C-H borylation reactions.<sup>1</sup> Catalytic variants were achieved only recently, as already briefly mentioned in the general introduction. Ohki and Tatsumi reported on the iron-catalyzed borylation of furans and thiophenes in the presence of a hydrogen acceptor,<sup>2</sup> while Kuang and Wang demonstrated that Fe<sub>2</sub>O<sub>3</sub> nanoparticles can also catalyze the borylation of arenes in the presence of more than a stoichiometric amount of *tert*-butyl peroxide.<sup>3</sup> A mechanistically unique C-H borylation using a Cu/Fe bimetallic catalyst under photochemical conditions was developed by Mankad.<sup>4</sup> The use of a bidentate phosphineligated iron complex in catalytic borylation reactions under photochemical conditions was reported by the group of Bontemps, Sortais, Sabo-Etienne and Darcel.<sup>5</sup> As a base-metal variant of a group 9 metal, Chirik reported on a cobalt-catalyzed C-H borylation that is applicable to a broad range of arenes and heteroarenes and subsequent studies revealed that cobalt has promising catalytic activity for C-H borylations.<sup>6</sup> Despite these advances, the full potential of base metals in promoting C-H borylation in a broader sense, to promote C-H activation, has still not yet been explored, especially in the case of metals other than iron<sup>7</sup> and cobalt.<sup>8</sup>

The use of a nickel complex in C-H activation reactions has long been known since the pioneering work in 1963, in which the cyclometallation of stoichiometric cyclopentadienyl nickel with azobenzene was examined.<sup>9</sup> Despite the early finding, the development of catalytic C-H functionalizations using nickel has only recently progressed. Chatani extensively studied the bidentate directing group strategy for effective nickel-catalyzed C-H functionalization.<sup>10</sup> Concerning substrates that do not have directing groups, the nickel catalyst can function effectively only for substrates that contain acidic C-H bonds such as those located on the highly fluorinated arenes, azoles, and activated pyridine derivatives.<sup>11</sup> Although co-catalytic systems using nickel and aluminum Lewis acids have

recently broaden the scope of such reactions to aromatic rings bearing carbonyl functionalities,<sup>12</sup> the nickel-catalyzed C-H functionalization of the simple benzene, to the best of our knowledge, is limited to only one example, involving the oxidation of benzene to phenol.<sup>13</sup> The author therefore investigated the C-H borylation of simple arenes using a nickel catalyst.

#### **1.2 Results and Discussion**

The author initially examined the nickel-catalyzed reaction of HBpin with benzene in the presence of a variety of ligands (Table 1). Phosphine ligand such as  $PCy_3$  did not lead to the formation of the desired borylated product **1** (Entry 2). In contrast, the use of IMes, an *N*-heterocyclic carbene (NHC) bearing a mesityl group, was found to generate **1** in 51% yield (Entry 3). Replacing the mesityl group with either the larger 2,6-<sup>*i*</sup>Pr<sub>2</sub>C<sub>6</sub>H<sub>2</sub> group (i.e., IPr) or the smaller methyl group (i.e., IMe) decreased the yield of **1** (Entries 4 and 5).

HBpin +	solvent	Ni(cod) <sub>2</sub> (3 mol % Ligand (3 mol %) NaO <sup>t</sup> Bu (6 mol %) 80 °C, 20 h	) Bpin 1
Entry	Ligand		Yield (%) <sup>b</sup>
1 2° 3 4 5 6 7 8 9 10 11 12	none PCy <sub>3</sub> IMes·HCl ( IPr·HCl (2,' IMe·HCl (R I <sup>/</sup> Pr·HBF <sub>4</sub> ( I <sup>/</sup> Bu·HCl (R ICy·HCl (R I(1-Ad)·HC I(2-Ad)·HC SICy·HCl BICy·HCl	$R = 2,4,6-Me_{3}C_{6}H_{2})$ $6^{-i}Pr_{2}C_{6}H_{3})$ $R = {}^{i}Pr)$ $R = {}^{i}Bu)$ $= cyclohexyl)$ $I (R = 1-adamantyl)$ $I (R = 2-adamantyl)$	5 0 51 16 22 33 50 75 (71) <sup>d</sup> 36 trace 2 69
Ligand: R-N CI	\+ N∼R Cị	y∽N <sub>∽</sub> N <sup>+</sup> Cy CĪ SICy·HCI	Cy-N_N-Cy CI BICy-HCI

Table 1. Nickel-Catalyzed Borylation of Benzene: Ligand Screening<sup>a</sup>

<sup>*a*</sup> Reaction conditions: HBpin (1.2 mmol), Ni(cod)<sub>2</sub> (0.036 mmol), ligand (0.036 mmol), NaO<sup>*t*</sup>Bu (0.072 mmol) and benzene (1.0 mL) in a screw-capped vial under N<sub>2</sub> at 80 °C for 20 h. <sup>*b*</sup> GC yield based on HBpin. <sup>*c*</sup> NaO<sup>*t*</sup>Bu was not added <sup>*d*</sup> Isolated yield.

Further experimentation with the substituent on the nitrogen of the NHC ligand revealed that the cyclohexyl-substituted derivative (i.e., ICy) was the most effective ligand among those tested, with **1** being formed in 71% isolated yield (Entry 8). Whereas the use of a saturated analog of ICy (i.e., SICy) led to a complete loss of catalytic activity (Entry 11), the benzohomolog of ICy (i.e., BICy) promoted the borylation of benzene with comparable efficiency (Entry 12).

 $B_2pin_2$  was also found to be a suitable boron source for this nickel-catalyzed borylation, forming **1** in a 125% yield based on  $B_2pin_2$  (Eq 1).

$$B_{2}pin_{2} + \bigcup_{solvent} Ni(cod)_{2} (5 mol \%)$$

$$B_{2}pin_{2} + \bigcup_{solvent} NaO^{t}Bu (15 mol \%)$$

$$B_{2}pin_{2} + \bigcup_{solvent} Bpin (1)$$

The Ni(cod)<sub>2</sub>/ICy-catalyzed borylation of substituted benzene derivatives was also investigated (Table 2). When toluene was used as the substrate, the borylation proceeded efficiently to give a mixture of regioisomers (Entry 1). The regioselectivity of the borylation was found to be primarily controlled by steric effects, forming the meta and para isomers as the major products, a result similar to those obtained using Ir<sup>14</sup>, Fe<sup>4</sup> or Co<sup>6a</sup> catalysts. The difference between the present work and these previously reported catalysts is that the borylation in the current study also occurred at the benzylic position. The observed product distribution was similar to that obtained using a rhodium-based catalyst<sup>15</sup> although the rhodium system allows the benzylic isomer to be formed as the major product. The borylation of ethylbenzene under these nickel-catalyzed conditions also proceeded to deliver four isomers (Entry 2). Disubstituted benzenes were generally much less reactive than mono-substituted benzenes under the current reaction conditions (Entry 3). The use of anisole as the substrate resulted in borylation products, with the ortho-borylated isomer as the major product (Entry 4). This regioselectivity was in sharp contrast to that observed in homogeneous Ir,<sup>3</sup> Fe<sup>4</sup> and Co<sup>6a</sup> systems, wherein the major products were the meta and para isomers. This result indicated that a methoxy group can serve as an ortho-directing group in the current nickel-catalyzed borylation reaction. Borylative cleavage of the C(aryl)-OMe bond<sup>16</sup> in anisole was also observed, albeit in low yield (7%). Unlike the Hartwig's hydroarylation system,<sup>17</sup> 1,3-bis(trifluoromethyl)benzene was not a good substrate in the present reaction (<10% yield). Halogen-substituted benzenes such as chloro, bromo and fluorobenzenes were found to be inapplicable since C-X bonds can react with the nickel species under these conditions.<sup>18</sup>

**Table 2** Nickel-Catalyzed Borylation of Various Arenes<sup>a</sup>



<sup>*a*</sup> Reaction conditions: HBpin (1.2 mmol), Ni(cod)<sub>2</sub> (0.035 mmol), ICy·HCl (0.035 mmol), NaO'Bu (0.070 mmol) and arene (1.0 mL) in a screw-capped vial under N<sub>2</sub> at 100 °C for 20 h. <sup>*b*</sup> Isolated yield based on HBpin. <sup>*c*</sup> At 80 °C. <sup>*d*</sup> *N*-Methylpyrrole (2.0 mmol) in methylcyclohexane (1.0 mL).

Heteroarenes were also examined as potential substrates for the nickel-catalyzed borylation reactions. Whereas furan, thiophene and pyridine could not be borylated under these conditions, pyrrole was found to be exceptionally reactive during this borylation, forming the 2-borylated product in a regioselective manner (Entry 5). Notably, the product was obtained in good yield even when the amount of pyrrole was reduced to 1.7 equivalents relative to HBpin.

The evident high reactivity of pyrrole prompted the author to examine indole derivatives in this nickel-catalyzed borylation reaction (Table 3). As expected, the borylation occurred efficiently using indole as the limiting agent and also took place regioselectively at the 2-position. In addition to *N*-methylindole (Entry 1), a bulkier *N*-butyl-substituted indole successfully underwent borylation without any decrease in the regioselectivity of the reaction (Entry 2). An *N*,*O*-acetal moiety remained intact,

Table 3 Nickel-Catalyzed Borylation of Indoles<sup>a</sup>



<sup>*a*</sup> Reaction conditions: indole (0.80 mmol), HBpin (1.2 mmol), Ni(cod)<sub>2</sub> (0.040 mmol), ICy·HCl (0.040 mmol), NaO<sup>*t*</sup>Bu (0.080 mmol), and methylcyclohexane (1.0 mL) in a screw-capped vial under N<sub>2</sub> at 80 °C for 16 h. <sup>*b*</sup> HBpin (1.2 equiv) was used. <sup>*c*</sup> Indole (0.72 mmol) and HBpin (0.98 mmol, 1.4 equiv) were used.

serving as a suitable protecting group in this reaction (Entry 3). In addition, *N*-benzyl groups were also tolerated, leaving the aryl group free from borylation (Entries 4 and 5). Installation of a boryl group at the sterically hindered 2-position of 3-substituted indoles was also possible (Entry 6). The reactivity and selectivity of this borylation was unaffected by the introduction of a methoxy group to the indole framework (Entry 7). Although a C(aryl)-F bond can be activated by a low valent nickel catalyst,<sup>18</sup>

fluoroindole underwent borylation with its fluoride moiety remaining intact (Entry 8). Azaindole also served as a good substrate for this borylation, generating the corresponding 2-borylated product (Entry 9).

Although Ni(cod)<sub>2</sub> was routinely used as the catalyst for this borylation during our exploratory studies, Ni(OAc)<sub>2</sub> was found to serve as a more useful catalyst precursor. A protocol using Ni(OAc)<sub>2</sub> allowed the successful implementation of this borylation on the gram-scale, highlighting the practical utility of the nickel-catalyzed system (eq. 2).



To obtain further insights into the mechanistic aspects of the nickel-catalyzed borylation, some preliminary mechanistic experiments were performed. Comparison of the initial rates of the borylation reactions of benzene and benzene- $d_6$  determined that  $k_{\rm H}/k_{\rm D}$  was 2.1 under these conditions, suggesting that cleavage of the C-H bond was involved in the turnover-limiting step. The nickel-catalyzed borylation of benzene or indole was also found to be completely inhibited when the reaction was attempted in the presence of an excess of mercury. In addition, filtration tests indicated that the liquid phase of the catalytic mixture did not contain catalytically active species (see experimental section for details). The currently available data indicate that the borylation examined here is likely mediated by a heterogeneous nickel species, although more detailed studies will be required for a definitive understanding of the nature of the catalysis.<sup>18</sup> It is worth noting that the use of slight stoichiometric excess of NaO'Bu relative to NHC+HCl was important for an efficient reaction, although the role of NaO'Bu was initially thought to involve the generation of a free carbene ligand from the corresponding imidazolium salt. When the reaction was performed in the absence of the NaO'Bu using free IMes, the yield dramatically decreased compared to the results when an IMes ligand generated from IMes•HCl and NaO'Bu was used (see experimental section for details). Additionally, a seminal report on the KO'Bu-catalyzed silvlation of heteroarenes<sup>19,20</sup> motivated the author to examine the borylation of indole in the presence of only NaO'Bu. No borylation product was detected under such conditions, indicating that the nickel species plays an essential role in this borylation reaction. It should also be noted that at the same time of our publication, Itami et al. independently reported nickel-catalyzed C-H borylation reaction using phosphine ligands and a catalytic amount of CsF.<sup>21</sup>

#### **1.3 Conclusion**

In summary, the findings reported here show, for the first time, that the direct borylation of arenes can be mediated by a nickel-based catalyst. Indoles are particularly reactive with this catalyst system, affording the corresponding 2-borylated products. The successful use of nickel is also noteworthy because nickel is rarely employed as a catalyst for the C-H bond activation process in the absence of a directing group.<sup>11-13</sup> The heterogeneous nature of the active species generated under the current conditions could be important in explaining the unprecedented reactivity of nickel,<sup>22</sup> although further investigations will be required for a complete understanding of the mechanism of this borylation.

#### **1.4 Experimental Section**

#### **1.4.1. General Information**

<sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a JEOL ECS-400 spectrometer (JEOL, Tokyo, Japan) or VARIAN UNITY INOVA-600 spectrometer in CDCl<sub>3</sub> with tetramethylsilane as an internal reference standard. Some NMR analysis were conducted using  $C_6D_6$  solvent and a residual  $C_6H_6$  peak was set as an internal reference standard ( $\delta = 7.15$  ppm). NMR data have been reported as follows: chemical shift ( $\delta$ ) in ppm, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, and m = multiplet), coupling constant (*J*) in Hz, and integration. NOE experiments were conducted with Delta noe\_1d\_dpfgse pulse sequence and detail experimental parameter was listed in the spectroscopic data section for each compound. Infrared spectra (IR) were obtained on a JASCO FT/IR-4000 spectrometer, and the absorptions have been reported in reciprocal centimeters with the following relative intensities: s (strong), m (medium), or w (weak). Mass spectra were obtained on a Shimadzu GCMS-QP 5000 or GCMS-QP 2010 instrument with an ionization voltage of 70 eV. High resolution mass spectra (HRMS) were obtained on a JEOL JMS-DX303(EI) and a Bruker micrOTOF II (APCI). Analytical gas chromatography (GC) was carried out on Shimadzu GC-2014, equipped with a flame ionization detector. Melting points were determined using a Stanford Research Systems OptiMelt. Column chromatography was performed over SiO<sub>2</sub> (Silycycle Silica Flash F60 (230-400 mesh).

#### 1.4.2. Materials

Ni(cod)<sub>2</sub> (Strem), Ni(OAc)<sub>2</sub>•4H<sub>2</sub>O (Wako), NaO'Bu (TCI), HBpin (TCI), B<sub>2</sub>pin<sub>2</sub> (TCI), PCy<sub>3</sub> (Aldrich), IMes•HCl (TCI), IPr•HCl (TCI), IMe•HCl (TCI), I<sup>'</sup>Pr•HBF<sub>4</sub> (TCI), I(1-Ad)•HCl (TCI), BICy•HCl (Aldrich) were obtained from commercial suppliers and used as received. I'Bu•HCl,<sup>23</sup> ICy•HCl,<sup>24</sup> SICy•HCl<sup>25</sup> and I(2-Ad)•HCl<sup>26</sup> were synthesized according to the literature procedure. Methylcyclohexane (TCI), *N*-methylpyrrole (TCI), anisole (Nacalai), and *o*-xylene (TCI) were purified by distillation over CaH<sub>2</sub> prior use. 1,3-Bis(trifluoromethyl)benzene (Wako), *N*-methylindole (TCI), anhydrous benzene (Wako), benzene-*d*<sub>6</sub> (Wako), Hg (Wako) were obtained from commercial suppliers and used as received. 1-Butyl-1*H*-indole, 1-benzyl-1*H*-indole, 1-(4-methoxybenzyl)-1*H*-indole, 1,3dimethyl-1*H*-indole, 1-(methoxymethyl)-1*H*-indole, 5-fluoro-1methyl-1*H*-indole, 1-methyl-7-azaindole, were synthesized by the reaction of the corresponding indole with alkyl halide according to literature procedures.<sup>27</sup>

**1.4.3.** General Procedures for Nickel-Catalyzed Borylation of Arenes and Indoles Method A: Procedure for the Nickel-Catalyzed Borylation of Benzene with Pinacolborane (Entry 8, Table 1)



In a glovebox, Ni(cod)<sub>2</sub> (9.9 mg, 0.035 mmol), ICy•HCl (9.7 mg, 0.036 mmol), NaO'Bu (6.4 mg, 0.07 mmol) and HBpin (159 mg, 1.2 mmol) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. Benzene (1.0 mL) was then added, and the cap was applied to seal the vial. The vial was stirred at 80 °C over a hot plate for 20 h and the resulting mixture was filtered through silica gel (eluting with 10 mL of hexane/AcOEt = 5/1). The filtrate was analyzed by gas chromatography (GC) using dodecane as an internal standard (75% GC yield). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/AcOEt = 20/1) to give 4,4,5,5-tetramethyl-2-phenyl-1,3,2-dioxaborolane as a white solid (179 mg, 71% based on HBpin).

Method B: Procedure for the Nickel-Catalyzed Borylation of *N*-Methylindole with Pinacolborane (Entry 1, Table 3)



In a glovebox, Ni(cod)<sub>2</sub> (12.4 mg, 0.045 mmol), ICy•HCl (11.7 mg, 0.044 mmol), NaO'Bu (7.1 mg, 0.074 mmol), 1-methylindole (109 mg, 0.82 mmol) and HBpin (160 mg, 1.25 mmol) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. Methylcyclohexane (1.0 mL) was then added, and the cap was applied. The vial was stirred at 80 °C over a hot plate for 20 h. The resulting mixture was filtered through silica gel (eluting with 10 mL of hexane/AcOEt = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/AcOEt = 20/1) to give 1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole as a white solid (165 mg, 78% based on 1-methylindole).

#### A Procedure for the Gram Scale Synthesis of 2-Borylated Indole without a Glovebox (Scheme 2)



To a two-necked flask equipped with a rubber septum and a three-way stopcock connected to a vacuum line and a  $N_2$  line, Ni(OAc)<sub>2</sub>•4H<sub>2</sub>O (76 mg, 0.31 mmol) was added. The flask was then heated by a

heat gun under vacuum until Ni(OAc)<sub>2</sub>•4H<sub>2</sub>O (initially light blue) completely changed color to a greenish yellow (ca. 1.5 min.). After the flask was cooled and refilled with N<sub>2</sub>, ICy•HCl (84.4 mg, 0.31 mmol) and NaO'Bu (50.8 mg, 0.53 mmol) were added under a N<sub>2</sub> flow. Following the addition of these solid reagents, the vessel was again placed under vacuum and its volume subsequently purged with N<sub>2</sub> three times. Following this, *N*-methylindole (1.16 g, 8.8 mmol), HBpin (2.44 g, 19.1 mmol) and methylcyclohexane (25 mL) were added via syringe through the rubber septum. The reaction mixture was heated at 80 °C for 14 h in an oil bath, during which time the loss of the initial reagent *N*-methylindole was monitored by GC. The reaction mixture was then poured onto silica gel and eluted with 100 mL of a hexane/AcOEt (10/1) solution. The filtrate was concentrated in vacuo and analytically pure 1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole was obtained quantitatively as a white solid (2.35 g, quant.).

#### **1.4.4. Spectroscopic Data for the Products**

#### 4,4,5,5-Tetramethyl-2-phenyl-1,3,2-dioxaborolane (entry 8, Table 1) [CAS: 24388-23-6].

Bpin Rf 0.60 (hexane/EtOAc = 5/1). White solid (179mg, 71%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.34 (s, 12H), 7.37 (dt, *J* = 0.8, 7.2 Hz, 2H), 7.46 (tt, *J* = 1.6 Hz, 6.8 Hz, 1H). 7.81 (dd, *J* = 1.2 Hz, 7.2 Hz, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.8, 83.7, 127.6, 131.2, 134.7. HRMS (EI): Calcd for C<sub>12</sub>H<sub>17</sub>BO<sub>2</sub> 204.1322, Found 204.1321.

#### Borylation of toluene (Entry 1, Table 2).



Method A was followed except that the reaction was conducted with HBpin (149 mg, 1.16 mmol) in toluene (1.0 mL) at 100 °C. After purification by flash column chromatography over silica gel (eluting with hexane/AcOEt = 20/1), a mixture of four isomers was obtained (163 mg, 65%, >99% purity by GC). GC/MS analysis revealed the formation of four isomers of the borylated products, all of which had an m/z of 218 (M<sup>+</sup>). The identity and ratio of each of the four isomers was determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>28</sup> The resonances specific to each isomer are as follows: 7.75 ppm (d, J = 7.6 Hz, 1H, ortho isomer, H<sub>a</sub>); 7.60-7.64 ppm (m, 2H, meta isomer, H<sub>b</sub> and H<sub>c</sub>), 7.71 ppm (d, J = 6.0 Hz, 1H, para isomer, H<sub>d</sub>); 2.29 ppm (s, 2H, benzylic isomer, H<sub>e</sub>). (See below.)



Method A was followed except that the reaction was conducted with HBpin (160 mg, 1.25 mmol) in ethylbenzene (1.0 mL) at 100 °C. After purification by flash column chromatography over silica gel (eluting with hexane/AcOEt = 20/1), a mixture of four isomers was obtained (145 mg, 50%, >99% purity by GC). GC/MS analysis revealed the formation of four isomers of the borylated products, all of which had an m/z of 232 (M<sup>+</sup>). The identities and proportions of the meta, para and benzylic isomers were determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>28b,29</sup> Because the ortho isomer was a new compound, an authentic sample was synthesized by the reaction of 2-ethylphenylboronic acid with pinacol (see below). The identity and ratio of the ortho isomer was determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with that of the authentic sample. The resonances specific to each isomer are as follows: 2.91 ppm (q, J = 8.0 Hz, 2H ortho isomer, H<sub>a</sub>); 7.62-7.65 ppm (m, 2H, meta isomer, H<sub>b</sub> and H<sub>c</sub>), 7.74 ppm (d, J = 8.0 Hz, 2H, para isomer, H<sub>d</sub>); 2.43 ppm (q, J = 8.0 Hz 1H, benzylic isomer, H<sub>e</sub>).



#### 2-(2-Ethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

A combination of *o*-ethylphenylboronic acid (300 mg, 2.0 mmol), pinacol (260 mg, 2.2 mmol) and Et<sub>2</sub>O (8 mL) were added to a round bottom flask, after which CaCl<sub>2</sub> (ca. 300 mg) was added and the reaction mixture was stirred overnight. The resulting suspension was filtered and the filtrate was concentrated in vacuo. The residue was purified by flash column chromatography over silica gel (eluting with hexane/AcOEt = 20/1) to give the title compound as a colorless oil (24 mg, 5%).

Rf 0.60 (hexane/EtOAc = 5/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.19 (t, *J* = 8.0 Hz, 3H), 1.34 (s, 12H), 2.91 (q, *J* = 8.0 Hz, 2H), 7.15-7.20 (m, 2H), 7.35 (t, *J* = 7.6 Hz, 1H), 7.76 (d, *J* = 7.2 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  17.3, 24.9, 28.9, 83.4, 124.9, 128.5, 131.0, 136.1, 151.5. IR (ATR): 2977 w, 2931 w, 2872 w, 1600 w, 1570 w, 1489 w, 1442 w, 1381 m, 1345 s, 1310 m, 1272 w, 1214 w, 1145 s, 1126 w, 1077 m, 1031 w, 962 w, 862 m, 830 w, 792 w, 755 m, 661 m. MS m/z (% relative intensity): 232 (M+, 11), 176 (11), 175 (100), 174 (33), 133 (59), 132 (87), 131 (45), 117 (36), 116 (11), 115 (27), 105 (32), 101 (63), 91 (19), 85 (23), 83 (28), 77 (11), 59 (15), 57 (17), 55 (13). HRMS (EI): Calcd for C<sub>14</sub>H<sub>21</sub>BO<sub>2</sub> 232.1635, Found 232.1630.

#### Borylation of *o*-xylene (Entry 3, Table 2).



Method A was followed except that the reaction was conducted with HBpin (150 mg, 1.17 mmol) in *o*-xylene (1.0 mL) at 100 °C. After purification by flash column chromatography over silica gel (eluting with hexane/AcOEt = 20/1), a mixture of three isomers was obtained (117 mg, 43%, >99% purity by GC). GC/MS analysis revealed the formation of three isomers of the borylated products, all of which had an m/z of 232 (M<sup>+</sup>). The identity and ratio of each of the four isomers was determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>28a,30</sup> The resonances specific to each isomer are as follows: 2.46 ppm (s, 3H, 3-isomer, H<sub>a</sub>); 7.54-7.58 ppm (m, 2H, 4-isomer, H<sub>b</sub> and H<sub>c</sub>), 1.22 ppm (s, 12H, benzylic isomer, H<sub>d</sub>). (See below.)



Borylation of anisole (Entry 4, Table 2).



Method A was followed except that the reaction was conducted with HBpin (160 mg, 1.25 mmol) in anisole (1.0 mL) at 100 °C. After purification by flash column chromatography over silica gel (eluting with hexane/AcOEt = 20/1), a mixture of three isomers of the borylated products and PhBpin were obtained (108 mg; PhBpin 7% and borylated anisole 33%; >99% purity by GC). GC/MS analysis revealed the formation of three isomers of the borylated products, all of which had an m/z of 218 (M<sup>+</sup>) and PhBpin had an m/z of 204 (M<sup>+</sup>). The identity and ratio of each of the borylated products was determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>14</sup> The resonances specific to each isomer are as follows: 7.67 ppm (dd, J = 1.6 Hz, 7.2 Hz, 1H ortho isomer, H<sub>a</sub>); 7.01 ppm (ddd, J = 0.8 Hz, 2.8 Hz, 8.0 Hz, 1H, meta isomer, H<sub>b</sub>), 7.75 ppm (d, J = 8.2 Hz, 2H, para isomer, H<sub>c</sub>); 7.81 ppm (dd, J = 1.6 Hz, 6.4 Hz, 2H, PhBpin, H<sub>d</sub>). (See below.)



Borylation of 1,3-bis(trifluoromethyl)benzene.



Method A was followed except that the reaction was conducted with HBpin (155 mg, 1.21 mmol) in 1,3-bis(trifluoromethyl)benzene (1.0 mL) at 80 °C. After purification by flash column chromatography over silica gel (eluting with hexane/AcOEt = 20/1), the product was obtained as a mixture with unidentifiable byproducts (44 mg, <10%, 84% purity by GC). HRMS analysis of this sample unambiguously determined the formation of the borylated product (Calcd for  $C_{14}H_{15}BF_6O_2$  340.1069, Found 340.1062). The identity of the borylated product was determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with that reported in the literature.<sup>31</sup> The resonances specific to the compound are as follows: 7.94 ppm (s, 1H, H<sub>a</sub>), 8.24 pp (s, 2H, H<sub>b</sub>).

## 1-Methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole (Entry 5, Table 2) [CAS: 850567-47-4].

Method A was followed except that the reaction was conducted with HBpin (160 mg, 1.25 mmol) in *N*-methylpyrrole (1.0 mL) at 80 °C.

Rf 0.49 (hexane/EtOAc = 5/1). White solid (177 mg, 68%). <sup>1</sup>H and <sup>13</sup>C NMR data in CDCl<sub>3</sub> were in agreement with those reported for 2-borylated pyrrole.<sup>32</sup> However, the title compound gradually decomposed in a CDCl<sub>3</sub> solution to form *N*-metylpyrrole. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.30 (s, 12H), 3.66 (s, *N*-methylpyrrole), 3.83 (s, 3H), 6.15-6.16 (m, 1H + *N*-methylpyrrole), 6.61 (s, *N*-methylpyrrole), 6.80-6.82 (m, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.5 (*N*-methylpyrrole), 24.7, 36.5, 83.0, 108.1 (*N*-methylpyrrole), 108.3, 121.6 (*N*-methylpyrrole), 121.8, 128.1. Pure spectra were obtained using C<sub>6</sub>D<sub>6</sub> as the solvent. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz):  $\delta$  1.09 (s, 12H), 3.51 (s, 3H), 6.29-6.30 (m, 1H), 6.51-6.52 (m, 1H), 7.32-7.34 (m, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz):  $\delta$  24.8, 36.2, 82.9, 109.1, 123.3. One carbon peak is overlapped with the peak of residual solvent. HRMS (EI): Calcd for C<sub>11</sub>H<sub>18</sub>BNO<sub>2</sub> 207.1431, Found 207.1431.

## 1-Methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (Entry 1, Table 3) [CAS: 596819-10-2].

Rf 0.54 (hexane/EtOAc = 5/1). White solid (165 mg, 78%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.36 (s, 12H), 3.97 (s, 3H), 7.06 (t, J = 7.2 Hz, 1H), 7.13 (s, 1H), 7.23-7.27 (m, 1H), 7.34 (d, J = 8.4 Hz, 1H), 7.64 (d, J = 8.4 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53

MHz):  $\delta$  24.9, 32.3, 83.8, 109.8, 114.3, 119.4, 121.7, 123.3, 127.9, 140.2. HRMS (EI): Calcd for C<sub>15</sub>H<sub>20</sub>BNO<sub>2</sub> 257.1587, Found 257.1584. All spectroscopic data were in agreement with the reported 2-borylated product.<sup>33</sup>

#### 1-Butyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (Entry 2, Table 3).

Method B was followed using 1-butyl-1H-indole (146 mg, 0.84 mmol) instead of 1-methylindole.

Rf 0.69 (hexane/EtOAc = 5/1). White solid (171 mg, 68%). Mp = 47-48 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.92 (t, *J* = 7.2 Hz, 3H), 1.29-1.34 (m, 2H), 1.36 (s, 12H), 1.74(t, *J* = 7.2 Hz, 2H), 4.41 (t, *J* = 7.2 Hz, 2H), 7.06 (t, *J* = 7.6 Hz, 1H), 7.12 (s, 1H), 7.23 (t, *J* = 7.6 Hz, 1H), 7.36 (d, *J* = 8.4 Hz, 1H), 7.63 (d, *J* = 7.6 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  14.0, 20.3, 24.9, 33.2, 45.3, 83.7, 110.6, 114.4, 119.2, 121.8, 123.0, 128.0, 139.4. IR (ATR): 3051 w, 2976 w, 2930 w, 2873 w, 1610 w, 1569 w, 1523 s, 1480 w, 1468 w, 1386 m, 1354 m, 1321 s, 1304 s, 1263 s, 1237 w, 1200 m, 1134 s, 1114 m, 1072 m, 1008 w, 964 w, 926 m, 858 m, 834 m, 803 m, 753 m, 732 s, 707 m, 690 s, 666m. MS m/z (% relative intensity): 299 (M+, 76), 257 (26), 256 (100), 156 (67), 130 (47). HRMS (EI): Calcd for C<sub>18</sub>H<sub>26</sub>BNO<sub>2</sub> 299.2057, Found 299.2053.



The regiochemistry of the compound was determined by NOE experiments in CDCl<sub>3</sub>. When signal H<sub>a</sub> at  $\delta$  = 7.12 (s, 1H, offset signal  $\delta$  = 7.109, mixing time 3.43 s) was irradiated, H<sub>b</sub> at  $\delta$  = 7.63 (d, *J* = 7.6 Hz, 1H) exhibited an enhancement by 9%. When signal H<sub>c</sub> at  $\delta$  = 4.41 (t, *J* = 7.2 Hz, 1H, offset signal  $\delta$  = 4.407, mixing time 0.443 s) was irradiated, H<sub>d</sub> at  $\delta$  = 7.36 (d, *J* = 8.4 Hz, 1H) exhibited an enhancement by 6%.

#### 1-(Methoxymethyl)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (Entry 3, Table 3).



Method B was followed using 1-(methoxymethyl)-1*H*-indole (127 mg, 0.79 mmol) instead of 1-methylindole.

Come Rf 0.57 (hexane/EtOAc = 5/1). White solid (153 mg, 68%). Mp = 52-53 °C <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz): δ 1.07 (s, 12H), 3.04 (s, 3H), 5.71 (s, 2H), 7.13 (m, 1H), 7.23 (dt, J = 1.2, 6.0 Hz, 1H), 7.53 (dd, J = 8.0 Hz, 1H), 7.55 (s, 1H), 7.63 (d, 8.0 Hz, 1H). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.37 (s, 12H), 3.22 (s, 3H), 5.81 (s, 2H), 7.13 (t, J = 7.6 Hz, 1H), 7.18 (s, 1H), 7.28 (dt, J = 1.2, 7.4 Hz, 1H), 7.53 (d, J = 8.0 Hz, 1H), 7.65 (d, J = 7.8 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz): δ 24.7, 55.0, 76.1, 83.7, 111.3, 117.0, 120.8, 122.1, 124.2, 129.3, 140.6. IR (ATR): 2979 w, 2933 w, 2363 w, 1723 w, 1612 w, 1573 w, 1528 s, 1482 w, 1369 m, 1346 m, 1322 s, 1305 s, 1263 s, 1224 m, 1191 w, 1164 m, 1137 s, 1101 m, 1081 s, 1050 m, 1010 w, 962 m, 913 w, 857 m, 833 m, 812 m, 738 s, 704 w, 688 s, 670 w. MS m/z (% relative intensity): 287 (M+, 100), 286 (27), 256 (77), 171 (27), 170 (20), 156 (68), 130 (39), 77 (20). HRMS (EI): Calcd for C<sub>16</sub>H<sub>22</sub>BNO<sub>3</sub> 287.1693, Found 287.1692.



The regiochemistry of the compound was determined by NOE experiments in CDCl<sub>3</sub>. When signal H<sub>a</sub> at  $\delta$  = 7.65 (d, *J* = 7.8 Hz, 1H, offset signal  $\delta$  = 7.651, mixing time 0.8168 s) was irradiated, H<sub>b</sub> at  $\delta$  = 7.18 (s, 1H) was enhanced by 1% and H<sub>c</sub> at  $\delta$  = 7.13 (t, *J* = 7.6 Hz, 1H) by 5%. When signal H<sub>d</sub> at  $\delta$  = 5.81 (s, 2H, offset signal  $\delta$  = 5.799, mixing time 0.8598 s) was irradiated, H<sub>e</sub>  $\delta$  = 7.53 (d, *J* = 8.0 Hz, 1H) was enhanced by 8%.

**1-Benzyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1***H***-indole (Entry 4, Table 3).** Method B was followed using 1-benzyl-1*H*-indole (150 mg, 0.73 mmol) instead of 1-methylindole.



Rf 0.43 (hexane/EtOAc = 5/1). White solid (199 mg, 82%). Mp = 125-127 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.26 (s, 12H), 5.66 (s, 2H), 7.03-7.09 (m, 3H), 7.17-7.22 (m, 5H), 7.30 (d, J = 8.4 Hz, 1H), 7.66 (d, J = 8.0 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.8, 49.0, 83.8, 110.4, 115.1, 119.6, 121.8, 123.5, 126.6, 126.9,

128.3, 128.4, 139.4, 139.9. IR (ATR): 3030 w, 2979 w, 2925 w, 2852 w, 1609 w, 1523 m, 1494 w, 1480 w, 1452 w, 1387 m, 1360 m, 1324 s, 1300 s, 1267 s, 1192 m, 1136 s, 1114 m, 1087 m, 1031 w, 1005 w, 984 m, 960 w, 926 w, 855 m, 829 m, 799 m, 772 w, 726 s, 690 s, 672. MS m/z (% relative intensity): 333 (M+, 54), 332 (30), 91 (100). HRMS (EI): Calcd for  $C_{21}H_{24}BNO_2$  333.1900, Found 333.1899.



The regiochemistry of the compound was determined by NOE experiments in CDCl<sub>3</sub>. When signal H<sub>a</sub> at  $\delta$  = 7.66 (d, *J* = 8.0 Hz, 1H, offset signal  $\delta$  = 7.673, mixing time 2.601 s) was irradiated, H<sub>b</sub> at  $\delta$  = 7.20 (s, 1H) exhibited an enhancement by 5% and H<sub>c</sub> at  $\delta$  = 7.06 (t, *J* = 7 Hz, 1H) enhanced by 12%. When signal H<sub>d</sub> at  $\delta$  = 5.66 (s, 2H, offset signal  $\delta$  = 5.659, mixing time 0.8 s) was irradiated, H<sub>e</sub> at  $\delta$  = 7.30 (d, *J* = 8.4 Hz, 1H) was enhanced by 7%.

## 1-(4-Methoxybenzyl)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (Entry 5, Table 3).



Method B was followed using 1-(4-methoxybenzyl)-1H-indole (174 mg, 0.73 mmol) instead of 1-methylindole.

Rf 0.54 (hexane/EtOAc = 5/1). White solid (201 mg, 76 %). Mp = 116 °C. <sup>1</sup>H MMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz):  $\delta$  1.04 (s, 12H), 3.19 (s, 3H), 5.55 (s, 2H), 6.59 (d, J= 8.8 Hz, 2H), 6.92 (d, J = 8.8 Hz, 2H), 7.10 (t, J = 8.2 Hz, 1H), 7.19 (t, J = 7.2 Hz, 1H), 7.25 (d, J = 8.4 Hz, 1H), 7.65 (s, 1H), 7.70 (d, J = 8.0 Hz, 1H). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.30 (s, 12H), 3.74 (s, 3H), 5.59 (s, 2H), 6.76 (d, J = 8.7 Hz, 2H), 7.02 (d, J = 8.8 Hz, 2H), 7.07 (t, J = 8.0 Hz, 1H), 7.18 (s, 1H), 7.19 (t, J = 7.6 Hz, 1H), 7.32 (d, J = 8.4 Hz, 1H), 7.65 (d, J = 7.8 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz):  $\delta$  24.6, 48.5, 54.5, 83.6, 110.8, 114.6, 116.1, 120.0, 122.2, 123.8, 127.8 (overlapped with the solvent peak), 129.0, 131.6, 140.4, 159.0. <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.7, 48.3, 55.1, 83.7, 110.3, 113.6, 114.9, 119.4, 121.6, 123.2, 127.7, 128.1, 131.3, 139.6, 158.4. IR (ATR): 2977 w, 1613w, 1514 s, 1481 w, 1462 w, 1441 w, 1382 m, 1354 m, 1321 m, 1295 s, 1269 s, 1249 s, 1193 m, 1174 m, 1134 s, 1085 m, 1038 m, 985 w, 960 w, 856 m, 830 m, 805 737 s, 704 w, 688 s, 672 w. MS m/z (% relative intensity): 363 (M+, 17), 121 (100). HRMS (EI): Calcd for C<sub>22</sub>H<sub>26</sub>BNO<sub>3</sub> 363.2006, Found 363.2005.



The regiochemistry of the compound was determined by NOE experiments in CDCl<sub>3</sub>. When signal H<sub>a</sub> at  $\delta$  = 7.65 (d, *J* = 7.8 Hz, 1H, offset signal  $\delta$  = 7.647, mixing time 2.379 s) was irradiated, H<sub>b</sub> at  $\delta$  = 7.18 (s, 1H) was enhanced by 7% and H<sub>c</sub> at  $\delta$  = 7.07 (t, *J* = 8.0 Hz, 1H) had an enhancement of 10%. When signal H<sub>d</sub> at  $\delta$  = 7.32 (d, *J* = 8.4 Hz, 1H, offset signal  $\delta$  = 7.317, mixing time 2.379 s) was irradiated, He at  $\delta$  = 5.59 (s, 2H) had an NOE enhancement of

6%.

#### 1,3-Dimethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (Entry 6, Table 3).



Method B was followed using 1,3-dimethyl-1*H*-indole (116 mg, 0.80 mmol) instead of 1-methylindole.

Me Rf 0.54 (hexane/EtOAc = 5/1). White solid (187mg, 87%). Mp = 88-89 °C. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz): δ 1.13 (s, 12H), 2.85 (s, 3H), 3.71 (s, 3H), 7.13 (d, J = 8.8 Hz, 1H), 7.19 (t, J = 7.6 Hz, 1H), 7.31 (dt, J = 0.8, 8.0 Hz, 1H), 7.74 (d, J = 7.6 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz): δ 10.7, 24.7, 32.0, 83.0, 109.8, 119.0, 120.3, 123.8, 125.6, 129.3, 140.5. IR (ATR): 3051 w, 2979 w, 2932 w, 2863 w, 1711 w, 1610 w, 1530 m, 1438 m, 1410 m, 1370 m, 1334 m, 1294 s, 1264 s, 1242 m, 1213 m, 1166 m, 1144 s, 1088 m, 1045 m, 1007 w, 962 m, 859 m, 835 m, 739 s, 714 w, 697 s, 672 w. MS m/z (% relative intensity): 272 (16), 271 (M+, 100), 270 (36), 189 (22), 188 (41), 171 (20), 170 (32), 145 (10), 144 (44), 115 (12), 77 (14). HRMS (EI): Calcd for C<sub>16</sub>H<sub>22</sub>BNO<sub>2</sub> 271.1744, Found 271.1745.

## 5-Methoxy-1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (Entry 7, Table3).

MeO M N Bpin m

Method B was followed using 5-methoxy-1-methyl-1*H*-indole (127 mg, 0.79 mmol) instead of 1-methylindole.

Me Rf 0.46 (hexane/EtOAc = 5/1). White solid (172 mg, 76%). Mp = 100-101 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.35 (s, 12H), 3.84 (s, 3H), 3.93 (s, 3H), 6.93 (dd, J = 2.0 Hz, 9.2 Hz, 1H), 7.02 (s, 1H), 7.05 (d, J = 2.4 Hz, 1H), 7.23 (d, J = 9.2 Hz, 1H). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz) δ 1.12 (s, 12H), 3.44 (s, 3H), 3.67 (s, 3H), 6.98 (d, J = 8.7 Hz, 1H), 7.06 (d, J = 2.3 Hz, 1H), 7.17 (dd, J = 2.4 Hz, 8.8 Hz, 1H), 7.56 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 24.9, 32.5, 55.9, 83.7, 102.1, 110.6, 113.5, 114.5, 128.1, 135.9, 154.0. IR (ATR): 2977 w, 1619 w, 1521 s, 1457 w, 1394 m, 1372 w, 1346 w, 1315 m, 1299 s, 1260 s, 1233 s, 1206 s, 1169 m, 1139 s, 1107 m, 1066 m, 1036 m, 963 m, 849 m, 832 s, 792 s, 706 w, 689 s, 677w. MS m/z (% relative intensity): 287 (M+, 100), 286 (25), 272 (41), 190 (17). HRMS (EI): Calcd for C<sub>16</sub>H<sub>22</sub>BNO<sub>3</sub> 287.1693, Found 287.1695.



The regiochemistry of the compound was determined by NOE experiments in C<sub>6</sub>D<sub>6</sub>. When signal H<sub>a</sub> at  $\delta$  = 7.06 (d, *J* = 2.3 Hz, 1H, offset signal  $\delta$  = 7.006, mixing time 1.52 s) was irradiated, H<sub>b</sub> at  $\delta$  = 7.56 (s, 1H) exhibited an enhancement by 9%, and H<sub>c</sub> at  $\delta$  = 3.44 (s, 3H) by 3 %.

#### 5-Fluoro-1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (Entry 8, Table 3).



Method B was followed using 5-fluoro-1-methyl-1*H*-indole (115 mg, 0.77 mmol) and 1.2 equivalent of HBpin (120 mg, 0.94 mmol).

Me Rf 0.49 (hexane/EtOAc = 5/1). White solid (173 mg, 82%). Mp = 116 °C. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 395.88 MHz) δ 1.09 (s, 12H), 3.56 (s, 3H), 6.77 (dd, J = 4.0 Hz, 8.8 Hz, 1H), 6.99 (dt, J = 2.4 Hz, 8.8 Hz, 1H), 7.26 (dd, J = 2.4 Hz, 9.2 Hz, 1H), 7.37 (d, J = 1.2 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 99.54 MHz): δ 24.7, 32.1, 83.6, 106.3 (d, J = 23 Hz), 110.6 (d, J = 10 Hz), 112.2 (d, J = 27 Hz), 115.0 (d, J = 5 Hz), 128.6 (d, J = 10 Hz), 137.4, 158.4 (d, J = 235 Hz). IR (ATR): 2982 m, 1523 s, 1458 w, 1390

m, 1372 m, 1317 m, 1295 m, 1254 s, 1181 m, 1137 s, 1065 m, 962 m, 861 m, 846 s, 787 s, 707 w, 690 s, 678 m. MS m/z (% relative intensity): 276 (16), 275 (M+, 100), 274 (28), 202 (20), 193 (20), 190 (32), 176 (19), 175 (43), 174 (30). HRMS (APCI): Calcd for [M+H]<sup>+</sup> C<sub>15</sub>H<sub>20</sub>BFNO<sub>2</sub> 276.1568, Found 276.1569.



The regiochemistry of the compound was determined by NOE experiments in C<sub>6</sub>D<sub>6</sub>. When signal H<sub>a</sub> at  $\delta$  = 7.37 (d, *J* = 1.2 Hz, 1H, offset signal  $\delta$  = 7.3116, mixing time 1.5 s) was irradiated, H<sub>b</sub> at  $\delta$  = 7.26 (dd, *J* = 2.4, 9.2 Hz, 1H) exhibited an enhancement of 2%.

#### 1-Methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-aza-1*H*-indole (Entry 9, Table 3).

Bpin N Me Method B was followed using 1-methyl-7-aza-1*H*-indole (105 mg, 0.80 mmol) and 1.2 equivalent of HBpin (120 mg, 0.94 mmol).

<sup>N</sup> Me Rf 0.14 (hexane/EtOAc = 5/1). White solid (116 mg, 56%). Mp = 66-68 °C. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz):  $\delta$  1.07 (s, 12H), 4.09 (s, 3H), 6.74 (dd, *J* = 4.6 Hz, 7.8 Hz, 1H), 7.38 (s, 1H), 7.59 (dd, *J* = 1.3 Hz, 7.8 Hz, 1H), 8.48 (dd, *J* = 1.3 Hz, 4.4 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz):  $\delta$  24.7, 30.9, 83.7, 112.9, 115.9, 120.2, 129.3, 145.3, 151.1. IR (ATR): 2978 w, 1592 w, 1564 w, 1521 s, 1461 m, 1380 m, 1323 s, 1254 s, 1214 w, 1135 s, 1109 m, 1061 m, 961 w, 910 w, 855 m, 834 m, 816 m, 774 s, 724 w, 707 w, 690 s, 668 m. MS m/z (% relative intensity): 258 (M+, 100), 257 (31), 185 (20), 173 (30), 159 (27), 158 (27) 157 (32), 132 (24), 131 (72). HRMS (EI): Calcd for C<sub>14</sub>H<sub>19</sub>BN<sub>2</sub>O<sub>2</sub> 258.1540, Found 258.1542.



The regiochemistry of the compound was determined by NOE experiments in C<sub>6</sub>D<sub>6</sub>. When signal H<sub>a</sub> at  $\delta$  = 7.59 (dd, *J* = 1.3, 7.8 Hz, 1H, offset signal  $\delta$  = 7.547, mixing time 2.7 s) was irradiated, H<sub>b</sub> at  $\delta$  = 7.38 (s, 1H) was enhanced by 3% and H<sub>c</sub> at  $\delta$  = 6.74 (dd, *J* = 4.6, 7.8 Hz, 1H) by 8%.

#### 1.4.5. Mechanistic Studies

#### 5-1. Kinetic Isotope Effect

The reaction progress was monitored by GC. Dodecane was added as an internal standard.

A typical procedure for kinetic studies. In a glovebox,  $Ni(cod)_2$  (9.6 mg, 0.035 mmol), ICy•HCl (9.2 mg, 0.034 mmol), NaO'Bu (6.4 mg, 0.067 mmol), HBpin (161 mg, 1.3 mmol) and dodecane (60 mg as internal standard) were added to a 10 mL-sample vial with a Teflon-sealed screw cap. Benzene (1.0 mL) was then added and the cap was applied. The vial was heated at 80 °C with stirring over a hot plate. At the indicated times, the vial was rapidly cooled with running water and an aliquot of the reaction mixture was taken from the vial in a glovebox. The aliquot was diluted with AcOEt and analyzed by GC. After sampling, the reaction mixture was heated again until the next sampling.

The amount of PhBpin in the reaction mixture was determined over time, and the results are shown below.



The total yield after 8 hours was 72%. These results showed that reaction monitoring by cooling did not deactivate the catalytic species. We observed that the reaction had an induction period of approximately 20 minutes at 80  $^{\circ}$ C.

**Measurement of kinetic isotope effect.** The initial rate up to 20% conversion was measured following the induction period, in which ca. 1 % of PhBpin was generated. The experiments were performed with benzene and benzene- $d_6$ . Each experimental trial was performed in duplicate.



The average rate with benzene was determined to be  $7.5 \times 10^{-5}$  M/s, while the average rate with benzene- $d_6$  was determined to be  $3.5 \times 10^{-5}$  M/s. The KIE for this reaction was 2.1.

#### 5-2. Test for Heterogeneous Catalyst



**Filtration test.** In a glovebox, Ni(cod)<sub>2</sub> (9.1 mg, 0.033 mmol), ICy•HCl (9.4 mg, 0.035 mmol), NaO'Bu (6.8 mg, 0.070 mmol), HBpin (163 mg, 1.3 mmol) and dodecane (59 mg as internal standard) were added to a 10 mL-sample vial with a Teflon-sealed screw cap. Benzene (1.0 mL) was then added and the cap was applied. The vial was heated at 80 °C with stirring over a hot plate. After 70 minutes, the vial was rapidly cooled under running water. An aliquot was taken from the mixture in a glovebox for GC analysis, which indicated that PhBpin was formed in 21% yield. The remainder of the reaction mixture was passed through alumina using 1 mL of benzene as the eluent. The filtrate was then transferred to a new 10 mL-sample vial and heated again for 13 h. The yield after 13 h was determined to be 21 %, thus demonstrating that the reaction did not proceed further following filtration.

Note: When the reaction was performed under condition A with added alumina (40 mg), 63% GC yield was found following a 19 h reaction time span. This result indicated that alumina did not have a significant negative effect on this reaction.



**Hg drop test.** In a glovebox, Ni(cod)<sub>2</sub> (10.5 mg, 0.038 mmol), ICy•HCl (10.2 mg, 0.038 mmol), NaO'Bu (6.4 mg, 0.067 mmol), HBpin (164 mg, 1.3 mmol), dodecane (64 mg as internal standard) and Hg (145 mg, 0.72 mmol) were added to a 10 mL-sample vial with a Teflon-sealed screw cap. Benzene (1.0 mL) was then added and the cap was applied. The vial was heated at 80 °C for 20 h with stirring over a hot plate. The resulting mixture was analyzed by GC and none of the target product was observed.

#### **5-3.** Diboron as a Borylating Reagent

This reaction was conducted with  $B_2pin_2$  as the boron source instead of HBpin.



In a glovebox, Ni(cod)<sub>2</sub> (7.5 mg, 0.027 mmol), ICy•HCl (7.4 mg, 0.028 mmol), NaO'Bu (7.2 mg, 0.074 mmol) and B<sub>2</sub>pin<sub>2</sub> (126 mg, 0.49 mmol) were added to a 10 mL-sample vial with a Teflon-sealed screw cap. Benzene (1.0 mL) was then added and the cap was applied. The vial was heated at 80 °C for 20 h with stirring over a hot plate. The resulting mixture was filtered through a silica gel column (eluting with 10 mL of hexane/AcOEt =5/1). The filtrate was analyzed by GC with dodecane as the internal standard and the target product was found in a 125% GC yield based on the molar amount of B<sub>2</sub>pin<sub>2</sub>.

#### 5-4. Effect of Base

Method A was followed using 3 mol% of the bases listed in the table. NaO'Bu was chosen as the optimal base for nickel-catalyzed C-H borylation.



The equivalent of base dramatically influenced the reaction efficiency in both cases using HBpin and B<sub>2</sub>pin<sub>2</sub>.



When the reaction was performed in the absence of the NaO'Bu using free IMes, the yield dramatically decreased compared to the results when an IMes ligand generated from IMes•HCl and NaO'Bu was used.



#### **1.5 References and Notes**

- (1) K. M. Waltz, X. He, C. Muhoro, J. F. Hartwig, J. Am. Chem. Soc. 1995, 117, 11357.
- (2) T. Hatanaka, Y. Ohki, K. Tatsumi, Chem.-Asian J. 2010, 5, 1657.
- (3) G. Yan, Y. Jiang, C. Kuang, S. Wang, H. Liu, Y. Zhang, J. Wang, *Chem. Commun.* **2010**, *46*, 3170.
- (4) T. J. Mazzacano, N. P. Mankad, J. Am. Chem. Soc. 2013, 135, 17258.
- (5) T. Dombray, C. G. Werncke, S. Jiang, M. Grellier, L. Vendier, S. Bontemps, J.-B. Sortais, S. Sabo-Etienne, C. Darcel, J. Am. Chem. Soc. 2015, 137, 4062.
- (6) a) J. V. Obligacion, S. P. Semproni, P. J. Chirik, J. Am. Chem. Soc. 2014, 136, 4133. b) J. V. Obligacion, S. P. Semproni, I. Pappas, P. J. Chirik, J. Am. Chem. Soc. 2016, 138, 10645. c) W. N. Palmer, J. V. Obligacion, I. Pappas, P. J. Chirik, J. Am. Chem. Soc. 2016, 138, 766. d) N. G. Léonard, M. J. Bezdek, P. J. Chirik, Organometallics 2016, in press (DOI: 10.1021/acs.organomet.6b00630). e) Manna, K.; Ji, P.; Lin, Z.; Greene, F. X.; Urban, A.; Thacker, N. C.; Lin, W. Nat. Commun. 2016, 7, 12610.
- (7) Selected reviews on C-H bond functionalization under Fe catalysis: a) E. Nakamura, N. Yoshikai, *J. Org. Chem.* 2010, 75, 6061. b) C.-L. Sun, B.-J. Li, Z.-J. Shi, *Chem. Rev.* 2011, 111, 1293. c) B. D. Sherry, A. Fürstner, *Acc. Chem. Res.* 2008, 41, 1500.
- (8) Selected reviews on C-H bond functionalization under Co catalysis: a) N. Yoshikai, *Bull. Chem. Soc. Jpn.* 2014, 87, 843. b) K. Gao, N. Yoshikai, *Acc. Chem. Res.* 2014, 47, 1208. c) L. Ackermann, *J. Org. Chem.* 2014, 79, 8948.
- (9) J. P. Kleiman, M. Dubeck, J. Am. Chem. Soc. 1963, 85, 1544.
- (10) A review: L. C. M. Castro, N. Chatani, Chem. Lett. 2015, 44, 410.
- (11) Reviews: a) Y. Nakao, *Chem. Rec.* 2011, *11*, 242. b) J. Yamaguchi, K. Muto, K. Itami, *Eur. J. Org. Chem.* 2013, 19. c) S. A. Johnson, *Dalton Trans.* 2015, *44*, 10905.
- (12) S. Okumura, S. Tang, T. Saito, K. Semba, S. Sakaki, Y. Nakao, J. Am. Chem. Soc. 2016, 138, 14699.
- (13) Y. Morimoto, S. Bunno, N. Fujieda, H. Sugimoto, S. Itoh, J. Am. Chem. Soc. 2015, 137, 5867.
- (14) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaura, N. R. Anastasi, J. F. Hartwig, J. Am. Chem. Soc. 2002, 124, 390.
- (15) S. Shimada, A. S. Batsanov, J. A. K. Howard, T. B. Marder, *Angew. Chem., Int. Ed.* **2001**, *40*, 2168.
- (16) a) K. Nakamura, M. Tobisu, N. Chatani, Org. Lett. 2015, 17, 6142. b) C. Zarate, R. Manzano, R. Martin, J. Am. Chem. Soc. 2015, 137, 6754.
- (17) J. S. Bair, Y. Schramm, A. G. Sergeev, E. Clot, O. Eisenstein and J. F. Hartwig, J. Am. Chem. Soc. 2014, 136, 13098.
- (18) a) X.-W. Liu, J. Echavarren, C. Zarate, R. Martin, J. Am. Chem. Soc. 2015, 137, 12470. b) T. Niwa, H. Ochiai, Y. Watanabe, T. Hosoya, J. Am. Chem. Soc. 2015, 137, 14313. c) J. Zhou, M. W. Kuntze-Fechner, R. Bertermann, U. S. D. Paul, J. H. J. Berthel, A. Friedrich, Z. Du, T. B. Marder, U. Radius, J. Am. Chem. Soc. 2016, 138, 5250.

- (19) A. A. Toutov, W.-B. Liu, K. N. Betz, A. Fedorov, B. M. Stoltz, R. H. Grubbs, *Nature* 2015, *518*, 80.
- (20) A related base effect in Ir-catalyzed C-H borylation: a) M. N. Eliseeva and L. T. Scott, J. Am. Chem. Soc. 2012, 134, 15169. b) T. Ohmura, T. Torigoe, M. Suginome, Chem. Commun. 2014, 50, 6333.
- (21) H. Zhang, S. Hagihara, K. Itami, Chem. Lett. 2015, 44, 779.
- (22) Unique reactivity of heterogeneous nickel species in C-O bond activation: A. G. Sergeev, J. D. Webb, J. F. Hartwig, J. Am. Chem. Soc. 2012, 134, 20226.
- (23) C. J. Serpell, J. Cookson, A. L. Thompson, C. M. Brown, P. D. Beer, *Dalton Trans.* 2013, 42, 1385.
- (24) E. A. Mistryukov, Mendeleev Commun. 2006, 16, 258.
- (25) N. Gürbüz, I. Özdemir, S. Demir, B. Çetinkaya, J. Mol. Cat. A Chem. 2004, 209, 23.
- (26) M. Tobisu, T. Morioka, A. Ohtsuki, N. Chatani, Chem. Sci. 2015, 6, 3410.
- (27) H. Huo, C. Fu, K.Harms, E. Meggers, J. Am. Chem. Soc. 2014, 136, 2990.
- (28) a) G. Yan, Y. Jiang, C. Kuang, S. Wang, H. Liu, Y. Zhang, J. Wang, *Chem. Commun.* 2010, 46, 3170. b) C. Pintaric, S. Olivero, Y. Gimbert, P. Y. Chavant, D. Duñach, *J. Am. Chem. Soc.* 2010, 132, 11825.
- (29) T. A. Boebel, J. F. Hartwig, Organometallics 2008, 27, 6013.
- (30) K. Endo, T. Ohkubo, T. Shibata, Org. Lett. 2011, 13, 3368.
- (31) K. Wilckens, D. Lentz, C. Czekelius, Organometallics 2011, 30, 1287.
- (32) A. D. Grosso, P. J. Singleton, C. A. Muryn, M. J. Ingleson, *Angew. Chem., Int. Ed.* **2011**, *50*, 2102.
- (33) K. Mertins, A. Zapf, M. Beller, J. Mol. Cat. A Chem. 2004, 207, 21.

#### Chapter 2

#### Platinum-Catalyzed Borylation of Aromatic C-H Bonds

#### **2.1 Introduction**

Based on the results for the nickel-catalyzed C-H borylation of arenes described in Chapter 1, the author became interested in the reactivity of its heavier congener platinum for the C-H borylation of arenes. When this study was initiated, there were no other reports in the literature describing the use of a platinum complex in a C-H borylation reaction. Additionally, catalytic C-H functionalization reactions of arenes using platinum catalysts have been much less studied<sup>1</sup> compared to reactions involving palladium and nickel catalysts, and only a few examples of C-H silylation,<sup>2</sup> intramolecular C-H amination,<sup>3</sup> hydroarylation<sup>4</sup> and arylation<sup>5</sup> can be found in the literature. In contrast, it is well known that platinum complexes are efficient catalysts for hydroboration and diboration reactions,<sup>6</sup> which have been postulated to be mediated by a platinum-boryl intermediate.<sup>7</sup> The author hypothesized that a platinum-boryl species could also mediate C-H borylation reactions if it were used with an appropriate choice of ligand and/or additive. Herein, an efficient borylation reaction of aromatic C-H bonds catalyzed by N-heterocyclic carbene (NHC)-ligated platinum catalysts bearing a pincer ligand can effect the C-H borylation of polyfluorinated arenes.<sup>8</sup>

#### 2.2 Results and Discussion

**C-H Borylation with Pt/NHC Complexes**. The author initially evaluated the performance of Karstedt's catalyst **4** [Pt<sub>2</sub>(dvtms)<sub>3</sub>, dvtms = 1,3-divinyltetramethyldisiloxane] for the reaction of benzene with bis(pinacolato)diboron (**1a**), and found that this system produced a small amount of the target borylated product **3a** (TON =5, Table1, Entry1). The effectiveness of NHCs as ligands in the nickel-catalyzed C-H borylation reaction described in Chapter 1 led the author to investigate the catalytic activity of Pt/NHC catalysts. Pt(NHC)(dvtms) complexes was chosen in this particular case because they are air stable and easy to handle platinum catalyst precursors, which can be prepared in two steps from H<sub>2</sub>PtCl<sub>6</sub>. The Pt(NHC)(dvtms) complexes have been reported to be effective catalysts for hydrosilylation.<sup>9</sup> In the current study, the introduction of NHCs generally led to an improvement in the turnover number (TON) values of the catalysts. For example, the IMes-ligated platinum complex

B₂pin	ha <b>+</b>	Pt catalyst (0.30 m	Pt catalyst (0.30 mol %)		
1a		<sup>↓</sup> 100 °C, 20 h		., -	пвріп
	2a		3a		
	(solve	ent)			
	Entry	Catalyst	NHC	TON <sup>b</sup>	
	1	$Pt_2(dvtms)_3$ (4)	-	5	
	2	4a	IMes	40	
	3	4b	IPr	0	
	4	<b>4c</b>	IMe	6	
	5	<b>4d</b>	I <sup>i</sup> Pr	69	
	6	<b>4</b> e	I <sup>t</sup> Bu	16	
	7	<b>4</b> f	I(1-Ad)	58	
	8	4g	ICy	157	
	9	<b>4h</b>	SICy	106	
	10	<b>4i</b>	BICy	102	
	11	$Pt(PPh_3)_4$	-	0	
	12 <sup>c</sup>	$4 + PCy_3$	-	2	
1	13 <sup>c</sup>	4 + 2,2'-bipyridine	-	13	L
	1	R Mo	Cy	Мо	
	/ N	Si Si	N ×	-Si	
	L.		⊢ Pť	) O	
		R Me <sub>2</sub>	Cy	Me <sub>2</sub>	
	<b>4</b> a:	R = 2.4.6-Me <sub>2</sub> C <sub>e</sub> H <sub>2</sub>	4h		
	4b:	$R = 2,6^{-i}Pr_2C_6H_3$	Çy	Mea	
	<b>4c</b> :	R = Me	ŗŃ, _ 🏷	-Si	
	4d:	R = Pr		,0 	
	4e: 4f· i	к = ъu R = 1-adamantvl		Me <sub>2</sub>	
	4g:	R = cyclohexyl	2, 4i		

Table 1. Platinum-Catalysts for the C-H Borylation of Benzene<sup>a</sup>

<sup>a</sup>Reaction conditions: **1a** (0.30 mmol), catalyst (0.90  $\mu$ mol) in benzene (1.0 mL) at 100 °C for 20 h. <sup>b</sup>TON: molar amount of **3a** formed per molar amount of catalyst. <sup>c</sup>**4** (6.0  $\mu$ mol) and PCy<sub>3</sub> or 2,2'-bipyridine (6.0  $\mu$ mol), if indicated, were used.

**4a** resulted in an 8-fold improvement in the TON to 40 (Entry 2), whereas the IPr complex **4b** gave none of the desired borylated product (Entry 3). Among the NHCs examined in this study, the ICy-ligated complex **4g** was determined to be optimal in terms of its TON for the borylation of benzene (Entry 8). In the presence of 2 mol% of **4g**, the borylated product **3a** was isolated in 93% yield (Scheme1). In contrast, phosphine and bipyridine ligands were much less effective than **4g** (Entries 11-13). Diboron reagent **1a** was determined the most effective boron source for this platinum-catalyzed C-H borylation. The use of other boron reagents such as pinacolborane (HBpin) or

bis(neopentylglycolato)diboron (B<sub>2</sub>nep<sub>2</sub>) gave insufficient results (1% and 6% yield, respectively, in the presence of 2 mol% 4g).

Scheme 1. Platinum-Catalyzed Borylation of Benzene



With the optimized conditions in hand, the author proceeded to investigate the borylation of a series of substituted arenes to examine the regioselectivity of this platinum-catalyzed borylation reaction (Figure 2). The borylation of toluene was conducted at 120 °C with 2 mol% of the ICy catalyst 4g, leading to the formation of a mixture of o-, m- and p-borylated products **3b** without any of the benzylic borylation product. The formation of a significant amount of the ortho-borylated isomer indicated that the present platinum catalyst system was not particular sensitive to steric effects. Catalyst 4g was also used to borylate anisole, trifluorobenzene and chlorobenzene, resulting in o-m- and p-borylated mixtures of the corresponding products 3c-3e, respectively, indicating that this catalyst system was insensitive to the electronic properties of the arene substrate. The borylation of *p*-xylene proceeded efficiently to give the 2-borylated product **3f**, as a single product. In the case of 4-tert-butyltoluene (2g), the borylation proceeded at the least hindered 2-position to give 3g. This platinum catalyst system was also used to borylate the C-H bond ortho to the isopropyl group in 2h, albeit in a low yield. m-Xylene was also borylated under these platinum-catalyzed conditions to give a mixture of regioisomers. Surprisingly, some borylation was observed at the sterically congested 2-position of m-xylene, in addition to borylation at the 4- and 5-positions (2:4:5 = 10:25:65, 3i). This reactivity contrasts sharply with the selectivity observed for iridium catalysts, where the least hindered 5-position is borylated exclusively.<sup>10</sup>

**Borylation of Sterically Congested C-H Bonds.** The high sensitivity of iridium catalysts to steric hindrance allows for the selective borylation of the least hindered C-H bond in an arene substrate. However, this sensitivity can become a problem for the borylation of sterically congested C-H bonds such as those of mesitylene. For instance, the borylation of mesitylene in the presence of an Ir/dtbpy catalyst occurs exclusively at the sterically more accessible benzylic C-H bond (eq 1). To the best of our knowledge, the only reported example of the  $C(sp^2)$ -H borylation of mesitylene involved the use
Figure 2. Platinum-Catalyzed Borylation of Substituted Arenes<sup>a</sup>



<sup>a</sup>Reaction conditions: arene (1.0 mL), **1a** (0.30 mmol), **4g** (2 mol%), 120 °C, 20 h. Yields are based on **1a**. The regioisomeric ratios were determined by <sup>1</sup>H NMR spectroscopy or GC analysis. <sup>b</sup>**4f** (2 mol%) was used instead of **4g**. <sup>c</sup>At 140 °C. <sup>d</sup>Isolated yield determined by the conversion of the product to the corresponding phenol.



of a heterogeneous iron catalyst (41% GC yield after 4 days).<sup>11</sup> Furthermore, the more sterically hindered substrate 1,3,5-triethylbenzene (**2m**) failed to react under the iridium-catalyzed conditions (eq 2). The development of new methods for the C-H borylation of hindered arenes is therefore highly desired because these systems are common structural motifs in a variety of different fields, including pharmaceuticals,<sup>12</sup> photocatalyst,<sup>13</sup> steric protection of reactive chemical species,<sup>14</sup> chiral catalysts,<sup>15</sup> and functional polymers<sup>16</sup> (Figure 3). The low sensitivity exhibited by our platinum catalyst towards the steric hindrance exerted by neighboring substituents prompted the author to examine the borylation of mesitylene. Pleasingly, catalyst **4g** allowed for the successful borylation of the sterically hindered  $C(sp^2)$ –H bond in mesitylene to give **3j** in 55% yield as the sole product (eq 3). In an attempt to improve the yield of this reaction, the author investigated the effect of the ligand (See experimental section for details about the effect of the ligand). In this regard, the IPr-bound platinum complex **4b** was found to be more effective than **4g**, leading to a considerable improvement in the yield of **3j** to 72%.

As shown in Entry 3 of Table 1, catalyst **4b** was totally ineffective for the borylation of benzene. NMR monitoring of the reaction using **4b** revealed that the dvtms ligand of **4b** does not decompose under the catalytic conditions at 100 °C, while the decomposition occurs at 120 °C. In contrast, the dvtms ligand of ICy-bound complex **4g** dissociates at 100 °C (see eq 6). Therefore, the nature of the ligand affects not only the catalytic activity for the borylation reaction, but also the efficiency of the generation of active catalytic species (i.e., the efficiency of the dissociation of dvtms), and both effects would lead to an improved yield of the borylated product.

Catalyst **4b** effectively borylated a series of 1,3,5-trisubstituted arenes to give the corresponding arylboronic ester bearing two *ortho* substituents (Figure 4). A *tert*-butyl group was sufficiently large to block borylation at its *ortho* position in **2k**, as evidenced by the exclusive formation of **3k**. Similarly, 1-isopropyl-3,5-dimethylbenzene (**2l**) was regioselectively borylated at its 4-position (the least hindered position) to give **3l**, although some borylation was also observed at the C-H bond *ortho* to the isopropyl group (positions 2:4 = 92:8), further highlighting that platinum is much less sensitive to steric hindrance than iridium. It is noteworthy that 1,3,5-triethylbenzene (**2m**) was successfully borylated through the activation of its  $C(sp^2)$ -H bonds, which were shielded by two *ortho*-ethyl groups to give the corresponding borylated product **3m**. The iridium/dtbpy catalyst was completely inactive against **2m** (eq 2). This result therefore clearly highlights the unique properties of platinum catalyst systems for the borylation of sterically hindered  $C(sp^2)$ -H bonds. Furthermore, several functional groups, including esters, amines, chlorides and fluorides were tolerated under these conditions, which allows access to a series of elaborate hindered arylboronic esters **3n-3q**.



Figure 3. Widespread Applications of Hindered Arenes

Figure 4. Platinum-Catalyzed Borylation of C-H Bonds in Sterically Congested Arenes<sup>a</sup>



<sup>a</sup>Reaction conditions: arene (1.0 mL), **1a** (0.30 mmol), **4b** (2 mol%), 120 °C, 20 h. Yields are based on **1a**. The regioisomeric ratios were determined by <sup>1</sup>H NMR spectroscopy. <sup>b</sup>At 140 °C, <sup>c</sup>The dechlorinated product **3j** was also formed (4%).

Although this platinum-catalyzed borylation requires an excess (ca. 20-35 equiv) of the arene substrate to obtain satisfactory levels of conversion, the unreacted substrate can be recovered quantitatively and used for another borylation reaction. However, the need for such a large excess of substrate clearly limits the application of this method to more elaborate substrates. To address this issue, the author conducted a series of optimization to identify much more practical conditions. The effect of the ligand was investigated in much greater detail using hindered arene 2k as a limiting agent in the presence of 1.0 equiv of 1a under neat conditions (Table 2). The IPr-ligated catalyst 4b was still active under these conditions and gave the borylated product **3k** in 51% yield (Entry 1). Although the yield of **3k** was moderate, **1a** was completely consumed, indicating that **1a** was also involved in some undesired pathways. There have been several reports pertaining to C-H borylation reactions, where the catalyst is deactivated through the C-H borylation of the ligand.<sup>17</sup> These reports led the author to investigate the use of IPr derivatives with fewer C-H bonds capable of being borylated. Whilst the use of an NHC bearing 1,3,5-triisopropylphenyl groups (IPr', 4j) was less effective than IPr (Entry 2), the use of IPr derivatives with modifications on the imidazole backbone, such as SIPr (4k), IPr<sup>Me</sup> (4l) and IPr<sup>AN</sup> (4m), all improved the yield of the borylated product 3k (Entries 3-5). These results therefore suggested that the higher efficiency of catalysts 4k-4m could be attributed to their larger size<sup>18</sup> or limited rotation around their C(aryl)-N bonds,<sup>19</sup> rather than suppressing the C-H borylation of the imidazole ring. To develop conditions applicable to various arene substrates, the author subsequently investigated the use of several solvents with platinum catalyst 4k (Entries 6-10). The borylation was found to be highly sensitive to the concentration of the aromatic substrate. For example, concentration of 0.3 and 3.0 M resulted in yields of 18 and 58% for the borylation of 2k in octane (Entries 6 and 7, respectively). Several other nonpolar solvents, such as cyclooctane, decane and triisopropylbenzene (TIPB) were also used without any discernible loss of catalytic activity, thereby expanding the scope of this borylation procedure to solid arene substrates. product **3q**-*di* was obtained as the major isomer when the borylation reaction was repeated (90% combined yield, 3q-mono:3q-di = 28:72). The new protocol developed above allowed for the borylation of several relatively complicated arenes, such as those containing menthol (3v) and borneol (3w) substructures. The protocol was capable of regioselectively borylating a silyl-substituted substrate to form an arene bearing both boron and silicon functionalities (**3**y), which are amenable to orthogonal functionalization (*vide infra*). Trifluoromethyl (3z), alkyl chloride (3aa) and TBDMS ether (3ab) groups were also found to be compatible with these reaction conditions. Carbonyl functionalities such as benzamide (3ac), anilide (3ad) and alkyl ester (3ae) groups were also tolerated under these conditions.



Table 2. Optimization towards Practical C-H Borylation of Hindered Arenes<sup>a</sup>

<sup>a</sup>Reaction conditions: **2k** (0.30 mmol), **1a** (0.30 mmol for Entries 1-5; 0.36 mmol in Entries 6-10), Pt(NHC)(dvtms) (3 mol%), 120 °C, 16 h. <sup>b</sup>Yields determined by GC based on **2k**. <sup>c</sup>Isolated yield. TIPB = 1,3,5-triisopropylbenzene.

It is noteworthy that this platinum-catalyzed methodology is not limited to the borylation of C-H bonds surrounded by two methyl groups. For example, C-H bonds between methyl and methoxy groups (**3af**), two ethyl groups (**3m**) and two *n*-propyl groups (**3ag**) were all successfully borylated under these modified conditions to form the corresponding products. Although the yields of these products were moderate, most of the unreacted substrates could be recovered in good yields. It was therefore possible to improve the yields of these borylated products by simply repeating the borylation reaction. For example, the elaborate boronic ester **3v** was successfully synthesized on a gram scale by simply repeating the borylation reaction (Scheme 2). Increasing the loading of the platinum catalyst did not generally lead to improvements in the yields of the borylated products. The origin of the unique



Figure 5. Substrate Scope for Sterically Congested Arenes under the Improved Conditions<sup>a</sup>

<sup>a</sup>Reaction conditions: **2** (0.30 mmol), **1a** (0.36 mmol) and **4k** (3 mol%) in TIPB (0.10 mL) at 120 °C for 16 h. Yields are the isolated yield based on **2**. Yields in parentheses refer to those of the recovered starting materials. <sup>b</sup> Octane (0.10 mL) was used instead of TIPB. <sup>c</sup>**2q** (0.15 mmol), **1a** (0.36 mmol), **4k** (6 mol%) were used and the reaction was repeated twice. <sup>d</sup> **2r** (0.60 mmol) and **1a** (0.30 mmol) were used; the yield is based on **1a**. See SI for details. <sup>e</sup>The reaction was repeated twice. See SI for details. <sup>f</sup> Isolated yield determined by the conversion of the product to the corresponding phenol.

regioselectivity by platinum catalysis currently remains elusive. Unlike iridium catalyst system, lack of the present platinum system in the activity toward  $C(sp^3)$ –H bond activation should be the key to the successful borylation of mesitylene derivatives.

Scheme 2. Gram-Scale Borylation of 3v



**Borylation of Fluorinated Arenes.** The introduction of a fluorine atom to the benzene ring had a profound effect on both the reactivity and selectivity of the platinum-catalyzed C-H borylation reaction. For example, the borylation of fluorobenzene (**5a**) proceeded efficiently even when the amount of the substrate was decreased to 5.0 equiv to give **6a** (eq 4). In contrast, the reaction of benzene under the same conditions resulted in insufficient yield of the borylated product (eq 5).



Notably, the C-H borylation of **5a** proceeded with a high level of *ortho* selectivity (73%), suggesting that fluorine atom exhibited an *ortho*-directing effect. When the borylation of fluorobenzene was performed with Ir/dtbpy catalyst, *meta*-borylated product was obtained as a major isomer (o/m/p ratio of 37:50:13). Fluorine has been reported to behave as a directing group in C-H borylation reactions involving rhodium-<sup>20</sup> and cobalt-<sup>21</sup> catalyzed systems, although the scope of this directing effect has not been extensively investigated. Takaya and Iwasawa reported a similar *ortho*-directing effect for fluorine when they investigated the C-H borylation of polyfluorinated benzene substrates with

platinum catalysts bearing a pincer ligand.<sup>8</sup> A similar activating effect was also observed for chloroarenes, as exemplified by the formation of **6c** and **6d** (Figure 6). Several 1,3-difluorobenzene derivatives were also borylated to give the 2-borylated products as the major isomers, although this position was the most congested among the C-H bonds available (**6e-6g**). Most notably, 1,2-difluorobenzene derivatives bearing a C-H bond *ortho* to a fluorine atom were selectively borylated at this position to form **6h** and **6i**, rather than being borylated at their much less hindered C-H bonds. The two difluorinated arenes **5j** and **5k** also showed high *ortho* selectivity to form the corresponding polysubstituted aryl boronic esters **6j** and **6k**. Although the isolated yields of these fluorinated phenylboronic esters were found to be modest because of their poor stability during chromatographic isolation, these borylated aromatic compounds. For example, **6j** was directly converted to the corresponding phenol without isolation. The observed directing effect probably results from the increased acidity of a C-H bond that locates at the *ortho* position of fluorine,<sup>22</sup> which facilitates the metalation process to occur at the *ortho* position of fluorine. The higher bond strength of carbon-metal bonds that are close to fluorine can also contribute to the observed *ortho* selectivity.<sup>23</sup>

Figure 6. Substrate Scope of Fluorinated Arenes<sup>a</sup>



<sup>a</sup>Reaction conditions: **1a** (0.30 mmol), **5** (5 equiv), **4g** (2 mol%), 80 °C, 20 h. Yields are based on **1a**. The regioisomeric ratios were determined by <sup>1</sup>H NMR spectroscopy or GC analysis. Yields in

parentheses refer to the NMR yields. <sup>b</sup>At 60 °C. <sup>c</sup>Isolated yield determined by the conversion of the product to the corresponding phenol.

**Borylation of 5-Membered Heteroarenes.** 5-Membered heteroarenes are generally considered to be highly reactive substrates for C-H borylation reactions.<sup>24</sup> A similar trend was also observed in the present platinum-catalyzed system. For example, *N*-methylindole **7a** was sufficiently reactive to undergo borylation in the presence of 2.0 equiv of **1a** and 2.0 mol% of platinum catalyst **4g** in heptane to afford the 2-borylated product **8a** in 87% isolated yield (Figure 7). Indoles bearing a variety of different substituents, including an ester (**7b**), ether (**7c**) and fluoride (**7d**) were also successfully borylated to give the corresponding 2-borylated indoles **8b-8d**, respectively. The exclusive formation of **8d** suggested that an indole C-H bond is more susceptible to borylation than a C-H bond *ortho* to a fluorine atom. Although pyrrole **7e** was also successfully borylated under our newly developed platinum-catalyzed conditions, the reaction needed to be conducted in the presence of a large excess of pyrrole to suppress the undesired diborylation reaction. Although 6-membered heteroarenes such as pyridine and quinoline were found to be unsuitable for this type of platinum catalysis, they were not detrimental to the C-H borylation of azoles, as evidenced by the successful formation of the borylated





<sup>a</sup>Reaction conditions: **7** (0.30 mmol), **1a** (1.2 equiv), **4g** (2 mol%), heptane (1.0 mL), 80 °C, 20 h. Yields are based on **7**. The regioisomeric ratios were determined by <sup>1</sup>H NMR spectroscopy or GC analysis. Yields in parentheses refer to the NMR yields. <sup>b</sup>**1a** (2.0 equiv) was used. <sup>c</sup>**4g** (4 mol%) was used. <sup>d</sup>N-Methylpyrrole **7e** (5.0 equiv to **1a**) was used. <sup>e</sup>Heteroarene was used as the solvent.

azaindole **8f**. Benzofuran **7g** was also borylated under these conditions, albeit with decreased reactivity and poor 2-/3-selectivity. Thiophenes and benzothiophenes served as good substrates for the platinum-catalyzed borylation and afforded the corresponding borylated products **8h-8k** in moderate to good yields.

**Mechanistic Considerations.** Several experiments were conducted to gain a deeper insight into the nature of the catalytically active platinum species involved in this C-H borylation reaction. It was previously proposed that the dvtms ligand in the platinum precursor decomposed prior to the formation of the catalytically active platinum species in hydrosilylation reactions involving Pt(NHC)(dvtms).<sup>9,25</sup> With this in mind, the author investigated the reaction of dvtms with **1a** in the presence of Pt(ICy)(dvtms) **4g** to examine the reactivity of dvtms under the catalytic conditions used in the current C-H borylation reaction. APCI-MS analysis revealed that the dvtms was completely consumed under the reaction conditions to give the corresponding tetraborylated product (eq 6). This observation clearly demonstrated that the dvtms ligand could dissociate from the platinum center via the diboration of its alkene moieties to generate a coordinatively unsaturated platinum species, which could subsequently react with **1a**. It has been reported that several platinum(0) complexes can react with diboron reagents to afford platinum(II) diboryl species. For example, platinum diboryl complexes bearing phosphine ligands have been isolated and characterized, and their catalytic activity towards the diboration of alkenes and alkynes have been reported in detail.<sup>7</sup>

It has also been reported that the Pt(NHC)(dvtms) complex, where NHC is I<sup>n</sup>Bu or a triazole-based carbene, can catalyze the diboration of alkynes and alkenes, most likely through the formation of a Pt(NHC)(Bpin)<sub>2</sub> species.<sup>6d,e</sup> It was also confirmed in our own hands that Pt(ICy)(dvtms) **4g** catalyzed the diboration of diphenylacetylene (eq 7). This observation suggested that **4g** could also generate a platinum diboryl species via its reaction with **1a**.

Scheme 2. Diboration of an Alkene and an Alkyne with Pt(ICy)(dvtms) Complex



Based on these experiments, the author proposed a possible reaction mechanism for the Pt(NHC)catalyzed borylation of arenes, which is shown in Scheme 3. The Pt(NHC)(dvtms) complex would be activated by its reaction with **1a** to give the platinum diboryl species **A** via the decomposition of the dvtms ligand. Complex **A** would then function as an active species to activate the C-H bond in the arenes. With regard to the subsequent C-H bond cleavage step, we considered two different mechanisms. The first of these mechanisms involve a  $\sigma$ -bond metathesis reaction between **A** and one of the Ar-H bonds of the substrate to form the Ar-Pt(II)-Bpin species **B**. The other mechanism would involve the oxidative addition of one of the C-H bonds to **A**, leading to the formation of the platinum(IV) species **B'**. It is noteworthy that  $\sigma$ -bond metathesis pathways have been proposed for rhodium- and iron-catalyzed C-H borylation reactions, based on DFT calculations,<sup>26</sup> whereas oxidative addition pathways have been proposed for iridium-catalyzed C-H borylation reactions.<sup>27</sup> However, there is currently no evidence to discriminate these two pathways. The subsequent reaction of **B** or **B'** with **1a** would result in the release of HBpin and ArBpin with the regeneration of **A**.<sup>28</sup>





The platinum-catalyzed borylation of benzene was monitored by GC and <sup>1</sup>H-NMR spectrometry and the results revealed that there was an induction period before the borylation started to occur. The decomposition of the dvtms ligand most likely occurred during the induction period to generate the catalytically active platinum species. Other experiments revealed that HBpin was generated during the borylation of benzene, and that this material remained unreacted (see experimental section for details). Initial rate experiments for the borylation of benzene revealed a large KIE value (4.8), which indicated that the cleavage of the C-H bond was most likely the turnover-limiting step of the catalytic cycle (Scheme 4). A similarly large KIE value (2.7) was also observed for the platinum-catalyzed borylation of *N*-methylindole under dilute conditions.

Scheme 4. KIE Experiments



**Synthetic Applications.** The current platinum-catalyzed borylation was found to be amenable to scale up, as exemplified by the gram-scale synthesis of **3k**, which could be used as a versatile building block for the introduction of different groups (Scheme 5). For example, the steric demand around the boron moiety in **3k** did not hamper its reactivity towards a standard palladium-catalyzed Suzuki-Miyaura cross-coupling reactions for the introduction of aryl (**9**) and alkyl (**10**) groups. Compound **3k** also afforded a hindered aniline **11** and a hindered phenol **12** via known amination<sup>29</sup> and oxidation methods, respectively.

Scheme 5. Gram-Scale Synthesis of 3k and Its Transformations



<sup>a</sup>Yield based on the recovered starting material.

Sequential C-H functionalization reactions were also evaluated in this study (Scheme 6). The rhodium-catalyzed silylation of *m*-xylene is known to occur at the least hindered 5-position to form 2y.<sup>30,31</sup> the subsequent borylation of 2y under our platinum-catalyzed conditions led to the activation of the C-H bond surrounded by two methyl groups to form 3y. The silyl and boryl groups in 3y could be converted to other functionalities in an orthogonal manner.<sup>30</sup> For example, the selective transformation of the boryl functionality in 3y to an aryl group was successfully accomplished under standard Suzuki-Miyaura reaction conditions whilst leaving the silyl group unchanged to form 13. Conversely, the silyl group in 3y was brominated without the loss of the Bpin group to generate 14. These examples further demonstrated the tolerance of our platinum catalyst towards steric demand, as well highlighting how it can be used in combination with existing methods to generate new synthetic strategy for the regioselective functionalization of hindered arenes.

Scheme 6. Sequential C-H Functionalization of *m*-Xylene and Divergent Functionalization of 3y



As demonstrated in Figure 4-7, the platinum catalysts developed in the current study afforded unique levels of regioselectivity that are different to those observed with other catalysts. The complementary regioselectivities of these different systems is best exemplified by the borylation of *p*-methylbenzoate **15** (Scheme 7). The use of the standard Ir/dtbpy catalyst led to a mixture of products, including the benzylic borylation product **16a** and the 3-borylated product **16b**.<sup>32</sup> Selective borylation at the 2-position was only achieved with an Ir/P(3,5-(CF<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>)<sub>3</sub> system, where the ester group of the substrate acted as an *ortho*-directing group to form **16c**.<sup>33</sup> Selective borylation at the benzylic position was possible with an iridium/phenanthroline system when silylborane was used as the borylating agent to deliver **16a**.<sup>32</sup> In contrast, the robust Pt catalyst developed in the current study was found to be insensitive to the steric hindrance suffered by the other systems to allow for the exclusive C-H borylation of **15** at its 3-position to provide **16b**.

Scheme 7. Comparison of Regioselectivity for the C-H Borylation of Methyl *p*-Methylbenzoate Using Different Catalysts



#### **2.3 Conclusion**

The author have developed a Pt/NHC complex-catalyzed C-H borylation of aromatic substrates. The use of a platinum complex in conjunction with NHC ligands, such as ICy and IPr, resulted in effective catalysts for the C-H borylation of a wide range of arenes and heteroarenes. Notably, this platinum system was found to be largely insensitive to steric hindrance, allowing for the borylation of the sterically hindered aromatic C-H bond in a series of mesitylene derivatives to give the corresponding phenylboronic ester derivatives bearing two *ortho* substituents. Further ligand screening determined that SIPr was as a suitable ligand for the borylation of sterically hindered arenes, whilst avoiding the need for a large excess of the substrate, thereby broadening the scope of this reaction. Another feature of this platinum catalyst system is that it exhibited a pronounced *ortho*-directing effect in the presence of fluorine substituents, providing facile access to a wide range of *ortho*-fluorophenylboronic esters directly from fluoroarenes. It is noteworthy that iridium catalyst system cannot be used in this way to achieve regioselective borylation of both types of substrates. It is therefore envisaged that the unique features of our platinum catalyst system for C-H borylation will allow for the development of new synthetic strategies for the regioselective functionalization of arenes, especially those with steric demand.

## 2.4 Experimental Section

## 2.4.1 General Information

<sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a JEOL ECS-400 spectrometer (JEOL, Tokyo, Japan) or VARIAN UNITY INOVA-600 spectrometer in either CDCl<sub>3</sub> with tetramethylsilane as an internal reference standard ( $\delta = 0.00$  ppm) or C<sub>6</sub>D<sub>6</sub> with a residual C<sub>6</sub>H<sub>6</sub> peak as an internal reference standard ( $\delta = 7.15$  ppm). NMR data have been reported as follows: chemical shift ( $\delta$ ) in ppm, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, and m = multiplet), coupling constant (*J*) in Hz, and integration. NOE experiments were conducted with Delta noe\_1d\_dpfgse pulse sequence and Delta noesy pulse sequence and detail experimental parameter was listed in the spectroscopic data section for each compound. Infrared spectra (IR) were obtained on a JASCO FT/IR-4000 spectrometer, and the absorptions have been reported in reciprocal centimeters with the following relative intensities: s (strong), m (medium), or w (weak). Mass spectra were obtained on a Shimadzu GCMS-QP 5000 or GCMS-QP 2010 instrument with an ionization voltage of 70 eV. High resolution mass spectra (HRMS) were obtained on a JEOL JMS-DX303 (EI) and a Bruker micrOTOF II (APCI). Analytical gas chromatography (GC) was carried out on Shimadzu GC-2014, equipped with a flame ionization detector. Melting points were determined using a Stanford Research Systems OptiMelt. Column chromatography was performed over SiO<sub>2</sub> (Silycycle Silica Flash F60 (230-400 mesh).

## 2.4.2 Materials

Pt<sub>2</sub>(dvtms)<sub>3</sub> (0.1 M, Aldrich), Pt(PPh<sub>3</sub>)<sub>4</sub> (TCI), H<sub>2</sub>PtCl<sub>6</sub>•nH<sub>2</sub>O (Furuya Metal), PCy<sub>3</sub> (Aldrich), dtbpy (Aldrich), [Ir(OMe)(cod)]<sub>2</sub> (TCI), HBpin (TCI) and B<sub>2</sub>pin<sub>2</sub> (TCI) were used as received. All arenes were obtained from commercial suppliers and used as received. *N*-Methylindole (TCI), *N*-methylpyrrole (Aldrich), 2-methoxythiophene (TCI), 2-methylthiophene (TCI), benzofuran (TCI), benzofuran (TCI), benzothiophene (TCI), and 5-chloro-3-methylbenzo[b]thiophene (TCI) were used as received. Other *N*-methylindoles used in this study were synthesized by the reaction of the corresponding indole with MeI according to the literature procedure.<sup>34</sup> (3,5-Dimethylphenyl)triethylsilane **2y** was synthesized according to the literature procedure<sup>35</sup> and all spectroscopic data were in agreement with the reported values.<sup>36</sup> 2-Chloro-5-methoxy-1,3-dimethylbenzene **2af** was synthesized according to the literature procedure.<sup>38</sup>

#### 1-Cyclopentyl-3,5-dimethylbenzene (2t, Figure 5).



To a dried two-necked flask equipped with a reflux condenser, Mg (264 mg, 11 mmol) and THF (10 mL) were added. I<sub>2</sub> chip was added to activate Mg. 3,5-Dimethylphenyl bromide (1.85 g, 10 mmol)

was added slowly, and the mixture was refluxed for 1 h. In another round-bottom flask, cyclopentyl bromide (1.49 g, 10 mmol), FeCl<sub>3</sub> (81 mg, 0.5 mmol), TMEDA (87 mg, 0.75 mmol) and THF (10 mL) were added. The Grignard solution was transferred to a THF solution of cyclopentyl bromide at 0 °C by cannulation. After the addition of Grignard reagent, the mixture was warmed to ambient temperature and the resulting mixture was stirred overnight. To the resulting mixture, H<sub>2</sub>O was added and extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with brine and dried over MgSO<sub>4</sub>. Concentrated mixture was filtered through a pad of silica gel (hexane as eluent). To remove homo-coupled biaryl byproduct, the material was further purified by GPC to give the title product in pure form (colorless oil, 460 mg, 26%).

Rf 0.86 (hexane/EtOAc = 5/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.54-1.88 (m, 6H), 1.92-2.12 (m, 2H), 2.29 (s, 6H), 2.81-3.00 (m, 1H), 6.82 (s, 1H), 6.86 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  21.3, 25.6, 34.6, 45.8, 124.9, 127.3, 137.6, 146.5. IR (ATR): 2950 s, 2915 m, 2866 m, 1604 m, 1452 m, 1375 w, 1036 w, 936 w, 888 w, 841 s, 702 s. MS m/z (% relative intensity): 174 (M<sup>+</sup>, 100), 159 (93), 145 (75), 133 (68), 132 (41), 131 (44), 119 (50), 117 (37), 115 (30), 106 (32), 105 (64), 91 (43). HRMS (EI): Calcd for C<sub>13</sub>H<sub>18</sub> 174.1409, Found 174.1408.

#### sec-Butyl 3,5-dimethylbenzoate (2u, Figure 5).



To a dried round-bottom flask, 3,5-dimethylbenzoic acid (450 mg, 3.0 mmol), DCC (680 mg, 3.3 mmol), DMAP (73 mg, 0.6 mmol), 2-propanol (490  $\mu$ L, 5.4 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (10 mL) were added. The mixture was stirred at ambient temperature overnight. To the resulting reaction mixture, H<sub>2</sub>O was added and extracted with CH<sub>2</sub>Cl<sub>2</sub> three times. Combined organic extracts were washed with brine and dried over MgSO<sub>4</sub>. Concentrated mixture was purified by flash column chromatography (hexane/EtOAc = 10/1) to give the title compound (clear oil, 404 mg, 65%).

Rf 0.71 (hexane/EtOAc = 5/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.97 (t, *J* = 7.6 Hz, 3H), 1.33 (d, *J* = 6.4 Hz, 3H), 1.60-1.82 (m, 2H), 2.36 (s, 6H), 5.08 (sextet, *J* = 6.4 Hz, 1H), 7.17 (s, 1H), 7.65 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  9.9, 19.6, 21.2, 29.0, 72.6, 127.2, 130.8, 134.3, 137.9, 166.6. IR (ATR): 2972 w, 2934 w, 1712 s, 1608 w, 1455 w, 1379 w, 1354 w, 1307 m, 1215 s, 1111 m, 1095 w, 1000 w, 929 w, 867 w, 767 m, 679 w. MS m/z (% relative intensity): 206 (M<sup>+</sup>, 20), 151(25), 150 (50), 134 (13), 133 (100), 107 (6), 106 (6), 105 (41), 103 (8), 91 (12), 79 (10), 77 (10). HRMS (EI): Calcd for C<sub>13</sub>H<sub>18</sub>O<sub>2</sub> 206.1307, Found 206.1305.

#### (1R,2S,5R)-2-Isopropyl-5-methylcyclohexyl 3,5-dimethylbenzoate (2v, Figure 5).



To a dried round-bottom flask, 3,5-dimethylbenzoic acid (450 mg, 3.0 mmol), DCC (680 mg, 3.3 mmol), DMAP (73 mg, 0.6 mmol), (-)-menthol (843 mg, 5.4 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (10 mL) were added. The mixture was stirred at ambient temperature overnight. To the resulting reaction mixture, H<sub>2</sub>O was added and extracted with CH<sub>2</sub>Cl<sub>2</sub> twice. Combined organic extracts were washed with water and brine and then dried over MgSO<sub>4</sub>. Concentrated mixture was purified by flash column chromatography (hexane/EtOAc = 100/1 to 12/1) to give the title compound (colorless viscous oil, 658 mg, 76%). Rf 0.73 (hexane/EtOAc = 5/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.79 (d, *J* = 6.9 Hz, 3H), 0.87- 0.97 (m, 1H), 0.92 (dd, *J* = 6.4, 2.8 Hz, 6H), 1.02-1.22 (m, 2H), 1.45-1.64 (m, 2H), 1.67-1.80 (m, 2H), 1.96 (quintet-d, *J* = 6.9, 2.7 Hz, 1H), 2.03-2.16 (m, 1H), 2.36 (s, 6H), 4.93 (td, *J* = 10.9, 4.4 Hz, 1H), 7.17 (s, 1H), 7.65 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  16.4, 20.8, 21.2, 22.0, 23.5, 26.4, 31.4, 34.3, 41.0, 47.2, 74.5, 127.2, 130.7, 134.3, 137.9, 166.4. IR (ATR): 2953 w, 2923 w, 2868 w, 1712 s, 1608 w, 1454 w, 1382 w, 1305 m, 1214 s, 1116 m, 1010 w, 982 w, 965 w, 865 w, 766 m, 678 w. MS m/z (% relative intensity): 151 (11), 139 (9), 138 (82), 134 (10), 133 (100), 105 (34), 96 (19), 95 (75), 94 (11), 82 (20), 81 (41), 79 (12). HRMS (EI): Calcd for C<sub>19</sub>H<sub>28</sub>O<sub>2</sub> 288.2089, Found 288.2091.

(1S,2R,4S)-1,7,7-Trimethylbicyclo[2.2.1]heptan-2-yl 3,5-dimethylbenzoate (2w, Figure 5).



To a dried round-bottom flask, 3,5-dimethylbenzoic acid (750 mg, 5.0 mmol), DCC (1.1 g, 5.5 mmol), DMAP (122 mg, 1.0 mmol), (-)-borneol (1.4 g, 9.0 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (17 mL) were added. The mixture was stirred at ambient temperature overnight. To the resulting reaction mixture, H<sub>2</sub>O was added and extracted with CH<sub>2</sub>Cl<sub>2</sub> twice. Combined organic extracts were washed with brine and then dried over MgSO<sub>4</sub>. Concentrated mixture was purified by flash column chromatography (hexane/EtOAc = 100/1 to 20/1) to give the title compound (colorless oil, 816 mg, 57%).

Rf 0.63 (hexane/EtOAc = 5/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.91 (s, 6H), 0.97 (s, 3H), 1.11 (dd, *J* = 14.0, 3.4 Hz, 1H), 1.28-1.48 (m, 2H), 1.73 (t, *J* = 4.6 Hz, 1H), 1.75-1.87 (m, 1H), 2.06-2.20 (m, 1H), 2.37 (s, 6H), 2.41-2.56 (m, 1H), 5.10 (ddd, *J* = 9.8, 3.4, 2.3 Hz, 1H), 7.18 (s, 1H), 7.66 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  13.6, 18.9, 19.7, 21.2, 27.4, 28.1, 36.9, 45.0, 47.8, 49.0, 80.3, 127.2, 130.8, 134.4, 137.9, 167.1. IR (ATR): 2952 w, 2878 w, 1714 s, 1608 w, 1453 w,

1380 w, 1313 s, 1215 s, 1116 m, 1022 m, 979 w, 865 w, 766 m, 678 w. MS m/z (% relative intensity): 286 (M<sup>+</sup>, 9), 136 (17), 133 (100), 121 (5), 105 (17), 93 (9), 79 (6). HRMS (EI): Calcd for C<sub>19</sub>H<sub>26</sub>O<sub>2</sub> 286.1933, Found 286.1934.

#### 3-Chloro-1,2,4,5-tetramethylbenzene (2x, Figure 5).



The title compound was synthesized according to the literature procedure.<sup>39</sup>

In a round-bottom flask, 1,2,4,5-tetramethylbenzene (2.68 g, 20 mmol) and NCS (1.34 g, 10 mmol) was dissolved in MeCN (20 mL) to give a heterogeneous mixture. TMSCl (130  $\mu$ L, 1.0 mmol) was added dropwise, and the heterogeneous mixture changed to homogeneous with heat generation. The mixture was stirred at ambient temperature overnight. To the resulting mixture, H<sub>2</sub>O was added and extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with 1M NaOH aq. and brine, and dried over MgSO<sub>4</sub>. Concentrated mixture was filtered through a pad of silica gel (hexane as eluent) to remove benzyl chlorides as byproducts. To remove unreacted starting material, the mixture was further purified by GPC to give the title product in pure form (white solid, 300 mg, 18%).

Rf 0.86 (hexane/EtOAc = 5/1). White solid. Mp = 49 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  2.24 (s, 6H), 2.30 (s, 6H), 6.85 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  16.8, 20.5, 129.4, 131.9, 134.6, 135.1. IR (ATR): 3007 w, 2969 w, 2923 w, 2862 w, 1554 w, 1473 m, 1455 m, 1391 m, 1273 w, 1188 m, 1000 s, 865 m, 822 s. MS m/z (% relative intensity): 170 (12), 168 (M<sup>+</sup>, 35), 153 (17), 133 (100), 117 (12), 115 (15), 105 (10), 91 (14). HRMS (EI): Calcd for C<sub>10</sub>H<sub>13</sub>Cl 168.0706, Found 168.0704.

#### 3,3,3-Trifluoropropyl 3,5-dimethylbenzoate (2z, Figure 5).



To a dried round-bottom flask, 3,5-dimethylbenzoic acid (600 mg, 4.0 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (14 mL) and a few drops of DMF was added. Oxalyl chloride (370  $\mu$ L, 4.4 mmol) was added dropwise resulting in gas evolution. The mixture was stirred for 30 min. 3,3,3-Trifluoropropan-1-ol (460 mg, 4.0 mmol), DMAP (10 mg, 0.080 mmol) and NEt<sub>3</sub> (1.1 mL, 8.0 mmol) were dissolved in CH<sub>2</sub>Cl<sub>2</sub> (2.0 mL) and the CH<sub>2</sub>Cl<sub>2</sub> solution was added to the reaction mixture. The mixture was stirred at ambient temperature overnight. To the resulting mixture, H<sub>2</sub>O was added and extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with 1M NaOH aq., 1M HCl aq. and brine and dried over MgSO<sub>4</sub>. Concentrated mixture was purified by flash column chromatography (hexane/EtOAc = 100/1 to 20/1) to give the title compound (colorless oil, 703 mg, 71%).

Rf 0.71 (hexane/EtOAc = 5/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  2.36 (s, 6H), 2.53-2.67 (qt, *J* = 10.5, 6.4 Hz, 2H), 4.52 (t, *J* = 6.4 Hz, 2H), 7.20 (s, 1H), 7.65 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  21.1, 33.47 (q, *J* = 29 Hz), 57.5 (q, *J* = 3.8 Hz), 125.8 (q, *J* = 277 Hz), 127.3, 129.4, 134.9, 138.1, 166.5. IR (ATR): 1721 s, 1609 w, 1387 w, 1347 w, 1311 m, 1256 m, 1209 s, 1154 s, 1134 s, 1117 s, 1067 m, 1008 m, 867 w, 844 w, 765 m. MS m/z (% relative intensity): 246 (M<sup>+</sup>, 100), 134 (12), 133 (100), 105 (43), 79 (12), 77 (14). HRMS (EI): Calcd for C<sub>12</sub>H<sub>13</sub>F<sub>3</sub>O<sub>2</sub> 246.0868, Found 246.0865.

#### 3-Chloropropyl 3,5-dimethylbenzoate (2aa, Figure 5).



The title compound was synthesized using the same procedure for **2z** except that 3-chloropropanol was used instead of 3,3,3-trifluoropropan-1-ol on a 10 mmol scale.

Rf 0.57 (hexane/EtOAc = 5/1). Colorless oil (1.69 g, 75%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  2.36 (quintet, J = 6.2 Hz, 2H), 2.36 (s, 6H), 3.70 (t, J = 6.4 Hz, 2H), 4.46 (t, J = 6.2 Hz, 2H), 7.19 (s, 1H), 7.64 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  21.1, 31.7, 41.3, 61.5, 127.2, 129.8, 134.7, 138.0, 166.7. IR (ATR): 2965 w, 2919 w, 1716 s, 1608 w, 1450 w, 1384 w, 1308 s, 1209 s, 1166 w,1152 w, 1116 m, 1061 w, 1020 m, 945 w, 866 s, 765 s, 678 w, 656 w. MS m/z (% relative intensity): 228 (6), 226 (M<sup>+</sup>, 19), 151 (8), 150 (74), 133 (100), 105 (54), 91 (9), 79 (13), 77 (14). HRMS (EI): Calcd for C<sub>12</sub>H<sub>15</sub>ClO<sub>2</sub> 226.0761, Found 226.0759.

#### tert-Butyl(3,5-dimethylphenoxy)dimethylsilane (2ab, Figure 5).



In a dried round-bottom flask, 3,5-dimethylphenol (1.2 g, 10 mmol) was dissolved in  $CH_2Cl_2$  (15 mL). TBDMSCl (1.9 g, 12.5 mmol), DMAP (24 mg, 0.2 mmol) and NEt<sub>3</sub> (2.6 mL, 19 mmol) were then added. The mixture was stirred at ambient temperature overnight. To the resulting mixture, H<sub>2</sub>O was added and extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with brine and dried over MgSO<sub>4</sub>. Concentrated mixture was purified by flash column chromatography (hexane/EtOAc = 200/1) to give the title compound (clear oil, 2.2 g, 93%).

Rf 0.78 (hexane/EtOAc = 5/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.18 (s, 6H), 0.98 (s, 9H), 2.25 (s, 6H), 6.46 (s, 2H), 6.59 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  -4.42, 18.2, 21.3, 25.7, 117.8, 123.0, 139.0, 155.5. IR (ATR): 2955 w, 2929 w, 2857 w, 1592 w, 1468 w, 1316 m, 1253 w, 1159 m, 1038 w, 961 w, 835 s, 778 s. MS m/z (% relative intensity): 237 (5), 236 (M<sup>+</sup>, 20), 180 (20),

179 (100), 163 (5), 149 (9), 105 (33), 75 (8). HRMS (EI): Calcd for C<sub>14</sub>H<sub>24</sub>OSi 236.1596, Found 236.1593.

#### Mesityl(piperidin-1-yl)methanone (2ac, Figure 5).



To a dried round-bottom flask, 2,4,6-tetramethylbenzoic acid (1.64 g, 10 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (14 mL) and a few drops of DMF was added. Oxalyl chloride (930  $\mu$ L, 11 mmol) was added dropwise resulting in gas evolution. The mixture was stirred for 1 h. The mixture was cooled to 0 °C and then piperidine (990  $\mu$ L, 10 mmol) and NEt<sub>3</sub> (1.1 mL, 8.0 mmol) were added. The mixture was warmed to ambient temperature and stirred overnight. To the resulting mixture, H<sub>2</sub>O was added and extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with 1M NaOH aq., 1M HCl aq. and brine and then dried over MgSO<sub>4</sub>. Concentrated mixture was treated with activated charcoal and removal of the charcoal by filtration gave the pure title compound. (pale-yellow oil, 2.24 g, 97%). Rf 0.16 (hexane/EtOAc = 5/1). Pale-yellow oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.38-1.55 (m, 2H), 1.57-1.75 (m, 4H), 2.22 (s, 6H), 2.27 (s, 3H), 3.08-3.22 (m, 2H), 3.69-3.85 (m, 2H), 6.84 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  18.9, 21.0, 24.5, 25.7, 26.6, 41.9, 47.0, 128.1, 133.4, 133.7, 137.7, 169.6. IR (ATR): 2933 w, 2854 w, 1628 s, 1438 s, 1375 w, 1271 s, 1240 m, 1184 w, 1098 m, 1027 w, 998 m,879 w, 851 m, 750 w. MS m/z (% relative intensity): 231 (M<sup>+</sup>, 25), 216 (31), 148 (11), 147 (100), 146 (32), 119 (19), 91 (12). HRMS (EI): Calcd for C<sub>15</sub>H<sub>21</sub>NO 231.1623, Found 231.1619.

#### *N*-Mesityl-*N*-methylacetamide (3ad, Figure 5).



In a dried round-bottom flask, *N*-mesitylacetamide (530 mg, 3.0 mmol) was dissolved in DMF (6.0 mL). NaH (60% in mineral oil, 150 mg, 3.6 mmol) was added portionwise. The mixture was stirred overnight. To the resulting mixture, H<sub>2</sub>O was added and extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with brine and dried over MgSO<sub>4</sub>. Concentrated mixture was purified by flash column chromatography (hexane/EtOAc = 10/1 to 5/1) to give the title compound (colorless oil, 354 mg, 62%).

Rf 0.45 (hexane/EtOAc = 1/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.73 (s, 3H), 2.16 (s, 6H), 2.30 (s, 3H), 3.11 (s, 3H), 6.93 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  17.5, 20.9, 21.2, 34.3, 129.5, 135.1, 137.8, 139.2, 171.0. IR (ATR): 2921 w, 1658 s, 1483 m, 1424 m, 1374 s, 1348 m, 1308 w, 1282 w,1133 w, 1078 w, 1033 w, 972 w, 854 w. MS m/z (% relative intensity): 191 (M<sup>+</sup>, 26), 176

(11), 148 (16), 134 (19), 91 (11), 56 (100). HRMS (EI): Calcd for  $C_{12}H_{17}NO$  191.1310, Found 191.1309.

Ethyl 4-(mesityloxy)butanoate (2ae, Figure 5).



In a dried round-bottom flask, NaH (60% in mineral oil, 440 mg, 11 mmol) was dispersed in DMF (20 mL). 2,4,6-Trimethylphenol (1.36 g, 10 mmol) was added slowly, and the mixture was stirred at ambient temperature for 1 h. Ethyl 4-bromobutanoate (2.34 g, 12 mmol) was slowly added and stirred for another 5.5 h. To the resulting mixture, H<sub>2</sub>O was added and extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with brine and dried over MgSO<sub>4</sub>. Concentrated mixture was dried in vacuo at 160 °C to remove unreacted alkyl bromide and further purified by flash column chromatography (hexane/EtOAc = 100/1 to 25/1) to give the title compound (colorless oil, 1.42 g, 57%).

Rf 0.67 (hexane/EtOAc = 5/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.27 (t, *J* = 7.1 Hz, 3H), 2.11 (quintet, *J* = 7.1 Hz, 2H), 2.22 (s, 6H), 2.23 (s, 3H), 2.59 (t, *J* = 7.3 Hz, 2H), 3.76 (t, *J* = 6.2 Hz, 2H), 4.15 (q, *J* = 7.2 Hz, 2H), 6.81 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  14.2, 16.1, 20.6, 25.7, 30.9, 60.4, 70.7, 129.3, 130.4, 133.0, 153.4, 173.4. IR (ATR): 2923 w, 1733 s, 1482 m, 1443 w, 1373 m, 1308 w, 1214 s, 1173 s, 1147 s, 1096 w, 1029 m, 922 w, 853 m. MS m/z (% relative intensity): 205 (16), 136 (13), 121 (18), 115 (79), 91 (16), 87 (100). HRMS (EI): Calcd for C<sub>15</sub>H<sub>22</sub>O<sub>3</sub> 250.1569, Found 250.1565.

#### 2.4.3 Synthesis of Pt(NHC)(dvtms) Complexes

Pt(NHC)(dvtms) complexes were synthesized by using slightly modified versions of the reported procedures, <sup>25a,40</sup> as follows.



**Pt(ICy)(dvtms)** (**4g).** H<sub>2</sub>PtCl<sub>6</sub>•nH<sub>2</sub>O (1.8 g, 3.4 mmol, estimated as hexahydrate), NaHCO<sub>3</sub> (3.5 g, 42 mmol), dvtms (3.5 g, 19 mmol) and degassed EtOH (10 mL) were added to an oven-dried round bottom flask. The mixture was heated at 80 °C for 1 h; the color of the solution changed from red to yellow. The mixture was stirred at room temperature (rt) for another 3 h. The resulting mixture was filtered through a Celite pad, and the pad was quickly washed with EtOH. The filtrate was concentrated in vacuo, and degassed benzene (3 mL) was added. The solution was again filtered through a Celite pad, and the pad was quickly washed with enconcentrated to give crude  $Pt_2(dvtms)_3$  as a yellow oil. ICy•HCl (1.1 g, 4.1 mmol, 1.2 equiv), <sup>*t*</sup>BuOK (0.68 g, 1.8 equiv) and degassed benzene

(10 mL) were added to the crude mixture, and the mixture was stirred for 3 h at rt. The resulting mixture was filtered through a pad of silicagel/Celite (1:1), and the pad was washed with hexane/Et<sub>2</sub>O (8:2, 100 mL). The filtrate was concentrated in vacuo to yield a pale-yellow solid, which was recrystallized from <sup>*i*</sup>PrOH/Et<sub>2</sub>O to give Pt(ICy)(dvtms) as a white solid (1.64 g, 79% based on H<sub>2</sub>PtCl<sub>6</sub>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  -0.25 (s, 6H), 0.33 (s, 6H), 1.09-1.33 (m, 6H), 1.47 (q, *J* = 12.3 Hz, 4H), 1.66 (d, *J* = 12.4 Hz, 2H), 1.73-1.98 (m, 12H), 2.17 (d, *J* = 10.5 Hz, <sup>2</sup>*J*<sub>Pt-H</sub> = 54 Hz, 2H), 4.21 (tt, *J* = 3.9 Hz, 12.4 Hz, 1H), 4.32 (tt, *J* = 3.9 Hz, 12.4 Hz, 1H), 7.01 (d, *J* = 1.3 Hz, <sup>4</sup>*J*<sub>Pt-H</sub> = 11.5 Hz, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  -1.89, 1.44, 25.3, 25.4, 33.4 (<sup>1</sup>*J*<sub>Pt-C</sub> = 119 Hz), 33.9, 40.3 (<sup>1</sup>*J*<sub>Pt-C</sub> = 158 Hz), 58.2 (<sup>3</sup>*J*<sub>Pt-C</sub> = 44 Hz), 58.4 (<sup>3</sup>*J*<sub>Pt-C</sub> = 42 Hz), 117.16 (<sup>3</sup>*J*<sub>Pt-C</sub> = 37 Hz), 117.21 (<sup>3</sup>*J*<sub>Pt-C</sub> = 37 Hz), 179.8 (<sup>1</sup>*J*<sub>Pt-C</sub> = 1390 Hz). HRMS (FAB): Calcd for C<sub>23</sub>H<sub>42</sub>N<sub>2</sub>OPtSi<sub>2</sub> 613.2483, Found 613.2501. The spectroscopic data were in agreement with the reported values.<sup>25a</sup>

Other Pt(NHC)(dvtms) complexes were synthesized by the same procedure using the corresponding NHC•HCl [IMes•HCl (TCI), IPr•HCl (TCI), I'Bu•HCl<sup>41</sup>, SICy•HCl<sup>42</sup>, BICy•HCl (Aldrich), IMe•HCl (TCI)] or NHC•HBF<sub>4</sub> [I(<sup>*i*</sup>Pr)•HBF<sub>4</sub> (TCI), I(Ad)•HBF<sub>4</sub><sup>43</sup>]. The spectroscopic date for **4a**, **4b**, **4c**, **4e**, **4f** and **4k** were in agreement with the reported values.(40-2)

**Pt**(**I**'**Pr**)(**dvtms**) (**4d**). Synthesis of the title compound was conducted on a 1.5 mmol scale based on H<sub>2</sub>PtCl<sub>6</sub>•nH<sub>2</sub>O to give 548 mg of Pt(I<sup>*i*</sup>Pr)(dvtms) as white solid (69%) <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ -0.29 (s, 6H), 0.33 (s, 6H), 1.30 (t, *J* = 6.9 Hz, 12H), 1.72-1.98 (m, 4H), 2.20 (d, *J* = 10 Hz, <sup>2</sup>*J*<sub>Pt-H</sub> = 54 Hz, 2H), 4.60-4.74 (m, 2H), 7.05(s, <sup>4</sup>*J*<sub>Pt-H</sub> = 12Hz, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ -2.21, 1.45, 23.1, 23.2, 33.6 (<sup>1</sup>*J*<sub>Pt-C</sub> = 119 Hz), 40.4 (<sup>1</sup>*J*<sub>Pt-C</sub> = 157 Hz), 50.9 (<sup>3</sup>*J*<sub>Pt-C</sub> = 46 Hz), 51.1 (<sup>3</sup>*J*<sub>Pt-C</sub> = 44 Hz), 116.8 (<sup>3</sup>*J*<sub>Pt-C</sub> = 37 Hz), 179.7, (<sup>1</sup>*J*<sub>Pt-C</sub> = 1370 Hz).

IR (ATR): 295w, 1424 w, 1368 w, 1297 w, 1246 m, 1213 m, 1171 m, 1131 w, 1002 w, 968 s, 901 w, 836 s, 781 s, 721 m, 707 m, 684 m. HRMS (FAB): Calcd for  $C_{17}H_{34}N_2OPtSi_2$  533.1857, Found 533.1851.

Pt(SICy)(dvtms) (4h). Synthesis of the title compound was conducted on a 1.4 mmol scale based on H<sub>2</sub>PtCl<sub>6</sub>•nH<sub>2</sub>O to give 583 mg of Pt(SICy)(dvtms) as a white solid (66%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ -0.28 (s, 6H), 0.32 (s, 6H), 0.94-1.05 (m, 2H), 1.11-1.23 (m, 4H), 1.30-1.43 (m, 4H), 1.54-1.74 (m, 12H), 1.83 (d, J = 13 Hz,  ${}^{2}J_{Pt-H} = 54$  Hz, 2H), 2.17 (d, J = 11 Hz,  ${}^{2}J_{Pt-H} = 54$  Hz, 2H), 3.56 (s, 4H), 3.88 (tt, J = 12, 3.2 Hz, 1H), 3.99 (tt, J = 12, 3.2 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ -2.03, 1.41, 25.25, 25.33, 25.4, 30.8, 30.9, 33.0 ( ${}^{1}J_{Pt-C} = 113$  Hz), 39.9 ( ${}^{1}J_{Pt-C} = 160$  Hz), 42.9 ( ${}^{3}J_{Pt-C} = 43$  Hz), 43.0 ( ${}^{3}J_{Pt-C} = 43$  Hz), 56.9 ( ${}^{3}J_{Pt-C} = 46$  Hz), 57.2 ( ${}^{3}J_{Pt-C} = 44$  Hz), 209.9 ( ${}^{3}J_{Pt-C} = 1330$  Hz). IR (ATR): 2934 m, 2855, w, 1482 m, 1440 m, 1290 m, 1271 m, 1237 s, 1168 m, 990 s, 893 w, 857 m, 829 s, 771 s, 703 m, 673 m. HRMS (FAB): Calcd for C<sub>23</sub>H<sub>44</sub>N<sub>2</sub>OPtSi<sub>2</sub> 615.2640, Found 615.2647.

Pt(BICy)(dvtms) (4i). Synthesis of the title compound was conducted on a 1.3 mmol scale based on H<sub>2</sub>PtCl<sub>6</sub>•nH<sub>2</sub>O to give 224 mg of Pt(BICy)(dvtms) as white solid (26%). The compound was further

purified by flash column chromatography over silica gel (eluting with hexane/AcOEt = 100/1) after the recrystallization from <sup>*i*</sup>PrOH.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ -0.21 (s, 6H), 0.36 (s, 6H), 1.20-1.38 (m, 7H), 1.65-2.03 (m, 13H), 2.09-2.34 (m, 6H), 4.79 (tt, J = 13, 3.4 Hz, 1H), 4.88 (tt, J = 13, 3.8 Hz, 1H), 7.15-7.19 (m, 2H), 7.56-7.62 (m, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ -1.86, 1.40, 25.40, 25.43, 25.5, 25.8, 30.6, 30.7, 34.4 (<sup>1</sup>*J*<sub>Pt-C</sub> = 115 Hz), 40.7 (<sup>1</sup>*J*<sub>Pt-C</sub> = 157 Hz), 59.7 (<sup>3</sup>*J*<sub>Pt-C</sub> = 46 Hz), 60.0 (<sup>3</sup>*J*<sub>Pt-C</sub> = 46 Hz), 111.8, 111.9, 120.9, 121.0, 134.3, 134.5, 196.3 (<sup>1</sup>*J*<sub>Pt-C</sub> = 1410 Hz). IR (ATR): 2929 w, 2857 w, 1476 w, 1450 w, 1409 w, 1374 m, 1341 w, 1293 m, 1239 m, 1168 w, 1054 w, 997 s, 909 w, 858 w, 834 s, 774 s, 736 s, 701 w. HRMS (FAB): Calcd for C<sub>27</sub>H<sub>44</sub>N<sub>2</sub>OPtSi<sub>2</sub> 663.2640, Found 663.2624.

## Pt(IPr')(dvtms) (4j)

IPr'•HCl was synthesized described below.



2,4,6-triisopropyllaniine<sup>44</sup> (950 mg, 4.33 mmol), and aqueous solution of glyoxal (8.8 M, 0.25 mL, 2.2 mmol) were dissolved in THF (10 mL) and stirred for 14 h at room temperature. To the resulting mixture, EtOAc and water were added. The organic layer was separated and washed with brine and water, and dried over MgSO<sub>4</sub>. The organic layer was concentrated *in vacuo* to give a diimine as a yellow solid (936 mg, ca. 94%). The diimine was used for the subsequent reaction without further purification.

Paraformaldehyde (137 mg, 2.36 mmol), diimine (920 mg, 2.00 mmol) and EtOAc (50 mL) were added to a 100 mL three-necked flask with a reflux condenser under N<sub>2</sub>. The mixture was refluxed for 1 h. TMSCl (0.30 mL, 260 mg, 2.39 mmol) was added to the reaction flask and the mixture was refluxed overnight. To the resulting mixture, EtOAc and water were added. The organic layer was separated and washed with brine and water and dried over MgSO<sub>4</sub>. The organic layer was concentrated *in vacuo* to give IPr'•HCl as a white solid (531 mg, 52%).

Mp = 237 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.19 (d, *J* = 6.9 Hz, 12H), 1.26 (d, *J* = 6.9 Hz, 12H), 1.28 (d, *J* = 6.9 Hz, 12H), 2.40 (septet, *J* = 6.9 Hz, 4H), 2.96 (d, *J* = 6.9 Hz, 2H), 7.14 (s, 4H), 8.16 (s, 2H), 9.43 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  23.7, 23.8, 24.7, 29.1, 34.4, 122.6, 127.2, 127.5, 137.7, 144.6, 153.0. IR (ATR): 2962 s, 2869 m, 2734 w, 1603 w, 1527 s, 1466 s, 1385 w, 1365 m, 1321 m, 1220 m, 1185 w, 1129 w, 1105 w, 1051 m, 957 m, 878 s, 767 s, 689 m. HRMS (FAB): Calcd for [M-HCl]<sup>+</sup> C<sub>33</sub>H<sub>48</sub>N<sub>2</sub> 472.3817, Found 472.3818.



Synthesis of the title Pt complex was conducted with the same procedure for **4g** except that IPr'•HCl was used instead of IPr<sup>Me</sup>•HCl on 0.12 mmol scale based on H<sub>2</sub>PtCl<sub>6</sub>•nH<sub>2</sub>O to give 58 mg of Pt(IPr')(dvtms) as a white solid (59%).

 $Mp > 210 \text{ °C (decomp.). }^{1}\text{H NMR (CDCl}_{3}, 399.78 \text{ MHz}): \delta -0.76 \text{ (s,} 6H), 0.13 (s, 6H), 1.12 (d,$ *J*= 6.9 Hz, 12H), 1.23 (d,*J*= 6.9 Hz, 12H), 1.24 (d,*J*= 6.9 Hz, 12H), 1.28-1.59 (m, 4H), 1.69 (d,*J* $= 11.5 Hz, <math>{}^{3}J_{\text{Pt-C}} = 27 \text{ Hz}, 2H$ ), 2.88 (septet, *J* = 6.9 Hz, 2H), 2.95 (septet, *J* = 6.9 Hz, 4H), 6.99 (s, 4H), 7.18 (s,  ${}^{5}J_{\text{Pt-C}} = 4.6 \text{ Hz}, 2H$ ).  ${}^{13}\text{C NMR}$  (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  -2.17, 1.60, 22.5, 24.0, 25.9, 28.3, 34.3, 35.1 ( ${}^{1}J_{\text{Pt-C}} = 59 \text{ Hz}$ ), 41.6 ( ${}^{1}J_{\text{Pt-C}} = 83 \text{ Hz}$ ), 121.3, 123.9 ( ${}^{3}J_{\text{Pt-C}} = 21 \text{ Hz}$ ), 134.4, 145.2, 149.6, 186.0. IR (ATR): 2960 m, 1471 w, 1401 w, 1383 w, 1364 w, 1317 m, 1243 m, 1179 w, 1130 w, 1052 w, 990 m, 949 w, 909 w, 860 m, 836 s, 781 s, 731 m, 699 m. HRMS (FAB): Calcd for C<sub>41</sub>H<sub>66</sub>N<sub>2</sub>OPtSi<sub>2</sub> 853.4361, Found 853.4359.

## Pt(IPr<sup>Me</sup>)(dvtms) (4l).

IPr<sup>Me</sup>•HCl was synthesized according to the literature procedure<sup>45</sup> with a slight modification.



To a dried round bottom flask, 2,6-diisopropylaniline (6.0 mL, 32 mmol) and diacetyl (1.4 mL, 16 mmol) and HCOOH (10 drops) were added. The yellow homogeneous mixture was sonicated for 10 min then the yellow solution turned into cloudy yellow mixture. MeOH (20 mL) were added to give homogeneous solution again and the solution was left for 1 h without stirring. Additional MeOH (10 mL) was added and the mixture was stirred vigorously. Soon yellow precipitate appeared and the mixture was stirred overnight. Filtration of the precipitate and washing with cold methanol gave the pure target diimine (2.2 g, 34%).

All spectroscopic data were in agreement with the reported values in the literature.<sup>45</sup>

The formation of the imidazolium ring was conducted according to the literature method.<sup>45</sup>

Synthesis of the title Pt complex was conducted on a 0.22 mmol scale based on H<sub>2</sub>PtCl<sub>6</sub>•nH<sub>2</sub>O to give 113 mg of Pt(IPr<sup>Me</sup>)(dvtms) as white solid (64%)



IMp > 205 °C (decomp.). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  -0.77 (s, 6H), 0.11 (s, 6H), 1.12 (d, *J* = 6.9 Hz, 12H), 1.23 (d, *J* = 6.9 Hz, 12H), 1.25-1.65 (m, 6H), 2.01 (s, 6H), 2.92 (septet, *J* = 6.9 Hz, 4H), 7.17 (d, *J* = 7.8 Hz, 4H), 7.34 (t, *J* = 7.8 Hz, 2H).

<sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ -2.39, 1.64, 11.0, 23.5, 24.9, 28.1, 35.4 ( ${}^{1}J_{Pt-C}$  = 59 Hz), 42.6 ( ${}^{1}J_{Pt-C}$  = 84 Hz), 123.9, 126.7, 129.1, 135.2, 146.2, 182.9

 $(^{1}J_{Pt-C} = 721 \text{ Hz})$ . HRMS (FAB): Calcd for C<sub>37</sub>H<sub>58</sub>N<sub>2</sub>OPtSi<sub>2</sub> 797.3735, Found 797.3733.

#### Pt(IPr<sup>AN</sup>)(dvtms) (4m).



IPr<sup>AN</sup>•HCl was synthesized according to the literature procedure.<sup>46</sup>

Synthesis of the title Pt complex was conducted with the same procedure for **4g** except that  $IPr^{AN}$ •HCl was used instead of  $IPr^{Me}$ •HCl on 0.47 mmol scale based on H<sub>2</sub>PtCl<sub>6</sub>•nH<sub>2</sub>O to give 153 mg of Pt( $IPr^{AN}$ )(dvtms) as an orange solid (36%). Mp > 215 °C (decomp.). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  -0.72 (s, 6H), 0.16 (s, 6H), 0.98 (d, *J* = 6.4 Hz, 12H), 1.26 (d, *J* = 6.8 Hz, 12H), 1.50 (dd, *J* = 13.5,

11.7 Hz,  ${}^{3}J_{Pt-C} = 25$  Hz, 2H), 1.71 (d, J = 13.3 Hz,  ${}^{3}J_{Pt-C} = 27$  Hz, 2H), 1.84 (d, J = 11.5 Hz,  ${}^{3}J_{Pt-C} = 28$  Hz, 2H), 3.19 (septet, J = 6.9 Hz, 4H), 6.84 (d, J = 6.9 Hz, 2H), 7.28 (d, J = 7.8 Hz, 4H), 7.33 (dd, J = 8.2, 6.9 Hz, 2H), 7.46 (t, J = 7.8 Hz, 2H), 7.68 (d, J = 7.8 Hz, 2H).  ${}^{13}$ C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  -2.17, 1.61, 23.1, 25.4, 28.6, 36.4, 42.6 ( ${}^{1}J_{Pt-C} = 84$  Hz), 121.3, 124.1, 126.9, 127.2, 127.4, 129.55, 129.60, 129.64, 135.4, 140.1, 145.9, 194.9.

IR (ATR): 2960 w, 2869 w, 1477 w, 1382 w, 1330 w, 1303 w, 1245 m, 1182 w, 1062 w, 988 m, 915 w, 859 w, 836 s, 820 s, 799 m, 781 s, 694 m. HRMS (FAB): Calcd for  $C_{45}H_{58}N_2OPtSi_2$  893.3735, Found 893.3731.

#### 2.4.4 Optimization Studies

Effect of the ligand for Pt-catalyzed borylation of benzene. Pt(NHC)(dvtms) (0.90 µmol) was dissolved into 10 mL of benzene to prepare Pt(NHC)(dvtms) solution ( $9.0 \times 10^{-4}$  M). B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol) and Pt(NHC)(dvtms) benzene solution (1.0 mL, 0.9 µmol, 0.3 mol%) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The vial was stirred at 100 °C for 20 h, and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/EtOAc = 5/1). The filtrate was analyzed by gas chromatography (GC) using dodecane as an internal standard. As a result, ICy-ligated complex 4g exhibited the highest TON among the complexes tested. The results are summarized in table 1.

**Optimization of the reaction of mesitylene.** Pt(NHC)(dvtms) (6.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol) and mesitylene (1.0 mL) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The vial was stirred at 120 °C for 20 h, and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/EtOAc = 5/1). The filtrate was analyzed by gas chromatography (GC) using dodecane as an internal standard. As a result, IPr-bound complex **4b** exhibited the highest activity.

Me	Me			(ULO) ( -[	Me		Me
Į		+ B <sub>2</sub> pin <sub>2</sub>		IHC)(dvtms) 2	moi%		Ĺ
	$\mathbf{Y}$	21 2	120	°C, 20 h		Ŷ	`Bpin
	Ме	1a				Me	
	2ј	0.3 mmol				Зј	
				GC yield (%)			
	Entry	Pt complex	NHC	product	recovered	B <sub>2</sub> pin <sub>2</sub>	
	1	4a	IMes	46	0		
	2	4b	IPr	72	0		
	3	4c	IMe	0	55		
	4	4d	l <sup>′</sup> Pr	17	41		
	5	4e	l <sup>t</sup> Bu	37	0		
	6	4f	IAd	66	0		
	7	4g	ICy	55	5		
	8	4h	SICy	18	31		
	9	4i	BICy	31	28		

#### 2.4.5 General Procedures for Platinum-Catalyzed Borylation of Arenes

# Method A: Procedure for the Platinum-Catalyzed Borylation of Benzene with Bis(pinacolato)diboron (3a, Scheme 1)



Pt(ICy)(dvtms) (3.6 mg, 5.9  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol) and benzene (1.0 mL) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The vial was stirred at 100 °C for 20 h, and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was analyzed by gas chromatography (GC) using dodecane as an internal standard (94% GC yield). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/ EtOAc = 25/1) to give 4,4,5,5-tetramethyl-2-phenyl-1,3,2-dioxaborolane as a white solid (57 mg, 93% based on B<sub>2</sub>pin<sub>2</sub>).

# Method B: Procedure for the Platinum-Catalyzed Borylation of 1,4-Difluorobenzene with Bis(pinacolato)diboron (6b, Figure 6)



Pt(ICy)(dvtms) (3.6 mg, 5.9  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol) and 1,4-difluorobenzene (154  $\mu$ L, 1.5 mmol) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The vial was stirred at 80 °C for 20 h and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was analyzed by NMR using 1,1,2,2-tetrachloroethane as an internal standard (86% NMR yield). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with

hexane/ EtOAc = 25/1) to give 2-(2,5-difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane as a colorless oil (33 mg, 46% based on B<sub>2</sub>pin<sub>2</sub>).

# Method C: Procedure for the Platinum-Catalyzed Borylation of *N*-Methylindole with Bis(pinacolato)diboron (8a, Figure 7)



Pt(ICy)(dvtms) (3.6 mg, 5.9  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (152 mg, 0.60 mmol), *N*-methylindole (40 mg, 0.30 mmol) and heptane (1.0 mL) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The vial was stirred at 100 °C for 20 h and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/ EtOAc = 25/1) to give 1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole as a white solid (69 mg, 87% based on *N*-methylindole).

## Method D: Procedure for the Platinum-Catalyzed Borylation of 2v with Bis(pinacolato)diboron (3v, Figure 5)



Pt(SIPr)(dvtms) (6.9 mg, 9.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol), **2v** (86 mg, 0.30 mmol) and 1,3,5triisopropylbenzen (0.1 mL) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The mixture was stirred at 120 °C for 16 h and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/ EtOAc = 100/1 to 50/1) to give the target borylated product as a white solid (63 mg, 51% based on **2v**) and 25 mg of unreacted starting **2v** was recovered (29%).

## 2.4.6 Spectroscopic Data of the Products

#### 4,4,5,5-Tetramethyl-2-phenyl-1,3,2-dioxaborolane (3a) [CAS: 24388-23-6].

Rf 0.51 (hexane/EtOAc = 5/1). White solid (57 mg, 93%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.35 (s, 12H), 7.37 (t, J = 7.6 Hz, 2H), 7.46 (t, J = 7.3 Hz, 1H), 7.81 (d, J = 7.1 Hz, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.8, 83.7, 127.6, 131.2, 134.7. HRMS (EI): Calcd for C<sub>12</sub>H<sub>17</sub>BO<sub>2</sub> 204.1322, Found 204.1321.

#### Borylation of toluene (3b, Figure 2).



Method A was used, except that the reaction was conducted in toluene (1.0 mL) at 120 °C. After purification by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1), a mixture of three isomers was obtained (59 mg, 90%, >99% purity by GC). GC/MS analysis showed the formation of three isomers of the borylated product, all of which had an m/z of 218 (M<sup>+</sup>). The identity and ratio of each of the three isomers were determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>10a,11</sup> The resonances specific to each isomer are as follows: 7.76 ppm (dd, J = 7.6, 1.4 Hz, 1H, ortho isomer, H<sub>a</sub>); 7.60-7.64 ppm (m, 2H, meta isomer, H<sub>b</sub> and H<sub>c</sub>), 7.71 ppm (d, J = 8.2 Hz, 1H, para isomer, H<sub>d</sub>). (see below.)



Borylation of anisole (3c, Figure 2).



Method A was used, except that the reaction was conducted in anisole (1.0 mL) at 120 °C. After purification by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1), a mixture of three isomers of the borylated product were obtained (57 mg, 82%, >99% purity by GC). GC/MS analysis showed the formation of three isomers of the borylated product, all of which had an m/z of 234 (M<sup>+</sup>). The identity and ratio of each of the borylated products were determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>10a</sup> The resonances specific to each isomer are as follows: 7.67 ppm (dd, J = 1.8 Hz, 7.3 Hz, 1H ortho isomer, H<sub>a</sub>); 7.01 ppm (ddd, J = 0.9 Hz, 2.7 Hz, 8.2 Hz, 1H, meta isomer, H<sub>b</sub>), 7.75 ppm (d, J = 8.7 Hz, 2H, para isomer, H<sub>c</sub>). (See below.)



Borylation of (trifluoromethyl)benzene (3d, Figure 2).



Method A was used, except that the reaction was conducted in (trifluoromethyl)benzene (1.0 mL) at 120 °C. After purification by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1), a mixture of two isomers of the borylated product were obtained (46 mg, 59%, >99% purity by GC). GC/MS analysis showed the formation of two isomers of the borylated product, all of which had an m/z of 272 (M<sup>+</sup>). The identity and ratio of each of the borylated products were determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>10a</sup> The resonances specific to each isomer are as follows:

meta-isomer;  $\delta$  7.48 (t, *J* = 7.4 Hz, 1H), 7.70 (d, *J* = 7.8 Hz, 1H), 7.97 (d, *J* = 7.3 Hz, 1H), 8.06 (s, 1H). para-isomer;  $\delta$  7.61 (d, *J* = 7.8 Hz, 2H), 7.91 (d, *J* = 7.8 Hz, 1H). (See below.)



Borylation of chlorobenzene (3e, Figure 2).



Method A was used, except that the reaction was conducted in chlorobenzene (1.0 mL) at 120 °C. After purification by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1), a mixture of three isomers of the borylated product and PhBpin was obtained (41 mg). GC/MS analysis showed the formation of three isomers of ClC<sub>6</sub>H<sub>4</sub>Bpin (m/z = 238) and PhBpin (m/z = 204). The identity and ratio of each of the borylated products were determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>47</sup> The resonances specific to each isomer are as follows: ortho-isomer,  $\delta$  7.68 (d, partially overlapped in the H<sub>c</sub> peak of meta-isomer, 1H, H<sub>a</sub>). meta-isomer,  $\delta$  7.66 (dt, *J* = 7.3, 1.1 Hz, 1 H, H<sub>c</sub>), 7.78 (d, *J* = 1.4 Hz, 1H, H<sub>b</sub>). para-isomer,  $\delta$  7.73 (d, *J* = 8.4 Hz, 2H, H<sub>d</sub>). PhBpin,  $\delta$  7.81 (dd, *J* = 8.0, 1.1 Hz, 2H, H<sub>e</sub>). <sup>1</sup>H NMR spectroscopic analysis showed that the yields of ClC<sub>6</sub>H<sub>4</sub>Bpin and ClC<sub>6</sub>H<sub>4</sub>Bpin were 55% and 3 %, respectively.



# 2-(2,5-Dimethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3f, Figure 2) [CAS: 356570-53-1].



Method A was used, except that the reaction was conducted in *p*-xylene (1.0 mL) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0  $\mu$ mol) instead of Pt(ICy)(dvtms).

129.8, 131.5, 133.9, 136.3, 141.7. HRMS (EI): Calcd for  $C_{14}H_{21}BO_2$  232.1635, Found 232.1628.

## 2-(5-(tert-Butyl)-2-methylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6g, Figure 2).



Method A was used, except that the reaction was conducted in 1-(tert-butyl)-4methylbenzene (1.0 mL) at 120 °C with Pt(IAd)(dvtms) (4.3 mg, 6.0 µmol) instead of Pt(ICy)(dvtms).

Rf 0.68 (hexane/EtOAc = 5/1). White solid (64 mg, 78%). Mp = 46-48 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.32 (s, 9H), 1.34 (s, 12H), 2.50 (s, 3H, H<sub>a</sub>), 7.10 (d, *J* = 8.0 Hz, 1H, H<sub>b</sub>), 7.35 (dd, *J* = 8.0, 2.3 Hz, 1H, H<sub>c</sub>), 7.77 (d, *J* = 2.3 Hz, 1H, H<sub>d</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  21.7, 24.9, 31.4, 34.3, 83.3, 127.9, 129.6, 132.4, 141.9, 147.1. IR (ATR): 2969 w, 1461 w, 1398 m, 1368 s, 1337 s, 1307 s, 1269 m, 1255 m, 1211 w, 1165 w, 1141 s, 1116 m, 1067 m, 995 s, 963 m, 926 w, 889 w, 853 m, 829 m, 781 w, 737 m, 685 m, 575 m. MS m/z (% relative intensity): 274 (M+, 13), 260 (17), 259 (100), 258 (24), 159 (22), 131 (14), 101 (14), 83 (13), 57 (18). HRMS (EI): Calcd for C<sub>17</sub>H<sub>27</sub>BO<sub>2</sub> 274.2104, Found 204.2101.



The regiochemistry of the compound was determined by NOE experiments in CDCl<sub>3</sub>. When the signal for H<sub>a</sub> at  $\delta = 2.50$  (s, 3H, offset signal  $\delta = 2.4937$ , mixing time 1 s) was irradiated, H<sub>b</sub> at  $\delta = 7.10$  (d, J = 7.8 Hz, 1H) was enhanced by 4%.

## 2,5-Diisopropylphenol (3h, Figure 2) [CAS: 35946-91-9].



The borylated product contained inseparable unidentified impurities in this specific case, therefore product was isolated in the form of the corresponding phenol after oxidative work-up.

Me Method A was followed, except that the reaction was conducted in 1,4diisopropylbenzene (1.0 mL) at 140 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0 µmol) instead of Pt(ICy)(dvtms). After filtration of the mixture through a pad of Celite, an aqueous solution of NaOH (1.0 M, 1.0 mL) and an aqueous solution of H<sub>2</sub>O<sub>2</sub> (30%, 0.5 mL) were added to the filtrate and the mixture was stirred for 30 min at ambient temperature. A small amount of KI was added to quench the residual H<sub>2</sub>O<sub>2</sub> and the mixture was extracted three times with AcOEt. The combined organic extracts were concentrated in vacuo and purified by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1) to give 2,5-diisopropylphenol as a colorless oil (18 mg, 34% based on B<sub>2</sub>pin<sub>2</sub>). Rf 0.40 (hexane/EtOAc = 5/1). Colorless oil (18 mg, 34%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.22 (d, J = 6.9 Hz, 6H), 1.25 (d, J = 6.9 Hz, 6H), 2.82 (septet, J = 6.9 Hz, 1H), 3.15 (septet, J = 6.9 Hz, 1H), 4.62 (br, 1H), 6.62 (d, J = 1.4 Hz, 1H), 6.78 (dd, J = 7.8, 1.4 Hz, 1H), 7.11 (d, J = 8.2 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  22.6, 23.9, 26.8, 33.5, 113.3, 119.0, 126.2, 131.5, 147.8, 152.5. HRMS (EI): Calcd for C<sub>12</sub>H<sub>18</sub>O 178.1358, Found 178.1355.

#### Borylation of *m*-xylene (3i, Figure 2).



Method A was used, except that the reaction was conducted in *m*-xylene (1.0 mL) at 120 °C. After purification by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1), a mixture of three isomers of the borylated product were obtained (44 mg, 63%, >99% purity by GC). GC/MS analysis showed the formation of three isomers of the borylated product, all of which had an m/z of 232 (M<sup>+</sup>). The identity and ratio of each of the borylated products were determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>10a,11,48</sup> The resonances specific to each isomer are as follows.

2-Borylated isomer: δ 2.39 (s, 6H), 6.94 (d, *J* = 7.8 Hz, 1H), 7.12 (t, *J* = 7.3 Hz, 2H).

4-Borylated isomer: δ 2.31 (s, 3H), 2.50 (s, 3H), 6.97-7.00 (m, 2H), 7.66 (d, *J* = 7.8 Hz, 1H).

5-Borylated isomer:  $\delta$  2.32 (s, 6H), 7.10 (d, J = 0.9 Hz, 1H), 7.44 (s, 2H).



## 2-(3,5-Dimethylbenzyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (eq 1) [CAS: 356570-54-2].

Me Bpin Me  $[Ir(OMe)(cod)]_2$  (2.0 mg, 3.0 µmol), 4,4'-di-tert-butyl-2,2'-bipyridine (1.7 mg, 6.0 µmol), B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol), and m-xylene (1.0 mL) were added to a 10 mL sample vial with a Teflon-sealed screwcap. The cap was used to seal the vial under a flow of N<sub>2</sub>. The mixture was stirred at 120 °C for 20 h and the resulting mixture

was filtered through silica gel (eluting with 7 mL of hexane/AcOEt = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1) to give 2-(3,5-dimethylbenzyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane as colorless oil (110 mg, 151% based on B<sub>2</sub>pin<sub>2</sub>).

Rf 0.51 (hexane/EtOAc = 5/1). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.24 (s, 12H), 2.22 (s, 2H), 2.26 (s, 6H), 6.76 (s, 1H), 6.80 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  21.3, 24.7, 83.3, 126.5, 126.8, 137.6, 138.3. HRMS (EI): Calcd for C<sub>15</sub>H<sub>23</sub>BO<sub>2</sub> 246.1791, Found 246.1788.

## 2-Mesityl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3j, eq 3) [CAS: 171364-84-4].

Me Method A was used except that the reaction was conducted in mesitylene (1.0 mL) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0 μmol) instead of Pt(ICy)(dvtms).

<sup>Bpin</sup> Rf 0.46 (hexane/EtOAc = 5/1). Colorless oil (53 mg, 72%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.37 (s, 12H), 2.24 (s, 3H), 2.37 (s, 6H), 6.77 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 24.12) (CDCl<sub>3</sub>, 24.12) (c, 14.12) (c, 14

100.53 MHz):  $\delta$  21.2, 22.2, 24.9, 83.4, 127.4, 138.9, 142.1. HRMS (EI): Calcd for C<sub>15</sub>H<sub>23</sub>BO<sub>2</sub> 246.1791, Found 246.1792.

Me

Мe

Me Me

Мe

Me

# 2-(4-(tert-Butyl)-2,6-dimethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3k, Figure 4) [CAS: 1701449-62-8].

Method A was used, except that the reaction was conducted in 1-(tert-butyl)-3,5dimethyl-benzene (1.0 mL) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0 μmol) instead of Pt(ICy)(dvtms).

 $M_{\text{Me}}$  Rf 0.63 (hexane/EtOAc = 5/1). White solid (68 mg, 78%). Mp = 169 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.27 (s, 9H), 1.36 (s, 12H), 2.41 (s, 6H), 6.97 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 22.6, 24.9, 31.2, 34.3, 83.4, 123.6, 141.9, 152.2. IR (ATR): 2961 w, 2866 w, 1604 w, 1546 w, 1436 w, 1406 w, 1368 s, 1332 s, 1300 s, 1275 m, 1213 w, 1138 s, 1109 w, 1060 m, 963 w, 859 m, 781 w, 739 w, 680 m. MS m/z (% relative intensity): 288 (M+, 25), 274 (18), 273 (100), 272 (26), 231 (16), 189 (16), 173 (34), 101 (16), 83 (20), 57 (50), 55 (15). HRMS (EI): Calcd for C<sub>18</sub>H<sub>29</sub>BO<sub>2</sub> 288.2261, Found 288.2265.

## 2-(4-Isopropyl-2,6-dimethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3l, Figure 4).

Method A was used, except that the reaction was conducted in 1-isopropyl-3,5dimethylbenzene (1.0 mL) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0 μmol) instead of Pt(ICy)(dvtms). The isomer ratio was determined to be 92:8 based on <sup>1</sup>H NMR. The spectroscopic data for major isomer **3l** are as follows.

Rf 0.69 (hexane/EtOAc = 5/1). White solid (48 mg, 59%). Mp = 59-61 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.19 (d, *J* = 6.9 Hz, 6H), 1.37 (s, 12H), 2.39 (s, 6H), 2.78 (septet, *J* = 6.9 Hz, 1H), 6.81 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  22.4, 23.8, 24.9, 34.0, 83.4, 124.8, 142.2, 150.0. IR (ATR): 2960 m, 2926 w, 2867 w, 1606 w, 1559 w, 1433 m, 1370 m, 1328 s, 1301 s, 1273 m, 1213 w, 1140 s, 1106 w, 1085 m, 1059 m, 1006 w, 962 w, 853 m, 834 w, 796 w, 739 w, 679 m, 579 w. MS m/z (% relative intensity): 274 (M+, 51), 259 (33), 217 (64), 216 (26), 175 (100), 174 (70), 173 (18), 159 (44), 131 (27), 105 (14), 101 (22), 91 (16), 83 (24), 55 (17). HRMS (EI): Calcd for C<sub>17</sub>H<sub>27</sub>BO<sub>2</sub> 274.2104, Found 274.2106.

Method D was followed except that the reaction was conducted with 1-isopropyl-3,5-dimethylbenzene (42 mg, 0.28 mmol) as a substrate. The unreacted starting material (8.3 mg, 20%) was recovered and the borylated product (47 mg, 47%) was obtained. The isomer ratio was determined to be 96:4 based on <sup>1</sup>H NMR.

Resonances assignable to minor isomer are as follows


CO<sub>2</sub>Et

Мe

Me

Me

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.23 (d, *J* = 6.9 Hz, 6H), 1.37 (s, 12H), 2.67 (s, 3H), 2.34 (s, 3H), 3.03 (quintet, *J* = 6.9 Hz, 1H), 6.78 (s, 1H), 6.88 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  21.5, 24.4, 24.9, 30.9, 33.6, 83.6, 122.0, 127.6. Three peaks were overlapped with those for major isomer. MS m/z (% relative intensity): 274 (M+, 51), 259 (33), 217 (33), 175 (43), 174 (51), 173 (63), 159 (96), 158 (27), 157

(25), 145 (20), 131 (45), 101 (29), 84 (100), 83 (29). HRMS (EI): Calcd for  $C_{17}H_{27}BO_2$  274.2104, Found 274.2102.

## 4,4,5,5-Tetramethyl-2-(2,4,6-triethylphenyl)-1,3,2-dioxaborolane (3m, Figure 4) [CAS: 1334181-19-9].

Me Method A was used, except that the reaction was conducted in 1,3,5-triethylbenzene (1.0 mL) at 140 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0  $\mu$ mol) instead of Pt(ICy)(dvtms). Rf 0.60 (hexane/EtOAc = 5/1). Colorless oil (60 mg, 68%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.17-1.25 (m, 9H), 1.37 (s, 12H), 2.57 (q, *J* = 7.5 Hz, 2H), 2.67 (q, *J* = 7.6 Hz, 4H), 6.83 (s, 2H).<sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  15.5, 16.9, 24.9, 28.9, 29.4, 83.5, 124.8, 145.4, 148.3. IR (ATR): 2965 w, 2932 w, 2871 w, 1607 w, 1557 w, 1459 w, 1422 w, 1371 m, 1337 s, 1296 s, 1247 w, 1213 w, 1142 s, 1109 w, 1068 s, 961 w, 857 s, 781 w, 733 w, 681 m, 577 w. MS m/z (% relative intensity): 288 (M+, 31), 232 (15), 231 (100), 230 (30), 203 (17), 189 (18), 188 (35), 187 (23), 173 (15), 159 (26), 131 (14), 117 (17), 101 (39), 84 (20), 83 (22). HRMS (EI): Calcd for C<sub>18</sub>H<sub>29</sub>BO<sub>2</sub> 288.2961, Found 288.2263.

### Ethyl 2,4,6-trimethyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3n, Figure 4).

Method A was used, except that the reaction was conducted in ethyl 2,4,6trimethylbenzoate (1.0 mL) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0  $\mu$ mol) instead of Pt(ICy)(dvtms).

<sup>11</sup> Rf 0.27 (hexane/EtOAc = 10/1). Colorless oil (59 mg, 62%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.35 (t, *J* = 7.1 Hz, 3H), 1.37 (s, 12H), 2.24 (s, 3H), 2.34 (s, 3H), 2.35 (s,

3H), 4.35 (q, J = 7.1 Hz, 2H), 6.81 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  14.4, 19.7, 19.8, 22.2, 25.1, 60.8, 83.9, 128.5, 131.9, 135.6, 138.3, 143.0, 170.7. IR (ATR): 2978 w, 2930 w, 2868 w, 1724 s, 1599 w, 1443 w, 1372 w, 1333 m, 1300 m, 1257 s, 1201 s, 1142 s, 1098 s, 1023 m, 857 m. MS m/z (% relative intensity): 319 (M+1, 11), 318 (M+, 54), 273 (66), 272 (54), 261 (55), 215 (100), 189 (35), 173 (36), 172 (61), 83 (40). HRMS (EI): Calcd for C<sub>18</sub>H<sub>27</sub>BO<sub>4</sub> 318.2002, Found 318.2002.

### N,N,2,4,6-Pentamethyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (30, Figure 4).

Me Method A was used, except that the reaction was conducted in N,N,2,4,6pentamethylaniline (1.0 mL) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0 µmol) instead of Pt(ICy)(dvtms).

Me Rf 0.46 (hexane/EtOAc = 10/1). White solid (50 mg, 58%). Mp = 90-91 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.38 (s, 12H), 2.21 (s, 3H), 2.29 (s, 3H), 2.31 (s, 3H), 2.75 (s, 6H), 6.75 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  18.6, 19.2, 21.7, 25.1, 42.6, 83.7, 130.2, 138.0, 138.4, 140.9,

146.9. IR (ATR): 2976 w, 2919 w, 2783 w, 1439 w, 1392 m, 1372 m, 1334 s, 1298 s, 1211 w, 1136 s, 1023 s, 858 s. MS m/z (% relative intensity): 290 (M+1, 17), 289 (M+, 100), 274 (36), 83 (38). HRMS (EI): Calcd for C<sub>17</sub>H<sub>28</sub>BNO<sub>2</sub> 289.2213, Found 289.2215.

### 2-(3-Chloro-2,4,6-trimethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3p, Figure 4).

Method A was used, except that the reaction was conducted in 2-chloro-1,3,5-Me trimethylbenzene (1.0 mL) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0 µmol) instead of Pt(ICy)(dvtms). Isolation by column chromatography gave the title compound as a Bpin colorless oil (67 mg). NMR analysis revealed that the sample contained 4% of a dechlorinated product **3j**. Thus, the yield of the borylated product was determined to be 76%.

Rf 0.43 (hexane/EtOAc = 10/1). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.38 (s, 12H), 2.31 (s, 3H), 2.32 (s, 3H), 2.42 (s, 3H), 6.86 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 205, 21.0, 21.7, 25.1, 84.0, 129.8, 132.2, 137.1, 139.1, 139.4. IR (ATR): 2978 w, 2926 w, 2866 w, 1436 w, 1375 m, 1323 s, 1299 s, 1211 w, 1143 s, 984 m, 857 m. MS m/z (% relative intensity): 282 (M+2, 14), 280 (M+, 39), 225 (33), 223 (100), 222 (36), 181 (48), 180 (81), 145 (34), 115 (30), 101 (31), 85 (30), 83 (30). HRMS (EI): Calcd for C<sub>15</sub>H<sub>22</sub>BClO<sub>2</sub> 280.1401, Found 280.1403.

### 2-(3-Fluoro-2,4,6-trimethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3q, Figure 4).

Method A was used, except that the reaction was conducted in 2-fluoro-1,3,5-Me trimethylbenzene (1.0 mL) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0 µmol) instead of Pt(ICy)(dvtms). Bpin

Rf 0.51 (hexane/EtOAc = 10/1). Colorless oil (59 mg, 74%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.38 (s, 12H), 2.19 (d, J = 2.3 Hz, 3H), 2.29 (d, J = 2.7 Hz, 3H), 2.31 (s, 3H), 6.76 (d, J = 6.9 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  14.2 (d, J = 5.8 Hz), 14.8 (d, J = 3.8 Hz), 21.6, 25.1, 83.9, 125.3 (d, J = 18.2 Hz), 127.9 (d, J = 15.3 Hz), 130.0 (d, J = 4.8 Hz), 136.8 (d, J = 3.8 Hz), 158.1 (d, J = 242 Hz). IR (ATR): 2979 w, 2929 w, 1442 w, 1412 m, 1372 m, 1329 s, 1301 s, 1273 m, 1217 w, 1166 m, 1142 s, 1021 m, 858 m. MS m/z (% relative intensity): 264 (M+, 47), 207 (100), 206 (37), 165 (48), 164 (92), 163 (34). HRMS (EI): Calcd for C<sub>15</sub>H<sub>22</sub>BFO<sub>2</sub> 264.1697, Found 264.1698.

### 4,4,5,5-Tetramethyl-2-(2,3,5,6-tetramethylphenyl)-1,3,2-dioxaborolane (3r, Figure 4).



Me

Me

Method A was used, except that the reaction was conducted in melted 1,2,4,5tetramethylbenzene (805 mg, 20 equiv) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0 µmol) instead of Pt(ICy)(dvtms).

Rf 0.34 (hexane/EtOAc = 10/1). White solid (59 mg, 76%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78) MHz): δ 1.39 (s, 12H), 2.16 (s, 6H), 2.23 (s, 6H), 6.88 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 19.0, 19.7, 25.2, 83.8, 132.4, 133.1, 136.7. IR (ATR): 2977 m, 2931 w, 1463 w, 1410 m, 1376 m, 1338 s, 1295 s, 1251 m, 1195 w, 1142 s, 1018 w, 962 w. MS m/z (% relative intensity): 260 (M+, 60), 259 (15), 204 (14), 203 (100), 202 (36), 188 (12), 161 (40), 160 (80), 159 (42), 145 (34), 119 (26), 117 (19), 115 (15), 105 (14), 101 (17), 91 (17), 85 (13), 83 (17), 55 (12). HRMS (EI): Calcd for C<sub>16</sub>H<sub>25</sub>BO<sub>2</sub> 261.1948, Found 260.1949.

## 4,4,5,5-Tetramethyl-2-(3,3',5,5'-tetramethyl-[1,1'-biphenyl]-4-yl)-1,3,2-dioxaborolane (3s, Figure 4).



Me

Bpin

Me

Bpin

Method A was used, except that the reaction was conducted in melted 3,3',5,5'-tetramethyl-1,1'-biphenyl (1.19 g, 19 equiv) at 120 °C with Pt(IPr)(dvtms) (4.6 mg, 6.0  $\mu$ mol) instead of Pt(ICy)(dvtms).

Me Rf 0.34 (hexane/EtOAc = 10/1). White solid (76 mg, 75%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.39 (s, 12H), 2.36 (s, 6H), 2.46 (s, 6H), 6.96 (s, 1H), 7.17 (s, 2H), 7.19 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 21.4, 22.4, 25.0, 83.6, 125.0, 125.4, 128.8, 138.0, 141.3, 142.1, 142.4. IR (ATR): 2977 w, 2921 w, 1602 m, 1552 w, 1438 w, 1370 m, 1329 s, 1302 s, 1274 w, 1145 s, 1110 w, 1061 m, 962 w.

MS m/z (% relative intensity): 338 (M+2, 18), 337 (M+1, 66), 336 (M+, 18), 279 (22), 237 (49), 236 (100), 235 (77). HRMS (EI): Calcd for C<sub>22</sub>H<sub>29</sub>BO<sub>2</sub> 336.2261, Found 336.2261.

# 2,2'-(5-Chloro-2,4,6-trimethyl-1,3-phenylene)bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolane) (3p-di, Figure 5).

Method D was followed except that the reaction was conducted with 2-chloro-1,3,5-<sup>Me</sup> trimethylbenzene (46 mg, 0.30 mmol) as a substrate. Monoborylated product **3pmono** (54 mg, 64%) and the title compound **3p-di** (27 mg, 23%) were obtained.

Me Rf 0.53 (hexane/EtOAc = 5/1). White solid (27 mg, 23%). Mp = 160-163 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.37 (s, 24H), 2.38 (s, 3H), 2.39 (s, 6H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  20.8, 22.0, 25.0, 83.8, 132.3, 139.6, 142.2. IR (ATR): 2978 m, 1425 w, 1373 m, 1337 m, 1300 s, 1267 m, 1141 s, 987 w, 857 w, 671 w. MS m/z (% relative intensity): 408 (9), 407 (9), 406 (M+, 26), 405 (12), 349 (12), 246 (24), 248 (16), 101 (100), 83 (35). HRMS (EI): Calcd for C<sub>21</sub>H<sub>33</sub>B<sub>2</sub>ClO<sub>4</sub> 406.2253, Found 406.2257.

# 2,2'-(5-Fluoro-2,4,6-trimethyl-1,3-phenylene)bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolane) (3q-di, Figure 5).



Me Rf 0.51 (hexane/EtOAc = 5/1). White solid (13 mg, 11%). Mp = 144-145 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.40 (s, 24H), 2.26 (d, *J* = 2.3 Hz, 6H), 2.39 (s, 3H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  14.2 (d, *J* = 4.8 Hz), 21.9, 24.9, 83.7, 128.4 (d, *J* = 17 Hz), 139.9 (d, *J* = 4.8 Hz), 157.6 (d, *J* = 240 Hz). IR (ATR): 2977 w, 1444 w, 1370 m, 1306 s, 1267 m, 1213 w, 1138 s, 1109 w, 1031 w, 1007 w, 990 w, 857 m, 831 w, 758 w, 711 w, 672 m. MS m/z (% relative intensity): 390 (M+, 38), 389 (18), 333 (16), 289 (14), 233 (34), 232 (25), 101 (100), 83 (36). HRMS (EI): Calcd for C<sub>21</sub>H<sub>33</sub>B<sub>2</sub>FO<sub>4</sub> 390.2549, Found 390.2548.

To obtain the title compound as a major product, the borylation was repeated twice as described below:

Pt(SIPr)(dvtms) (6.9 mg, 9.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol), fluoromesitylene (21 mg, 0.15 mmol) and octane (0.10 mL) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The mixture was stirred at 120 °C for 16 h and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was concentrated in vacuo and again filtered through a pad of silica gel (eluting with 7 mL of hexane/ EtOAc = 20/1) to give colorless filtrate. The filtrate was concentrated in vacuo and transferred to another vial for 2<sup>nd</sup> borylation. To the reaction mixture, Pt(SIPr)(dvtms) (6.9 mg, 9.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol) and octane (0.10 mL) were added, and the mixture was stirred at 120 °C for 16 h. The resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/ EtOAc = 100/1 to 50/1) to give the target diborylated product **3q-di** as a white solid (38 mg, 65%). The monoborylated product was formed in 25% yield based on <sup>1</sup>H-NMR analysis.

### 2-(4-Cyclopentyl-2,6-dimethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3t, Figure 5).



Bpin

Method D was followed except that the reaction was conducted with 1-cyclopentyl-3,5-dimethylbenzene (50 mg, 0.29 mmol) as a substrate. 42% of the starting material remained unreacted based on <sup>1</sup>H NMR analysis.

Me Rf 0.72 (hexane/EtOAc = 5/1). Colorless oil (45 mg, 53%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.36 (s, 12H), 1.45-1.60 (m, 2H), 1.60-1.71 (m, 2H), 1.71-1.89 (m, 2H), 1.89-2.08 (m, 2H), 2.38 (s, 6H), 2.87 (quintet, *J* = 8.6 Hz, 1H), 6.83 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  22.4, 24.9, 25.6, 34.4, 45.8, 83.4, 125.5, 142.1, 147.7. IR (ATR): 2952 w, 2867 w, 1607 w, 1436 m, 1370 m, 1330 s, 1299 s, 1274 m, 1213 w, 1142 s, 1108 w, 1066 s, 962 w, 858 m, 733 w, 677 w. MS m/z (% relative intensity): 301 (20), 300 (M+, 94), 299 (23), 285 (18), 244 (16), 243 (100), 242 (39), 201 (68), 200 (79), 199 (22), 175 (88), 158 (17), 157 (17), 131 (21), 101 (21), 83 (20), 69 (17). HRMS (EI): Calcd for C<sub>19</sub>H<sub>29</sub>BO<sub>2</sub> 300.2261, Found 300.2260.

Small amount of a regioisomer was also formed in 5% (<sup>1</sup>H-NMR).

Me Resonances assignable to this isomer are as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.37 (s, 12H), 2.26 (s, 3H), 2.34 (s, 3H), 3.01 (quintet, J = 8.0 Hz, 1H), 6.77 (s, 1H), 6.88 (s, 1H). Other peaks were overlapped with those of the major isomer.

<sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 21.5, 22.0, 24.9, 25.7, 35.2, 46.1, 83.6, 122.6, 127.6, 138.8, 140.9, 150.0.

MS m/z (% relative intensity): 301 (19), 300 (M+, 89), 299 (22), 259 (18), 217 (33), 216 (84), 215 (48), 200 (58), 199 (100), 198 (41), 185 (23), 173 (22), 172 (43), 171 (39), 159 (30), 158 (15), 157 (40), 146 (22), 145 (43), 133 (23), 131 (23), 129 (22), 101 (34), 85 (43), 84 (98), 83(, 26), 69 (22). HRMS (EI): Calcd for C<sub>19</sub>H<sub>29</sub>BO<sub>2</sub> 300.2261, Found 300.2255.

sec-Butyl 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3u, Figure 5).



Method D was followed except that the reaction was conducted with secbutyl 3,5-dimethylbenzoate (67 mg, 0.32 mmol) as a substrate. The unreacted starting material was recovered (28 mg, 42%).

<sup>h</sup>e Rf 0.63 (hexane/EtOAc = 5/1). White solid (44 mg, 41%). Mp = 62-63 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.96 (t, *J* = 7.6 Hz, 3H), 1.32 (d, *J* = 6.0 Hz, 3H), 1.42 (d, *J* = 14.7 Hz, 12H), 1.60-1.82 (m, 2H), 2.43 (s, 6H), 5.07 (sextet, *J* = 6.3 Hz, 1H), 7.60 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  9.8, 19.6, 22.1, 24.9, 28.9, 72.6, 84.0, 127.1, 131.2, 141.8, 166.7. IR (ATR): 2974 m, 2931 m, 1708 s, 1560 w, 1445 w, 1370 m, 1335 s, 1295 s, 1207 s, 1147 m, 1107 s, 1061 m, 1026 m, 997 m, 963 w, 943 w, 895 w, 882 w, 854 m, 807 w, 769 m, 732 w, 698 w, 679 m. MS m/z (% relative intensity): 332 (M+, 28), 317 (12), 276 (48), 275 (90), 274 (31), 260 (21), 259 (100), 258 (24), 233 (44), 232 (21), 219 (97), 218 (54), 201 (93), 200 (23), 177 (51), 176 (32), 159 (26), 131 (23), 105 (22), 101 (33), 83 (33), 57 (21), 56 (34). HRMS (EI): Calcd for C<sub>19</sub>H<sub>29</sub>BO<sub>4</sub> 332.2159, Found 332.2161.

### (*1R,2S,5R*)-2-Isopropyl-5-methylcyclohexyl 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3v, Figure 5).



Method D was followed except that the reaction was conducted with (1R,2S,5R)-2-isopropyl-5-methylcyclohexyl 3,5-dimethylbenzoate (86 mg, 0.30 mmol) as a substrate. The unreacted starting material was recovered (25 mg, 29%).

Me Rf 0.60 (hexane/EtOAc = 5/1). White solid (63 mg, 51%). Mp = 88-89 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 0.78 (d, J = 6.9 Hz, 3H), 0.88-0.96 (m, 1H), 0.92 (t, J = 6.4 Hz, 6H), 1.02-1.20 (m, 2H), 1.39 (s, 12H), 1.46-1.59 (m, 2H), 1.72 (d, J = 11.4 Hz, 2H), 1.93 (quintet, J = 6.9, 2.8 Hz, 1H), 2.10 (d, J = 11.9 Hz, 1H), 2.43 (s, 6H), 4.92 (td, J = 10.9, 4.3 Hz, 1H), 7.60 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 16.5, 20.8, 22.1, 23.6, 24.9, 26.4, 31.4, 34.3, 41.0, 47.2, 74.6, 84.0, 127.1, 131.2, 141.9, 166.5.

IR (ATR): 2959 w, 2866 w, 1708 s, 1443 w, 1407 w, 1370 m, 1331 m, 1296 s, 1210 s, 1182 w, 1141 m, 1123 s, 1062 m, 1037 w, 1010 w, 984 w, 962 m, 915 w, 851 m, 813 w, 771 m, 679 w. MS m/z (% relative intensity): 399 ([M-Me]+, 4), 277 (27), 276 (10), 260 (10), 259 (56), 258 (14), 219 (34), 218 (16), 177 (10), 138 (100), 133 (20), 123 (24), 96 (14), 95 (58), 83 (21), 82 (16), 81 (33), 55 (12). HRMS (EI): Calcd for  $C_{25}H_{39}BO_4$  414.2941, Found 414.2940.

## (*1S*,2*R*,4*S*)-1,7,7-Trimethylbicyclo[2.2.1]heptan-2-yl 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3w, Figure 5).



Method D was followed except that the reaction was conducted with (1S,2R,4S)-1,7,7-trimethylbicyclo[2.2.1]heptan-2-yl 3,5-dimethylbenzoate (86 mg, 0.30 mmol) as a substrate. 46 mg of unreacted starting material was recovered (54%).

Rf 0.45 (hexane/EtOAc = 5/1). White solid (43 mg, 35%). Mp = 146-147 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.905 (s, 3H), 0.912 (s, 3H), 0.96 (s, 3H), 1.11 (dd, *J* = 13.7, 3.2 Hz, 1H), 1.29-1.41 (m, 2H), 1.39 (s, 12H), 1.73 (t, *J* = 4.3 Hz, 1H), 1.74-1.86 (m, 1H), 2.05-2.22 (m, 1H), 2.40-2.51 (m, 1H), 2.44

(s, 6H), 5.09 (d, J = 10.1 Hz, 1H), 7.61 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  13.6, 18.9, 19.7, 22.1, 24.9, 27.4, 28.1, 36.8, 45.0, 47.9, 49.1, 80.3, 84.0, 127.1, 131.2, 141.9, 167.2. IR (ATR): 2958 w, 1708 s, 1440 w, 1374 w, 1340 w, 1304 s, 1213 s, 1144 m, 1122 s, 1064 m, 1024 m, 963 w, 857 m, 768 m, 673 w. MS m/z (% relative intensity): 412 (M+, 6), 260 (16), 259 (100), 258 (25), 136 (11), 83 (6). HRMS (EI): Calcd for C<sub>25</sub>H<sub>37</sub>BO<sub>4</sub> 412.2785, Found 412.2789.

To improve the conversion, the reaction was repeated twice as described below:

Pt(SIPr)(dvtms) (6.9 mg, 9.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol), **2w** (95 mg, 0.33 mmol) and TIPB (0.10 mL) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The mixture was stirred at 120 °C for 16 h and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was concentrated in vacuo and again filtered through a pad of silica gel (eluting with 7 mL of hexane/ EtOAc = 10/1) to give colorless filtrate. The filtrate was concentrated in vacuo and transferred to another vial for 2<sup>nd</sup> borylation. To the reaction mixture, Pt(SIPr)(dvtms) (6.9 mg, 9.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol) and TIPB (0.10 mL) were added, and the mixture was stirred at 120 °C for 16 h. The resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/ EtOAc = 100/1 to 50/1) to give the target borylated product as a white solid (93 mg, 68%). The unreacted starting material was recovered (22 mg, 23%).

#### 2-(4-Chloro-2,3,5,6-tetramethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3x, Figure 5).

CI Me Me Bpin Method D was followed except that the reaction was conducted with 3-chloro-1,2,4,5-tetramethylbenzene (51mg, 0.30 mmol) as a substrate. 31% of the starting material remained unreacted based on <sup>1</sup>H-NMR analysis.

Me Rf 0.74 (hexane/EtOAc = 5/1). White solid (39 mg, 45%). Mp = 130-131 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.40 (s, 12H), 2.289 (s, 6H), 2.292 (s, 6H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  17.1, 20.5, 25.2, 84.0, 131.5, 136.5, 137.7. IR (ATR): 2977 m, 1555 w, 1453 w, 1408 m, 1372 m, 1337 s, 1296 s, 1256 m, 1185 m, 1140 s, 1020 s, 963 m, 853 s, 684 m. MS m/z (% relative intensity): 296 (19), 295 (15), 294 (M+, 57), 293 (14), 239 (33), 238 (24), 237 (100), 236 (34), 196 (29), 195 (45), 194 (83), 193 (24), 175 (39), 159 (27), 101 (19), 83 (15). HRMS (EI): Calcd for C<sub>16</sub>H<sub>24</sub>BClO<sub>2</sub> 294.1558, Found 294.1560.

# (3,5-Dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)triethylsilane (3y, Figure 5).



Method D was followed except that the reaction was conducted with (3,5dimethylphenyl)triethylsilane (66 mg, 0.30 mmol) as a substrate. The unreacted starting material was recovered (46 mg, 69%).

Rf 0.63 (hexane/EtOAc = 5/1). White solid (27 mg, 26%). Mp = 81-82 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.74 (q, *J* = 7.9 Hz, 6H), 0.93 (t, *J* = 7.8 Hz, 9H), 1.38 (s, 12H), 2.39 (s, 6H), 7.05 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  3.23, 7.38, 22.3, 25.0, 83.6, 132.2, 138.4, 140.6. IR (ATR): 2955

m, 2875 w, 1435 m, 1369 s, 1333 s, 1305 s, 1259 w, 1213 w, 1142 s, 1109 w, 1058 s, 1014 m, 962 w, 884 m, 857 m, 790 w, 719 s, 700 m, 677 w. MS m/z (% relative intensity): 346 (M+, 10), 318 (23), 317 (90), 316 (22), 290 (20), 289 (87), 288 (21), 262 (21), 261 (100), 260 (24), 189 (12), 161 (17). HRMS (EI): Calcd for C<sub>20</sub>H<sub>35</sub>BO<sub>2</sub>Si 346.2499, Found 346.2499.

To improve the conversion, the reaction was repeated twice as described below:

Pt(SIPr)(dvtms) (6.9 mg, 9.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol), (3,5-dimethylphenyl)triethylsilane (86 mg, 0.30 mmol) and octane (0.10 mL) were added to a 10 mL-sample vial with a Teflonsealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The mixture was stirred at 120 °C for 20 h and the resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was concentrated in vacuo and again filtered through a pad of silica gel (eluting with 7 mL of hexane/ EtOAc = 10/1) to give colorless filtrate. The filtrate was concentrated in vacuo and transferred to another vial for 2<sup>nd</sup> borylation. To the reaction mixture, Pt(SIPr)(dvtms) (6.9 mg, 9.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol) and octane (0.10 mL) were added, and the mixture was stirred at 120 °C for 20 h. The resulting mixture was filtered through silica gel (eluting with 7 mL of hexane/ EtOAc = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/ EtOAc = 100/1 to 50/1) to give the target borylated product as a white solid (56 mg, 54%). The unreacted starting material was recovered (24 mg, 36%).

## 3,3,3-Trifluoropropyl 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3z, Figure 5).



Method D was followed except that the reaction was conducted with 3,3,3-trifluoropropyl 3,5-dimethylbenzoate (70 mg, 0.28 mmol) as a substrate. The unreacted starting material was recovered (27 mg, 39%).

<sup>h</sup>e Rf 0.51 (hexane/EtOAc = 5/1). Colorless oil (56 mg, 53%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.40 (s, 12H), 2.43 (s, 6H), 2.59 (qt, *J* = 10.5, 6.4 Hz, 2H), 4.51 (t, *J* = 6.4 Hz, 2H), 7.60 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  22.1, 25.0, 33.5 (q, *J* = 30 Hz), 57.6 (q, *J* = 3.8 Hz), 84.2, 125.9 (q, *J* = 278 Hz), 127.31, 129.9, 142.0, 166.6. IR (ATR): 2978 w, 1722 m, 1436 w, 1373 m, 1339 m, 1300 s, 1256 m, 1206 s, 1123 s, 1060 m, 1009 w, 962 w, 898 w, 855 m, 813 w, 768 w, 668 w. MS m/z (% relative intensity): 372 (M+, 21), 316 (15), 315 (100), 314 (46), 273 (54), 272 (38), 201 (42), 159 (26), 131 (15), 101 (17), 83 (16). HRMS (EI): Calcd for C<sub>18</sub>H<sub>24</sub>BF<sub>3</sub>O<sub>4</sub> 372.1720, Found 372.1722.

## 3-Chloropropyl 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3aa, Figure 5).



Method D was followed except that the reaction was conducted with 3chloropropyl 3,5-dimethylbenzoate (65 mg, 0.29 mmol) as a substrate. The starting material was recovered (41 mg, 62%). Rf 0.49 (hexane/EtOAc = 5/1). Colorless oil (39 mg, 55%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.40 (s, 12H), 2.23 (quintet, *J* = 6.4 Hz, 2H), 2.43 (s, 6H), 3.70 (t, *J* = 6.4 Hz, 2H), 4.45 (t, *J* = 6.0 Hz, 2H), 7.59 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  22.1, 25.0, 31.8, 41.4, 61.5, 84.1, 127.1, 130.3, 142.0, 166.8. IR (ATR): 2976 w, 1718 m, 1439 w, 1371 w, 1337 m, 1298 s, 1207 s, 1143 m, 1123 s, 1063 m, 1024 w, 961 w, 897 w, 855 m, 809 w, 768 m, 663 w. MS m/z (% relative intensity): 354 (8), 353 (6), 352 (M+, 21), 351 (5), 297 (32), 296 (28), 295 (100), 294 (41), 259 (23), 255 (10), 253 (32), 252 (17), 201 (53), 200 (13), 176 (40), 175 (16), 159 (15), 131 (19), 105 (15), 101 (20), 85 (12), 83 (18). HRMS (EI): Calcd for C<sub>18</sub>H<sub>26</sub>BClO<sub>4</sub> 352.1613, Found 352.1612.

### *tert*-Butyl(3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)dimethylsilane (3ab, Figure 5).



Method D was followed except that the reaction was conducted with *tert*butyl(3,5-dimethylphenoxy)dimethylsilane (71 mg, 0.30 mmol) as a substrate. The starting material was recovered (14 mg, 20%).

Rf 0.79 (hexane/EtOAc = 5/1). White solid (59 mg, 55%). Mp = 103-104 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.16 (s, 6H), 0.96 (s, 9H), 1.37 (s, 12H), 2.35 (s, 6H), 6.44 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  -4.4, 18.2, 22.4, 24.9, 25.7, 83.4, 118.6, 144.1, 156.5. IR (ATR): 2957 w, 2930 w, 2858 w, 1596 m, 1469 w, 1441 w, 1369 m, 1330 s, 1306 s, 1254 m, 1161 s, 1144 s, 1039 m, 961 w, 861 s, 844 s, 804 w, 781 m, 673 w. MS m/z (% relative intensity): 362 (M+, 32), 306 (16), 305 (48), 206 (18), 205 (100), 204 (24), 163 (17), 83 (59). HRMS (EI): Calcd for C<sub>20</sub>H<sub>35</sub>BO<sub>3</sub>Si 362.2449, Found 362.2447.

### Piperidin-1-yl(2,4,6-trimethyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanone (3ac, Figure 5).



Method D was followed except that the reaction was conducted with mesityl(piperidin-1-yl)methanone (69 mg, 0.30 mmol) as a substrate. The starting material was recovered (25 mg, 37%).

Rf 0.23 (hexane/EtOAc = 1/1). Colorless oil (45 mg, 42%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.34 (s, 12H), 1.40-1.54 (m, 2H), 1.56-1.73 (m, 4H), 2.18 (s, 3H), 2.29 (s, 3H), 2.35 (s, 3H), 3.11 (dd, *J* = 6.6, 4.4 Hz, 2H), 3.62-3.91 (m, 2H), 6.81 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  19.1, 22.0, 24.6, 24.9, 24.9, 25.7, 26.6, 41.8, 47.0, 83.7, 128.4, 133.8, 134.5, 137.1, 141.7, 169.9. IR (ATR): 2977 w, 2934 w, 2856 w, 1627 s, 1440 s, 1394 m, 1372 m, 1329 s, 1298 s, 1268 s, 1210 m, 1141 s, 1107 m, 1026 m, 988 m, 958 w, 920 w, 856 m, 831 w, 730 m, 671 w. MS m/z (% relative intensity): 357 (M+, 18), 342 (35), 274 (17), 273 (100), 272 (61), 145 (11). HRMS (EI): Calcd for C<sub>21</sub>H<sub>32</sub>BNO<sub>3</sub> 357.2475, Found 357.2477.





Because the target borylated product was inseparable from the starting material by chromatography, the borylated product was isolated after oxidition to the corresponding phenol.

After the borylation of N-mesityl-N-methylacetamide (57 mg, 0.3 mmol) according to Method D, the reaction mixture was treated with NaOH aq. (1 M, 0.5 mL) and  $H_2O_2$  (30%, 0.60 g) in THF (1.0 mL) for 30 min. The mixture was then quenched with Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> aq, and the mixture was extracted with Et<sub>2</sub>O three times. The combined organic extracts were washed with brine and dried over MgSO<sub>4</sub>. Concentrated reaction mixture was purified by flash column chromatography (eluting with hexane/EtOAc = 5/1 to 1/1) to give the title compound as a white solid (19 mg, 30 %). 39% of the starting material remained unreacted based on <sup>1</sup>H NMR analysis.

Rf 0.09 (hexane/EtOAc = 5/1). White solid (19 mg, 30%). Mp = 149-150 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.73 (s, 3H), 2.11 (s, 6H), 2.25 (s, 3H), 3.11 (s, 3H), 6.90 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  10.7, 15.9, 16.9, 21.3, 34.6, 121.1, 123.1, 126.6, 130.1, 140.2, 151.1, 171.2. IR (ATR): 3285 w, 2922 w, 1632 s, 1482 m, 1442 m, 1383 m, 1307 m, 1247 m, 1220 w, 1180 m, 1129 w, 1051 m, 974 w, 941 w, 915 w, 869 w, 810 w, 743 w. MS m/z (% relative intensity): 207 (M+, 33), 192 (8), 164 (15), 150 (13), 91 (8), 56 (100). HRMS (EI): Calcd for C<sub>12</sub>H<sub>17</sub>NO<sub>2</sub> 207.1259, Found 207.1257.

### Ethyl 4-(2,4,6-trimethyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)butanoate (3ae, Figure 5).



Method D was followed except that the reaction was conducted with ethyl 4-(mesityloxy)butanoate (73 mg, 0.29 mmol) as a substrate. The unreacted starting material was recovered (32 mg, 45%).

Rf 0.45 (hexane/EtOAc = 5/1). Colorless oil (42 mg, 39%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.26 (t, *J* = 7.1 Hz, 3H), 1.38 (s, 12H), 2.09 (quintet, *J* = 6.9 Hz, 2H), 2.19 (s, 3H), 2.29 (s, 3H), 2.30 (s, 3H), 2.57 (t, *J* = 7.3 Hz, 2H), 3.69 (t, *J* = 6.2 Hz, 2H), 4.15 (q, *J* = 7.2 Hz, 2H), 6.77 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  14.2, 15.6, 16.2, 21.7, 25.0, 25.7, 31.0, 60.4, 70.7, 83.6, 130.0, 131.7, 134.2, 137.0, 153.2, 173.4. IR (ATR): 2977 w, 2930 w, 1734 s, 1441 w, 1409 m, 1372 m, 1327 s, 1296 s, 1272 m, 1226 m, 1167 s, 1142 s, 1107 w, 1024 s, 959 w, 922 w, 858 m, 698 w, 672 w. MS m/z (% relative intensity): 376 (M+, 1), 331 (6), 116 (7), 115 (100), 87 (57). HRMS (EI): Calcd for C<sub>21</sub>H<sub>33</sub>BO<sub>5</sub> 376.2421, Found 376.2419.

### 2-(3-Chloro-6-methoxy-2,4-dimethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3af, Figure 5).



Method D was followed except that the reaction was conducted with 2-chloro-5methoxy-1,3-dimethylbenzene (51 mg, 0.30 mmol) as a substrate. The starting material was recovered (35 mg, 69%).

 $^{1}_{OMe}$  Rf 0.51 (hexane/EtOAc = 5/1). White solid (26 mg, 29%). Mp = 102-103 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.38 (s, 12H), 2.35 (s, 3H), 2.36 (s, 3H), 3.73 (s, 3H), 6.55 (s, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 20.4, 21.5, 24.74, 55.8, 83.9, 110.4, 126.7, 138.1, 139.8, 160.5. IR (ATR): 2979 w, 1597 w, 1454 w, 1373 m, 1326 s, 1303 s, 1272 w, 1205 m, 1143 s, 1098 m, 1080 m, 1007 w, 982 w, 960 w, 855 m, 832 m,674 w. MS m/z (% relative intensity): 298 (34), 297 (23), 296 (M+, 100),

295 (23), 241 (28), 240 (19), 239 (84), 238 (32), 225 (17), 224 (17), 223 (50), 222 (14), 180 (18), 167 (21), 155 (16), 153 (28), 85 (16). HRMS (EI): Calcd for C<sub>15</sub>H<sub>22</sub>BClO<sub>3</sub> 296.1351, Found 296.1354.

#### 4,4,5,5-Tetramethyl-2-(2,4,6-tripropylphenyl)-1,3,2-dioxaborolane (3ag, Figure 5).



Method D was followed except that the reaction was conducted with 1,3,5-tripropylbenzene (61 mg, 0.30 mmol) as a substrate and in octane (0.10 mL) instead of TIPB. The unreacted starting material was recovered (43mg, 70%). Rf 0.63 (hexane/EtOAc = 5/1). Colorless oil (29 mg, 29%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.89 (t, *J* = 7.8 Hz, 3H), 0.93 (t, *J* = 7.3 Hz, 6H), 1.37 (s, 12H), 1.52-1.64 (m, 6H), 2.49 (t, *J* = 7.8 Hz, 2H), 2.59 (t, *J* = 7.8 Hz, 4H), 6.77 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  13.8, 14.1, 24.5, 25.0, 26.0, 38.0, 38.7, 83.4, 126.3, 143.3, 146.6. IR (ATR): 2958 m, 2929 m, 2869 w, 1607 w, 1465 w, 1420 w, 1372 m, 1335 s, 1297 m, 1213 w, 1143 s, 1111 w, 1084 w, 1063 m, 962 w, 857 m, 680 w. MS m/z (% relative intensity): 331 (10), 330 (M+, 46), 329 (11), 302 (15), 288 (15), 274 (19), 273 (100), 272 (30), 231 (37), 230 (30), 229 (24), 216 (20), 201 (44), 187 (22), 173 (15), 159 (14), 131 (12), 117 (14), 101 (36), 83 (27). HRMS (EI): Calcd for C<sub>21</sub>H<sub>35</sub>BO<sub>2</sub> 330.2730, Found 330.2732.

#### Borylation of fluorobenzene (6a, eq 4).



Method B was followed except that the reaction was conducted in fluorobenzene (141  $\mu$ L, 1.5 mmol). NMR analysis before the purification by column chromatography showed that the borylated products were produced in 84% (*o:m:p* = 73:20:7).

After purification by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1), a mixture of three isomers of the borylated products were obtained (35 mg, 53%, *o:m:p* = 74:21:5, >99% purity by GC). GC/MS analysis revealed the formation of three isomers of the borylated product, all of which had an m/z of 222 (M<sup>+</sup>). The identity and ratio of each of the borylated products were determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>49</sup> The resonances specific to each isomer are as follows:

ortho-isomer,  $\delta$  7.03 (t, J = 8.7 Hz, 1H + 2H of *p*-isomer), 7.13 (t, J = 7.3 Hz, 1H + 1H of *m*-isomer), 7.4-7.5 (m, 1H + 1H of *m*-isomer), 7.74 (m, 1H).

meta-isomer, δ 7.11–7.16 (m, overlapped, 1H), 7.31-7.37 (m, 1H), 7.41-7.47 (m, overlapped, 1H), 7.57 (d, *J* = 7.3 Hz, 1H).

para-isomer, 7.00-7.09 (m, overlapped, 1H), 7.80 (dd, J = 7.6, 2.3, 1H). (see below.)



For comparison, the iridium-catalyzed borylation of fluorobenzene was investigated. [Ir(OMe)(cod)]<sub>2</sub> (2.0 mg, 3.0 µmol), 4,4'-di-tert-butyl-2,2'-bipyridine (1.7 mg, 6.0 µmol), B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol) and fluorobenzene (141 µL, 1.5 mmol) were added to a 10 mL-sample vial with a Teflon-sealed screwcap. The cap was used to seal the vial under a flow of N<sub>2</sub>. The mixture was stirred at 60 °C for 20 h and the resulting mixture was filtered through silica gel (eluted with 7 mL of hexane/AcOEt = 5/1). The filtrate was analyzed by NMR spectroscopy using 1,1,2,2-tetrachloroethane as an internal standard (137% NMR yield, *o:m:p* = 41:48:11). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluted with hexane/AcOEt = 25/1) to give a mixture

of three isomers of the borylated products (63 mg, 94% based on  $B_2pin_2$ , o:m:p = 37:50:13, >99% purity by GC).



X : parts per Million : Proton

## 2-(2,5-Difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6b, Figure 6) [CAS: 408492-25-1].

 $\begin{array}{l} {\sf F} \\ {\sf F}$ 

### 2-(2,5-Dichlorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6c, Figure 6) [CAS: 479411-91-1].



Method B was used, except that the reaction was conducted using *p*-dichlorobenzene (221 mg, 1.5 mmol) instead of 1,4-difluorobenzene (154  $\mu$ L, 1.5 mmol). NMR analysis before the purification by column chromatography showed that a 75% yield

of the borylated product was formed.

Rf 0.54 (hexane/EtOAc = 5/1). White solid (38 mg, 47%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.37 (s, 12H), 7.23-7.34 (m, 2H), 7.65 (d, *J* = 2.3 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.8, 84.5, 130.7, 131.7, 132.1, 136.0, 137.7. HRMS (EI): Calcd for C<sub>12</sub>H<sub>15</sub>BCl<sub>2</sub>O<sub>2</sub> 272.0542, Found 272.0536.

## 2-(2,4-Dichloro-6-fluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6d, Figure 6) [CAS: 1192002-89-3].

 $CI \longrightarrow F_{Bpin}$  Method B was used, except that the reaction was conducted using 1,3-dichloro-5fluorobenzene (247 mg, 1.5 mmol) instead of 1,4-difluorobenzene. NMR analysis before purification by column chromatography showed that a 76% yield of the borylated product was formed.

Rf 0.51 (hexane/EtOAc = 5/1). colorless oil (42 mg, 47%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.39 (s, 12H), 6.97 (dd, J = 8.2, 1.4 Hz, 1H), 7.17 (t, J = 0.9 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.7, 85.0, 114.4 (d, J = 28 Hz), 125.1 (d, J = 2.9 Hz), 136.9 (d, J = 12 Hz), 139.2 (d, J = 12 Hz), 165.3 (d, J = 251 Hz). IR (ATR): 2980 w, 1598 m, 1563 w, 1538 w, 1471 w, 1420 w, 1390 m, 1370 m, 1330 s, 1271 w, 1213 w, 1166 w, 1141 s, 1101 s, 1061 w, 1042 w, 962 w, 931 m, 850 m, 837 s, 785 w, 670 w. MS m/z (% relative intensity): 294 (1), 292 (5), 290 (M+, 9), 277 (28), 276 (11), 275 (42), 257 (28), 255 (83), 254 (22), 233 (10), 231 (14), 215 (13), 213 (38), 212 (11), 194 (12), 192 (47), 191 (21), 190 (77), 189 (24), 155 (10), 131 (10), 115 (14), 111 (17), 109 (50), 85 (100), 75 (10), 74 (12), 69 (16), 59 (65), 58 (80), 57 (40), 55 (13). HRMS (EI): Calcd for C<sub>12</sub>H<sub>14</sub>BCl<sub>2</sub>FO<sub>2</sub> 290.0448, Found 290.0438.

## 4,4,5,5-Tetramethyl-2-(2,4,6-trifluorophenyl)-1,3,2-dioxaborolane (supplementary data) [CAS: 325143-04-2].

Rf 0.54 (hexane/EtOAc = 5/1). White solid (35 mg, 44%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.37 (s, 12H), 6.61 (t, *J* = 8.5 Hz, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.7, 84.2, 100.1, (ddd, *J* = 33, 24, 3.3 Hz), 165.0 (dd, *J* = 252, 16Hz), 167.3 (dt, *J* = 252 Hz, 15 Hz). HRMS (EI): Calcd for C<sub>12</sub>H<sub>14</sub>BF<sub>3</sub>O<sub>2</sub> 258.1039, Found 258.1038.

### 2-(3-Bromo-2,4,6-trifluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6e, Figure 6).

F F F Bpin

Method B was used, except that the reaction was conducted using in 2-bromo-1,3,5trifluorobenzene (176  $\mu$ L, 1.5 mmol) instead of 1,4-difluorobenzene at 60 °C. NMR analysis before purification by column chromatography showed that a 74% yield of the borylated product was formed.

Rf 0.49 (hexane/EtOAc = 5/1). Colorless oil (53 mg, 53%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.38 (s, 12H), 6.73 (td, *J* = 8.6 Hz, 2.0 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.8, 84.6, 93.6 (ddd, *J* = 38 Hz, 22 Hz, 7.2 Hz), 101.1 (ddd, *J* = 33 Hz, 26 Hz, 4.3 Hz), 161.9 (ddd, *J* = 253 Hz, 15 Hz, 6.7 Hz), 163.7 (ddd, *J* = 253 Hz, 16 Hz, 7.3 Hz), 165.9 (dt, *J* = 252 Hz, 15 Hz). IR (ATR): 2981 w, 1610 m,

1582 w, 1468 w, 1419 m, 1371 w, 1331 s, 1272 m, 1213 w, 1140 s, 1101 m, 1040 s, 962 w, 854 m, 836 m, 781 w, 743 w, 694 w, 673 w, 620 w. MS m/z (% relative intensity): 338 (M+2, 10), 336 (M+, 9), 323 (36), 322 (12), 321 (37), 279 (33), 278 (12), 277 (35), 276 (24), 274 (16), 239 (13), 236 (16), 234 (15), 232 (16), 85 (79), 69 (18), 59 (78), 58 (100), 57 (47). HRMS (EI): Calcd for C<sub>12</sub>H<sub>13</sub>BBrF<sub>3</sub>O<sub>2</sub> 336.0144, Found 336.0145.

### 2-(2,6-Difluoro-4-methoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6f, Figure 6) [CAS: 1111096-19-5].

Rf 0.43 (hexane/EtOAc = 5/1). Colorless oil (42 mg, 51%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.36 (s, 12H), 3.80 (s, 3H), 6.36-6.46 (m, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.7, 55.7, 83.7, 97.7 (d, *J* = 32 Hz), 163.9 (t, *J* = 14 Hz), 168.0 (dd, *J* = 250 Hz, 16 Hz). IR (ATR): 2978 w, 1629 m, 1572 m, 1461 w, 1431 m, 1377 m, 1349 m, 1333 s, 1272 m, 1198 m, 1138 s, 1090 m, 1033 m, 1020 m, 962 w, 856 m, 835 m, 738 w, 669 m, 613 w. MS m/z (% relative intensity): 270 (M+, 60), 269 (16), 255 (70), 254 (18), 212 (21), 211 (57), 210 (31), 197 (26), 185 (43), 181 (18), 171 (66), 170 (100), 169 (24), 156 (35), 141 (19), 140 (24), 105 (33), 85 (79), 77 (18), 75 (25), 69 (28), 59 (40), 58 (31), 57 (33). HRMS (EI): Calcd for C<sub>13</sub>H<sub>17</sub>BF<sub>2</sub>O<sub>3</sub> 270.1239, Found 270.1237.

### Borylation of *m*-difluorobenzene (6g, Figure 6).



Method B was used, except that the reaction was conducted using *m*-difluorobenzene (118  $\mu$ L, 1.5 mmol) instead of 1,4-difluorobenzene at 60 °C. NMR analysis before purification by column chromatography showed that a 61% yield (2:4:5 = 72:17:trace) of the borylated products was formed. After purification by flash column chromatography over silica gel (eluting with hexane/AcOEt = 25/1), a mixture of two isomers of the borylated products was obtained (24 mg, 34%, 2:4 = 83:17, >99% purity by GC). GC/MS analysis revealed the formation of three isomers of the borylated products, all of which had an m/z of 240 (M<sup>+</sup>). The identity and ratio of each of the borylated products was determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>50</sup> The resonances specific to each isomer are as follows:

2-borylated isomer,  $\delta$  6.84 (t, *J* = 7.8 Hz, 2H), 7.35 (m, 1H).

4-borylated isomer,  $\delta$  6.84, (td, J = 9.2 Hz, 2.3 Hz, 1H), 7.73 (q, J = 7.6 Hz, 1H).



# 2-(2,3-Difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6h, Figure 6) [CAS: 1073339-17-9].

Bpin Method B was used, except that the reaction was conducted at 60 °C using 1,2difluorobenzene (133  $\mu$ L, 1.5 mmol) instead of 1,4-difluorobenzene. NMR analysis before purification by column chromatography showed that a 70% yield of the borylated product was formed. A trace amount of 4-borylated product was observed by GCMS (GC area ratio 3-isomer:4isomer = 99:1)

Rf 0.51 (hexane/EtOAc = 5/1). Colorless oil (30 mg, 42%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.37 (s, 12H), 7.04-7.10 (m, 1H), 7.19-7.28 (m, 1H), 7.44-7.49 (m, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.9, 84.3, 120.2 (d, *J* = 17 Hz), 124.0 (dd, *J* = 4.7 Hz, 5.7 Hz), 131.1 (dd, *J* = 6.7 Hz, 3.8 Hz), 150.5 (dd, *J* = 249 Hz, 15 Hz), 154.6 (dd, *J* = 252 Hz, 12 Hz). HRMS (EI): Calcd for C<sub>12</sub>H<sub>15</sub>BF<sub>2</sub>O<sub>2</sub> 240.1133, Found 240.1141.

### 2-(2,3-Difluoro-4-methoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6i, Figure 6).

Bpin F F OMe Method B was used, except that the reaction was conducted at 80 °C using 1,2-difluoro-3-methoxybenzene (216 mg, 1.5 mmol) instead of 1,4-difluorobenzene. NMR analysis before purification by column chromatography showed that a 74% yield of the borylated product was formed. GC analysis of the crude reaction mixture revealed that three isomers

were formed in a ratio of 93/6/1 (The identity of the minor isomers was confirmed by GCMS). Spectroscopic data of the major isomer is shown below.

Rf 0.26 (hexane/EtOAc = 10/1). Colorless oil (25 mg, 31%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.35 (s, 12H), 3.92 (s, 3H), 6.73 (ddd, J = 8.2, 6.9, 1.4 Hz, 1H), 7.41 (ddd, J = 8.2, 6.0, 2.3 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.9, 56.7, 84.0, 108.2 (d, J = 2.9 Hz), 130.7 (dd, J = 9.1, 5.3 Hz), 141.0 (dd, J = 248, 16 Hz), 151.9, 155.6 (dd, J = 252 9.6 Hz). IR (ATR): 2979 w, 2938 w, 1628 m,

F + H + 10% F + H + 3.3%

1466 m, 1358 s, 1328 w, 1292 m, 1218 m, 1144 m, 1092 s, 972 m, 911 m. MS m/z (% relative intensity): 270 (M+, 54), 255 (39), 185 (50), 171 (43), 170 (100), 156 (34), 85 (56), 75 (37), 59 (37).

HRMS (EI): Calcd for C<sub>13</sub>H<sub>17</sub>BF<sub>2</sub>O<sub>3</sub> 270.1239, Found 270.1242.

The structure of the major isomer was unambiguously determined by NOE experiments, in which the following correlations were observed.

### 2-(2,4-Difluoro-3-methoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6j, Figure 6).



Method B was used, except that the reaction was conducted at 80 °C using 1,3-difluoro-2-methoxybenzene (216 mg, 1.5 mmol) instead of 1,4-difluorobenzene. NMR analysis before purification by column chromatography showed that an 86% yield of the borylated product was formed. GC analysis of the crude reaction mixture revealed that

two isomers were formed in a ratio of 92/8. Because of the unstable nature of the title compound, it was unable to isolate **29** in pure form by column chromatography [Rf 0.29 (hexane/EtOAc = 10/1)]. The identity of **29** was unambiguously confirmed by the following spectroscopic data:

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.36 (s, 12H), 3.97 (s, 3H), 6.88 (ddd, *J* = 10, 8.7, 1.4 Hz, 1H), 7.41 (dt, *J* = 8.3, 6.4 Hz, 1H).

HRMS (EI): Calcd for C<sub>13</sub>H<sub>17</sub>BF<sub>2</sub>O<sub>3</sub> 270.1239, Found 270.1241.

Isolated yield was determined by directly converting the crude mixture to the corresponding phenol derivative after oxidative work-up. Thus, an aqueous solution of NaOH (1.0 M, 0.30 mL) and an aqueous solution of H<sub>2</sub>O<sub>2</sub> (30%, 0.30 mL) were added to the crude reaction mixture at 0 °C and the mixture was stirred for 30 min at ambient temperature. A small amount of KI was added to quench the residual H<sub>2</sub>O<sub>2</sub> and the mixture was extracted three times with AcOEt. The combined organic extracts were concentrated in vacuo and purified by flash column chromatography over silica gel (eluted with hexane  $\rightarrow$  hexane/AcOEt = 4/1) to give 2,4-difluoro-3-methoxyphenol (**29-Ox**) as a colorless oil (24 mg, 50% yield based on B<sub>2</sub>pin<sub>2</sub>).



Rf 0.14 (hexane/EtOAc = 10/1). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.19 (s, 3H), 5.16 (bs, 1H), 6.63 (td, J = 9.2, 4.6 Hz, 1H), 6.77 (ddd, J = 11, 9.2, 2.3 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 62.0 (d, J = 2.9 Hz), 100.2 (d, J = 26 Hz), 109.8 (d, J = 9.6 Hz), 111.2 (dd, J = 21, 4.3 Hz), 140.9 (d, J = 13 Hz), 144.7 (d, J = 239 Hz), 149.6 (d, J = 241 Hz). IR (ATR): 3358 br, 2952 w, 1611 w,

1496 s, 1435 w, 1357 m, 1254 m, 1216 w, 1140 m, 1059 s, 1000 s, 949 w. MS m/z (% relative intensity): 160 (M+, 59), 145 (25), 117 (100), 88 (11), 71 (15), 69 (39), 57 (10). HRMS (EI): Calcd for  $C_7H_6F_2O_2$  160.0336, Found 160.0336.

### 2-(3-Chloro-2,4-difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (6k, Figure 6).

Bpin Method B was used, except that the reaction was conducted at 60 °C using 2-chloro-1,3difluorobenzene (223 mg, 1.5 mmol) instead of 1,4-difluorobenzene. NMR analysis before purification by column chromatography showed that a 82% yield of the borylated product was formed. GC analysis of the crude reaction mixture revealed that two isomers were formed in a ratio of 95/4 (The identity of the minor isomers was confirmed by GCMS). Spectroscopic data of the major isomer is shown below.

Rf 0.40 (hexane/EtOAc = 10/1). Colorless oil (49 mg, 60%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.36 (s, 12H), 6.94-6.99 (m, 1H), 7.59-7.64 (m, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.9, 84.4, 112.0 (dd, *J* = 20, 3.8 Hz), 117.5 (d, *J* = 22 Hz), 134.9 (t, *J* = 9.6 Hz), 161.1 (d, *J* = 213 Hz), 163.6 (d, *J* = 207 Hz). IR (ATR): 2980 w, 2932 w, 1610 m, 1584 w, 1427 m, 1384 m, 1355 s, 1264 m, 1225 w, 1143 s, 1110 m, 1015 m, 963 m, 863 m. MS m/z (% relative intensity): 274 (M+, 20), 259 (82), 215 (63), 212 (30), 177 (31), 175 (84), 174 (36), 170 (52), 109 (34), 93 (39), 85 (92), 59 (87), 58 (100), 57 (64). HRMS (EI): Calcd for C<sub>12</sub>H<sub>14</sub>BClF<sub>2</sub>O<sub>2</sub> 274.0743, Found 274.0742.

### 1-Methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (8a, Figure 7) [CAS: 596819-10-2].

Rf 0.63 (hexane/EtOAc = 5/1). White solid (69 mg, 87%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.36 (s, 12H), 3.97 (s, 3H), 7.08 (td, *J* = 6.9 Hz, 0.9 Hz, 1H), 7.13 (d, *J* = 0.9 Hz, 1H), 7.25 (td, *J* = 8.2 Hz, 0.9 Hz, 1H), 7.34 (dd, *J* = 8.2 Hz, 0.9 Hz, 1H), 7.64 (dd, *J* = 8.2 Hz, 0.9 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.8, 32.2, 83.7, 109.7, 114.2, 119.2, 121.5, 123.4, 127.8, 140.1. HRMS (EI): Calcd for C<sub>15</sub>H<sub>20</sub>BNO<sub>2</sub> 257.1587, Found 257.1584.

## Methyl 1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-5-carboxylate (8b, Figure7).



Method C was used, except that the reaction was conducted using methyl 1methyl-1H-indole-5-carboxylate (56 mg, 0.30 mmol), 4 mol % of Pt(ICy)(dvtms) (7.2 mg, 1.2 μmol) and B<sub>2</sub>pin<sub>2</sub> (92 mg, 0.36 mmol). Rf 0.34 (hexane/EtOAc = 5/1). White solid (59 mg, 63%). Mp = 145-146 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.37 (s, 12H), 3.93 (s, 3H), 3.98 (s, 3H), 7.21 (s, 1H), 7.34 (d, *J* = 8.7 Hz, 1H), 7.95 (dd, *J* = 8.7 Hz, 1.6 Hz, 1H), 8.40 (d, *J* = 0.9 Hz, 1H). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz):  $\delta$  1.09 (s, 12H), 3.51 (s, 3H), 3.61 (s, 3H), 6.95 (d, *J* = 8.7 Hz, 1H), 7.50 (d, *J* = 0.9 Hz, 1H), 8.31 (dd, *J* = 8.9, 1.6 Hz, 1H), 8.71 (d, *J* = 0.9 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.9, 32.6, 51.9, 84.0, 109.4, 115.9, 121.5, 124.4, 124.9, 127.4, 142.4, 168.3. IR (ATR): 2977 w, 2952 w, 1709 m, 1610 w, 1533 m, 1452 m, 1438 m, 1389 m, 1371 m, 1348 m, 1305 s, 1266 m, 1248 s, 1213 s, 1194 m, 1155 m, 1137 s, 1109 m, 1086 m, 1064 m, 973 w, 963 m, 904 w, 862 w, 846 w, 830 w, 806 w, 768 w, 734 m, 705 w, 688 m, 669 m, 625 w, 607 w. MS m/z (% relative intensity): 316 (18), 315 (M+, 100), 314 (26), 284 (21), 230 (29),



215 (11), 202 (18), 184 (22), 156 (12), 77 (13). HRMS (CI): Calcd for  $[C_{17}H_{22}BNO_4 + H]$  316.1720, Found 316.1728. The regiochemistry of the compound was determined by NOE experiments in C<sub>6</sub>D<sub>6</sub>. When the signal for H<sub>a</sub> at  $\delta = 8.71$  (d, J = 0.9 Hz, 1H, offset signal  $\delta = 8.655$ , mixing time 1 s) was irradiated, H<sub>b</sub> at  $\delta = 7.50$  (d, J = 0.9 Hz, 1H) was enhanced by 2%.

5-Methoxy-1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (8c, Figure 7) [CAS: 1256360-41-4].



Method C was used, except that the reaction was conducted with 5-methoxy-1methyl-1H-indole (48 mg, 0.30 mmol).

Me Rf 0.51 (hexane/EtOAc = 5/1). White solid (67 mg, 79%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz): δ 1.36 (s, 12H), 3.84 (s, 3H), 3.94 (s, 3H), 6.93 (dd, J = 9.2 Hz, 2.3 Hz, 1H), 7.03 (s, 1H), 7.05 (d, J = 2.8 Hz, 1H), 7.22 (d, J = 9.2 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): δ 24.8, 32.3, 55.8, 83.6, 102.1, 110.4, 113.4, 114.3, 127.9, 135.8, 153.9. HRMS (CI): Calcd for [C<sub>16</sub>H<sub>22</sub>BNO<sub>3</sub> + H] 288.1771, Found 288.1782.

### 5-Fluoro-1-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (8d, Fig. 2D) [CAS: 1683582-67-3].



Method C was used, except that the reaction was conducted with 5-fluoro-1methyl-1H-indole (42 mg, 0.28 mmol), 4 mol% of Pt(ICy)(dvtms) (7.2 mg, 0.012 mmol) and B<sub>2</sub>pin<sub>2</sub> (92 mg, 0.36 mmol).

Rf 0.51 (hexane/EtOAc = 5/1). White solid (44 mg, 56%). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 395.88 MHz)  $\delta$  1.09 (s, 12H), 3.56 (s, 3H), 6.78 (dd, *J* = 8.7 Hz, 4.1 Hz, 1H), 6.99 (td, *J* = 9.0 Hz, 2.4 Hz, 1H), 7.27 (dd, *J* = 9.6 Hz, 2.3 Hz, 1H), 7.36 (s, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz):  $\delta$  24.8, 32.1, 83.7, 106.3 (d, *J* = 22 Hz), 110.7 (d, *J* = 9.6 Hz), 112.2 (d, *J* = 27 Hz), 115.0 (d, *J* = 4.8 Hz), 128.6 (d, *J* = 11Hz), 137.4, 158.4 (d, *J* = 233 Hz). HRMS (EI): Calcd for C<sub>15</sub>H<sub>19</sub>BFNO<sub>2</sub> 275.1493, Found 275.1493.

1,3-Dimethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole (supplementary data) [CAS: 1683582-66-2].

Me Method C was used using 1,3-dimethyl-1H-indole (45 mg, 0.31 mmol).

Rf 0.49 (hexane/EtOAc = 5/1). White solid (73 mg, 87%). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz): δ 1.10 (s, 12H), 2.84 (s, 3H), 3.72 (s, 3H), 7.13 (d, J = 8.7 Hz, 1H), 7.19 (t, J = 7.6 Hz, 1H), 7.31 (dt, J = 8.2 Hz, 0.9 Hz, 1H), 7.73 (dd, J = 7.8, 0.9 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz): δ 10.6, 24.7, 31.9, 82.9, 109.7, 118.9, 120.2, 123.7, 125.5, 129.2, 140.5. HRMS (EI): Calcd for C<sub>16</sub>H<sub>22</sub>BNO<sub>2</sub> 271.1744, Found 271.1742.

## 1-Methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole (8e, Figure 7) [CAS: 850567-47-4].

Method C was used, except that the reaction was conducted with 1-methylpyrrole (133  $\mu$ L, 1.5 mmol, 5 equiv.) and B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol).

Rf 0.14 (hexane/EtOAc = 5/1). White solid (47 mg, 77% based on B<sub>2</sub>pin<sub>2</sub>). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz):  $\delta$  1.01 (s, 12H), 3.51 (s, 3H), 6.30 (dd, *J* = 3.7 Hz, 2.7 Hz, 1H), 6.51-6.52 (m, 1H), 7.33 (dd, *J* = 1.3 Hz, 3.7 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz):  $\delta$  24.8, 36.2, 82.9, 109.1, 123.3, one carbon peak is overlapped with residual solvent peaks. <sup>13</sup>C NMR(CDCl<sub>3</sub>, 100.53 MHZ)  $\delta$  24.8, 36.6, 83.0, 108.3, 121.9, 128.2. HRMS (EI): Calcd for C<sub>11</sub>H<sub>18</sub>BNO<sub>2</sub> 207.1431, Found 207.1431. A trace amount of a diborylated product was detected by GC-MS analysis of the crude reaction mixture (m/z = 333).

## 1-Methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-aza-1*H*-indole (8f, Figure 7) [CAS: 912331-71-6].

Method C was used, except that the reaction was conducted with 1-methyl-7-aza-1*H*-<sup>Bpin</sup> indole (39 mg, 0.29 mmol), 4 mol % of Pt(ICy)(dvtms) (7.2 mg, 0.012 mmol) and  $B_2pin_2$  (92 mg, 0.36 mmol).

Rf 0.14 (hexane/EtOAc = 5/1). White solid (47 mg, 62%). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz):  $\delta$  1.08 (s, 12H), 4.09 (s, 3H), 6.75 (dd, *J* = 4.6 Hz, 7.8 Hz, 1H), 7.37 (s, 1H), 7.60 (dd, *J* = 7.8 Hz, 1.8 Hz, 1H), 8.48 (dd, *J* = 4.6 Hz, 1.4 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz):  $\delta$  24.5, 30.6, 83.5, 112.7, 115.6, 120.0, 129.1, 145.1, 150.9. HRMS (EI): Calcd for C<sub>14</sub>H<sub>19</sub>BN<sub>2</sub>O<sub>2</sub> 258.1540, Found 258.1537.

### 2-(Benzofuran-2-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (8g, Figure 7) [CAS: 402503-13-3].



Мe

Method C was used, except that the reaction was conducted with benzofuran (35 mg, 0.30 mmol), 4 mol% of Pt(ICy)(dvtms) (7.2 mg, 0.012 mmol) and B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol).

The product was obtained as a mixture of 2- and 3- borylated isomers (**35-2B**: **35-3B** = 75:25). The ratio of the isomers was determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>51</sup>

Rf 0.51 (hexane/EtOAc = 5/1). White solid (34 mg, 47%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.37 (s, 12H of **35-3B**), 1.39 (s, 12H of **35-2B**), 7.23 (t, *J* = 7.3 Hz, 1H of **35-2B**), 7.25-7.30 (m, 2H of **35-3B**), 7.34 (td, *J* = 7.8, 0.9 Hz, 1H of **35-2B**), 7.40 (s, 1H of **35-2B**), 7.50 (dd, *J* = 8.2, 1.8 Hz, 1H of **35-2B**), 7.57 (d, *J* = 8.2 Hz, 1H of **35-2B**), 7.63 (d, *J* = 7.3 Hz, 1H of **35-3B**), 7.91-7.93 (m, 1H of **35-3B**), 7.95 (s, 1H of **35-3B**). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.8 (**35-2B**), 24.9 (**35-3B**), 83.5 (**35-3B**), 84.7 (**35-2B**), 111.0 (**35-3B**), 112.0 (**35-2B**), 119.5 (**35-2B**), 121.9 (**35-2B**), 122.7 (**35-2B**), 122.87 (**35-3B**), 122.94 (**35-3B**), 124.2 (**35-3B**), 125.9 (**35-2B**), 127.5 (**35-2B**), 153.7 (**35-3B**), 157.5 (**35-2B**). HRMS (EI): Calcd for C<sub>14</sub>H<sub>17</sub>BO<sub>3</sub> 244.1271, Found 244.1276.

# 4,4,5,5-Tetramethyl-2-(5-methylthiophen-2-yl)-1,3,2-dioxaborolane (8h, Figure 7) [CAS: 476004-80-5].

Method A was used, except that the reaction was conducted in 2-methylthiophene (1 mL).

Rf 0.49 (hexane/EtOAc = 5/1). Colorless oil (47 mg, 70%). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz):  $\delta$  1.08 (s, 12H), 2.09 (s, 3H), 6.61 (dd, *J* = 0.9 Hz, 3.3 Hz, 1H), 7.77 (d, *J* = 3.2 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz):  $\delta$  14.8, 24.5, 83.5, 127.1, 138.0, 147.5. HRMS (EI): Calcd for C<sub>11</sub>H<sub>17</sub>BO<sub>2</sub>S 224.1042, Found 224.1045.

## 2-(5-Methoxythiophen-2-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (8i, Figure 7) [CAS: 596819-12-4]].

Method A was used, except that the reaction was conducted in 2-methoxythiophene (1 mL).

Rf 0.49 (hexane/EtOAc = 5/1). colorless oil (45 mg, 64%).<sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MeO s Hz):  $\delta$  1.32 (s, 12H), 3.92 (s, 3H), 6.30 (d, *J* = 4.1 Hz, 1H), 7.33 (d, *J* = 3.7 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.7, 60.3, 83.8, 106.1, 136.5, 172.8. IR (ATR): 2977 w,2933 w, 1546 w, 1479 s, 1422 s, 1389 m, 1362 s, 1333 m, 1298 m, 1271 w, 1239 m, 1208 m, 1140 s, 1111 w, 1060 s, 987 m, 955 w, 853 m, 830 w, 778 m, 724 w, 682 w, 660 s, 574 m. MS m/z (% relative intensity): 241 (15), 240 (M+, 100), 239 (31), 225 (20), 197 (34), 181 (16), 180 (75), 179 (20), 167 (19), 155 (41), 141 (63), 140 (95), 139 (35), 125 (32), 124 (15), 115 (15), 98 (16), 97 (28), 85 (20), 83 (52), 59 (25), 57 (18), 55 (31). HRMS (EI): Calcd for C<sub>11</sub>H<sub>17</sub>BO<sub>3</sub>S 240.0991, Found 240.0998.

The regiochemistry of the compound was determined by NOE experiments in CDCl<sub>3</sub>. When the signal



for H<sub>a</sub> at  $\delta$  = 7.33 (d, *J* = 3.7 Hz, 1H, offset signal  $\delta$  = 7.3093, mixing time 1 s) was irradiated, H<sub>b</sub> at  $\delta$  = 6.30 (d, *J* = 4.1 Hz, 1H) was enhanced by 2%. When the signal for H<sub>b</sub> at  $\delta$  = 6.30 (d, *J* = 4.1 Hz, 1H, offset signal  $\delta$  = 6.2818, mixing time 1 s) was irradiated, H<sub>c</sub>  $\delta$  = 3.92 (s, 3H) was enhanced by 1%.

2-(Benzo[b]thiophen-2-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (8j, Figure 7) [CAS: 376584-76-8]].



Method C was used, except that the reaction was conducted with benzo[b]thiophene (40 mg, 0.30 mmol), 4 mol% of Pt(ICy)(dvtms) (7.2 mg, 0.012 mmol) and B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol).

The product was obtained as a mixture of 2- and 3-borylated isomers (**38-2B**: **38-3B** = 92:8). The ratio of the isomers was determined by comparing the <sup>1</sup>H NMR spectrum of the product mixture with those reported in the literature.<sup>52</sup>

Rf 0.49 (hexane/EtOAc = 5/1). white solid (34 mg, 44%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.38 (s, 12H of **38-2B**, 12H of **38-3B**), 7.33-7.39 (m, 2H of **38-2B**, 2H of **38-3B**), 7.84-7.92 (m, 3H of **38-2B**, 1H of **38-3B**), 8.07 (s, 1H of **38-3B**), 8.37 (d, *J* = 7.8 Hz, 1H of **38-3B**). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz): **38-2B**;  $\delta$  24.8, 84.4, 122.5, 124.1, 124.4, 125.3, 134.5, 140.4, 143.7. HRMS (EI): Calcd for C<sub>14</sub>H<sub>17</sub>BO<sub>2</sub>S 260.1042, Found 260.1042.

### 2-(5-Chloro-3-methylbenzo[b]thiophen-2-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (8k, Figure 7).



Method C was used, except that the reaction was conducted with 5-chloro-3methylbenzo[b]thiophene (55 mg, 0.30 mmol) and  $B_2pin_2$  (92 mg, 0.36 mmol). Rf 0.49 (hexane/EtOAc = 5/1). White solid (59 mg, 64%). Mp = 141-142 °C. <sup>1</sup>H

NMR (C<sub>6</sub>D<sub>6</sub>, 399.78 MHz):  $\delta$  1.06 (s, 12H), 2.50 (s, 3H), 7.03 (dd, J = 8.5, 2.1 Hz, 1H), 7.17 (d, J = 8.7 Hz, 1H), 7.62 (d, J = 1.8 Hz, 1H). <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>, 100.53 MHz):  $\delta$  13.6, 24.5, 83.8, 122.4, 123.7, 125.8, 130.2, 141.6, 142.5, 143.1. IR (ATR): 2975 w, 1557 w, 1526 m, 1437 w, 1361 s, 1306 s, 1263 m, 1248 m, 1207 w, 1166 w, 1136 s, 1105 s, 1077 m, 1029 m, 975 m, 947 w, 867 m, 848 s, 806 s, 780 w, 692 w, 671 s, 608 m, 586 w, 557 w, 544 w, 534 w. MS m/z (% relative intensity): 310 (34), 309 (23), 308 (M+, 100), 307 (23), 253 (25), 252 (19), 251 (71), 250 (36), 222 (42), 216 (19), 210 (38), 209 (44), 208(100), 207 (39), 181 (17), 173 (27), 149 (18), 85 (27), 59 (26), 57 (15). HRMS (CI): Calcd for [C<sub>15</sub>H<sub>18</sub>BClO<sub>2</sub>S + H]309.0887, Found 309.0876.

### 7. Synthetic Applications

#### 7-1. Scale-up Experiment (Scheme 2)



To a dried two-necked flask equipped with a reflux condenser,  $B_2pin_2$  (1.25 g, 4.9 mmol), Pt(SIPr)(dvtms) (63 mg, 0.082 mmol), 2v (1.17 g, 4.1 mmol) and octane (1.4 mL) were added. The mixture was heated at 120 °C for 28 h. The resulting mixture was filtered through a column of silica gel (eluting with 7 mL of hexane/EtOAc = 5/1) to give the target borylated product (104 mg) and a mixture of the target product and unreacted starting material. The mixture was again transferred to another dried two-necked flask,  $B_2pin_2$  (1.25 g, 4.9 mmol), Pt(SIPr)(dvtms) (63 mg, 0.082 mmol) and octane (1.4 mL) were added. The reaction mixture was heated at 120 °C for 34 h. The resulting mixture was purified by column chromatography to give the target borylated product (1.17 g) and unreacted starting material (129 mg, 11%). As a result of two borylation, 1.28 g of the target product was obtained as a white solid (1.28g, 76%).

#### 7.2.1 Gram-Scale Synthesis of 3k and Its Transformations (3k, Scheme 5)



Pt(IPr)(dvtms) (91 mg, 0.12 mmol), B<sub>2</sub>pin<sub>2</sub> (1.50 g, 5.9 mmol) and 1-(tert-butyl)-3,5-dimethylbenzene (20 mL) were added to a 100 mL three-necked flask with a condenser under a N<sub>2</sub> atmosphere. The mixture was stirred at 120 °C for 20 h, and the resulting mixture was purified by flash column chromatography over silica gel (eluted with hexane to hexane/AcOEt = 95/5) to give 2-(4-(tert-butyl)-2,6-dimethylphenyl)-4,4,5,5-tetramethyl -1,3,2-dioxaborolane as a white solid (1.40 g, 82% based on B<sub>2</sub>pin<sub>2</sub>).

### 7.2.2 Synthetic Elaboration of the Borylated Product

#### Methyl 4'-(tert-butyl)-2',6'-dimethyl-[1,1'-biphenyl]-4-carboxylate (9, Scheme 5).



To a 10 mL sample vial with a Teflon-sealed screwcap, **3k** (86 mg, 0.30 mmol), methyl 4-iodobenzoate (87 mg, 0.33 mmol),  $Pd(OAc)_2$  (3.3 mg, 0.015 mmol), SPhos (12 mg, 0.030 mmol), toluene (1.0 mL), and an aqueous solution (0.50 mL) of K<sub>3</sub>PO<sub>4</sub> (160 mg, 0.75 mmol) was added under an atmosphere of nitrogen. The resulting mixture was stirred at 120

°C for 12 h under an atmosphere of nitrogen. The mixture was then partitioned between  $Et_2O$  and  $H_2O$ , and the organic layer was separated. The extract was washed with brine, dried (MgSO<sub>4</sub>) and concentrated. The residue was purified by column chromatography (SiO<sub>2</sub>, hexane/EtOAc = 95/5 to 85/15), followed by GPC, to give the title compound as a colorless oil (48 mg, 54%).

Rf 0.43 (hexane/EtOAc = 10/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.35 (s, 9H), 2.02 (s, 6H), 3.94 (s, 3H), 7.13 (s, 2H), 7.25 (d, *J* = 8.7 Hz, 2H), 8.09 (d, *J* = 8.7 Hz, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub> 100.53 MHz):  $\delta$  21.2, 31.5, 34.5, 52.2, 124.6, 128.6, 129.6, 129.8, 135.2, 138.1, 146.6, 150.4, 167.3.

IR (ATR): 2954 w, 1724 s, 1610 w, 1435 m, 1272 s, 1175 w, 1112 m, 1100 m, 1005 w. MS m/z (% relative intensity): 296 (M<sup>+</sup>, 26), 282 (21), 281 (100), 111 (24). HRMS (EI): Calcd for  $C_{20}H_{24}O_2$  296.1776, Found 296.1770.

#### 2-Allyl-5-(tert-butyl)-1,3-dimethylbenzene (10, Scheme 5).

To a 10 mL sample vial with a Teflon-sealed screwcap, **3k** (86 mg, 0.30 mmol), allyl chloride (69 mg, 0.90 mmol), Pd(dba)<sub>2</sub> (8.6 mg, 0.015 mmol), and MeOH (1.0 mL) was added, and the mixture was stirred at room temperature for 10 min. KF (52 mg, 0.90 mmol) was then added, and the resulting mixture was stirred at

room temperature for 18 h under an atmosphere of nitrogen. The mixture was then partitioned between  $Et_2O$  and  $H_2O$ , and the organic layer was separated. The extract was washed with brine, dried (MgSO<sub>4</sub>) and concentrated. The residue was purified by column chromatography (SiO<sub>2</sub>, hexane to hexane/EtOAc = 95/5) to give the title compound as a colorless oil (49 mg, 81%).

Rf 0.80 (hexane/EtOAc = 10/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.29 (s, 9H), 2.29 (s, 6H), 3.36 (dt, *J* = 5.5, 1.8 Hz, 2H), 4.90 (dq, *J* = 17.4 Hz, 1.8 Hz, 1H), 4.98 (dq, *J* = 10.1, 1.8 Hz, 1H), 5.88 (ddt, *J* = 17.4, 10.1, 5.5 Hz, 1H), 7.03 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub> 100.53 MHz):  $\delta$  20.1, 31.4, 33.5, 34.1, 114.8, 125.0, 133.1, 135.5, 136.1, 148.6. IR (ATR): 2961 s, 2867 m, 1729 m, 1637 w, 1609 w, 1575 w, 1485 m, 1455 m, 1361 m, 1274 s, 1233 w, 1191 w, 1113 m, 1100 m, 995 m, 910 s, 868 s. MS m/z (% relative intensity): 202 (M<sup>+</sup>, 29), 188 (16), 187 (100), 145 (18), 133 (19), 131 (11), 129 (11), 128 (11). HRMS (EI): Calcd for C<sub>15</sub>H<sub>22</sub> 202.1722, Found 202.1720.

#### 4-(tert-Butyl)-2,6-dimethylaniline (11, Scheme 5) [CAS: 42014-60-8].



Me

Me

Me

To a mixture of **3k** (86 mg, 0.30 mmol) and  $H_2NOSO_3H$  (85 mg, 0.75 mmol) in CH<sub>3</sub>CN (2.5 mL), an aqueous solution of NaOH (1 M, 2.5 mL) was added. The resulting mixture was stirred at room temperature under an atmosphere of nitrogen for 12 h. The mixture was then partitioned between Et<sub>2</sub>O and H<sub>2</sub>O, and the organic

layer was separated. The extract was washed with brine, dried (MgSO<sub>4</sub>) and concentrated. The residue was purified by column chromatography (SiO<sub>2</sub>, hexane/EtOAc = 95/5 to 85/15) to give the title compound as a colorless oil (17 mg, 34%). The starting **3k** was also recovered as a colorless solid (55 mg, 64%).

Rf 0.26 (hexane/EtOAc = 10/1). Colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.28 (s, 9H), 2.19 (s, 6H), 2.50-4.55 (br, 2H), 6.97 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub> 100.53 MHz):  $\delta$  17.9, 31.6, 33.7, 121.4, 125.1, 140.2, 140.8. HRMS (EI): Calcd for C<sub>12</sub>H<sub>19</sub>N 177.1517, Found 177.1518.

### 4-(tert-Butyl)-2,6-dimethylphenol (12, Scheme 5) [CAS: 879-97-0].



To a solution of **3k** (60 mg, 0.21 mmol) in EtOH (3.0 mL), an aqueous solution of  $H_2O_2$  (30%, 1.0 mL) and an aqueous solution of NaOH (1M, 1.0 mL) were added sequentially. After being stirred at room temperature for 2 h, the mixture was concentrated in vacuo, and the residue was partitioned between EtOAc and an

aqueous solution of HCl (1 M). The separated organic layer was washed with brine, dried (MgSO<sub>4</sub>)

and concentrated. The residue was purified by column chromatography (SiO<sub>2</sub>, hexane/EtOAc = 95/5 to 85/15) to give the title compound as a colorless solid (35 mg, 93%).

Rf 0.31 (hexane/EtOAc = 10/1). White solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.28 (s, 9H), 2.24 (s, 6H), 4.48 (br, 1H), 6.99 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub> 100.53 MHz):  $\delta$  16.2, 31.6, 33.9, 122.3, 125.5, 142.9, 149.9. HRMS (EI): Calcd for C<sub>12</sub>H<sub>18</sub>O 178.1358, Found 309. 1358.

### 7.3. Orthogonal Functionalization of 3y(3',4'-Dimethoxy-2,6-dimethyl-[1,1'-biphenyl]-4-yl)triethylsilane (13, Scheme 7).



To a 10 mL-sample vial with a Teflon-sealed screwcap, **3y** (15 mg, 0.043 mmol), Pd(OAc)<sub>2</sub> (0.5 mg, 0.0022 mmol), SPhos (1.8 mg, 0.0043 mmol), K<sub>3</sub>PO<sub>4</sub> (27 mg, 0.13 mmol) and 4-bromo-1,2dimethoxybenzene (11 mg, 0.052 mmol) were added. After adding H<sub>2</sub>O (0.3 mL) and 1,2dimethoxyethane (0.3 mL), the cap was applied to seal the vial under the flow of N<sub>2</sub>. The mixture was stirred at 100 °C until GC and TLC analysis showed full conversion of starting **3y** (4.5 h). To the resulting reaction mixture, H<sub>2</sub>O was added and the aqueous phase was extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with brine and dried over MgSO<sub>4</sub>. The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/EtOAc = 100/1 to 50/1) to give the title compound as a colorless oil (11 mg, 71%).

Rf 0.46 (hexane/EtOAc = 5/1). Colorless oil (11 mg, 71%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  0.81 (q, J = 7.9 Hz, 6H), 1.01 (t, J = 7.8 Hz, 9H), 2.07 (s, 6H), 3.85 (s, 3H), 3.93 (s, 3H), 6.65-6.74 (m, 2H), 6.93 (d, J = 8.2 Hz, 1H), 7.21 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  3.42, 7.53, 20.9, 55.8, 55.9, 111.0, 112.2, 121.0, 133.1, 133.7, 135.4, 135.9, 142.1, 147.5, 148.7. IR (ATR): 2952 m, 2909 m, 2874 m, 1512 s. 1461 m, 1403 w, 1320 w, 1294 w, 1242 s, 1214 m, 1172 m, 1136 s, 1028 m, 970 w, 861 w, 841 w, 808 w, 767 w, 733 s, 719 s, 700 m. MS m/z (% relative intensity): 357 (13), 356 (M+, 45), 327 (52), 299 (71), 272 (23), 271 (100), 135 (44), 114 (12). HRMS (EI): Calcd for C<sub>22</sub>H<sub>32</sub>O<sub>2</sub>Si 356.2172, Found 356.2172.

#### 2-(4-Bromo-2,6-dimethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (14, Scheme 7).



In a 10 mL-sample vial, 3y (21 mg, 0.060 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (0.3 mL) under N<sub>2</sub> atmosphere. The solution was cooled to 0 °C in an ice bath. To the cooled solution, Br<sub>2</sub> solution (11

mg, 0.069 mmol, dissolved in 0.3 mL of  $CH_2Cl_2$ ) was added dropwise. The resulting mixture was stirred at 0 °C for 1 h. Since TLC showed that starting **3y** remained, additional Br<sub>2</sub> solution (5.0 mg, 0.031 mmol, dissolved in 0.3 mL of  $CH_2Cl_2$ ) were added and the mixture was stirred for another 30 min. To the resulting mixture, Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> aq was added and the mixture was extracted with Et<sub>2</sub>O three times. Combined organic extracts were washed with brine and dried over MgSO<sub>4</sub>. The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/AcOEt = 200/1 to 100/1) to give the title compound as a white solid (16 mg, 83%).

Rf 0.60 (hexane/EtOAc = 5/1). White solid (16 mg, 83%). Mp = 101-103 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.38 (s, 12H), 2.36 (s, 6H), 7.11 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  22.0, 24.9, 83.9, 123.5, 129.3, 144.0. IR (ATR): 2977 m, 1575 m, 1443 w, 1401 w, 1369 s, 1334 s, 1301 s, 1272 w, 1211 w, 1164 w, 1145 m, 1111 m, 1063 m, 851 m, 677 m. MS m/z (% relative intensity): 312 (44), 310 (M+, 43), 255 (94), 254 (50), 253 (100), 252 (43), 213 (57), 212 (99), 211 (77), 210 (98), 209 (24), 174 (69), 131 (38), 105 (26), 101 (27), 83 (23). HRMS (EI): Calcd for C<sub>14</sub>H<sub>20</sub>BBrO<sub>2</sub> 310.0740, Found 310.0737.





Pt(SIPr)(dvtms) (6.9 mg, 9.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (91 mg, 0.36 mmol), methyl 4-methylbenzoate (49 mg, 0.30 mmol) and 1,3,5-triisopropylbenzene (0.10 mL) were added to a 10 mL-sample vial with a Teflonsealed screwcap. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The mixture was stirred at 120 °C for 20 h and the resulting mixture was filtered through a pad of silica gel (eluting with 7 mL of hexane/EtOAc = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography over silica gel (eluting with hexane/AcOEt = 100/1 to 50/1) to give the target borylated product as a white solid (35 mg, 42%) and 26 mg of unreacted starting material was recovered (52%). The recovered starting material was transferred to another vial for repeating the borylation. Pt(SIPr)(dvtms) (4.9 mg, 6.4  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (51 mg, 0.20 mmol) and 1,3,5-triisopropylbenzen (0.1 mL) were added. The mixture was stirred at 120 °C for 16 h and the resulting mixture was filtered through a pad of silica gel (eluting with 7 mL of hexane/AcOEt = 5/1). The filtrate was concentrated in vacuo and purified by flash column (0.1 mL) were added. The mixture was stirred at 120 °C for 16 h and the resulting mixture was filtered through a pad of silica gel (eluting with 7 mL of hexane/AcOEt = 5/1). The filtrate was concentrated in vacuo and purified by flash column chromatography to give the target borylated product (14 mg, 17% based on starting arene) and 7.9 mg of unreacted starting material was recovered. As a result of two borylation, 59% of the target product was obtained and 16% of unreacted starting was recovered.

Rf 0.49 (hexane/EtOAc = 5/1). White solid (49 mg, 59%). Mp = 158-159 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.34 (s, 12H), 2.56 (s, 3H), 3.88 (s, 3H), 7.21 (d, *J* = 7.8 Hz, 1H), 7.95 (dd, *J* = 8.2, 1.8 Hz, 1H), 8.39 (d, *J* = 1.4 Hz, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  22.4, 24.9, 51.9, 83.7, 126.7, 129.9, 131.8, 137.0, 150.4, 167.3. IR (ATR): 2976 w, 1713 s, 1602 w, 1403 m, 1368 m, 1343 s, 1316 w, 1280 s, 1255 s, 1134 s, 1116 s, 1068 m, 1005 w, 969 m, 934 w, 855 m, 759 m, 733 w, 680 w, 654 m. MS m/z (% relative intensity): 276 (M+, 10), 261 (33), 245 (21), 233 (35), 219 (100), 218 (55), 201 (14), 177 (87), 176 (56), 145 (55), 144 (15), 117 (43), 85 (25), 59 (20). HRMS (EI): Calcd for C<sub>15</sub>H<sub>21</sub>BO<sub>4</sub> 276.1533, Found 276.1530.

The regiochemistry of the compound was determined by NOESY experiments.



### 8. Mechanistic Studies

8-1. Diborylation of dvtms with Pt(ICy)dvtms catalyst.



To a 10 mL-sample vial with a Teflon-sealed screwcap,  $B_2pin_2$  (127 mg, 0.50 mmol), Pt(ICy)dvtms (3.7 mg, 0.0060 mmol), dvtms (37 mg, 0.20 mmol) and octane (0.5 mL) were added. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The mixture was stirred at 100 °C for 16 h. The resulting mixture was analyzed by TLC and GC to show full conversion of starting dvtms. An aliquot of the reaction mixture was taken and diluted with 0.5 mL of EtOAc for APCI-MS analysis. APCI-MS analysis showed the target tetraborylated product exists in the reaction mixture. HRMS(APCI): Calcd for [C<sub>32</sub>H<sub>66</sub>B<sub>4</sub>O<sub>9</sub>Si<sub>2</sub>+H] 695.4713, Found 695.4717.

8-2. Diborylation of diphenylacetylene with Pt(ICy)dvtms catalyst.



To a 10 mL-sample vial with a Teflon-sealed screwcap, diphenylacetylene (36 mg, 0.20 mmol), Pt(ICy)dvtms (3.7 mg, 0.0060 mmol), B<sub>2</sub>pin<sub>2</sub> (51 mg, 0.20 mmol) and octane (0.5 mL) were added. The cap was applied to seal the vial under the flow of N<sub>2</sub>. The mixture was stirred at 100 °C for 16 h. The resulting mixture was filtered through a pad of silica gel (eluting with 20 mL of hexane/EtOAc = 5/1). The filtrate was concentrated in vacuo to give a white solid. The solid was washed with minimal amount of hexane to give the pure target product as a white solid (54 mg, 62%) Mp = 178 °C <sup>-1</sup>H NMR (CDCl<sub>2</sub> 399.78

Rf 0.46 (hexane/EtOAc = 5/1). White solid (54 mg, 62%). Mp = 178 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 399.78 MHz):  $\delta$  1.32 (s, 24H), 6.83-6.98 (m, 4H), 6.98-7.16 (m, 6H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100.53 MHz):  $\delta$  24.9, 84.0, 125.6, 127.4, 129.3, 141.3. HRMS (EI): Calcd for C26H34B2O4 432.2643, Found 432.2643. All spectroscopic data were in agreement with the reported *cis*-adduct.<sup>53</sup>

### 8-3. Kinetic Isotope Effect

The reaction progress was monitored by GC. Dodecane was added as an internal standard.

**Typical procedure for kinetic studies.** Pt(ICy)(dvtms) (3.6 mg, 5.9 µmol),  $B_2pin_2$  (76 mg, 0.30 mmol), dodecane (30 mg, internal standard) and benzene (1.0 mL) were added to a 10 mL sample vial with a Teflon-sealed screwcap. The cap was used to seal the vial under a flow of N<sub>2</sub>. The mixture was stirred at 100 °C. At specified times, the vial was rapidly cooled with running water and an aliquot of the reaction mixture was taken. The aliquot was diluted with EtOAc and analyzed by GC. After sampling, the reaction mixture was heated again until the next sampling. The amounts of PhBpin and B<sub>2</sub>pin<sub>2</sub> in the reaction mixture were determined over time; the results are shown below.



The total yield after 8 h was 99 %. These results show that reaction monitoring by cooling did not deactivate the catalytic species.

#### Measurement of KIE.



The reaction monitoring was also performed in benzene- $d_6$ . The initial rate after 20 min in benzene and benzene- $d_6$  were compared. Each experimental trial was performed in duplicate.



The average rate in benzene was  $8.3 \times 10^{-5}$  M/s, and that in benzene- $d_6$  was  $1.7 \times 10^{-5}$  M/s. The KIE for this reaction was 4.8.

### Typical procedure for kinetic studies for *N*-methylindole.

*N*-Methylindole-*d* **7a**-*d* was synthesized according to the literature procedure<sup>54</sup> and the deuterium content was determined to be 95% by <sup>1</sup>H NMR.



To an NMR tube with a Teflon-sealed screwcap, Pt(ICy)(dvtms) (1.2 mg, 2.0  $\mu$ mol), B<sub>2</sub>pin<sub>2</sub> (51 mg, 0.20 mmol), *N*-methylindole (13 mg, 0.10 mmol), 1,2,4,5-tetramethylbenzene (5.0 mg, as internal standard) and octane-*d*<sub>18</sub> (0.50 mL) were added. The cap was used to seal the vial under a flow of N<sub>2</sub>. The mixture was heated at 100 °C by heat air in a NMR spectrometer and the reaction progress was monitored by <sup>1</sup>H-NMR each 5 min. The amounts of borylated products **8a** in the reaction mixture were determined over time. The same experiment was performed with *N*-methylindole-*d* **7a**-*d* and each initial rates after 10 min were compared.



The rate with *N*-methylindole was  $1.8 \times 10^{-5}$  M/s, and that with *N*-methylindole-*d* was  $6.6 \times 10^{-6}$  M/s. The KIE for this reaction was 2.7.

#### 8-3. Reactivity of HBpin

When the reaction was conducted with HBpin instead of B2pin2, 1% PhBpin was formed.



Moreover, HBpin was detected by <sup>11</sup>B-NMR analysis of the crude reaction mixture of the Pt-catalyzed borylation of benzene-*d*<sub>6</sub> using B<sub>2</sub>pin<sub>2</sub>.



**Experimental procedure**. Pt(ICy)(dvtms) (1.2 mg, 0.0020 mmol), B<sub>2</sub>pin<sub>2</sub> (51 mg, 0.20 mmol) and benzene- $d_6$  (1.0 mL) were added to a 10 mL sample vial with a Teflon-sealed screwcap. The cap was used to seal the vial under a flow of N<sub>2</sub>. The vial was stirred at 100 °C over a hot plate for 20 h. The resulting reaction mixture was directly transferred to an NMR test tube and <sup>11</sup>B-NMR spectra were recorded. The <sup>11</sup>B-NMR spectra showed three peaks, at  $\delta = 31.2$  (PhBpin), 28.4 (DBpin), and 21.7 ppm (HOBpin),<sup>55</sup> which could be generated by the reaction of HBpin with residual H<sub>2</sub>O) (external reference: 15% BF<sub>3</sub>•EtO<sub>2</sub> in CDCl<sub>3</sub>). After the NMR analysis, GC analysis was conducted, and showed that 63 % PhBpin was produced and B<sub>2</sub>pin<sub>2</sub> was consumed completely.

The amount of unreacted HBpin was quantified by conducting hydroborylation<sup>56</sup> with styrene after the Pt-catalyzed borylation of benzene with B<sub>2</sub>pin<sub>2</sub>.



Pt(SICy)(dvtms) (4.0 mg, 0.0065 mmol, 2 mol%), B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol), dodecane (45 mg, as internal standard) and benzene (1.0 mL) were added to a 10 mL sample vial with a Teflon-sealed screwcap. The cap was used to seal the vial under a flow of N<sub>2</sub>. The mixture was stirred at 100 °C for 20 h. An aliquot of the reaction mixture was taken under a flow of N<sub>2</sub>. The aliquot was diluted with AcOEt and analyzed by GC (PhBpin 99%, full conversion of B<sub>2</sub>pin<sub>2</sub>). To the reaction mixture, [IrCl(cod)]<sub>2</sub> (4.0 mg, 0.0060 mmol, 2 mol%), dppe (6.2 mg, 0.015 mmol, 5 mol %), styrene (50  $\Box$ L, 0.44 mmol, 1.5 equiv) and CH<sub>2</sub>Cl<sub>2</sub> (1 mL) were added.

The vial was capped again and the reaction mixture was stirred for 24h at ambient temperature. The resulting mixture was filtered through silica gel (eluted with 7 mL of hexane/AcOEt = 5/1). The filtrate was analyzed by NMR using 1,1,2,2-tetrachloroethane as an internal standard, and showed that 4,4,5,5-tetramethyl-2-phenethyl-1,3,2-dioxaborolane was produced in 65% NMR yield, based on B<sub>2</sub>pin<sub>2</sub>. The resonances specific to the compounds are as follows:  $\delta$  = 2.75 ppm (t, *J* = 8.2 Hz, 2H, hydroborylated product H<sub>b</sub>); 7.81 (d, *J* = 8.0 Hz, 2H, PhBpin H<sub>a</sub>).



#### 8-4. Mercury Poisoning Test

The reaction was performed in the presence of excess mercury. The reaction was not inhibited by mercury, indicating that a homogeneous platinum species is involved as an active catalyst.



Pt(ICy)(dvtms) (5.5 mg, 0.90  $\mu$ mol) was dissolved in benzene (10 mL) to prepare 9.0 × 10<sup>-4</sup> M Pt(NHC)(dvtms) solution. B<sub>2</sub>pin<sub>2</sub> (76 mg, 0.30 mmol), mercury (42 mg, 0.21 mmol, 233 equiv to catalyst) and 1.0 mL of 9.0 × 10<sup>-4</sup> M Pt(ICy)(dvtms) benzene solution (0.9  $\mu$ mol, 0.3 mol%) were added to a 10 mL sample vial with a Teflon-sealed screwcap. The cap was used to seal the vial under a flow of N<sub>2</sub>. The vial was stirred at 100 °C for 20 h, and the resulting mixture was filtered through silica gel (eluted with 7 mL of hexane/AcOEt = 5/1). The filtrate was analyzed by GC using dodecane as an internal standard; 32% PhBpin was produced (TON = 106).

#### 2.5 References and Notes

- (1) J. A. Labinger, *Chem. Rev.* **2016**, in press (DOI: 10.1021/acs.chemrev.6b00583).
- (2) a) N. Tsukada, J. F. Hartwig, J. Am. Chem. Soc. 2005, 127, 5022. b) M. Murata, N. Fukuyama, J. Wada, S. Watanabe, Y. Masuda, Chem. Lett. 2007, 36, 910.
- (3) M. Yamamoto, S. Matsubara, *Chem. Lett.* **2007**, *36*, 172.
- (4) a) C. Jia, D. Piao, J. Oyadama, W. Lu, K. Tsugio, Y. Fujiwara, *Science*. 2000, 287, 1992. b) D. Karshtedt, A. T. Bell, T. D. Tilley, *Organometallics* 2004, 23, 4169. Hydroalylation via attack of arenes onto an alkyne π-activated by a platinum catalyst have also been reported: c) N. Chatani, H. Inoue, T. Ikeda, S. Murai, *J. Org. Chem.* 2000, 65, 4913.
- (5) A. M. Wagner, A. J. Hickman, M. S. Sanford, J. Am. Chem. Soc. 2013, 135, 15710.
- (6) Reviews: a) T. Ishiyama, N. Miyaura, *Chem. Rec.* 2004, *3*, 271. b) E. C. Neeve, S. J. Geier, I. A. I. Mkhalid, S. A. Westcott, T. B. Marder, *Chem. Rev.* 2016, *116*, 9091. c) J. Takaya, N. Iwasawa, *ACS Catal.* 2012, *2*, 1993. Pt/NHC-catalyzed diboration d) V. Lillo, J. Mata, J. Ramírez, E. Peris, E. Fernandez, *Organometallics* 2006, *25*, 5829. e) C. Pubill-Ulldemolins, C. Bo, J. A. Mata, E. Fernández, *Chem. An Asian J.* 2010, *5*, 261.
- (7) a) C. N. Iverson, M. R. Smith, J. Am. Chem. Soc. 1995, 117, 4403. b) T. Ishiyama, N. Matsuda, M. Murata, F. Ozawa, A. Suzuki, N. Miyaura, Organometallics 1996, 15, 713. c) G. Lesley, P. Nguyen, N. J. Taylor, T. B. Marder, A. J. Scott, W. Clegg, N. C. Norman, Organometallics 1996, 15, 5137 d) C. N. Iverson, M. R. Smith, Organometallics 1996, 15, 5155.
- (8) J. Takaya, S. Ito, H. Nomoto, N. Saito, N. Kirai, N. Iwasawa, Chem. Commun. 2015, 51, 17662.
- I. E. Markó, S. Stérin, O. Buisine, G. Mignani, P. Branlard, B. Tinant, J.-P. Declercq, *Science*. 2002, 298, 204.
- (10) Seminal works: a) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaura, N. R. Anastasi, J. F. Hartwig, J. Am. Chem. Soc. 2002, 124, 390. b) J.-Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka, M. R. Smith, Science. 2002, 295, 305.
- (11) G. Yan, Y. Jiang, C. Kuang, S. Wang, H. Liu, Y. Zhang, J. Wang, *Chem. Commun.* **2010**, *46*, 3170.
- (12) a) E. Eccles, K. Martensson, S. C. Chem. *Cur. Med. Res. Opin.* 2010, 26, 889. b) T. W. von Geldern, A. S. Tasker, B. K. Sorensen, M. Winn, B. G. Szczepankiewicz, D. B. Dixon, W. J. Chiou, L. Wang, J. L. Wessale, A. Adler, K. C. Marsh, B. Nguyen, T. J. Opgenorth, *J. Med. Chem.* 1999, 42, 3668. c) Y. L. Chen, R. S. Obach, J. Braselton, M. L. Corman, J. Forman, J. Freeman, R. J. Gallaschun, R. Mansbach, A. W. Schmidt, J. S. Sprouse, III. F. D. Tingley, E. Winston, D. W. Schulz, *J. Med. Chem.* 2008, *51*, 1385.
- M. Hoshino, H. Uekusa, A. Tomita, S. Koshihara, T. Sato, S. Nozawa, S. Adachi, K. Ohkubo, H. Kotani, S. Fukuzumi, *J. Am. Chem. Soc.* 2012, *134*, 4569.
- (14) M. N. Hopkinson, C. Richter, M. Schedler, F. Glorius, Nature 2014, 510, 485.
- (15) O. El-Sepelgy, S. Haseloff, S. K. Alamsetti, C. Schneider, Angew. Chem., Int. Ed. 2014, 53, 7923.
- (16) C. de Gracia Lux, S. Joshi-Barr, T. Nguyen, E. Mahmoud, E. Schopf, N. Fomina, A. Almutairi, J. Am. Chem. Soc. 2012, 134, 15758.
- (17) J. V. Obligacion, S. P. Semproni, I. Pappas, P. J. Chirik, J. Am. Chem. Soc. 2016, 138, 10645.

- (18) In general, functionalization or reduction of imidazolium backbone increases the buried volume compared to unsubstituted imidazolium carbenes: a) H. Clavier, A. Correa, L. Cavallo, E. C. Escudero-Adán, J. Benet-Buchholz, A. M. Z. Slawin, S. P. Nolan, *Eur. J. Inorg. Chem.* 2009, 2009, 1767. b) C. A. Urbina-Blanco, X. Bantreil, H. Clavier, A. M. Z. Slawin, S. P. Nolan, *Beilstein J. Org. Chem.* 2010, *6*, 1120. c) T. Dröge, F. Glorius, *Angew. Chem., Int. Ed.* 2010, *49*, 6940.
- (19) A. Perfetto, V. Bertolasi, C. Costabile, V. Paradiso, T. Caruso, P. Longo, F. Grisi, *RSC Adv.* 2016, *6*, 95793.
- (20) a) M. A. Esteruelas, M. Oliván, A. Vélez, *Organometallics* 2015, 34, 1911. b) S. I. Kalläne, M. Teltewskoi, T. Braun, B. Braun, *Organometallics* 2015, 34, 1156.
- (21) J. V. Obligacion, S. P. Semproni, P. J. Chirik, J. Am. Chem. Soc. 2014, 136, 4133.
- (22) a) M. Schlosser, E. Marzi, *Chem. A Eur. J.* 2005, *11*, 3449. b) K. Shen, Y. Fu, J. N. Li, L. Liu, Q. X. Guo, *Tetrahedron* 2007, *63*, 1568.
- (23) a) E. Clot, C. Mégret, O. Eisenstein, R. N. Perutz, J. Am. Chem. Soc. 2009, 131, 7817. b) M. E. Evans, C. L. Burke, S. Yaibuathes, E. Clot, O. Eisenstein, W. D. Jones, C. Gerhardt, J. Am. Chem. Soc 2009, 131, 13464.
- (24) a) J. Takagi, K. Sato, J. F. Hartwig, T. Ishiyama, N. Miyaura, *Tetrahedron Lett.* 2002, *43*, 5649.
  b) M. A. Larsen, J. F. Hartwig, *J. Am. Chem. Soc.* 2014, *136*, 4287.
- (25) a) G. Berthon-Gelloz, O. Buisine, J. F. Brière, G. Michaud, S. Stérin, G. Mignani, B. Tinant, J. P. Declercq, D. Chapon, I. E. Markó, *J. Organomet. Chem.* 2005, 690, 6156. b) B. G. Guillaume, B. De Bruin, B. Tinant, I. E. Markó, *Angew. Chem., Int. Ed.* 2009, 48, 3161.
- (26) a) X. Wan, X. Wang, Y. Luo, S. Takami, M. Kubo, A. Miyamoto, *Organometallics* 2002, 21, 3703. b) W. H. Lam, Z. Lin, *Organometallics* 2003, 22, 473. c) C. E. Webster, Y. Fan, M. B. Hall, D. Kunz, J. F. Hartwig, *J. Am. Chem. Soc.* 2003, 125, 858. d) J. F. Hartwig, K. S. Cook, M. Hapke, C. D. Incarvito, Y. Fan, C. E. Webster, M. B. Hall, *J. Am. Chem. Soc.* 2005, 127, 2538.
- (27) a) H. Tamura, H. Yamazaki, H. Sato, S. Sakaki, J. Am. Chem. Soc. 2003, 125, 16114. b) T. M. Boller, J. M. Murphy, M. Hapke, T. Ishiyama, N. Miyaura, J. F. Hartwig, J. Am. Chem. Soc. 2005, 127, 14263.
- (28) Studies on reductive elimination processes from Pt-boryl species: a) H. Braunschweig, R. Bertermann, P. Brenner, M. Burzler, R. D. Dewhurst, K. Radacki, F. Seeler, *Chem. A Eur. J.* 2011, *17*, 11828. b) H. Braunschweig, P. Brenner, R. D. Dewhurst, F. Guethlein, J. O. C. Jimenez-Halla, K. Radacki, J. Wolf, L. Zöllner, *Chem. A Eur. J.* 2012, *18*, 8605. c) H. Braunschweig, A. Damme, *Chem. Commun.* 2013, *49*, 5216.
- (29) S. Voth, J. W. Hollett, J. A. McCubbin, J. Org. Chem. 2015, 80, 2545.
- (30) C. Cheng, J. F. Hartwig, Science 2014, 343, 853.
- (31) K. Lee, D. Katsoulis, J. Choi, ACS Catal. 2016, 6, 1493.
- (32) M. A. Larsen, C. V. Wilson, J. F. Hartwig, J. Am. Chem. Soc. 2015, 137, 8633.
- (33) T. Ishiyama, H. Isou, T. Kikuchi, N. Miyaura, Chem. Commun. 2010, 46, 159.
- (34) a) H. Huo, C. Fu, K. Harms, E. Meggers, J. Am. Chem. Soc. 2014, 136, 2990. b) T. Guney, J. J. Lee, G. A. Kraus, Org. Lett. 2014, 16, 1124.

- (35) S. Fujii, Y, Miyajima, H. Masuno, H. Kagechika, J. Med. Chem. 2013, 56, 160.
- (36) Y. Yamanoi, J. Org. Chem. 2005, 70, 9607.
- (37) L. Wu, E. Drinkel, F. Gaggia, S. Capolicchio, A. Linden, L. Falivene, L. Cavallo, R. Dorta, *Chem. Eur. J.* **2011**, *17*, 12886.
- (38) T. Liu, X. Zhao, Q. Shen, L. Lu, Tetrahedron 2012, 68, 6535.
- (39) T. Maibunkaew, C. Thongsornkleeb, J. Tummatorn, A. Bunrit, S. Ruchirawat, *Synlett* **2014**, *25*, 1769.
- (40) B. D. Karstedt, N. Y. Scotia, US patent 1970, US3715334 A. 6156.
- (41) C. J. Serpell, J. Cookson, A. L. Thompson, C. M. Brown, P. D. Beer, *Dalton Trans.* 2013, 42, 1385.
- (42) N. Gürbüz, I. Özdemir, S. Demir, B. Çetinkaya, J. Mol. Cat. A Chem. 2004, 209, 23.
- (43) H. Richter, H. Schwertfeger, P. R. Schreiner, R. Fröhlich, F. Glorius, Synlett 2009, 2, 193.
- (44) J. Liu, Y. Li, Y. Li, N. Hu, J. Appl. Polym. Sci. 2008, 109, 700.
- (45) B. R. Van Ausdall, J. L. Glass, K. M. Wiggins, A. M. Aarif, J. Louie, J. Org. Chem. 2009, 74, 7935.
- (46) J. Jiang, H. Zhu, Y. Shen, T. Tu, Org. Chem. Front. 2014, 1, 1172.
- (47) a) T. A. Boebel, J. F. Hartwig, *Organometallics* 2008, 27, 6013. b) S. Kawamorita, H. Ohmiya, K. Hara, A. Fukuoka, M. Sawamura, *J. Am. Chem. Soc.* 2009, *131*, 5058.
- (48) S. Kawamorita, H. Ohmiya, T. Iwai, M. Sawamura, Angew. Chem. Int. Ed. 2011, 50, 8363.
- (49) L. D. Marciasini, N. Richy, M. Vaultier, M. Pucheault, Adv. Synth. Catal. 2013, 355, 1083.
- (50) G. A. Chotana, M. A. Rak, M. R. Smith III, J. Am. Chem. Soc. 2005, 127, 10539.
- (51) Y. Maegawa, S. Inagaki, Dalton Trans. 2015, 44, 13007.
- (52) 3-Isomer: H. Nakagawa, S. Kawai, T. Nakashima, T. Kawai, *Org. Lett.* 2009, *11*, 1475. 2-Isomer:
  A. J. Sandee, C. K. Williams, N. R. Evans, J. E. Davies, C. E. Boothby, R. H. Friend, A. B. Holmes, *J. Am. Chem. Soc.* 2004, *126*, 7041.
- (53) N. Nakagawa, T. Hatakeyama, M. Nakamura, Chem. Eur. J. 2015, 21, 4257.
- (54) M. V. Leskinen, A. Madarász, K.-T. Yip, A. Vuorinen, I. Pápai, A. J. Neuvonen, P. M. Pihko, J. Am. Chem. Soc. 2014, 136, 6453.
- (55) H. F. Bettinger, M. Fithaus, H. Bornemann, I. M. Oppel, Angew. Chem. Int. Ed. 2008, 47, 4744.
- (56) Y. Yamamoto, R. Fujikawa, T. Umemoto, N. Miyaura, Tetrahedron 2004, 60, 10695.

#### Conclusion

The research reported in this thesis was directed at the nickel- and platinum-catalyzed borylation of aromatic C-H bonds. The present study revealed that nickel and platinum complexes have promising catalytic activities for the C-H borylation of arenes. Their reactivity and selectivity were investigated for the purpose of comparison with the well-established iridium systems.

In chapter 1, the nickel-catalyzed borylation reaction of aromatic C-H bonds is described. The use of an NHC ligand is essential for an efficient reaction and a variety of indole derivatives could be selectively borylated at the 2-position. The protocol can be applied to the gram-scale borylation of indoles using the low cost, readily available Ni(OAc)<sub>2</sub>. This study demonstrated that an earth abundant base metal can mediate cutting-edge C-H functionalization reactions and will help to address recent concerns regarding the limited availability of precious metals.

In chapter 2, the platinum-catalyzed borylation of aromatic C-H bonds is described. A platinum/ICy complex effectively borylated C-H bonds in aromatic substrates. One unique feature of the platinum systems is their robustness to steric hindrance, enabling the borylation of C-H bonds that are located at sterically congested positions. Further ligand screening determined that the SIPr ligand as an optimal ligand for the borylation of sterically hindered arenes under practical conditions without the need for a large excess of the substrates. Another feature of the platinum catalyst is that a fluorine functionality exhibited an *ortho*-directing effect, providing facile access to a wide range of *ortho*-fluorinated phenylboronic ester from fluoroarenes. These unique selectivities are not attainable when the established iridium catalyst system is used. Some synthetic applications that were made possible by the characteristic features of the current platinum system were demonstrated in this study.

Catalytic borylation of C-H bonds is now undoubtedly recognized as one of the most powerful methods for introducing a synthetically versatile boron functionality, therefore the development of catalysts with new reactivities and/or selectivities are highly desired for further progress in C-H borylation and the next breakthrough in undirected C-H functionalization. The knowledge and findings obtained through this study indicates the possibility of nickel and platinum as promising candidates to be investigated for use as C-H borylation catalysts in the future and will contribute to the advancement of C-H functionalization chemistry.

### **List of Publications**

- Nickel-Catalyzed Borylation of Arenes and Indoles via C-H Bond Cleavage Takayuki Furukawa, Mamoru Tobisu, Naoto Chatani *Chem. Commun.* 2015, *51*, 6508.
- (2) C-H Functionalization at Sterically Congested Positions by the Platinum-Catalyzed Borylation of Arenes

Takayuki Furukawa, Mamoru Tobisu, Naoto Chatani J. Am. Chem. Soc. **2015**, 137, 12211.

C-H Borylation by Platinum Catalysis
 Takayuki Furukawa, Mamoru Tobisu, Naoto Chatani
 *Bull. Chem. Soc. Jpn.* 2017, *in press.*
## **Supplementary List of Publications**

- Modular Synthesis of Phenanthridine Derivatives by Oxidative Cyclization of 2-Isocyanobiphenyls with Organoboron Reagents Mamoru Tobisu, Keika Koh, Takayuki Furukawa, Naoto Chatani Angew. Chem. Int. Ed. 2012, 51, 11363.
- Visible Light-Mediated Direct Arylation of Arenes and Heteroarenes Using Diaryliodonium Salts in the Presence and Absence of a Photocatalyst Mamoru Tobisu, Takayuki Furukawa, and Naoto Chatani *Chem. Lett.* 2013, 42, 1203.
- Nickel-Catalyzed Borylation of Aryl and Benzyl 2-Pyridyl Ethers: A Method for Converting a Robust *ortho*-Directing Group
  Mamoru Tobisu, Jiangning Zhao, Hirotaka Kinuta, Takayuki Furukawa, Takuya Igarashi and Naoto Chatani
  *Adv. Synth. Catal.* 2016, 358, 2417.
- (4) Nickel/N-Heterocyclic Carbene-Catalyzed Suzuki-Miyaura Type Cross-Coupling of Aryl Carbamates

Akimichi Ohtsuki, Kousuke Yanagisawa, Takayuki Furukawa, Mamoru Tobisu and Naoto Chatani *J. Org. Chem.* **2016**, *81*, 9409.

 (5) A Conia-Ene-Type Cyclization Under Basic Conditions Enables an Efficient Synthesis of (-)-Lycoposerramine R
 E Lie Lie Lie Lie Lie Did T

Felix Hartrampf, Takayuki Furukawa, Dirk Trauner

Angew. Chem. Int. Ed. 2017, 56, 893.

(6) Nickel-Mediated Decarbonylation of Simple Unstrained Ketones through the Cleavage of Carbon–Carbon Bonds

Toshifumi Morioka, Akihiro Nishizawa, Takayuki Furukawa, Mamoru Tobisu and Naoto Chatani *J. Am. Chem. Soc.* **2017**, *in press*.