

Title	Notes on trivial source modules
Author(s)	Tsushima, Yukio
Citation	Osaka Journal of Mathematics. 1995, 32(2), p. 475-482
Version Type	VoR
URL	<a href="https://doi.org/10.18910/6298">https://doi.org/10.18910/6298</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## NOTES ON TRIVIAL SOURCE MODULES

Dedicated to Professor S. Endo on his 60th birthday

YUKIO TSUSHIMA

(Received July 8, 1993)

### 1. Introduction

Let  $G$  be a finite group and  $k$  an algebraically closed field of characteristic  $p$ . An indecomposable  $kG$ -module with a vertex  $Q$  is said to be a *weight module* if its Green correspondent with respect to  $(G, Q, N_G(Q))$  is simple. Let  $B$  be a block of  $kG$ . Alperin [1] conjectured that the number of the weight modules belonging to  $B$  equals that of the simple modules in  $B$ . If this is the case and a defect group of  $B$  a TI set, then it can be shown under some additional assumption that the socles of weight modules are simple, which in turn determine the isomorphism classes of the weight modules; this holds if  $G$  is a simple group with a cyclic Sylow  $p$ -subgroup. This rather surprising property has been known to hold for finite groups of Lie type of characteristic  $p$ . However little is known about general properties of weight modules. In the final section we shall study solvable groups that have only simple weight modules.

Throughout this paper  $G$  denotes a finite group and  $k$  an algebraically closed field of prime characteristic  $p$ . For a  $kG$ -module  $M$ ,  $\text{hd}(M)$ ,  $\text{soc}(M)$  and  $P(M)$  denote the head, socle and projective cover of  $M$  respectively. If  $N$  is a  $kG$ -module,  $N|M$  indicates that  $N$  is isomorphic to a direct summand of  $M$ , and  $(N, M)$  denotes the multiplicity of  $N$  as a summand of  $M$ . We fix a block  $B$  of  $kG$  and let  $D$  be its defect group.  $\text{IRR}(B)$  denotes a full set of non-isomorphic simple modules in  $B$ ,  $l(B)$  its cardinality and  $\text{WM}(B)$  a full set of non-isomorphic weight modules belonging to  $B$ . Let  $f$  be the Green correspondence with respect  $(G, D, H)$ , where  $H = N_G(D)$ . If  $\text{WM}(B|D)$  denotes the subset of  $\text{WM}(B)$  consisting of the weight modules with vertices  $D$  and  $b$  the Brauer correspondent of  $B$  in  $kH$ , then  $f$  induces a bijection between  $\text{WM}(B|D)$  and  $\text{IRR}(b)$ .

The author thanks the referee for improving the proof of Proposition 4 below.

### 2. Weight modules over blocks with TI defect groups

To begin with, we quote the following as a preliminary lemma.

**Lemma 1** (Robinson [8]). *Let  $T$  be a subgroup of  $G$ . Let  $M$  (resp.  $N$ ) be*

a simple  $kG$  (resp.  $kT$ )-module. Then we have  $(P(M), N^G) = (P(N), M_{|H})$ .

Throughout this section  $D$  is assumed to be a non-trivial TI subgroup of  $G$ , i.e.,  $D \cap D^x = 1$  if  $x \in G \setminus H$ . Let  $\text{IRR}(B) = \{M_1, \dots, M_r\}$ ,  $\text{IRR}(b) = \{W_1, \dots, W_e\}$  and  $n_i = \dim_k W_i$ . We set  $\text{WM}(B|D) = \{V_i = f^{-1}(W_i); 1 \leq i \leq e\}$ . Note that  $\text{WM}(B) = \text{WM}(B|D)$ . In fact, let  $V \in \text{WM}(B)$  and  $Q = vx(V)$ . We may assume that  $D \supset Q$ . If  $D > Q$ , then  $N_D(Q) > Q$ . On the other hand, since  $D$  is a TI set, it follows that  $H \supset N_G(Q)$  and hence  $N_D(Q)$  is normal in  $N_G(Q)$ . So,  $N_G(Q)/Q$  fails to have a block of defect zero. This is a contradiction, since  $f(V)$  is simple and projective as an  $N_G(Q)/Q$ -module.

**Lemma 2.**  $M_{i|H} = f(M_i) \oplus N_i$ , where  $N_{i|D}$  is projective and  $f(M_i)_{|D}$  has no projective summand.

Proof. If  $L$  is an indecomposable component of  $N_i$  with vertex  $P$ , then  $P$  lies in  $\mathfrak{Y}(D, H)$ , where

$$\mathfrak{Y}(D, H) = \{Q; Q \subset D^x \cap H, x \in G \setminus H\}.$$

By the Mackey decomposition theorem we have

$$(L \otimes_P kH)_{|D} = \bigoplus_{y \in P \setminus H/D} (L \otimes_P y) \otimes_{P^y \cap D} kD.$$

There is  $x \in G \setminus H$  such that  $P \subset D^x \cap H$ . Hence for any  $y \in H$ , we have

$$P^y \cap D \subset D^{xy} \cap D \cap H = 1, \text{ as } xy \in G \setminus H.$$

Therefore  $(L \otimes_P kH)_{|D}$  is projective. Since  $L|L \otimes_P kH$ ,  $L_{|D}$  is also projective.

We next show that  $f(M_i)_{|D}$  is projective-free. Actually, this is a general fact. Note that  $f(M_i)$  belongs to  $b$  and  $b$  has the normal defect group  $D$ . So, it suffices to show that if  $L$  is a non-projective indecomposable  $b$ -module, then  $L_{|D}$  is projective-free. But since  $L$  is  $D$ -projective, this is a routine work, using Mackey decomposition.

**Lemma 3.**  $\text{Hom}_{kG}(M_i, V_j) \simeq \text{Hom}_{kH}(f(M_i), W_j)$  for all  $i, j$ .

Proof. There is an isomorphism

$$\text{Hom}_{kG}(M_i, V_j) / \text{Tr}_{\mathfrak{X}}^G(M_i, V_j) \simeq \text{Hom}_{kH}(f(M_i), W_j) / \text{Tr}_{\mathfrak{X}}^H(f(M_i), W_j),$$

where  $\mathfrak{X} = \mathfrak{X}(D, H) = \{Q; Q \subset D^x \cap D, x \in G \setminus H\}$ . However, since  $D$  is a TI set, we have  $\mathfrak{X} = \{1\}$ . And if  $M$  and  $V$  are non-projective indecomposable and if one of them is simple, then  $\text{Tr}_1^G(M, V) = 0$ , whence the result follows.

**Proposition 4.** *Let  $\varepsilon$  be the block idempotent of  $B$ . Then we have*

$$(k_D)^G \varepsilon \simeq \bigoplus_{i=1}^e n_i V_i \oplus \bigoplus_{i=1}^r a_i P(M_i), \text{ with } a_i = (kD, M_{iD}).$$

Proof. Let

$$k[H/D] = \sum_{i=1}^m n_i W_i$$

be an indecomposable decomposition. Note that no  $W_j$  belongs to  $b$  if  $j \geq e+1$ . Since  $D$  is a TI set, we have

$$W_i^G = f^{-1}(W_i) \oplus (\text{projectives}).$$

Moreover we know by Green's theorem that  $V_j = f^{-1}(W_j)$  does not belong to  $B$  if  $j \geq e+1$ . Thus

$$(k_D)^G = (k_B^H)^G = k[H/D]^G = \bigoplus_{i=1}^e n_i V_i \oplus \bigoplus_{j=e+1}^m n_j V_j \oplus (\text{projectives}),$$

whence we have

$$(k_D)^G \varepsilon = \bigoplus_{i=1}^e n_i V_i \oplus \bigoplus_{i=1}^r a_i P(M_i), \text{ with } a_i \geq 0$$

and by Lemma 1,  $a_i = (kD, M_{iD})$  for  $i = 1, 2, \dots, r$ .

**Theorem 5.** *Assume that  $D$  is a TI set and that  $\text{hd}(f(M_i))$  is simple for all  $i$ . Then we have the following:*

- (1)  $l(B) \geq l(b)$ ;
- (2) *the equality sign in the above holds if and only if  $\text{soc}(V_i)$  is simple for all  $i$  ( $1 \leq i \leq e$ ), in which case we have that*

$$\text{soc}(V_i) \simeq \text{soc}(V_j) \text{ if and only if } V_i \simeq V_j.$$

Proof. From the assumption we may set  $\text{hd}(f(M_i)) = W_{\tau(i)}$  ( $1 \leq i \leq r$ ,  $1 \leq \tau(i) \leq e$ ). By lemma 3 we find easily that

- (i)  $M_i | \text{soc}(V_{\tau(i)})$  with multiplicity one.
- (ii) If  $M_i | \text{soc}(V_j)$ , then  $j = \tau(i)$ .

Now, the second assertion yields that the map

$$\tau: \{1, 2, \dots, r\} \rightarrow \{1, 2, \dots, e\}$$

is a surjection. In fact, for an arbitrary  $V_j$ , take  $M_i$  such that  $M_i | \text{soc}(V_j)$ . Then  $j = \tau(i)$ . Thus  $\tau$  is surjective. In particular, we have that  $r \geq e$ .

To show the second part of the theorem, suppose that  $\text{soc}(V_j)$  is simple for all  $j$ . Then  $M_i = \text{soc}(V_{\tau(i)})$  and hence  $\tau$  is a bijection. Therefore we have  $r = e$ . If, conversely,  $r = e$ , then  $\tau$  is a bijection. This implies by (ii) above that  $\text{soc}(V_i)$  must be simple and  $V_i \simeq V_j$  if and only if  $\text{soc}(V_i) \simeq \text{soc}(V_j)$ .

REMARK 1. If the Alperin conjecture is true, we always have  $l(B) = l(b)$  when  $D$  is a TI set.

### 3. Weight modules for the symmetric group $S_p$

In this section we assume that  $G = S_p$  is the symmetric group on  $p$  letters. If  $D$  is a Sylow  $p$ -subgroup of  $G$ , then  $D$  has order  $p$  and  $C_G(D) = D, H/D \simeq (Z/(p))^*$ , the group of units of  $Z/(p)$ . In particular, it follows that  $b = kH$  is the block of  $kH$ . Let us write

$$\text{IRR}(b) = \{W_0, \dots, W_{p-2}\}, \text{ where } \dim_k W_i = 1 \ (0 \leq i \leq p-2).$$

If  $B$  denotes the principal block of  $G$ , then  $B$  is a unique block of  $kG$  of positive defect and  $l(B) = p-1$ . The decomposition matrix of  $B$  is known. It can be displayed as follows, see James [5].

	$\varphi_0$	$\varphi_1$	$\varphi_2$	$\dots$	$\varphi_{p-2}$
$\chi_0 = (p) = 1_G$	1				
$\chi_1 = (p-1, 1)$	1	1			<b>0</b>
$\chi_2 = (p-2, 1^2)$		1	1		
$\vdots$			$\ddots$	$\ddots$	
$\chi_{p-2} = (2, 1^{p-2})$	<b>0</b>			1	1
$\chi_{p-1} = (1^p)$					1

Since  $\chi_i = \varphi_{i-1} + \varphi_i$  and  $\deg \chi_i = {}_{p-1}C_i$ , we find via induction that  $\deg \varphi_i = {}_{p-2}C_i$  ( $0 \leq i \leq p-2$ ). So we can label the simple modules in  $B$  such that

$$\text{IRR}(B) = \{M_0, \dots, M_{p-2}\}, \text{ with } m_i = \dim_k M_i = {}_{p-2}C_i.$$

Here we note the following facts on binomial coefficients  ${}_n C_i$ .

**Lemma 6.**

$$(1) \quad m_i = {}_{p-2}C_i = \begin{cases} i + 1 \pmod p, & \text{if } i \text{ is even;} \\ p - i - 1 \pmod p, & \text{if } i \text{ is odd.} \end{cases}$$

(2) Suppose that  $n \geq 4$ . If  $2 \leq i \leq n-2$ , then  ${}_n C_i \geq n+2$ .

Now, since  $H/D$  is abelian, every principal indecomposable module over  $kH$  has dimension  $p$ , and thus every non-projective indecomposable module has dimension smaller than  $p$ . In particular it follows that  $\dim_k f(M_i) < p$ . By Lemma 2, we can write

$$M_{i|D} = f(M_i)_{|D} \oplus a_i kD.$$

For  $i=0, 1, p-3$  or  $p-2$ , we have that  $m_i < p$  and so  $a_i = 0$ . This is true for all  $i$ , provided  $p \leq 5$ . Suppose  $p > 5$ . If  $2 \leq i \leq p-4$ , then  $m_i \geq p$  by Lemma 6(2) and hence  $a_i > 0$ . This, together with Lemma 6(1) yields that  $\dim_k f(M_i) = i+1$  or  $p-i-1$  according as whether  $i$  is even or odd ( $2 \leq i \leq p-4$ ). Thus we have:

$$a_i = \begin{cases} (m_i - i - 1)/p, & \text{if } i \text{ is even;} \\ (m_i - (p - i - 1))/p, & \text{if } i \text{ is odd.} \end{cases}$$

Now we have the following result by Lemma 1 and Proposition 4.

**Proposition 7.** Let  $WM(B) = \{V_0, \dots, V_{p-2}\}$ , where  $V_i = f^{-1}(W_i)$ , and let  $\{U_j; 1 \leq j \leq q\}$  be the set of simple  $kG$ -modules belonging to the blocks of defect zero. Then we have

$$(k_D)^G \simeq \bigoplus_{i=0}^{p-2} V_i \oplus \bigoplus_{i=2}^{p-4} a_i P(M_i) \oplus \bigoplus_{i=1}^q (\dim_k U_i / p) U_i.$$

**4. Socles of weight modules**

In view of Theorem 5, it seems to be natural to consider the following situation:

(#) Every weight module belonging to  $B$  has a simple socle, and for  $U, V \in WM(B)$ , we have

$$\text{soc}(U) \simeq \text{soc}(V) \text{ if and only if } U \simeq V.$$

We first remark that

**Proposition 8.** If  $G$  is a simple group with a cyclic Sylow  $p$ -subgroup, the condition (#) holds for every block  $B$ .

In fact we know that a Sylow  $p$ -subgroup is a TI set (Blau[2]) and that  $l(B) = l(b)$ , hence the result follows from Theorem 5.

On the other hand, we have the following, as is shown on pp.370–371 in

Alperin [1].

**Proposition 9** (Alperin). *Let  $G$  be a finite group of Lie type of characteristic  $p$ . Then the condition (#) holds for every block  $B$ .*

Before proceeding let us recall that a simple module is a weight module if and only if it has trivial source (Okuyama [7]).

Now, for the rest of this paper we assume that  $G$  is solvable. In this case the Alperin conjecture has been proved by Okuyama.

**DEFINITION.** A solvable group  $G$  is said to be  $p'$ -supersolvable if all of its chief composition factors of order prime to  $p$  are cyclic.

**Proposition 10.** *If  $G$  is  $p'$ -supersolvable, every simple module has trivial source. Hence  $WM(B) = IRR(B)$  for every block  $B$ .*

**Proof.** Let  $G$  be a counter-example of minimum order and let  $V$  be a simple  $kG$ -module with source not isomorphic to  $k$ . Let  $K$  be a maximal abelian normal  $p'$ -subgroup of  $G$  and  $W$  a simple summand of  $V_{|K}$ . By Fong's reduction and the minimality of  $G$ ,  $W$  must be  $G$ -invariant. So  $W$  is faithful as  $K$ -module and hence  $K$  must be central. If  $O_p(G/K) = 1$ ,  $G/K$  has a cyclic normal  $p'$ -subgroup, say  $M/K$ . Then  $M$  is abelian, contradicting the choice of  $K$ . Thus  $O_p(G/K) > 1$ , which implies that  $O_p(G) > 1$ , since  $K$  is central. This is a contradiction. The second statement is clear since the number of weight modules belonging to  $B$  equals  $l(B)$ .

Now we give a definition:

**DEFINITION.** A finite group is said to be a CR1-group if all of its characteristic abelian subgroups are cyclic.

We say that the group  $G$  involves a group  $T$  provided there are subgroups  $L \triangleright M$  of  $G$  such that  $L/M \simeq T$ . For a prime number  $q$ , let us denote a Sylow  $q$ -subgroup of  $G$  by  $G_q$ .

**Theorem 11.** *Let  $G$  be a solvable group and suppose that  $G_q$  involves no non-abelian CR1-group for each prime  $q$  different from  $p$ . Then the conclusion of Proposition 10 holds.*

**Proof.** We shall show that every simple  $kG$ -module has trivial source by the induction on the order of  $G$ . We may assume  $O_p(G) = 1$ . Let  $K$  be the Fitting subgroup of  $G$ , so we have that  $C_G(K) \subset K$ . If  $K$  is cyclic,  $\text{Aut}(K)$  is abelian and

so is  $G/K$ . Thus  $G$  is supersolvable and the result follows from Proposition 10. If  $K$  is non-cyclic, our assumption implies that  $G$  has a non-cyclic abelian normal  $q$ -subgroup, say  $L$ , for some prime  $q$ . Let  $V$  be a simple  $kG$ -module and  $W$  a simple summand of  $V|_L$ . If the inertial group of  $W$  is proper, the result follows by induction. If  $W$  is  $G$ -invariant,  $N = \text{Ker}(W)$  is a non-trivial normal subgroup of  $G$ . Then we get the result by applying the inductive hypothesis to  $G/N$ .

REMARK 2. The CR1- $q$ -groups are classified (Gorenstein [3], Chap. 5). In particular, a non-abelian CR1- $q$ -group contains  $D_3$  or  $Q_3$  if  $q=2$ , while it contains  $M(q)$  if  $q$  is odd, where

$$M(q) = \langle x, y, z; x^q = y^q = z^q = 1, [x, z] = [y, z] = 1, [x, y] = z \rangle,$$

which has order  $q^3$  and exponent  $q$ .

REMARK 3. One may show that the following  $q$ -group  $Q$  involves no non-abelian CR1- $q$ -group:

$$Q = \langle x, y; x^{q^a} = y^{q^b} = 1, x^y = x^{1+q^{a-1}} \rangle,$$

where  $a \geq 2$ ,  $b \geq 1$ , and  $a \geq 3$  if  $q=2$ .

In fact every proper subgroup of  $Q$  is abelian (cf. Huppert [4] III, Aufgaben 22). So it suffices to show that  $Q$  has no factor group isomorphic to  $D_3$ ,  $Q_3$  or  $M(q)$ , which will be easily done.

REMARK 4. Let  $G = \langle \sigma \rangle$  be the semidirect product, where  $\sigma$  is an automorphism of the quaternion group  $Q_3$  of order 3. Then  $kG$  has a simple module whose source is not trivial, where  $k$  is of characteristic 3. On the other hand, if  $G$  is the symmetric group  $S_4$ , every simple  $kG$ -module has trivial source,  $k$  being the same as above. In both groups the Sylow 2-subgroups are CR1-groups.

---

#### References

- [1] J.L. Alperin: *Weights for finite groups*, Proc. Symposia in Pure Math. **47** (1987), 369–379, A.M.S.
- [2] H.I. Blau: *On trivial intersection of cyclic Sylow subgroups*, Proc. Amer. Math. Soc. **94** (1985), 572–576.
- [3] D. Gorenstein: *Finite Groups*, Harper & Row, New York, 1968.
- [4] B. Huppert: *Endliche Gruppen I*, Springer-Verlag, Berlin, 1967.
- [5] G.D. James: *The Representation Theory of the Symmetric Groups*, Lect. notes in Math. **682** (1978), Springer-Verlag.
- [6] H. Nagao and Y. Tsushima: *Representations of Finite Groups*, Academic Press, Boston, 1989.
- [7] T. Okuyama: *Module correspondence in finite groups*, Hokkaido Math. J. **10** (1981), 299–318.
- [8] G.R. Robinson: *On projective summands of induced modules*, J. Alg. **122** (1989), 106–111.



Department of Mathematics  
Osaka City University  
558 Osaka, Japan