



Title	LIE ALGEBRAS CONSTRUCTED WITH LIE MODULES AND THEIR POSITIVELY AND NEGATIVELY GRADED MODULES
Author(s)	Sasano, Nagatoshi
Citation	Osaka Journal of Mathematics. 2017, 54(3), p. 533-568
Version Type	VoR
URL	https://doi.org/10.18910/66999
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

LIE ALGEBRAS CONSTRUCTED WITH LIE MODULES AND THEIR POSITIVELY AND NEGATIVELY GRADED MODULES

NAGATOSHI SASANO

(Received December 10, 2015, revised July 29, 2016)

Abstract

In this paper, we shall give a way to construct a graded Lie algebra $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ from a standard pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ which consists of a Lie algebra \mathfrak{g} which has a non-degenerate invariant bilinear form B_0 and \mathfrak{g} -modules (ρ, V) and $\mathcal{V} \subset \text{Hom}(V, F)$ all defined over a field F with characteristic 0. In general, we do not assume that these objects are finite-dimensional. We can embed the objects $\mathfrak{g}, \rho, V, \mathcal{V}$ into $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. Moreover, we construct specific positively and negatively graded modules of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. Finally, we give a chain rule on the embedding rules of standard pentads.

1. Introduction

A standard quadruplet is a quadruplet of the form $(\mathfrak{g}, \rho, V, B_0)$, where \mathfrak{g} is a finite-dimensional reductive Lie algebra, (ρ, V) a finite-dimensional representation of \mathfrak{g} and B_0 a non-degenerate symmetric invariant bilinear form on \mathfrak{g} all defined over the complex number field \mathbb{C} , which satisfies the conditions that ρ is faithful and completely reducible and that V does not have a non-zero invariant element. In [8], the author proved that any standard quadruplet $(\mathfrak{g}, \rho, V, B_0)$ has a graded Lie algebra, denoted by $L(\mathfrak{g}, \rho, V, B_0) = \bigoplus_{n \in \mathbb{Z}} V_n$, such that $V_0 \cong \mathfrak{g}$, $V_1 \cong V$ and $V_{-1} \cong \text{Hom}(V, \mathbb{C})$ (see [8, Theorem 2.11]). That is, any finite-dimensional reductive Lie algebra and its finite-dimensional faithful and completely reducible representation can be embedded into some (finite or infinite-dimensional) graded Lie algebra. We call a graded Lie algebra of the form $L(\mathfrak{g}, \rho, V, B_0)$ the Lie algebra associated with a standard quadruplet. Some well-known Lie algebras correspond to some standard quadruplet, for example, finite-dimensional semisimple Lie algebras and loop algebras. Moreover, the bilinear form B_0 can be also embedded into $L(\mathfrak{g}, \rho, V, B_0)$, i.e. there exists a non-degenerate symmetric invariant bilinear form on $L(\mathfrak{g}, \rho, V, B_0)$ whose restriction to $V_0 \times V_0$ coincides with B_0 (see [8, Proposition 3.2]). By the way, H. Rubenthaler obtained some similar results in [7] using the Kac theory in [2].

The first purpose of this paper is to extend the theory of standard quadruplets to the cases where the objects are infinite-dimensional. For this, we need to consider pentads $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ instead of quadruplets, where \mathfrak{g} is a finite or infinite-dimensional Lie algebra, $\rho : \mathfrak{g} \otimes V \rightarrow V$ a representation of \mathfrak{g} on a finite or infinite-dimensional vector space V , \mathcal{V} a \mathfrak{g} -submodule of $\text{Hom}(V, F)$, B_0 a non-degenerate invariant bilinear form on \mathfrak{g} all defined over a field F with characteristic 0. In general, we do not assume that B_0 is symmetric. We define the notion of *standard pentads* by the existence of a linear map

$\Phi_\rho : V \otimes \mathcal{V} \rightarrow \mathfrak{g}$ satisfying $B_0(a, \Phi_\rho(v \otimes \phi)) = \langle \rho(a \otimes v), \phi \rangle$ for any $a \in \mathfrak{g}$, $v \in V$ and $\phi \in \mathcal{V}$. A standard quadruplet $(\mathfrak{g}, \rho, V, B_0)$ can be naturally regarded as a standard pentad $(\mathfrak{g}, \rho, V, \text{Hom}(V, \mathbb{C}), B_0)$, and, thus, we can say that the notion of standard pentads is an extension of the notion of standard quadruplets. Then, by a similar argument to the argument in [8], we can construct a graded Lie algebra from an arbitrary standard pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ denoted by $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) = \bigoplus_{n \in \mathbb{Z}} V_n$ such that the objects $\mathfrak{g}, \rho, V, \mathcal{V}$ can be embedded into it. We call such a graded Lie algebra a *Lie algebra associated with a standard pentad*. This is the first main result of this paper. Of course, the graded Lie algebra associated with a standard quadruplet $(\mathfrak{g}, \rho, V, B_0)$ is isomorphic to the graded Lie algebra associated with a standard pentad $(\mathfrak{g}, \rho, V, \text{Hom}(V, \mathbb{C}), B_0)$. Moreover, if the bilinear form B_0 of $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is symmetric, then B_0 can be also embedded into $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, i.e. there exists a non-degenerate symmetric invariant bilinear form B_L on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ whose restriction to $V_0 \times V_0$ coincides with B_0 .

When B_0 is symmetric, we can expect that a Lie algebra of the form $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ (not necessary finite-dimensional) and its representation can be embedded into some graded Lie algebra using B_L . The second purpose is to construct positively graded modules and negatively graded modules of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ which can be embedded into some graded Lie algebra under some assumptions. In general, it is known that for any graded Lie algebra $\mathfrak{l} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{l}_n$ and \mathfrak{l}_0 -module U , there exists a positively (respectively negatively) graded \mathfrak{l} -module such that the base space (respectively top space) is the given \mathfrak{l}_0 -module U (see [9, Theorem 1.2]). In this paper, we shall try to construct such $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -modules from a \mathfrak{g} -module (π, U) using a similar way to the construction of a Lie algebra associated with a standard pentad. Precisely, we inductively construct a positively (respectively negatively) graded $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module $(\tilde{\pi}^+, \tilde{U}^+)$, $\tilde{U}^+ = \bigoplus_{m \geq 0} U_m^+$ (respectively $(\tilde{\pi}^-, \tilde{U}^-)$, $\tilde{U}^- = \bigoplus_{m \leq 0} U_m^-$) such that the “base space” U_0^+ (respectively the “top space” U_0^-) is the given \mathfrak{g} -module U . In general, the modules \tilde{U}^+ and \tilde{U}^- are infinite-dimensional. We shall try to embed $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and its module of the form \tilde{U}^+ into some graded Lie algebra. If we assume that B_0 is symmetric and that U has a \mathfrak{g} -submodule \mathcal{U} of $\text{Hom}(U, F)$ such that $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ is a standard pentad, then we can embed the objects $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and \tilde{U}^+ into some graded Lie algebra. Precisely, under these assumptions, we have that a pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$ is also standard, and, thus, we can embed the objects $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, \tilde{U}^+ , \tilde{U}^- into the graded Lie algebra $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$. In this situation, we have a “chain rule” of the Lie algebras associated with a standard pentad. This is the second main result of this paper.

This paper consists of three sections.

In section 2, we shall study the Lie algebras associated with a standard pentad. First, in section 2.1, we define the notion of standard pentads (see Definition 2.2) and construct a graded Lie algebra from a standard pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, which is denoted by $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) = \bigoplus_{n \in \mathbb{Z}} V_n$ (see Theorem 2.15). In section 2.2, we consider some properties of Lie algebras of the form $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ such that B_0 is symmetric. In these cases, we can also embed the bilinear form B_0 into $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, i.e. we can obtain a non-degenerate symmetric invariant bilinear form on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ whose restriction to $V_0 \times V_0$ coincides with B_0 (see Proposition 2.18). Moreover, the Lie algebra $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ can be characterized by the transitivity and the existence of such a bilinear form (see Theorem 2.20). Finally,

we give two lemmas on derivations on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ (see Lemmas 2.37 and 2.38).

In section 3, we shall study positively and negatively graded modules of a Lie algebra of the form $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. First, in sections 3.1 and 3.2, we shall construct positively graded $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module and negatively graded $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module from a \mathfrak{g} -module (π, U) , i.e. we shall give another proof of [9, Theorem 1.2] in the special cases where the graded Lie algebra is of the form $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. In section 3.1, we construct a family of \mathfrak{g} -modules $\{U_m^+\}_{m \geq 0}$ (respectively $\{U_m^-\}_{m \leq 0}$) from the pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and the \mathfrak{g} -module (π, U) by induction. In section 3.2, we define a structure of positively (respectively negatively) graded $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module on $\tilde{U}^+ := \bigoplus_{m \geq 0} U_m^+$ (respectively $\tilde{U}^- := \bigoplus_{m \leq 0} U_m^-$). We call this positively (respectively negatively) graded module of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ the *positive extension* (respectively *negative extension*) of U with respect to $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ (see Theorems 3.12 and 3.14). These modules are transitive and characterized by their transitivity (see Theorem 3.17). In sections 3.3 and 3.4, we try to construct a standard pentad which contains a Lie algebra of the form $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and its module of the form \tilde{U}^+ . For this, we need to assume that B_0 is symmetric and that U is embedded into some standard pentad $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$. In section 3.3, for the \mathfrak{g} -submodule \mathcal{U} of $\text{Hom}(U, F)$, we shall extend the canonical pairing $U \times \mathcal{U}$ to $\tilde{U}^+ \times \tilde{\mathcal{U}}^-$. Moreover, in section 3.4, we shall construct the Φ -map of $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ from the Φ -map of the pentad $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ inductively. Consequently, under the assumptions that $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ are standard pentads and that their bilinear form B_0 is symmetric, we can embed the Lie algebra $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and its module \tilde{U}^+ into a standard pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$. Finally, in section 3.5, we consider the graded Lie algebra $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ under the situation of sections 3.3 and 3.4. From the constructions of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, \tilde{U}^+ and $\tilde{\mathcal{U}}^-$, we can expect that this graded Lie algebra is written using the data $\mathfrak{g}, \rho, V, \mathcal{V}, B_0$ and U, \mathcal{U} . Indeed, we have the following result on the structures of Lie algebras:

$$L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L) \simeq L(\mathfrak{g}, \rho \oplus \pi, V \oplus U, \mathcal{V} \oplus \mathcal{U}, B_0)$$

up to grading. This is a chain rule in the theory of standard pentads (see Theorem 3.26).

NOTATION 1.1. In this paper, we regard a representation ρ of a Lie algebra \mathfrak{l} on V as a linear map $\rho : \mathfrak{l} \otimes V \rightarrow V$ which satisfies that

$$\rho([a, b] \otimes v) = \rho(a \otimes \rho(b \otimes v)) - \rho(b \otimes \rho(a \otimes v))$$

for any $a, b \in \mathfrak{l}$ and $v \in V$.

DEFINITION 1.2. In this paper, we say that a Lie algebra \mathfrak{l} is a \mathbb{Z} -graded Lie algebra or simply a *graded Lie algebra* if and only if there exist vector subspaces \mathfrak{l}_n of \mathfrak{l} for all $n \in \mathbb{Z}$ such that:

- $\mathfrak{l} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{l}_n$ and $[\mathfrak{l}_n, \mathfrak{l}_m] \subset \mathfrak{l}_{n+m}$ for any $n, m \in \mathbb{Z}$,
- \mathfrak{l} is generated by $\mathfrak{l}_{-1} \oplus \mathfrak{l}_0 \oplus \mathfrak{l}_1$.

In general, we do not assume that each \mathfrak{l}_n is finite-dimensional (cf. [2, Definition 1]).

Moreover, if \mathfrak{l} satisfies the following two conditions, we say that \mathfrak{l} is transitive (see [2, Definition 2]):

- for $x \in \mathfrak{l}_i$, $i \geq 0$, $[x, \mathfrak{l}_{-1}] = \{0\}$ implies $x = 0$,

- for $x \in I_i$, $i \leq 0$, $[x, I_1] = \{0\}$ implies $x = 0$.

DEFINITION 1.3. In this paper, we say that a module (ϖ^+, W) , $W = \bigoplus_{m \geq 0} W_m$ (respectively (ϖ^-, W) , $W = \bigoplus_{m \leq 0} W_m$) of a graded Lie algebra $\bigoplus_{n \in \mathbb{Z}} I_n$ is positively graded (respectively negatively graded) when $\varpi^+(I_n \otimes W_m) \subset W_{n+m}$ (respectively $\varpi^-(I_n \otimes W_m) \subset W_{n+m}$) for any n, m (cf. [9, Definition 0.1]), and, moreover, we say that a positively graded module (ϖ^+, W) (respectively a negatively graded module (ϖ^-, W)) is transitive when the following condition holds (cf. [9, Definition 1.1]):

$$\begin{aligned} & \text{for } w \in W_m, m \geq 1, \varpi^+(V_{-1} \otimes w) = \{0\} \text{ implies } w = 0 \\ & (\text{respectively for } w \in W_m, m \leq -1, \varpi^-(V_1 \otimes w) = \{0\} \text{ implies } w = 0). \end{aligned}$$

NOTATION 1.4. In this paper, we denote the set of all natural numbers, integers and complex numbers by \mathbb{N} , \mathbb{Z} and \mathbb{C} respectively. We denote the set of matrices of size $n \times m$ ($n, m \in \mathbb{N}$) whose entries are belong to a ring R by $M(n, m; R)$, the unit matrix and the zero matrix of size n by I_n and O_n respectively. Moreover, δ_{kl} stands for the Kronecker delta, $\text{Tr}(A)$ stands for the trace of a square matrix A .

2. Standard pentads and corresponding Lie algebras

2.1. Standard pentads. Let us start with the definitions of Φ -map and standard pentads.

DEFINITION 2.1 (Φ -map, cf. [8, Definition 1.9]). Let F be a field with characteristic 0. Let \mathfrak{g} be a Lie algebra with non-degenerate invariant bilinear form B_0 , $\rho : \mathfrak{g} \otimes V \rightarrow V$ a representation of \mathfrak{g} on a vector space V and \mathcal{V} a \mathfrak{g} -submodule of $\text{Hom}(V, F)$ all defined over F . We denote the canonical pairing between V and $\text{Hom}(V, F)$ by $\langle \cdot, \cdot \rangle$ and the canonical representation of \mathfrak{g} on \mathcal{V} by ϱ . Then, if a pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ has a linear map $\Phi_\rho : V \otimes \mathcal{V} \rightarrow \mathfrak{g}$ which satisfies an equation

$$(2.1) \quad B_0(a, \Phi_\rho(v \otimes \phi)) = \langle \rho(a \otimes v), \phi \rangle = -\langle v, \varrho(a \otimes \phi) \rangle$$

for any $a \in \mathfrak{g}$, $v \in V$ and $\phi \in \mathcal{V}$, we call it a Φ -map of the pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. Moreover, when a pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ has a Φ -map, we define a linear map $\Psi_\rho : \mathcal{V} \otimes V \rightarrow \mathfrak{g}$ by:

$$(2.2) \quad B_0(a, \Psi_\rho(\phi \otimes v)) = \langle v, \varrho(a \otimes \phi) \rangle = -\langle \rho(a \otimes v), \phi \rangle.$$

We call this map Ψ_ρ a Ψ -map of $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$.

In general, a pentad might not have a Φ -map. If a pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ has a Φ -map, then the equation (2.1) determines the linear map Φ_ρ uniquely. Moreover, we have an equation

$$\Phi_\rho(v \otimes \phi) + \Psi_\rho(\phi \otimes v) = 0$$

for any $v \in V$ and $\phi \in \mathcal{V}$.

DEFINITION 2.2 (Standard pentads). We retain to use the notation of Definition 2.1. If a pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ satisfies the following conditions, we call it a *standard pentad*:

$$(2.3) \quad \text{the restriction of } \langle \cdot, \cdot \rangle \text{ to } V \times \mathcal{V} \text{ is non-degenerate,}$$

$$(2.4) \quad \text{there exists a } \Phi\text{-map from } V \otimes \mathcal{V} \text{ to } \mathfrak{g}.$$

Lemma 2.3. *Under the notation of Definitions 2.1 and 2.2, we have the following claims:*

- (2.5) *if V is finite-dimensional, then a vector space \mathcal{V} satisfying (2.3) coincides with $\text{Hom}(V, F)$,*
- (2.6) *if \mathfrak{g} is finite-dimensional, then any pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ satisfies the condition (2.4).*

In particular, if both \mathfrak{g} and V are finite-dimensional, then any quadruplet $(\mathfrak{g}, \rho, V, B_0)$ can be naturally regarded as a standard pentad $(\mathfrak{g}, \rho, V, \text{Hom}(V, F), B_0)$.

Proof. The claim (2.5) is clear. Let us show the claim (2.6). If \mathfrak{g} is finite-dimensional, then the dual space of \mathfrak{g} can be identified with \mathfrak{g} . Precisely, if \mathfrak{g} is finite-dimensional, then any linear map $f : \mathfrak{g} \rightarrow F$ corresponds to some element $A \in \mathfrak{g}$ such that

$$f(a) = B_0(a, A)$$

for any $a \in \mathfrak{g}$. Thus, for any $v \in V$ and $\phi \in \mathcal{V}$, there exists an element of \mathfrak{g} which corresponds to a linear map $\mathfrak{g} \rightarrow F$ defined by

$$a \mapsto \langle \rho(a \otimes v), \phi \rangle.$$

It means that the pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ has the Φ -map. □

REMARK 2.4. If V is infinite-dimensional, then a submodule \mathcal{V} of $\text{Hom}(V, F)$ satisfying the condition (2.3) does not necessary coincide with $\text{Hom}(V, F)$.

REMARK 2.5. In general, a Lie algebra \mathfrak{g} and its module (ρ, V) might not have a \mathfrak{g} -submodule $\mathcal{V} \subset \text{Hom}(V, F)$ and a bilinear form B_0 such that a pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is standard.

EXAMPLE 2.6. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$, K be the Killing form on \mathfrak{g} and $\mathcal{L}(\mathfrak{g}) = \mathbb{C}[t, t^{-1}] \otimes \mathfrak{g}$ be the loop algebra (see [3, Ch.7]). Let $K_{\mathcal{L}}$ be a bilinear form on $\mathcal{L}(\mathfrak{g})$ defined by:

$$K_{\mathcal{L}}(t^n \otimes X, t^m \otimes Y) := \delta_{n+m,0} K(X, Y).$$

Clearly, the bilinear form $K_{\mathcal{L}}$ is non-degenerate and invariant. Thus, we can regard $\mathcal{L}(\mathfrak{g})$ itself as a $\mathcal{L}(\mathfrak{g})$ -submodule of $\text{Hom}(\mathcal{L}(\mathfrak{g}), \mathbb{C})$ via the non-degenerate invariant bilinear form $K_{\mathcal{L}}$. Then, a pentad $(\mathcal{L}(\mathfrak{g}), \text{ad}, \mathcal{L}(\mathfrak{g}), \mathcal{L}(\mathfrak{g}), K_{\mathcal{L}})$, where ad stands for the adjoint representation, is standard. In fact, we have the condition (2.3) clearly, and, we can identify the bracket product $\mathcal{L}(\mathfrak{g}) \times \mathcal{L}(\mathfrak{g}) \rightarrow \mathcal{L}(\mathfrak{g})$ with the Φ -map of $(\mathcal{L}(\mathfrak{g}), \text{ad}, \mathcal{L}(\mathfrak{g}), \mathcal{L}(\mathfrak{g}), K_{\mathcal{L}})$, denoted by Φ_{ad}^1 .

However, a pentad $(\mathcal{L}(\mathfrak{g}), \text{ad}, \mathcal{L}(\mathfrak{g}), \text{Hom}(\mathcal{L}(\mathfrak{g}), \mathbb{C}), K_{\mathcal{L}})$ is not standard since it does not have the Φ -map. In fact, if we assume that this pentad might have the Φ -map, denoted by Φ_{ad}^2 , and put

$$H_0 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad X_0 := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad Y_0 := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \in \mathfrak{g},$$

$$\phi_{Y_0} \in \text{Hom}(\mathcal{L}(\mathfrak{g}), \mathbb{C}), \quad \langle t^n \otimes X, \phi_{Y_0} \rangle := K(Y_0, X),$$

then an element $\Phi_{\text{ad}}^2((1 \otimes X_0) \otimes \phi_{Y_0}) \in \mathcal{L}(\mathfrak{g})$ satisfies the equation

$$\begin{aligned}
(2.7) \quad K_{\mathcal{L}}(t^n \otimes H_0, \Phi_{\text{ad}}^2((1 \otimes X_0) \otimes \phi_{Y_0})) &= \langle [t^n \otimes H_0, 1 \otimes X_0], \phi_{Y_0} \rangle \\
&= \langle t^n \otimes 2X_0, \phi_{Y_0} \rangle \\
&= K(Y_0, 2X_0) \\
&= 8
\end{aligned}$$

for any $n \in \mathbb{Z}$. The Lie algebra $\mathcal{L}(\mathfrak{g})$ does not have an element satisfying (2.7) for any $n \in \mathbb{Z}$, and, thus, the pentad $(\mathcal{L}(\mathfrak{g}), \text{ad}, \mathcal{L}(\mathfrak{g}), \text{Hom}(\mathcal{L}(\mathfrak{g}), \mathbb{C}), K_{\mathcal{L}})$ does not have the Φ -map.

On the Φ -map and Ψ -map of a standard pentad, we have similar properties to ones of the Φ -map and Ψ -map of a standard quadruplet (see [8]).

Proposition 2.7. *The Φ -map and the Ψ -map of a standard quadruplet $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ are homomorphisms of Lie modules. (cf. [8, Proposition 1.3]).*

Proof. We can prove it by the same way to [8, Proposition 1.3]. \square

Definition 2.8. Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad. For each element $v \in V$ and $\phi \in \mathcal{V}$, we define linear maps $\Phi_{\rho, v} \in \text{Hom}(\mathcal{V}, \mathfrak{g})$ and $\Psi_{\rho, \phi} \in \text{Hom}(V, \mathfrak{g})$ by:

$$\Phi_{\rho, v}(\psi) := \Phi_{\rho}(v \otimes \psi), \quad \Psi_{\rho, \phi}(u) := \Psi_{\rho}(\phi \otimes u)$$

for any $u \in V$ and $\psi \in \mathcal{V}$. Moreover, we define the following linear maps:

$$\begin{aligned}
\Phi_{\rho}^{\circ} : V &\rightarrow \text{Hom}(\mathcal{V}, \mathfrak{g}) & \Psi_{\rho}^{\circ} : \mathcal{V} &\rightarrow \text{Hom}(V, \mathfrak{g}) \\
v &\mapsto \Phi_{\rho, v}, & \phi &\mapsto \Psi_{\rho, \phi}.
\end{aligned}$$

To simplify, we denote $\Phi_{\rho, v}(\psi)$ and $\Psi_{\rho, \phi}(u)$ by $v(\psi)$ and $\phi(u)$ respectively.

Definition 2.9. Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad. Put $V_0 := \mathfrak{g}$, $V_1 := V$ and $V_{-1} := \mathcal{V}$ and denote the canonical representations of \mathfrak{g} on V_0 and $V_{\pm 1}$ by ρ_0 and $\rho_{\pm 1}$. We define homomorphisms of \mathfrak{g} -modules p_0 and q_0 by:

$$\begin{aligned}
p_0 : V_1 \otimes V_0 &\rightarrow V_1 \\
v_1 \otimes a &\mapsto -\rho_1(a \otimes v_1), \\
q_0 : V_{-1} \otimes V_0 &\rightarrow V_{-1} \\
\phi_{-1} \otimes b &\mapsto -\rho_{-1}(b \otimes \phi_{-1}).
\end{aligned}$$

Moreover, we define homomorphisms of \mathfrak{g} -modules p_1 and q_{-1} by:

$$\begin{aligned}
p_1 : V_1 \otimes V_1 &\rightarrow \text{Hom}(V_{-1}, V_1) \\
v_1 \otimes u_1 &\mapsto (\eta_{-1} \mapsto \rho_1(v_1(\eta_{-1}) \otimes u_1) - \rho_1(u_1(\eta_{-1}) \otimes v_1)), \\
q_{-1} : V_{-1} \otimes V_{-1} &\rightarrow \text{Hom}(V_1, V_{-1}) \\
\phi_{-1} \otimes \psi_{-1} &\mapsto (\xi_1 \mapsto \rho_{-1}(\phi_{-1}(\xi_1) \otimes \psi_{-1}) - \rho_{-1}(\psi_{-1}(\xi_1) \otimes \phi_{-1})),
\end{aligned}$$

where $v_1(\eta_{-1}) \in V_0$ and $\phi_{-1}(\xi_1) \in V_0$ stand for $\Phi_{\rho, v_1}(\eta_{-1})$ and $\Psi_{\rho, \phi_{-1}}(\xi_1)$ respectively.

Moreover, suppose that $i \geq 2$ and there exist \mathfrak{g} -modules (ρ_{i-1}, V_{i-1}) and (ρ_{-i+1}, V_{-i+1}) and homomorphisms of \mathfrak{g} -modules $p_{i-1} : V_1 \otimes V_{i-1} \rightarrow \text{Hom}(V_{-1}, V_{i-1})$ and $q_{-i+1} : V_{-1} \otimes V_{-i+1} \rightarrow \text{Hom}(V_1, V_{-i+1})$. Then, we put $V_i := \text{Im } p_{i-1}$, $V_{-i} := \text{Im } q_{-i+1}$ and define linear maps p_i, q_{-i} by:

$$\begin{aligned}
p_i &: V_1 \otimes V_i \rightarrow \text{Hom}(V_{-1}, V_i) \\
v_1 \otimes u_i &\mapsto (\eta_{-1} \mapsto \rho_i(v_1(\eta_{-1}) \otimes u_i) + p_{i-1}(v_1 \otimes u_i(\eta_{-1}))), \\
q_{-i} &: V_{-1} \otimes V_{-i} \rightarrow \text{Hom}(V_1, V_{-i}) \\
\phi_{-1} \otimes \psi_{-i} &\mapsto (\xi_1 \mapsto \rho_{-i}(\phi_{-1}(\xi_1) \otimes \psi_{-i}) + q_{-i+1}(\phi_{-1} \otimes \psi_{-i}(\xi_1))),
\end{aligned}$$

where $u_i(\eta_{-1}) \in V_{i-1}$ and $\psi_{-i}(\xi_1) \in V_{-i+1}$ are the images of η_{-1} and ξ_1 via u_i and ψ_{-i} respectively. Then, the linear maps p_i and q_{-i} are homomorphisms of \mathfrak{g} -modules (cf. [8, Proposition 1.10]). We denote the images of p_i and q_{-i} by V_{i+1} and V_{-i-1} and the canonical representations of \mathfrak{g} on V_{i+1} and V_{-i-1} by ρ_{i+1} and ρ_{-i-1} respectively. Thus, inductively, we obtain \mathfrak{g} -modules V_n and representations ρ_n of \mathfrak{g} on V_n for all $n \in \mathbb{Z}$. We call V_n the n -graduation of $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$.

REMARK 2.10. For any $v_1 \in V_1$ and $\phi_{-1} \in V_{-1}$, we have

$$\begin{aligned}
p_1(v_1 \otimes v_1)(\eta_{-1}) &= \rho_1(v_1(\eta_{-1}) \otimes v_1) - \rho_1(v_1(\eta_{-1}) \otimes v_1) = 0, \\
q_{-1}(\phi_{-1} \otimes \phi_{-1})(\xi_1) &= \rho_{-1}(\phi_{-1}(\xi_1) \otimes \phi_{-1}) - \rho_{-1}(\phi_{-1}(\xi_1) \otimes \phi_{-1}) = 0.
\end{aligned}$$

In general, we do not assume that ρ and ϱ are surjective, i.e. we do not assume that $V_1 = \text{Im } p_0$ and $V_{-1} = \text{Im } q_0$. In particular cases where these linear maps are surjective, we have the following proposition.

Proposition 2.11. *If $\rho : \mathfrak{g} \otimes V \rightarrow V$ and $\varrho : \mathfrak{g} \otimes \mathcal{V} \rightarrow \mathcal{V}$ are surjective, then Φ_ρ° and Ψ_ρ° are injective, and, thus, V and \mathcal{V} can be regarded as \mathfrak{g} -submodules of $\text{Hom}(V_{-1}, V_0)$ and $\text{Hom}(V_1, V_0)$ respectively.*

Proof. To show this proposition, we use the condition (2.3). Let us show that the linear map Φ_ρ° is injective. We take an arbitrary element $v \in V$ which satisfies that $\Phi_{\rho,v} = 0$. Then we have

$$(2.8) \quad 0 = B_0(a, \Phi_{\rho,v}(\phi)) = \langle \rho(a \otimes v), \phi \rangle = -\langle v, \varrho(a \otimes \phi) \rangle$$

for all $a \in \mathfrak{g}$ and $\phi \in \mathcal{V}$. By the condition (2.3) and the assumption that ϱ is surjective, we have that $v = 0$. Therefore, we obtain that Φ_ρ° is injective. Similarly, we can show that Ψ_ρ° is injective. \square

DEFINITION 2.12. We define the following bilinear maps

$$[\cdot, \cdot]_n^0 : V_0 \times V_n \rightarrow V_n, \quad [\cdot, \cdot]_n^1 : V_1 \times V_n \rightarrow V_{n+1}, \quad [\cdot, \cdot]_n^{-1} : V_{-1} \times V_n \rightarrow V_{n-1}$$

by:

$$\begin{aligned}
[a_0, z_n]_n^0 &:= \rho_n(a_0 \otimes z_n), \\
[x_1, z_n]_n^1 &:= \begin{cases} p_n(x_1 \otimes z_n) & (n \geq 0) \\ -z_n(x_1) & (n \leq -1) \end{cases}, \\
[y_{-1}, z_n]_n^{-1} &:= \begin{cases} -z_n(y_{-1}) & (n \geq 1) \\ q_n(y_{-1} \otimes z_n) & (n \leq 0) \end{cases}
\end{aligned}$$

where $a_0 \in V_0$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$ and $z_n \in V_n$. Note that $z_n(x_1)$ stands for $\Psi_{\rho, z_{-1}}(x_1)$ when $n = -1$ and the image of x_1 via $z_n \in \text{Hom}(V_1, V_{n+1})$ when $n \leq -2$. Moreover, for $i \geq 1$, we define the following bilinear maps

$$[\cdot, \cdot]_n^{i+1} : V_{i+1} \times V_n \rightarrow V_{i+n+1}, \quad [\cdot, \cdot]_n^{-i-1} : V_{-i-1} \times V_n \rightarrow V_{-i+n-1}$$

by:

$$(2.9) \quad [p_i(x_1 \otimes z_i), w_n]_n^{i+1} := [x_1, [z_i, w_n]_n^i]_{i+n}^1 - [z_i, [x_1, w_n]_n^1]_{n+1}^i \\ (x_1 \in V_1, z_i \in V_i, w_n \in V_n)$$

and

$$(2.10) \quad [q_{-i}(y_{-1} \otimes \omega_{-i}), w_n]_n^{-i-1} := [y_{-1}, [\omega_{-i}, w_n]_n^{-i}]_{-i+n}^{-1} - [\omega_{-i}, [y_{-1}, w_n]_n^{-1}]_{n-1}^{-i} \\ (y_{-1} \in V_{-1}, \omega_{-i} \in V_{-i}, w_n \in V_n)$$

inductively. Then the bilinear maps (2.9) and (2.10) are well-defined. It can be shown by the same argument to the argument of [8, Propositions 2.5 and 2.6]. Consequently, we can define a bilinear map $[\cdot, \cdot]_m^n : V_n \times V_m \rightarrow V_{n+m}$ for any $n, m \in \mathbb{Z}$.

DEFINITION 2.13. For a standard pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, we denote a direct sum of its n -graduations by $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, i.e.

$$L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) := \bigoplus_{n \in \mathbb{Z}} V_n.$$

Moreover, we define a bilinear map $[\cdot, \cdot] : L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) \times L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) \rightarrow L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ by

$$(2.11) \quad [x_n, y_m] := [x_n, y_m]_m^n$$

for any $n, m \in \mathbb{Z}$, $x_n \in V_n$ and $y_m \in V_m$.

Proposition 2.14. *This bilinear map $[\cdot, \cdot]$ satisfies the following equations*

$$(2.12) \quad [x, y] + [y, x] = 0,$$

$$(2.13) \quad [x, [y, z]] + [z, [x, y]] + [y, [z, x]] = 0$$

for any $x, y, z \in L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$.

Proof. We can prove it by the same argument to the argument of [8, Propositions 2.9 and 2.10]. \square

As a corollary of Proposition 2.14, we have the following theorem immediately.

Theorem 2.15 (Lie algebra associated with a standard pentad). *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad over a field F with characteristic 0. Then the vector space $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) = \bigoplus_{n \in \mathbb{Z}} V_n$ is a graded Lie algebra with a bracket product $[\cdot, \cdot]$ defined in Definition 2.13. We call this graded Lie algebra the Lie algebra associated with $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ (cf. [8, Theorem 2.11]).*

REMARK 2.16. Note that we can prove Theorem 2.15 without the assumption that the bilinear form B_0 is symmetric.

Note that $V_0 = \mathfrak{g}$ and that the V_0 -modules V_0, V_1, V_{-1} are isomorphic to $\mathfrak{g}, V, \mathcal{V}$ respectively. In this sense, we can say that the objects $\mathfrak{g}, (\rho, V), (\varrho, \mathcal{V})$ can be embedded into $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$.

In particular, when ρ and ϱ are faithful and surjective, we have a similar result on the structure of a graded Lie algebra of the form $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ to the result which is obtained by H. Rubenthaler in [7, Proposition 3.4.2]. We can show the following proposition by Proposition 2.11 immediately.

Proposition 2.17. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad. If both $\rho : \mathfrak{g} \otimes V \rightarrow V$ and $\varrho : \mathfrak{g} \otimes \mathcal{V} \rightarrow \mathcal{V}$ are faithful and surjective, then the graded Lie algebra $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is transitive.*

2.2. Standard pentads with a symmetric bilinear form. In the previous section, we proved that for any standard pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, there exists a graded Lie algebra such that \mathfrak{g}, ρ, V and \mathcal{V} can be embedded into it. In this section, we discuss cases where B_0 is symmetric. In these cases, we can also embed B_0 into $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and we can obtain some useful properties.

Proposition 2.18. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad such that B_0 is symmetric. We define a symmetric bilinear form B_L on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ inductively as follows:*

$$\begin{cases} B_L(a, b) = B_0(a, b), \\ B_L(v, \phi) = \langle v, \phi \rangle, \\ B_L(p_i(v_1 \otimes u_i), q_{-i}(\phi_{-1} \otimes \psi_{-i})) = B_L(u_i, [q_{-i}(\phi_{-1} \otimes \psi_{-i}), v_1]), \\ B_L(x_n, y_m) = 0 \end{cases}$$

for any $a, b \in V_0, v \in V, \phi \in \mathcal{V}, i \geq 1, v_1 \in V_1, \phi_{-1} \in V_{-1}, u_i \in V_i, \psi_{-i} \in V_{-i}, n, m \in \mathbb{Z}, n + m \neq 0, x_n \in V_n$ and $y_m \in V_m$. Then B_L is a non-degenerate symmetric invariant bilinear form on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ (cf. [8, Proposition 3.2]).

Proof. Note that it is clear that the restriction of B_L to $V_0 \times V_0$ and $V_1 \times V_{-1}$ is well-defined. Let us show the well-definedness of B_L on $V_2 \times V_{-2}$. For any $v_1, u_1 \in V_1$ and $\phi_{-1}, \psi_{-1} \in V_{-1}$, we have

$$\begin{aligned} (2.14) \quad B_L(u_1, [q_{-1}(\phi_{-1} \otimes \psi_{-1}), v_1]) &= B_L(u_1, [[\phi_{-1}, v_1], \psi_{-1}] + [\phi_{-1}, [\psi_{-1}, v_1]]) \\ &= \langle u_1, [[\phi_{-1}, v_1], \psi_{-1}] + [\phi_{-1}, [\psi_{-1}, v_1]] \rangle \\ &= B_0([\phi_{-1}, v_1], \psi_{-1}(u_1)) - B_0([\psi_{-1}, v_1], \phi_{-1}(u_1)) \\ &= B_0([\phi_{-1}, v_1], \psi_{-1}(u_1)) - B_0(\phi_{-1}(u_1), [\psi_{-1}, v_1]) \\ &\quad \text{(by the assumption that } B_0 \text{ is symmetric)} \\ &= B_0([v_1, \phi_{-1}], u_1(\psi_{-1})) - B_0(u_1(\phi_{-1}), [v_1, \psi_{-1}]) \\ &= \langle [[v_1, \phi_{-1}], u_1] + [v_1, [u_1, \phi_{-1}]], \psi_{-1} \rangle \\ &= B_L([p_1(v_1 \otimes u_1), \phi_{-1}], \psi_{-1}). \end{aligned}$$

Thus, if $v_1^1, \dots, v_1^l, u_1^1, \dots, u_1^l \in V_1$ and $\phi_{-1}^1, \dots, \phi_{-1}^k, \psi_{-1}^1, \dots, \psi_{-1}^k \in V_{-1}$ satisfy equations

$$\sum_{s=1}^l p_1(v_1^s \otimes u_1^s) = 0, \quad \sum_{t=1}^k q_{-1}(\phi_{-1}^t \otimes \psi_{-1}^t) = 0,$$

then

$$\sum_{s=1}^l B_L(u_1^s, [q_{-1}(\phi_{-1} \otimes \psi_{-1}), v_1^s]) = \sum_{s=1}^l B_L([p_1(v_1^s \otimes u_1^s), \phi_{-1}], \psi_{-1}) = 0,$$

$$\sum_{t=1}^k B_L(u_1, [q_{-1}(\phi_{-1}^t \otimes \psi_{-1}^t), v_1]) = 0$$

for any $v_1, u_1 \in V_1$ and $\phi_{-1}, \psi_{-1} \in V_{-1}$, that is, we have the well-definedness of B_L on $V_2 \times V_{-2}$. This $B_L|_{V_2 \times V_{-2}}$ is \mathfrak{g} -invariant. Moreover, by a similar argument, we have the well-definedness of B_L on $V_i \times V_{-i}$ for each $i \geq 3$ by induction (see [8, section 1.2]). Consequently, we can show the well-definedness of B_L on the whole $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and that B_L is non-degenerate symmetric invariant by the same argument as the argument in [8, section 1.2 and Proposition 3.2]. \square

REMARK 2.19. We need the assumption that B_0 is symmetric to show that the bilinear form B_L is $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -invariant. Precisely, we need this assumption to show an equation

$$B_L(v_1, [\phi_{-1}, a]) = B_L([v_1, \phi_{-1}], a)$$

for any $a \in V_0, v_1 \in V_1, \phi_{-1} \in V_{-1}$.

Under the assumption that B_0 is symmetric, the graded Lie algebra is characterized by the existence of such a bilinear form. The following is a proposition concerning the “universality” and “uniqueness” of Lie algebras associated with a standard pentad with a symmetric bilinear form.

Theorem 2.20. *Let $\mathfrak{Q} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{Q}_n$ be a graded Lie algebra which has a non-degenerate symmetric invariant bilinear form $B_{\mathfrak{Q}}$. If \mathfrak{Q} and $B_{\mathfrak{Q}}$ satisfy the following conditions, then a pentad $(\mathfrak{Q}_0, \text{ad}, \mathfrak{Q}_1, \mathfrak{Q}_{-1}, B_{\mathfrak{Q}}|_{\mathfrak{Q}_0 \times \mathfrak{Q}_0})$ is standard and \mathfrak{Q} is isomorphic to $L(\mathfrak{Q}_0, \text{ad}, \mathfrak{Q}_1, \mathfrak{Q}_{-1}, B_{\mathfrak{Q}}|_{\mathfrak{Q}_0 \times \mathfrak{Q}_0})$:*

$$(2.15) \quad \mathfrak{Q}_{i+1} = [\mathfrak{Q}_1, \mathfrak{Q}_i], \mathfrak{Q}_{-i-1} = [\mathfrak{Q}_{-1}, \mathfrak{Q}_{-i}] \text{ for all } i \geq 1,$$

$$(2.16) \quad \text{the restriction of } B_{\mathfrak{Q}} \text{ to } \mathfrak{Q}_i \times \mathfrak{Q}_{-i} \text{ is non-degenerate for any } i \geq 0,$$

where ad stands for the adjoint representation of \mathfrak{Q} on itself (cf. [8, Proposition 3.3]).

Proof. First of all, let us check that the pentad $(\mathfrak{Q}_0, \text{ad}, \mathfrak{Q}_1, \mathfrak{Q}_{-1}, B_{\mathfrak{Q}}|_{\mathfrak{Q}_0 \times \mathfrak{Q}_0})$ is standard. By (2.16), we can obtain that $B_{\mathfrak{Q}}|_{\mathfrak{Q}_0 \times \mathfrak{Q}_0}$ is non-degenerate and that \mathfrak{Q}_1 and \mathfrak{Q}_{-1} satisfy the condition (2.3). It is easy to show that we can identify the restriction of the bracket product $[\cdot, \cdot]$ of \mathfrak{Q} to $\mathfrak{Q}_1 \times \mathfrak{Q}_{-1} \rightarrow \mathfrak{Q}_0$ with the Φ -map of the pentad $(\mathfrak{Q}_0, \text{ad}, \mathfrak{Q}_1, \mathfrak{Q}_{-1}, B_{\mathfrak{Q}}|_{\mathfrak{Q}_0 \times \mathfrak{Q}_0})$. Thus, the condition (2.4) holds.

We denote the n -graduation of $(\mathfrak{Q}_0, \text{ad}, \mathfrak{Q}_1, \mathfrak{Q}_{-1}, B_{\mathfrak{Q}}|_{\mathfrak{Q}_0 \times \mathfrak{Q}_0})$ by $(\mathfrak{Q})_n$ for any $n \in \mathbb{Z}$ and a bilinear form on $L(\mathfrak{Q}_0, \text{ad}, \mathfrak{Q}_1, \mathfrak{Q}_{-1}, B_{\mathfrak{Q}}|_{\mathfrak{Q}_0 \times \mathfrak{Q}_0})$ obtained in Proposition 2.18 by $(B)_{\mathfrak{Q}}$. Let $\sigma_0 : (\mathfrak{Q})_0 \rightarrow \mathfrak{Q}_0$ and $\sigma_{\pm 1} : (\mathfrak{Q})_{\pm 1} \rightarrow \mathfrak{Q}_{\pm 1}$ be the identity maps respectively. Then the linear maps σ_0 and $\sigma_{\pm 1}$ satisfy the following equations:

$$(2.17) \quad [\sigma_0(a), \sigma_{\pm 1}(x_{\pm 1})] = \sigma_{\pm 1}([a, x_{\pm 1}]),$$

$$(2.18) \quad [\sigma_1(x_1), \sigma_{-1}(x_{-1})] = \sigma_0([x_1, x_{-1}])$$

for any $a \in (\mathfrak{V})_0$ and $x_{\pm 1} \in (\mathfrak{V})_{\pm 1}$. Indeed, the equation (2.17) is clear, and, we have

$$(2.19) \quad \begin{aligned} B_{\mathfrak{V}}(\sigma_0(b), [\sigma_1(x_1), \sigma_{-1}(x_{-1})]) &= B_{\mathfrak{V}}([\sigma_0(b), \sigma_1(x_1)], \sigma_{-1}(x_{-1})) = B_{\mathfrak{V}}(\sigma_1([b, x_1]), \sigma_{-1}(x_{-1})) \\ &= (B)_{\mathfrak{V}}([b, x_1], x_{-1}) = (B)_{\mathfrak{V}}(b, [x_1, x_{-1}]) = B_{\mathfrak{V}}(\sigma_0(b), \sigma_0([x_1, x_{-1}])) \end{aligned}$$

for any $b \in (\mathfrak{V})_0$. Thus, we can obtain the equation (2.18).

For each $i \geq 1$, we define linear maps $\sigma_{i+1} : (\mathfrak{V})_{i+1} \rightarrow \mathfrak{V}_{i+1}$ and $\sigma_{-i-1} : (\mathfrak{V})_{-i-1} \rightarrow \mathfrak{V}_{-i-1}$ by:

$$(2.20) \quad \sigma_{i+1} : p_i(x_1 \otimes z_i) \mapsto [\sigma_1(x_1), \sigma_i(z_i)],$$

$$(2.21) \quad \sigma_{-i-1} : q_{-i}(x_{-1} \otimes z_{-i}) \mapsto [\sigma_{-1}(x_{-1}), \sigma_{-i}(z_{-i})]$$

for any $x_{\pm 1} \in (\mathfrak{V})_{\pm 1}$ and $z_{\pm i} \in (\mathfrak{V})_{\pm i}$ inductively. Note that it follows from (2.17) that the linear maps σ_1 and σ_{-1} on $\rho(\mathfrak{g} \otimes V)$ and $\varrho(\mathfrak{g} \otimes \mathcal{V})$ defined by the same equations as (2.20) and (2.21) where $i = 0$ coincide with the identity maps respectively. We can prove that the linear maps σ_n ($n \in \mathbb{Z}$) are well-defined and satisfy

$$(2.22) \quad [\sigma_0(a), \sigma_n(z_n)] = \sigma_n([a, z_n]),$$

$$(2.23) \quad [\sigma_{\pm 1}(x_{\pm 1}), \sigma_n(z_n)] = \sigma_{n\pm 1}([x_{\pm 1}, z_n])$$

for any $n \in \mathbb{Z}$, $a \in (\mathfrak{V})_0$, $x_{\pm 1} \in (\mathfrak{V})_{\pm 1}$ and $z_n \in (\mathfrak{V})_n$ by a similar argument to the argument of [8, Proposition 3.3]. Then a linear map $\sigma : L(\mathfrak{V}_0, \text{ad}, \mathfrak{V}_1, \mathfrak{V}_{-1}, B_{\mathfrak{V}} |_{\mathfrak{V}_0 \times \mathfrak{V}_0}) \rightarrow \mathfrak{V}$ defined by

$$(2.24) \quad \sigma(z_n) := \sigma_n(z_n),$$

where $n \in \mathbb{Z}$ and $z_n \in (\mathfrak{V})_n \subset L(\mathfrak{V}_0, \text{ad}, \mathfrak{V}_1, \mathfrak{V}_{-1}, B_{\mathfrak{V}} |_{\mathfrak{V}_0 \times \mathfrak{V}_0})$, is an isomorphism of Lie algebras. We can also prove this by a similar argument to the argument of [8, Proposition 3.3]. \square

As a corollary of Theorem 2.20, we can say that the theory of standard pentads is an extension of the theory of standard quadruplets.

Proposition 2.21. *Let $(\mathfrak{g}, \rho, V, B_0)$ be a standard quadruplet (see [8, Definition 1.9]). Then the Lie algebra $L(\mathfrak{g}, \rho, V, B_0)$ associated with $(\mathfrak{g}, \rho, V, B_0)$ (see [8, Theorem 2.11]) is isomorphic to the Lie algebra $L(\mathfrak{g}, \rho, V, \text{Hom}(V, \mathbb{C}), B_0)$.*

Definition 2.22. Let $(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1)$ and $(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$ be standard pentads. We say that these pentads are *equivalent* if and only if there exists an isomorphism of Lie algebras $\tau : \mathfrak{g}^1 \rightarrow \mathfrak{g}^2$, linear isomorphisms $\sigma : V^1 \rightarrow V^2$, $\varsigma : \mathcal{V}^1 \rightarrow \mathcal{V}^2$ and a non-zero element $c \in F$ such that

$$(2.25) \quad \sigma(\rho^1(a^1 \otimes x^1)) = \rho^2(\tau(a^1) \otimes \sigma(x^1)),$$

$$(2.26) \quad \varsigma(\varrho^1(a^1 \otimes y^1)) = \varrho^2(\tau(a^1) \otimes \varsigma(y^1)),$$

$$(2.27) \quad \langle x^1, y^1 \rangle^1 = \langle \sigma(x^1), \varsigma(y^1) \rangle^2,$$

$$(2.28) \quad B_0^1(a^1, b^1) = cB_0^2(\tau(a^1), \tau(b^1))$$

where $a^1, b^1 \in \mathfrak{g}^1$, $x^1 \in V^1$, $y^1 \in \mathcal{V}^1$ and $\langle \cdot, \cdot \rangle^i$ stands for the pairing between V^i and \mathcal{V}^i ($i = 1, 2$). We denote this equivalence relation by

$$(2.29) \quad (\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1) \simeq (\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2).$$

REMARK 2.23. Note that if V is finite-dimensional, then linear isomorphisms τ, σ satisfying (2.25) induce a linear isomorphism from $\mathcal{V}^1 = \text{Hom}(V^1, F)$ to $\mathcal{V}^2 = \text{Hom}(V^2, F)$ satisfying (2.26) and (2.27).

Proposition 2.24. *If standard pentads $(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1)$ and $(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$ are equivalent, then the Lie algebras associated with them are isomorphic, i.e. we have*

$$(2.30) \quad L(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1) \simeq L(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$$

(cf. [8, Proposition 3.6]).

Proof. We denote the n -graduation of $(\mathfrak{g}^i, \rho^i, V^i, \mathcal{V}^i, B_0^i)$ by V_n^i ($i = 1, 2$) for all $n \in \mathbb{Z}$ and the bilinear forms on $L(\mathfrak{g}^i, \rho^i, V^i, \mathcal{V}^i, B_0^i)$ defined in Proposition 2.18 by B_L^i ($i = 1, 2$). Under the notation of Definition 2.22, we define linear maps $\sigma_0 := \tau : V_0^1 \rightarrow V_0^2$, $\sigma_1 := \frac{1}{c}\sigma : V_1^1 \rightarrow V_1^2$ and $\sigma_{-1} := \varsigma : V_{-1}^1 \rightarrow V_{-1}^2$. Then, these linear maps σ_0 and $\sigma_{\pm 1}$ satisfy the same equations as (2.17) and (2.18). In fact, the equation (2.17) is clear, and, we have

$$\begin{aligned} B_0^2(\sigma_0(a_0^1), [\sigma_1(x_1^1), \sigma_{-1}(y_{-1}^1)]) &= B_L^2(\sigma_1([a_0^1, x_1^1]), \sigma_{-1}(y_{-1}^1)) \\ &= \frac{1}{c} B_L^1([a_0^1, x_1^1], y_{-1}^1) = \frac{1}{c} B_0^1(a_0^1, [x_1^1, y_{-1}^1]) = B_0^2(\sigma_0(a_0^1), \sigma_0([x_1^1, y_{-1}^1])) \end{aligned}$$

for any $a_0^1 \in V_0^1$, $x_1^1 \in V_1^1$ and $y_{-1}^1 \in V_{-1}^1$. Thus, we have the equation (2.18). Then, by the same argument as the argument in proof of Theorem 2.20, we can construct an isomorphism of Lie algebras from $L(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1)$ to $L(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$. \square

REMARK 2.25. The converse of Proposition 2.22 is not true. In fact, we have an example of two non-equivalent pentads such that the corresponding Lie algebras are isomorphic (see [8, pp. 398–399]).

DEFINITION 2.26. Let $(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1)$ and $(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$ be standard pentads. Let $\rho^1 \boxplus \rho^2$ and $\varrho^1 \boxplus \varrho^2$ be representations of $\mathfrak{g}^1 \oplus \mathfrak{g}^2$ on $V^1 \oplus V^2$ and $\mathcal{V}^1 \oplus \mathcal{V}^2$ defined by:

$$\begin{aligned} (\rho^1 \boxplus \rho^2)((a^1, a^2) \otimes (v^1, v^2)) &:= (\rho^1(a^1 \otimes v^1), \rho^2(a^2 \otimes v^2)), \\ (\varrho^1 \boxplus \varrho^2)((b^1, b^2) \otimes (\phi^1, \phi^2)) &:= (\varrho^1(b^1 \otimes \phi^1), \varrho^2(b^2 \otimes \phi^2)) \end{aligned}$$

where $a^i, b^i \in \mathfrak{g}^i$, $v^i \in V^i$, $\phi^i \in \mathcal{V}^i$ ($i = 1, 2$). Let $B_0^1 \oplus B_0^2$ be a bilinear form on $\mathfrak{g}^1 \oplus \mathfrak{g}^2$ defined by:

$$(2.31) \quad (B_0^1 \oplus B_0^2)((a^1, a^2), (b^1, b^2)) := B_0^1(a^1, b^1) + B_0^2(a^2, b^2)$$

where $a^i, b^i \in \mathfrak{g}^i$ ($i = 1, 2$). Then, clearly, a pentad $(\mathfrak{g}^1 \oplus \mathfrak{g}^2, \rho^1 \boxplus \rho^2, V^1 \oplus V^2, \mathcal{V}^1 \oplus \mathcal{V}^2, B_0^1 \oplus B_0^2)$ is also a standard pentad. We call it a *direct sum* of $(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1)$ and $(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$ and denote it by $(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1) \oplus (\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$.

Proposition 2.27. *Let $(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1)$ and $(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$ be standard pentads. Then the Lie algebra $L((\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1) \oplus (\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2))$ is isomorphic to $L(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1) \oplus L(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$ (cf. [8, Proposition 3.9]).*

Proof. We retain to use the notation of Proposition 2.24. Then, we have the following \mathbb{Z} -grading of $L(\mathfrak{g}^1, \rho^1, V^1, \mathcal{V}^1, B_0^1) \oplus L(\mathfrak{g}^2, \rho^2, V^2, \mathcal{V}^2, B_0^2)$:

$$(2.32) \quad L(g^1, \rho^1, V^1, \mathcal{V}^1, B_0^1) \oplus L(g^2, \rho^2, V^2, \mathcal{V}^2, B_0^2) = \bigoplus_{n \in \mathbb{Z}} (V_n^1 \oplus V_n^2).$$

By Theorem 2.20, we have our claim. \square

DEFINITION 2.28. Let $(g, \rho, V, \mathcal{V}, B_0)$ be a standard pentad. We say that $(g, \rho, V, \mathcal{V}, B_0)$ is *decomposable* if and only if there exist standard pentads $(a, \rho_a, V_a, \mathcal{V}_a, B_{0,a})$ and $(b, \rho_b, V_b, \mathcal{V}_b, B_{0,b})$ such that

$$(2.33) \quad (\dim a + \dim V_a)(\dim b + \dim V_b) \neq 0,$$

$$(2.34) \quad (g, \rho, V, \mathcal{V}, B_0) \simeq (a, \rho_a, V_a, \mathcal{V}_a, B_{0,a}) \oplus (b, \rho_b, V_b, \mathcal{V}_b, B_{0,b}).$$

If $(g, \rho, V, \mathcal{V}, B_0)$ is not decomposable, we say that $(g, \rho, V, \mathcal{V}, B_0)$ is *indecomposable*.

DEFINITION 2.29. Let $(g, \rho, V, \mathcal{V}, B_0)$ be a standard pentad. We say that $(g, \rho, V, \mathcal{V}, B_0)$ is *reducible* if and only if there exist an ideal a of g and g -submodules V_a and \mathcal{V}_a of V and \mathcal{V} satisfying that:

$$(2.35) \quad \{0\} \neq \mathcal{V}_a \oplus a \oplus V_a \subsetneq \mathcal{V} \oplus a \oplus V,$$

$$(2.36) \quad \rho(a \otimes V), \rho(g \otimes V_a) \subset V_a \text{ and } \varrho(a \otimes \mathcal{V}), \varrho(g \otimes \mathcal{V}_a) \subset \mathcal{V}_a,$$

$$(2.37) \quad \Phi_\rho(V_a \otimes \mathcal{V}), \Phi_\rho(V \otimes \mathcal{V}_a) \subset a.$$

And, we say that $(g, \rho, V, \mathcal{V}, B_0)$ is *irreducible* if and only if it is not reducible.

REMARK 2.30. If a standard pentad is irreducible, then it is indecomposable.

Proposition 2.31. Let $(g, \rho, V, \mathcal{V}, B_0)$ be an irreducible standard pentad. Then the representations $\rho : g \otimes V \rightarrow V$, $\varrho : g \otimes \mathcal{V} \rightarrow \mathcal{V}$ and the Φ -map $\Phi_\rho : V \otimes \mathcal{V} \rightarrow g$ are surjective.

Proof. If $\varrho(g \otimes \mathcal{V}) \oplus \Phi_\rho(V \otimes \mathcal{V}) \oplus \rho(g \otimes V) = \{0\}$, it follows that $\dim \mathcal{V} = \dim g = \dim V = 0$ from the assumption that $(g, \rho, V, \mathcal{V}, B_0)$ is irreducible. In particular, we have $\varrho(g \otimes \mathcal{V}) = \mathcal{V} = \{0\}$ and $\rho(g \otimes V) = V = \{0\}$. If $\varrho(g \otimes \mathcal{V}) \oplus \Phi_\rho(V \otimes \mathcal{V}) \oplus \rho(g \otimes V) \neq \{0\}$, since it satisfies the conditions (2.36) and (2.37), we have $\varrho(g \otimes \mathcal{V}) \oplus \Phi_\rho(V \otimes \mathcal{V}) \oplus \rho(g \otimes V) = \mathcal{V} \oplus g \oplus V$. \square

Proposition 2.32. Let $(g, \rho, V, \mathcal{V}, B_0)$ be an irreducible standard pentad whose representation ρ is faithful and denote the Lie algebra associated with it by $L(g, \rho, V, \mathcal{V}, B_0) = \bigoplus_{n \in \mathbb{Z}} V_n$. Let N (respectively M) be an integer such that V_{N+1} is not $\{0\}$ (respectively V_{-M-1} is not $\{0\}$). Then for any non-zero element $z_N \in V_N$ (respectively $\omega_{-M} \in V_{-M}$), there exists an element $x_1 \in V_1$ such that $[x_1, z_N] \neq 0$ (respectively $y_{-1} \in V_{-1}$ such that $[y_{-1}, \omega_{-M}] \neq 0$) (cf. [8, Proposition 3.11]).

Proof. When $N \leq -1$, we have our claim by Propositions 2.11, 2.17 and 2.31. When $N = 0$, we have our claim by the assumption that ρ is faithful. Assume that $N \geq 1$, $V_{N+1} \neq \{0\}$ and put $a_N := \{a_N \in V_N \mid [x_1, a_N] = 0 \text{ for any } x_1 \in V_1\}$ and $a_n := \{a_n \in V_n \mid [x_1, a_n] \in a_{n+1} \text{ for any } x_1 \in V_1\}$ for $n \leq N-1$ inductively. Then a_n is a V_0 -submodule of V_n for each n , i.e. $[V_0, a_n] \subset a_n$, and, we have that $[V_{\pm 1}, a_n] \subset a_{n \pm 1}$ for any $n \in \mathbb{Z}$ (see [8, the proof of Proposition 3.11]). In particular, $a_{-1} \oplus a_0 \oplus a_1$ satisfies the conditions (2.36) and (2.37). If $a_{-1} \oplus a_0 \oplus a_1 = \mathcal{V} \oplus g \oplus V$, then we have $a_N = V_N$ and a contradiction to the assumption that $V_{N+1} \neq \{0\}$. Thus we have $a_1 = \{0\}$, and, thus, $a_2 = \{0\}, \dots, a_N = \{0\}$ by the transitivity of $L(g, \rho, V, \mathcal{V}, B_0)$. Similarly, we have our result for M such that $V_{-M-1} \neq \{0\}$. \square

Proposition 2.33. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be an irreducible standard pentad whose representation ρ is faithful. If the Lie algebra $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is finite-dimensional, then $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is simple (cf. [8, Proposition 3.12]). Moreover, if $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is defined over \mathbb{C} and $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is a finite-dimensional simple Lie algebra, then a triplet (\mathfrak{g}, ρ, V) corresponds to some prehomogeneous vector space of parabolic type (see [8, Theorem 3.13]).*

Proof. We can show this by Proposition 2.32 and the same argument to the argument of [8, Proposition 3.12 and Theorem 3.13]. \square

A prehomogeneous vector space of parabolic type (abbrev. a PV of parabolic type) is a PV which can be obtained from a \mathbb{Z} -graded finite-dimensional semisimple Lie algebra. PVs of parabolic type are classified by H. Rubenthaler (see [4, 5, 6]).

EXAMPLE 2.34. Let $m \geq 2$ and $\mathfrak{g} = \mathfrak{gl}_1(\mathbb{C}) \oplus \mathfrak{sl}_m(\mathbb{C})$, $\rho = \Lambda_1$ a representation of \mathfrak{g} on \mathbb{C}^m defined by

$$\Lambda_1((a, A) \otimes v) := av + Av \quad (a \in \mathfrak{gl}_1, A \in \mathfrak{sl}_m, v \in V),$$

$B_0 = \kappa_m$ a bilinear form on \mathfrak{g} defined by

$$\kappa_m((a, A), (a', A')) := \frac{m}{m+1}aa' + \text{Tr}(AA') \quad (a, a' \in \mathfrak{gl}_1, A, A' \in \mathfrak{sl}_m).$$

Then, a pentad $(\mathfrak{g}, \rho, V, \text{Hom}(V, \mathbb{C}), B_0) = (\mathfrak{gl}_1 \oplus \mathfrak{sl}_m, \Lambda_1, \mathbb{C}^m, \mathbb{C}^m, \kappa_m)$ is a standard pentad which has a $(m^2 + 2m)$ -dimensional graded simple Lie algebra $L(\mathfrak{gl}_1 \oplus \mathfrak{sl}_m, \Lambda_1, \mathbb{C}^m, \mathbb{C}^m, \kappa_m) = V_{-1} \oplus V_0 \oplus V_1$ (see [8, Example 1.14]). This Lie algebra $L(\mathfrak{gl}_1 \oplus \mathfrak{sl}_m, \Lambda_1, \mathbb{C}^m, \mathbb{C}^m, \kappa_m)$ is isomorphic to \mathfrak{sl}_{m+1} . Indeed, from the classification of PVs of parabolic type (see [4, 5, 6]) and the dimension of $L(\mathfrak{g}, \rho, V, \text{Hom}(V, \mathbb{C}), B_0)$, it is isomorphic to \mathfrak{sl}_{m+1} .

EXAMPLE 2.35. Put $\mathfrak{g} := \mathfrak{gl}_1(\mathbb{C}) \oplus \mathfrak{gl}_1(\mathbb{C}) \oplus \mathfrak{sl}_2(\mathbb{C})$, $V := \mathbb{C}^2 = M(2, 1; \mathbb{C})$, $\mathcal{V} := \mathbb{C}^2$ and define representations $\rho : \mathfrak{g} \otimes V \rightarrow V$, $\varrho : \mathfrak{g} \otimes \mathcal{V} \rightarrow \mathcal{V}$ by:

$$\rho((a, b, A) \otimes v) := bv + Av, \quad \varrho((a, b, A) \otimes \phi) := -b\phi - {}^tA\phi$$

for any $(a, b, A) \in \mathfrak{g}$, $v \in V$, $\phi \in \mathcal{V}$. We can identify \mathcal{V} with $\text{Hom}(V, \mathbb{C})$ via the following bilinear map $\langle \cdot, \cdot \rangle_V : V \times \mathcal{V} \rightarrow \mathbb{C}$ defined by:

$$\langle v, \phi \rangle_V := {}^tv\phi.$$

Let B_0 be a bilinear form on \mathfrak{g} defined by:

$$B_0((a, b, A), (a', b', A')) := \frac{3}{4}aa' + bb' + \frac{1}{2}(ab' + a'b) + \text{Tr}(AA').$$

Then, a pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is a standard pentad whose Φ -map is given by:

$$\Phi_\rho(v \otimes \phi) = (-{}^tv\phi, \frac{3}{2}{}^tv\phi, v'\phi - \frac{1}{2}{}^tv\phi I_2).$$

The Lie algebra $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is isomorphic to $\mathfrak{gl}_1 \oplus \mathfrak{sl}_3$. Indeed, if we put $\mathfrak{g}_V^1 := \mathbb{C} \cdot (1, 0, O_2)$, $\mathfrak{g}_V^2 := \mathbb{C} \cdot (-\frac{2}{3}, 1, O_2) \oplus \mathfrak{sl}_2$, then we have

$$\begin{aligned} L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) &\simeq L((\mathfrak{g}_V^1, \rho|_{\mathfrak{g}_V^1}, \{0\}, \{0\}, B_0|_{\mathfrak{g}_V^1 \times \mathfrak{g}_V^1}) \oplus (\mathfrak{g}_V^2, \rho|_{\mathfrak{g}_V^2}, V, \mathcal{V}, B_0|_{\mathfrak{g}_V^2 \times \mathfrak{g}_V^2})) \\ &\simeq \mathfrak{g}_V^1 \oplus L(\mathfrak{g}_V^2, \rho|_{\mathfrak{g}_V^2}, V, \mathcal{V}, B_0|_{\mathfrak{g}_V^2 \times \mathfrak{g}_V^2}) \simeq \mathfrak{g}_V^1 \oplus \mathcal{V} \oplus \mathfrak{g}_V^2 \oplus V \\ &\simeq \mathfrak{gl}_1 \oplus \mathfrak{sl}_3 \end{aligned}$$

from Example 2.34. Moreover, under this identification, the bilinear form B_L on $L(\mathfrak{g}_V^2, \rho|_{\mathfrak{g}_V^2}, V, \mathcal{V}, B_0|_{\mathfrak{g}_V^2 \times \mathfrak{g}_V^2})$ is given by $B_L(\hat{A}, \hat{A}') = \text{Tr}(\hat{A}\hat{A}')$ ($\hat{A}, \hat{A}' \in \mathfrak{sl}_3$). In fact, if we put

$$h := (0, 0, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}) \in \mathfrak{g}_V^2,$$

then $B_0(h, h) = 2$. On the other hand, we can obtain $\text{Tr}(\text{ad } h \text{ ad } h) = 12$, where ad stands for the adjoint representation of $L(\mathfrak{g}_V^2, \rho|_{\mathfrak{g}_V^2}, V, \mathcal{V}, B_0|_{\mathfrak{g}_V^2 \times \mathfrak{g}_V^2})$, by a direct calculation. Since any non-degenerate invariant bilinear form on \mathfrak{sl}_3 is a scalar multiple of the Killing form, we can obtain that B_L is 1/6 times the Killing form of \mathfrak{sl}_3 , i.e. $B_L(\hat{A}, \hat{A}') = \text{Tr}(\hat{A}\hat{A}')$.

Proposition 2.36. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad whose representation ρ is faithful. Under this assumption, the pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is irreducible if and only if the Lie algebra $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ does not have a non-zero proper graded ideal.*

Proof. Assume that $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is reducible. Under the notation of Definition 2.29, we put $\mathfrak{a}_{-1} := \mathcal{V}_{\mathfrak{a}}$, $\mathfrak{a}_0 := \mathfrak{a}$, $\mathfrak{a}_1 := \mathcal{V}_{\mathfrak{a}}$. Moreover, we put $\mathfrak{a}_n := [V_1, \mathfrak{a}_{n-1}]$ for all $n \geq 2$ and $\mathfrak{a}_m := [V_{-1}, \mathfrak{a}_{m+1}]$ for all $m \leq -2$ inductively. Then a direct sum $\mathfrak{A} := \bigoplus_{n \in \mathbb{Z}} \mathfrak{a}_n$ is a non-zero proper graded ideal of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. In fact, by the assumption that $[V_i, \mathfrak{a}_j] \subset \mathfrak{a}_{i+j}$ for any $-1 \leq i, j, i+j \leq 1$, we can easily show that $[V_0, \mathfrak{A}], [V_{\pm 1}, \mathfrak{A}] \subset \mathfrak{A}$ by induction. Since $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is generated by V_0 and $V_{\pm 1}$, we have $[L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \mathfrak{A}] \subset \mathfrak{A}$. Thus, \mathfrak{A} is a graded ideal. Since $\{0\} \neq \mathfrak{a}_{-1} \oplus \mathfrak{a}_0 \oplus \mathfrak{a}_1 \subsetneq \mathcal{V} \oplus \mathfrak{g} \oplus V$, we have $\{0\} \neq \mathfrak{A} \subsetneq L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$.

Conversely, assume that $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is irreducible. Let $\mathfrak{b} = \sum_{n \in \mathbb{Z}} (\mathfrak{b} \cap V_n)$ be a non-zero graded ideal of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and put $\mathfrak{b}_n := \mathfrak{b} \cap V_n$. Then, by Proposition 2.32, we can obtain that $\mathfrak{b}_0 \neq \{0\}$. In fact, since $\mathfrak{b} \neq \{0\}$, there exists an integer $n \in \mathbb{Z}$ and a non-zero element $z_n \in \mathfrak{b}_n$. For example, if $n \geq 1$, then there exist n elements $y_{-1}^1, \dots, y_{-1}^n \in V_{-1}$ such that $[y_{-1}^n, [\dots, [y_{-1}^1, z_n] \dots]] \in \mathfrak{b}_0 \setminus \{0\}$. Since $\mathfrak{b}_{-1} \oplus \mathfrak{b}_0 \oplus \mathfrak{b}_1$ satisfies the conditions (2.36) and (2.37), it coincides with $V_{-1} \oplus V_0 \oplus V_1$, and, thus, $\mathfrak{b} = L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. \square

The following lemmas are to construct a derivation on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. They are used in Theorem 3.26.

Lemma 2.37. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad, $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) = \bigoplus_{n \in \mathbb{Z}} V_n$ be the Lie algebra associated with it. Let $\alpha_i : V_i \rightarrow V_i$ ($i = 0, \pm 1$) be linear maps which satisfy*

$$(2.38) \quad \alpha_{i+j}([a_i, b_j]) = [\alpha_i(a_i), b_j] + [a_i, \alpha_j(b_j)]$$

for any $-1 \leq i, j, i+j \leq 1$ and elements $a_i \in V_i$, $b_j \in V_j$. Then, there exists a linear map $\alpha : L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) \rightarrow L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ such that α is a derivation on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and its restriction to V_i ($i = 0, \pm 1$) coincides with α_i .

Proof. First, let us construct linear maps $\alpha_i : V_i \rightarrow V_i$ for all $i \in \mathbb{Z}$ by induction. Let $i \geq 1$ and assume that the integer i satisfies the condition that we have linear maps $\alpha_j : V_j \rightarrow V_j$ for all $0 \leq j \leq i$ which satisfy the following equations:

$$\begin{aligned} \alpha_j([a_0, b_j]) &= [\alpha_0(a_0), b_j] + [a_0, \alpha_j(b_j)], \\ \alpha_j([x_1, b_{j-1}]) &= [\alpha_1(x_1), b_{j-1}] + [x_1, \alpha_{j-1}(b_{j-1})], \\ \alpha_{j-1}([y_{-1}, b_j]) &= [\alpha_{-1}(y_{-1}), b_j] + [y_{-1}, \alpha_j(b_j)] \end{aligned}$$

for any $0 \leq j \leq i$, $a_0 \in V_0$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $b_j \in V_j$ and $b_{j-1} \in V_{j-1}$. By the assumption

(2.38), when $i = 1$ the given linear maps $\alpha_0, \alpha_{\pm 1}$ satisfy these equations. Then we define a linear map $\alpha_{i+1} : V_{i+1} \rightarrow V_{i+1}$ by:

$$(2.39) \quad \alpha_{i+1}([x_1, b_i]) := [\alpha_1(x_1), b_i] + [x_1, \alpha_i(b_i)]$$

for any $x_1 \in V_1$ and $b_i \in V_i$. Let us check the well-definedness of α_{i+1} . In fact, for any $y_{-1} \in V_{-1}$, $x_1 \in V_1$ and $b_i \in V_i$, we have

$$(2.40) \quad \begin{aligned} [y_{-1}, [\alpha_1(x_1), b_i] + [x_1, \alpha_i(b_i)]] &= [y_{-1}, [\alpha_1(x_1), b_i]] + [y_{-1}, [x_1, \alpha_i(b_i)]] \\ &= [[y_{-1}, \alpha_1(x_1)], b_i] + [\alpha_1(x_1), [y_{-1}, b_i]] + [[y_{-1}, x_1], \alpha_i(b_i)] + [x_1, [y_{-1}, \alpha_i(b_i)]] \\ &= [\alpha_0([y_{-1}, x_1]), b_i] - [[\alpha_{-1}(y_{-1}), x_1], b_i] + [\alpha_1(x_1), [y_{-1}, b_i]] \\ &\quad + [[y_{-1}, x_1], \alpha_i(b_i)] + [x_1, \alpha_{i-1}([y_{-1}, b_i])] - [x_1, [\alpha_{-1}(y_{-1}), b_i]] \\ &= \alpha_i([y_{-1}, x_1], b_i) + \alpha_i([x_1, [y_{-1}, b_i]]) - [\alpha_{-1}(y_{-1}), [x_1, b_i]] \\ &= \alpha_i([y_{-1}, [x_1, b_i]]) - [\alpha_{-1}(y_{-1}), [x_1, b_i]]. \end{aligned}$$

Thus, if $x_1^1, \dots, x_1^l \in V_1$ and $b_i^1, \dots, b_i^l \in V_i$ satisfy $\sum_{s=1}^l [x_1^s, b_i^s] = 0$, then we have

$$\sum_{s=1}^l [y_{-1}, [\alpha_1(x_1^s), b_i^s] + [x_1^s, \alpha_i(b_i^s)]] = 0$$

for any $y_{-1} \in V_{-1}$. Therefore, we have $\sum_{s=1}^l ([\alpha_1(x_1^s), b_i^s] + [x_1^s, \alpha_i(b_i^s)]) = 0$ and the well-definedness of α_{i+1} . Moreover, α_{i+1} satisfies the following equations:

$$(2.41) \quad \alpha_{i+1}([a_0, b_{i+1}]) = [\alpha_0(a_0), b_{i+1}] + [a_0, \alpha_{i+1}(b_{i+1})],$$

$$(2.42) \quad \alpha_{i+1}([x_1, b_i]) = [\alpha_1(x_1), b_i] + [x_1, \alpha_i(b_i)],$$

$$(2.43) \quad \alpha_i([y_{-1}, b_{i+1}]) = [\alpha_{-1}(y_{-1}), b_{i+1}] + [y_{-1}, \alpha_{i+1}(b_{i+1})]$$

for any $a_0 \in V_0$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $b_i \in V_i$ and $b_{i+1} \in V_{i+1}$. In fact, for any $a_0 \in V_0$, $x_1 \in V_1$, and $b_i \in V_i$, we have

$$(2.44) \quad \begin{aligned} \alpha_{i+1}([a_0, [x_1, b_i]]) &= \alpha_{i+1}([a_0, x_1], b_i) + \alpha_{i+1}([x_1, [a_0, b_i]]) \\ &= [\alpha_1([a_0, x_1]), b_i] + [[a_0, x_1], \alpha_i(b_i)] + [\alpha_1(x_1), [a_0, b_i]] + [x_1, \alpha_i([a_0, b_i])] \\ &= [[\alpha_0(a_0), x_1], b_i] + [[a_0, \alpha_1(x_1)], b_i] + [[a_0, x_1], \alpha_i(b_i)] \\ &\quad + [\alpha_1(x_1), [a_0, b_i]] + [x_1, [\alpha_0(a_0), b_i]] + [x_1, [a_0, \alpha_i(b_i)]] \\ &= [\alpha_0(a_0), [x_1, b_i]] + [a_0, [\alpha_1(x_1), b_i]] + [a_0, [x_1, \alpha_i(b_i)]] \\ &= [\alpha_0(a_0), [x_1, b_i]] + [a_0, \alpha_{i+1}([x_1, b_i])]. \end{aligned}$$

Thus, we can obtain the equation (2.41). The equation (2.42) is clear. The equation (2.43) follows from (2.40). Thus, inductively, we can obtain linear maps α_i for all $i \geq 0$, and, similarly, we can construct linear maps $\alpha_{-i} : V_{-i} \rightarrow V_{-i}$ for all $i \geq 0$. Consequently, we have linear maps $\alpha_n : V_n \rightarrow V_n$ for all $n \in \mathbb{Z}$ which satisfy

$$(2.45) \quad \alpha_n([a_0, b_n]) = [\alpha_0(a_0), b_n] + [a_0, \alpha_n(b_n)],$$

$$(2.46) \quad \alpha_{n+1}([x_1, b_n]) = [\alpha_1(x_1), b_n] + [x_1, \alpha_n(b_n)],$$

$$(2.47) \quad \alpha_{n-1}([y_{-1}, b_n]) = [\alpha_{-1}(y_{-1}), b_n] + [y_{-1}, \alpha_n(b_n)]$$

for any $a_0 \in V_0$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$ and $b_n \in V_n$.

We define a linear map $\alpha : L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) \rightarrow L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ by:

$$(2.48) \quad \alpha(a_n) := \alpha_n(a_n)$$

for any $n \in \mathbb{Z}$ and $a_n \in V_n$. Then α is a derivation of Lie algebras. In fact, we can show the following equation

$$(2.49) \quad \alpha([a_n, b_m]) = [\alpha(a_n), b_m] + [a_n, \alpha(b_m)]$$

for any $n, m \in \mathbb{Z}$, $a_n \in V_n$ and $b_m \in V_m$ by the equations (2.45), (2.46), (2.47) inductively. \square

Lemma 2.38. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad and α be a derivation on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. If α satisfies the equation*

$$(2.50) \quad B_L(\alpha(z), \omega) = -B_L(z, \alpha(\omega))$$

for any $z = z_n \in V_n$ ($n = 0, \pm 1$) and $\omega \in L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, then we have the same equation for any $z, \omega \in L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$.

Proof. We argue our claim in the cases where $z = z_n \in V_n$ for some n and prove it by induction on n . Suppose that $n \geq 0$. If $n = 0, 1$, then our claim follows from the assumption. Suppose that $n \geq 2$. Then, by the induction hypothesis, we have

$$\begin{aligned} B_L(\alpha([x_1, z_{n-1}]), \omega) &= B_L([\alpha(x_1), z_{n-1}], \omega) + B_L([x_1, \alpha(z_{n-1})], \omega) \\ &= -B_L(z_{n-1}, [\alpha(x_1), \omega]) - B_L(\alpha(z_{n-1}), [x_1, \omega]) \\ &= -B_L(z_{n-1}, [\alpha(x_1), \omega]) + B_L(z_{n-1}, \alpha([x_1, \omega])) \\ &= B_L(z_{n-1}, [x_1, \alpha(\omega)]) \\ &= -B_L([x_1, z_{n-1}], \alpha(\omega)) \end{aligned}$$

for any $x_1 \in V_1$, $z_{n-1} \in V_{n-1}$. Since $V_n = [V_1, V_{n-1}]$, we have our claim for n . Thus, by induction, we have our claim for all $n \geq 0$. Similarly, we can show our claim for $n \leq -1$. \square

3. Graded modules of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$

3.1. A construction of vector spaces \tilde{U}^+ and \tilde{U}^- . As mentioned in section 1, the purpose of this and the next section is to construct a positively graded module and a negatively graded module of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ from a given \mathfrak{g} -module U , which will be denoted by \tilde{U}^+ and \tilde{U}^- . First, we construct \tilde{U}^+ and \tilde{U}^- as vector spaces by induction.

DEFINITION 3.1. Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad and $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) = \bigoplus_{n \in \mathbb{Z}} V_n$ be the Lie algebra associated with it. Let $\pi : \mathfrak{g} \otimes U \rightarrow U$ be a representation of $\mathfrak{g} = V_0$ on a vector space U over F . We put $U_0^+ = U_0^- := U$, $\pi_0^+ = \pi_0^- := \pi$ and define linear maps $r_0^+ : V_1 \otimes U_0^+ \rightarrow \text{Hom}(V_{-1}, U_0^+)$ and $r_0^- : V_{-1} \otimes U_0^- \rightarrow \text{Hom}(V_1, U_0^-)$ by:

$$(3.1) \quad \begin{aligned} r_0^+ : V_1 \otimes U_0^+ &\rightarrow \text{Hom}(V_{-1}, U_0^+) \\ x_1 \otimes u_0 &\mapsto (\eta_{-1} \mapsto \pi_0^+([\eta_{-1}, x_1] \otimes u_0)), \end{aligned}$$

$$(3.2) \quad \begin{aligned} r_0^- : V_{-1} \otimes U_0^- &\rightarrow \text{Hom}(V_1, U_0^-) \\ y_{-1} \otimes u_0 &\mapsto (\xi_1 \mapsto \pi_0^-([\xi_1, y_{-1}] \otimes u_0)). \end{aligned}$$

Proposition 3.2. *The maps r_0^+ and r_0^- are homomorphisms of \mathfrak{g} -modules.*

Proof. We prove for r_0^+ . For any elements $a \in \mathfrak{g}$, $x_1 \in V_1$, $\eta_{-1} \in V_{-1}$ and $u_0 \in U_0^+$, we have

$$\begin{aligned} r_0^+([a, x_1] \otimes u_0 + x_1 \otimes \pi_0^+(a \otimes u_0))(\eta_{-1}) &= \pi_0^+([\eta_{-1}, [a, x_1]] \otimes u_0) + \pi_0^+([\eta_{-1}, x_1] \otimes \pi_0^+(a \otimes u_0)) \\ &= \pi_0^+([a, [\eta_{-1}, x_1]] \otimes u_0) - \pi_0^+([[a, \eta_{-1}], x_1] \otimes u_0) + \pi_0^+([\eta_{-1}, x_1] \otimes \pi_0^+(a \otimes u_0)) \\ &= \pi_0^+(a \otimes \pi_0^+([\eta_{-1}, x_1] \otimes u_0)) - \pi_0^+([[a, \eta_{-1}], x_1] \otimes u_0) \\ &= \pi_0^+(a \otimes (r_0^+(x_1 \otimes u_0)(\eta_{-1}))) - r_0^+(x_1 \otimes u_0)([a, \eta_{-1}]). \end{aligned}$$

Thus r_0^+ is a homomorphism of \mathfrak{g} -modules. Similarly, we can prove that r_0^- is a homomorphism of \mathfrak{g} -modules. \square

It follows from Proposition 3.2 that the linear spaces $U_1^+ := \text{Im } r_0^+$ and $U_{-1}^- := \text{Im } r_0^-$ have the canonical \mathfrak{g} -module structures. We denote these canonical representations by π_1^+ and π_{-1}^- respectively. Moreover, we inductively construct \mathfrak{g} -modules U_2^+, U_3^+, \dots by using the following proposition.

Proposition 3.3. *Assume that there exist \mathfrak{g} -modules (ϖ^+, W^+) , (ϖ^-, W^-) and \mathfrak{g} -module homomorphisms $\lambda^+ : V_1 \otimes W^+ \rightarrow \text{Hom}(V_{-1}, W^+)$ and $\lambda^- : V_{-1} \otimes W^- \rightarrow \text{Hom}(V_1, W^-)$. We put $\hat{W}^+ := \text{Im } \lambda^+$, $\hat{W}^- := \text{Im } \lambda^-$ and denote the canonical representations of \mathfrak{g} on them by $\hat{\varpi}^+$ and $\hat{\varpi}^-$ respectively. Then the following linear maps are \mathfrak{g} -module homomorphisms:*

$$(3.3) \quad \begin{aligned} \hat{\lambda}^+ : V_1 \otimes \hat{W}^+ &\rightarrow \text{Hom}(V_{-1}, \hat{W}^+) \\ x_1 \otimes \hat{w}^+ &\mapsto (\eta_{-1} \mapsto \hat{\varpi}^+([\eta_{-1}, x_1] \otimes \hat{w}^+) + \lambda^+(x_1 \otimes \hat{w}^+(\eta_{-1}))), \end{aligned}$$

$$(3.4) \quad \begin{aligned} \hat{\lambda}^- : V_{-1} \otimes \hat{W}^- &\rightarrow \text{Hom}(V_1, \hat{W}^-) \\ y_{-1} \otimes \hat{w}^- &\mapsto (\xi_1 \mapsto \hat{\varpi}^-([\xi_1, y_{-1}] \otimes \hat{w}^-) + \lambda^-(y_{-1} \otimes \hat{w}^-(\xi_1))). \end{aligned}$$

Proof. We can prove it by a similar argument to the argument of [8, Proposition 1.10]. Take any elements $a \in \mathfrak{g}$, $x_1 \in V_1$, $\eta_{-1} \in V_{-1}$ and $\hat{w}^+ \in \hat{W}^+$. Then we have

$$\begin{aligned} &(\hat{\lambda}^+([a, x_1] \otimes \hat{w}^+) + \hat{\lambda}^+(x_1 \otimes \hat{\varpi}^+(a \otimes \hat{w}^+)))(\eta_{-1}) \\ &= \hat{\varpi}^+([\eta_{-1}, [a, x_1]] \otimes \hat{w}^+) + \lambda^+([a, x_1] \otimes \hat{w}^+(\eta_{-1})) \\ &\quad + \hat{\varpi}^+([\eta_{-1}, x_1] \otimes \hat{\varpi}^+(a \otimes \hat{w}^+)) + \lambda^+(x_1 \otimes (\hat{\varpi}^+(a \otimes \hat{w}^+)(\eta_{-1}))) \\ &= \hat{\varpi}^+([a, [\eta_{-1}, x_1]] \otimes \hat{w}^+) + \hat{\varpi}^+([\eta_{-1}, x_1] \otimes \hat{\varpi}^+(a \otimes \hat{w}^+)) + \lambda^+([a, x_1] \otimes \hat{w}^+(\eta_{-1})) \\ &\quad + \lambda^+(x_1 \otimes \hat{\varpi}^+(a \otimes \hat{w}^+(\eta_{-1}))) - \hat{\varpi}^+([[a, \eta_{-1}], x_1] \otimes \hat{w}^+) - \lambda^+(x_1 \otimes \hat{w}^+([a, \eta_{-1}])) \\ &= \hat{\varpi}^+(a \otimes \hat{\varpi}^+([\eta_{-1}, x_1] \otimes \hat{w}^+)) + \hat{\varpi}^+(a \otimes \lambda^+(x_1 \otimes \hat{w}^+(\eta_{-1}))) \\ &\quad - \hat{\varpi}^+([[a, \eta_{-1}], x_1] \otimes \hat{w}^+) - \lambda^+(x_1 \otimes \hat{w}^+([a, \eta_{-1}])) \\ &= \hat{\varpi}^+(a \otimes \hat{\lambda}^+(x_1 \otimes \hat{w}^+)(\eta_{-1})) - \hat{\lambda}^+(x_1 \otimes \hat{w}^+)([a, \eta_{-1}]). \end{aligned}$$

Thus, $\hat{\lambda}^+$ is a homomorphism of \mathfrak{g} -modules. By the same way, we can prove that $\hat{\lambda}^-$ is also a \mathfrak{g} -module homomorphism. \square

DEFINITION 3.4. Suppose that $j \geq 1$ and there exist \mathfrak{g} -modules (π_{j-1}^+, U_{j-1}^+) , $(\pi_{-j+1}^-, U_{-j+1}^-)$ and homomorphisms of \mathfrak{g} -modules $r_{j-1}^+ : V_1 \otimes U_{j-1}^+ \rightarrow \text{Hom}(V_{-1}, U_{j-1}^+)$ and $r_{-j+1}^- : V_{-1} \otimes$

$U_{-j+1}^- \rightarrow \text{Hom}(V_1, U_{-j+1}^-)$. Put $U_j^+ := \text{Im } r_{j-1}^+$ and $U_j^- := \text{Im } r_{-j+1}^-$. Then we define linear maps r_j^+ and r_j^- by:

$$(3.5) \quad r_j^+ : V_1 \otimes U_j^+ \rightarrow \text{Hom}(V_1, U_j^+)$$

$$x_1 \otimes u_j^+ \mapsto (\eta_1 \mapsto \pi_j^+([\eta_{-1}, x_1] \otimes u_j^+) + r_{j-1}^+(x_1 \otimes u_j^+(\eta_{-1}))),$$

$$(3.6) \quad r_j^- : V_{-1} \otimes U_j^- \rightarrow \text{Hom}(V_1, U_j^-)$$

$$y_{-1} \otimes u_j^- \mapsto (\xi_1 \mapsto \pi_j^-([\xi_1, y_{-1}] \otimes u_j^-) + r_{-j+1}^-(y_{-1} \otimes u_j^-(\xi_1))).$$

Then, by Proposition 3.3, r_j^+ and r_j^- are homomorphisms of \mathfrak{g} -modules. We denote by U_{j+1}^+ and U_{-j-1}^- the images of r_j^+ and r_j^- and the canonical representations of \mathfrak{g} on U_{j+1}^+ and U_{-j-1}^- by π_{j+1}^+ and π_{-j-1}^- respectively. Moreover, we put

$$U_{-j}^+ := \{0\}, \quad U_j^- := \{0\}$$

for $j \geq 1$. We denote the zero representations of \mathfrak{g} on U_{-j}^+ and U_j^- by π_{-j}^+ and π_j^- for all $j \geq 1$. Thus, inductively, we obtain \mathfrak{g} -modules (π_m^+, U_m^+) , (π_m^-, U_m^-) for all $m \in \mathbb{Z}$. Under these notation, we define linear spaces \tilde{U}^+ and \tilde{U}^- by:

$$(3.7) \quad \tilde{U}^+ := \bigoplus_{m \in \mathbb{Z}} U_m^+, \quad \tilde{U}^- := \bigoplus_{m \in \mathbb{Z}} U_m^-.$$

Throughout this paper, we use these notation.

3.2. A construction of representations of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ on \tilde{U}^+ and \tilde{U}^- . In this section, we define $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module structures on vector spaces \tilde{U}^+ and \tilde{U}^- constructed in (3.7). For this, we start with the following definition.

DEFINITION 3.5. We define the following linear maps:

$$\pi_{0,m}^+ : V_0 \otimes U_m^+ \rightarrow U_m^+, \quad \pi_{1,m}^+ : V_1 \otimes U_m^+ \rightarrow U_{m+1}^+, \quad \pi_{-1,m}^+ : V_{-1} \otimes U_m^+ \rightarrow U_{m-1}^+$$

by:

$$(3.8) \quad \pi_{0,m}^+(a \otimes u_m^+) := \pi_m^+(a \otimes u_m^+) \quad (m \in \mathbb{Z}),$$

$$(3.9) \quad \pi_{1,m}^+(x_1 \otimes u_m^+) := \begin{cases} r_m^+(x_1 \otimes u_m^+) & (m \geq 0) \\ 0 & (m \leq -1) \end{cases},$$

$$(3.10) \quad \pi_{-1,m}^+(y_{-1} \otimes u_m^+) := \begin{cases} u_m^+(y_{-1}) & (m \geq 1) \\ 0 & (m \leq 0) \end{cases}$$

where $a \in V_0$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$ and $u_m^+ \in U_m^+$.

By the above definition, we can obtain the following proposition.

Proposition 3.6. *Under the above notation, we have the following equations:*

$$(3.11) \quad \pi_{1,m}^+([x_1, a] \otimes u_m^+) = \pi_{1,m}^+(x_1 \otimes \pi_{0,m}^+(a \otimes u_m^+)) - \pi_{0,m+1}^+(a \otimes \pi_{1,m}^+(x_1 \otimes u_m^+)),$$

$$(3.12) \quad \pi_{-1,m}^+([y_{-1}, a] \otimes u_m^+) = \pi_{-1,m}^+(y_{-1} \otimes \pi_{0,m}^+(a \otimes u_m^+)) - \pi_{0,m-1}^+(a \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+)),$$

$$(3.13) \quad \pi_{1,m-1}^+(x_1 \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+)) = \pi_{0,m}^+([x_1, y_{-1}] \otimes u_m) + \pi_{-1,m+1}^+(y_{-1} \otimes \pi_{1,m}^+(x_1 \otimes u_m)).$$

Proof. Let us show (3.13). The equations (3.11) and (3.12) can be shown similarly. If $m \leq -1$, then (3.13) is clear. If $m = 0$, then the left hand side equals to 0. For the right hand

side, we have

$$\begin{aligned} & \pi_{0,0}^+([x_1, y_{-1}] \otimes u_0) + \pi_{-1,1}^+(y_{-1} \otimes \pi_{1,0}^+(x_1 \otimes u_0)) \\ &= \pi_0^+([x_1, y_{-1}] \otimes u_0^+) + r_0^+(x_1 \otimes u_0^+)(y_{-1}) = \pi_0^+([x_1, y_{-1}] \otimes u_0^+) + \pi_0^+([y_{-1}, x_1] \otimes u_0^+) = 0. \end{aligned}$$

Thus we have (3.13) when $m = 0$. For $m \geq 1$, the equation (3.13) follows from definition. \square

DEFINITION 3.7. We define the following linear maps for $i \geq 1$ inductively:

$$(3.14) \quad \pi_{i+1,m}^+ : V_{i+1} \otimes U_m^+ \rightarrow U_{i+m+1}^+$$

$$p_i(x_1 \otimes z_i) \otimes u_m^+ \mapsto \pi_{1,i+m}^+(x_1 \otimes \pi_{i,m}^+(z_i \otimes u_m^+)) - \pi_{i,m+1}^+(z_i \otimes \pi_{1,m}^+(x_1 \otimes u_m^+)),$$

$$(3.15)$$

$$\pi_{-i-1,m}^+ : V_{-i-1} \otimes U_m^+ \rightarrow U_{-i+m-1}^+$$

$$\begin{aligned} q_{-i}(y_{-1} \otimes \omega_{-i}) \otimes u_m^+ &\mapsto \pi_{-1,-i+m}^+(y_{-1} \otimes \pi_{-i,m}^+(\omega_{-i} \otimes u_m^+)) \\ &\quad - \pi_{-i,m-1}^+(\omega_{-i} \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+)). \end{aligned}$$

Note that the linear maps $\pi_{0,m}^+, \pi_{\pm 1,m}^+$ defined in Definition 3.5 satisfy the same equations as (3.14) and (3.15) in the cases where $i = 0$ by Proposition 3.6. For $i \geq 1$, we must show the well-definedness of Definition 3.7. To prove it, let us show the following two propositions.

Proposition 3.8. (The well-definedness of $\pi_{i+1,m}^+$ given in (3.14)) Suppose that $i \geq 0$. Suppose that the linear map $\pi_{i,m}^+$ defined in (3.14) is well-defined for any $m \in \mathbb{Z}$ and satisfies the following equations:

$$(3.16) \quad \pi_{0,i+m}^+(a \otimes \pi_{i,m}^+(z_i \otimes u_m^+)) = \pi_{i,m}^+([a, z_i] \otimes u_m^+) + \pi_{i,m}^+(z_i \otimes \pi_{0,m}^+(a \otimes u_m^+)),$$

$$(3.17) \quad \pi_{i,m-1}^+(z_i \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+)) = \pi_{i-1,m}^+([z_i, y_{-1}] \otimes u_m^+) + \pi_{-1,i+m}^+(y_{-1} \otimes \pi_{i,m}^+(z_i \otimes u_m^+)).$$

If $x_1^1, \dots, x_1^l \in V_1$ and $z_i^1, \dots, z_i^l \in V_i$ satisfy $\sum_{s=1}^l p_i(x_1^s \otimes z_i^s) = 0$, then we have

$$(3.18) \quad \sum_{s=1}^l (\pi_{1,i+m}^+(x_1^s \otimes \pi_{i,m}^+(z_i^s \otimes u_m^+)) - \pi_{i,m+1}^+(z_i^s \otimes \pi_{1,m}^+(x_1^s \otimes u_m^+))) = 0$$

for all $m \in \mathbb{Z}$ and $u_m^+ \in U_m^+$. In particular, we can obtain the well-definedness of the linear map $\pi_{i+1,m}^+$ defined in (3.14) for any $m \in \mathbb{Z}$. Moreover, the linear maps $\pi_{i+1,m}^+$ ($m \in \mathbb{Z}$) satisfy the following equations:

$$(3.19) \quad \begin{aligned} & \pi_{0,i+m+1}^+(a \otimes \pi_{i+1,m}^+(z_{i+1} \otimes u_m^+)) \\ &= \pi_{i+1,m}^+([a, z_{i+1}] \otimes u_m^+) + \pi_{i+1,m}^+(z_{i+1} \otimes \pi_{0,m}^+(a \otimes u_m^+)), \end{aligned}$$

$$(3.20) \quad \begin{aligned} & \pi_{i+1,m-1}^+(z_{i+1} \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+)) \\ &= \pi_{i,m}^+([z_{i+1}, y_{-1}] \otimes u_m^+) + \pi_{-1,i+m+1}^+(y_{-1} \otimes \pi_{i+1,m}^+(z_{i+1} \otimes u_m^+)). \end{aligned}$$

Proof. We argue by induction on i . For $i = 0$, our claim follows from Proposition 3.6. Suppose that $i \geq 1$. We fix i and argue (3.18) by induction on m . First, if $m \leq -1$, then the equation (3.18) is clear. If $m \geq 0$, then we have

(3.21)

$$\begin{aligned}
& \pi_{-1,i+m+1}^+(y_{-1} \otimes \pi_{1,i+m}^+(x_1 \otimes \pi_{i,m}^+(z_i \otimes u_m^+)) - \pi_{i,m+1}^+(z_i \otimes \pi_{1,m}^+(x_1 \otimes u_m^+))) \\
&= \pi_{0,i+m}^+([y_{-1}, x_1] \otimes \pi_{i,m}^+(z_i \otimes u_m^+)) + \pi_{1,i+m-1}^+(x_1 \otimes \pi_{-1,i+m}^+(y_{-1} \otimes \pi_{i,m}^+(z_i \otimes u_m^+))) \\
&\quad - \pi_{i-1,m+1}^+([y_{-1}, z_i] \otimes \pi_{1,m}^+(x_1 \otimes u_m^+)) - \pi_{i,m}^+(z_i \otimes \pi_{-1,m+1}^+(y_{-1} \otimes \pi_{1,m}^+(x_1 \otimes u_m^+))) \\
&= \pi_{0,i+m}^+([y_{-1}, x_1] \otimes \pi_{i,m}^+(z_i \otimes u_m^+)) + \pi_{1,i+m-1}^+(x_1 \otimes \pi_{i-1,m}^+([y_{-1}, z_i] \otimes u_m^+)) \\
&\quad + \pi_{1,i+m-1}^+(x_1 \otimes \pi_{i,m-1}^+(z_i \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+))) - \pi_{i-1,m+1}^+([y_{-1}, z_i] \otimes \pi_{1,m}^+(x_1 \otimes u_m^+)) \\
&\quad - \pi_{i,m}^+(z_i \otimes \pi_{0,m}^+([y_{-1}, x_1] \otimes u_m^+)) - \pi_{i,m}^+(z_i \otimes \pi_{1,m-1}^+(x_1 \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+))) \\
&= \pi_{i,m}^+([y_{-1}, x_1], z_i] \otimes u_m^+) + \pi_{i,m}^+([x_1, [y_{-1}, z_i]] \otimes u_m^+) \\
&\quad + \pi_{1,i+m-1}^+(x_1 \otimes \pi_{i,m-1}^+(z_i \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+))) - \pi_{i,m}^+(z_i \otimes \pi_{1,m-1}^+(x_1 \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+))) \\
&= \pi_{i,m}^+([y_{-1}, [x_1, z_i]] \otimes u_m^+) \\
&\quad + \pi_{1,i+m-1}^+(x_1 \otimes \pi_{i,m-1}^+(z_i \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+))) - \pi_{i,m}^+(z_i \otimes \pi_{1,m-1}^+(x_1 \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+)))
\end{aligned}$$

for any $x_1 \in V_1$, $z_i \in V_i$, $y_{-1} \in V_{-1}$ and $u_m^+ \in U_m^+$. By the induction hypotheses on i and m , if we take elements $x_1^1, \dots, x_1^l \in V_1$ and $z_i^1, \dots, z_i^l \in V_i$ satisfying $\sum_{s=1}^l p_i(x_1^s \otimes z_i^s) = 0$, then we have

(3.22)

$$\sum_{s=1}^l \pi_{i,m}^+([x_1^s, z_i^s], y_{-1}] \otimes u_m^+) = 0 \quad (\text{by the induction hypothesis on } i),$$

(3.23)

$$\begin{aligned}
& \sum_{s=1}^l (\pi_{1,i+m-1}^+(x_1^s \otimes \pi_{i,m-1}^+(z_i^s \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+))) - \pi_{i,m}^+(z_i^s \otimes \pi_{1,m-1}^+(x_1^s \otimes \pi_{-1,m}^+(y_{-1} \otimes u_m^+)))) \\
&= 0 \quad (\text{by the induction hypothesis on } m).
\end{aligned}$$

Thus, we have

$$(3.24) \quad \sum_{s=1}^l \pi_{-1,i+m+1}^+(y_{-1} \otimes \pi_{1,i+m}^+(x_1^s \otimes \pi_{i,m}^+(z_i^s \otimes u_m^+)) - \pi_{i,m+1}^+(z_i^s \otimes \pi_{1,m}^+(x_1^s \otimes u_m^+))) = 0$$

from (3.21). Since $i + m + 1 \geq 1$, we can obtain that

(3.25)

$$\begin{aligned}
& \sum_{s=1}^l (\pi_{1,i+m}^+(x_1^s \otimes \pi_{i,m}^+(z_i^s \otimes r_{m+1}(y_{-1} \otimes u_{m+1}^+))) - \pi_{i,m+1}^+(z_i^s \otimes \pi_{1,m}^+(x_1 \otimes r_{m+1}(y_{-1} \otimes u_{m+1}^+)))) \\
&= 0 \in U_{i+m+1}^+ \subset \text{Hom}(V_{-1}, U_{i+m}^+).
\end{aligned}$$

Therefore we can obtain the well-definedness of the linear map $\pi_m^{i+1} : V_{i+1} \otimes U_m^+ \rightarrow U_{i+m+1}^+$ given in (3.14) for any m .

In order to complete the proof, we must show the equations (3.20) and (3.19). Let us show (3.20). Under the above notation, for any $m \in \mathbb{Z}$, we have

$$\begin{aligned}
& \pi_{0,i+m+1}^+(a \otimes \pi_{i+1,m}^+(p_i(x_1 \otimes z_i) \otimes u_m^+)) \\
&= \pi_{0,i+m+1}^+(a \otimes \pi_{1,i+m}^+(x_1 \otimes \pi_{i,m}^+(z_i \otimes u_m^+))) - \pi_{0,i+m+1}^+(a \otimes \pi_{i,m+1}^+(z_i \otimes \pi_{1,m}^+(x_1 \otimes u_m^+))) \\
&= \pi_{1,i+m}^+([a, x_1] \otimes \pi_{i,m}^+(z_i \otimes u_m^+)) + \pi_{1,i+m}^+(x_1 \otimes \pi_{0,i+m}^+(a \otimes \pi_{i,m}^+(z_i \otimes u_m^+)))
\end{aligned}$$

$$\begin{aligned}
& -\pi_{i,m+1}^+([a, z_i] \otimes \pi_{1,m}^+(x_1 \otimes u_m^+)) - \pi_{i,m+1}^+(z_i \otimes \pi_{0,m+1}^+(a \otimes \pi_{1,m}^+(x_1 \otimes u_m^+))) \\
& = \pi_{1,i+m}^+([a, x_1] \otimes \pi_{i,m}^+(z_i \otimes u_m^+)) + \pi_{1,i+m}^+(x_1 \otimes \pi_{i,m}^+([a, z_i] \otimes u_m^+)) \\
& \quad + \pi_{1,i+m}^+(x_1 \otimes \pi_{i,m}^+(z_i \otimes \pi_{0,m}^+(a \otimes u_m^+))) - \pi_{i,m+1}^+([a, z_i] \otimes \pi_{1,m}^+(x_1 \otimes u_m^+)) \\
& \quad - \pi_{i,m+1}^+(z_i \otimes \pi_{1,m}^+([a, x_1] \otimes u_m^+)) - \pi_{i,m+1}^+(z_i \otimes \pi_{1,m}^+(x_1 \otimes \pi_{0,m}^+(a \otimes u_m^+))) \\
& = \pi_{i+1,m}^+([a, x_1], z_i \otimes u_m^+) + \pi_{i+1,m}^+([x_1, [a, z_i]] \otimes u_m^+) + \pi_{i+1,m}^+([x_1, z_i] \otimes \pi_{0,m}^+(a \otimes u_m^+)) \\
& = \pi_{i+1,m}^+([a, p_i(x_1 \otimes z_i)] \otimes u_m^+) + \pi_{i+1,m}^+(p_i(x_1 \otimes z_i) \otimes \pi_{0,m}^+(a \otimes u_m^+)).
\end{aligned}$$

Thus we have (3.20). The equation (3.19) follows from (3.21). This completes the proof. \square

Proposition 3.9. (The well-definedness of $\pi_{-i-1,m}^+$ given in (3.15)) Suppose that $i \geq 0$. Suppose that the linear map $\pi_{-i,m}^+$ defined in (3.15) is well-defined for any $m \in \mathbb{Z}$ and satisfies the following equations:

(3.26)

$$\pi_{0,-i+m}^+(a \otimes \pi_{-i,m}^+(\omega_{-i} \otimes u_m^+)) = \pi_{-i,m}^+([a, \omega_{-i}] \otimes u_m^+) + \pi_{-i,m}^+(\omega_{-i} \otimes \pi_{0,m}^+(a \otimes u_m^+)),$$

(3.27)

$$\pi_{-i,m+1}^+(\omega_{-i} \otimes \pi_{1,m}^+(x_1 \otimes u_m^+)) = \pi_{-i+1,m}^+([\omega_{-i}, x_1] \otimes u_m^+) + \pi_{-i+1,m}^+(x_1 \otimes \pi_{-i,m}^+(\omega_{-i} \otimes u_m^+)).$$

If $y_{-1}^1, \dots, y_{-1}^l \in V_{-1}$ and $\omega_{-i}^1, \dots, \omega_{-i}^l \in V_{-i}$ satisfy $\sum_{s=1}^l q_{-i}(y_{-1}^s \otimes \omega_{-i}^s) = 0$, then we have

$$(3.28) \quad \sum_{s=1}^l (\pi_{-1,-i+m}^+(y_{-1}^s \otimes \pi_{-i,m}^+(\omega_{-i}^s \otimes u_m^+)) - \pi_{-i,m-1}^+(\omega_{-i}^s \otimes \pi_{-1,m}^+(y_{-1}^s \otimes u_m^+))) = 0$$

for all $m \in \mathbb{Z}$ and $u_m^+ \in U_m^+$. In particular, we can obtain the well-definedness of the linear map $\pi_{-i-1,m}^+$ defined in (3.15) for any $m \in \mathbb{Z}$. Moreover, the maps $\pi_{-i-1,m}^+$ ($m \in \mathbb{Z}$) satisfy the following equations:

(3.29)

$$\begin{aligned}
& \pi_{0,-i+m-1}^+(a \otimes \pi_{-i-1,m}^+(\omega_{-i-1} \otimes u_m^+)) \\
& = \pi_{-i-1,m}^+([a, \omega_{-i-1}] \otimes u_m^+) + \pi_{-i-1,m}^+(\omega_{-i-1} \otimes \pi_{0,m}^+(a \otimes u_m^+)),
\end{aligned}$$

(3.30)

$$\begin{aligned}
& \pi_{-i-1,m+1}^+(\omega_{-i-1} \otimes \pi_{1,m}^+(x_1 \otimes u_m^+)) \\
& = \pi_{-i,m}^+([\omega_{-i-1}, x_1] \otimes u_m^+) + \pi_{-i+1,m-1}^+(x_1 \otimes \pi_{-i-1,m}^+(\omega_{-i-1} \otimes u_m^+)).
\end{aligned}$$

Proof. If $i = 0$, then our claim immediately follows from the definition. Suppose that $i \geq 1$. We fix i and discuss by induction on m . If $m \leq 0$, the equation (3.28) is clear. Suppose that $m \geq 1$. Then, for any $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $\omega_{-i} \in V_{-i}$ and $u_{m-1}^+ \in U_{m-1}^+$, we have

(3.31)

$$\begin{aligned}
& \pi_{-1,-i+m}^+(y_{-1} \otimes \pi_{-i,m}^+(\omega_{-i} \otimes \pi_{1,m-1}^+(x_1 \otimes u_{m-1}^+))) \\
& \quad - \pi_{-i,m-1}^+(\omega_{-i} \otimes \pi_{-1,m}^+(y_{-1} \otimes \pi_{1,m-1}^+(x_1 \otimes u_{m-1}^+))) \\
& = \pi_{-1,-i+m}^+(y_{-1} \otimes \pi_{-i+1,m-1}^+([\omega_{-i}, x_1] \otimes u_{m-1}^+)) \\
& \quad + \pi_{-1,-i+m}^+(y_{-1} \otimes \pi_{-i+1,m-1}^+(x_1 \otimes \pi_{-i,m-1}^+(\omega_{-i} \otimes u_{m-1}^+))) \\
& \quad - \pi_{-i,m-1}^+(\omega_{-i} \otimes \pi_{0,m-1}^+([y_{-1}, x_1] \otimes u_{m-1}^+)) \\
& \quad - \pi_{-i,m-1}^+(\omega_{-i} \otimes \pi_{1,m-2}^+(x_1 \otimes \pi_{-1,m-1}^+(y_{-1} \otimes u_{m-1}^+))) \\
& = \pi_{-1,-i+m}^+(y_{-1} \otimes \pi_{-i+1,m-1}^+([\omega_{-i}, x_1] \otimes u_{m-1}^+)) - \pi_{-i,m-1}^+(\omega_{-i} \otimes \pi_{0,m-1}^+([y_{-1}, x_1] \otimes u_{m-1}^+))
\end{aligned}$$

$$\begin{aligned}
& + \pi_{0,-i+m-1}^+([y_{-1}, x_1] \otimes \pi_{-i,m-1}^+(\omega_{-i} \otimes u_{m-1}^+)) \\
& + \pi_{1,-i+m-2}^+(x_1 \otimes \pi_{-1,-i+m-1}^+(y_{-1} \otimes \pi_{-i,m-1}^+(\omega_{-i} \otimes u_{m-1}^+))) \\
& - \pi_{-i+1,m-2}^+([\omega_{-i}, x_1] \otimes \pi_{-1,m-1}^+(y_{-1} \otimes u_{m-1}^+)) \\
& - \pi_{1,-i+m-2}^+(x_1 \otimes \pi_{-i,m-2}^+(\omega_{-i} \otimes \pi_{-1,m-1}^+(y_{-1} \otimes u_{m-1}^+))) \\
& = -\pi_{-i,m-1}^+([\omega_{-i}, x_1], y_{-1}] \otimes u_{m-1}^+) + \pi_{-i,m-1}^+([y_{-1}, x_1], \omega_{-i}] \otimes u_{m-1}^+) \\
& + \pi_{1,-i+m-2}^+(x_1 \otimes \pi_{-1,-i+m-1}^+(y_{-1} \otimes \pi_{-i,m-1}^+(\omega_{-i} \otimes u_{m-1}^+))) \\
& - \pi_{1,-i+m-2}^+(x_1 \otimes \pi_{-i,m-2}^+(\omega_{-i} \otimes \pi_{-1,m-1}^+(y_{-1} \otimes u_{m-1}^+))) \\
& = \pi_{-i,m-1}^+([y_{-1}, \omega_{-i}], x_1] \otimes u_{m-1}^+) \\
& + \pi_{1,-i+m-2}^+(x_1 \otimes \pi_{-1,-i+m-1}^+(y_{-1} \otimes \pi_{-i,m-1}^+(\omega_{-i} \otimes u_{m-1}^+))) \\
& - \pi_{1,-i+m-2}^+(x_1 \otimes \pi_{-i,m-2}^+(\omega_{-i} \otimes \pi_{-1,m-1}^+(y_{-1} \otimes u_{m-1}^+))).
\end{aligned}$$

By the induction hypotheses on i and m , if we take elements $y_{-1}^1, \dots, y_{-1}^l \in V_{-1}$ and $\omega_{-i}^1, \dots, \omega_{-i}^l \in V_{-i}$ satisfying $\sum_{s=1}^l q_{-i}(y_{-1}^s \otimes \omega_{-i}^s) = 0$, then we have

$$(3.32) \quad \sum_{s=1}^l \pi_{-i,m-1}^+([y_{-1}^s, \omega_{-i}^s], x_1] \otimes u_{m-1}^+) = 0 \quad (\text{by the induction hypothesis on } i),$$

$$\begin{aligned}
(3.33) \quad & \sum_{s=1}^l (\pi_{1,-i+m-2}^+(x_1 \otimes \pi_{-1,-i+m-1}^+(y_{-1} \otimes \pi_{-i,m-1}^+(\omega_{-i} \otimes u_{m-1}^+))) \\
& - \pi_{1,-i+m-2}^+(x_1 \otimes \pi_{-i,m-2}^+(\omega_{-i} \otimes \pi_{-1,m-1}^+(y_{-1} \otimes u_{m-1}^+)))) = 0 \\
& (\text{by the induction hypothesis on } m).
\end{aligned}$$

Thus, we have

$$\begin{aligned}
(3.34) \quad & \sum_{s=1}^l (\pi_{-1,-i+m}^+(y_{-1}^s \otimes \pi_{-i,m}^+(\omega_{-i}^s \otimes \pi_{1,m-1}^+(x_1 \otimes u_{m-1}^+))) \\
& - \pi_{-i,m-1}^+(\omega_{-i}^s \otimes \pi_{-1,m}^+(y_{-1}^s \otimes \pi_{1,m-1}^+(x_1 \otimes u_{m-1}^+)))) = 0
\end{aligned}$$

from (3.31). Since $\pi_{1,m-1}^+ : V_1 \otimes U_{m-1}^+ \rightarrow U_m^+$ is surjective, we can obtain the equation (3.28). Therefore we can obtain the well-definedness of the linear map $\pi_{-i-1,m}^+ : V_{-i-1} \otimes U_m^+ \rightarrow U_{-i+m-1}^+$ given in (3.15) for any m .

The equation (3.29) can be shown by a similar way to the proof of Proposition 3.8. Moreover, the equation (3.30) follows from (3.31). \square

DEFINITION 3.10. By the above propositions, Propositions 3.8 and 3.9, we define a linear map $\tilde{\pi}^+ : L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) \otimes \tilde{U}^+ \rightarrow \tilde{U}^+$ by:

$$\tilde{\pi}^+(z_n \otimes u_m^+) := \pi_{n,m}^+(z_n \otimes u_m^+)$$

where $n, m \in \mathbb{Z}$, $z_n \in V_n$ and $u_m^+ \in U_m^+$.

This linear map $\tilde{\pi}^+$ satisfies the following equations:

$$\begin{aligned}
& \tilde{\pi}^+([a, z_n] \otimes u_m^+) = \tilde{\pi}^+(a \otimes \tilde{\pi}^+(z_n \otimes u_m^+)) - \tilde{\pi}^+(z_n \otimes \tilde{\pi}^+(a \otimes u_m^+)), \\
& \tilde{\pi}^+([x_1, z_n] \otimes u_m^+) = \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(z_n \otimes u_m^+)) - \tilde{\pi}^+(z_n \otimes \tilde{\pi}^+(x_1 \otimes u_m^+)), \\
& \tilde{\pi}^+([y_{-1}, z_n] \otimes u_m^+) = \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(z_n \otimes u_m^+)) - \tilde{\pi}^+(z_n \otimes \tilde{\pi}^+(y_{-1} \otimes u_m^+))
\end{aligned}$$

for any $n, m \in \mathbb{Z}$, $a \in V_0$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $z_n \in V_n$ and $u_m^+ \in U_m^+$. Moreover, we have the following proposition on $\tilde{\pi}^+$.

Proposition 3.11. *The map $\tilde{\pi}^+$ satisfies the following equation:*

$$(3.35) \quad \tilde{\pi}^+([x, y] \otimes u) = \tilde{\pi}^+(x \otimes \tilde{\pi}^+(y \otimes u)) - \tilde{\pi}^+(y \otimes \tilde{\pi}^+(x \otimes u))$$

for any $x, y \in L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $u^+ \in \tilde{U}^+$.

Proof. To prove our claim, it is sufficient to show the case where $x = z_n \in V_n$ for some $n \in \mathbb{Z}$. We argue by induction on n .

Assume that $n \geq 0$. For $n = 0, 1$, our result has been shown. For $n \geq 2$. We can assume that $z_n = p_{n-1}(x_1 \otimes z_{n-1})$ for some $x_1 \in V_1$ and $z_{n-1} \in V_{n-1}$ without loss of generality. Then, by the induction hypothesis, we have

$$\begin{aligned} \tilde{\pi}^+([p_{n-1}(x_1 \otimes z_{n-1}), y] \otimes u^+) &= \tilde{\pi}^+([x_1, [z_{n-1}, y]] \otimes u^+) - \tilde{\pi}^+([z_{n-1}, [x_1, y]] \otimes u^+) \\ &= \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+([z_{n-1}, y] \otimes u^+)) - \tilde{\pi}^+([z_{n-1}, y] \otimes \tilde{\pi}^+(x_1 \otimes u^+)) \\ &\quad - \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+([x_1, y] \otimes u^+)) + \tilde{\pi}^+([x_1, y] \otimes \tilde{\pi}^+(z_{n-1} \otimes u^+)) \\ &= \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+(y \otimes u^+))) - \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(y \otimes \tilde{\pi}^+(z_{n-1} \otimes u^+))) \\ &\quad - \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+(y \otimes \tilde{\pi}^+(x_1 \otimes u^+))) + \tilde{\pi}^+(y \otimes \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+(x_1 \otimes u^+))) \\ &\quad - \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(y \otimes u^+))) + \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+(y \otimes \tilde{\pi}^+(x_1 \otimes u^+))) \\ &\quad + \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(y \otimes \tilde{\pi}^+(z_{n-1} \otimes u^+))) - \tilde{\pi}^+(y \otimes \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(z_{n-1} \otimes u^+))) \\ &= \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+(y \otimes u^+))) - \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(y \otimes u^+))) \\ &\quad - \tilde{\pi}^+(y \otimes \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(z_{n-1} \otimes u^+))) + \tilde{\pi}^+(y \otimes \tilde{\pi}^+(z_{n-1} \otimes \tilde{\pi}^+(x_1 \otimes u^+))) \\ &= \tilde{\pi}^+([x_1, z_{n-1}] \otimes \tilde{\pi}^+(y \otimes u^+)) - \tilde{\pi}^+(y \otimes \tilde{\pi}^+([x_1, z_{n-1}] \otimes u^+)) \\ &= \tilde{\pi}^+(p_{n-1}(x_1 \otimes z_{n-1}) \otimes \tilde{\pi}^+(y \otimes u^+)) - \tilde{\pi}^+(y \otimes \tilde{\pi}^+(p_{n-1}(x_1 \otimes z_{n-1}) \otimes u^+)). \end{aligned}$$

Thus, we have our result for any $n \geq 0$.

Similarly, we can obtain our result for any $n \leq -1$. This completes the proof. \square

From Proposition 3.11, we have the following theorem.

Theorem 3.12. *The vector space $\tilde{U}^+ = \bigoplus_{m \in \mathbb{Z}} U_m^+ = \bigoplus_{m \geq 0} U_m^+$ has a structure of a positively graded $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module whose representation is $\tilde{\pi}^+$. We call the module $(\tilde{\pi}^+, \tilde{U}^+)$ the positive extension of U with respect to a standard pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. (This is a special case of [9, Theorem 1.2].)*

By the same argument, we can obtain a negatively graded Lie module of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$.

DEFINITION 3.13. We define the following linear maps:

$$\pi_{0,m}^- : V_0 \otimes U_m^- \rightarrow U_m^-, \quad \pi_{1,m}^- : V_1 \otimes U_m^- \rightarrow U_{m+1}^-, \quad \pi_{-1,m}^- : V_{-1} \otimes U_m^- \rightarrow U_{m-1}^-$$

by:

$$(3.36) \quad \pi_{0,m}^-(a \otimes u_m^-) := \pi_m^-(a \otimes u_m^-),$$

$$(3.37) \quad \pi_{1,m}^-(x_1 \otimes u_m^-) := \begin{cases} 0 & (m \geq 0) \\ u_m^-(x_1) & (m \leq -1) \end{cases},$$

$$(3.38) \quad \pi_{-1,m}^-(y_{-1} \otimes u_m^-) := \begin{cases} 0 & (m \geq 1) \\ r_m^-(y_{-1} \otimes u_m^-) & (m \leq 0) \end{cases}$$

where $m \in \mathbb{Z}$, $a \in V_0$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$ and $u_m^- \in U_m^-$.

Theorem 3.14. *The vector space $\tilde{U}^- = \bigoplus_{m \in \mathbb{Z}} U_m^- = \bigoplus_{m \leq 0} U_m^-$ has a structure of a negatively graded $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module whose representation is $\tilde{\pi}^-$. We call the module $(\tilde{\pi}^-, \tilde{U}^-)$ the negative extension of U with respect to a standard pentad $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. (This is a special case of [9, Theorem 1.2].)*

Note that an arbitrary module of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is not necessary written in the form of \tilde{U}^+ or \tilde{U}^- . For example, the adjoint representation of a loop algebra $L(\mathfrak{sl}_2, \text{ad}, \mathfrak{sl}_2, \mathfrak{sl}_2, K_{\mathfrak{sl}_2}) = \mathcal{L}(\mathfrak{sl}_2(\mathbb{C})) = \mathbb{C}[t, t^{-1}] \otimes \mathfrak{sl}_2(\mathbb{C}) = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}t^n \otimes \mathfrak{sl}_2$, where $K_{\mathfrak{sl}_2}$ is the Killing form of \mathfrak{sl}_2 , cannot be written in the form of positively or negatively graded module. Indeed, $\mathcal{L}(\mathfrak{sl}_2(\mathbb{C}))$ does not have a non-zero element which commutes with any element of the form $t \otimes X$ or $t^{-1} \otimes X$ ($X \in \mathfrak{sl}_2$).

Proposition 3.15. *Under the notation of Theorems 3.12 and 3.14, $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -modules \tilde{U}^+ and \tilde{U}^- have the following properties:*

$$(3.39) \quad \tilde{U}^+ \text{ and } \tilde{U}^- \text{ are transitive,}$$

$$(3.40) \quad \tilde{U}^+ \text{ and } \tilde{U}^- \text{ are } L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)\text{-irreducible if and only if } U = U_0^+ = U_0^- \text{ is } \mathfrak{g}\text{-irreducible.}$$

(This is a special case of [9, Theorem 1.1].)

Proof. By the definition, we can show (3.39) immediately.

Let us show (3.40). Assume that U is an irreducible \mathfrak{g} -module. Let \underline{W} be an arbitrary non-zero $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -submodule of \tilde{U}^+ . Then we have that $\underline{W} \cap U_0^+ \neq \{0\}$ (cf. [9, Corollary 1.2]). In fact, take a non-zero element $\underline{w} \in \underline{W}$. Then there exist integers $0 \leq m_1 < \dots < m_k$ and $\underline{w}_{m_1} \in \underline{W} \cap U_{m_1}^+, \dots, \underline{w}_{m_k} \in \underline{W} \cap U_{m_k}^+$ such that $\underline{w} = \underline{w}_{m_1} + \dots + \underline{w}_{m_k}$. Since \tilde{U}^+ is transitive, we can take $y_{-1}^1, \dots, y_{-1}^{m_k} \in V_{-1}$ such that $0 \neq \tilde{\pi}^+(y_{-1}^1 \otimes \dots \otimes \tilde{\pi}^+(y_{-1}^{m_k} \otimes w) \dots) \in \underline{W} \cap U_0^+$. By the assumption that U is irreducible, we have $\underline{W} \cap U_0^+ = U$. Since \tilde{U}^+ is generated by $U = U_0^+$ and V_0, V_1 , we have that \underline{W} coincides with \tilde{U}^+ .

Conversely, assume that \tilde{U}^+ is an irreducible $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module. Take a non-zero \mathfrak{g} -submodule W of U . Then a submodule \underline{W} of \tilde{U}^+ which is generated by V_0, V_1, W is a non-zero $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -submodule of \tilde{U}^+ . Thus, $\underline{W} = \tilde{U}^+$, and, in particular, $W = \underline{W} \cap U_0^+ = U$. Similarly, we can show (3.40) for the negative extension \tilde{U}^- . \square

EXAMPLE 3.16. We retain to use the notations of Example 2.35. Put $U := \mathbb{C}$ and define a representation $\pi : \mathfrak{g} \otimes U \rightarrow U$ by:

$$\pi((a, b, A) \otimes u) := au$$

for any $u \in U$. Then, the positive extension \tilde{U}^+ of U with respect to $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is 3-dimensional irreducible representation of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) = V_{-1} \oplus V_0 \oplus V_1 \simeq \mathfrak{gl}_1 \oplus \mathfrak{sl}_3$. In fact, for any $v \in V_1 = V$, $\phi \in V_{-1} = \mathcal{V}$ and $u \in U$, we have

$$\tilde{\pi}^+(\phi \otimes \tilde{\pi}^+(v \otimes u)) = -\tilde{\pi}^+(\Phi_\rho(v \otimes \phi) \otimes u) = -\pi((-{}^t v \phi, \frac{3}{2} {}^t v \phi, v {}^t \phi - \frac{1}{2} {}^t v \phi I_2) \otimes u) = {}^t v \phi u.$$

Thus, the element $\tilde{\pi}^+(v \otimes u)$ can be identified with $uv \in V_1 = V$ via $\langle \cdot, \cdot \rangle_V$, in particular, U_1^+ is 2-dimensional. Moreover, we have

$$\begin{aligned} & \tilde{\pi}^+(\phi \otimes \tilde{\pi}^+(v' \otimes \tilde{\pi}^+(v \otimes u))) \\ &= -\tilde{\pi}^+((-{}^t v' \phi, \frac{3}{2} {}^t v' \phi, v' {}^t \phi - \frac{1}{2} {}^t v' \phi I_2) \otimes \tilde{\pi}^+(v \otimes u)) + \tilde{\pi}^+(v' \otimes \tilde{\pi}^+(\phi \otimes \tilde{\pi}^+(v \otimes u))) \\ &= -\tilde{\pi}^+({}^t v' \phi \cdot v \otimes u) - \tilde{\pi}^+({}^t v \phi \cdot v' \otimes u) + \tilde{\pi}^+(v \otimes {}^t v' \phi u) + \tilde{\pi}^+(v' \otimes {}^t v \phi u) \\ &= 0 \end{aligned}$$

for any $v, v' \in V_1$, $\phi \in V_{-1}$ and $u \in U$. Therefore, the positive extension $\tilde{U}^+ = U_0^+ \oplus U_1^+$ is a 3-dimensional irreducible representation (see Proposition 3.15).

The positive and negative extensions of U are characterized by the transitivity.

Theorem 3.17. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad. Let $(\underline{\pi}, \underline{U}) = (\underline{\pi}, \bigoplus_{m \geq 0} \underline{U}_m)$ (respectively $(\underline{\omega}, \underline{\mathcal{U}}) = (\underline{\omega}, \bigoplus_{m \leq 0} \underline{\mathcal{U}}_m)$) be a positively graded Lie module (respectively a negatively graded Lie module) of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$. If the $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -module $(\underline{\pi}, \underline{U})$ (respectively $(\underline{\omega}, \underline{\mathcal{U}})$) is transitive and generated by V_0 , V_1 and \underline{U}_0 (respectively generated by V_0 , V_{-1} and $\underline{\mathcal{U}}_0$), then \underline{U} is isomorphic to the positive extension of \underline{U}_0 with respect to $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ (respectively $\underline{\mathcal{U}}$ is isomorphic to the negative extension of $\underline{\mathcal{U}}_0$ with respect to $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$). (This is a special case of [9, Theorem 1.2].)*

Proof. We denote the positive extension of \underline{U}_0 with respect to $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ by

$$\widetilde{\underline{U}}_0^+ = \bigoplus_{m \geq 0} (\underline{U}_m)^+$$

and the canonical representation of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ on $\widetilde{\underline{U}}_0^+$ by $\widetilde{\pi}^+$. Note that $(\underline{U})_0^+ = \underline{U}_0$. We let $\tau_0 : (\underline{U})_0^+ \rightarrow \underline{U}_0$ be the identity map on $(\underline{U})_0^+ = \underline{U}_0$ and define linear maps $\tau_i : (\underline{U})_i^+ \rightarrow \underline{U}_i$ by

$$\tau_i(r_{i-1}^+(x_1 \otimes \underline{u}_{i-1}^+)) := \underline{\pi}(x_1 \otimes \tau_{i-1}(\underline{u}_{i-1}^+))$$

for $i \geq 1$ and any $x_1 \in V_1$ and $\underline{u}_{i-1}^+ \in (\underline{U})_{i-1}^+$ inductively. These τ_i 's are well-defined and satisfy the following equation:

$$(3.41) \quad \tau_{i+j}(\widetilde{\pi}^+(a_j \otimes \underline{u}_i^+)) = \underline{\pi}(a_j \otimes \tau_i(\underline{u}_i^+))$$

for $j = 0, \pm 1$ and any $a_j \in V_j$, $\underline{u}_i^+ \in (\underline{U})_i^+$. Let us show it by induction on i . It is clear that the equation (3.41) holds when $i = 0$ and $j = 0, -1$. In order to show the equation (3.41) for $i = 0$ and $j = 1$, let us show that τ_1 is well-defined. Take an arbitrary element $y_{-1} \in V_{-1}$, then we have

$$\begin{aligned} (3.42) \quad & \underline{\pi}(y_{-1} \otimes \underline{\pi}(x_1 \otimes \tau_0(\underline{u}_0^+))) = \underline{\pi}([y_{-1}, x_1] \otimes \tau_0(\underline{u}_0^+)) + \underline{\pi}(x_1 \otimes \underline{\pi}(y_{-1} \otimes \tau_0(\underline{u}_0^+))) \\ &= \underline{\pi}([y_{-1}, x_1] \otimes \tau_0(\underline{u}_0^+)) = \tau_0(\widetilde{\pi}^+([y_{-1}, x_1] \otimes \underline{u}_0^+)) = \tau_0(\widetilde{\pi}^+(y_{-1} \otimes r_0^+(x_1 \otimes \underline{u}_0^+))). \end{aligned}$$

Thus, if $x_1^1, \dots, x_1^l \in V_1$ and $\underline{u}_0^{+1}, \dots, \underline{u}_0^{+l} \in (\underline{U})_0^+$ satisfy $\sum_{s=1}^l r_0^+(x_1^s \otimes \underline{u}_0^{+s}) = 0$, then we have

$$\sum_{s=1}^l \underline{\pi}(y_{-1} \otimes \underline{\pi}(x_1^s \otimes \tau_0(\underline{u}_0^{+s}))) = 0$$

for any $y_{-1} \in V_{-1}$. Since $(\underline{\pi}, \underline{U})$ is transitive, it follows that $\sum_{s=1}^l \underline{\pi}(x_1^s \otimes \tau_0(\underline{u}_0^{+s})) = 0$, and, thus, we have the well-definedness of τ_1 . By the equation (3.42), we can obtain the equation (3.41) where $i = 0$ and $j = 1$.

Let $i \geq 1$ and assume that τ_0, \dots, τ_i are well-defined and that τ_i satisfies the equation (3.41) for $j = 0, -1$. Then for any $y_{-1} \in V_{-1}$, we have

$$\begin{aligned}
 (3.43) \quad & \underline{\pi}(y_{-1} \otimes \underline{\pi}(x_1 \otimes \tau_i(\underline{u}_i^+))) = \underline{\pi}([y_{-1}, x_1] \otimes \tau_i(\underline{u}_i^+)) + \underline{\pi}(x_1 \otimes \underline{\pi}(y_{-1} \otimes \tau_i(\underline{u}_i^+))) \\
 & = \tau_i(\underline{\pi}^+([y_{-1}, x_1] \otimes \underline{u}_i^+)) + \tau_i(\underline{\pi}^+(x_1 \otimes \underline{\pi}^+(y_{-1} \otimes \underline{u}_i^+))) \\
 & = \tau_i(\underline{\pi}^+(y_{-1} \otimes \underline{\pi}^+(x_1 \otimes \underline{u}_i^+))) = \tau_i(\underline{\pi}^+(y_{-1} \otimes r_0^+(x_1 \otimes \underline{u}_i^+))).
 \end{aligned}$$

Thus, by the same argument to the argument of the case where $i = 0$ and $j = 1$, we have the well-definedness of τ_{i+1} , i.e. τ_i satisfies the equation (3.41) for $j = 1$, and that τ_{i+1} satisfies the equation (3.41) for $j = -1$. Moreover, by a similar argument to the argument of (3.43), we have that τ_{i+1} satisfies the equation (3.41) for $j = 0$. Therefore, by induction on i , we can obtain the well-definedness of τ_i and the equation (3.41) for all $i \geq 0$ and $j = 0, \pm 1$.

We define a linear map $\tau : \widetilde{U}_0^+ \rightarrow \underline{U}$ by

$$(3.44) \quad \tau(\underline{u}_i^+) := \tau_i(\underline{u}_i^+)$$

for any $i \geq 0$ and $\underline{u}_i^+ \in (U)_i^+$. This τ is an isomorphism of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -modules. In fact, by the assumption that \underline{U} is generated by V_1 and \underline{U}_0 , we have the surjectivity of τ . Moreover, by the equation (3.41) in the cases where $i \geq 1$ and $j = -1$ and the definition of τ_0 and the transitivity of the positive extension of \widetilde{U}_0^+ , we have the injectivity of τ . Thus, τ is bijective. Moreover, since $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is generated by $V_0, V_{\pm 1}$, it follows that τ is a homomorphism of $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -modules from the equation (3.41). Therefore \underline{U} is isomorphic to \widetilde{U}_0^+ as $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -modules.

By the same argument, we can prove our claim for $(\underline{U}, \underline{U})$. □

As an application of Theorem 3.17, we have the following proposition.

Proposition 3.18. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ be a standard pentad and U, W (respectively \mathcal{U}, \mathcal{W}) be \mathfrak{g} -modules. Then the positive extension of $U \oplus W$ (respectively the negative extension of $\mathcal{U} \oplus \mathcal{W}$) with respect to $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ is isomorphic to a direct sum of positive extensions of U and W (respectively negative extensions of \mathcal{U} and \mathcal{W}) with respect to $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, i.e.*

$$(\widetilde{U \oplus W})^+ \simeq \widetilde{U}^+ \oplus \widetilde{W}^+ \quad (\text{respectively } (\widetilde{\mathcal{U} \oplus \mathcal{W}})^- \simeq \widetilde{\mathcal{U}}^- \oplus \widetilde{\mathcal{W}}^-).$$

3.3. A pairing between $(\tilde{\pi}^+, \tilde{U}^+)$ and $(\tilde{\omega}^-, \tilde{U}^-)$. In the previous section, we constructed positively and negatively graded $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -modules. Next, let us try to embed these modules into some graded Lie algebra. For this, we need to embed $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(\tilde{\pi}^+, \tilde{U}^+)$ into some standard pentad. However, as mentioned in Remark 2.5, the objects $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and \tilde{U}^+ might not have a submodule of $\text{Hom}(\tilde{U}^+, F)$ and a bilinear form on $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ satisfying the conditions (2.3) and (2.4). In the present and the next sections, we only consider the cases where B_0 is symmetric and U has a submodule $\mathcal{U} \subset \text{Hom}(U, F)$ such that $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ is standard. Then, we can show that a pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$ is standard. First, in this section, we consider the negative extension \tilde{U}^- of \mathcal{U} and construct a non-degenerate invariant bilinear form $\tilde{U}^+ \times \tilde{U}^- \rightarrow F$ under the assumption (2.3) inductively (cf. [9, Remark 1.4]). In the next section, we shall construct the Φ -map of

the pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$.

DEFINITION 3.19. Let $(\tilde{\pi}^+, \tilde{U}^+)$ and $(\tilde{\pi}^-, \tilde{U}^-)$, $\mathcal{U} \subset \text{Hom}(U, F)$ be \mathfrak{g} -modules such that the restriction of the canonical pairing $\langle \cdot, \cdot \rangle_0 : U \times \mathcal{U} \rightarrow F$ is non-degenerate, and, let \tilde{U}^+ and \tilde{U}^- be the positive and negative extensions of U and \mathcal{U} respectively. We define a bilinear map $\langle \cdot, \cdot \rangle_0^0$ by:

$$(3.45) \quad \begin{aligned} \langle \cdot, \cdot \rangle_0^0 : U_0^+ \times \mathcal{U}_0^- &\rightarrow F \\ (u_0^+, w_0^-) &\mapsto \langle u_0^+, w_0^- \rangle_0. \end{aligned}$$

Moreover, for $i \geq 1$, we define a bilinear map $\langle \cdot, \cdot \rangle_{-i}^i$ by:

$$(3.46) \quad \begin{aligned} \langle \cdot, \cdot \rangle_{-i}^i : U_i^+ \times \mathcal{U}_{-i}^- &\rightarrow F \\ (r_{i-1}^+(x_1 \otimes u_{i-1}^+), r_{-i+1}^-(y_{-1} \otimes w_{-i+1}^-)) &\mapsto -\langle \tilde{\pi}^+(y_{-1} \otimes r_{i-1}^+(x_1 \otimes u_{i-1}^+)), w_{-i+1}^- \rangle_{-i+1}^{i-1} \end{aligned}$$

inductively.

The well-definedness of Definition 3.19 can be obtained by the following proposition.

Proposition 3.20. *Let $j \geq 0$. Assume that the bilinear map $\langle \cdot, \cdot \rangle_{-j}^j$ defined in (3.46) is well-defined and satisfies the following equations:*

$$(3.47) \quad \langle \tilde{\pi}^+(a \otimes u_j^+), w_{-j}^- \rangle_{-j}^j + \langle u_j^+, \tilde{\pi}^-(a \otimes w_{-j}^-) \rangle_{-j}^j = 0,$$

$$(3.48) \quad \langle \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(x_1 \otimes u_j^+)), w_{-j}^- \rangle_{-j}^j = \langle u_j^+, \tilde{\pi}^-(x_1 \otimes \tilde{\pi}^-(y_{-1} \otimes w_{-j}^-)) \rangle_{-j}^j$$

$$(3.49) \quad \langle \tilde{\pi}^+(x_1 \otimes u_{j-1}^+), w_{-j}^- \rangle_{-j}^j = \begin{cases} -\langle u_{j-1}^+, \tilde{\pi}^-(x_1 \otimes w_{-j}^-) \rangle_{-j+1}^{j-1} & (j \geq 1) \\ 0 & (j = 0) \end{cases}$$

for any $a \in \mathfrak{g} = V_0 \subset L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $u_{j-1}^+ \in U_{j-1}^+$, $u_j^+ \in U_j^+$ and $w_{-j}^- \in \mathcal{U}_{-j}^-$. Then the bilinear map $\langle \cdot, \cdot \rangle_{-j-1}^{j+1}$ defined in (3.46) is also well-defined and satisfies the following equations:

$$(3.50) \quad \langle \tilde{\pi}^+(a \otimes u_{j+1}^+), w_{-j-1}^- \rangle_{-j-1}^{j+1} + \langle u_{j+1}^+, \tilde{\pi}^-(a \otimes w_{-j-1}^-) \rangle_{-j-1}^{j+1} = 0,$$

$$(3.51) \quad \langle \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(x_1 \otimes u_{j+1}^+)), w_{-j-1}^- \rangle_{-j-1}^{j+1} = \langle u_{j+1}^+, \tilde{\pi}^-(x_1 \otimes \tilde{\pi}^-(y_{-1} \otimes w_{-j-1}^-)) \rangle_{-j-1}^{j+1}$$

$$(3.52) \quad \langle \tilde{\pi}^+(x_1 \otimes u_j^+), w_{-j-1}^- \rangle_{-j-1}^{j+1} = -\langle u_j^+, \tilde{\pi}^-(x_1 \otimes w_{-j-1}^-) \rangle_{-j}^j$$

for any $a \in \mathfrak{g} = V_0 \subset L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $u_j^+ \in U_j^+$, $u_{j+1}^+ \in U_{j+1}^+$ and $w_{-j-1}^- \in \mathcal{U}_{-j-1}^-$.

Proof. First, we let $j = 0$. It is clear that $\langle \cdot, \cdot \rangle_0^0$ satisfies (3.47) and (3.49). Let us show that $\langle \cdot, \cdot \rangle_0^0$ satisfies (3.48). Indeed, under the above notation, we have

$$(3.53) \quad \begin{aligned} \langle \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(x_1 \otimes u_0^+)), w_0^- \rangle_0^0 &= \langle \tilde{\pi}^+([y_{-1}, x_1] \otimes u_0^+), w_0^- \rangle_0^0 = \langle u_0^+, \tilde{\pi}^-([x_1, y_{-1}] \otimes w_0^-) \rangle_0^0 \\ &= \langle u_0^+, \tilde{\pi}^-(x_1 \otimes \tilde{\pi}^-(y_{-1} \otimes w_0^-)) \rangle_0^0. \end{aligned}$$

Thus, the bilinear map $\langle \cdot, \cdot \rangle_0^0$ satisfies the assumptions of Proposition 3.20.

Next, let us show that the bilinear map $\langle \cdot, \cdot \rangle_{-1}^1$ is well-defined. Take arbitrary natural numbers $\nu, \mu \in \mathbb{N}$ and elements $x_1^1, \dots, x_1^\nu \in V_1$, $u_0^{+,1}, \dots, u_0^{+,\nu} \in U_0^+$, $y_{-1}^1, \dots, y_{-1}^\mu \in V_{-1}$,

$w_0^{-,1}, \dots, w_0^{-,\mu} \in \mathcal{U}_0^-$ satisfying

$$\sum_{s=1}^{\nu} r_0^+(x_1^s \otimes u_0^{+,s}) = 0, \quad \sum_{t=1}^{\mu} r_0^-(y_{-1}^t \otimes w_0^{-,t}) = 0.$$

Then, for any $y_{-1} \in V_{-1}$, $w_0^- \in \mathcal{U}_0^-$, $x_1 \in V_1$ and $u_0^+ \in U_0^+$, we have

$$(3.54) \quad \left\langle \sum_{s=1}^{\nu} r_0^+(x_1^s \otimes u_0^{+,s})(y_{-1}), w_0^- \right\rangle_0^0 = 0,$$

and, by the equation (3.53), we have

$$(3.55) \quad \sum_{t=1}^{\mu} \langle r_0^+(x_1 \otimes u_0^+)(y_{-1}^t), w_0^{-,t} \rangle_0^0 = \sum_{t=1}^{\mu} \langle u_0^+, r_0^-(y_{-1}^t \otimes w_0^{-,t})(x_1) \rangle_0^0 = 0.$$

By (3.54) and (3.55), we can obtain that $\langle \cdot, \cdot \rangle_{-1}^1$ is well-defined.

Let us consider properties of $\langle \cdot, \cdot \rangle_{-1}^1$. By (3.53), we have that $\langle \cdot, \cdot \rangle_{-1}^1$ satisfies

$$(3.56) \quad \langle \tilde{\pi}^+(x_1 \otimes u_0^+), w_{-1}^- \rangle_{-1}^1 = -\langle u_0^+, \tilde{\pi}^-(x_1 \otimes w_{-1}^-) \rangle_{-1}^1$$

for any $x_1 \in V_1$, $u_0^+ \in U_0^+$ and $w_{-1}^- \in \mathcal{U}_{-1}^-$, i.e. $\langle \cdot, \cdot \rangle_{-1}^1$ satisfies the equation (3.52). Moreover, $\langle \cdot, \cdot \rangle_{-1}^1$ satisfies the equations (3.50) and (3.51). In fact, for all $a \in V_0$, $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $u_0^+ \in U_0^+$ and $w_0^- \in \mathcal{U}_0^-$, we have

$$(3.57) \quad \begin{aligned} \langle \tilde{\pi}^+(a \otimes r_0^+(x_1 \otimes u_0^+)), r_0^-(y_{-1} \otimes w_0^-) \rangle_{-1}^1 &= -\langle \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(a \otimes \tilde{\pi}^+(x_1 \otimes u_0^+))), w_0^- \rangle_0^0 \\ &= -\langle \tilde{\pi}^+(a \otimes \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(x_1 \otimes u_0^+))), w_0^- \rangle_0^0 + \langle \tilde{\pi}^+([a, y_{-1}] \otimes \tilde{\pi}^+(x_1 \otimes u_0^+)), w_0^- \rangle_0^0 \\ &= \langle \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(x_1 \otimes u_0^+)), \tilde{\pi}^-(a \otimes w_0^-) \rangle_0^0 - \langle \tilde{\pi}^+(x_1 \otimes u_0^+), \tilde{\pi}^-([a, y_{-1}] \otimes w_0^-) \rangle_{-1}^1 \\ &= -\langle \tilde{\pi}^+(x_1 \otimes u_0^+), \tilde{\pi}^-(y_{-1} \otimes \tilde{\pi}^-(a \otimes w_0^-)) \rangle_{-1}^1 - \langle \tilde{\pi}^+(x_1 \otimes u_0^+), \tilde{\pi}^-([a, y_{-1}] \otimes w_0^-) \rangle_{-1}^1 \\ &= -\langle r_0^+(x_1 \otimes u_0^+), \tilde{\pi}^-(a \otimes r_0^-(y_{-1} \otimes w_0^-)) \rangle_{-1}^1. \end{aligned}$$

Thus $\langle \cdot, \cdot \rangle_{-1}^1$ satisfies (3.50). And, from (3.56) and (3.57), we have

$$(3.58) \quad \begin{aligned} \langle \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(x_1 \otimes u_1^+)), w_{-1}^- \rangle_{-1}^1 &= \langle \tilde{\pi}^+([y_{-1}, x_1] \otimes u_1^+), w_{-1}^- \rangle_{-1}^1 + \langle \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(y_{-1} \otimes u_1^+)), w_{-1}^- \rangle_{-1}^1 \\ &= -\langle u_1^+, \tilde{\pi}^-([y_{-1}, x_1] \otimes w_{-1}^-) \rangle_{-1}^1 - \langle \tilde{\pi}^+(y_{-1} \otimes u_1^+), \tilde{\pi}^-(x_1 \otimes w_{-1}^-) \rangle_0^0 \\ &= \langle u_1^+, \tilde{\pi}^-([x_1, y_{-1}] \otimes w_{-1}^-) \rangle_{-1}^1 + \langle u_1^+, \tilde{\pi}^-(y_{-1} \otimes \tilde{\pi}^-(x_1 \otimes w_{-1}^-)) \rangle_{-1}^1 \\ &= \langle u_1^+, \tilde{\pi}^-(x_1 \otimes \tilde{\pi}^-(y_{-1} \otimes w_{-1}^-)) \rangle_{-1}^1 \end{aligned}$$

for any $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $u_1^+ \in U_1^+$ and $w_{-1}^- \in \mathcal{U}_{-1}^-$. Thus $\langle \cdot, \cdot \rangle_{-1}^1$ satisfies (3.51).

We let $j \geq 1$. Suppose that the bilinear map $\langle \cdot, \cdot \rangle_{-j}^j$ is well-defined and satisfies the equations (3.47), (3.48) and (3.49). Let us show the well-definedness of $\langle \cdot, \cdot \rangle_{-j-1}^{j+1}$. Take arbitrary natural numbers $\nu, \mu \in \mathbb{N}$ and elements $x_1^1, \dots, x_1^\nu \in V_1$, $u_j^{+,1}, \dots, u_j^{+,\nu} \in U_0^+$, $y_{-1}^1, \dots, y_{-1}^\mu \in V_{-1}$, $w_{-j}^{-,1}, \dots, w_{-j}^{-,\mu} \in \mathcal{U}_0^-$ satisfying

$$(3.59) \quad \sum_{s=1}^{\nu} r_j^+(x_1^s \otimes u_j^{+,s}) = 0, \quad \sum_{t=1}^{\mu} r_{-j}^-(y_{-1}^t \otimes w_{-j}^{-,t}) = 0.$$

Then, by the equation (3.48) and the same argument to the argument of (3.54) and (3.55), we have the following equations:

$$(3.60) \quad \left\langle \sum_{s=1}^v r_j^+(x_1^s \otimes u_j^{+,s})(y_{-1}), w_{-j}^- \right\rangle_{-j}^j = 0, \quad \sum_{t=1}^{\mu} \langle r_j^+(x_1 \otimes u_j^+)(y_{-1}^t), w_{-j}^- \rangle_{-j}^j = 0.$$

Thus, we have that the bilinear map $\langle \cdot, \cdot \rangle_{-j-1}^{j+1}$ is well-defined.

From the equation (3.48), we have

$$(3.61) \quad \langle \tilde{\pi}^+(x_1 \otimes u_j^+), w_{-j-1}^- \rangle_{-j-1}^{j+1} = -\langle u_j^+, \tilde{\pi}^-(x_1 \otimes w_{-j-1}^-) \rangle_{-j}^j$$

for any $x_1 \in V_1$, $u_j^+ \in U_j^+$ and $w_{-j-1}^- \in \mathcal{U}_{-j-1}^-$. We can show that the bilinear map $\langle \cdot, \cdot \rangle_{-j-1}^{j+1}$ satisfies the equation (3.52) from the equation (3.61) and that it also satisfies the equations (3.50) and (3.51) by the same argument to the argument of the case where $j = 0$. \square

By Proposition 3.20, we can obtain pairings $\langle \cdot, \cdot \rangle_{-j}^j$ for all $j \geq 0$ inductively. Then, we can define a pairing between $(\tilde{\pi}^+, \tilde{U}^+)$ and $(\tilde{\omega}^-, \tilde{\mathcal{U}}^-)$.

DEFINITION 3.21. We define a bilinear map $\langle \cdot, \cdot \rangle : \tilde{U}^+ \times \tilde{\mathcal{U}}^- \rightarrow F$ by:

$$(3.62) \quad \langle u_n^+, w_{-m}^- \rangle := \begin{cases} \langle u_n^+, w_{-n}^- \rangle_{-n}^n & (n = m) \\ 0 & (n \neq m) \end{cases}$$

for any $n, m \geq 0$, $u_n^+ \in U_n^+ \subset \tilde{U}^+$ and $w_{-m}^- \in \mathcal{U}_{-m}^- \subset \tilde{\mathcal{U}}^-$.

By Definition 3.19 and Proposition 3.20, we have that $\langle \cdot, \cdot \rangle$ satisfies

$$(3.63) \quad \langle \tilde{\pi}^+(z_j \otimes \tilde{u}^+), \tilde{w}^- \rangle = -\langle \tilde{u}^+, \tilde{\pi}^-(z_j \otimes \tilde{w}^-) \rangle$$

for $j = 0, \pm 1$ and any $z_j \in V_j$, $\tilde{u}^+ \in \tilde{U}^+$, $\tilde{w}^- \in \tilde{\mathcal{U}}^-$.

Proposition 3.22. *The bilinear form $\langle \cdot, \cdot \rangle : \tilde{U}^+ \times \tilde{\mathcal{U}}^- \rightarrow F$ is non-degenerate and $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -invariant (cf. [9, Definition 1.4 and Remark 1.4]).*

Proof. First, let us show that the bilinear form $\langle \cdot, \cdot \rangle$ is non-degenerate. For this, it is sufficient to show that the bilinear map $\langle \cdot, \cdot \rangle_{-j}^j : U_j^+ \times \mathcal{U}_{-j}^- \rightarrow F$ is non-degenerate for each $j \geq 0$. We show it by induction on j . For $j = 0$, it follows that $\langle \cdot, \cdot \rangle_0^0$ is non-degenerate from the assumption. For $j + 1$, we take an element $u_{j+1}^+ \in U_{j+1}^+$ which satisfies $\langle u_{j+1}^+, r_{-j}^-(y_{-1} \otimes w_{-j}^-) \rangle_{-j-1}^{j+1} = 0$ for any $y_{-1} \in V_{-1}$ and $w_{-j}^- \in \mathcal{U}_{-j}^-$. Then, we have

$$0 = \langle u_{j+1}^+, r_{-j}^-(y_{-1} \otimes w_{-j}^-) \rangle_{-j-1}^{j+1} = -\langle \tilde{\pi}^+(y_{-1} \otimes u_{j+1}^+), w_{-j}^- \rangle_{-j}^j = -\langle u_{j+1}^+(y_{-1}), w_{-j}^- \rangle_{-j}^j.$$

By the induction hypothesis that $\langle \cdot, \cdot \rangle_{-j}^j$ is non-degenerate, we can obtain that $u_{j+1}^+(y_{-1}) = 0$ for any $y_{-1} \in V_{-1}$, and, thus, we have $u_{j+1}^+ = 0 \in U_{j+1}^+ \subset \text{Hom}(V_{-1}, U_j^+)$. Similarly, we can show that an element $w_{-j-1}^- \in \mathcal{U}_{-j-1}^-$ which satisfies $\langle r_j^+(x_1 \otimes u_j^+), w_{-j-1}^- \rangle_{-j-1}^{j+1} = 0$ for any $x_1 \in V_1$ and $u_j^+ \in U_j^+$ is 0 by (3.63). Summarizing the above argument, we can obtain that the map $\langle \cdot, \cdot \rangle_{-j-1}^{j+1}$ is non-degenerate. Therefore, by induction, we can obtain that the bilinear map $\langle \cdot, \cdot \rangle : \tilde{U}^+ \times \tilde{\mathcal{U}}^- \rightarrow F$ is non-degenerate.

Next, let us show that the bilinear map $\langle \cdot, \cdot \rangle$ is $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -invariant. For this, it is sufficient to show that the following equation holds:

$$(3.64) \quad \langle \tilde{\pi}^+(x_j \otimes u_n^+), w_{-n-j}^- \rangle_{-n-j}^{n+j} + \langle u_n^+, \tilde{\pi}^-(x_j \otimes w_{-n-j}^-) \rangle_{-n}^n = 0$$

for any $j, n \in \mathbb{Z}$, $x_j \in V_j$, $u_n \in U_n^+$ and $w_{-n-j}^- \in \mathcal{U}_{-n-j}^-$. We shall show it by induction on j . Assume that $j \geq 0$. For $j = 0, 1$, the equation (3.64) follows from (3.63) immediately. For $j + 1$, by induction hypothesis, we have

$$(3.65) \quad \begin{aligned} & \langle \tilde{\pi}^+([v_1, x_j] \otimes u_n^+), w_{-n-j-1}^- \rangle_{-n-j-1}^{n+j+1} \\ &= \langle \tilde{\pi}^+(v_1 \otimes \tilde{\pi}^+(x_j \otimes u_n^+)), w_{-n-j-1}^- \rangle_{-n-j-1}^{n+j+1} - \langle \tilde{\pi}^+(x_j \otimes \tilde{\pi}^+(v_1 \otimes u_n^+)), w_{-n-j-1}^- \rangle_{-n-j-1}^{n+j+1} \\ &= -\langle \tilde{\pi}^+(x_j \otimes u_n^+), \tilde{\pi}^-(v_1 \otimes w_{-n-j-1}^-) \rangle_{-n-j}^{n+j} + \langle \tilde{\pi}^+(v_1 \otimes u_n^+), \tilde{\pi}^-(x_j \otimes w_{-n-j-1}^-) \rangle_{-n-1}^{n+1} \\ &= \langle u_n^+, \tilde{\pi}^-(x_j \otimes \tilde{\pi}^-(v_1 \otimes w_{-n-j-1}^-)) \rangle_{-n}^n - \langle u_n^+, \tilde{\pi}^-(v_1 \otimes \tilde{\pi}^-(x_j \otimes w_{-n-j-1}^-)) \rangle_{-n}^n \\ &= -\langle u_n^+, \tilde{\pi}^-([v_1, x_j] \otimes w_{-n-j-1}^-) \rangle_{-n}^n \end{aligned}$$

for any $n \in \mathbb{Z}$, $x_1 \in V_1$, $v_j \in V_j$, $u_n^+ \in U_n^+$ and $w_{-n-j-1}^- \in \mathcal{U}_{-n-j-1}^-$. Thus, by induction, we can show the equation (3.64) for all $j \geq 0$. Similarly, we can obtain the equation (3.64) for all $j \leq 0$. Thus, we have the equation (3.64) for all $j \in \mathbb{Z}$. Therefore the bilinear map $\langle \cdot, \cdot \rangle : \tilde{U}^+ \times \tilde{U}^- \rightarrow F$ is $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -invariant. \square

By Proposition 3.22, we can regard \tilde{U}^- as an $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ -submodule of $\text{Hom}(\tilde{U}^+, F)$.

3.4. The Φ -map between $(\tilde{\pi}^+, \tilde{U}^+)$ and $(\tilde{\pi}^-, \tilde{U}^-)$. We retain to assume that a pentad $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ is standard and the bilinear form B_0 is symmetric. As I proved in section 3.3, a pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$ satisfies the condition (2.3). Let us construct the Φ -map of the pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$ and show that it is standard.

DEFINITION 3.23. Assume that pentads $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ are standard and that B_0 is symmetric. We define a linear map $\tilde{\Phi}_0^0 : U_0^+ \otimes \mathcal{U}_0^- \rightarrow V_0$ as:

$$(3.66) \quad \tilde{\Phi}_0^0(u_0^+ \otimes w_0^-) := \Phi_\pi(u_0^+ \otimes w_0^-)$$

where $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $u_0^+ \in U_0^+$, $w_0^- \in \mathcal{U}_0^-$ and Φ_π is the Φ -map of $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$.

Moreover, for each $i \geq 0$, we inductively define a linear map $\tilde{\Phi}_0^{i+1} : U_{i+1}^+ \otimes \mathcal{U}_0^- \rightarrow V_{i+1}$ by:

$$(3.67) \quad \tilde{\Phi}_0^{i+1}(r_i^+(x_1 \otimes u_i^+) \otimes w_0^-) := [x_1, \tilde{\Phi}_0^i(u_i^+ \otimes w_0^-)],$$

where $x_1 \in V_1$, $y_{-1} \in V_{-1}$, $u_i^+ \in U_i^+$ and $w_0^- \in \mathcal{U}_0^-$.

Assume that an integer $j \geq 0$ satisfies a condition that we have linear maps $\tilde{\Phi}_{-j}^k : U_k^+ \otimes \mathcal{U}_{-j}^- \rightarrow V_{k-j}$ for all $k \geq 0$. Then, for any $k \geq 0$, we define a linear map $\tilde{\Phi}_{-j-1}^k : U_k^+ \otimes \mathcal{U}_{-j-1}^- \rightarrow V_{k-j-1}$ by:

$$(3.68) \quad \begin{aligned} & \tilde{\Phi}_{-j-1}^k(u_k^+ \otimes r_{-j}^-(y_{-1} \otimes w_{-j}^-)) \\ &:= \begin{cases} [y_{-1}, \tilde{\Phi}_{-j}^0(u_0^+ \otimes w_{-j}^-)] & (k = 0) \\ [y_{-1}, \tilde{\Phi}_{-j}^k(u_k^+ \otimes w_{-j}^-)] - \tilde{\Phi}_{-j}^{k-1}(\tilde{\pi}^+(y_{-1} \otimes u_k^+) \otimes w_{-j}^-) & (k \geq 1) \end{cases} \end{aligned}$$

where $y_{-1} \in V_{-1}$, $u_k^+ \in U_k^+$ and $w_{-j}^- \in \mathcal{U}_{-j}^-$.

Consequently, we can define linear maps $\tilde{\Phi}_{-j}^i : U_i^+ \otimes \mathcal{U}_{-j}^- \rightarrow V_{i-j}$ for all $i, j \geq 0$.

Proposition 3.24. *The linear map $\tilde{\Phi}_{-j}^i$ is well-defined and satisfies the following equation:*

$$(3.69) \quad B_L(a_{-i+j}, \tilde{\Phi}_{-j}^i(u_i^+ \otimes w_{-j}^-)) = \langle \tilde{\pi}^+(a_{-i+j} \otimes u_i^+), w_{-j}^- \rangle$$

for any $i, j \geq 0$, $a_{-i+j} \in V_{-i+j}$, $u_i^+ \in U_i^+$ and $w_{-j}^- \in \mathcal{U}_{-j}^-$.

Proof. Let us show that the linear maps defined by the equations (3.66), (3.67) and (3.68) satisfy our claim by induction. First, let us show that the linear map $\tilde{\Phi}_0^{i+1}$ ($i \geq 0$) defined in (3.67) is well-defined by induction on i . For $i = 0$, under the above notation, we have

$$(3.70) \quad B_L(a_{-1}, [\tilde{\Phi}_0^0(u_0^+ \otimes w_0^-)]) = B_L([a_{-1}, x_1], \tilde{\Phi}_0^0(u_0^+ \otimes w_0^-)) = \langle \tilde{\pi}^+([a_{-1}, x_1] \otimes u_0^+), w_0^- \rangle \\ = \langle r_0^+(x_1 \otimes u_0^+)(a_{-1}), w_0^- \rangle = \langle \tilde{\pi}^+(a_{-1} \otimes r_0^+(x_1 \otimes u_0^+)), w_0^- \rangle$$

for any $a_{-1} \in V_{-1}$. Thus, if $x_1^1, \dots, x_1^l \in V_1$ and $u_0^{+,1}, \dots, u_0^{+,l} \in U_0^+$ satisfy $\sum_{s=1}^l r_0^+(x_1^s \otimes u_0^{+,s}) = 0$, then we have

$$(3.71) \quad \sum_{s=1}^l B_L(a_{-1}, [\tilde{\Phi}_0^0(u_0^{+,s} \otimes w_0^-)]) = 0$$

for any $a_{-1} \in V_{-1}$. Since the restriction of B_L to $V_{-1} \times V_1$ is non-degenerate, we have

$$(3.72) \quad \sum_{s=1}^l [x_1^s, \tilde{\Phi}_0^0(u_0^{+,s} \otimes w_0^-)] = 0,$$

and, thus, the map $\tilde{\Phi}_0^1$ is well-defined. The equation (3.69) follows from (3.70).

For $i \geq 1$, under the notation of (3.67), we have

$$(3.73) \quad B_L(a_{-i-1}, [\tilde{\Phi}_0^i(u_i^+ \otimes w_0^-)]) = B_L([a_{-i-1}, x_1], \tilde{\Phi}_0^i(u_i^+ \otimes w_0^-)) \\ = \langle \tilde{\pi}^+([a_{-i-1}, x_1] \otimes u_i^+), w_0^- \rangle \\ = \langle \tilde{\pi}^+(a_{-i-1} \otimes \tilde{\pi}^+(x_1 \otimes u_i^+)), w_0^- \rangle - \langle \tilde{\pi}^+(x_1 \otimes \tilde{\pi}^+(a_{-i-1} \otimes u_i^+)), w_0^- \rangle \\ = \langle \tilde{\pi}^+(a_{-i-1} \otimes r_i^+(x_1 \otimes u_i^+)), w_0^- \rangle$$

by the induction hypothesis for any $a_{-i-1} \in V_{-i-1}$. Thus, by the same argument to the argument of the case where $i = 0$, we have the well-definedness of $\tilde{\Phi}_0^{i+1}$ and that $\tilde{\Phi}_0^{i+1}$ satisfies the equation (3.69). Therefore, by induction, we can obtain our claim on $\tilde{\Phi}_0^{i+1}$ for all $i \geq 0$.

Let us show that the linear maps defined in (3.68) are well-defined. We assume that an integer $i \geq 0$ satisfies the condition that we have linear maps $\tilde{\Phi}_{-i}^k : U_k^+ \otimes \mathcal{U}_{-i}^- \rightarrow V_{k-i}$ for all $k \geq 0$ which satisfy the equation (3.69). When $i = 0$, it has been shown that this assumption holds. Then, we can show the well-definedness of the linear maps $\tilde{\Phi}_{-1}^k$ ($k \geq 0$) by induction on k . When $k = 0$, we can show that $\tilde{\Phi}_{-1}^0$ is well-defined and satisfies (3.69) by a similar argument to the argument of (3.67). When $k \geq 1$, we have

$$(3.74) \quad B_L(a_{-k+1}, [\tilde{\Phi}_0^k(u_k^+ \otimes w_0^-)]) - \tilde{\Phi}_0^{k-1}(\tilde{\pi}^+(y_{-1} \otimes u_k^+) \otimes w_0^-) \\ = B_L([a_{-k+1}, y_{-1}], \tilde{\Phi}_0^k(u_k^+ \otimes w_0^-)) - B_L(a_{-k+1}, \tilde{\Phi}_0^{k-1}(\tilde{\pi}^+(y_{-1} \otimes u_k^+) \otimes w_0^-)) \\ = \langle \tilde{\pi}^+([a_{-k+1}, y_{-1}] \otimes u_k^+), w_0^- \rangle - \langle \tilde{\pi}^+(a_{-k+1} \otimes \tilde{\pi}^+(y_{-1} \otimes u_k^+)), w_0^- \rangle \\ = -\langle \tilde{\pi}^+(y_{-1} \otimes \tilde{\pi}^+(a_{-k+1} \otimes u_k^+)), w_0^- \rangle \\ = \langle \tilde{\pi}^+(a_{-k+1} \otimes u_k^+), \tilde{\pi}^-(y_{-1} \otimes w_0^-) \rangle = \langle \tilde{\pi}^+(a_{-k+1} \otimes u_k^+), r_0^-(y_{-1} \otimes w_0^-) \rangle$$

for any $k \geq 1$ and $a_{-k+1} \in V_{-k+1}$ under the notation of (3.68). Thus, by a similar argument to the argument of (3.67), we have the well-definedness of $\tilde{\Phi}_{-1}^k$ for all $k \geq 1$ and that $\tilde{\Phi}_{-1}^k$ satisfies the equation (3.69). For $i \geq 1$, by the same argument to the argument of the case where $i = 0$, we have the well-definedness of $\tilde{\Phi}_{-i-1}^k$ for all $k \geq 0$ and that $\tilde{\Phi}_{-i-1}^k$ satisfies the equation (3.69). Thus, by induction, we have linear maps $\tilde{\Phi}_{-j}^i$ for all $i, j \geq 0$ which satisfies the equation (3.69). This completes the proof. \square

As a corollary of Propositions 3.22 and 3.24, we have the following theorem.

Theorem 3.25. *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ be standard pentads and assume that B_0 is symmetric. Then a pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ is also a standard pentad whose Φ -map, denoted by $\tilde{\Phi}_{\pi}^+$, is defined by:*

$$(3.75) \quad \tilde{\Phi}_{\pi}^+(u_i^+ \otimes w_{-j}^-) := \tilde{\Phi}_{-j}^i(u_i^+ \otimes w_{-j}^-)$$

for any $i, j \geq 0$, $u_i^+ \in U_i^+$ and $w_{-j}^- \in \mathcal{U}_{-j}^-$, where $\tilde{\Phi}_{-j}^i$ is the linear map defined in Definition 3.23.

3.5. Chain rule. Under the assumptions of sections 3.3 and 3.4, let us construct the Lie algebra associated with a standard pentad of the form $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, U^+, \mathcal{U}^-, B_L)$. To find the structure of the Lie algebra $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, U^+, \mathcal{U}^-, B_L)$, we give the following theorem.

Theorem 3.26 (chain rule). *Let $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ be standard pentads. Assume that B_0 is symmetric. Then a pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ is also a standard pentad and the Lie algebra associated with it is isomorphic to $L(\mathfrak{g}, \rho \oplus \pi, V \oplus U, \mathcal{V} \oplus \mathcal{U}, B_0)$, i.e. we have*

$$(3.76) \quad L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L) \simeq L(\mathfrak{g}, \rho \oplus \pi, V \oplus U, \mathcal{V} \oplus \mathcal{U}, B_0)$$

as Lie algebras up to grading.

Proof. Note that the pentad $(\mathfrak{g}, \rho \oplus \pi, V \oplus U, \mathcal{V} \oplus \mathcal{U}, B_0)$ is a standard pentad whose Φ -map $\Phi_{\rho \oplus \pi}$ is given by:

$$\Phi_{\rho \oplus \pi}((v, u) \otimes (\phi, \psi)) = \Phi_{\rho}(v \otimes \phi) + \Phi_{\pi}(u \otimes \psi)$$

where $v \in V$, $\phi \in \mathcal{V}$, $u \in U$, $\psi \in \mathcal{U}$ and Φ_{ρ} and Φ_{π} are the Φ -maps of the pentads $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(\mathfrak{g}, \pi, U, \mathcal{U}, B_0)$ respectively. It has been already shown in Theorem 3.25 that the pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ is standard. We denote the n -graduations of $(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ by V_n and $(\tilde{U}^+)_n$, i.e.

$$(3.77) \quad L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0) = \bigoplus_{n \in \mathbb{Z}} V_n, \quad L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L) = \bigoplus_{m \in \mathbb{Z}} (\tilde{U}^+)_m.$$

Moreover, we denote $(\tilde{U}^+)_1$ and $(\tilde{U}^+)_{-1}$ by:

$$(3.78) \quad (\tilde{U}^+)_1 = \bigoplus_{i \geq 0} U_i^+, \quad (\tilde{U}^+)_{-1} = \bigoplus_{j \geq 0} \mathcal{U}_{-j}^-.$$

Denote a bilinear form on $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ defined in Definition 2.18 by \bar{B}_L . By Lemmas 2.37 and 2.38, we can define derivations α and β on $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+,$

$U^+, \mathcal{U}^-, B_L)$ which satisfy

$$(3.79) \quad \alpha(v_n) = nv_n, \quad \alpha(u_i^+) = iu_i^+, \quad \alpha(w_{-j}^-) = -jw_{-j}^-, \quad \beta(\tilde{u}_m^+) = m\tilde{u}_m^+$$

and

$$(3.80) \quad \overline{B}_L(\alpha(\bar{z}), \bar{\omega}) + \overline{B}_L(\bar{z}, \alpha(\bar{\omega})) = \overline{B}_L(\beta(\bar{z}), \bar{\omega}) + \overline{B}_L(\bar{z}, \beta(\bar{\omega})) = 0$$

for any $n, m \in \mathbb{Z}$, $i, j \geq 0$, $v_n \in V_n$, $u_i^+ \in U_i^+ \subset (\tilde{U}^+)_1$, $w_{-j}^- \in \mathcal{U}_{-j}^- \subset (\tilde{U}^+)^-_{-1}$, $\tilde{u}_m^+ \in (\tilde{U}^+)_m$ and $\bar{z}, \bar{\omega} \in L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$. Since $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ is generated by $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(\tilde{U}^+)_{\pm 1}$ and since $L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0)$ and $(\tilde{U}^+)_{\pm 1}$ are generated by $V_0, V_{\pm 1}$, $U = U_0^+$ and $\mathcal{U} = \mathcal{U}_0^-$, we have that $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ is generated by $V_0, V_{\pm 1}, U_0^+$ and \mathcal{U}_0^- . Put

$$W_{(n,m)} := \left\{ \bar{X} \in L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L) \mid \alpha(\bar{X}) = n\bar{X}, \beta(\bar{X}) = m\bar{X} \right\}$$

for any $n, m \in \mathbb{Z}$. Then we can easily show that all eigenvalues of α and β are integers by induction and that $[W_{(n,m)}, W_{(k,l)}] \subset W_{(n+k, m+l)}$. Thus, we can obtain the following \mathbb{Z} -grading of $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ induced by the eigenspace decomposition of $\gamma := \alpha + \beta$:

$$L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L) = \bigoplus_{k \in \mathbb{Z}} \left(\bigoplus_{n+m=k} W_{(n,m)} \right).$$

If we put $W_k^\gamma := \left\{ \bar{X} \mid \gamma(\bar{X}) = k\bar{X} \right\}$, then we have $W_k^\gamma = \bigoplus_{n+m=k} W_{(n,m)}$ and, thus, we can obtain the following \mathbb{Z} -grading of $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$:

$$L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L) = \bigoplus_{k \in \mathbb{Z}} W_k^\gamma.$$

In particular,

$$(3.81) \quad W_0^\gamma = V_0, \quad W_1^\gamma = V_1 \oplus U_0^+, \quad W_{-1}^\gamma = V_{-1} \oplus \mathcal{U}_0^-.$$

We can easily show that $W_{k+1}^\gamma = [W_1^\gamma, W_k^\gamma]$, $W_{-k-1}^\gamma = [W_{-1}^\gamma, W_{-k}^\gamma]$ for all $k \geq 1$ and that the restriction of \overline{B}_L to $W_k^\gamma \times W_{-k}^\gamma$ is non-degenerate for any $k \in \mathbb{Z}$ from (3.80). Therefore, by Theorem 2.20, we have the isomorphism (3.76). \square

EXAMPLE 3.27. We retain to use the notations of Examples 2.35 and 3.16. Put $\mathcal{U} := \mathbb{C}$ and define a representation $\varpi : \mathfrak{g} \otimes \mathcal{U} \rightarrow \mathcal{U}$ and a bilinear map $\langle \cdot, \cdot \rangle_U : U \times \mathcal{U} \rightarrow \mathbb{C}$ by:

$$\varpi((a, b, A) \otimes w) := -aw, \quad \langle u, w \rangle_U := uw.$$

We can identify \mathcal{U} with $\text{Hom}(U, \mathbb{C})$ via $\langle \cdot, \cdot \rangle_U$. Then pentads $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{\mathcal{U}}^-, B_L)$ and $(\mathfrak{g}, \rho \oplus \pi, V \oplus U, \mathcal{V} \oplus \mathcal{U}, B_0)$ are standard. Let us show that the Lie algebra $L(\mathfrak{g}, \rho \oplus \pi, V \oplus U, \mathcal{V} \oplus \mathcal{U}, B_0)$ is isomorphic to \mathfrak{sl}_4 . Put elements

$$H_0 := \begin{pmatrix} \frac{5}{4} & & & \\ & \frac{1}{4} & & \\ & & \frac{-3}{4} & \\ & & & \frac{-3}{4} \end{pmatrix}, \quad H_1 := \begin{pmatrix} \frac{3}{4} & & & \\ & \frac{-1}{4} & & \\ & & \frac{-1}{4} & \\ & & & \frac{-1}{4} \end{pmatrix}, \quad H_2 := \begin{pmatrix} \frac{1}{2} & & & \\ & \frac{1}{2} & & \\ & & \frac{-1}{2} & \\ & & & \frac{-1}{2} \end{pmatrix} \in \mathfrak{sl}_4.$$

Then we can obtain a \mathbb{Z} -grading of \mathfrak{sl}_4 by the eigenspace decomposition of $\text{ad } H_0$:

$$(3.82) \quad \mathfrak{sl}_4 = \bigoplus_{i=-2}^2 \mathfrak{l}_i \quad (\mathfrak{l}_i := \{X \in \mathfrak{sl}_4 \mid [H_0, X] = iX\}).$$

In particular,

$$\mathfrak{l}_0 = \left\{ \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & & \\ 0 & 0 & A & \end{pmatrix} \middle| a, b \in \mathbb{C}, A \in \mathfrak{gl}_2, a + b + \text{Tr}(A) = 0 \right\} \simeq \mathfrak{gl}_1 \oplus \mathfrak{gl}_1 \oplus \mathfrak{sl}_2,$$

$$\mathfrak{l}_1 = \left\{ \begin{pmatrix} 0 & u & 0 & 0 \\ 0 & 0 & v_1 & v_2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \middle| u, v_1, v_2 \in \mathbb{C} \right\}, \quad \mathfrak{l}_{-1} = \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 \\ \psi & 0 & 0 & 0 \\ 0 & \phi_1 & 0 & 0 \\ 0 & \phi_2 & 0 & 0 \end{pmatrix} \middle| \psi, \phi_1, \phi_2 \in \mathbb{C} \right\}.$$

Then, we have that $\mathfrak{l}_0 \simeq \mathbb{C}H_1 \oplus \mathbb{C}H_2 \oplus \mathfrak{sl}_2$ and that the restriction of a bilinear form T , defined by $T(X, X') := \text{Tr}(XX')$ ($X, X' \in \mathfrak{sl}_4$), to $\mathfrak{l}_0 \times \mathfrak{l}_0$ satisfies:

$$T|_{\mathfrak{l}_0 \times \mathfrak{l}_0}((a, b, A), (a', b', A')) = \frac{3}{4}aa' + bb' + \frac{1}{2}(ab' + a'b) + \text{Tr}(AA'),$$

where $a, a' \in \mathbb{C}H_1$, $b, b' \in \mathbb{C}H_2$, $A, A' \in \mathfrak{sl}_2$. Thus, we can easily show that the grading (3.82) and the Killing form of \mathfrak{sl}_4 , denoted by $K_{\mathfrak{sl}_4}$, satisfy the assumptions of Theorem 2.20 and that a pentad $(\mathfrak{l}_0, \text{ad}, \mathfrak{l}_1, \mathfrak{l}_{-1}, K_{\mathfrak{sl}_4}|_{\mathfrak{l}_0 \times \mathfrak{l}_0})$ is equivalent to $(\mathfrak{g}, \rho \oplus \pi, V \oplus U, \mathcal{V} \oplus \mathcal{U}, B_0)$ (cf. [4, 5, 6, the theory of prehomogeneous vector spaces of parabolic type]). Thus, by Theorems 2.20 and 3.26, we have

$$L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L) \simeq L(\mathfrak{g}, \rho \oplus \pi, V \oplus U, \mathcal{V} \oplus \mathcal{U}, B_0) \simeq \mathfrak{sl}_4.$$

In this case, we can directly check that the Lie algebra $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$ is isomorphic to \mathfrak{sl}_4 using Examples 2.34, 2.35 and 3.16. In fact, by the results of Examples 2.35 and 3.16, we have that the pentad $(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$ is equivalent to the pentad $(\mathfrak{gl}_1 \oplus \mathfrak{sl}_3, \Lambda_1, \mathbb{C}^3, \mathbb{C}^3, \kappa_3)$, which is defined in Example 2.34. Thus, we have that the Lie algebra $L(L(\mathfrak{g}, \rho, V, \mathcal{V}, B_0), \tilde{\pi}^+, \tilde{U}^+, \tilde{U}^-, B_L)$ is isomorphic to \mathfrak{sl}_4 .

References

- [1] N. Bourbaki: Lie groups and Lie algebra. Springer, Berlin, 1989.
- [2] V.G. Kac: *Simple irreducible graded Lie algebras of finite growth*, Math. USSR-Izvestija vol.2 (1968), 1271–1311.
- [3] V.G. Kac: Infinite dimensional Lie algebras, third edition, Cambridge University Press, Cambridge, 1990.
- [4] H. Rubenthaler: *Espaces préhomogènes de type parabolique*, Lect. Math. Kyoto Univ. **14** (1982), 189–221.
- [5] H. Rubenthaler: *Espaces préhomogènes de type parabolique*, Thèse d'Etat, Université de Strasbourg, 1982.
- [6] H. Rubenthaler: Algèbres de Lie et espaces préhomogènes (Travaux en Cours), Hermann, Paris, 1992.
- [7] H. Rubenthaler: *Graded Lie algebras associated to a representation of a quadratic algebra*, arXiv: 1410.0031v2 (2014).
- [8] N. Sasano: *Lie algebras generated by Lie modules*, Kyushu J. Math. **68** (2014), 377–403.
- [9] G. Shen: *Graded modules of graded Lie algebras of Cartan type (II)-positive and negative graded modules*, Sci. Senia Ser. A **29** (1986), 1009–1019.

Institute of Mathematics-for-Industry
Kyushu University
744, Motoooka, Nishi-ku
Fukuoka 819-0395
Japan
e-mail: n-sasano@imi.kyushu-u.ac.jp