



Title	Diffusion processes and a class of Markov chains related to population genetics
Author(s)	Sato, Ken-iti
Citation	Osaka Journal of Mathematics. 1976, 13(3), p. 631-659
Version Type	VoR
URL	https://doi.org/10.18910/6856
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

DIFFUSION PROCESSES AND A CLASS OF MARKOV CHAINS RELATED TO POPULATION GENETICS

KEN-ITI SATO*

(Received July 21, 1975)

1. Introduction

We investigate convergence of sequences of Markov chains induced by direct product branching processes, which are defined by Karlin and McGregor [7] with the intention of unified treatment of Markov chains in population genetics. The induced Markov chains that we deal with in this paper have d types ($d \geq 2$) with equal fertility (that is, selection does not occur), and mutation and migration are allowed for. Let \mathbf{R}^{d-1} be the $(d-1)$ -dimensional Euclidean space and let \mathbf{K} be the set of $x = (x_1, \dots, x_{d-1}) \in \mathbf{R}^{d-1}$ such that $x_1 \geq 0, \dots, x_{d-1} \geq 0, 1 - \sum_{i=1}^{d-1} x_i \geq 0$. Under some conditions, we prove convergence of the Markov chains (suitably normalized and interpolated) to the diffusion process on \mathbf{K} with diffusion coefficient $a(x) = (a_{pq}(x))_{p,q=1, \dots, d-1}$ and drift coefficient $b(x) = (b_p(x))_{p=1, \dots, d-1}$ of the form

$$(1.1) \quad a_{pp}(x) = \sigma^2 x_p (1 - x_p),$$

$$(1.2) \quad a_{pq}(x) = -\sigma^2 x_p x_q \quad (p \neq q),$$

$$(1.3) \quad b_p(x) = \sum_{i=1}^{d-1} x_i \alpha_{ip} + (1 - \sum_{i=1}^{d-1} x_i) \alpha_{dp} + (1 - x_p) \mu_p - x_p \mu_p'.$$

Here $\sigma^2, \alpha_{pq}, \mu_p, \mu_p'$ are constants satisfying $\sigma^2 > 0, \alpha_{pq} \geq 0$ ($p \neq q$), $\alpha_{pp} \leq 0, \sum_{q=1}^d \alpha_{pq} = 0, \mu_p \geq 0, \mu_p' = (\sum_{i=1}^d \mu_i) - \mu_p$. In some sense, α_{pq} ($p \neq q$) is the intensity of mutation from type p to type q and μ_p is the intensity of immigration of type p . Our conditions consist of two sorts. The first is some regularity of the branching process with immigration, which induces the Markov chain, imposed on the distributions of the number of offspring and of the number of immigrants. The second is uniqueness, in the sense of martingale problem, of the diffusion process on \mathbf{K} associated with $a(x)$ and $b(x)$. We conjecture that the uniqueness always holds, but we do not yet have the proof. We give a proof of the uniqueness in some special cases including the following:

* Present address: College of Liberal Arts, Kanazawa University, Kanazawa.

- (i) general one-dimensional case ($d-1=1$);
- (ii) no mutation and no migration ($b(x)=0$);
- (iii) migration allowed, but no mutation ($\alpha_{pq}=0$);
- (iv) α_{pq} ($p \neq q$) depends on q alone.

Consequently, the convergence is established in the above cases. Our method applies also to some Markov chain models involving selection, which will be treated in another paper.

Our results have connection with diffusion approximation to genetics model Markov chains, which is not given a rigorous justification but is a powerful tool in population genetics (see Kimura [8], Crow and Kimura [2]).

In the one-dimensional case, Feller [3] considers convergence of Wright's model. Karlin and McGregor [7] make the assertion, without proof, of the convergence of the induced Markov chains under the condition $d-1=1$, $\alpha_{12}>0$, $\alpha_{21}>0$, $\mu_1=\mu_2=0$. They prove in [6] convergence to the same diffusion for a certain birth-and-death process model. Kushner [9] gives an invariance principle related to Section 3 of this paper and mentions an application to a genetics model, but he deals with convergence in the space D . As for the convergence of the corresponding eigenvalues, we give a detailed analysis in [11].

In Section 2 we will formulate our assumptions and results. The proof will be given in the subsequent three sections. Namely, we establish in Section 3 a general invariance principle in a form convenient for us, prove in Section 4 that the moments of the transition probabilities of the induced Markov chains have the desired properties and, in Section 5, check the uniqueness of the solution of the martingale problem in some cases. The most painstaking part is the derivation of the asymptotic form of the moments of the transition probabilities in Section 4. Our main tool is the powerful saddle point method, which is used in [11] in finding asymptotic behavior of the eigenvalues.

I would like to express my hearty thanks to Nobuyuki Ikeda and Shinzo Watanabe for their valuable advice. The uniqueness proof of Theorem 2.4 in the case of $d-1=2$ is due to S. Watanabe.

2. Assumptions and results

Let \mathbf{Z}_+^d be the set of d -dimensional lattice points with nonnegative coordinates. For each positive integer N , let $\{Z^{(N)}(n)=(Z_1^{(N)}(n), \dots, Z_d^{(N)}(n)); n=0, 1, 2, \dots\}$ be a d type branching process with stationary immigration. That is, $\{Z^{(N)}(n)\}$ is a Markov chain taking values in \mathbf{Z}_+^d and there exist generating functions $f_{N,p}(s_1, \dots, s_d)$, $p=1, \dots, d$, and $g_N(s_1, \dots, s_d)$ of distributions in \mathbf{Z}_+^d such that, for any $j=(j_1, \dots, j_d)$ and $k=(k_1, \dots, k_d)$ in \mathbf{Z}_+^d ,

$$P(Z^{(N)}(n+1) = k | Z^{(N)}(n) = j) = \text{coefficient of } s_1^{k_1} \cdots s_d^{k_d} \text{ in} \\ g_N(s_1, \dots, s_d) \prod_{p=1}^d f_{N,p}(s_1, \dots, s_d)^{j_p}.$$

We make the following assumptions.

Assumptions 2.1. (i) $f_{N,p}$ is of the form

$$f_{N,p}(s_1, \dots, s_d) = \sum_{n=0}^{\infty} c_n \left(\sum_{q=1}^d \alpha_{pq}^{(N)} s_q \right)^n \quad \text{for } N \geq 1, 1 \leq p \leq d,$$

where c_n and $\alpha_{pq}^{(N)}$ satisfy the following conditions.

(ii) $\{c_n\}$ is a probability distribution in \mathbb{Z}_+ independent of N and p with $c_0 > 0$ and with maximum span 1 (that is, there is no pair of $\gamma > 1$ and δ such that $\sum_{n=\gamma+\delta}^{\infty} c_n = 1$). Let $a = \sum_{n=0}^{\infty} n c_n$ (mean), $f(w) = \sum_{n=0}^{\infty} c_n w^n$ (generating function), $M(w) = \sum_{n=0}^{\infty} c_n e^{nw}$ (moment generating function), $F(w) = M(w)e^{-w}$, $b = \sup \{w; M(w) < \infty\}$. Then one of the following holds:

- (a) $1 < a \leq +\infty$;
- (b) $a = 1$ and $b > 0$;
- (c) $a < 1$ and $\lim_{w \uparrow b} F'(w) > 0$.

(iii) $\{\alpha_{pq}^{(N)}\}$ is of the form $\alpha_{pq}^{(N)} = \alpha_{pq}/N$ ($p \neq q$) and $\alpha_{pp}^{(N)} = 1 + (\alpha_{pp}/N)$ for all sufficiently large N , where $\{\alpha_{pq}\}$ is independent of N and satisfies $\alpha_{pq} \geq 0$ ($p \neq q$), $\alpha_{pp} \leq 0$, $\sum_{q=1}^d \alpha_{pq} = 0$.

As is remarked in [11], b is positive in the case (c). If $a < 1$, $b > 0$ and $\lim_{w \uparrow b} M(w) = \infty$, then (c) holds. It is easy to prove that (ii) implies the existence of a unique $\beta \in (-\infty, b)$ such that $F'(\beta) = 0$. β is negative, zero, positive in the cases (a), (b), (c), respectively. Let $K(w) = \log M(w)$ for $w < b$. Then $K'(\beta) = 1$ and $K''(\beta) > 0$. See [11], Lemma 2.1. Let $\sigma^2 = K''(\beta)$. If we define an associated distribution $\{\hat{c}_n\}$ of $\{c_n\}$ by $\hat{c}_n = c_n e^{n\beta} / M(\beta)$, then $\{\hat{c}_n\}$ has mean 1 and variance σ^2 .

Assumption 2.2. g_N is independent of N , that is,

$$g_N(s_1, \dots, s_d) = g(s_1, \dots, s_d) = \sum_{k \in \mathbb{Z}_+^d} b_k s_1^{k_1} \dots s_d^{k_d}$$

for some distribution $\{b_k\}$ in \mathbb{Z}_+^d . Moreover, g satisfies $g(e^{\beta+\varepsilon}, \dots, e^{\beta+\varepsilon}) < \infty$ for some $\varepsilon > 0$.

Note that the last condition on g is automatically satisfied in the case (a). Assumption 2.1 implies that reproduction of offspring by one individual of type p is made in two steps—first it produces independently a random number of children of the same type p according to the distribution $\{c_n\}$, and then, each child has a chance of mutation to type q ($p \neq q$) with probability α_{pq}/N .

Assumption 2.2 implies that the immigration probability is independent of N . Hence dependence of the law of $\{Z^{(N)}(n)\}$ on N lies only in $\{\alpha_{pq}^{(N)}\}$.

Let $\{\hat{b}_k\}$ be an associated distribution of $\{b_k\}$ defined by

$$\hat{b}_k = b_k e^{c_{k_1} + \dots + c_{k_d} \beta} / g(e^\beta, \dots, e^\beta) \quad \text{for } k = (k_1, \dots, k_d) \in \mathbf{Z}_+^d.$$

Let (μ_1, \dots, μ_d) be the mean of $\{\hat{b}_k\}$, $\mu_p = \sum_{k \in \mathbf{Z}_+^d} k_p \hat{b}_k$, and let $\mu_p' = (\sum_{q=1}^d \mu_q) - \mu_p$.

These are important characteristics in our discussion.

Let us define the induced Markov chain. Let $\mathbf{J}(N)$ be the set of points

$j = (j_1, \dots, j_d) \in \mathbf{Z}_+^d$ such that $\sum_{p=1}^d j_p = N$. For $j, k \in \mathbf{J}(N)$ let

$$(2.1) \quad P_{jk}^{(N)} = P(Z^{(N)}(n+1) = k | Z^{(N)}(n) = j, Z^{(N)}(n+1) \in \mathbf{J}(N)).$$

Clearly,

$$(2.2) \quad P_{jk}^{(N)} = \frac{1}{A_1(N)} (\text{coefficients of } s_1^{k_1} \dots s_d^{k_d} \text{ in } g(s_1, \dots, s_d) \prod_{p=1}^d f(\sum_{q=1}^d \alpha_{pq}^{(N)} s_q)^{j_p})$$

where

$$(2.3) \quad A_1(N) = P(Z^{(N)}(n+1) \in \mathbf{J}(N) | Z^{(N)}(n) = j) \\ = \text{coefficient of } w^N \text{ in } g(w, \dots, w) f(w)^N.$$

Assumption 2.1 guarantees $A_1(N) > 0$ for large N . See (4.25) in a later section. Hence we can define $P_{jk}^{(N)}$ for all $j, k \in \mathbf{J}(N)$ if N is large. Let $\{X^{(N)}(n) = (X_1^{(N)}(n), \dots, X_d^{(N)}(n)); n=0, 1, \dots\}$ be a Markov chain defined on a probability space $(\Omega^{(N)}, \mathcal{Q}^{(N)})$, taking values in $\mathbf{J}(N)$ with one-step transition probability $P_{jk}^{(N)}$. The initial distribution is given arbitrarily. $X^{(N)}(n)$ is the *induced Markov chain* of Karlin and McGregor. The fact that this includes various genetics models as special cases is shown in Karlin [5].

Since the sum of components is N , we can consider the induced Markov chain as a Markov chain on a $(d-1)$ -dimensional state space. We normalize and interpolate this chain as follows:

$$(2.4) \quad Y^{(N)}(t) = \left(\frac{1}{N} X_1^{(N)}(n), \dots, \frac{1}{N} X_{d-1}^{(N)}(n) \right) \quad \text{for } t = \frac{n}{N},$$

$$(2.5) \quad Y^{(N)}(t) = (n+1-Nt) Y^{(N)}\left(\frac{n}{N}\right) + (Nt-n) Y^{(N)}\left(\frac{n+1}{N}\right) \\ \text{for } \frac{n}{N} \leq t \leq \frac{n+1}{N}.$$

Then $\{Y^{(N)}(t); 0 \leq t < \infty\}$ is a continuous process taking values in \mathbf{K} (\mathbf{K} is defined in Section 1). Let Ω be the space of continuous paths $\omega: [0, \infty) \rightarrow \mathbf{K}$, endowed with the topology of uniform convergence on compact subsets of $[0, \infty)$.

There is a complete separable metric compatible with this topology. Let $x(t, \omega) = \omega(t)$. Let \mathcal{M} be the topological σ -algebra of Ω and \mathcal{M}_t be the σ -algebra generated by $x(s)$, $s \leq t$. Let $P^{(N)}$ be the probability measure on (Ω, \mathcal{M}) that $(\Omega^{(N)}, Q^{(N)}, Y^{(N)}(t); 0 \leq t < \infty)$ induces. That is, $P^{(N)}(B) = Q^{(N)}(\varphi^{-1}(B))$, $B \in \mathcal{M}$, where φ is the measurable mapping from $\Omega^{(N)}$ into Ω defined by $\varphi(\omega^{(N)}) = \omega$, $\omega(t) = Y^{(N)}(t, \omega^{(N)})$. Obviously, $P^{(N)}$ is determined by the initial distribution of $(X^{(N)}(n))$ and its transition probability $(P_{jk}^{(N)})$. What we would like to prove is the convergence of the sequence of the probability measures $(P^{(N)})$.

Define $a(x) = (a_{pq}(x))_{p, q=1, \dots, d-1}$ and $b(x) = (b_p(x))_{p=1, \dots, d-1}$ by (1.1)–(1.3) and consider the following martingale problem: given $x \in K$, to find a probability measure P_x on (Ω, \mathcal{M}) such that $P_x(x(0)=x)=1$ and, for each $\theta \in \mathbf{R}^{d-1}$, $(M_\theta(t), \mathcal{M}_t, P_x; 0 \leq t < \infty)$ is a martingale, where $M_\theta(t)$ is defined by

$$(2.6) \quad M_\theta(t) = \exp \{ \langle \theta, x(t) - x(0) \rangle - \int_0^t \langle \theta, b(x(u)) \rangle du - \frac{1}{2} \int_0^t \langle \theta, a(x(u)) \theta \rangle du \}.$$

Here \langle, \rangle denotes the inner product. We call this problem *the martingale problem on K for a, b starting from x* , or, for short, *the martingale problem (K, a, b, x)* . Such a problem was originated by Stroock and Varadhan [12], but the above problem is different from theirs on the point of the restriction of the state space.

We will prove the following results.

Theorem 2.1. *For any set of σ^2 , α_{pq} , μ_p , the martingale problem (K, a, b, x) has a solution for each $x \in K$.*

Theorem 2.2. *Suppose that the solution P_x of the martingale problem (K, a, b, x) is unique. Let $Y^{(N)}(0) = x^{(N)}$, which is non-random. If $x^{(N)} \rightarrow x$, then the sequence of the measures $\{P^{(N)}\}$ weakly converges to P_x , that is, for each bounded, continuous, real function $\xi(\omega)$ on Ω , we have*

$$(2.7) \quad \int_\Omega \xi(\omega) P^{(N)}(d\omega) \rightarrow \int_\Omega \xi(\omega) P_x(d\omega), \quad N \rightarrow \infty.$$

It is well known that the weak convergence of $P^{(N)}$ to P_x implies (2.7) for any bounded, measurable, real function $\xi(\omega)$ on Ω whose discontinuity is a set of P_x measure 0. Sojourn times and hitting times are included in applications in many cases. See Billingsley [1].

The above two theorems reduce the point to the uniqueness of the solution of the martingale problem (K, a, b, x) . We conjecture that the uniqueness holds, and the following theorems give the proof in some cases.

Theorem 2.3. *Let $d-1=1$. Then, for each $x \in K$, the solution of the martingale problem (K, a, b, x) is unique, and hence the conclusion of Theorem 2.2 holds.*

Theorem 2.4. *Suppose that if $p > q$ and $p' > q$ then $\alpha_{pq} = \alpha_{p'q}$. Then, for each $x \in K$, the solution of the martingale problem (K, a, b, x) is unique, and the conclusion of Theorem 2.2 holds.*

Note that Theorem 2.4 covers the cases (ii), (iii), (iv) in Section 1.

If the martingale problem (K, a, b, x) has a unique solution P_x for every $x \in K$, then $(x(t), \mathcal{M}_t, P_x; x \in K)$ is a strong Markov process, to which the process $\{Y^{(N)}(t); 0 \leq t < \infty\}$ converges as $N \rightarrow \infty$. In case of $d-1=1$, it is a diffusion process on the interval $[0, 1]$ with backward Kolmogorov equation

$$(2.8) \quad \frac{\partial u}{\partial t} = \frac{\sigma^2}{2} x(1-x) \frac{\partial^2 u}{\partial x^2} + \{-x(\alpha_{12} + \mu_2) + (1-x)(\alpha_{21} + \mu_1)\} \frac{\partial u}{\partial x}.$$

If $d-1=2$, then it is a diffusion process on the triangular region K with backward Kolmogorov equation

$$(2.9) \quad \begin{aligned} \frac{\partial u}{\partial t} = & \frac{\sigma^2}{2} \left\{ x_1(1-x_1) \frac{\partial^2 u}{\partial x_1^2} - 2x_1x_2 \frac{\partial^2 u}{\partial x_1 \partial x_2} + x_2(1-x_2) \frac{\partial^2 u}{\partial x_2^2} \right\} \\ & + \{-x_1(\alpha_{12} + \alpha_{13} + \mu_2 + \mu_3) + x_2\alpha_{21} + (1-x_1-x_2)\alpha_{31} + (1-x_1)\mu_1\} \frac{\partial u}{\partial x_1} \\ & + \{-x_2(\alpha_{21} + \alpha_{23} + \mu_1 + \mu_3) + x_1\alpha_{12} + (1-x_1-x_2)\alpha_{32} + (1-x_2)\mu_2\} \frac{\partial u}{\partial x_2}. \end{aligned}$$

Properties of the boundaries for these diffusion processes will be discussed at the end of Section 5.

3. An invariance principle

In this section K denotes an arbitrary compact set in R^l and Ω denotes the space of continuous paths $\omega: [0, \infty) \rightarrow K$. The topology of Ω and the σ -algebras $\mathcal{M}, \mathcal{M}_t$ are defined as in Section 2. Let $a(x) = (a_{pq}(x))_{p,q=1,\dots,l}$ be a continuous, symmetric, nonnegative-definite $l \times l$ -matrix defined on K and $b(x) = (b_p(x))_{p=1,\dots,l}$ be a continuous l -vector defined on K . By the martingale problem (K, a, b, x) we mean the problem to find a probability measure P_x on (Ω, \mathcal{M}) such that $P_x(x(0)=x)=1$ and, for each $\theta \in R^l$, $(M_\theta(t), \mathcal{M}_t, P_x; 0 \leq t < \infty)$ is a martingale, where $M_\theta(t)$ is defined by (2.6). Suppose that, on a probability space $(\Omega^{(N)}, Q^{(N)})$, time homogeneous Markov chains $\{\bar{Y}^{(N)}(n); n=0, 1, \dots\}$ taking values in K are given. Let $\Pi^{(N)}(x, dy)$ be the one-step transition probability of $\{\bar{Y}^{(N)}(n)\}$. Let $\bar{Y}^{(N)}(0) = x_0^{(N)}$ (non-random). Let $K(N)$ be a Borel subset of K such that $Q^{(N)}(\bar{Y}^{(N)}(n) \in K(N)) = 1$, and suppose that $\Pi^{(N)}(x, dy)$ is

defined for $x \in K(N)$ and $dy \subset K(N)$. Define $Y^{(N)}(t)$ by

$$\begin{aligned} Y^{(N)}(t) &= \bar{Y}^{(N)}(n) \quad \text{for } t = n/N, \\ Y^{(N)}(t) &= (n+1-Nt)\bar{Y}^{(N)}(n) + (Nt-n)\bar{Y}^{(N)}(n+1) \\ &\quad \text{for } n/N \leq t \leq (n+1)/N. \end{aligned}$$

Let $P^{(N)}$ be the probability measure induced on (Ω, \mathcal{M}) by the process $\{Y^{(N)}(t); 0 \leq t < \infty\}$.

We will prove the following theorems. By convergence of measures we mean weak convergence.

Theorem 3.1. *Suppose that the following conditions are satisfied:*

- (i) $\lim_{N \rightarrow \infty} \sup_{x \in K(N)} N \Pi^{(N)}(x, \{y; |y-x| > \varepsilon\}) = 0 \quad \text{for every } \varepsilon > 0.$
- (ii) $\lim_{N \rightarrow \infty} \sup_{x \in K(N)} |N \int_{K(N)} (y_p - x_p) \Pi^{(N)}(x, dy) - b_p(x)| = 0$
for $p = 1, \dots, l.$
- (iii) $\lim_{N \rightarrow \infty} \sup_{x \in K(N)} |N \int_{K(N)} (y_p - x_p)(y_q - x_q) \Pi^{(N)}(x, dy) - a_{pq}(x)| = 0$
for $p, q = 1, \dots, l.$
- (iv) $\lim_{N \rightarrow \infty} x_0^{(N)} = x_0.$

Then, the sequence $\{P^{(N)}\}$ is relatively compact and the limit of any convergent subsequence is a solution of the martingale problem (K, a, b, x_0) .

REMARK. An obvious sufficient condition for (i) is

$$(i)' \quad \lim_{N \rightarrow \infty} \sup_{x \in K(N)} N \int_{K(N)} |y-x|^{2+\delta} \Pi^{(N)}(x, dy) = 0 \quad \text{for some } \delta > 0.$$

Theorem 3.2. *If (i)–(iv) hold and if, moreover, uniqueness of the solution P_{x_0} of the martingale problem (K, a, b, x_0) holds, then $P^{(N)}$ converges to P_{x_0} .*

These theorems are close to results of Stroock and Varadhan [12]. The difference lies in the following three points. First, the state space is a set K , not R' . Secondly, $a(x)$ may be degenerate. Thirdly, $\Pi^{(N)}(x, dy)$ need not be defined on the whole of K . The compactness of the state space makes the proof of the invariance principle easier. But the restriction to K of the state space gives to Theorem 3.1 applicability to the proof of the existence of the solution of the martingale problem (K, a, b, x) . Note that, while the martingale problem (R', a, b, x) is proved to have a solution for general bounded continuous a, b , the solution of the problem (K, a, b, x) exists only in some special degenerate cases.

Theorem 3.2 is an obvious consequence of Theorem 3.1. Theorem 3.1 can be proved by an appropriate modification of [12]. We give the proof here for completeness.

Lemma 3.1. $\{P^{(N)}\}$ is relatively compact if (i) and the following two conditions are satisfied:

$$\begin{aligned} \text{(ii)'} \quad & \sup_N \sup_{x \in K(N)} |N \int_{K(N)} (y_p - x_p) \Pi^{(N)}(x, dy)| < \infty \quad \text{for } p = 1, \dots, l, \\ \text{(iii)'} \quad & \sup_N \sup_{x \in K(N)} |N \int_{K(N)} (y_p - x_p)(y_q - x_q) \Pi^{(N)}(x, dy)| < \infty \\ & \text{for } p, q = 1, \dots, l. \end{aligned}$$

Proof. For each finite T , the restriction of $P^{(N)}$ to \mathcal{M}_T can be viewed as a probability measure on $C([0, T], \mathbf{K})$, the space of continuous paths $[0, T] \rightarrow \mathbf{K}$. For T fixed, this restriction of $\{P^{(N)}\}$ is relatively compact by [12], Lemma 10.2. Let $\{P^{(n_N)}\}$ be a subsequence of $\{P^{(N)}\}$. Using the diagonal procedure, one can choose a subsequence $\{P^{(m_N)}\}$ of $\{P^{(n_N)}\}$ such that if, for some T , $\xi_T(\omega)$ is a bounded, continuous, real function on Ω determined by the value $x(t, \omega)$, $t \leq T$, then $\int_{\Omega} \xi_T(\omega) P^{(m_N)}(d\omega)$ is convergent. Let $\rho(\omega, \omega') = \sum_{n=1}^{\infty} 2^{-n} (1 \wedge \max_{0 \leq t \leq n} |\omega(t) - \omega'(t)|)$. Then ρ is a metric compatible with the topology of Ω . For each $\omega \in \Omega$, let $\omega_T(t) = \omega(t \wedge T)$. Then we see that $\sup_{\omega \in \Omega} \rho(\omega, \omega_T) \rightarrow 0$ as $T \rightarrow \infty$. Hence $\int_{\Omega} \xi(\omega) P^{(m_N)}(d\omega)$ is convergent for every bounded, ρ -uniformly continuous, real function $\xi(\omega)$. This suffices for the convergence of $P^{(m_N)}$.

Proof of Theorem 3.1. Relative compactness of $\{P^{(N)}\}$ follows from Lemma 3.1. Let P be the limit of a subsequence of $\{P^{(N)}\}$. What we have to prove is that P is a solution of the martingale problem (\mathbf{K}, a, b, x_0) . For simplicity of notations, we assume that $P^{(N)}$ converges to P . Since $\{\omega; |x(0) - x_0| \leq \varepsilon\}$ is a closed set, we have

$$P(|x(0) - x_0| \leq \varepsilon) \geq \limsup_{N \rightarrow \infty} P^{(N)}(|x(0) - x_0| \leq \varepsilon) = 1,$$

and $P(x(0) = x_0) = 1$ follows. Fix $\theta \in \mathbf{R}'$ and write $M(t)$ for $M_{\theta}(t)$. Let

$$\begin{aligned} \varphi^{(N)}(x) &= \log \left\{ \int_{K(N)} e^{\langle \theta, y - x \rangle} \Pi^{(N)}(x, dy) \right\}, \\ \bar{M}^{(N)}(n) &= \exp \left\{ \langle \theta, \bar{Y}^{(N)}(n) - \bar{Y}^{(N)}(0) \rangle - \sum_{m=0}^{n-1} \varphi^{(N)}(\bar{Y}^{(N)}(m)) \right\}, \\ M^{(N)}(t) &= \exp \left\{ \langle \theta, x(t) - x(0) \rangle - N \int_0^t \varphi^{(N)} \left(x \left(\frac{[Nu]}{N} \right) \right) du \right\}, \end{aligned}$$

where $[Nu]$ is the greatest integer not exceeding Nu , and let $\bar{\mathcal{M}}_n^{(N)}$ be the σ -algebra generated by $\{\bar{Y}^{(N)}(m); m \leq n\}$. It follows from the Markov property of $\{\bar{Y}^{(N)}(n)\}$ that $(\bar{M}^{(N)}(n), \bar{\mathcal{M}}_n^{(N)}, Q^{(N)}; n=0, 1, \dots)$ is a martingale. Hence $(M^{(N)}(n/N), \mathcal{M}_{n/N}, P^{(N)}; n=0, 1, \dots)$ is a martingale, because

$$M^{(N)}\left(\frac{n}{N}\right) = \exp \left\{ \langle \theta, x\left(\frac{n}{N}\right) - x(0) \rangle - \sum_{m=0}^{n-1} \varphi^{(N)}\left(x\left(\frac{m}{N}\right)\right) \right\}.$$

We will prove that

$$(3.1) \quad \lim_{N \rightarrow \infty} E^{P^{(N)}}\left(\xi \cdot M^{(N)}\left(\frac{[Nt]+1}{N}\right)\right) = E^P(\xi \cdot M(t))$$

for every bounded, continuous, \mathcal{M}_t -measurable function $\xi(\omega)$. Here E^P and $E^{P^{(N)}}$ denote expectation with respect to P and $P^{(N)}$, respectively. Suppose, for a moment, that (3.1) is proved. If $s < t$ and ξ is bounded, continuous, \mathcal{M}_s -measurable, then

$$E^{P^{(N)}}\left(\xi \cdot M^{(N)}\left(\frac{[Ns]+1}{N}\right)\right) = E^{P^{(N)}}\left(\xi \cdot M^{(N)}\left(\frac{[Nt]+1}{N}\right)\right)$$

by the martingale property, and, letting $N \rightarrow \infty$, we get

$$E^P(\xi \cdot M(s)) = E^P(\xi \cdot M(t)).$$

This is just the martingale property of $(M(t), \mathcal{M}_t, P)$ and the proof would be complete.

In order to prove (3.1), we will first show

$$(3.2) \quad \lim_{N \rightarrow \infty} \sup_{x \in K^{(N)}} |N \left(\int_{K^{(N)}} e^{\langle \theta, y-x \rangle} \Pi^{(N)}(x, dy) - 1 \right) - \langle \theta, b(x) \rangle - \frac{1}{2} \langle \theta, a(x) \theta \rangle| = 0.$$

We have

$$\begin{aligned} \int e^{\langle \theta, y-x \rangle} \Pi^{(N)}(x, dy) - 1 &= \langle \theta, y-x \rangle \Pi^{(N)}(x, dy) \\ &+ \frac{1}{2} \langle \theta, y-x \rangle^2 \Pi^{(N)}(x, dy) \\ &+ \int_{|y-x| \leq \varepsilon} \langle \theta, y-x \rangle^2 \delta(y-x) \Pi^{(N)}(x, dy) + B \cdot \Pi^{(N)}(x, V_\varepsilon(x)) \end{aligned}$$

where $V_\varepsilon(x) = \{y; |y-x| > \varepsilon\}$, $\delta(y-x) \rightarrow 0$ as $|y-x| \rightarrow 0$, and B is a bounded function of N, x, ε . Let $f_\varepsilon(N) = \sup_{x \in K^{(N)}} N \Pi^{(N)}(x, V_\varepsilon(x))$. Then $f_\varepsilon(N) \rightarrow 0$ as $N \rightarrow \infty$ for each fixed $\varepsilon > 0$ by the assumption (i). Using the assumptions (ii) and (iii) we see that the absolute value in (3.2) is bounded by

$$g(N) + C \sup_{|y-x| \leq \varepsilon} |\delta(y-x)| + Cf_{\varepsilon}(N)$$

for every $\varepsilon > 0$ with $g(N) \rightarrow 0$ and a constant C . Choosing ε_N which decreases to zero sufficiently slowly, we get $f_{\varepsilon_N}(N) \rightarrow 0$, and hence (3.2). Noting that

$$\varphi^{(N)}(x) = \int e^{\langle \theta, y-x \rangle} \Pi^{(N)}(x, dy) - 1 + o\left(\int e^{\langle \theta, y-x \rangle} \Pi^{(N)}(x, dy) - 1\right),$$

we get

$$(3.3) \quad \lim_{N \rightarrow \infty} \sup_{x \in K(N)} |N\varphi^{(N)}(x) - \langle \theta, b(x) \rangle - \frac{1}{2} \langle \theta, a(x)\theta \rangle| = 0$$

from (3.2).

Let $\Gamma^{(N)} = \{\omega; x(n/N, \omega) \in K(N) \text{ for } n=0, 1, \dots\}$. If $\Lambda \in \mathcal{M}_T$ and if Λ is compact when it is viewed as a subset of $C([0, T], K)$, then, for any $t < T$

$$(3.4) \quad \lim_{N \rightarrow \infty} \sup_{\omega \in \Gamma^{(N)} \cap \Lambda} |M^{(N)}\left(\frac{[Nt]+1}{N}, \omega\right) - M(t, \omega)| = 0.$$

In fact, let us write $([Nt]+1)/N = t_N$ and notice that

$$M^{(N)}(t_N, \omega) = \exp \left\{ \langle \theta, x(t_N) - x(0) \rangle - N \int_0^{t_N} \varphi^{(N)}\left(x\left(\frac{[Nu]}{N}\right)\right) du \right\}$$

and

$$\begin{aligned} & N \int_0^{t_N} \varphi^{(N)}\left(x\left(\frac{[Nu]}{N}\right)\right) du - \int_0^t \langle \theta, b(x(u)) \rangle du - \frac{1}{2} \int_0^t \langle \theta, a(x(u))\theta \rangle du \\ &= \int_0^{t_N} \{N\varphi^{(N)}\left(x\left(\frac{[Nu]}{N}\right)\right) - \langle \theta, b\left(x\left(\frac{[Nu]}{N}\right)\right) \rangle - \frac{1}{2} \langle \theta, a\left(x\left(\frac{[Nu]}{N}\right)\right)\theta \rangle\} du \\ &+ \int_0^{t_N} \left\{ \langle \theta, b\left(x\left(\frac{[Nu]}{N}\right)\right) \rangle - b(x(u)) \right\} + \frac{1}{2} \langle \theta, (a\left(x\left(\frac{[Nu]}{N}\right)\right) - a(x(u)))\theta \rangle du \\ &+ \int_t^{t_N} \{ \langle \theta, b(x(u)) \rangle + \frac{1}{2} \langle \theta, a(x(u))\theta \rangle \} du. \end{aligned}$$

Uniformly in $\omega \in \Lambda \cap \Gamma^{(N)}$, the first integral in the right-hand side tends to zero by (3.3), and so does the second integral by the uniform continuity of a, b , and also the third integral tends to zero by the boundedness of a, b . Note that $|x([Nu]/N, \omega) - x(u, \omega)| \rightarrow 0$ uniformly in $u < T$ and $\omega \in \Lambda$.

Now let us prove (3.1). Let

$$\Delta = \sup_N \sup_{\omega \in \Gamma^{(N)}} |\xi(\omega) \left(M^{(N)}\left(\frac{[Nt]+1}{N}, \omega\right) - M(t, \omega) \right)|,$$

which is finite by (3.3). Given $\varepsilon > 0$, choose $\Lambda \in \mathcal{M}_T$ compact as a subset of $C([0, T], K)$ in such a way that $\inf_N P^{(N)}(\Lambda) > 1 - (\varepsilon/\Delta)$. This can be done since $\{P^{(N)}\}$ is tight by Lemma 3.1. Then,

$$\begin{aligned}
 & |E^{P^{(N)}}\left(\xi \cdot M^{(N)}\left(\frac{[Nt]+1}{N}\right)\right)-E^P(\xi \cdot M(t))| \leq \Delta(1-P^{(N)}(\Lambda)) \\
 & +\|\xi\| \int_{\Lambda \cap \Gamma^{(N)}}\left|M^{(N)}\left(\frac{[Nt]+1}{N}\right)-M(t)\right| d P^{(N)} \\
 & +\left|E^{P^{(N)}}(\xi \cdot M(t))-E^P(\xi \cdot M(t))\right| \\
 & < 3 \varepsilon
 \end{aligned}$$

if N is large. Here we have used $P^{(N)} \rightarrow P$ in the third term and (3.4) in the second term. The proof is complete.

4. Asymptotic estimate of the moments of the induced Markov chains and proof of Theorems 2.1 and 2.2

All notations in this section are the same as in Section 2. In particular, $\{X^{(N)}(n)\}$ is the induced Markov chain on $J(N)$. In order to prove Theorems 2.1 and 2.2, we apply Theorems 3.1 and 3.2 with $l=d-1$ and

$$\bar{Y}^{(N)}(n)=\left(\frac{1}{N} X_1^{(N)}(n), \cdots, \frac{1}{N} X_{d-1}^{(N)}(n)\right) .$$

Let $J^0(N)$ be the set of points $j=(j_1, \cdots, j_{d-1}) \in Z_+^{d-1}$ such that $\sum_{p=1}^{d-1} j_p \leq N$ and let

$K(N)=\left\{\frac{j}{N} ; j \in J^0(N)\right\}$. For $j \in J^0(N)$, let

$$(4.1) \quad b_p^{(N)}\left(\frac{j}{N}\right)=N \sum_{k \in J^{(N)}}\left(\frac{k_p-j_p}{N}-\frac{j_p}{N}\right) P_{jk}^{(N)},$$

$$(4.2) \quad a_{pq}^{(N)}\left(\frac{j}{N}\right)=N \sum_{k \in J^{(N)}}\left(\frac{k_p-j_p}{N}-\frac{j_p}{N}\right)\left(\frac{k_q-j_q}{N}-\frac{j_q}{N}\right) P_{jk}^{(N)},$$

$$(4.3) \quad e_p^{(N)}\left(\frac{j}{N}\right)=N \sum_{k \in J^{(N)}}\left(\frac{k_p-j_p}{N}-\frac{j_p}{N}\right)^4 P_{jk}^{(N)} .$$

Here we have identified $j=(j_1, \cdots, j_{d-1}) \in J^0(N)$ with $(j_1, \cdots, j_{d-1}, 1-\sum_{p=1}^{d-1} j_p) \in J(N)$. $b_p(x)$ and $a_{pq}(x)$ are defined by (1.1)–(1.3).

Lemma 4.1. Suppose that for $p, q=1, \cdots, d-1$

$$(4.4) \quad \lim _{N \rightarrow \infty} \sup _{j \in J^0(N)}\left|b_p^{(N)}\left(\frac{j}{N}\right)-b_p\left(\frac{j}{N}\right)\right|=0,$$

$$(4.5) \quad \lim _{N \rightarrow \infty} \sup _{j \in J^0(N)}\left|a_{pq}^{(N)}\left(\frac{j}{N}\right)-a_{pq}\left(\frac{j}{N}\right)\right|=0,$$

$$(4.6) \quad \lim _{N \rightarrow \infty} \sup _{j \in J^0(N)} e_p^{(N)}\left(\frac{j}{N}\right)=0 .$$

Then Theorems 2.1 and 2.2 follow.

Proof. Clearly (4.4) and (4.5) imply Conditions (ii) and (iii) of Theorem 3.1. Since

$$\left(\sum_{p=1}^{d-1} (k_p - j_p)^2\right)^2 \leq (d-1) \sum_{p=1}^{d-1} (k_p - j_p)^4,$$

(4.6) implies (i)' for $\delta=2$. Hence, if $Y^{(N)}(0)=x_0^{(N)} \rightarrow x_0$, then $\{P^{(N)}\}$ is relatively compact and the limit of any convergent subsequence of $\{P^{(N)}\}$ is a solution of the martingale problem (\mathbf{K}, a, b, x_0) . Now, using any branching process with immigration satisfying Assumptions 2.1 and 2.2 (given any set of σ^2 , α_{pq} , μ_p , we can find such a process), we see that the martingale problem (\mathbf{K}, a, b, x) has at least one solution. That Theorem 2.2 follows is a consequence of Theorem 3.2.

Our task in this section is to prove the estimate (4.4)–(4.6). We will prove the following stronger asymptotic formulas :

$$(4.7) \quad b_p^{(N)}\left(\frac{j}{N}\right) = b_p\left(\frac{j}{N}\right) + O\left(\frac{1}{N}\right) \quad \text{uniformly in } j \in \mathbf{J}^0(N),$$

$$(4.8) \quad a_{pq}^{(N)}\left(\frac{j}{N}\right) = a_{pq}\left(\frac{j}{N}\right) + O\left(\frac{1}{N}\right) \quad \text{uniformly in } j \in \mathbf{J}^0(N),$$

$$(4.9) \quad e_p^{(N)}\left(\frac{j}{N}\right) = O\left(\frac{1}{N}\right) \quad \text{uniformly in } j \in \mathbf{J}^0(N).$$

What we mean by (4.7) is

$$\sup_N \sup_{j \in \mathbf{J}^0(N)} N |b_p^{(N)}\left(\frac{j}{N}\right) - b_p\left(\frac{j}{N}\right)| < \infty.$$

We use the phrase “uniformly in $j \in \mathbf{J}^0(N)$ ” in this meaning. In order to prove (4.7)–(4.9) we have to make many estimations of coefficients of power series. Our tool for this is Lemma 4.3 below which is proved by the saddle point method. Naturally the proof of (4.9) is the most complicated. This condition guarantees that the limit process is a diffusion process (that is, sample functions are continuous).

Given $N \geq 1$ and $j \in \mathbf{J}(N)$, let

$$(4.10) \quad G(s_1, \dots, s_d) = \sum_{h \in \mathbf{J}(N)} P_{j_h}^{(N)} s_1^{h_1} \dots s_d^{h_d},$$

$$(4.11) \quad \Phi(w, s_1, \dots, s_d) = \prod_{p=1}^d f(ws_p)^{j_p},$$

$$(4.12) \quad \Psi(w, s_1, \dots, s_d) = \Phi(w, \sum_{l=1}^d \alpha_{1l}^{(N)} s_l, \dots, \sum_{l=1}^d \alpha_{dl}^{(N)} s_l),$$

$$(4.13) \quad \Theta(w, s_1, \dots, s_d) = \Psi(w, s_1, \dots, s_d) g(ws_1, \dots, ws_d).$$

Since we have

$$P_{jk}^{(N)} = A_1(N)^{-1} \text{ (coefficient of } w^N s_1^{k_1} \cdots s_d^{k_d} \text{ in } \Theta(w, s_1, \dots, s_d))$$

by (2.2), we see that

$$(4.14) \quad G(s_1, \dots, s_d) = A_1(N)^{-1} \text{ (coefficient of } w^N \text{ in } \Theta(w, s_1, \dots, s_d)).$$

Let us denote

$$D_{p_1 \dots p_m} = \frac{\partial^m}{\partial s_{p_1} \cdots \partial s_{p_m}},$$

which operates on functions of (s_1, \dots, s_d) or of (w, s_1, \dots, s_d) .

Let

$$C_{p_1 \dots p_m} = D_{p_1 \dots p_m} G(1, \dots, 1), \quad C_{p_1 \dots p_m}^* = \sum_{k \in J(N)} k_{p_1} \cdots k_{p_m} P_{jk}^{(N)}.$$

Lemma 4.2.

$$\begin{aligned} C_p^* &= C_p, \quad C_{pq}^* = C_{pq} \quad (p \neq q), \quad C_{pp}^* = C_{pp} + C_p, \\ C_{ppp}^* &= C_{ppp} + 3C_{pp} + C_p, \quad C_{pppp}^* = C_{pppp} + 6C_{ppp} + 7C_{pp} + C_p. \end{aligned}$$

Proof is easy and omitted.

Functions $M(w)$ and $K(w)$ are defined in Section 2. $M(w)$ extends to an analytic function $M(z)$ of complex z with $\operatorname{Re} z < b$. $K(w)$ extends to $K(z)$ analytic in a neighborhood of β . Define κ_n by

$$(4.15) \quad K(z) = \sum_{n=0}^{\infty} \kappa_n (z - \beta)^n.$$

Note that $\kappa_0 = K(\beta)$, $\kappa_1 = 1$, $\kappa_2 = \sigma^2/2$. Let

$$(4.16) \quad L(s_1, \dots, s_d) = g(e^{s_1}, \dots, e^{s_d}).$$

By Assumption 2.2, $L(s_1, \dots, s_d)$ can be extended to $L(z_1, \dots, z_d)$ for complex z_1, \dots, z_d with $\operatorname{Re} z_l < \beta + \varepsilon$ ($l=1, \dots, d$). Similarly $D_{p_1 \dots p_m} L(s_1, \dots, s_d)$ can be extended to $D_{p_1 \dots p_m} L(z_1, \dots, z_d)$.

Lemma 4.3. Let

$$(4.17) \quad \hat{A}(N) = \frac{1}{2\pi i} \int_{\beta - i\pi}^{\beta + i\pi} M(z)^{N-r} \tilde{M}(z) e^{-Nz} dz, \quad N \geq r,$$

where the integral is along the line segment from $\beta - i\pi$ to $\beta + i\pi$, r is a fixed integer, and $\tilde{M}(z)$ is a bounded continuous function on the segment. Suppose $\tilde{M}(z)$ is analytic in a neighborhood of $z = \beta$ and let

$$M(z)^{-r} \tilde{M}(z) = \sum_{n=0}^{\infty} \tilde{\rho}_n (z - \beta)^n$$

there. Then,

$$\tilde{A}(N) = \Delta_N(\tilde{\rho}_0 + N^{-1}\tilde{a} + O(N^{-2})) \quad \text{as } N \rightarrow \infty,$$

where

$$\Delta_N = \frac{e^{N(K(\beta) - \beta)}}{\sigma\sqrt{2\pi}\sqrt{N}},$$

$$\tilde{a} = \tilde{\rho}_0\left(\frac{3\kappa_4}{\sigma^4} - \frac{15\kappa_3^2}{2\sigma^6}\right) + \tilde{\rho}_1\frac{3\kappa_3}{\sigma^4} - \tilde{\rho}_2\frac{1}{\sigma^2}.$$

Proof is omitted, since this is part of Lemma 3.1 of Sato [11].

Let

$A_\nu(N)$ = coefficient of w^N in a power series $h_\nu(w)g(w, \dots, w)$,

where we define $h_\nu(w)$ as follows:

$$\begin{aligned} h_1(w) &= f(w)^N, \\ h_2(w) &= f(w)^{N-1}f'(w)w, \\ h_3(w) &= f(w)^{N-2}f'(w)^2w^2, \\ h_4(w) &= f(w)^{N-1}f''(w)w^2, \\ h_5(w) &= f(w)^{N-3}f'(w)^3w^3, \\ h_6(w) &= f(w)^{N-2}f'(w)f''(w)w^3, \\ h_7(w) &= f(w)^{N-1}f'''(w)w^3, \\ h_8(w) &= f(w)^{N-4}f'(w)^4w^4, \\ h_9(w) &= f(w)^{N-3}f'(w)^2f''(w)w^4, \\ h_{10}(w) &= f(w)^{N-2}f''(w)^2w^4, \\ h_{11}(w) &= f(w)^{N-2}f'(w)f'''(w)w^4, \\ h_{12}(w) &= f(w)^{N-1}f''''(w)w^4. \end{aligned}$$

Note that $A_1(N)$ is the same that is defined in (2.3). Further, let

$$A_{\nu, p_1 \dots p_m}(N) = \text{coefficient of } w^N \text{ in } h_\nu(w)D_{p_1 \dots p_m}g(w, \dots, w)w^m.$$

All functions which appear above are power series of w with non-negative coefficients and the convergence radii are bigger than e^β . Hence they are convergent at e^z with $\text{Re } z = \beta$.

Lemma 4.4. For each ν, m, p_1, \dots, p_m , we have

$$(4.18) \quad A_\nu(N) = \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} h_\nu(e^z)g(e^z, \dots, e^z)e^{-Nz}dz,$$

$$(4.19) \quad A_{\nu, p_1 \dots p_m}(N) = \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} h_\nu(e^z)D_{p_1 \dots p_m}g(e^z, \dots, e^z)e^{mz}e^{-Nz}dz,$$

and Lemma 4.3 applies.

Proof. We need only use Fubini's theorem to get the integral representation (4.18), (4.19). It is easy to see that they are of the form (4.17) satisfying the conditions in Lemma 4.3.

Given $N \geq 1$ and $j \in J(N)$, we write $x = N^{-1}j$. Hence $x_p = N^{-1}j_p$ for $1 \leq p \leq d$ and $\sum_{p=1}^d x_p = 1$. Sometimes we write $y_p = \sum_{i=1}^d x_i \alpha_{ip}$. We use the notation $(m)_n = m(m-1) \cdots (m-n+1)$.

Lemma 4.5.

$$(4.20) \quad C_p = j_p + \sum_{i=1}^d x_i \alpha_{ip} + \mu_p - x_p \sum_{i=1}^d \mu_i + O(N^{-1}) \text{ uniformly in } j \in J(N),$$

and

$$(4.21) \quad b_p^{(N)}(x) = \sum_{i=1}^d x_i \alpha_{ip} + \mu_p - x_p \sum_{i=1}^d \mu_i + O(N^{-1}) \text{ uniformly in } j \in J(N),$$

that is, (4.7) holds.

Proof. It follows from (4.14) that

$$C_p = D_p G(1, \dots, 1) = A_1(N)^{-1} (\text{coefficient of } w^N \text{ in } D_p \Theta(w, 1, \dots, 1)).$$

From (4.11)—(4.13) we have

$$\begin{aligned} D_p \Theta(w, s_1, \dots, s_d) &= D_p \Psi(w, s_1, \dots, s_d) g(ws_1, \dots, ws_d) \\ &\quad + \Psi(w, s_1, \dots, s_d) D_p g(ws_1, \dots, ws_d) w, \\ D_p \Psi(w, s_1, \dots, s_d) &= \sum_i D_i \Phi(w, \sum_m \alpha_{im}^{(N)} s_m, \dots, \sum_m \alpha_{dm}^{(N)} s_m) \alpha_{ip}^{(N)}, \\ D_i \Phi(w, s_1, \dots, s_d) &= f(ws_1)^{j_1} \cdots \{j_i f(ws_i)^{j_i-1} f'(ws_i) w\} \cdots f(ws_d)^{j_d}. \end{aligned}$$

Noting that

$$\begin{aligned} \Psi(w, 1, \dots, 1) &= \Phi(w, 1, \dots, 1) = f(w)^N, \\ \sum_i j_i \alpha_{ip}^{(N)} &= j_p + y_p, \end{aligned}$$

we have

$$\begin{aligned} D_i \Phi(w, 1, \dots, 1) &= j_i h_2(w), \quad D_p \Psi(w, 1, \dots, 1) = (j_p + y_p) h_2(w), \\ D_p \Theta(w, 1, \dots, 1) &= (j_p + y_p) h_2(w) g(w, \dots, w) + h_1(w) D_p g(w, \dots, w) w. \end{aligned}$$

Hence

$$(4.22) \quad C_p = (j_p + y_p) \frac{A_2(N)}{A_1(N)} + \frac{A_{1p}(N)}{A_1(N)}.$$

Let us estimate $A_1(N)$, $A_2(N)$, $A_{1p}(N)$. By Lemma 4.4,

$$A_1(N) = \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^N L(z, \dots, z) e^{-Nz} dz.$$

Define ρ_n by the expansion

$$(4.23) \quad L(z, \dots, z) = \sum_{n=0}^{\infty} \rho_n (z-\beta)^n \text{ in a neighborhood of } \beta.$$

We have $\rho_0 = g(e^\beta, \dots, e^\beta) > 0$ and

$$(4.24) \quad \rho_1 = \rho_0 \sum_{i=1}^d \mu_i.$$

Thus

$$(4.25) \quad A_1(N) = \Delta_N(\rho_0 + N^{-1}a_1 + O(N^{-2})),$$

$$(4.26) \quad a_1 = \rho_0 \left(\frac{3\kappa_4}{\sigma^4} - \frac{15\kappa_3^2}{2\sigma^6} \right) + \rho_1 \frac{3\kappa_3}{\sigma^4} - \rho_2 \frac{1}{\sigma^2}$$

by Lemma 4.3. Similarly,

$$A_2(N) = \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-1} M'(z) L(z, \dots, z) e^{-Nz} dz$$

since $f'(e^z)e^z = M'(z)$. We have

$$\begin{aligned} M(z)^{-1} M'(z) L(z, \dots, z) &= K'(z) L(z, \dots, z) \\ &= \rho_0 + (\sigma^2 \rho_0 + \rho_1)(z-\beta) + (3\kappa_3 \rho_0 + \sigma^2 \rho_1 + \rho_2)(z-\beta)^2 + \dots \end{aligned}$$

by (4.15) and (4.23), and hence

$$(4.27) \quad A_2(N) = \Delta_N(\rho_0 + N^{-1}a_2 + O(N^{-2})), \quad a_2 = a_1 - \rho_1$$

by Lemma 4.3 and (4.26). Further,

$$A_{1p}(N) = \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^N D_p L(z, \dots, z) e^{-Nz} dz$$

since $D_p L(s_1, \dots, s_d) = D_p g(e^{s_1}, \dots, e^{s_d}) e^{s_p}$. Noting that $D_p L(\beta, \dots, \beta) = \rho_0 \mu_p$, we get

$$(4.28) \quad A_{1p}(N) = \Delta_N(\rho_0 \mu_p + O(N^{-1})).$$

Using (4.25), (4.27), (4.28) in (4.22), we obtain

$$C_p = (j_p + y_p)(1 + N^{-1} \rho_0^{-1}(a_2 - a_1) + O(N^{-2})) + \mu_p + O(N^{-1}) \text{ uniformly,}$$

from which (4.20) follows since $\rho_0^{-1}(a_2 - a_1) = -\sum_{l=1}^d \mu_l$. Note that j_p is at most N and y_p is bounded. Since $b_p^{(N)}(x) = C_p - j_p$, we get (4.21) from (4.20),

Lemma 4.6.

$$(4.29) \quad C_{pq} = j_p j_q - N^{-1} j_p j_q (\sigma^2 + 2 \sum_{l=1}^d \mu_l) + j_p \sum_{l=1}^d x_l \alpha_{lq} + j_q \sum_{l=1}^d x_l \alpha_{lp} \\ + j_p \mu_q + j_q \mu_p + O(1) \quad \text{for } p \neq q,$$

$$(4.30) \quad C_{pp} = j_p^2 + j_p (\sigma^2 - 1 + 2 \mu_p + 2 \sum_{l=1}^d x_l \alpha_{lp} - x_p (\sigma^2 + 2 \sum_{l=1}^d \mu_l)) + O(1),$$

$$(4.31) \quad a_{pq}^{(N)}(x) = -\sigma^2 x_p x_q + O(N^{-1}) \quad \text{for } p \neq q,$$

$$(4.32) \quad a_{pp}^{(N)}(x) = \sigma^2 x_p (1 - x_p) + O(N^{-1}).$$

All O signs here are uniform in $j \in \mathbf{J}(N)$.

Proof. We can prove

$$(4.33) \quad C_{pq} = \{(j_p + y_p)(j_q + y_q) - \sum_{l=1}^d j_l \alpha_{lp}^{(N)} \alpha_{lq}^{(N)}\} \frac{A_3(N)}{A_1(N)} + \sum_{l=1}^d j_l \alpha_{lp}^{(N)} \alpha_{lq}^{(N)} \frac{A_4(N)}{A_1(N)} \\ + (j_p + y_p) \frac{A_{2q}(N)}{A_1(N)} + (j_q + y_q) \frac{A_{2p}(N)}{A_1(N)} + \frac{A_{1pq}(N)}{A_1(N)}.$$

where $p=q$ is not excluded. In fact,

$$C_{pq} = A_1(N)^{-1} (\text{coefficient of } w^N \text{ in } D_{pq} \Theta(w, 1, \dots, 1)),$$

and $D_{pq} \Theta(w, 1, \dots, 1)$ is obtained as follows:

$$D_{pq} \Psi(w, s_1, \dots, s_d) = \sum_{l,m} D_{lm} \Phi(w, \sum_n \alpha_{ln}^{(N)} s_n, \dots, \sum_n \alpha_{dn}^{(N)} s_n) \alpha_{lp}^{(N)} \alpha_{mq}^{(N)}, \\ D_{lm} \Phi(w, s_1, \dots, s_d) = f(ws_1)^{j_1} \dots \{j_l f(ws_l)^{j_l-1} f'(ws_l) w\} \dots \\ \{j_m f(ws_m)^{j_m-1} f'(ws_m) w\} \dots f(ws_d)^{j_d} \quad \text{if } l < m, \\ D_{ll} \Phi(w, s_1, \dots, s_d) = f(ws_1)^{j_1} \dots \{(j_l)_2 f(ws_l)^{j_l-2} f'(ws_l)^2 w^2 \\ + j_l f(ws_l)^{j_l-1} f''(ws_l) w^2\} \dots f(ws_d)^{j_d},$$

and hence

$$D_{lm} \Phi(w, 1, \dots, 1) = j_l j_m h_3(w) \quad \text{if } l \neq m \\ D_{ll} \Phi(w, 1, \dots, 1) = (j_l)_2 h_3(w) + j_l h_4(w).$$

Thus,

$$D_{pq} \Psi(w, 1, \dots, 1) = \{ \sum_{l,m} j_l j_m \alpha_{lp}^{(N)} \alpha_{mq}^{(N)} - \sum_l j_l \alpha_{lp}^{(N)} \alpha_{lq}^{(N)} \} h_3(w) \\ + \sum_l j_l \alpha_{lp}^{(N)} \alpha_{lq}^{(N)} h_4(w).$$

Since

$$\begin{aligned}
D_{pq}\Theta(w, 1, \dots, 1) &= D_{pq}\Psi(w, 1, \dots, 1)g(w, \dots, w) \\
&+ D_p\Psi(w, 1, \dots, 1)D_qg(w, \dots, w)w + D_q\Psi(w, 1, \dots, 1)D_pg(w, \dots, w)w \\
&+ \Psi(w, 1, \dots, 1)D_{pq}g(w, \dots, w)w^2
\end{aligned}$$

and the terms except the first in the right-hand side are treated in the proof of Lemma 4.5, (4.33) follows.

Having

$$(4.34) \quad \sum_i j_i \alpha_{ip}^{(N)} \alpha_{iq}^{(N)} = \begin{cases} O(1) & \text{if } p \neq q \\ j_p + O(1) & \text{if } p = q \end{cases}$$

in mind, let us estimate the right-hand side of (4.33). We have

$$A_3(N) = \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-2} M'(z)^2 L(z, \dots, z) e^{-Nz} dz$$

and, in a neighborhood of β ,

$$\begin{aligned}
M(z)^{-2} M'(z)^2 L(z, \dots, z) &= K'(z)^2 L(z, \dots, z) \\
&= \rho_0 + (2\sigma^2 \rho_0 + \rho_1)(z - \beta) + (6\kappa_3 \rho_0 + \sigma^4 \rho_0 + 2\sigma^2 \rho_1 + \rho_2)(z - \beta)^2 + \dots
\end{aligned}$$

Hence

$$(4.35) \quad A_3(N) = \Delta_N(\rho_0 + N^{-1}a_3 + O(N^{-2})), \quad a_3 = a_1 - \sigma^2 \rho_0 - 2\rho_1$$

by Lemma 4.3 and (4.26). Next,

$$A_4(N) = \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-1} (M''(z) - M'(z)) L(z, \dots, z) e^{-Nz} dz$$

by $f''(e^z)e^{2z} = M''(z) - M'(z)$, and

$$\begin{aligned}
M(z)^{-1} (M''(z) - M'(z)) L(z, \dots, z) \\
= (K'(z)^2 + K''(z) - K'(z)) L(z, \dots, z) = \sigma^2 \rho_0
\end{aligned}$$

at $z = \beta$. Hence

$$(4.36) \quad A_4(N) = \Delta_N(\sigma^2 \rho_0 + O(N^{-1})).$$

Also $A_{2p}(N)$ and $A_{1pq}(N)$ have analogous expression by Lemma 4.4.

Thus

$$\begin{aligned}
(4.37) \quad A_{2p}(N) &= \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-1} M'(z) D_p L(z, \dots, z) e^{-Nz} dz \\
&= \Delta_N(\rho_0 \mu_p + O(N^{-1}))
\end{aligned}$$

since $M(z)^{-1} M'(z) D_p L(z, \dots, z) = K'(z) D_p L(z, \dots, z) = \rho_0 \mu_p$ at $z = \beta$, and

$$(4.38) \quad A_{1pq}(N) = \Delta_N O(1).$$

It follows from (4.25), (4.33)—(4.38) that

$$\begin{aligned} C_{pq} &= (j_p j_q + j_p y_q + j_q y_p)(1 - N^{-1}(\sigma^2 + 2\rho_0^{-1}\rho_1)) + j_p \mu_q + j_q \mu_p + O(1) \\ &\quad \text{for } p \neq q, \\ C_{pp} &= (j_p^2 + 2j_p y_p - j_p)(1 - N^{-1}(\sigma^2 + 2\rho_0^{-1}\rho_1)) + j_p \sigma^2 + 2j_p \mu_p + O(1). \end{aligned}$$

Hence we get (4.29), (4.30) by (4.24). Since

$$\begin{aligned} a_{pq}^{(N)}(x) &= N^{-1}(C_{pq} - j_p C_q - j_q C_p + j_p j_q) \quad \text{for } p \neq q, \\ a_{pp}^{(N)}(x) &= N^{-1}(C_{pp} + C_p - 2j_p C_p + j_p^2), \end{aligned}$$

(4.31) and (4.32) follow from (4.20), (4.29), (4.30).

Lemma 4.7.

$$(4.39) \quad C_{ppp} = j_p^3 + 3j_p^2(\sigma^2 - 1 + \mu_p + \sum_{l=1}^d x_l \alpha_{lp} - x_p(\sigma^2 + \sum_{l=1}^d \mu_l)) + O(N)$$

uniformly in $J(N)$.

Proof. Clearly

$$(4.40) \quad C_{ppp} = U_1 + 3U_2 + 3U_3 + U_4$$

where U_1, U_2, U_3 and U_4 are $A_1(N)^{-1}$ times coefficients of w^N in $D_{ppp} \Psi(w, 1, \dots, 1)g(w, \dots, w)$, $D_{pp} \Psi(w, 1, \dots, 1)D_p g(w, \dots, w)w$, $D_p \Psi(w, 1, \dots, 1)D_p g(w, \dots, w)w^2$, and $\Psi(w, 1, \dots, 1)D_{ppp} g(w, \dots, w)w^3$, respectively. We have

$$\begin{aligned} D_{ppp} \Psi(w, s_1, \dots, s_d) &= \sum_{l, m, n} D_{lmn} \Phi(w, \sum_r \alpha_{lr}^{(N)} s_r, \dots, \sum_r \alpha_{dr}^{(N)} s_r) \\ &\quad \alpha_{lp}^{(N)} \alpha_{mp}^{(N)} \alpha_{np}^{(N)}, \end{aligned}$$

$$\begin{aligned} D_{lmn} \Phi(w, 1, \dots, 1) &= j_l j_m j_n h_5(w) \quad \text{if } l, m, n \text{ are all different,} \\ D_{lln} \Phi(w, 1, \dots, 1) &= (j_l^2 j_n - j_l j_n) h_5(w) + j_l j_n h_6(w) \quad \text{if } l \neq n \end{aligned}$$

and

$$D_{lll} \Phi(w, 1, \dots, 1) = (j_l^3 - 3j_l^2 + 2j_l) h_5(w) + 3(j_l^2 - j_l) h_6(w) + j_l h_7(w),$$

because

$$\begin{aligned} D_{lll} \Phi(w, s_1, \dots, s_d) &= f(ws_1)^{j_1} \dots \{(j_l)_3 f(ws_l)^{j_l-3} f'(ws_l)^3 w^3 \\ &\quad + 3(j_l)_2 f(ws_l)^{j_l-2} f'(ws_l) f''(ws_l) w^3 + j_l f(ws_l)^{j_l-1} f'''(ws_l) w^3\} \dots f(ws_d)^{j_d}. \end{aligned}$$

Hence,

$$\begin{aligned} U_1 &= A_1(N)^{-1} (\text{coefficient of } w^N \text{ in } \sum_{l, m, n} D_{lmn} \Phi(w, 1, \dots, 1) \\ &\quad g(w, \dots, w) \alpha_{lp}^{(N)} \alpha_{mp}^{(N)} \alpha_{np}^{(N)}) \\ &= (S_1 - 3S_2 + 2S_3) \frac{A_5(N)}{A_1(N)} + (3S_2 - 3S_3) \frac{A_6(N)}{A_1(N)} + S_3 \frac{A_7(N)}{A_1(N)}, \end{aligned}$$

where

$$\begin{aligned} S_1 &= \sum_{l, m, n} j_l j_m j_n \alpha_{lp}^{(N)} \alpha_{mp}^{(N)} \alpha_{np}^{(N)} = (j_p + y_p)^3 = j_p^3 + 3j_p^2 y_p + O(N), \\ S_2 &= \sum_{l, n} j_l j_n (\alpha_{lp}^{(N)})^2 \alpha_{np}^{(N)} = j_p^2 + O(N), \\ S_3 &= \sum_l j_l (\alpha_{lp}^{(N)})^3 = O(N). \end{aligned}$$

As in the proof of Lemmas 4.5 and 4.6, U_2 , U_3 and U_4 are expressed as follows:

$$\begin{aligned} U_2 &= \{(j_p + y_p)^2 - \sum_l j_l (\alpha_{lp}^{(N)})^2\} \frac{A_{3p}(N)}{A_1(N)} + \sum_l j_l (\alpha_{lp}^{(N)})^2 \frac{A_{4p}(N)}{A_1(N)}, \\ U_3 &= (j_p + y_p) \frac{A_{2pp}(N)}{A_1(N)}, \quad U_4 = \frac{A_{1ppp}(N)}{A_1(N)}. \end{aligned}$$

Hence

$$(4.41) \quad C_{ppp} = (j_p^3 + 3j_p^2 y_p - 3j_p^2) \frac{A_5(N)}{A_1(N)} + 3j_p^2 \frac{A_6(N)}{A_1(N)} + 3j_p^2 \frac{A_{3p}(N)}{A_1(N)} + O(N)$$

by (4.40) and Lemma 4.4. We have

$$\begin{aligned} (4.42) \quad A_5(N) &= \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-3} M'(z)^3 L(z, \dots, z) e^{-Nz} dz \\ &= \Delta_N(\rho_0 + N^{-1} a_5 + O(N^{-2})), \quad a_5 = a_1 - 3\sigma^2 \rho_0 - 3\rho_1 \end{aligned}$$

by Lemma 4.3 and (4.26), since

$$\begin{aligned} M^{-3} M'^3 L = K'^3 L &= \rho_0 + (3\sigma^2 \rho_0 + \rho_1)(z - \beta) + (3\sigma^4 \rho_0 + 9\kappa_3 \rho_0 \\ &\quad + 3\sigma^2 \rho_1 + \rho_2)(z - \beta)^2 + \dots \end{aligned}$$

in a neighborhood of β . Similarly

$$\begin{aligned} (4.43) \quad A_6(N) &= \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-2} M'(z)(M''(z) - M'(z)) L(z, \dots, z) e^{-Nz} dz \\ &= \Delta_N(\sigma^2 \rho_0 + O(N^{-1})) \end{aligned}$$

since $M^{-2} M'(M'' - M') L = K'(K'^2 + K'' - K') L = \sigma^2 \rho_0$ at $z = \beta$, and

$$\begin{aligned} (4.44) \quad A_{3p}(N) &= \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-2} M'(z)^2 D_p L(z, \dots, z) e^{-Nz} dz \\ &= \Delta_N(\rho_0 \mu_p + O(N^{-1})) \end{aligned}$$

as in (4.37). It follows from (4.25), (4.41)–(4.44) that

$$C_{ppp} = (j_p^3 + 3j_p^2 y_p - 3j_p^2)(1 - 3N^{-1}(\sigma^2 + \rho_0^{-1} \rho_1)) + 3j_p^2 \sigma^2 + 3j_p^2 \mu_p + O(N),$$

which is (4.39).

Lemma 4.8.

$$(4.45) \quad C_{pppp} = j_p^4 + 2j_p^3(3\sigma^2 - 3 + 2\mu_p + 2 \sum_{l=1}^d x_l \alpha_{lp} - x_p(3\sigma^2 + 2 \sum_{l=1}^d \mu_l)) + O(N^2)$$

uniformly in $j \in I(N)$.

Proof. This time we have

$$C_{pppp} = V_1 + 4V_2 + 6V_3 + 4V_4 + V_5,$$

where V_1, \dots, V_5 are $A_1(N)^{-1}$ times coefficients of w^N in $D_{pppp}\Psi(w, 1, \dots, 1)g(w, \dots, w)$, $D_{ppp}\Psi(w, 1, \dots, 1)D_{pg}(w, \dots, w)w$, $D_{pp}\Psi(w, 1, \dots, 1)D_{pg}(w, \dots, w)w^2$, $D_p\Psi(w, 1, \dots, 1)D_{ppg}(w, \dots, w)w^3$, $\Psi(w, 1, \dots, 1)D_{pppg}(w, \dots, w)w^4$, respectively. The most involved part is

$$(4.46) \quad D_{pppp}\Psi(w, 1, \dots, 1) = \sum_{l, m, n, r} D_{lmnr}\Phi(w, 1, \dots, 1) \alpha_{lp}^{(N)} \alpha_{mp}^{(N)} \alpha_{np}^{(N)} \alpha_{rp}^{(N)}$$

and we have

$$\begin{aligned} D_{lmnr}\Phi(w, 1, \dots, 1) &= j_l j_m j_n j_r h_8(w) \quad \text{if } l, m, n, r, \text{ are all different,} \\ D_{ltnr}\Phi(w, 1, \dots, 1) &= (j_l^2 j_n j_r - j_l j_n j_r) h_8(w) + j_l j_n j_r h_9(w) \\ &\quad \text{if } l \neq n \neq r \neq l, \\ D_{ltnn}\Phi(w, 1, \dots, 1) &= (j_l^2 j_n^2 - j_l^2 j_n - j_l j_n^2 + j_l j_n) h_8(w) \\ &\quad + (j_l^2 j_n + j_l j_n^2 - 2j_l j_n) h_9(w) + j_l j_n h_{10}(w) \quad \text{if } l \neq n, \\ D_{lltr}\Phi(w, 1, \dots, 1) &= (j_l^3 j_r - 3j_l^2 j_r + 2j_l j_r) h_8(w) + 3(j_l^2 j_r - j_l j_r) h_9(w) \\ &\quad + j_l j_r h_{11}(w) \quad \text{if } l \neq r, \\ D_{lllt}\Phi(w, 1, \dots, 1) &= (j_l^4 - 6j_l^3 + 11j_l^2 - 6j_l) h_8(w) + 6(j_l^3 - 3j_l^2 + 2j_l) h_9(w) \\ &\quad + 3(j_l^2 - j_l) h_{10}(w) + 4(j_l^2 - j_l) h_{11}(w) + j_l h_{12}(w). \end{aligned}$$

The last one comes from the expression

$$\begin{aligned} D_{lllt}\Phi(w, s_1, \dots, s_d) &= f(ws_1)^{j_1} \dots \{(j_l)_4 f(ws_l)^{j_l-4} f'(ws_l)^4 w^4 \\ &\quad + 6(j_l)_3 f(ws_l)^{j_l-3} f'(ws_l)^2 f''(ws_l) w^4 + 3(j_l)_2 f(ws_l)^{j_l-2} f''(ws_l)^2 w^4 \\ &\quad + 4(j_l)_2 f(ws_l)^{j_l-2} f'(ws_l) f'''(ws_l) w^4 + j_l f(ws_l)^{j_l-1} f''''(ws_l) w^4\} \dots f(ws_d)^{j_d} \end{aligned}$$

Let $\sum^{(1)}$, $\sum_{(l=m)}^{(2)}$, $\sum_{(l=m, n=r)}^{(3)}$, $\sum_{(l=m=n)}^{(4)}$, $\sum^{(5)}$ denote the summations over $\{(l, m, n, r); l, m, n, r \text{ are different from each other}\}$, $\{(l, m, n, r); l=m \neq n \neq r \neq l\}$, $\{(l, m, n, r); l=m \neq n=r\}$, $\{(l, m, n, r); l=m=n \neq r\}$, $\{(l, m, n, r); l=m=n=r\}$, respectively. Decompose the right-hand side of (4.46) so that

$$\begin{aligned} \sum_{l, m, n, r} &= \sum^{(1)} + (\sum_{(l=m)}^{(2)} + \sum_{(l=n)}^{(2)} + \sum_{(l=r)}^{(2)} + \sum_{(m=n)}^{(2)} + \sum_{(m=r)}^{(2)} + \sum_{(n=r)}^{(2)}) \\ &\quad + (\sum_{(l=m, n=r)}^{(3)} + \sum_{(l=n, m=r)}^{(3)} + \sum_{(l=r, m=n)}^{(3)}) \\ &\quad + (\sum_{(l=m=n)}^{(4)} + \sum_{(l=m=r)}^{(4)} + \sum_{(l=n=r)}^{(4)} + \sum_{(m=n=r)}^{(4)}) + \sum^{(5)}, \end{aligned}$$

and use the above formulas. Then, by Lemma 4.4, we get

$$\begin{aligned} V_1 &= (\sum_i j_i \alpha_{i_p}^{(N)})^4 \frac{A_8(N)}{A_1(N)} + 6(\sum_i j_i (\alpha_{i_p}^{(N)})^2) (\sum_i j_i \alpha_{i_p}^{(N)})^2 \\ &\quad \frac{A_9(N) - A_8(N)}{A_1(N)} + O(N^2) \\ &= (j_p^4 + 4j_p^3 y_p - 6j_p^3) \frac{A_8(N)}{A_1(N)} + 6j_p^3 \frac{A_9(N)}{A_1(N)} + O(N^2) \end{aligned}$$

after some reflection. By argument quite similar to the proof of Lemmas 4.5, 4.6, 4.7, we get

$$\begin{aligned} V_2 &= (S_1 - 3S_2 + 2S_3) \frac{A_{5p}(N)}{A_1(N)} + (3S_2 - 3S_3) \frac{A_{6p}(N)}{A_1(N)} + S_3 \frac{A_{7p}(N)}{A_1(N)} \\ &= j_p^3 \frac{A_{5p}(N)}{A_1(N)} + O(N^2), \\ V_3 &= O(N^2), \quad V_4 = O(N), \quad V_5 = O(1). \end{aligned}$$

Hence,

$$(4.47) \quad C_{pppp} = (j_p^4 + 4j_p^3 y_p - 6j_p^3) \frac{A_8(N)}{A_1(N)} + 6j_p^3 \frac{A_9(N)}{A_1(N)} + 4j_p^3 \frac{A_{5p}(N)}{A_1(N)} + O(N^2).$$

We have

$$\begin{aligned} (4.48) \quad A_8(N) &= \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-4} M'(z)^4 L(z, \dots, z) e^{-Nz} dz \\ &= \Delta_N(\rho_0 + N^{-1}a_8 + O(N^{-2})), \quad a_8 = a_1 - 6\sigma^2\rho_0 - 4\rho_1 \end{aligned}$$

since

$$\begin{aligned} M^{-4}M'^4 L &= K'^4 L = \rho_0 + (4\sigma^2\rho_0 + \rho_1)(z - \beta) + (12\kappa_3\rho_0 + 6\sigma^4\rho_0 + 4\sigma^2\rho_1 \\ &\quad + \rho_2)(z - \beta)^2 + \dots \end{aligned}$$

in a neighborhood of β ,

$$\begin{aligned} (4.49) \quad A_9(N) &= \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-3} M'(z)^2 (M''(z) - M'(z)) L(z, \dots, z) e^{-Nz} dz \\ &= \Delta_N(\sigma^2\rho_0 + O(N^{-1})) \end{aligned}$$

since $M^{-3}M'^2(M'' - M')L = K'^2(K'^2 + K'' - K')L = \sigma^2\rho_0$ at β , and

$$\begin{aligned} (4.50) \quad A_{5p}(N) &= \frac{1}{2\pi i} \int_{\beta-i\pi}^{\beta+i\pi} M(z)^{N-3} M'(z)^3 D_p L(z, \dots, z) e^{-Nz} dz \\ &= \Delta_N(\rho_0 \mu_p + O(N^{-1})) \end{aligned}$$

as in (4.37). By (4.25), (4.47)–(4.50) we have

$$C_{pppp} = (j_p^4 + 4j_p^3 y_p - 6j_p^3)(1 - N^{-1}(6\sigma^2 + 4\rho_0^{-1}\rho_1)) + 6j_p^3 \sigma^2 + 4j_p^3 \mu_p + O(N^2),$$

and the proof of (4.45) is complete.

Lemma 4.9.

$$(4.51) \quad e_p^{(N)}(x) = O(N^{-1}) \quad \text{uniformly in } j \in J(N).$$

Proof. We have

$$\begin{aligned} e_p^{(N)}(x) &= N^{-3}(C_{pppp}^* - 4j_p C_{ppp}^* + 6j_p^2 C_{pp}^* - 4j_p^3 C_p^* + j_p^4) \\ &= N^{-3}\{C_{pppp} + (-4j_p + 6)C_{ppp} + (6j_p^2 - 12j_p + 7)C_{pp} \\ &\quad + (-4j_p^3 + 6j_p^2 - 4j_p + 1)C_p + j_p^4\} \end{aligned}$$

by Lemma 4.2. Use Lemmas 4.5—4.8 and substitute (4.20), (4.30), (4.39), (4.45) for C_p , C_{pp} , C_{ppp} , C_{pppp} . Then, all terms cancel except terms of magnitude $O(N^{-1})$, and (4.51) follows.

Now Lemmas 4.5, 4.6 and 4.9 say that (4.7), (4.8) and (4.9) hold. Hence the proof of Theorems 2.1 and 2.2 is complete.

5. Uniqueness of the solution of the martingale problem and proof of Theorems 2.3 and 2.4

All notations in this section are the same as in Section 2. By Theorems 2.1 and 2.2, convergence of the interpolated normalization of the induced Markov chains is proved if the solution of the martingale problem (\mathbf{K}, a, b, x) is unique. This uniqueness problem is a problem on some stochastic differential equations, as the following lemma says.

Lemma 5.1. *Let $c(x) = (c_{pq}(x))$ be a bounded, Borel measurable $(d-1) \times (d-1)$ -matrix on \mathbf{K} satisfying*

$$(5.1) \quad c(x)c(x)' = a(x),$$

where $c(x)'$ is the transpose of $c(x)$. Consider a stochastic differential equation

$$(5.2) \quad \begin{cases} dX_p(t) = \sum_{q=1}^{d-1} c_{pq}(X(t)) dB_q(t) + b_p(X(t)) dt, & p = 1, \dots, d-1 \\ X(0) = x \end{cases}$$

where $X(t) = (X_1(t), \dots, X_{d-1}(t))$ is a process taking values in \mathbf{K} and $B(t) = (B_1(t), \dots, B_{d-1}(t))$ is a $(d-1)$ -dimensional Brownian motion. Then, uniqueness of the solution of the martingale problem (\mathbf{K}, a, b, x) is equivalent to uniqueness, in the sense of probability measure induced in (Ω, \mathcal{M}) , of the solution of (5.2).

Proof. This is proved essentially by Stroock and Varadhan [13]. Priouret [10] contains a detailed account.

We choose $c(x) = (c_{pq}(x))_{p,q=1,\dots,d-1}$ as a lower triangular matrix. Let K^0 and ∂K be the interior and the boundary of K and let, for $x \in K^0$,

$$\begin{aligned}
 c_{pq}(x) &= 0 \quad \text{for } q > p \\
 c_{11}(x) &= \sigma x_1^{1/2} (1-x_1)^{1/2}, \\
 c_{21}(x) &= -\sigma x_2 x_1^{1/2} (1-x_1)^{-1/2}, \quad c_{22}(x) = \sigma x_2^{1/2} (1-x_1-x_2)^{1/2} (1-x_1)^{-1/2}, \\
 c_{31}(x) &= -\sigma x_3 x_1^{1/2} (1-x_1)^{-1/2}, \quad c_{32}(x) = -\sigma x_3 x_2^{1/2} (1-x_1)^{-1/2} (1-x_1-x_2)^{-1/2}, \\
 (5.3) \quad c_{33}(x) &= \sigma x_3^{1/2} (1-x_1-x_2-x_3)^{1/2} (1-x_1-x_2)^{-1/2}, \\
 c_{41}(x) &= -\sigma x_4 x_1^{1/2} (1-x_1)^{-1/2}, \quad c_{42}(x) = -\sigma x_4 x_2^{1/2} (1-x_1)^{-1/2} (1-x_1-x_2)^{-1/2}, \\
 c_{43}(x) &= -\sigma x_4 x_3^{1/2} (1-x_1-x_2)^{-1/2} (1-x_1-x_2-x_3)^{-1/2}, \\
 c_{44}(x) &= \sigma x_4^{1/2} (1-x_1-x_2-x_3-x_4)^{1/2} (1-x_1-x_2-x_3)^{-1/2}, \\
 &\dots\dots\dots
 \end{aligned}$$

Lemma 5.2. $c(x)$ defined above on K^0 extends continuously to K . Denote the extension by the same $c(x)$. Then

$$(5.4) \quad a_{pq}(x) = \sum_{r=1}^{p \wedge q} c_{pr}(x) c_{qr}(x),$$

that is, (5.1) holds.

Proof. Let $x \in \partial K$. We define $c_{pq}(x)$ by the formulas in (5.3) when negative powers of 0 do not appear, and define $c_{pq}(x) = 0$ when negative powers of 0 appear. Then all c_{pq} are continuous on K , since we have

$$\begin{aligned}
 (5.5) \quad 0 &\geq c_{21}(x) \geq -\sigma x_1^{1/2} (1-x_1)^{1/2}, \quad 0 \leq c_{22}(x) \leq \sigma (1-x_1)^{1/2}, \\
 0 &\geq c_{31}(x) \geq -\sigma x_1^{1/2} (1-x_1)^{1/2}, \quad 0 \geq c_{32}(x) \geq -\sigma (1-x_1-x_2)^{1/2}, \\
 0 &\leq c_{33}(x) \leq \sigma (1-x_1-x_2)^{1/2}, \\
 &\dots\dots\dots
 \end{aligned}$$

for $x \in K^0$. In order to show (5.4), we may assume $x \in K^0$. Let

$$1-x_1 = y_1, \quad 1-x_1-x_2 = y_2, \quad 1-x_1-x_2-x_3 = y_3, \dots$$

Then,

$$\begin{aligned}
 c_{11} &= \sigma (x_1 y_1)^{1/2}, \\
 c_{21} &= -\sigma x_2 \left(\frac{x_1}{y_1} \right)^{1/2} = -\sigma x_2 \left(\frac{1}{y_1} - 1 \right)^{1/2}, \quad c_{22} = \sigma \left(\frac{x_2 y_2}{y_1} \right)^{1/2}, \\
 c_{31} &= -\sigma x_3 \left(\frac{x_1}{y_1} \right)^{1/2} = -\sigma x_3 \left(\frac{1}{y_1} - 1 \right)^{1/2}, \\
 c_{32} &= -\sigma x_3 \left(\frac{x_2}{y_1 y_2} \right)^{1/2} = -\sigma x_3 \left(\frac{1}{y_2} - \frac{1}{y_1} \right)^{1/2}, \quad c_{33} = \sigma \left(\frac{x_3 y_3}{y_2} \right)^{1/2}, \\
 &\dots\dots\dots
 \end{aligned}$$

and (5.4) is obtained by direct calculation.

Henceforth $c(x)$ denotes the one defined by (5.3) and Lemma 5.2.

Lemma 5.3. *Let $d-1=1$. Then the solution of the stochastic differential equation (5.2) is pathwise unique. That is, if $X(t)$ and $\tilde{X}(t)$ are solutions of (5.2) with a common one-dimensional Brownian motion $B(t)$, then $X(t)=\tilde{X}(t)$ almost surely.*

Proof. In this case (5.2) is a single stochastic differential equation

$$dX(t) = c_{11}(X(t))dB(t) + b_1(X(t))dt, \quad X(0) = x.$$

c_{11} is extended on \mathbf{R}^1 to a Hölder continuous function with exponent $1/2$ and b_1 is extended to a Lipschitz continuous function on \mathbf{R}^1 . Hence the theorem of Yamada and S. Watanabe [15] applies.

Lemma 5.4. *If $x, y \in K$ and $x_l = y_l$ for $l=1, \dots, p-1$, then*

$$(5.6) \quad |c_{pq}(x) - c_{pq}(y)| \leq 2\sigma |x_p - y_p|^{1/2}.$$

Proof. Let $q=1 < p$. If $1-\varepsilon \leq x_1 \leq 1$, then $0 \geq c_{p1}(x) \geq -\sigma\varepsilon^{1/2}$ and $0 \geq c_{p1}(y) \geq -\sigma\varepsilon^{1/2}$ by (5.5). If $1-\varepsilon > x_1 \geq 0$, then

$$|c_{p1}(x) - c_{p1}(y)| = \sigma |x_p - y_p| x_1^{1/2} (1-x_1)^{-1/2} \leq \sigma \varepsilon^{-1/2} |x_p - y_p|.$$

Hence

$$|c_{p1}(x) - c_{p1}(y)| \leq \sigma(\varepsilon^{-1/2} |x_p - y_p| + \varepsilon^{1/2})$$

in any case. Since ε is arbitrary we may choose $\varepsilon = |x_p - y_p|^{1/2}$ and get (5.6) for $q=1 < p$. The proof for $1 < q < p$ is similar. In case $p=q$, we let $1-x_1-\dots-x_{p-1}=1-y_1-\dots-y_{p-1}=\xi$ and have

$$\begin{aligned} |c_{pp}(x) - c_{pp}(y)| &\leq \sigma \xi^{-1/2} (|x_p^{1/2}(\xi - x_p)^{1/2} - y_p^{1/2}(\xi - x_p)^{1/2}| \\ &\quad + |y_p^{1/2}(\xi - x_p)^{1/2} - y_p^{1/2}(\xi - y_p)^{1/2}|) \\ &\leq \sigma \xi^{1/2} ((\xi - x_p)^{1/2} |x_p - y_p|^{1/2} + y_p^{1/2} |x_p - y_p|^{1/2}) \leq 2\sigma |x_p - y_p|^{1/2}, \end{aligned}$$

since

$$|\eta^{1/2} - \zeta^{1/2}| \leq |\eta - \zeta|^{1/2}.$$

Lemma 5.5. *Let $d-1 > 1$ and suppose that $b(x)$ satisfies the condition in Theorem 2.4. Then the solution of the stochastic differential equation (5.2) is pathwise unique.*

Proof. By the assumption $b_p(x)$ is a linear function only of x_1, \dots, x_p . As is seen from (5.3), $c_{pq}(x)$ is also a function of x_1, \dots, x_p . Suppose that $X(t)=(X_1(t), \dots, X_{d-1}(t))$ and $\tilde{X}(t)=(\tilde{X}_1(t), \dots, \tilde{X}_{d-1}(t))$ are solutions of (5.2)

taking values in \mathbf{K} with a common Brownian motion $B(t)=(B_1(t), \dots, B_{d-1}(t))$. First, we get $X_1(t)=\tilde{X}_1(t)$ a.s. as in Lemma 5.3, since the equation for $X_1(t)$ does not involve $X_2(t), \dots, X_{d-1}(t)$. Let us prove $X_p(t)=\tilde{X}_p(t)$ a.s. for all p by induction. Suppose $X_l(t)=\tilde{X}_l(t)$ a.s. for $l=1, \dots, p-1$. The following proof is essentially the same as part of Yamada and S. Watanabe [15]. Choose nonnegative C^2 functions $\varphi_n(\xi)$ on \mathbf{R}^1 such that $\varphi_n(\xi)$ increases to $|\xi|$ as $n \rightarrow \infty$, $\varphi_n(\xi)=0$ in a neighborhood of 0, $\varphi_n'(\xi)$ is bounded, and $0 \leq \varphi_n''(\xi) \leq 2n^{-1}|\xi|^{-1}$. Since

$$X_p(t) - \tilde{X}_p(t) = \sum_{i=1}^p \int_0^t \{c_{pq}(X(s)) - c_{pq}(\tilde{X}(s))\} dB_q(s) \\ + \int_0^t \{b_p(X(s)) - b_p(\tilde{X}(s))\} ds,$$

we get

$$\varphi_n(X_p(t) - \tilde{X}_p(t)) = \text{stochastic integral} + \frac{1}{2} \int_0^t \varphi_n''(X_p(s) - \tilde{X}_p(s)) \\ \sum_{i=1}^p \{c_{pq}(X(s)) - c_{pq}(\tilde{X}(s))\}^2 ds + \int_0^t \varphi_n'(X_p(s) - \tilde{X}_p(s)) \{b_p(X(s)) - b_p(\tilde{X}(s))\} ds.$$

by the Ito formula. We have $\{c_{pq}(X(s)) - c_{pq}(\tilde{X}(s))\}^2 \leq 4\sigma^2 |X_p(s) - \tilde{X}_p(s)|$ by Lemma 5.4 and hence

$$E[\varphi_n(X_p(t) - \tilde{X}_p(t))] \leq \frac{1}{2} \int_0^t \frac{2}{n} 4p\sigma^2 ds + CE \left[\int_0^t |X_p(s) - \tilde{X}_p(s)| ds \right], \\ C = \text{const.}$$

Going to the limit as $n \rightarrow \infty$ we get

$$E|X_p(t) - \tilde{X}_p(t)| \leq C \int_0^t E|X_p(s) - \tilde{X}_p(s)| ds.$$

By iteration it follows that $E|X_p(t) - \tilde{X}_p(t)| = 0$. Hence $X_p(t) = \tilde{X}_p(t)$, a.s.

Proof of Theorems 2.3 and 2.4. Yamada and S. Watanabe [15] prove that pathwise uniqueness of the solution of a stochastic differential equation implies uniqueness in the sense of probability measure induced in (Ω, \mathcal{M}) . Hence Theorems 2.3 and 2.4 follow from Lemmas 5.3 and 5.5, respectively.

Properties of the limiting diffusions

Let $d-1=1$. We have proved that the martingale problem (\mathbf{K}, a, b, x) has a unique solution (Theorem 2.3). It is a one-dimensional diffusion process on $[0, 1]$ with backward Kolmogorov equation (2.8). Let $\lambda_1 = \alpha_{21} + \mu_1$ and $\lambda_2 = \alpha_{12} + \mu_2$. In Feller's boundary classification into four types (regular, pure exit, pure entrance, natural), the boundary 0 is pure exit, regular, pure entrance according as $\lambda_1=0$, $0 < \lambda_1 < \sigma^2/2$, $\lambda_1 \geq \sigma^2/2$, respectively. Similarly, the boundary 1 is pure exit, regular, pure entrance according as $\lambda_2=0$, $0 < \lambda_2 < \sigma^2/2$, $\lambda_2 \geq \sigma^2/2$,

respectively. The proof is a standard argument. In case the boundary is pure exist, it is a trap. This is a consequence of the general theory, but this is clear also from the uniqueness of the solution since in case $\lambda_1=0$ the process standing still at 0 is a solution. If $\lambda_1>0$, it is shown by the method of S. Watanabe [14], p. 459, that $E_x[\int_0^\infty \chi_{\{0\}}(x(t))dt]=0$ for every $x\in[0, 1]$, where $\chi_{\{0\}}$ is the indicator function of the set $\{0\}$. Therefore, in case the boundary is regular, the limiting diffusion has reflecting boundary condition there, while, in case of pure entrance boundary, the limiting diffusion starting from there immediately enters the interior and never returns.

Let $d-1=2$ and suppose that the uniqueness holds for the martingale problem (\mathbf{K}, a, b, x) for every x . Thus $(x(t), \mathcal{M}_t, P_x; x\in\mathbf{K})$ is a diffusion process on the triangular region $\mathbf{K}=\{(x_1, x_2); x_1\geq 0, x_2\geq 0, 1-x_1-x_2\geq 0\}$ and its backward equation is (2.9). Let us examine its boundary properties. Since all the three sides of \mathbf{K} are similar, we examine $\Gamma=\{(x_1, 0); 0<x_1<1\}$. Following Hasminsky [4] and S. Watanabe [14], we define regular and repulsive boundary points and unattainable and pure entrance boundary segments as follows. Let $x\in\Gamma$ and let U be a neighborhood in \mathbf{K} of x having positive distance with $\partial\mathbf{K}-\Gamma$. For $\eta>0$, let $U_\eta=U\cap\{(x_1, x_2); x_2>\eta\}$, $T_\eta(\omega)$ =first leaving time of U_η , and $T(\omega)=\lim_{\eta\downarrow 0} T_\eta(\omega)$. Let $\Gamma_\eta^x(\omega)$ be the set of all limit points of $x(t, \omega)$ when $t\uparrow T(\omega)$. x is called *regular* if, for every U and for every neighborhood V in \mathbf{K} of x , we have

$$\lim_{y\in\mathbf{K}^0, y\rightarrow x} P_y(\Gamma_\eta^x(\omega)\subset V\cap\Gamma)=1.$$

x is called *repulsive*, if, for some U and for some $\eta>0$, we have

$$\liminf_{y\in\mathbf{K}^0, y\rightarrow x} P_y(\Gamma_\eta^x(\omega)\subset(\partial U)_\eta)<1.$$

Here $(\partial U)_\eta=\partial U\cap\{y=(y_1, y_2); y_2<\eta\}$, ∂U being the boundary in \mathbf{R}^2 of U . Let Σ be an open interval in Γ . Σ is called *unattainable* if for every $x\in\Sigma$ there exists a neighborhood U in \mathbf{K} of x such that

$$P_y(\Gamma_\eta^x(\omega)\cap\Gamma \text{ is empty})=1 \quad \text{for every } y\in U\cap\mathbf{K}^0.$$

Σ is called a *pure entrance boundary segment* if Σ is unattainable and if every $x\in\Sigma$ has a neighborhood U such that $P_y(T_U<\infty)=1$ for every $y\in U$. Here T_U is the first leaving time of U .

Define Σ_1 and Σ_2 as follows:

$$\begin{aligned} \Sigma_1 &= \{(x_1, 0)\in\Gamma; x_1(\alpha_{12}-\alpha_{32})+\alpha_{32}+\mu_2<\sigma^2/2\}, \\ \Sigma_2 &= \{(x_1, 0)\in\Gamma; x_1(\alpha_{12}-\alpha_{32})+\alpha_{32}+\mu_2>\sigma^2/2\}. \end{aligned}$$

Σ_1 and Σ_2 are open intervals or empty.

Theorem 5.1. (i) *Every point in Σ_1 is regular. If $\alpha_{12}=\alpha_{32}=\mu_2=0$, then $\Sigma_1=\Gamma$ and, after hitting Γ , the process moves in $\{(x_1, 0); 0 \leq x_1 \leq 1\}$ following the backward equation*

$$\frac{\partial u}{\partial t} = \frac{\sigma^2}{2} x_1(1-x_1) \frac{\partial^2 u}{\partial x_1^2} + \{-x_1(\alpha_{13} + \mu_3) + (1-x_1)(\alpha_{31} + \mu_1)\} \frac{\partial u}{\partial x_1}$$

with the behavior at $(0, 0)$ and $(1, 0)$ described in the case $d-1=1$. If at least one of α_{12} , α_{32} and μ_2 is positive, then

$$(5.7) \quad \int_0^\infty \chi_{\Sigma_1}(x(t)) dt = 0 \quad a.s.$$

and, for any open subinterval Σ of Σ_1 , the process starting from K^0 hits Σ in a finite time with positive probability.

(ii) Σ_2 is a pure entrance boundary segment and every point in Σ_2 is repulsive.

Proof. Regularity of the points in Σ_1 , repulsiveness of the points in Σ_2 and unattainability of Σ_2 are applications of Hasminsky's tests [4]. The property described in (i) in case $\alpha_{12}=\alpha_{32}=\mu_2=0$ is a consequence of the assumed uniqueness of the solution of the martingale problem. The proof of the other properties is similar to the discussion by S. Watanabe [14] of two-dimensional diffusion processes with branching property. In proving (5.7) we use the expression

$$\begin{cases} dX_1(t) = c_{11}^*(X(t)) dB_1^*(t) + c_{12}^*(X(t)) dB_2^*(t) + b_1(X(t)) dt \\ dX_2(t) = c_{22}^*(X(t)) dB_2^*(t) + b_2(X(t)) dt \end{cases}$$

where $B^*(t) = (B_1^*(t), B_2^*(t))$ is a two-dimensional Brownian motion and

$$\begin{aligned} c_{11}^*(x) &= \sigma x_1^{1/2} (1-x_1-x_2)^{1/2} (1-x_2)^{-1/2}, & c_{12}^*(x) &= -\sigma x_1 x_2^{1/2} (1-x_2)^{-1/2}, \\ c_{22}^*(x) &= \sigma x_2^{1/2} (1-x_2)^{1/2}. \end{aligned}$$

TOKYO UNIVERSITY OF EDUCATION

References

- [1] P. Billingsley: Convergence of probability measures, John Wiley, New York, 1968.
- [2] J.F. Crow and M. Kimura: An introduction to population genetics theory, Harper and Row, New York, 1970.
- [3] W. Feller: Diffusion processes in genetics, Proc. Second Berkeley Symp. Math. Stat. Prob., University of California Press, Berkeley and Los Angeles, 1951, 227-246.

- [4] R.Z. Hasminsky: *Diffusion processes and elliptic differential equations degenerating at the boundary of the region*, Teor. Veroyatnost. Primenen. **3** (1958), 430–451. (Russian)
- [5] S. Karlin: *A first course in stochastic processes*, Academic Press, New York, 1966.
- [6] S. Karlin and J. McGregor: *On a genetics model of Moran*, Proc. Camb. Phil. Soc. **58** (1962), 299–311.
- [7] S. Karlin and J. McGregor: *Direct product branching processes and related Markov chains*, Proc. Nat. Acad. Sci. USA **51** (1964), 598–602.
- [8] M. Kimura: *Some problems of stochastic processes in genetics*, Ann. Math. Stat. **28** (1957), 882–901.
- [9] H.J. Kushner: *On the weak convergence of interpolated Markov chains to a diffusion*, Ann. Probability **2** (1974), 40–50.
- [10] P. Priouret: *Processus de diffusion et equations differentielles stochastiques*, Ecole d'Eté de Probabilités de Saint-Flour III-1973, Lecture Notes in Math. 390, Springer, 1974, 37–113.
- [11] K. Sato: *Asymptotic properties of eigenvalues of a class of Markov chains induced by direct product branching processes*, J. Math. Soc. Japan **28** (1976), 192–211.
- [12] D.W. Stroock and S.R.S. Varadhan: *Diffusion processes with continuous coefficients*, I, Comm. Pure Appl. Math. **22** (1969), 345–400; II, *ibid.*, 479–530.
- [13] D.W. Stroock and S.R.S. Varadhan: *On the support of diffusion processes with application to the strong maximum principle*, Proc. Sixth Berkeley Symp. Math. Stat. Prob., Vol. 3, University of California Press, Berkeley and Los Angeles, 1972, 333–359.
- [14] S. Watanabe: *On two dimensional Markov processes with branching property*, Trans. Amer. Math. Soc. **136** (1969), 447–466.
- [15] T. Yamada and S. Watanabe: *On the uniqueness of solutions of stochastic differential equations*, J. Math. Kyoto Univ. **11** (1971), 155–167.

Added in proof. S. N. Ethier (*A class of degenerate diffusion processes occurring in population genetics*, Comm. Pure Appl. Math. **29** (1976), 483–493) gives a proof to our conjecture on the uniqueness of the solution of the martingale problem (K, a, b, x) in general dimensions. Thus the assumption in our Theorem 2.2 is always satisfied.

