

Title	Deformable flat tori in $S^3$ with constant mean curvature
Author(s)	Kitagawa, Yoshihisa
Citation	Osaka Journal of Mathematics. 2003, 40(1), p. 103-119
Version Type	VoR
URL	<a href="https://doi.org/10.18910/6858">https://doi.org/10.18910/6858</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## DEFORMABLE FLAT TORI IN $S^3$ WITH CONSTANT MEAN CURVATURE

To the memory of Professor Shukichi Tanno

YOSHIHISA KITAGAWA

(Received August 20, 2001)

### 1. Introduction

In 1975, Yau [9, p. 87] posed the problem of the classification of flat tori in the unit 3-sphere  $S^3$ . Concerning this problem, the author established a method for constructing all the flat tori in  $S^3$  ([2]), and obtained some results on flat tori in  $S^3$  ([1], [3], [4], [5]). In this paper, using this method, we study isometric deformations of flat tori isometrically immersed in  $S^3$  with constant mean curvature, and we obtain the classification of undeformable flat tori in  $S^3$ .

For positive constants  $R_1$  and  $R_2$  satisfying  $R_1^2 + R_2^2 = 1$ , let  $F: \mathbb{R}^2 \rightarrow S^3$  be an isometric immersion given by

$$(1.1) \quad F(x_1, x_2) = \left( R_1 \cos \frac{x_1}{R_1}, R_1 \sin \frac{x_1}{R_1}, R_2 \cos \frac{x_2}{R_2}, R_2 \sin \frac{x_2}{R_2} \right),$$

and  $G_0$  a lattice of  $\mathbb{R}^2$  defined by

$$(1.2) \quad G_0 = \{(2\pi R_1 n_1, 2\pi R_2 n_2) : n_1, n_2 \in \mathbb{Z}\}.$$

If  $G$  is a lattice of  $\mathbb{R}^2$  such that  $G \subset G_0$ , then we obtain a flat torus  $\mathbb{R}^2/G$  and an isometric immersion

$$(1.3) \quad F/G: \mathbb{R}^2/G \rightarrow S^3$$

with constant mean curvature. Conversely, every flat torus isometrically immersed in  $S^3$  with constant mean curvature is obtained in this way. Note that the immersion  $F/G$  is the composition of the covering map  $\mathbb{R}^2/G \rightarrow \mathbb{R}^2/G_0$  and the embedding  $F/G_0: \mathbb{R}^2/G_0 \rightarrow S^3$ . In [3] the author studied isometric deformations of  $F/G_0$ , and proved the following theorem.

---

2000 *Mathematics Subject Classification* : Primary 53C42.

Partly supported by the Grant-in-Aid for Scientific Research, The Ministry of Education, Science, Sports and Culture, Japan, (No. 12640059).

**Theorem 1.1.** *If  $f_t: \mathbb{R}^2/G_0 \rightarrow S^3$ ,  $t \in \mathbb{R}$ , is a smooth one parameter family of isometric immersions with  $f_0 = F/G_0$ , then for each  $t \in \mathbb{R}$  there exists an isometry  $A_t: S^3 \rightarrow S^3$  such that  $f_t = A_t \circ f_0$ .*

This theorem says that every isometric deformation of  $F/G_0$  is trivial. On the other hand, there are many lattices  $G \subset G_0$  such that the immersion  $F/G$  is deformable. Let  $W_+(n)$  and  $W_-(n)$  be lattices of  $\mathbb{R}^2$  defined by

$$(1.4) \quad W_{\pm}(n) = \{(2\pi R_1 n_1, 2\pi R_2 n_2) : n_1 \pm n_2 \in n\mathbb{Z}\} \subset G_0.$$

Then we can show that if  $G \subset W_+(n)$  or  $G \subset W_-(n)$  for some integer  $n \geq 2$ , the immersion  $F/G$  is deformable (Theorem 3.1). Here, we give the following definition.

**DEFINITION.** For immersions  $f_1: M_1 \rightarrow S^3$  and  $f_2: M_2 \rightarrow S^3$ , we write  $f_1 \equiv f_2$ , and we say “ $f_1$  is congruent to  $f_2$ ” if there exist an isometry  $A: S^3 \rightarrow S^3$  and a diffeomorphism  $\rho: M_1 \rightarrow M_2$  such that  $A \circ f_1 = f_2 \circ \rho$ . An isometric immersion  $f: M \rightarrow S^3$  is said to be *deformable* if it admits an isometric deformation  $f_t: M \rightarrow S^3$  ( $t \in \mathbb{R}$ ,  $f_0 = f$ ) such that  $f_0 \not\equiv f_1$ .

The assertion of Theorem 3.1 leads us to the problem of finding all the lattices  $G \subset G_0$  such that the immersion  $F/G$  is deformable. In this paper we study this problem, and prove the following theorem.

**Theorem 1.2.** *Let  $G$  be a lattice of  $\mathbb{R}^2$  such that  $G \subset G_0$ . Then the immersion  $F/G$  is deformable if and only if there exists an integer  $n \geq 2$  such that  $G \subset W_+(n)$  or  $G \subset W_-(n)$ .*

Furthermore, as a corollary of this theorem, we obtain the following classification of undeformable flat tori isometrically immersed in  $S^3$ .

**Theorem 1.3.** *Let  $f: M \rightarrow S^3$  be an isometric immersion of a flat torus  $M$  into the unit sphere  $S^3$ . Then the following statements are equivalent.*

- (1) *Every isometric deformation of  $f$  is trivial.*
- (2) *There exist positive constants  $R_1$  and  $R_2$  with  $R_1^2 + R_2^2 = 1$  such that  $f$  is congruent to the immersion  $F/G$  given by (1.3), where the lattice  $G$  satisfies  $G \not\subset W_+(n)$  and  $G \not\subset W_-(n)$  for all integers  $n \geq 2$ .*

**REMARK.** Let  $G$  be a lattice of  $\mathbb{R}^2$  generated by the following two vectors

$$u = (2\pi R_1 u_1, 2\pi R_2 u_2), v = (2\pi R_1 v_1, 2\pi R_2 v_2), \quad u_i, v_i \in \mathbb{Z}.$$

Then it is easy to see that the following statements are equivalent.

- (1)  $G \not\subset W_+(n)$  and  $G \not\subset W_-(n)$  for all integers  $n \geq 2$ .
- (2)  $g.c.d(u_1 + u_2, v_1 + v_2) = g.c.d(u_1 - u_2, v_1 - v_2) = 1$ .

The outline of this paper is as follows. In Section 2 we study the geometry of a flat torus  $M_\gamma \subset S^3$  which is the inverse image under the Hopf fibration  $S^3 \rightarrow S^2$  of a closed curve  $\gamma$  in  $S^2$ . In Section 3 we show that if  $\gamma$  is an  $n$ -fold circle in  $S^2$  ( $n \geq 2$ ), then the flat torus  $M_\gamma \subset S^3$  is deformable (Lemma 3.2). Using this lemma, we obtain Theorem 3.1. In Section 4 we prove Theorems 1.2 and 1.3. The key ingredient in the proof of Theorem 1.2 is Lemma 4.1 which is obtained by using a method developed in [2]. The assertion of Theorem 1.3 follows from the main result of [5] which states that every flat torus isometrically immersed in  $S^3$  with nonconstant mean curvature is deformable. In the final section we prove Theorem 5.1. This theorem, which is used in the proof of Lemma 3.2, ensures the existence of certain deformation of an  $n$ -fold circle in  $S^2$  for  $n \geq 2$ .

### 2. Hopf tori in $S^3$

In this section we study the geometry of a flat torus in  $S^3$  constructed by using the Hopf fibration  $S^3 \rightarrow S^2$ . We start with the description of the Hopf fibration by using the group structure of  $S^3$ . Let  $SU(2)$  be the group of all  $2 \times 2$  unitary matrices with determinant 1. Its Lie algebra  $\mathfrak{su}(2)$  consists of all  $2 \times 2$  skew Hermitian matrices of trace 0. We define a positive definite inner product  $\langle \cdot, \cdot \rangle$  on  $\mathfrak{su}(2)$  by

$$\langle u, v \rangle = -\frac{1}{2} \text{trace}(uv), \quad u, v \in \mathfrak{su}(2).$$

The inner product  $\langle \cdot, \cdot \rangle$  is invariant under the adjoint action  $\text{Ad}: SU(2) \rightarrow \text{Aut}(\mathfrak{su}(2))$ . We set

$$e_1 = \begin{bmatrix} 0 & \sqrt{-1} \\ \sqrt{-1} & 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad e_3 = \begin{bmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{bmatrix}.$$

Then  $\{e_1, e_2, e_3\}$  is an orthonormal basis of  $\mathfrak{su}(2)$  such that

$$[e_1, e_2] = 2e_3, \quad [e_2, e_3] = 2e_1, \quad [e_3, e_1] = 2e_2,$$

where  $[\cdot, \cdot]$  is the Lie bracket on  $\mathfrak{su}(2)$ . For  $i = 1, 2, 3$ , we denote by  $E_i$  the left invariant vector field on  $SU(2)$  corresponding to  $e_i$ , and we endow  $SU(2)$  with a bi-invariant Riemannian metric  $\langle \cdot, \cdot \rangle$  satisfying  $\langle E_i, E_j \rangle = \delta_{ij}$ . Then  $SU(2)$  is a Riemannian manifold isometric to the unit sphere  $S^3$ . Henceforth, we identify  $S^3$  with  $SU(2)$ . Let  $S^2$  be the unit sphere in  $\mathfrak{su}(2)$  given by  $S^2 = \{u \in \mathfrak{su}(2) : |u| = 1\}$ , and  $p: S^3 \rightarrow S^2$  the Hopf fibration given by

$$p(a) = \text{Ad}(a)e_3.$$

The vector field  $E_3$  is tangent to the fibers of the Hopf fibration. For  $X, Y \in T_a S^3$ , it follows that

$$(2.1) \quad \langle p_* X, p_* Y \rangle = 4\{\langle X, Y \rangle - \langle X, E_3 \rangle \langle Y, E_3 \rangle\},$$

$$(2.2) \quad p_*(D_X E_3) = -J(p_* X),$$

where  $D$  denotes the Riemannian connection on  $S^3$ , and  $J$  denotes the almost complex structure on  $S^2$  defined by  $J(v) = [u, v]/2$  for  $v \in T_u S^2$ . We identify the unit tangent bundle of  $S^2$  with the subset  $US^2 \subset S^2 \times S^2$  defined by

$$US^2 = \{(u, v) \in S^2 \times S^2 : \langle u, v \rangle = 0\}.$$

Here, the canonical projection  $p_1: US^2 \rightarrow S^2$  is given by  $p_1(u, v) = u$ . Furthermore, we define a double covering  $p_2: S^3 \rightarrow US^2$  by

$$p_2(a) = (\text{Ad}(a)e_3, \text{Ad}(a)e_1).$$

Let  $\gamma: \mathbb{R} \rightarrow S^2$  be a  $2\pi$ -periodic regular curve in  $S^2$ . Using the Hopf fibration  $p: S^3 \rightarrow S^2$ , we construct a 2-dimensional torus  $M_\gamma$  and an immersion  $f_\gamma: M_\gamma \rightarrow S^3$  by

$$M_\gamma = \{(e^{is}, a) \in S^1 \times S^3 : \gamma(s) = p(a)\}, \quad f_\gamma(e^{is}, a) = a,$$

where  $S^1$  denotes the unit circle in  $\mathbb{C}$ . The immersion  $f_\gamma$  induces a flat Riemannian metric on  $M_\gamma$  (see [8]). So we obtain a flat torus  $M_\gamma$  and an isometric immersion

$$f_\gamma: M_\gamma \rightarrow S^3.$$

The immersion  $f_\gamma$  is called the *Hopf torus* corresponding to  $\gamma$ .

In the rest of this section we describe the Riemannian structure of  $M_\gamma$  and the second fundamental form of  $f_\gamma$ . Let  $L(\gamma)$  be the length of  $\gamma$  and  $K(\gamma)$  the total geodesic curvature of  $\gamma$ , that is,

$$L(\gamma) = \int_0^{2\pi} |\gamma'(s)| ds, \quad K(\gamma) = \int_0^{2\pi} k_\gamma(s) |\gamma'(s)| ds,$$

where  $k_\gamma(s)$  denotes the geodesic curvature of  $\gamma(s)$  given by

$$k_\gamma(s) = \frac{\langle \gamma''(s), J(\gamma'(s)) \rangle}{|\gamma'(s)|^3}.$$

We now consider the curve  $\hat{\gamma}: \mathbb{R} \rightarrow US^2$  given by

$$(2.3) \quad \hat{\gamma}(s) = \left( \gamma(s), \frac{\gamma'(s)}{|\gamma'(s)|} \right),$$

and denote by  $I(\gamma)$  the element of the homology group  $H_1(US^2)$  represented by the closed curve  $\hat{\gamma} \mid [0, 2\pi]$ . Note that  $H_1(US^2) \cong \mathbb{Z}_2$ . Let  $c(s)$  be a lift of the curve  $\hat{\gamma}(s)$  with respect to the double covering  $p_2: S^3 \rightarrow US^2$ . Since  $p_2(-a) = p_2(a)$ , we obtain

$$(2.4) \quad c(s + 2\pi) = \begin{cases} c(s) & \text{if } I(\gamma) = 0, \\ -c(s) & \text{if } I(\gamma) = 1. \end{cases}$$

We set

$$(2.5) \quad \Omega(\gamma) = \begin{cases} K(\gamma) & \text{if } I(\gamma) = 0, \\ K(\gamma) + 2\pi & \text{if } I(\gamma) = 1, \end{cases}$$

and define  $W(\gamma)$  to be the lattice of  $\mathbb{R}^2$  generated by the following two vectors

$$(2.6) \quad v_1 = \left( \frac{L(\gamma)}{2}, \frac{\Omega(\gamma)}{2} \right), \quad v_2 = (0, 2\pi).$$

Then the Riemannian structure of  $M_\gamma$  is given by the following

**Lemma 2.1.** *The flat torus  $M_\gamma$  is isometric to  $\mathbb{R}^2/W(\gamma)$ .*

To establish the lemma we consider the covering  $\varphi: \mathbb{R}^2 \rightarrow M_\gamma$  defined by

$$(2.7) \quad \varphi(s, \tau) = (e^{is}, \bar{\gamma}(s) \exp(\tau e_3)),$$

where  $\bar{\gamma}(s)$  is a curve in  $S^3$  such that  $p(\bar{\gamma}(s)) = \gamma(s)$  and  $\langle \bar{\gamma}'(s), E_3 \rangle = 0$ . Then it follows from (2.1) that  $|\gamma'(s)| = |p_*\bar{\gamma}'(s)| = 2|\bar{\gamma}'(s)|$ . So we obtain

$$(2.8) \quad \varphi^* g_\gamma = \frac{1}{4} |\gamma'(s)|^2 ds^2 + d\tau^2,$$

where  $g_\gamma$  denotes the Riemannian metric on  $M_\gamma$ . Let  $\rho: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a diffeomorphism given by

$$\rho(s, \tau) = \left( \frac{1}{2} \int_0^s |\gamma'(s)| ds, \tau \right),$$

and  $\Phi: \mathbb{R}^2 \rightarrow M_\gamma$  a covering map defined by

$$(2.9) \quad \Phi(x_1, x_2) = \varphi(\rho^{-1}(x_1, x_2)).$$

Since  $\rho^*(dx_1^2 + dx_2^2) = \varphi^* g_\gamma$ , the map  $\Phi$  is a Riemannian covering, and so the assertion of Lemma 2.1 follows from the lemma below.

**Lemma 2.2.** *For  $x, x' \in \mathbb{R}^2$ ,  $\Phi(x) = \Phi(x')$  if and only if  $x' - x \in W(\gamma)$ .*

Proof. We set  $x = \rho(s, \tau)$  and  $x' = \rho(s', \tau')$ . Then we obtain

$$(2.10) \quad \Phi(x) = (e^{is}, \bar{\gamma}(s) \exp(\tau e_3)), \quad \Phi(x') = (e^{is'}, \bar{\gamma}(s') \exp(\tau' e_3)).$$

Since  $p(c(s)) = p(\bar{\gamma}(s))$ , there exists a real valued function  $\mu(s)$  such that

$$(2.11) \quad c(s) = \bar{\gamma}(s) \exp(\mu(s) e_3).$$

On the other hand, it follows from [4, Lemma2.2] that the curve  $c(s)$  satisfies

$$(2.12) \quad c(s)^{-1} c'(s) = \frac{1}{2} |\gamma'(s)| (e_2 + k_\gamma(s) e_3).$$

Since  $\langle \bar{\gamma}^{-1}(s) \bar{\gamma}'(s), e_3 \rangle = 0$ , (2.11) and (2.12) imply  $\mu'(s) = (1/2) k_\gamma(s) |\gamma'(s)|$ . Hence

$$(2.13) \quad \mu(s + 2\pi) - \mu(s) = \int_0^{2\pi} \frac{1}{2} k_\gamma(s) |\gamma'(s)| ds = \frac{1}{2} K(\gamma).$$

Using (2.4), (2.5), (2.11) and (2.13), we obtain  $\bar{\gamma}(s + 2\pi) = \bar{\gamma}(s) \exp\{-(1/2)\Omega(\gamma)e_3\}$ . So it follows from (2.10) that  $\Phi(x) = \Phi(x')$  if and only if there exist  $m_1, m_2 \in \mathbb{Z}$  satisfying

$$(2.14) \quad s' - s = 2m_1\pi, \quad \tau' - \tau = \frac{m_1}{2}\Omega(\gamma) + 2m_2\pi.$$

Since  $x' - x = \{(1/2) \int_s^{s'} |\gamma'(s)| ds, \tau' - \tau\}$ , we see that (2.14) is equivalent to

$$x' - x = \left( \frac{m_1}{2} L(\gamma), \frac{m_1}{2} \Omega(\gamma) + 2m_2\pi \right).$$

This completes the proof. □

We now deal with the second fundamental form of the immersion  $f_\gamma: M_\gamma \rightarrow S^3$ . Let  $\xi$  be a unit normal vector field of  $f_\gamma$  such that

$$p_* \xi(e^{is}, a) = 2n(s), \quad (e^{is}, a) \in M_\gamma,$$

where  $n(s) = J(\gamma'(s))/|\gamma'(s)|$ , and let  $h_\gamma$  denote the second fundamental form of the immersion  $f_\gamma$  with respect to  $\xi$ . Then

**Lemma 2.3.**  $\varphi^* h_\gamma = (1/2) k_\gamma(s) |\gamma'(s)|^2 ds^2 - |\gamma'(s)| ds d\tau$ .

Proof. We set

$$f = f_\gamma \circ \varphi, \quad X = \frac{\partial f}{\partial s}, \quad Y = \frac{\partial f}{\partial \tau}.$$

Since  $\langle \bar{\gamma}', E_3 \rangle = 0$ , it follows from [2, Lemma 3.3] that

$$p_*(D_X X) = p_*(D_{\bar{\gamma}'} \bar{\gamma}') = \nabla_{\gamma'} \gamma',$$

where  $\nabla$  denotes the Riemannian connection on  $S^2$ . Hence (2.1) implies

$$(2.15) \quad \begin{aligned} h_\gamma \left( \frac{\partial \varphi}{\partial s}, \frac{\partial \varphi}{\partial s} \right) &= \langle D_X X, \xi(\varphi) \rangle = \frac{1}{4} \langle p_*(D_X X), p_* \xi(\varphi) \rangle \\ &= \frac{1}{4} \langle \nabla_{\gamma'} \gamma', 2n \rangle = \frac{1}{2} \langle \gamma'', n \rangle = \frac{1}{2} k_\gamma |\gamma'|^2. \end{aligned}$$

Since  $Y(s, \tau) = E_3(f(s, \tau))$ , it follows from (2.2) that  $p_*(D_X Y) = p_*(D_X E_3) = -J(p_* X) = -J(\gamma')$ . Hence

$$(2.16) \quad \begin{aligned} h_\gamma \left( \frac{\partial \varphi}{\partial s}, \frac{\partial \varphi}{\partial \tau} \right) &= \langle D_X Y, \xi(\varphi) \rangle = \frac{1}{4} \langle p_*(D_X Y), p_* \xi(\varphi) \rangle \\ &= \frac{1}{4} \langle -J(\gamma'), 2n \rangle = -\frac{1}{2} |\gamma'|. \end{aligned}$$

Since the integral curves of the vector field  $E_3$  are geodesics in  $S^3$ , we see that  $D_Y Y = 0$ . Hence

$$(2.17) \quad h_\gamma \left( \frac{\partial \varphi}{\partial \tau}, \frac{\partial \varphi}{\partial \tau} \right) = \langle D_Y Y, \xi(\varphi) \rangle = 0.$$

The assertion of Lemma 2.3 follows from (2.15)–(2.17).  $\square$

Using (2.8) and Lemma 2.3, we obtain

$$(2.18) \quad |H_\gamma(\varphi(s, \tau))| = |k_\gamma(s)|,$$

where  $H_\gamma$  denotes the mean curvature vector field of the immersion  $f_\gamma$ .

### 3. Isometric deformations of $F/G$

Let  $W_\pm(n)$  denote the lattices of  $\mathbb{R}^2$  defined by (1.4). In this section we show the following theorem.

**Theorem 3.1.** *Let  $G$  be a lattice of  $\mathbb{R}^2$  such that  $G \subset G_0$ . If  $G \subset W_+(n)$  or  $G \subset W_-(n)$  for some integer  $n \geq 2$ , the isometric immersion  $F/G$  given by (1.3) is deformable.*

To establish the theorem above we need some lemmas. For each integer  $n \geq 1$ , let  $\gamma: \mathbb{R} \rightarrow S^2$  be a  $2\pi$ -periodic regular curve defined by

$$(3.1) \quad \gamma(s) = (\cos \theta \cos ns)e_1 + (\cos \theta \sin ns)e_2 + (\sin \theta)e_3,$$



where  $\theta$  is a constant such that

$$(3.2) \quad \frac{R_2^2 - R_1^2}{2R_1R_2} = \tan \theta, \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2}.$$

Note that the geodesic curvature of  $\gamma$  satisfies

$$(3.3) \quad k_\gamma(s) = \tan \theta.$$

We now consider the Hopf torus  $f_\gamma: M_\gamma \rightarrow S^3$  corresponding to  $\gamma$ . Then

**Lemma 3.2.** *The immersion  $f_\gamma$  is deformable for  $n \geq 2$ .*

*Proof.* Since  $n \geq 2$ , it follows that there exists a smooth one parameter family of  $2\pi$ -periodic regular curves  $\gamma_t: \mathbb{R} \rightarrow S^2$ ,  $t \in \mathbb{R}$ , such that  $\gamma_0 = \gamma$  and

$$(3.4) \quad L(\gamma_t) = L(\gamma), \quad K(\gamma_t) = K(\gamma),$$

$$(3.5) \quad k_{\gamma_t}(0) \neq \tan \theta \quad \text{for all } t \neq 0.$$

The existence of  $\gamma_t$  as above will be established in the final section (Theorem 5.1). Let  $\tilde{\gamma}_t: \mathbb{R} \rightarrow S^3$ ,  $t \in \mathbb{R}$  be a smooth one parameter family of curves in  $S^3$  such that  $p(\tilde{\gamma}_t(s)) = \gamma_t(s)$  and  $\langle \tilde{\gamma}'_t(s), E_3 \rangle = 0$ , and  $\Phi_t: \mathbb{R}^2 \rightarrow M_{\gamma_t}$  the Riemannian covering map defined in the same way as (2.9). Then, by Lemma 2.2,  $\Phi_t$  induces the isometry

$$\tilde{\Phi}_t: \mathbb{R}^2/W(\gamma_t) \rightarrow M_{\gamma_t}.$$

Since  $I(\gamma_t) = I(\gamma)$ , it follows from (3.4) that  $W(\gamma_t) = W(\gamma)$ . So, by setting  $f_t = f_{\gamma_t} \circ \tilde{\Phi}_t \circ \tilde{\Phi}_0^{-1}$ , we obtain a smooth one parameter family of isometric immersions  $f_t: M_\gamma \rightarrow S^3$ ,  $t \in \mathbb{R}$ , such that  $f_0 = f_\gamma$ . Let  $H_t$  denote the mean curvature vector field of  $f_t$ . Then it follows from (2.18) and (3.5) that there exists a point  $a \in M_\gamma$  such that  $|H_1(a)| \neq |\tan \theta|$ . On the other hand, (3.3) implies that  $|H_0(x)| = |\tan \theta|$  for all  $x \in M_\gamma$ . Hence  $f_0 \not\equiv f_1$ , and so the immersion  $f_\gamma$  is deformable.  $\square$

**Lemma 3.3.** *The immersions  $F/W_+(n)$  and  $F/W_-(n)$  are deformable for  $n \geq 2$ .*

*Proof.* We first note that  $F/W_+(n) \equiv F/W_-(n)$ . So, by Lemma 3.2, it is sufficient to show that  $f_\gamma \equiv F/W_+(n)$ . Let  $\Phi: \mathbb{R}^2 \rightarrow M_\gamma$  be the Riemannian covering defined by (2.9), and  $\tilde{f}_\gamma: \mathbb{R}^2 \rightarrow S^3$  an isometric immersion given by  $\tilde{f}_\gamma = f_\gamma \circ \Phi$ . We denote by  $\tilde{h}$  the second fundamental form of the immersion  $\tilde{f}_\gamma$ . Then it follows from Lemma 2.3 and (3.3) that

$$(3.6) \quad \tilde{h} = 2 \tan \theta dx_1^2 - 2 dx_1 dx_2.$$

Let  $T$  be an isometry of  $\mathbb{R}^2$  given by

$$T(x_1, x_2) = (R_2x_1 + R_1x_2, -R_1x_1 + R_2x_2).$$

Then it follows from (3.2) and (3.6) that

$$T^*\tilde{h} = \frac{R_2}{R_1} dx_1^2 - \frac{R_1}{R_2} dx_2^2.$$

Hence the isometric immersions  $\tilde{f}_\gamma \circ T: \mathbb{R}^2 \rightarrow S^3$  and  $F: \mathbb{R}^2 \rightarrow S^3$  have the same second fundamental form. So it follows from the fundamental theorem of the theory of surfaces that there exists an isometry  $A: S^3 \rightarrow S^3$  satisfying

$$(3.7) \quad \tilde{f}_\gamma \circ T = A \circ F.$$

On the other hand, (3.1) implies

$$L(\gamma) = 2n\pi \cos \theta, \quad K(\gamma) = 2n\pi \sin \theta, \quad I(\gamma) = \begin{cases} 0 & \text{if } n \text{ is even,} \\ 1 & \text{if } n \text{ is odd.} \end{cases}$$

So, by (2.6), the lattice  $W(\gamma)$  is generated by

$$v_1 = (n\pi \cos \theta, n\pi \sin \theta + n\pi), \quad v_2 = (0, 2\pi).$$

Hence we obtain  $W(\gamma) = \{n_1\xi_1 + n_2\xi_2 : n_1 + n_2 \in n\mathbb{Z}\}$ , where

$$\xi_1 = (\pi \cos \theta, \pi \sin \theta - \pi), \quad \xi_2 = (\pi \cos \theta, \pi \sin \theta + \pi).$$

By (3.2) the vectors  $\xi_1$  and  $\xi_2$  can be written as

$$\xi_1 = 2\pi R_1(R_2, -R_1), \quad \xi_2 = 2\pi R_2(R_1, R_2).$$

This shows that  $T(W_+(n)) = W(\gamma)$ , and so it follows from Lemma 2.2 that there exists a diffeomorphism  $\tilde{T}: \mathbb{R}^2/W_+(n) \rightarrow M_\gamma$  satisfying  $\Phi \circ T = \tilde{T} \circ q$ , where  $q$  denotes the canonical projection of  $\mathbb{R}^2$  onto  $\mathbb{R}^2/W_+(n)$ . Therefore (3.7) implies that  $f_\gamma \circ \tilde{T} = A \circ F/W_+(n)$ .  $\square$

By the lemma above the assertion of Theorem 3.1 follows from the following

**Lemma 3.4.** *Let  $W$  be a lattice of  $\mathbb{R}^2$  such that  $W \subset G_0$  and the immersion  $F/W$  is deformable. Then for each lattice  $G \subset W$  the immersion  $F/G$  is deformable.*

*Proof.* By the assumption, the isometric immersion  $F/W: \mathbb{R}^2/W \rightarrow S^3$  admits an isometric deformation  $f_t: \mathbb{R}^2/W \rightarrow S^3$  such that  $f_0 \neq f_1$ . Then the mean curvature

vector field of  $f_t$ , denoted by  $H_t$ , satisfies

$$(3.8) \quad \begin{cases} |H_0(x)| = |R_2^2 - R_1^2|/2R_1R_2 & \text{for all } x \in \mathbb{R}^2/W, \\ |H_1(a)| \neq |R_2^2 - R_1^2|/2R_1R_2 & \text{for some } a \in \mathbb{R}^2/W. \end{cases}$$

We now consider the canonical projection  $q: \mathbb{R}^2/G \rightarrow \mathbb{R}^2/W$  and an isometric deformation of  $F/G$  given by  $\tilde{f}_t = f_t \circ q$ . Then (3.8) implies that  $\tilde{f}_0 \not\cong \tilde{f}_1$ , and so the immersion  $F/G$  is deformable.  $\square$

#### 4. Proof of main theorems

In this section we give the proof of Theorems 1.2 and 1.3. Consider the map  $\sigma: G_0 \rightarrow G_0$  defined by

$$\sigma(2\pi R_1 n_1, 2\pi R_2 n_2) = (2\pi R_1 n_2, 2\pi R_2 n_1).$$

The following lemma is the key ingredient in the proof of Theorem 1.2.

**Lemma 4.1.** *Let  $G$  be a lattice of  $\mathbb{R}^2$  such that  $G \subset G_0$ . If  $F_t: \mathbb{R}^2 \rightarrow S^3$ ,  $t \in \mathbb{R}$ , is a  $G$ -invariant isometric deformation of the immersion  $F$ , then the deformation  $F_t$  is  $\sigma(G)$ -invariant.*

*Proof.* Let  $v \in G$ . Then it is sufficient to show that

$$(4.1) \quad F_t(x + \sigma(v)) = F_t(x) \quad \text{for all } t \in \mathbb{R}.$$

Since  $F_t: \mathbb{R}^2 \rightarrow S^3$  is an isometric immersion, it follows from [7] that there exists a diffeomorphism  $T_t: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that

$$(4.2) \quad |\partial_i T_t| = 1, \quad h_t(\partial_i T_t, \partial_i T_t) = 0 \quad \text{for } i = 1, 2,$$

where  $h_t$  denotes the second fundamental form of  $F_t$ . We may assume that the map  $(t, s_1, s_2) \mapsto T_t(s_1, s_2)$  is smooth and

$$(4.3) \quad T_t(0, 0) = (0, 0), \quad T_0(s_1, s_2) = (R_1(s_1 - s_2), R_2(s_1 + s_2)).$$

By (4.2) we obtain  $\partial_1 \partial_2 T_t = (0, 0)$ . So it follows from  $T_t(0, 0) = (0, 0)$  that

$$(4.4) \quad T_t(s_1, s_2) = T_t(s_1, 0) + T_t(0, s_2).$$

We set

$$(4.5) \quad (l_1(t), l_2(t)) = T_t^{-1}(v), \quad z(t) = T_t(l_1(t), 0).$$

Then  $v = T_0(l_1(0), l_2(0)) = (R_1(l_1(0) - l_2(0)), R_2(l_1(0) + l_2(0)))$ , and so we obtain

$$v + \sigma(v) = 2z(0).$$

Since  $F_t$  is  $G$ -invariant, the relation above implies

$$(4.6) \quad F_t(x + \sigma(v)) = F_t(x + 2z(0)).$$

Let  $p_t: \mathbb{R}^2 \rightarrow \mathbb{R}^2/G$  be a covering given by  $p_t = p \circ T_t$ , where  $p: \mathbb{R}^2 \rightarrow \mathbb{R}^2/G$  denotes the canonical projection. Since  $p(v) = p(0, 0)$ , it follows from (4.3) and (4.5) that  $p_t(l_1(t), l_2(t)) = p_t(0, 0)$ . So there exists a diffeomorphism  $\varphi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $p_t \circ \varphi = p_t$  and  $\varphi(0, 0) = (l_1(t), l_2(t))$ . Then  $T_t(\varphi(s_1, s_2)) - T_t(s_1, s_2) \in G$ , and so it follows from (4.5) that

$$(4.7) \quad T_t(\varphi(s_1, s_2)) = T_t(s_1, s_2) + v.$$

We now consider an immersion  $\tilde{F}_t: \mathbb{R}^2 \rightarrow S^3$  defined by  $\tilde{F}_t = F_t \circ T_t$ . Then, by (4.2), we see that the immersion  $\tilde{F}_t$  is a FAT. Here, we refer the reader to [2, p. 460] for the definition of FAT. Furthermore, it follows from (4.7) that  $\tilde{F}_t \circ \varphi = \tilde{F}_t$ . Therefore, [2, Theorem 2.3] implies

$$(4.8) \quad \varphi(s_1, s_2) = (s_1 + l_1(t), s_2 + l_2(t)).$$

In particular, we obtain  $\tilde{F}_t(s_1 + l_1(t), s_2 + l_2(t)) = \tilde{F}_t(s_1, s_2)$ , and so it follows from [2, Theorem 3.9, Lemma 5.5] that

$$(4.9) \quad \tilde{F}_t(s_1 + 2l_1(t), s_2) = \tilde{F}_t(s_1, s_2).$$

On the other hand, combining (4.7) and (4.8), we obtain

$$(4.10) \quad T_t(s_1 + l_1(t), s_2 + l_2(t)) = T_t(s_1, s_2) + v.$$

Using (4.4), (4.5) and (4.10), we see that

$$\begin{aligned} T_t(s_1 + l_1(t), s_2) &= T_t(s_1 + l_1(t), l_2(t)) + T_t(0, s_2) - T_t(0, l_2(t)) \\ &= T_t(s_1, s_2) + v - T_t(0, l_2(t)) = T_t(s_1, s_2) + z(t). \end{aligned}$$

This implies  $T_t(s_1 + 2l_1(t), s_2) = T_t(s_1, s_2) + 2z(t)$ , and so it follows from (4.9) that

$$(4.11) \quad F_t(x + 2z(t)) = F_t(x).$$

By (4.6) and (4.11), we see that (4.1) follows from the assertion that  $z(t) = z(0)$  for all  $t \in \mathbb{R}$ . To establish this assertion, suppose that there exists  $t_0$  such that  $z(t_0) \neq z(0)$ .

Since the set of all points  $z(t) \in \mathbb{R}^2$  is not countable, there exists  $a \in \mathbb{R}$  such that  $z(a)$  is not contained in the countable set  $\bigcup_{n=1}^{\infty} \{x/2n : x \in G\}$ . Let  $f_a: \mathbb{R}^2/G \rightarrow S^3$  be an immersion defined by the relation  $f_a \circ p = F_a$ , and  $\{y_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R}^2/G$  given by  $y_n = p(2nz(a))$ . Then it follows from (4.11) that  $f_a(y_m) = f_a(y_n)$ . Furthermore, as  $2nz(a) \notin G$  for all  $n \geq 1$ , we obtain  $y_m \neq y_n$  ( $m \neq n$ ). So, using the fact that  $f_a$  is locally injective, we see that the sequence  $\{y_n\}_{n=1}^{\infty}$  has no convergent subsequence. This contradicts the fact that  $\mathbb{R}^2/G$  is compact. Hence we obtain  $z(t) = z(0)$  for all  $t \in \mathbb{R}$ .  $\square$

We now recall the lattices  $W_{\pm}(n)$  given by (1.4), and for each lattice  $G \subset G_0$  we consider the lattice

$$G + \sigma(G) = \{u + \sigma(v) : u, v \in G\}.$$

Then we obtain

**Lemma 4.2.** *Let  $G$  be a lattice of  $\mathbb{R}^2$  such that  $G \subset G_0$ . If  $G \not\subset W_+(n)$  and  $G \not\subset W_-(n)$  for all integers  $n \geq 2$ , then  $G + \sigma(G) = G_0$ .*

*Proof.* Since  $G + \sigma(G) \subset G_0$ , it is sufficient to show that  $G_0 \subset G + \sigma(G)$ . Let  $u$  and  $v$  be generators of the lattice  $G$ . Since  $u, v \in G_0$ , we can write as

$$u = (2\pi R_1 u_1, 2\pi R_2 u_2), \quad v = (2\pi R_1 v_1, 2\pi R_2 v_2), \quad u_i, v_i \in \mathbb{Z}.$$

For each integer  $n \geq 2$ , using the assumption  $G \not\subset W_+(n)$ , we see that there exist  $k, l \in \mathbb{Z}$  such that the integer  $n$  does not divide  $k(u_1 + u_2) + l(v_1 + v_2)$ , and so  $n$  is not a common divisor for  $u_1 + u_2$  and  $v_1 + v_2$ . Hence the greatest common divisor for  $u_1 + u_2$  and  $v_1 + v_2$  is equal to 1. Similarly, using the assumption that  $G \not\subset W_-(n)$  for all  $n \geq 2$ , we see that the greatest common divisor for  $u_1 - u_2$  and  $v_1 - v_2$  is equal to 1. Hence there exist  $p, q, r, s \in \mathbb{Z}$  such that

$$(4.12) \quad p(u_1 + u_2) + q(v_1 + v_2) = 1, \quad r(u_1 - u_2) + s(v_1 - v_2) = 1.$$

We now consider the elements  $a, b \in G$  given by

$$a = pu + qv, \quad b = ru + sv.$$

Then it follows from (4.12) that

$$\begin{aligned} b - (ru_2 + sv_2)(a + \sigma(a)) &= (2\pi R_1, 0), \\ a - (pu_1 + qv_1)(b - \sigma(b)) &= (0, 2\pi R_2). \end{aligned}$$

So the lattice  $G + \sigma(G)$  contains  $(2\pi R_1, 0)$  and  $(0, 2\pi R_2)$ . Hence  $G_0 \subset G + \sigma(G)$ .  $\square$

**Lemma 4.3.** *Let  $G$  be a lattice of  $\mathbb{R}^2$  such that  $G \subset G_0$ . If  $G + \sigma(G) = G_0$ , then every isometric deformation of  $F/G$  is trivial.*

*Proof.* Let  $f_t: \mathbb{R}^2/G \rightarrow S^3$ ,  $t \in \mathbb{R}$ , be an isometric deformation of  $F/G$ . Then we obtain a  $G$ -invariant isometric deformation of  $F$  given by  $F_t = f_t \circ p$ , where  $p$  denotes the canonical projection of  $\mathbb{R}^2$  onto  $\mathbb{R}^2/G$ . Since  $G + \sigma(G) = G_0$ , it follows from Lemma 4.1 that each  $F_t$  is  $G_0$ -invariant, and so we obtain

$$F_t/G_0: \mathbb{R}^2/G_0 \rightarrow S^3, \quad t \in \mathbb{R}$$

which is an isometric deformation of the embedding  $F/G_0: \mathbb{R}^2/G_0 \rightarrow S^3$ . Then Theorem 1.1 implies that for each  $t \in \mathbb{R}$  there exists an isometry  $A_t: S^3 \rightarrow S^3$  satisfying  $F_t/G_0 = A_t \circ (F/G_0)$ . Let  $q: \mathbb{R}^2/G \rightarrow \mathbb{R}^2/G_0$  denote the canonical projection. Since  $f_t = (F_t/G_0) \circ q$ , we obtain

$$A_t \circ f_0 = A_t \circ (F/G_0) \circ q = (F_t/G_0) \circ q = f_t.$$

Hence the isometric deformation  $f_t$  is trivial. □

*Proof of Theorem 1.2.* To establish Theorem 1.2, it is sufficient to show the converse of Theorem 3.1. Suppose that  $G \not\subset W_+(n)$  and  $G \not\subset W_-(n)$  for all integers  $n \geq 2$ . Then it follows from Lemmas 4.2 and 4.3 that every isometric deformation of  $F/G$  is trivial. In particular, the isometric immersion  $F/G$  is not deformable. This shows the converse of Theorem 3.1. □

*Proof of Theorem 1.3.* By Lemmas 4.2 and 4.3, it is easy to see that (2)  $\Rightarrow$  (1). We now show that (1)  $\Rightarrow$  (2). Recall the main result of [5] which states that every flat torus isometrically immersed in  $S^3$  with nonconstant mean curvature is deformable. Hence, the assumption (1) implies that the mean curvature of the immersion  $f: M \rightarrow S^3$  must be constant. So there exist positive constants  $R_1$  and  $R_2$  with  $R_1^2 + R_2^2 = 1$  such that  $f$  is congruent to the immersion  $F/G$  given by (1.3). Then  $F/G$  is not deformable, and so it follows from Theorem 1.2 that the lattice  $G$  satisfies  $G \not\subset W_+(n)$  and  $G \not\subset W_-(n)$  for all integers  $n \geq 2$ . □

### 5. Deformations of circles in $S^2$

For each  $2\pi$ -periodic regular curve  $\gamma(s)$  in  $S^2$ , we recall the following notation.

$$L(\gamma) = \int_0^{2\pi} |\gamma'(s)| ds, \quad K(\gamma) = \int_0^{2\pi} k_\gamma(s) |\gamma'(s)| ds,$$

where  $k_\gamma(s)$  denotes the geodesic curvature of  $\gamma(s)$  given by

$$k_\gamma(s) = \frac{\langle \gamma''(s), J(\gamma'(s)) \rangle}{|\gamma'(s)|^3}.$$

In this section we prove the following theorem which was used in the proof of Lemma 3.2.

**Theorem 5.1.** *For each integer  $n \geq 2$ , let  $\gamma: \mathbb{R} \rightarrow S^2$  be a  $2\pi$ -periodic regular curve defined by  $\gamma(s) = (\cos \theta \cos ns)e_1 + (\cos \theta \sin ns)e_2 + (\sin \theta)e_3$ , where  $\theta$  is a constant satisfying  $-\pi/2 < \theta < \pi/2$ . Then there exists a smooth one parameter family of  $2\pi$ -periodic regular curves  $\gamma_t: \mathbb{R} \rightarrow S^2$ ,  $-\delta < t < \delta$ , such that*

- (1)  $\gamma_0 = \gamma$ ,
- (2)  $L(\gamma_t) = L(\gamma)$ ,  $K(\gamma_t) = K(\gamma)$ ,
- (3)  $k_{\gamma_t}(0) \neq \tan \theta$  for all  $t \neq 0$ .

We first show the following lemma which proves the assertion of Theorem 5.1 in the case of  $\theta = 0$ .

**Lemma 5.2.** *For each integer  $n \geq 2$ , let  $\alpha: \mathbb{R} \rightarrow S^2$  be a  $2\pi$ -periodic regular curve defined by  $\alpha(s) = (\cos ns)e_1 + (\sin ns)e_2$ . Then there exists a smooth one parameter family of  $2\pi$ -periodic regular curves  $\alpha_t: \mathbb{R} \rightarrow S^2$ ,  $-\epsilon < t < \epsilon$ , such that*

- (1)  $\alpha_0 = \alpha$ ,
- (2)  $L(\alpha_t) = L(\alpha)$ ,  $K(\alpha_t) = K(\alpha)$ ,
- (3)  $k_{\alpha_t}(0) \neq 0$  for all  $t \neq 0$ .

*Proof.* Let  $v_1(s)$  and  $v_2(s)$  be  $2\pi$ -periodic functions defined by

$$(5.1) \quad v_1(s) = \cos s, \quad v_2(s) = \cos ms, \quad \text{where } m = 2n + 1.$$

For each  $x = (x_1, x_2) \in \mathbb{R}^2$ , we consider the curve  $q_x: \mathbb{R} \rightarrow S^2$  given by

$$q_x(s) = \cos \left( \sum_{i=1}^2 x_i v_i(s) \right) \alpha(s) + \sin \left( \sum_{i=1}^2 x_i v_i(s) \right) e_3.$$

Note that  $q_x(s + 2\pi) = q_x(s)$ , and

$$q_o(s) = \alpha(s), \quad \text{where } o = (0, 0).$$

So there exists an open neighborhood  $V$  of the origin  $o \in \mathbb{R}^2$  such that for each  $x \in V$  the curve  $q_x$  is regular. We consider the smooth functions  $\bar{L}, \bar{K}: V \rightarrow \mathbb{R}$  given by

$$\bar{L}(x) = L(q_x) = \int_0^{2\pi} |q'_x(s)| ds, \quad \bar{K}(x) = K(q_x) = \int_0^{2\pi} k_{q_x}(s) |q'_x(s)| ds.$$

Since  $v_i(s + \pi) = -v_i(s)$ , we obtain  $|q'_x(s + \pi)| = |q'_x(s)|$  and  $k_{q_x}(s + \pi) = -k_{q_x}(s)$ . Therefore

$$(5.2) \quad \bar{K}(x) = 0.$$

Since  $q_o: \mathbb{R} \rightarrow S^2$  is a geodesic, the origin  $o \in \mathbb{R}^2$  is a critical point for the smooth function  $\bar{L}$ . The Hessian of  $\bar{L}$  at the critical point  $o$  is given by

$$\frac{\partial^2 \bar{L}}{\partial x_i \partial x_j}(o) = \frac{1}{n} \int_0^{2\pi} (v'_i(s)v'_j(s) - n^2 v_i(s)v_j(s)) ds.$$

So it follows from (5.1) that

$$(5.3) \quad \frac{\partial^2 \bar{L}}{\partial x_1 \partial x_1}(o) = \frac{1 - n^2}{n} \pi, \quad \frac{\partial^2 \bar{L}}{\partial x_2 \partial x_2}(o) = \frac{m^2 - n^2}{n} \pi, \quad \frac{\partial^2 \bar{L}}{\partial x_1 \partial x_2}(o) = 0.$$

Since  $n > 1$  and  $m = 2n + 1$ , the index of  $\bar{L}$  at the critical point  $o$  is equal to  $-1$ . Hence the Lemma of Morse [6, p. 6] implies that there exists a local coordinate system  $(y_1, y_2)$  in a neighborhood  $U$  of the origin  $o$  such that

$$(5.4) \quad \bar{L}(x) = \bar{L}(o) - y_1(x)^2 + y_2(x)^2, \quad y_1(o) = y_2(o) = 0.$$

For a sufficiently small  $\epsilon > 0$ , let  $x(t) = (x_1(t), x_2(t))$ ,  $-\epsilon < t < \epsilon$ , be a smooth curve in  $U$  defined by

$$(5.5) \quad y_1(x(t)) = t, \quad y_2(x(t)) = t,$$

and we consider the smooth one parameter family of  $2\pi$ -periodic regular curves  $\alpha_t: \mathbb{R} \rightarrow S^2$ ,  $-\epsilon < t < \epsilon$  given by  $\alpha_t = q_{x(t)}$ . Then it follows from (5.2), (5.4) and (5.5) that

$$\alpha_0 = q_o = \alpha, \quad L(\alpha_t) = \bar{L}(x(t)) = \bar{L}(o), \quad K(\alpha_t) = \bar{K}(x(t)) = 0.$$

This implies the assertions (1) and (2). Since the geodesic curvature of  $\alpha_t$  satisfies  $k_{\alpha_t} = \langle \alpha''_t, J(\alpha'_t) \rangle / |\alpha'_t|^3$ , we obtain

$$(5.6) \quad k_{\alpha_t}(0) = \frac{\varphi(t)}{n^2 \cos^2(x_1(t) + x_2(t))},$$

where  $\varphi(t) = n^2 \cos(x_1(t) + x_2(t)) \sin(x_1(t) + x_2(t)) - x_1(t) - m^2 x_2(t)$ . Note that

$$(5.7) \quad \varphi(0) = 0, \quad \varphi'(0) = (n^2 - 1)x'_1(0) + (n^2 - m^2)x'_2(0).$$

Differentiating the relation  $\bar{L}(x(t)) = \bar{L}(o)$  and using (5.3), we obtain

$$(5.8) \quad (n^2 - 1)x'_1(0)^2 + (n^2 - m^2)x'_2(0)^2 = 0.$$



If  $\varphi'(0) = 0$ , it follows from (5.7) and (5.8) that  $x_1'(0) = x_2'(0) = 0$  which is a contradiction. Hence  $\varphi'(0) \neq 0$ . So the assertion (3) follows from (5.6).  $\square$

**Proof of Theorem 5.1.** Let  $\alpha_t: \mathbb{R} \rightarrow S^2$ ,  $-\epsilon < t < \epsilon$ , be a smooth one parameter family of  $2\pi$ -periodic regular curves satisfying the conditions (1)–(3) of Lemma 5.2, and  $n_t$  a unit normal vector field along  $\alpha_t$  given by  $n_t(s) = J(\alpha_t'(s))/|\alpha_t'(s)|$ . Consider the curve  $\gamma_t: \mathbb{R} \rightarrow S^2$  given by  $\gamma_t(s) = (\cos \theta)\alpha_t(s) + (\sin \theta)n_t(s)$ . Then it follows from the relation  $n_t'(s) = -k_{\alpha_t}(s)\alpha_t'(s)$  that

$$(5.9) \quad \gamma_t'(s) = (\cos \theta - k_{\alpha_t}(s) \sin \theta)\alpha_t'(s).$$

Since  $k_{\alpha_0}(s) = 0$  and  $\cos \theta > 0$ , there exists a positive number  $\delta$  such that

$$\cos \theta - k_{\alpha_t}(s) \sin \theta > 0 \quad \text{for } |t| < \delta.$$

So it follows that  $\gamma_t: \mathbb{R} \rightarrow S^2$ ,  $-\delta < t < \delta$ , is a smooth one parameter family of  $2\pi$ -periodic regular curves. Hence it is sufficient to show that the family  $\gamma_t$  satisfies (1)–(3) of Theorem 5.1. Using (1) of Lemma 5.2, we obtain  $n_0(s) = e_3$ . This implies the assertion (1). On the other hand, the geodesic curvature of  $\gamma_t$  satisfies

$$(5.10) \quad k_{\gamma_t}(s) = \frac{\sin \theta + k_{\alpha_t}(s) \cos \theta}{\cos \theta - k_{\alpha_t}(s) \sin \theta}.$$

By (5.9) and (5.10) we obtain

$$L(\gamma_t) = \cos \theta L(\alpha_t) - \sin \theta K(\alpha_t), \quad K(\gamma_t) = \sin \theta L(\alpha_t) + \cos \theta K(\alpha_t).$$

So the assertion (2) follows from (2) of Lemma 5.2. Furthermore, using (3) of Lemma 5.2 and (5.10), we see that  $k_{\gamma_t}(0) \neq \tan \theta$  for all  $t \neq 0$ . This implies the assertion (3).  $\square$

---

### References

- [1] K. Enomoto, Y. Kitagawa and J.L. Weiner: *A rigidity theorem for the Clifford tori in  $S^3$* , Proc. A.M.S. **124** (1996), 265–268.
- [2] Y. Kitagawa: *Periodicity of the asymptotic curves on flat tori in  $S^3$* , J. Math. Soc. Japan, **40** (1988), 457–476.
- [3] Y. Kitagawa: *Rigidity of the Clifford tori in  $S^3$* , Math. Z., **198** (1988), 591–599.
- [4] Y. Kitagawa: *Embedded flat tori in the unit 3-sphere*, J. Math. Soc. Japan, **47** (1995), 275–296.
- [5] Y. Kitagawa: *Isometric deformations of a flat torus in the 3-sphere with nonconstant mean curvature*, Tohoku Math. J. **52** (2000), 283–298.
- [6] J. Milnor: *Morse theory*, Princeton University Press, Princeton, NJ, 1963.
- [7] J.D. Moore: *Isometric immersions of space forms in space forms*, Pacific J. Math., **40** (1972), 157–166.

- [8] U. Pinkall: *Hopf tori in  $S^3$* , *Invent. math.*, **81** (1985), 379–386.
- [9] S.T. Yau: *Submanifolds with constant mean curvature II*, *Amer. J. Math.*, **97** (1975), 76–100.

Department of Mathematics  
Utsunomiya University  
Utsunomiya 321-8505, Japan  
e-mail: kitagawa@cc.utsunomiya-u.ac.jp