

|              |   |
|--------------|---|
| Title        | On regular rings whose maximal right quotient rings are type I <sub>f</sub> |
| Author(s)    | Kutami, Mamoru; Inoue, Ichiro   |
| Citation     | Osaka Journal of Mathematics. 1995, 32(2), p. 501-512                       |
| Version Type | VoR   |
| URL          | <a href="https://doi.org/10.18910/6952">https://doi.org/10.18910/6952</a>   |
| rights       |   |
| Note         |   |

***Osaka University Knowledge Archive : OUKA***

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## ON REGULAR RINGS WHOSE MAXIMAL RIGHT QUOTIENT RINGS ARE TYPE $I_f$

MAMORU KUTAMI AND ICHIRO INOUE

(Received July 26, 1993)

### Introduction

This paper is written about the property (DF) on regular rings whose maximal right quotient rings are Type  $I_f$ . Hereafter regular rings whose maximal right quotient rings are Type  $I_f$  are said to satisfy (\*). The property (DF) is very important property when we study on regular rings satisfying (\*), and it was treated in the paper [5] written by the first author, where (DF) for a ring  $R$  is defined as that if the direct sum of any two directly finite projective  $R$ -modules is always directly finite. In the above paper, the equivalent condition that a regular ring  $R$  of bounded index satisfies (DF) was discovered and called (#). Stillmore, we proved that the condition (DF) is equivalent to (#) for regular rings whose primitive factor rings are artinian in the paper [6]. Then we have the problem that (DF) is equivalent (#) for regular rings satisfying (\*) or not, where the condition (\*) is weaker than one that primitive factor rings are artinian.

In §2, we shall prove Theorem 2.4. This is important, and using this, Theorem 2.5 (i.e. if  $R$  is a regular ring satisfying (\*) and  $k$  is any positive integer, then  $kP$  is directly finite for every directly finite projective  $R$ -module  $P$ ) is proved. Moreover, we shall solve the above problem in Theorem 2.11.

In §3, we shall consider some applications of Theorem 2.11. We prove Theorem 3.3 that if  $R$  is a regular ring satisfying (\*) whose maximal right quotient ring of  $R$  satisfies (DF), then so does  $R$ . Though it is clear that a regular rings satisfying (\*) which has a nonzero essential socle satisfies (DF), we can prove that, for regular rings satisfying (\*), the condition having a nonzero essential socle is not equivalent to (#) in Example 3.4. Next, we shall consider that  $(\Pi_1^\infty R)/(\oplus R)$  satisfies (DF) or not for a regular ring  $R$  satisfying (\*). This problem is a generalization of Example 3.4, and we prove that, for a regular ring  $R$  of bounded index,  $(\Pi_1^\infty R)/(\oplus R)$  satisfies (DF) (Theorem 3.9).

Throughout this paper,  $R$  is a ring with identity and  $R$ -modules are unitary right  $R$ -modules.

### 1. Definitions and notations

DEFINITION 1. A ring  $R$  is (Von Neumann) *regular* provided that for every  $x \in R$  there exists  $y \in R$  such that  $xyx = x$ .

NOTE. Every projective modules over regular rings have the exchange property.

DEFINITION 2. A module  $M$  is *directly finite* provided that  $M$  is not isomorphic to a proper direct summand of itself. If  $M$  is not directly finite, then  $M$  is said to be *directly infinite*. A ring  $R$  is said to be *directly finite* (resp. *directly infinite*) if so is  $R$  as an  $R$ -module.

DEFINITION 3. The *index* of a nilpotent element  $x$  in a ring  $R$  is the least positive integer such that  $x^n = 0$  (In particular, 0 is nilpotent of index 1). The *index* of a two-sided ideal  $J$  of  $R$  is the supremum of the indices of all nilpotent elements of  $J$ .

If this supremum is finite, then  $J$  is said to have *bounded index*. If  $J$  does not have bounded index,  $J$  is said to be *index  $\infty$* .

NOTE. Let  $R$  be a regular ring with index  $\infty$ . Then using [3, the proof of Lemma 2], there exists a family  $\{A_n\}_{n=1}^{\infty}$  of independent right ideals of  $R$  such that  $A_n$  contains a direct sum of  $n$  nonzero pairwise isomorphic right ideals. Therefore  $R$  has a family  $\{e_{ij}\}_{i,j=1,2,\dots}$  of idempotents such that

$$e_{21}R \simeq e_{22}R$$

$$e_{31}R \simeq e_{32}R \simeq e_{33}R$$

.....

, where  $e_{ij} = 0$  ( $i < j$ ), and  $\{e_{i1}, \dots, e_{ii}\}$  are orthogonal for all  $i$ .

DEFINITION 4. A ring  $R$  has (DF) if the direct sum of two directly finite projective  $R$ -modules is directly finite.

DEFINITION 5. A regular ring  $R$  is *abelian* provided all idempotents in  $R$  are central.

DEFINITION 6. A ring  $R$  satisfies (\*) if every nonzero two-sided ideal of  $R$  contains a nonzero two-sided ideal of bounded index.

DEFINITION 7. A ring  $R$  is *unit-regular* provided that for each  $x \in R$  there is a unit  $u \in R$  such that  $xux = x$ .

NOTE. Every finitely generated projective module over a unit-regular ring has the cancellation property ([2, Theorem 4.14]).

DEFINITION 8. Let  $e$  be an idempotent in a regular ring  $R$ . Then  $e$  is called an *abelian idempotent* (of  $R$ ) whenever the ring  $eRe$  is abelian.

DEFINITION 9. Let  $e$  be an idempotent in a regular right self-injective ring  $R$ . Then  $e$  is *faithful* (in  $R$ ) if 0 is the only central idempotent of  $R$  which is orthogonal to  $e$ . A regular right self-injective ring  $R$  is said to be *Type I* provided that it contains a faithful abelian idempotent, and  $R$  is *Type  $I_f$*  if  $R$  is *Type I* and directly finite.

NOTE. It is well-known from [4, Theorem 2] and [2, Lemma 7.17] that a regular ring  $R$  satisfies (\*) if and only if the maximal right quotient ring of  $R$  is *Type  $I_f$* .

NOTE. Let  $R$  be a regular ring satisfying (\*). If  $P$  is a finitely generated projective  $R$ -module, then  $\text{End}_R(P)$  is a regular ring satisfying (\*).

Proof. Choose a positive integer  $n$  and an idempotent matrix  $e \in M_n(R)$  such that  $e(nR_R) \simeq P$ . Then  $\text{End}_R(P) \simeq eM_n(R)e$ . Using [2, Corollary 10.5], we see that  $eM_n(Q(R))e \simeq Q(eM_n(R)e)$  is *Type  $I_f$* , where  $Q(R)$  is the maximal right quotient of  $R$ . Since  $eM_n(R)e \leq_e Q(eM_n(R)e)$  as an  $eM_n(R)e$ -module, we have that  $eM_n(R)e$  satisfies (\*), and so has  $\text{End}_R(P)$ .

NOTATIONS. Let  $A$ ,  $B$  and  $A_i$  ( $i \in I$ ) be  $R$ -modules, and  $k$  be a positive integer. Take  $x \in \prod A_i$ . Then we have some notations as following.

$A < B$  ;  $A$  is a submodule of  $B$ .

$A \lesssim B$  ;  $B$  has a submodule isomorphic to  $A$ .

$A < \oplus B$  ;  $A$  is a direct summand of  $B$ .

$A \lesssim \oplus B$  ;  $B$  has a direct summand isomorphic to  $A$ .

$A <_e B$  ;  $A$  is an essential submodule of  $B$ .

$A \lesssim_e B$  ;  $B$  has an essential submodule isomorphic to  $A$ .

$kA$  ; the  $k$ -copies of  $A$ .

$x(i)$  ; the  $i$ -th component of  $x$ .

$Q(R)$  ; the maximal right quotient ring of  $R$ .

## 2. The property (DF) for regular rings satisfying (\*)

**Lemma 2.1** ([2, Theorem 6.6]). *Let  $R$  be a regular ring whose primitive factor rings are artinian. Then  $R$  satisfies (\*).*

**Lemma 2.2.** *Let  $R$  be a regular ring satisfying (\*). Then there exist abelian regular rings  $\{S_t\}_{t \in T}$  and orthogonal central idempotents  $\{e_t\}_{t \in T}$  of  $R$  such that  $R_R \lesssim_e [\prod M_{n(t)}(S_t)]_R \oplus M_{n(t)}(S_t) \lesssim R$  and  $e_t R = M_{n(t)}(S_t)$ . Therefore  $Q(R) \simeq \prod M_{n(t)}(Q(S_t))$ .*

*Proof.* This theorem follows from [2, Lemma 7.17 and the proof of Theorem 7.18].

**Lemma 2.3.** *Let  $R$  be a regular ring of bounded index and  $P$  be a finitely generated projective  $R$ -module. Then  $P$  can not contain a family  $\{A_1, A_2, \dots\}$  of nonzero finitely generated submodules such that  $A_i \supsetneq A_{i+1}$  and  $iA_i \lesssim P$  for each  $i = 1, 2, \dots$ .*

*Proof.* By [2, Corollary 7.13], we see that  $\text{End}_R(P)$  has bounded index. Note the claim in the proof of [2, Theorem 6.6], and applying [5, Lemma 5] to  $\text{End}_R(P)$ , we see that this lemma holds.

**Theorem 2.4.** *Let  $R$  be a regular ring satisfying (\*), and  $P$  be a projective  $R$ -module with a cyclic decomposition  $P = \bigoplus_{i \in I} P_i$ . Then the following conditions (a)~(d) are equivalent:*

- (a)  $P$  is directly infinite.
- (b) There exists a nonzero cyclic projective  $R$ -module  $X$  such that  $\aleph_0 X \lesssim P$ .
- (c) There exists a nonzero cyclic projective  $R$ -module  $X$  such that  $X \lesssim \bigoplus_{i \in I - \{i_1, \dots, i_n\}} P_i$  for any finite subset  $\{i_1, \dots, i_n\}$  of  $I$ .
- (d) There exists a nonzero cyclic projective  $R$ -module  $X$  such that  $\aleph_0 X \lesssim \bigoplus P$ .

*Proof.* It is clear that (a)  $\rightarrow$  (b) and (c)  $\rightarrow$  (d)  $\rightarrow$  (a) hold, hence we shall prove that (b)  $\rightarrow$  (c) holds. We may assume  $\bigoplus M_{n(t)}(S_t) < R_R <_e [\prod M_{n(t)}(S_t)]_R$  for some set of abelian regular rings  $\{S_t\}_{t \in T}$  by Lemma 2.2. Now we assume that (b) holds, hence there exists a nonzero principal right ideal  $X$  of  $R$  such that  $\aleph_0 X \lesssim P$ . Let  $\{i_1, \dots, i_n\}$  be a subset of  $I$  and set  $I' = I - \{i_1, \dots, i_n\}$ . Since  $\bigoplus M_{n(t)}(S_t) < R_R <_e [\prod M_{n(t)}(S_t)]_R$ , there exists  $t' \in T$  such that  $Y = [(\prod_{t \neq t'} 0) \times M_{n(t')}(S_{t'})] \cap X \neq 0$ . By the property of regular ring, it is clear that  $Y$  is a principal right ideal of  $R$ . Then  $\aleph_0 Y \lesssim P$ , hence  $Y \lesssim \bigoplus P$ . Thus for each  $i \in I$ , we have decompositions  $P_i = P_i^1 \oplus P_i^{(1)}$  and  $Y \simeq P_{i_1}^1 \oplus \dots \oplus P_{i_n}^1 \oplus (\bigoplus_{i \in I'} P_i)$ . Set  $(\prod_{t \neq t'} 0) \times M_{n(t')}(S_{t'}) = S$ , and then there exists a central idempotent  $e$  in  $R$  such that  $eR = S$ .

Note that  $S$  is a regular ring of bounded index. It is clear that

$$Y \otimes_R S \simeq (P_{i_1}^1 \otimes_R S) \oplus \dots \oplus (P_{i_n}^1 \otimes_R S) \oplus [\bigoplus_{i \in I'} (P_i^1 \otimes_R S)]$$

and  $2Y \otimes_R S \lesssim P \otimes_R S$ . Since  $S$  is unit-regular,  $Y \otimes_R S$  has the cancellation property. Hence

$$Y \otimes_R S_S \lesssim \oplus (P_{i_1}^{(1)} \otimes_R S) \oplus \cdots \oplus (P_{i_n}^{(1)} \otimes_R S) \oplus [\oplus_{i \in I'} (P_i^{(1)} \otimes_R S_S)]$$

Thus for each  $i$ , we obtain that  $P_i^{(1)} \otimes_R S_S = \bar{P}_i^2 \oplus \bar{P}_i^{(2)}$  for each  $i \in I$  and

$$Y \otimes_R S_S \simeq \bar{P}_{i_1}^2 \oplus \cdots \oplus \bar{P}_{i_n}^2 \oplus (\oplus_{i \in I'} \bar{P}_i^2).$$

Continuing this procedure, we have that  $\bar{P}_i^{(m)} = \bar{P}_i^{m+1} \oplus \bar{P}_i^{(m+1)}$  and

$$Y \otimes_R S_S \simeq \bar{P}_{i_1}^{m+1} \oplus \cdots \oplus \bar{P}_{i_n}^{m+1} \oplus (\oplus_{i \in I'} \bar{P}_i^{m+1})$$

for each  $i \in I$  and each positive integer  $m$ .

Now we set  $A_m = \bar{P}_{i_1}^m \oplus \cdots \oplus \bar{P}_{i_n}^m$ , where  $A_1 = (P_{i_1}^{(1)} \otimes_R S) \oplus \cdots \oplus (P_{i_n}^{(1)} \otimes_R S)$ . Then  $A_1 \lesssim \oplus A_2 \oplus (\oplus_{i \in I'} \bar{P}_i^2)$ , hence there exist a direct summand  $B_2$  of  $A_2$  and a direct summand  $Q_i^2$  of  $\bar{P}_i^2$  such that  $A_1 \simeq B_2 \oplus (\oplus_{i \in I'} Q_i^2)$ . Continuing this procedure, we obtain a family  $\{B_1, B_2, \dots\}$  ( $A_1 = B_1$ ) of finitely generated projective  $S$ -submodules of  $(P_{i_1} \oplus \cdots \oplus P_{i_n}) \otimes_R S$  such that  $B_m \gtrsim B_{m+1}$  and  $mB_m \lesssim nS$  for all  $m$ . By Lemma 2.3, there exists a positive integer  $k$  such that  $B_m = 0$  for all  $m (\geq k)$ . Thus we have that  $A_1 \simeq (\oplus_{i \in I'} Q_i^2) \oplus \cdots \oplus (\oplus_{i \in I'} Q_i^k)$  and  $Y \otimes_R S_S \simeq (\oplus_{i \in I'} Q_i^1) \oplus \cdots \oplus (\oplus_{i \in I'} Q_i^k)$ . Noting that  $0 \neq Y < S$ , we have that  $Y_R \lesssim \oplus_{i \in I'} P_i$ .

**Corollary 2.5.** *Let  $R$  be a regular ring satisfying  $(*)$ . Then  $R$  contains no infinite direct sums of nonzero pairwise isomorphic right ideals. Hence  $R$  is directly finite.*

*Proof.* From Lemma 2.2, we may assume that,  $R_R < {}_e \Pi M_{n(t)}(S_t)$  for some abelian regular rings  $\{S_t\}_{t \in T}$ . Set  $T = \Pi M_{n(t)}(S_t)$ . Now we assume that  $R$  contains a direct sum of nonzero pairwise isomorphic right ideals, and so there exists a nonzero idempotent  $e$  of  $R$  such that  $0 \neq \aleph_0(eR) \lesssim R_R$ . Then  $\aleph_0(eR) \otimes_R T \lesssim R \otimes_R T$ , and so  $\aleph_0(eT) \lesssim T$ , which contradicts to Theorem 2.4 because  $T$  is a directly finite regular ring satisfying  $(*)$ .

**Theorem 2.6.** *Let  $R$  be a regular ring satisfying  $(*)$  and  $k$  be a positive integer. If  $P$  is a directly finite projective  $R$ -module, then so is  $kP$ .*

*Proof.* We may assume that  $\oplus M_{n(t)}(S_t) < R_R < {}_e [\Pi M_{n(t)}(S_t)]_R$  for some abelian regular rings  $\{S_t\}_{t \in T}$ , and let  $P = \oplus_{i \in I} P_i$  be a cyclic decomposition of  $P$ . It is sufficient to prove that this theorem holds in case  $k=2$ . Assume that  $2P$  is directly infinite. Then Theorem 2.4 follows that there exists a nonzero principal right ideal  $X$  of  $R$  such that  $X \lesssim \oplus_{i \in I - \{i_1, \dots, i_n\}} 2P_i$  for any finite subset  $\{i_1, \dots, i_n\}$  of  $I$ . By the proof of Theorem 2.4, we may assume that exists  $t'$  of  $T$  such that  $X < (\prod_{t \neq t'} 0) \times M_{n(t')}(S_{t'}) = S$ . For any finite subset  $\{i_1, \dots, i_n\}$  of  $I$ , we have that  $0 \neq X \otimes_R S_S \lesssim \oplus_{i \in I - \{i_1, \dots, i_n\}} (2P_i \otimes_R S)$ . Since  $S$  is a regular ring of bounded index, we see that  $2(P \otimes_R S)_S$  is directly infinite by Theorem 2.4 and so  $(P \otimes_R S)_S$  is directly

infinite by [5, Theorem 4]. Moreover, using Theorem 2.4 again, there exists a nonzero principal right ideal  $Y$  of  $S$  such that  $Y \lesssim \bigoplus_{i \in I - \{i_1, \dots, i_n\}} (P_i \otimes_R S_S)$  for any finite subset  $\{i_1, \dots, i_n\}$  of  $I$ . Considering  $Y$  as an  $R$ -module,  $0 \neq Y_R \lesssim \bigoplus_{i \in I - \{i_1, \dots, i_n\}} P_i$ . Therefore  $P$  is directly infinite, and so this theorem is complete.

**Corollary 2.7.** *Let  $R$  be a regular ring satisfying (\*). Then every finitely generated projective  $R$ -module is directly finite.*

*Proof.* It is clear by Corollary 2.5 and Theorem 2.6.

**Corollary 2.8.** *Let  $R$  be a regular ring satisfying (\*).*

(a)  $M_n(R)$  is directly finite for all positive integer  $n$ , and so  $M_n(R)$  contains no infinite direct sums of nonzero pairwise isomorphic right ideals.

(b) If  $P$  and  $Q$  are finitely generated projective  $R$ -modules, then  $P \oplus Q$  is directly finite.

*Proof.* (a)  $R$  is a regular ring satisfying (\*), and hence so is  $M_n(R)$ . Therefore Corollary 2.5 shows that (a) holds. (b) follows from Corollary 2.7.

**NOTE.** In [1], Chuang and Lee have shown that there exists a regular ring satisfying (\*) which is not unit-regular. Our Corollary 2.8 gives a partially solution for open problems 1 and 9 in Goodearl's book ([2]).

**DEFINITION.** Let  $R$  be a regular ring and  $P$  be a projective  $R$ -module. We call that  $P$  satisfies (#) provided that, for each nonzero finitely generated submodule  $I$  of  $P$  and any family  $\{A_1, B_1, \dots\}$  of submodules of  $P$  with

$$\begin{aligned} I &= A_1 \oplus B_1, \\ A_i &= A_{2i} \oplus B_{2i}, \\ B_i &= A_{2i+1} \oplus B_{2i+1} \quad \text{for each } i = 1, 2, \dots, \end{aligned}$$

there exists a nonzero projective  $R$ -module  $X$  such that  $X \lesssim \bigoplus_{i=m}^{\infty} A_i$  or  $X \lesssim \bigoplus_{i=m}^{\infty} B_i$  for any positive integer  $m$ .

**Lemma 2.9** ([5, Lemma 6]). *Let  $P$  be a nonzero finitely generated projective module over a regular ring  $R$ , and set  $T = \text{End}_R(P)$ . Then the following conditions are equivalent:*

- (a)  $P$  satisfies (#).
- (b)  $T$  satisfies (#) as a  $T$ -module.

**Lemma 2.10** ([5, Lemma 7]). *Let  $P$  be a nonzero finitely generated projective*

module over a regular ring  $R$ , and set  $T = \text{End}_R(P)$ . Then the following conditions are equivalent:

- (a)  $R$  satisfies (#) as an  $R$ -module.
- (b) All nonzero finitely generated projective  $R$ -modules satisfy (#).
- (c) For any positive integer  $k$ ,  $kR$  satisfies (#).
- (d) There exists a positive integer  $k$  such that  $kR$  satisfies (#).

**Theorem 2.11.** *Let  $R$  be a regular ring satisfying (\*). Then the following conditions are equivalent:*

- (a)  $R$  has (DF).
- (b)  $R$  satisfies (#) as an  $R$ -module.
- (c) For any nonzero finitely generated projective  $R$ -module  $P$ ,  $\text{End}_R(P)$  has (DF).
- (d) For any positive integer  $k$ ,  $M_k(R)$  has (DF).
- (e) There exists a positive integer  $k$  such that  $M_k(R)$  satisfies (DF).

*Proof.* Note that  $\text{End}_R(P)$  is a regular ring with (\*). [5, Theorem 8] was proved only using [5, Theorem 2]. Now [5, Theorem 2] holds on a regular ring satisfying (\*) by Theorem 2.4. Hence we see that this theorem holds under this condition using the similar proof of [5, Theorem 8] (Note that the unit-regularity is not needed).

### 3. Some applications

**Lemma 3.1.** *Let  $R$  be a regular ring satisfying (\*), and let  $\{e_i\}$  be a set of nonzero orthogonal central idempotents of  $R$  such that  $\oplus e_i R_R <_e R_R$ . Then  $R$  has (DF) if and only if  $e_i R$  has (DF) for all  $i$ .*

*Proof.* Note that  $e_i R$  is a ring direct summand of  $R$ . It is clear from Theorem 2.11 that “only if” part holds. We shall prove that “if” part holds. Let  $I$  be a nonzero direct summand of  $R$ , and so  $e_i R \cap I \neq 0$  for some  $i$ . Setting  $J = e_i R \cap I$ ,  $J$  is a principal right ideal of both  $R$  and  $e_i R$ . We consider decompositions

$$\begin{aligned} I &= A_1 \oplus B_1 \\ A_j &= A_{2j} \oplus B_{2j} \\ B_j &= A_{2j+1} \oplus B_{2j+1} \quad \text{for each } j=1,2,\dots, \end{aligned}$$

and so there exist decompositions of  $J$  such that

$$\begin{aligned} J &= C_1 \oplus D_1 \\ C_j &= C_{2j} \oplus D_{2j} \end{aligned}$$



$$D_j = C_{2j+1} \oplus D_{2j+1}$$

$$C_j \lesssim \oplus A_j \quad \text{and} \quad D_j \lesssim \oplus B_j \quad \text{for each } j=1,2,\dots$$

By the assumption, there exists a nonzero cyclic projective  $e_i R$ -module  $X$  such that  $X \lesssim \oplus_{j=m}^{\infty} C_j$  or  $X \lesssim \oplus_{j=m}^{\infty} D_j$  for each positive integer  $m$ . Hence  $X \otimes_{R e_i} R \lesssim \oplus_{j=m}^{\infty} (C_j \otimes_{R e_i} R)$  or  $X \otimes_{R e_i} R \lesssim \oplus_{j=m}^{\infty} (D_j \otimes_{R e_i} R)$ . Note that  $\oplus_{j=m}^{\infty} (C_j \otimes_{R e_i} R) \lesssim \oplus_{j=m}^{\infty} A_j$  and  $\oplus_{j=m}^{\infty} (D_j \otimes_{R e_i} R) \lesssim \oplus_{j=m}^{\infty} B_j$ . Therefore  $X \otimes_{R e_i} R \lesssim \oplus_{j=m}^{\infty} A_j$  or  $X \otimes_{R e_i} R \lesssim \oplus_{j=m}^{\infty} B_j$ . Since  $X \otimes_{R e_i} R \neq 0$ , this lemma has proved by Theorem 2.11.

**Lemma 3.2** ([6, Proposition 2.1]). *Let  $R$  be an abelian regular ring. If  $Q(R)$  has (DF), then so has  $R$ .*

**Theorem 3.3.** *Let  $R$  be a regular ring satisfying (\*). If  $Q(R)$  has (DF), then so does  $R$ .*

*Proof.* By Lemma 2.2, we may assume that there exists a set  $\{S_t\}$  of abelian regular rings such that  $R_R <_e [\Pi M_{n(t)}(S_t)]$ . Then  $Q(R) = \Pi M_{n(t)} Q(S_t)$ . Assume that  $Q(R)$  has (DF), then so does  $M_{n(t)}(Q(S_t))$  for all  $t$  by Lemma 3.1. Moreover, Theorem 2.11 shows that  $Q(S_t)$  also has (DF), hence so has  $S_t$  by Lemma 3.2. Thus  $M_{n(t)}(S_t)$  also has (DF) by Theorem 2.11. There exists the set  $\{e_t\}$  of orthogonal central idempotents of  $R$  such that  $e_t R = M_{n(t)}(S_t) \times [\Pi_{t \neq t'} 0]$  and  $\oplus e_t R <_e R_R$ . Therefore  $R$  has (DF) by Lemma 3.1.

Now we shall give an example of a regular ring with a zero socle satisfying (\*) which has (DF), as following.

**EXAMPLE 3.4.** Let  $F$  be a field, and set  $R = \Pi_{i=1}^{\infty} F_i (F_i = F)$  and  $\bar{R} = R / \text{soc}(R)$ . Then  $\bar{R}$  is a regular ring satisfying (\*) which has (DF).

*Proof.* Since it is clear that  $\bar{R}$  is a regular ring satisfying (\*), we shall prove that  $\bar{R}$  has (DF) using Theorem 2.11. Let  $\Psi$  be the natural map from  $R$  to  $\bar{R}$ , and let  $I$  be a nonzero direct summand of  $\bar{R}$  with following decompositions:

$$I = A_1 \oplus B_1$$

$$A_i = A_{2i} \oplus B_{2i}$$

$$B_i = A_{2i+1} \oplus B_{2i+1} \quad \text{for } i=1,2,\dots$$

Now assume that there does not exist  $\{C_j\}$  ( $C_j = A_j$  for some  $j$ ) which is an infinite subset of  $\{A_i\}_{i=1}^{\infty}$  such that  $C_j > C_{j+1}$  and  $C_j \neq 0$  for all  $j$ . Let  $\{D_p\}$  ( $D_p = A_i$  for some  $i$ ) be an infinite decreasing sequence of  $\{A_i\}$ , and so there exists a positive integer  $p'$  such that  $D_p = 0$  ( $p \leq p'$ ). Hence  $0 = D_{p'} = A_{i_1}$  for some

$i_1$ . Thus  $B_{i_1} \neq 0$ . Next, we take  $\{E_q\}$  ( $E_q = A_i$  for some  $i$ ) which is an infinite decreasing sequence of  $\{A_{ij}\}$ , where  $E_q < B_{i_1}$  and  $B_{k_q} < B_{i_1}$  ( $A_{k_q} = E_q$ ) for all positive integer  $q$ . Similarly, there exists a positive integer  $q'$  such that  $E_{q'} = 0$  ( $q' \leq q$ ). Hence there exists a positive integer  $i_2$  ( $i_2 > i_1$ ) such that  $E_{q'} = A_{i_2} = 0$ . Therefore  $B_{i_2} \neq 0$  and  $B_{i_1} > B_{i_2}$ . Continuing this procedure, we can get an infinite set  $\{B_{i_k}\}$  such that  $\{B_i\} \supset \{B_{i_k}\}$  and  $B_{i_k} \neq 0$  for all  $k$ . From the above, we may assume that there exists an infinite decreasing sequence  $\{C_j\}$  such that  $\{A_i\} \supset \{C_j\}$ ,  $C_j > C_{j+1}$  and  $C_j \neq 0$  for all  $j$ .

We have a set  $\{e_j\}$  of idempotents of  $R$  such that  $\Psi(e_j R) = C_j$  and  $e_j R \geq e_{j+1} R$  for all  $j$ . We take an idempotent  $f_1 (\in e_1 R)$  with  $\dim_F(f_1 R) = 1$ . Next we take an idempotent  $f_2 (\in e_2 R)$  such that  $\dim_F(f_2 R) = 1$  and  $f_1 f_2 = 0$ . Continuing this procedure, we can take a set  $\{f_j\}$  of orthogonal idempotents of  $R$ . Set  $e = \bigvee f_j$ , and then  $\Psi(e) \neq 0$ . We have that  $eR = J \oplus (eR \cap e_j R)$  and  $J \subset \bigoplus F_i$  for some right ideal  $J$ . Noting that  $J \otimes_R \bar{R} = 0$ , we have that

$$\begin{aligned} 0 \neq \Psi(e)\bar{R} &\simeq eR \otimes_R \bar{R} \\ &\simeq [J \oplus (eR \cap e_j R)] \otimes_R \bar{R} \\ &\lesssim e_j R \otimes_R \bar{R} \\ &\simeq C_j \quad \text{for all } j. \end{aligned}$$

Therefore  $0 \neq \Psi(e)\bar{R} \lesssim \bigoplus_{i=m}^{\infty} A_i$  for any positive integer  $m$ . Hence  $\bar{R}$  has (DF) by Theorem 2.11.

By Example 3.4, we have a problem that, for any regular ring  $S$ ,  $R = (\Pi_1^{\infty} S) / (\bigoplus S)$  satisfies (DF) or not. Example 3.5 shows that, even if  $S$  satisfies (\*),  $R$  does not satisfy (\*). Therefore we shall give the necessary and sufficient condition for that  $R$  satisfies (\*), and we solve the above problem under this condition.

**EXAMPLE 3.5.** Let  $F$  be a field and set  $S = \Pi_{n=1}^{\infty} M_n(F)$ ,  $\bar{S} = S / (\bigoplus M_n(F))$ ,  $T = \Pi_{i=1}^{\infty} S_i$  ( $S_i = S$ ) and  $R = T / (\bigoplus S_i)$ . Then  $S$  satisfies (\*), but  $R$  does not satisfy (\*).

**Proof.** It is clear that  $S$  satisfies (\*). Therefore we shall show that  $R$  does not satisfy (\*). Set a central idempotent  $e (\in T)$  as following:

$$e(n) = (0, \dots, 0, \underbrace{\begin{bmatrix} 1 & \\ & 1 \end{bmatrix}}_{\lfloor n-1 \rfloor}, 0, 0, \dots),$$

where  $e(n) \in S_n$ .

Let  $\Phi$  be the natural map from  $S$  to  $\bar{S}$ , and  $\rho$  be the natural map from  $T$  to  $R$ . Set  $\Psi = \rho|_{eT}$ . Noting that  $e(n)S_n \simeq M_n(F)$ , we have  $eT \simeq S$ . Hence there exists a ring

isomorphism  $\kappa$  from  $eT$  to  $S$ . Now, we define a ring homomorphism  $\alpha$  from  $\Psi(e)R$  to  $\bar{S}$  as following; for each  $x \in \Psi(e)R$ , we take any element  $y$  of  $\Psi^{-1}(x)$  and set  $\alpha(x) = \Phi\kappa(y)$ .

$$\begin{array}{ccc} \Psi(e)R & \xrightarrow{\alpha} & \bar{S} \\ \uparrow \psi & & \uparrow \Phi \\ eT & \xrightarrow{\kappa} & S \end{array}$$

Similarly we define a ring homomorphism  $\beta$  from  $\bar{S}$  to  $\Psi(e)R$ . Then we have that  $\beta\alpha = 1_{\Psi(e)R}$  and  $\alpha\beta = 1_{\bar{S}}$ . Hence  $\alpha$  and  $\beta$  are isomorphisms. Therefore  $\Psi(e)R \simeq \bar{S}$ . Let  $I$  be a nonzero two-sided ideal of  $\bar{S}$ , and so  $I = J/(\oplus S_i)$  for some nonzero two-sided ideal of  $S$  which contains  $\oplus S_i$ . There exists  $0 \neq x \in J - (\oplus S_i)$  with  $x(i) \neq 0$  for almost all  $i$ . Since  $S_i x(i) S_i = M_i(F)$  has index  $i$ , there exists a nonzero central idempotent  $e(i)$  of  $M_i(F)$  which  $S_i x(i) S_i$  has index  $i$ . Therefore  $SxS$  does not have bounded index, and so does not  $J/(\oplus S_i)$ . Therefore  $\bar{S}$  does not satisfy  $(*)$ , and hence so does not  $\Psi(e)R$ . Thus  $R$  does not satisfy  $(*)$ .

**Lemma 3.6.** *Let  $R$  be a ring, and  $e, f$  be idempotents of  $R$ . Then  $eR \simeq fR$  if and only if there exist  $u$  and  $v$  of  $R$  such that  $vu = e$  and  $uv = f$ .*

**Lemma 3.7.** *Let  $S$  be a regular ring which has index  $\infty$ , and set  $R = (\prod_{i=1}^{\infty} S_i) / (\oplus S_i)$  ( $S_i = S$ ). Then  $R$  has an infinite direct sum of nonzero pairwise isomorphic right ideals.*

*Proof.* Let  $\Psi$  be the natural map from  $\prod_{i=1}^{\infty} S_i$  to  $R$ . Since  $S$  has index  $\infty$ , there exists a set of idempotents  $\{e_{ij}\}_{i,j=1,2,\dots}$  as following:

$$\begin{aligned} e_{11}S \\ e_{21}S \simeq e_{22}S \\ e_{31}S \simeq e_{32}S \simeq e_{33}S \\ \dots \end{aligned}$$

, where  $e_{ij} = 0$  ( $i < j$ ) and  $\{e_{i1}, \dots, e_{ii}\}$  are nonzero orthogonal for all  $i$ . For all positive integer  $m$ , we take idempotents  $\{f_m\}$  such that  $f_m(k) = e_{km}$  for all positive integer  $k$ . Since  $e_{k1}S \simeq e_{k2}S$  for all  $k$ , there exist  $u_k$  and  $v_k$  of  $S$  such that  $u_kv_k = e_{k2}$  and  $v_ku_k = e_{k1}$  by Lemma 3.6. Set  $u$  and  $v$  of  $\prod_{i=1}^{\infty} S_i$  such that  $u(k) = u_k$  and  $v(k) = v_k$ . Then  $uv = f_2$  and  $vu = f_1 - e$ , where  $e$  is an idempotent with  $e(1) = e_{11}$  and  $e(k) = 0$  ( $k \neq 1$ ). Hence  $(f_1 - e)(\prod S_i) \simeq f_2(\prod S_i)$  and  $(f_1 - e)(\prod S_i) \cap f_2(\prod S_i) = 0$ . Therefore we see from Lemma 3.6 that  $\Psi(f_1 - e)R \simeq \Psi(f_2)R$  and  $\Psi(f_1 - e)R \cap \Psi(f_2)R = 0$ . Since  $\Psi(f_1 - e)R = \Psi(f_1)R$ , we have that  $\Psi(f_1)R \simeq \Psi(f_2)R$  and  $\Psi(f_1)R \cap \Psi(f_2)R = 0$ .

$=0$ . Continuing this procedure, for all positive integers  $i$  and  $j$ ,  $\Psi(f_i)R \simeq \Psi(f_j)R$  and  $\Psi(f_i) \cap \Psi(f_j)R = 0$  ( $i \neq j$ ). Thus  $R$  has an infinite direct sum of nonzero pairwise isomorphic right ideals.

**Theorem 3.8.** *Let  $S$  be a regular ring, and set  $R = (\prod_{i=1}^{\infty} S_i) / (\oplus S_i)$  ( $S_i = S$ ). Then the following conditions are equivalent:*

- (a)  $R$  satisfies (\*).
- (b)  $R$  is a regular ring whose primitive factor rings are artinian.
- (c)  $R$  has bounded index.
- (d)  $R$  contains no infinite direct sums of nonzero pairwise isomorphic right ideals.
- (e)  $S$  has bounded index.

**Proof.** It is clear by Lemma 3.7 that (d)  $\rightarrow$  (e)  $\rightarrow$  (c)  $\rightarrow$  (b)  $\rightarrow$  (a) hold. (a)  $\rightarrow$  (d) follows from Corollary 2.5. Therefore this theorem is complete.

**Theorem 3.9.** *Let  $S$  be a regular ring of bounded index. Set  $R = (\prod_{n=1}^{\infty} S_n) / (\oplus S_n)$  ( $S_n = S$ ). Then  $R$  has (DF).*

**Proof.** Set  $\prod_{n=1}^{\infty} S_n = T$ , and let  $\Psi$  be the natural map from  $T$  to  $R$ . Let  $I$  be a nonzero direct summand of  $R$  with following decompositions:

$$\begin{aligned} I &= A_1 \oplus B_1 \\ A_i &= A_{2i} \oplus B_{2i} \\ B_i &= A_{2i+1} \oplus B_{2i+1} \quad \text{for } i = 1, 2, \dots \end{aligned}$$

Similarly to the proof of Example 3.4, we may assume that there exists an infinite subset  $\{C_j\}$  of  $\{A_i\}$  ( $C_j = A_i$  for some  $i$ ) such that  $C_j > C_{j+1}$  and  $C_j \neq 0$  for all positive integer  $j$ . We have the set of idempotents  $\{e_j\}$  of  $T$  such that  $\Psi(e_j T) = C_j$  and  $e_j T > e_{j+1} T$ . Set  $J_n = S_n \times (\prod_{i \neq n} 0)$ . Then,  $J_{n_1} \cap e_1 T \neq 0$  for some positive integer  $n_1$ . There exists a nonzero idempotent  $f_1 \in T$  such that  $f_1 T = J_{n_1} \cap e_1 T$ . Next we have a nonzero idempotent  $f_2 \in T$  for some  $n_2$  ( $> n_1$ ) such that  $f_2 R = J_{n_2} \cap e_2 R$ . Continuing this procedure, we have the set  $\{f_j\}$  of orthogonal idempotents of  $T$ . Now, we set an idempotent  $g$  of  $T$  as following;

$$\begin{aligned} g(n_j) &= f_j(n_j) = e_j(n_j) \\ g(k) &= 0 \quad (k \notin \{n_j\}). \end{aligned}$$

Put  $K_j = f_1 T \oplus \dots \oplus f_{j-1} T$  for all  $j$ . Then  $gT = K_j \oplus (gT \cap e_j T)$ . Noting  $K_j \otimes_T R = 0$ , we have that

$$\begin{aligned}
0 \neq \Psi(g)R &\simeq gT \otimes_T R \\
&\simeq [K_j \oplus (gT \cap e_j T)] \otimes_T R \\
&\simeq (gT \cap e_j T) \otimes_T R \\
&\lesssim e_j T \otimes_T R \\
&\simeq C_j \quad \text{for all } j.
\end{aligned}$$

From the above, we have that  $\Psi(g)R \lesssim \bigoplus_{i=m}^{\infty} A_i$  for any positive integer  $m$ . Therefore  $R$  has (DF) by Theorem 2.11.

---

#### References

- [1] C.-L. Chuang and P.-H. Lee: *On regular subdirect products of simple artinian ring*. Pacific J. Math. **142** (1990), 17–21.
- [2] K.R. Goodearl: *Von Neumann regular rings*, Kreiger, Florida, 1991.
- [3] K.R. Goodearl and J. Moncasi: *Cancellation of finitely generated modules over regular rings*, Osaka J. Math. **26** (1989), 679–685.
- [4] H. Kambara and S. Kobayahi: *On regular self-injective rings*, Osaka J. Math. **22** (1985), 71–79.
- [5] M. Kutami: *Projective modules over regular rings of bounded index*, Math. J. Okayama Univ. **30** (1988), 53–62.
- [6] M. Kutami and I. Inoue: *The property (DF) for regular rings whose primitive factor rings are artinian*, Math. J. Okayama Univ. **35** (1993), 169–179.

Department of Mathematics  
Yamaguchi University  
Yoshida, Yamaguchi 753  
Japan