

Title	Remark on the rank of elliptic curves
Author(s)	Nikolaev, Igor
Citation	Osaka Journal of Mathematics. 46(2) P.515-P.527
Issue Date	2009-06
Text Version	publisher
URL	https://doi.org/10.18910/7110
DOI	10.18910/7110
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

REMARK ON THE RANK OF ELLIPTIC CURVES

IGOR NIKOLAEV

(Received August 17, 2007, revised February 15, 2008)

Abstract

A covariant functor on the elliptic curves with complex multiplication is constructed. The functor takes values in the noncommutative tori with real multiplication. A conjecture on the rank of an elliptic curve is formulated.

Introduction

A. Let $0 < \theta < 1$ be an irrational number given by the regular continued fraction

$$\theta = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}} = [a_0, a_1, a_2, \dots].$$

Consider an AF -algebra, \mathbb{A}_θ , defined by the Bratteli diagram of Fig. 1, where a_i indicate the multiplicity of the edges of the graph. (For a definition of the AF -algebras and their Bratteli diagrams, we refer the reader to [2], or §1.2.) For the simplicity, we shall say that \mathbb{A}_θ is a noncommutative torus. Note that the classical definition of a noncommutative torus is slightly different but equivalent from the standpoint of the K -theory [3], [7], [13]. The \mathbb{A}_θ is said to have real multiplication, if θ is a quadratic irrationality. Recall that the noncommutative tori $\mathbb{A}_\theta, \mathbb{A}_{\theta'}$ are stably isomorphic whenever $\mathbb{A}_\theta \otimes \mathcal{K} \cong \mathbb{A}_{\theta'} \otimes \mathcal{K}$, where \mathcal{K} is the C^* -algebra of the compact operators. It is known that $\mathbb{A}_\theta, \mathbb{A}_{\theta'}$ are stably isomorphic if and only if $\theta' \equiv \theta \pmod{GL(2, \mathbb{Z})}$, i.e. $\theta' = (a\theta + b)/(c\theta + d)$, where $a, b, c, d \in \mathbb{Z}$ and $ad - bc = \pm 1$.

B. Let $\Lambda = \omega_1\mathbb{Z} + \omega_2\mathbb{Z}$ be a lattice in the complex plane \mathbb{C} . Recall that Λ defines an elliptic curve $E(\mathbb{C}): y^2 = 4x^3 - g_2x - g_3$ via the complex analytic map $\mathbb{C}/\Lambda \rightarrow E(\mathbb{C})$ given by the formula $z \mapsto (\wp(z, \Lambda), \wp'(z, \Lambda))$, where $g_2 = 60 \sum_{\omega \in \Lambda^\times} \omega^{-4}$, $g_3 = 140 \sum_{\omega \in \Lambda^\times} \omega^{-6}$, $\Lambda^\times = \Lambda - \{0\}$ and

$$\wp(z, \Lambda) = \frac{1}{z^2} + \sum_{\omega \in \Lambda^\times} \left(\frac{1}{(z - \omega)^2} - \frac{1}{\omega^2} \right)$$




Fig. 1. The AF-algebra \mathbb{A}_θ .

is the Weierstrass \wp function. We identify the elliptic curves $E(\mathbb{C})$ with the complex tori \mathbb{C}/Λ . If $\tau = \omega_2/\omega_1$, then $E_\tau(\mathbb{C}), E_{\tau'}(\mathbb{C})$ are isomorphic whenever $\tau' \equiv \tau \pmod{GL(2, \mathbb{Z})}$. The endomorphism ring $End(\mathbb{C}/\Lambda)$ is isomorphic either to \mathbb{Z} or to an order in the imaginary quadratic number field k [14]. In the second case, we say that the elliptic curve has a complex multiplication and denote such a curve by E_{CM} .

C. Consider the cubic $E_\lambda: y^2 = x(x-1)(x-\lambda)$, $\lambda \in \mathbb{C} - \{0, 1\}$. The j -invariant of E_λ is given by the formula $j(E_\lambda) = 2^6(\lambda^2 - \lambda + 1)^3 \lambda^{-2}(\lambda - 1)^{-2}$. To find λ corresponding to an elliptic curve with the complex multiplication, one has to solve the polynomial equation $j(E_{CM}) = j(E_\lambda)$ with respect to λ . Since $j(E_{CM})$ is an algebraic integer, $\lambda_{CM} \in K$, where K is an algebraic extension (of the degree at most six) of the field $\mathbb{Q}(j(E_{CM}))$. Thus, each E_{CM} is isomorphic to the cubic $y^2 = x(x-1)(x-\lambda_{CM})$ defined over the field K . The Mordell-Weil theorem says that the set of the K -rational points of E_{CM} is a finitely generated abelian group, whose rank we shall denote by $rk(E_{CM})$.

D. Let \mathcal{E} be a category whose objects are elliptic curves and the arrows are isomorphisms of the elliptic curves. Likewise, let \mathcal{A} be a category whose objects are noncommutative tori and the arrows are stable isomorphisms of the noncommutative tori. Our main goals can be expressed as follows.

OBJECTIVES. (i) to construct a functor (if any) $F: \mathcal{E} \rightarrow \mathcal{A}$, which maps isomorphic elliptic curves to the stably isomorphic noncommutative tori; (ii) to study the range of F on the elliptic curves with complex multiplication and (iii) to interpret the invariants of the stable isomorphism classes of the noncommutative tori in terms of the arithmetic invariants of the elliptic curves.

In the course of this note, we were able to obtain an answer to (i) and (ii), while (iii) generates a conjecture. Namely, a covariant non-injective functor $F: \mathcal{E} \rightarrow \mathcal{A}$, which maps isomorphic elliptic curves to the stably isomorphic noncommutative tori, is constructed (Lemma 1). It is proved that F sends the elliptic curves with complex multiplication to the noncommutative tori with real multiplication (Theorem 1). Finally, a conjecture on the rank of an elliptic curve with the complex multiplication is formulated (§3). The functor F has been studied by Kontsevich [5] (e.g. §1.39), Manin [6], Polishchuk [8]–[11], Polishchuk-Schwarz [12], Soibelman [15] and [16], Soibelman-Vologodsky [17], Taylor [18] and [19] et al. Our terminology is freely and gratefully borrowed from the above works.

E. The existence and properties of F are part of a Hodge theory for the measured foliations on a closed surface. Such a theory has been developed by Hubbard and Masur [4], who were inspired by the works of Thurston [20]. We shall give in §1 a brief account of the Hubbard-Masur-Thurston theory and the explicit formulas for the functor F . At the heart of the construction is a diagram:

$$F: \mathcal{E} \xrightarrow{h} \mathbb{R}^2 \xrightarrow{\pi} \mathbb{R}P^1 \cong \mathcal{A},$$

where h is a bijection and π is a projection map. For the sake of brevity, let $Isom(E) = \{E' \in \mathcal{E} \mid E' \cong E\}$ be the isomorphism class of an elliptic curve E , $h(Isom(E)) = \mu_E(\mathbb{Z} + \theta_E\mathbb{Z}) := \mathfrak{m}_E \subset \mathbb{R}$ be a \mathbb{Z} -module and $F(E) = \mathbb{A}_{\theta_E}$. A summary of our results can be formulated as follows.

Lemma 1. *Let $\varphi: E \rightarrow E'$ be an isogeny of the elliptic curves. Then $\theta' \equiv \theta \pmod{M_2(\mathbb{Z})}$, where $M_2(\mathbb{Z})$ is an integer matrix of the rank 2. In particular, F maps the isomorphic elliptic curves to the stably isomorphic noncommutative tori.*

Theorem 1. *Let $E \in Isom(E_{CM})$. Then there exists an h , such that:*

- (i) \mathfrak{m}_E is a full module in the real quadratic number field;
- (ii) \mathfrak{m}_E is an invariant of the class $Isom(E_{CM})$.

In particular, θ_E is a quadratic irrationality.

The structure of the note is as follows. In Section 1, we introduce the notation and some preliminary facts. Lemma 1 and Theorem 1 are proved in Section 2. In Section 3, a conjecture on the rank of an elliptic curve is formulated.

1. Preliminaries

This section contains a summary of measured foliations, AF -algebras and the functor F . The reader is encouraged to consult [2] (operator algebras) and [4] (measured foliations and Teichmüller space) for a systematic account.

1.1. Measured foliations and $T(g)$.

A. A measured foliation, \mathcal{F} , on a surface X is a partition of X into the singular points x_1, \dots, x_n of order k_1, \dots, k_n and the regular leaves (1-dimensional submanifolds). On each open cover U_i of $X - \{x_1, \dots, x_n\}$ there exists a non-vanishing real-valued closed 1-form ϕ_i such that

- (i) $\phi_i = \pm\phi_j$ on $U_i \cap U_j$;
- (ii) at each x_i there exists a local chart $(u, v): V \rightarrow \mathbb{R}^2$ such that for $z = u + iv$, it holds $\phi_i = \text{Im}(z^{k_i/2} dz)$ on $V \cap U_i$ for some branch of $z^{k_i/2}$.

The pair (U_i, ϕ_i) is called an atlas for the measured foliation \mathcal{F} . Finally, a measure μ is assigned to each segment $(t_0, t) \in U_i$, which is transverse to the leaves of \mathcal{F} , via the integral $\mu(t_0, t) = \int_{t_0}^t \phi_i$. The measure is invariant along the leaves of the foliation \mathcal{F} , hence the name.

B. Let S be a Riemann surface, and $q \in H^0(S, \Omega^{\otimes 2})$ a holomorphic quadratic differential on S . The lines $\operatorname{Re} q = 0$ and $\operatorname{Im} q = 0$ define a pair of measured foliations on R , which are transversal to each other outside the set of singular points. The set of singular points is common to the both foliations and coincides with the zeroes of q . The above measured foliations are said to represent the vertical and horizontal trajectory structure of q , respectively.

C. Let $T(g)$ be the Teichmüller space of the topological surface X of genus g , i.e. the space of complex structures on X . Consider the vector bundle $p: \mathcal{Q} \rightarrow T(g)$ over $T(g)$ whose fiber above a point $S \in T(g)$ is the vector space $H^0(S, \Omega^{\otimes 2})$. Given non-zero $q \in \mathcal{Q}$ above S , one can consider the horizontal measured foliation $\mathcal{F}_q \in \Phi_X$ of the quadratic differential q , where Φ_X is the space of (equivalence classes of) measured foliations on X . If $\{0\}$ is the zero section of \mathcal{Q} , the above construction defines a map $\mathcal{Q} - \{0\} \rightarrow \Phi_X$. For any $\mathcal{F} \in \Phi_X$, let $E_{\mathcal{F}} \subset \mathcal{Q} - \{0\}$ be the fiber above \mathcal{F} . In other words, $E_{\mathcal{F}}$ is a subspace of the holomorphic quadratic differentials, whose horizontal trajectory structure coincides with the measured foliation \mathcal{F} .

Theorem ([4]). *The restriction $E_{\mathcal{F}} \rightarrow T(g)$ of p to $E_{\mathcal{F}}$ is a homeomorphism.*

D. Let Φ_X be the space of measured foliations on the topological surface X . Following Douady and Hubbard [1], we shall consider a coordinate system on Φ_X , suitable for the construction of the functor F . For clarity, let us make a generic assumption that $q \in H^0(S, \Omega^{\otimes 2})$ is a holomorphic quadratic differential with the simple zeroes only. We wish to construct a Riemann surface of \sqrt{q} , which is a double cover of S with the ramification over the zeroes of q . Such a surface, denoted by \tilde{S} , is unique and has an advantage of carrying a holomorphic differential ω , such that $\omega^2 = q$. Denote by $\pi: \tilde{S} \rightarrow S$ a covering projection. The vector space $H^0(\tilde{S}, \Omega)$ splits into the direct sum $H_{\text{even}}^0(\tilde{S}, \Omega) \oplus H_{\text{odd}}^0(\tilde{S}, \Omega)$ in view of the involution π^{-1} of \tilde{S} , and the vector space $H^0(S, \Omega^{\otimes 2}) \cong H_{\text{odd}}^0(\tilde{S}, \Omega)$. Let $H_1^{\text{odd}}(\tilde{S})$ be an odd part of the homology of \tilde{S} relatively the zeroes of q . Consider a pairing $H_1^{\text{odd}}(\tilde{S}) \times H^0(S, \Omega^{\otimes 2}) \rightarrow \mathbb{C}$, defined by the integration $(\gamma, q) \mapsto \int_{\gamma} \omega$. Take the associated map $\psi_q: H^0(S, \Omega^{\otimes 2}) \rightarrow \operatorname{Hom}(H_1^{\text{odd}}(\tilde{S}); \mathbb{C})$ and let $h_q = \operatorname{Re} \psi_q$.

Theorem ([1]). *The map $h_q: H^0(S, \Omega^{\otimes 2}) \rightarrow \operatorname{Hom}(H_1^{\text{odd}}(\tilde{S}); \mathbb{R})$ is an \mathbb{R} -isomorphism.*

Since each $\mathcal{F} \in \Phi_X$ is the vertical foliation $Re q = 0$ for a $q \in H^0(S, \Omega^{\otimes 2})$, the theorem implies that $\Phi_X \cong Hom(H_1^{\text{odd}}(\tilde{S}); \mathbb{R})$. By the formulas for the relative homology:

$$H_1^{\text{odd}}(\tilde{S}) \cong \mathbb{Z}^n, \quad \text{where } n = \begin{cases} 6g - 6, & \text{if } g \geq 2, \\ 2, & \text{if } g = 1. \end{cases}$$

Thus, if $\{\gamma_1, \dots, \gamma_n\}$ is a basis in $H_1^{\text{odd}}(\tilde{S})$, the reals $\lambda_i = \int_{\gamma_i} Re \omega$ are natural coordinates in the space Φ_X [1].

1.2. AF-algebras.

A. The C^* -algebra is an algebra A over \mathbb{C} with a norm $a \mapsto \|a\|$ and an involution $a \mapsto a^*$ such that it is complete with respect to the norm and $\|ab\| \leq \|a\| \|b\|$ and $\|a^*a\| = \|a\|^2$ for all $a, b \in A$. If A is commutative, then the Gelfand theorem says that A is isomorphic to the C^* -algebra $C_0(X)$ of continuous complex-valued functions on a locally compact Hausdorff space X . For otherwise, A represents a noncommutative topological space X .

B. Let A be a C^* -algebra deemed as a noncommutative topological space. One can ask when two such topological spaces A, A' are homeomorphic? To answer the question, let us recall the topological K -theory. If X is a (commutative) topological space, denote by $V_{\mathbb{C}}(X)$ an abelian monoid consisting of the isomorphism classes of the complex vector bundles over X endowed with the Whitney sum. The abelian monoid $V_{\mathbb{C}}(X)$ can be made to an abelian group, $K(X)$, using the Grothendieck completion. The covariant functor $F: X \rightarrow K(X)$ is known to map the homeomorphic topological spaces X, X' to the isomorphic abelian groups $K(X), K(X')$. Let now A, A' be the C^* -algebras. If one wishes to define a homeomorphism between the noncommutative topological spaces A and A' , it will suffice to define an isomorphism between the abelian monoids $V_{\mathbb{C}}(A)$ and $V_{\mathbb{C}}(A')$ as suggested by the topological K -theory. The rôle of the complex vector bundle of degree n over the C^* -algebra A is played by a C^* -algebra $M_n(A) = A \otimes M_n$, i.e. the matrix algebra with the entries in A . The abelian monoid $V_{\mathbb{C}}(A) = \bigcup_{n=1}^{\infty} M_n(A)$ replaces the monoid $V_{\mathbb{C}}(X)$ of the topological K -theory. Therefore, the noncommutative topological spaces A, A' are homeomorphic, if $V_{\mathbb{C}}(A) \cong V_{\mathbb{C}}(A')$ are isomorphic abelian monoids. The latter equivalence is called a *stable isomorphism* of the C^* -algebras A and A' and is formally written as $A \otimes \mathcal{K} \cong A' \otimes \mathcal{K}$, where $\mathcal{K} = \bigcup_{n=1}^{\infty} M_n$ is the C^* -algebra of compact operators. Roughly speaking, the stable isomorphism between the C^* -algebras A and A' means that A and A' are homeomorphic as the noncommutative topological spaces.

C. Let A be a unital C^* -algebra and $V(A)$ be the union (over n) of projections in the $n \times n$ matrix C^* -algebra with entries in A . Projections $p, q \in V(A)$ are equivalent if there exists a partial isometry u such that $p = u^*u$ and $q = uu^*$. The equivalence class

of projection p is denoted by $[p]$. The equivalence classes of orthogonal projections can be made to a semigroup by putting $[p]+[q]=[p+q]$. The Grothendieck completion of this semigroup to an abelian group is called a K_0 -group of algebra A . Functor $A \rightarrow K_0(A)$ maps a category of unital C^* -algebras into the category of abelian groups so that projections in algebra A correspond to a positive cone $K_0^+ \subset K_0(A)$ and the unit element $1 \in A$ corresponds to an order unit $u \in K_0(A)$. The ordered abelian group (K_0, K_0^+, u) with an order unit is called a *dimension group*.

D. An *AF-algebra* (approximately finite C^* -algebra) is defined to be the norm closure of an ascending sequence of the finite dimensional C^* -algebras M_n 's, where M_n is the C^* -algebra of the $n \times n$ matrices with the entries in \mathbb{C} . Here the index $n = (n_1, \dots, n_k)$ represents a semi-simple matrix algebra $M_n = M_{n_1} \oplus \dots \oplus M_{n_k}$. The ascending sequence mentioned above can be written as

$$M_1 \xrightarrow{\varphi_1} M_2 \xrightarrow{\varphi_2} \dots,$$

where M_i are the finite dimensional C^* -algebras and φ_i the homomorphisms between such algebras. The set-theoretic limit $A = \lim M_n$ has a natural algebraic structure given by the formula $a_m + b_k \rightarrow a + b$; here $a_m \rightarrow a$, $b_k \rightarrow b$ for the sequences $a_m \in M_m$, $b_k \in M_k$. The homomorphisms φ_i can be arranged into a graph as follows. Let $M_i = M_{i_1} \oplus \dots \oplus M_{i_k}$ and $M_{i'} = M_{i'_1} \oplus \dots \oplus M_{i'_k}$ be the semi-simple C^* -algebras and $\varphi_i: M_i \rightarrow M_{i'}$ the homomorphism. One has the two sets of vertices V_{i_1}, \dots, V_{i_k} and $V_{i'_1}, \dots, V_{i'_k}$ joined by the a_{rs} edges, whenever the summand M_{i_r} contains a_{rs} copies of the summand $M_{i'_s}$ under the embedding φ_i . As i varies, one obtains an infinite graph called a *Bratteli diagram* of the *AF*-algebra.

E. By \mathbb{A}_θ we denote an *AF*-algebra given by the Bratteli diagram of Fig. 1. It is known that $K_0(\mathbb{A}_\theta) \cong \mathbb{Z}^2$ and $K_0^+(\mathbb{A}_\theta) = \{(p, q) \in \mathbb{Z}^2 \mid p + \theta q \geq 0\}$. The *AF*-algebras $\mathbb{A}_\theta, \mathbb{A}_{\theta'}$ are stably isomorphic, i.e. $\mathbb{A}_\theta \otimes \mathcal{K} \cong \mathbb{A}_{\theta'} \otimes \mathcal{K}$, if and only if $\mathbb{Z} + \theta\mathbb{Z} = \mathbb{Z} + \theta'\mathbb{Z}$ as the subsets of \mathbb{R} .

1.3. The functor F .

A. The Hubbard-Masur theory (§1.1) has been treated in a general setting so far. From now on, we switch to the case $g = 1$ (complex torus). Notice that $S = \tilde{S} \cong T^2$, since every holomorphic quadratic differential q on the complex torus is the square of a holomorphic differential ω .




Fig. 2. The measured foliation \mathcal{F} on $T^2 = \mathbb{R}^2/\mathbb{Z}^2$.

B. Let $\phi = \text{Re } \omega$ be a 1-form defined by ω . Since ω is holomorphic, ϕ is a closed 1-form on T^2 . The \mathbb{R} -isomorphism $h_q: H^0(S, \Omega) \rightarrow \text{Hom}(H_1(T^2); \mathbb{R})$, as explained, is given by the formulas:

$$\begin{cases} \lambda_1 = \int_{\gamma_1} \phi, \\ \lambda_2 = \int_{\gamma_2} \phi, \end{cases}$$

where $\{\gamma_1, \gamma_2\}$ is a basis in the first homology group of T^2 . We further assume that, after a proper choice of the basis, λ_1, λ_2 are positive real numbers.

C. Denote by Φ_{T^2} the space of measured foliations on T^2 . Each $\mathcal{F} \in \Phi_{T^2}$ is (measure) equivalent to a foliation by a family of the parallel lines of a slope θ and the invariant (transverse) measure μ (Fig. 2).

We use the notation \mathcal{F}_θ^μ for such a foliation. There exists a simple relationship between the reals (λ_1, λ_2) and (θ, μ) . Indeed, the closed 1-form $\phi = \text{Const}$ defines a measured foliation, \mathcal{F}_θ^μ , so that

$$\begin{cases} \lambda_1 = \int_{\gamma_1} \phi = \int_0^1 \mu \, dx \\ \lambda_2 = \int_{\gamma_2} \phi = \int_0^1 \mu \, dy \end{cases}, \quad \text{where } \frac{dy}{dx} = \theta.$$

By the integration:

$$\begin{cases} \lambda_1 = \int_0^1 \mu \, dx = \mu, \\ \lambda_2 = \int_0^1 \mu \theta \, dx = \mu \theta. \end{cases}$$

Thus, one gets $\mu = \lambda_1$ and $\theta = \lambda_2/\lambda_1$.

D. Recall that the Hubbard-Masur theory (§1.1.C) establishes a homeomorphism $h: T_S(1) \rightarrow \Phi_{T^2}$, where $T_S(1) \cong \mathbb{H} = \{\tau: \text{Im } \tau > 0\}$ is the Teichmüller space of the torus. Denote by ω_N an invariant (Néron) differential of the complex torus $\mathbb{C}/(\omega_1\mathbb{Z} + \omega_2\mathbb{Z})$.

It is well known that $\omega_1 = \int_{\gamma_1} \omega_N$ and $\omega_2 = \int_{\gamma_2} \omega_N$, where γ_1 and γ_2 are the meridians of the torus. Let π be a projection acting by the formula $(\theta, \mu) \mapsto \theta$. An explicit formula for the functor $F: \mathcal{E} \rightarrow \mathcal{A}$ is given by the composition: $F = \pi \circ h$, where h is the Hubbard-Masur homeomorphism. In other words, one gets the following (explicit) correspondence between the complex and noncommutative tori:

$$E_\tau = E_{(\int_{\gamma_2} \omega_N) / (\int_{\gamma_1} \omega_N)} \xrightarrow{h} \mathcal{F}_{(\int_{\gamma_2} \phi) / (\int_{\gamma_1} \phi)} \xrightarrow{h} \mathbb{A}_{(\int_{\gamma_2} \phi) / (\int_{\gamma_1} \phi)} = \mathbb{A}_\theta,$$

where $E_\tau = \mathbb{C} / (\mathbb{Z} + \tau\mathbb{Z})$.

2. Proof

2.1. Proof of Lemma 1. Let $\varphi: E_\tau \rightarrow E_{\tau'}$ be an isogeny of the elliptic curves. The action of φ on the homology basis $\{\gamma_1, \gamma_2\}$ of T^2 is given by the formulas:

$$(1) \quad \begin{cases} \gamma'_1 = a\gamma_1 + b\gamma_2, \\ \gamma'_2 = c\gamma_1 + d\gamma_2, \end{cases} \quad \text{where} \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Z}).$$

Recall that the functor $F: \mathcal{E} \rightarrow \mathcal{A}$ is given by the formula:

$$(2) \quad \tau = \frac{\int_{\gamma_2} \omega_N}{\int_{\gamma_1} \omega_N} \mapsto \theta = \frac{\int_{\gamma_2} \phi}{\int_{\gamma_1} \phi},$$

where ω_N is an invariant differential on E_τ and $\phi = \operatorname{Re} \omega$ is a closed 1-form on T^2 .

(i) From the left-hand side of (2), one obtains

$$(3) \quad \begin{cases} \omega'_1 = \int_{\gamma'_1} \omega_N = \int_{a\gamma_1 + b\gamma_2} \omega_N = a \int_{\gamma_1} \omega_N + b \int_{\gamma_2} \omega_N = a\omega_1 + b\omega_2, \\ \omega'_2 = \int_{\gamma'_2} \omega_N = \int_{c\gamma_1 + d\gamma_2} \omega_N = c \int_{\gamma_1} \omega_N + d \int_{\gamma_2} \omega_N = c\omega_1 + d\omega_2, \end{cases}$$

and therefore $\tau' = \int_{\gamma'_2} \omega_N / \int_{\gamma'_1} \omega_N = (c + d\tau) / (a + b\tau)$.

(ii) From the right-hand side of (2), one obtains

$$(4) \quad \begin{cases} \lambda'_1 = \int_{\gamma'_1} \phi = \int_{a\gamma_1 + b\gamma_2} \phi = a \int_{\gamma_1} \phi + b \int_{\gamma_2} \phi = a\lambda_1 + b\lambda_2, \\ \lambda'_2 = \int_{\gamma'_2} \phi = \int_{c\gamma_1 + d\gamma_2} \phi = c \int_{\gamma_1} \phi + d \int_{\gamma_2} \phi = c\lambda_1 + d\lambda_2, \end{cases}$$

and therefore $\theta' = \int_{\gamma'_2} \phi / \int_{\gamma'_1} \phi = (c + d\theta) / (a + b\theta)$. Comparing (i) and (ii), one gets the conclusion of the first part of Lemma 1. To prove the second part, recall that

the invertible isogeny is an isomorphism of the elliptic curves. In this case $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathbb{Z})$ and $\theta' = \theta \pmod{GL_2(\mathbb{Z})}$. Therefore F sends the isomorphic elliptic curves to the stably isomorphic noncommutative tori. The second part of Lemma 1 is proved.

It follows from the proof that $F: \mathcal{E} \rightarrow \mathcal{A}$ is a covariant functor. Indeed, F preserves the morphisms and does not reverse the arrows: $F(\varphi_1\varphi_2) = \varphi_1\varphi_2 = F(\varphi_1)F(\varphi_2)$ for any pair of the isogenies $\varphi_1, \varphi_2 \in \text{Mor}(\mathcal{E})$. □

2.2. Proof of Theorem 1. The following lemma will be helpful.

Lemma 2. *Let $\mathfrak{m} \subset \mathbb{R}$ be a module of the rank 2, i.e. $\mathfrak{m} = \mathbb{Z}\lambda_1 + \mathbb{Z}\lambda_2$, where $\theta = \lambda_2/\lambda_1 \notin \mathbb{Q}$. If $\mathfrak{m}' \subseteq \mathfrak{m}$ is a submodule of the rank 2, then $\mathfrak{m}' = k\mathfrak{m}$, where either:*

- (i) $k \in \mathbb{Z} - \{0\}$ and $\theta \in \mathbb{R} - \mathbb{Q}$, or
- (ii) k and θ are the irrational numbers of a quadratic number field.

Proof. Any rank 2 submodule of m can be written as $\mathfrak{m}' = \lambda'_1\mathbb{Z} + \lambda'_2\mathbb{Z}$, where

$$(5) \quad \begin{cases} \lambda'_1 = a\lambda_1 + b\lambda_2 \\ \lambda'_2 = c\lambda_1 + d\lambda_2 \end{cases} \quad \text{and} \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{Z}).$$

(i) Let us assume that $b \neq 0$. Let $\Delta = (a + d)^2 - 4(ad - bc)$ and $\Delta' = (a + d)^2 - 4bc$. We shall consider the following cases.

CASE 1. $\Delta > 0$ and $\Delta \neq m^2$, $m \in \mathbb{Z} - \{0\}$. The real number k can be determined from the equations:

$$(6) \quad \begin{cases} \lambda'_1 = k\lambda_1 = a\lambda_1 + b\lambda_2, \\ \lambda'_2 = k\lambda_2 = c\lambda_1 + d\lambda_2. \end{cases}$$

Since $\theta = \lambda_2/\lambda_1$, one gets the equation $\theta = (c\theta + d)/(a\theta + b)$ by taking a ratio of the two equations above. A quadratic equation for θ writes as $b\theta^2 + (a - d)\theta - c = 0$. The discriminant of the equation coincides with Δ and therefore there exist the real roots $\theta_{1,2} = (a - d \pm \sqrt{\Delta})/(2c)$. Moreover, $k = a + b\theta = a + (b/(2c))(a - d \pm \sqrt{\Delta})$. Since Δ is not the square of an integer, k and θ are the irrationalities of the quadratic number field $\mathbb{Q}(\sqrt{\Delta})$.

CASE 2. $\Delta > 0$ and $\Delta = m^2$, $m \in \mathbb{Z} - \{0\}$. Note that $\theta = (a - d \pm |m|)/(2c)$ is a rational number. Since θ does not satisfy the rank assumption of the lemma, the case should be omitted.

CASE 3. $\Delta = 0$. The quadratic equation has a double root $\theta = (a - d)/(2c) \in \mathbb{Q}$. This case leads to a module of the rank 1, which is contrary to an assumption of the lemma.

CASE 4. $\Delta < 0$ and $\Delta' \neq m^2$, $m \in \mathbb{Z} - \{0\}$. Let us define a new basis $\{\lambda''_1, \lambda''_2\}$ in \mathfrak{m}' so that

$$(7) \quad \begin{cases} \lambda''_1 = \lambda'_1, \\ \lambda''_2 = -\lambda'_2. \end{cases}$$

Then:

$$(8) \quad \begin{cases} \lambda''_1 = a\lambda_1 + b\lambda_2, \\ \lambda''_2 = -c\lambda_1 - d\lambda_2, \end{cases}$$

and $\theta = \lambda''_2/\lambda''_1 = (-c - d\theta)/(a + b\theta)$. The quadratic equation for θ has the form $b\theta^2 + (a + d)\theta + c = 0$, whose discriminant is $\Delta' = (a + d)^2 - 4bc$. Let us show that $\Delta' > 0$. Indeed, $\Delta = (a + d)^2 - 4(ad - bc) < 0$ and the evident inequality $-(a - d)^2 \leq 0$ have the same sign, and we shall add them up. After an obvious elimination, one gets $bc < 0$. Therefore Δ' is a sum of the two positive integers, which is always a positive integer. Thus, there exist the real roots $\theta_{1,2} = (-a - d \pm \sqrt{\Delta'})/(2b)$. Moreover, $k = a + b\theta = (1/2)(a - d \pm \sqrt{\Delta'})$. Since Δ' is not the square of an integer, k and θ are the irrational numbers in the quadratic field $\mathbb{Q}(\sqrt{\Delta'})$.

CASE 5. $\Delta < 0$ and $\Delta' = m^2$, $m \in \mathbb{Z} - \{0\}$. Note that $\theta = (-a - d \pm |m|)/(2b)$ is a rational number. Since θ does not satisfy the rank assumption of the lemma, the case should be omitted.

(ii) Assume that $b = 0$.

CASE 1. $a - d \neq 0$. The quadratic equation for θ degenerates to a linear equation $(a - d)\theta + c = 0$. The root $\theta = c/(d - a) \in \mathbb{Q}$ does not satisfy the rank assumption again, and we omit the case.

CASE 2. $a = d$ and $c \neq 0$. It is easy to see, that the set of the solutions for θ is an empty set.

CASE 3. $a = d$ and $c = 0$. Finally, in this case all coefficients of the quadratic equation vanish, so that any $\theta \in \mathbb{R} - \mathbb{Q}$ is a solution. Note that in the view of (6), $k = a = d \in \mathbb{Z}$. Thus, one gets case (i) of the lemma. Since there are no other possibilities left, Lemma 2 is proved. \square

Lemma 3. *Let E be an elliptic curve with a complex multiplication and h the Hubbard-Masur map, which acts by the formulas of §1.3.D. Consider a module $h(\text{Isom}(E)) = \mu_E(\mathbb{Z} + \theta_E\mathbb{Z}) := \mathfrak{m}_E$. Then:*

- (i) θ_E is a quadratic irrationality,
- (ii) $a\mu_E \in \mathbb{Q}$ (up to a choice of h).

Proof. (i) Since E has a complex multiplication, $\text{End}(E) > \mathbb{Z}$. In particular, there exists a nontrivial endomorphism φ , i.e. an endomorphism which is not the multiplication by $k \in \mathbb{Z}$. By Lemma 1, φ defines a submodule \mathfrak{m}'_E of the rank 2 of the

module m_E . By Lemma 2, $m'_E = km_E$ for a $k \in \mathbb{R}$. Since φ is a nontrivial endomorphism, $k \notin \mathbb{Z}$. Thus, the option (i) of Lemma 2 is excluded. Therefore, by the item (ii) of Lemma 2, θ_E is a quadratic irrationality.

(ii) Recall that $E_{\mathcal{F}} \subset \mathcal{Q} - \{0\}$ is the space of holomorphic differentials on the complex torus, whose horizontal trajectory structure is equivalent to given measured foliation $\mathcal{F} = \mathcal{F}_{\theta}^{\mu}$. We shall vary $\mathcal{F}_{\theta}^{\mu}$, thus varying the Hubbard-Masur homeomorphism $h = h(\mathcal{F}_{\theta}^{\mu}): E_{\mathcal{F}} \rightarrow T(1)$. Namely, consider a 1-parameter continuous family of such maps $h = h_{\mu}$, where $\theta = Const$ and $\mu \in \mathbb{R}$. Recall that $\mu_E = \lambda_1 = \int_{\gamma_1} \phi$, where $\phi = Re \omega$ and $\omega \in E_{\mathcal{F}}$. The family h_{μ} generates a family $\omega_{\mu} = h_{\mu}^{-1}(C)$, where C is a fixed point in $T(1)$. Denote by ϕ_{μ} and λ_1^{μ} the corresponding families of the closed 1-forms and their periods, respectively. By the continuity, λ_1^{μ} takes on a rational value for a $\mu = \mu'$. (Actually, every neighborhood of μ_0 contains such a μ' .) Thus, $\mu_E \in \mathbb{Q}$ for the Hubbard-Masur homeomorphism $h = h_{\mu'}$. □

Lemma 3 implies (i) of Theorem 1. To prove (ii), notice that when $E_1, E_2 \in Isom(E_{CM})$, the respective modules $m_1 = m_2$. It follows from the fact that an isomorphism between the elliptic curves corresponds to a change of basis in the module m (Lemma 1). Theorem 1 is proved. □

3. Arithmetic complexity of the noncommutative tori

Let \mathbb{A}_{θ} be the noncommutative torus with a real multiplication. Since θ is a quadratic irrationality, the regular continued fraction of θ is eventually periodic:

$$(9) \quad \theta = [a_0, a_1, \dots, \overline{a_{k+1}, \dots, a_{k+p}}],$$

where $\overline{a_{k+1}, \dots, a_{k+p}}$ is the minimal period of the continued fraction.

DEFINITION 1. Let us call the number $c(\mathbb{A}_{\theta}) = p$ an arithmetic complexity of the noncommutative torus with real multiplication.

Lemma 4. *The number $c(\mathbb{A}_{\theta})$ is an invariant of the stable isomorphism class of the noncommutative torus \mathbb{A}_{θ} .*

Proof. It follows from Lemma 1 that $\mathbb{A}_{\theta}, \mathbb{A}_{\theta'}$ are stably isomorphic if and only if $\theta' = \theta \bmod GL(2, \mathbb{Z})$. By the main property of the regular continued fractions, the expansion of θ and θ' must coincide, except possibly a finite number of the entries. Since the continued fraction of θ is eventually periodic, so must be the continued fraction of θ' . Moreover, the minimal periods of θ, θ' must coincide as well as their lengths. Thus $c(\mathbb{A}_{\theta'}) = c(\mathbb{A}_{\theta})$. □

EXAMPLE 1. Let us find an arithmetic complexity of the noncommutative torus $\mathbb{A}_{3\sqrt{6}}$. The continued fraction expansion of $3\sqrt{6} = \sqrt{54}$ is $[7; \overline{2, 1, 6, 1, 2, 14}]$. Since the continued fraction is six-periodic, we have $c(\mathbb{A}_{3\sqrt{6}}) = 6$.

It is very useful to think of the normalized period $(1, a_{k+2}/a_{k+1}, \dots, a_{k+p}/a_{k+1})$ of \mathbb{A}_θ as coordinates of the ‘rational points’ of the noncommutative torus, taken up to a cyclic permutation. In the sense, such points are the generators of an abelian group of all rational points of \mathbb{A}_θ modulo the points of a finite order.

Conjecture 1. $c(\mathbb{A}_{\theta_{ECM}}) = rk(E_{CM}) + 1$.

ACKNOWLEDGMENTS. I am grateful to Lawrence D. Taylor for the helpful remarks regarding the manuscript. The referee’s suggestions are kindly acknowledged and incorporated in the text.

References

- [1] A. Douady and J. Hubbard: *On the density of Strebel differentials*, Invent. Math. **30** (1975), 175–179.
- [2] E.G. Effros: *Dimensions and C*-algebras*, CBMS Regional Conference Series in Mathematics **46**, Conf. Board Math. Sci., Washington, D.C., 1981.
- [3] E.G. Effros and C.L. Shen: *Approximately finite C*-algebras and continued fractions*, Indiana Univ. Math. J. **29** (1980), 191–204.
- [4] J. Hubbard and H. Masur: *Quadratic differentials and foliations*, Acta Math. **142** (1979), 221–274.
- [5] M. Kontsevich: *Hodge structures in non-commutative geometry*, (Notes by Ernesto Lupercio) XI Solomon Lefschetz Memorial Lecture Series: in Contemp. Mathematics, arXiv:0801.4760v1.
- [6] Yu.I. Manin: *Real multiplication and noncommutative geometry*; in The Legacy of Niels Henrik Abel Springer, Berlin, 2004, 685–727, arXiv:math/0202109.
- [7] M. Pimsner and D. Voiculescu: *Imbedding the irrational rotation C*-algebra into an AF-algebra*, J. Operator Theory **4** (1980), 201–210.
- [8] A. Polishchuk: *Classification of holomorphic vector bundles on noncommutative two-tori*, Doc. Math. **9** (2004), 163–181, arXiv:math/0308136.
- [9] A. Polishchuk: *Noncommutative two-tori with real multiplication as noncommutative projective varieties*, J. Geom. Phys. **50** (2004), 162–187, arXiv:math/0212306.
- [10] A. Polishchuk: *Analogues of the exponential map associated with complex structures on non-commutative two-tori*, Pacific J. Math. **226** (2006), 153–178, arXiv:math/0404056.
- [11] A. Polishchuk: *Quasicoherent sheaves on complex noncommutative two-tori*, arXiv:math/0506571.
- [12] A. Polishchuk and A. Schwarz: *Categories of holomorphic vector bundles on noncommutative two-tori*, Comm. Math. Phys. **236** (2003), 135–159, arXiv:math/0211262.
- [13] M.A. Rieffel: *C*-algebras associated with irrational rotations*, Pacific J. Math. **93** (1981), 415–429.
- [14] J.H. Silverman: *Advanced Topics in the Arithmetic of Elliptic Curves*, Graduate Texts in Mathematics **151**, Springer, New York, 1994.
- [15] Y. Soibelman: *Quantum tori, mirror symmetry and deformation theory*, Lett. Math. Phys. **56** (2001), 99–125, arXiv:math/0011162.

- [16] Y. Soibelman: *Mirror symmetry and noncommutative geometry of A_∞ -categories*, J. Math. Phys. **45** (2004), 3742–3757.
- [17] Y. Soibelman and V. Vologodsky: *Noncommutative compactifications and elliptic curves*, Int. Math. Res. Not. (2003), 1549–1569. arXiv:math/0205117.
- [18] L.D. Taylor: *A nonstandard approach to real multiplication*, arXiv:math/0612184.
- [19] L.D. Taylor: *Line bundles over quantum tori*, arXiv:math/0612186.
- [20] W.P. Thurston: *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc. (N.S.) **19** (1988), 417–431.

The Fields Institute for Mathematical Sciences
Toronto, ON
Canada
e-mail: igor.v.nikolaev@gmail.com