



Title	Decay of solutions of the wave equation in the exterior of two convex obstacles
Author(s)	Ikawa, Mitsuru
Citation	Osaka Journal of Mathematics. 1982, 19(3), p. 459-509
Version Type	VoR
URL	<a href="https://doi.org/10.18910/7113">https://doi.org/10.18910/7113</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## DECAY OF SOLUTIONS OF THE WAVE EQUATION IN THE EXTERIOR OF TWO CONVEX OBSTACLES

MITSURU IKAWA

(Received June 30, 1980)

**1. Introduction.** Let  $\mathcal{O}_1$  and  $\mathcal{O}_2$  be compact and convex sets in  $\mathbf{R}^3$  with smooth boundary  $\Gamma_1$  and  $\Gamma_2$  respectively. Suppose that

$$\mathcal{O}_1 \cap \mathcal{O}_2 = \phi.$$

Let us set

$$\mathcal{O} = \mathcal{O}_1 \cup \mathcal{O}_2, \quad \Gamma = \Gamma_1 \cup \Gamma_2.$$

Consider a mixed problem

$$(P) \quad \begin{cases} \frac{\partial^2 u}{\partial t^2} - \sum_{j=1}^3 \frac{\partial^2 u}{\partial x_j^2} = 0 & \text{in } \Omega \times (0, \infty) \\ Bu = 0 & \text{on } \Gamma \times (0, \infty) \\ u(x, 0) = u_0(x) \\ \frac{\partial u}{\partial t}(x, 0) = u_1(x) \end{cases}$$

where  $\Omega = \mathbf{R}^3 - \mathcal{O}$ . Here we treat as the boundary operator  $B$  the following two operators:

$$B_1 u = u$$

and

$$B_2 u = \frac{\partial u}{\partial n} + \sigma(x)u$$

where  $n$  denotes the unit outer normal of  $\Gamma$  and  $\sigma(x)$  is a real valued  $C^\infty$ -function defined on  $\Gamma$ .

Concerning the initial data the compatibility condition is always assumed, that is, for  $u_0, u_1 \in C^\infty(\bar{\Omega})$  defining successively  $u_j \in C^\infty(\bar{\Omega})$ ,  $j=2, 3, \dots$  by the formula

$$u_j = \Delta u_{j-2}$$

the condition

$$Bu_j = 0 \quad \text{on } \Gamma$$

is satisfied for all  $j$ .

About the obstacles  $\mathcal{O}_1, \mathcal{O}_2$  we set the condition

(SC) the Gaussian curvature of  $\Gamma$  never vanishes.

Under this condition we will consider the decay of solutions of (P). Let us set

$$\Omega_R = \Omega \cap \{x; |x| \leq R\}.$$

**Theorem 1.** Suppose that  $B=B_1$ , that is, the Dirichlet boundary is posed. Then there exists a constant  $\alpha_1 > 0$  determined by  $\mathcal{O}$  only with the following properties: Let  $R$  and  $\kappa$  be positive numbers and let  $m$  be a positive integer. For any  $u_0, u_1$  such that

$$\bigcup_{j=0}^1 \text{supp } u_j \subset \bar{\Omega}_\kappa$$

we have an estimate

$$(1.1) \quad \sum_{|\gamma| \leq m} \sup_{x \in \Omega_R} |D_{x,t}^\gamma u(x, t)| \leq C_{R,\kappa,m} e^{-\alpha_1 t} \{ \|u_0\|_{m+5, L^2(\Omega)} + \|u_1\|_{m+4, L^2(\Omega)} \},$$

where  $C_{R,\kappa,m}$  is a positive constant depending on  $R, \kappa$  and  $m$  but independent of  $u_0, u_1$ .

For  $P \in \mathbf{R}^3 - \Gamma$ , let us set

$$\Gamma_{jP} = \{x; x \in \Gamma_j, \{P + l\vec{P}x; l > 1\} \subset \Omega\}.$$

DEFINITION. We say that  $\sigma(x)$  satisfies the condition A when there exists a finite number of  $P_j \in \mathcal{O}_1, j=1, 2, \dots, n_1$  and  $Q_j \in \mathcal{O}_2, j=1, 2, \dots, n_2$  such that

$$\bigcup_{j=1}^{n_1} \Gamma_{1P_j} \supset \Gamma_1, \quad \bigcup_{j=1}^{n_2} \Gamma_{2Q_j} \supset \Gamma_2$$

and

$$\begin{aligned} \sigma(x) &< (n_1 + n_2)^{-1} \sum_j \frac{\vec{P}_j x \cdot n(x)}{|P_j - x|^2} && \text{on } \Gamma_1 \\ \sigma(x) &< (n_1 + n_2)^{-1} \sum_j \frac{\vec{Q}_j x \cdot n(x)}{|Q_j - x|^2} && \text{on } \Gamma_2 \end{aligned}$$

where  $\sum_j$  means the summation of  $j$  such that  $x \in \Gamma_{1P_j}$  or  $x \in \Gamma_{2Q_j}$ .

**Theorem 2.** Suppose that  $B=B_2$ , that is, the third kind boundary condition is posed. If  $\sigma(x)$  satisfies the condition A there exists a constant  $\alpha_2 > 0$  determined by  $\mathcal{O}$  and  $\sigma$  with the following properties: Let  $R$  and  $\kappa$  be positive numbers and let  $m$  be a positive integer. For any  $u_0, u_1$  such that

$$\bigcup_{j=0}^1 \text{supp } u_j \subset \bar{\Omega}_\kappa$$

we have an estimate

$$(1.2) \quad \sum_{|\gamma| \leq m} \sup_{x \in \Omega_R} |D_{x,t}^\gamma u(x, t)| \\ \leq C_{R,\kappa,m} e^{-\alpha_2 t} \{ \|u_0\|_{m+5, L^2(\Omega)} + \|u_1\|_{m+4, L^2(\Omega)} \}$$

where  $C_{R,\kappa,m}$  is a positive constant depending on  $R, \kappa, m$  but independent of  $u_0, u_1$ .

It seems to us that until now studies on the uniform decay of the solutions of the wave equation in the exterior domain are made mainly concerning the existence of a function  $p(t)$  such that

$$(*) \quad \begin{cases} E(u, R, t) \leq p(t) E(u, \infty, 0) \\ p(t) \rightarrow 0 \quad \text{as } t \rightarrow \infty \end{cases}$$

for all  $u_0, u_1 \in C_0(\Omega_\kappa)$ , where

$$E(u, R, t) = \frac{1}{2} \int_{\Omega_R} (|\nabla u(x, t)|^2 + |u_t(x, t)|^2) dx.$$

About the necessary condition on the obstacle for the existence of  $p(t)$  we know the work of Ralston [12]. Roughly speaking [12] shows that if the obstacle admits a trapped ray there is no  $p(t)$  verifying (\*).<sup>1)</sup>

If  $\mathcal{O}$  consists of two obstacles  $\mathcal{O}_1, \mathcal{O}_2$  there is always a trapped ray. Indeed, let  $a_j \in \mathcal{O}_j, j=1, 2$  be the points such that

$$|a_1 - a_2| = \inf_{\substack{x \in \mathcal{O}_1 \\ y \in \mathcal{O}_2}} |x - y|$$

Then  $\overrightarrow{a_1 a_2}$  is perpendicular to  $\Gamma_1$  at  $a_1$  and  $\overrightarrow{a_2 a_1}$  perpendicular to  $\Gamma_2$  at  $a_2$ . Therefore the ray starting at  $a_1$  in the direction  $\overrightarrow{a_1 a_2}$  hits  $\Gamma_2$  at  $a_2$ , and is reflected in the direction  $\overrightarrow{a_2 a_1}$ . Then the reflected ray hits  $\Gamma_1$  at  $a_1$ , and is reflected in the direction  $\overrightarrow{a_1 a_2}$ . Namely, the ray plies between  $a_1$  and  $a_2$  and never goes to the infinity. This shows that  $\mathcal{O}$  is trapping. Then we see that the estimate of the type (\*) never holds. Therefore, when  $\mathcal{O}$  consists of two obstacles, in order to estimate the uniform rate of the decay of solutions it is necessary to consider the decay in a weaker form than (\*). Indeed, Walker [13] shows that

$$\lim_{t \rightarrow \infty} p_{\alpha,R}(t) = 0$$

holds for any  $\alpha > 0$ , where

$$p_{\alpha,R}(t) = \sup_{\{u_0, u_1\} \in C_0^\infty(\mathcal{Q}_R)} E(u, R, t)^{1/2} (\|u_0\|_{H_0^{1+\alpha}(\mathcal{Q}_R)} + \|u_1\|_{H_0^\alpha(\mathcal{Q}_R)})^{-1}.$$

1) Morawetz, Ralston and Strauss [11] shows that when there is no trapped ray there exists  $p(t)$  verifying (\*) under some additional conditions.

But we cannot obtain from its proof any more informations about qualitative nature of  $p_{a,R}(t)$  for general obstacles.

We would like to remark about the condition SC. If only the convexity of obstacles is assumed, we can not expect in general the estimate of the form (1.1) or (1.2). Namely in the appendix we will present convex obstacles  $\mathcal{O}_1$  and  $\mathcal{O}_2$  such that

$$\overline{\lim}_{t \rightarrow \infty} p_{m,R}(t) t^{m(1+\varepsilon)} > 0$$

holds for all positive integer  $m$  and positive constant  $\varepsilon$  (Theorem A).

The essential part of the proof of the theorems is the construction of an asymptotic solution for an oscillatory boundary data. There, we estimate carefully the decay of the amplitude function while repeating the reflections.

## 2. Reduction of the problem

Let the supports of  $u_0, u_1$  be contained in  $\bar{\Omega}_\kappa$ . Take  $v_0, v_1 \in C_0^\infty(\mathbf{R}^3)$  so that

$$(2.1) \quad v_j(x) = u_j(x) \quad \text{in } \Omega, \quad j = 0, 1$$

and

$$(2.2) \quad \|v_j\|_{m, L^2(\mathbf{R}^3)} \leq C_m \|u_j\|_{m, L^2(\Omega)}$$

holds. Remark that the constant  $C_m$  depends on  $\Omega$  and  $m$ , but is independent of  $u_j$ . Let  $w(x, t)$  be the solution of the Cauchy problem

$$(2.3) \quad \begin{cases} \square w = 0 & \text{in } \mathbf{R}^3 \times (0, \infty) \\ w(x, 0) = v_0(x) \\ \frac{\partial w}{\partial t}(x, 0) = v_1(x). \end{cases}$$

Since

$$\text{supp } v \subset \{(x); |x| \leq \kappa\}$$

the Huygens' principle assures that

$$(2.4) \quad \text{supp } w \subset \{(x, t); t - \kappa \leq |x| \leq t + \kappa\}.$$

Let us set

$$-Bw(x, t) \mid_{\Gamma \times (0, \infty)} = h(x, t).$$

If we extend  $h$  to  $t < 0$  by setting  $h(x, t) = 0$  for  $t < 0$ , the compatibility condition of  $u_0, u_1$  means that

$$h \in C^\infty(\Gamma \times \mathbf{R}).$$

Combining  $\mathcal{O} \in \{x; |x| \leq d_1\}$  for some  $d_1$  and (2.4) we have

$$(2.5) \quad \text{supp } h \subset \Gamma \times (0, \kappa + d_1).$$

And it follows from the energy estimate of  $w$  that

$$(2.6) \quad \|h\|_{m+1/2, L^2(\Gamma \times \mathbf{R})} \leq C_m \{ \|v_0\|_{m+1+\epsilon} + \|v_1\|_{m+\epsilon} \},$$

where

$$\epsilon = \begin{cases} 1 & \text{if } B = B_2 \\ 0 & \text{if } B = B_1. \end{cases}$$

Consider the boundary value problem with parameter  $p = \mu + ik$  for boundary data  $g \in C^\infty(\Gamma)$

$$(2.7) \quad \begin{cases} (\Delta - p^2)u = 0 & \text{in } \Omega \\ Bu = g & \text{on } \Gamma. \end{cases}$$

It is well known that there exists  $\mu_0 > 0$  such that for  $\mu \geq \mu_0$  (2.7) has a solution uniquely in  $H^2(\Omega)$  and an estimate

$$\|u\|_{m+2, L^2(\Omega)} \leq C_m \sum_{j=0}^m |p|^{m-j} \|g\|_{j+3/2-\epsilon, L^2(\Gamma)}$$

holds for  $m=0, 1, 2, \dots$ . Define  $U(p)$  a mapping from  $C^\infty(\Gamma)$  into  $C^\infty(\bar{\Omega})$  by

$$U(p)g = u(x).$$

Then  $U(p)$  is analytic in  $\text{Re } p \geq \mu_0$  as  $\mathcal{L}(C^\infty(\Gamma), C^\infty(\bar{\Omega}))$ -valued function, where  $\mathcal{L}(C^\infty(\Gamma), C^\infty(\bar{\Omega}))$  denotes the set of all continuous linear mappings from  $C^\infty(\Gamma)$  into  $C^\infty(\bar{\Omega})$ .

**Theorem 2.1.** *Suppose that  $\mathcal{O}$  satisfies the condition SC and that  $\sigma$  verifies the condition A when  $B=B_2$ . Then  $U(p)$  can be prolonged analytically into a region containing*

$$\{p: \text{Re } p \geq -\alpha\}$$

for some  $\alpha > 0$ . And we have for  $\text{Re } p \geq -\alpha$

$$(2.8) \quad \begin{aligned} & \sup_{x \in \bar{\Omega}_R} \sum_{|\gamma| \leq m} |D_x^\gamma(U(p)g)(x)| \\ & \leq C_{m,R} \sum_{j=0}^{m+4} (|k|+1)^{m+3-j} \|g\|_{j, L^2(\Gamma)}, \quad \forall g \in C^\infty(\Gamma). \end{aligned}$$

Note that the solution of the problem

$$\begin{cases} \square z(x, t) = 0 & \text{in } \Omega \times \mathbf{R} \\ Bz(x, t) = h(x, t) & \text{on } \Gamma \times \mathbf{R} \\ \text{supp } z \subset \bar{\Omega} \times (0, \infty) \end{cases}$$

is represented as

$$z(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{(ik+\mu)t} (U(ik+\mu) \hat{h}(\cdot, ik+\mu))(x) dk,$$

where  $\mu \geq \mu_0$  and

$$\hat{h}(x, p) = \int_{-\infty}^{\infty} e^{-pt} h(x, t) dt.$$

By using Theorem 2.1 the path of the integration can be changed to  $\operatorname{Re} p = -\alpha$ , namely we have

$$z(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{(ik-\alpha)t} (U(ik-\alpha) \hat{h}(\cdot, ik-\alpha))(x) dx.$$

With the aid of the estimate (2.8) we obtain immediately from this formula

$$\begin{aligned} (2.9) \quad & \sum_{|\gamma| \leq m} \sup_{x \in \Omega_R} |D_{x,t}^\gamma z(x, t)| \\ & \leq C_{m,R} e^{-\alpha t} \int_{-\infty}^{\infty} \sum_{j=0}^{m+3} (|k|+1)^{m+3-j} \|\hat{h}(\cdot, ik-\alpha)\|_{j, L^2(\Gamma)} dk \\ & \leq C_{m,R} e^{-\alpha t} \|\hat{h}\|_{m+4, L^2(\Gamma \times \mathbb{R})}. \end{aligned}$$

Since the solution  $u(x, t)$  of (P) is represented as

$$u(x, t) = w(x, t) + z(x, t)$$

we have Theorems 1 and 2 from (2.4), (2.6) and (2.9).

Concerning the analytic continuation the following results hold without the condition SC:

- (i) For the Dirichlet boundary condition  $U(p)$  can be prolonged analytically into a region containing  $\{p; \operatorname{Re} p \geq 0\}$ .
- (ii) For the third kind boundary condition  $U(p)$  can be prolonged into a region containing  $\{p; \operatorname{Re} p \geq 0, p \notin [0, \mu_0]\}^2$ . In the rest of this paper we will show the following

**Theorem 2.2.** Suppose that  $\mathcal{O}$  satisfies the condition SC. Let  $a_j \in \Gamma_j, j=1, 2$  be the points such that  $|a_1 - a_2| = \operatorname{dis}(\mathcal{O}_1, \mathcal{O}_2)$  and let us denote by  $K_{ij}, j=1, 2$  the principal curvatures of  $\Gamma_i$  at  $a_i$ . Set

$$c_0 = \frac{1}{d_0} \log(1 + 2d_0 K)$$

where  $d_0 = \operatorname{dis}(\mathcal{O}_1, \mathcal{O}_2)$  and  $K = \inf_{i,j=1,2} K_{ij}$ . Then for any  $\varepsilon > 0$ , there exists  $k_\varepsilon > 0$

2) See, for example, about (i) Chapter V of Lax-Phillips [6], Chapter 8 of Mizohata [9] and about (ii) Asakura [1].

such that  $U(p)$  can be prolonged into

$$\{\mu + ik; \mu \geq -(c_0 - \varepsilon), |k| \geq k_\varepsilon\}.$$

Moreover an estimate

$$\begin{aligned} & \sum_{|\gamma| \leq m} \sup_{x \in \Omega_R} |D_x^\gamma(U(p)g)(x)| \\ & \leq C_{\varepsilon, R, m} \sum_{j=0}^{m=3} (|k| + 1)^{m+3-j} \|g\|_{j, L^2(\Gamma)}, \quad \forall g \in C^\infty(\Gamma) \end{aligned}$$

holds.

Then in order to show Theorem 2.1, if we admit Theorem 2.2, it remains to prove that  $p \in [0, \infty)$  is not the generalized eigenvalue of (2.7) for  $B = B_2$ , that is, there is no non-trivial solution of (2.7) for  $g=0$  verifying

$$(2.10) \quad |x| |u(x)| + |x|^2 |\nabla u(x)| \leq C \quad \text{as } |x| \rightarrow \infty.$$

Therefore we show

**Lemma 2.3.** *Suppose that  $\sigma$  satisfies the condition A. Then for  $p \in [0, \infty)$  there is no non-trivial solution of (2.7) for  $g=0$  verifying (2.10).*

*Proof.* Suppose that  $u(x)$  is a non-trivial solution of (2.7) for  $g=0$  verifying (2.10). Then by the integration by parts of  $(\Delta - p^2)u(x) \cdot \overline{u(x)}$  it follows that

$$\int_{\Gamma} \sigma(x) |u(x)|^2 ds = \int_{\Omega} |\nabla u(x)|^2 dx + p^2 \int_{\Omega} |u|^2 dx.$$

Therefore if we derive from the assumption on  $\sigma$  the estimate

$$(2.11) \quad \int_{\Gamma} \sigma |u|^2 dS \leq (1 - \varepsilon) \int_{\Omega} |\nabla u|^2 dx$$

for some  $\varepsilon > 0$ ,  $u$  must be identically zero. Suppose that  $P \in \mathcal{O}_1$ . For  $x \in \Gamma_{1P}$ ,

setting  $\omega = \frac{\vec{Px}}{|\vec{Px}|}$ ,  $r(\omega) = |\vec{xP}|$

$$u(x) = - \int_{r(\omega)}^{\infty} \frac{\partial}{\partial s} u(s\omega) ds = - \int_{r(\omega)}^{\infty} (\omega \cdot \nabla u)(s\omega) ds.$$

Then

$$\begin{aligned} |u(x)|^2 & \leq \int_{r(\omega)}^{\infty} \frac{ds}{s^2} \int_{r(\omega)}^{\infty} |\nabla u(s\omega)|^2 s^2 ds \\ & = r(\omega)^{-1} \int_{r(\omega)}^{\infty} |\nabla u(s\omega)|^2 s^2 ds. \end{aligned}$$

Setting  $\omega_{1P} = \left\{ \frac{\vec{Px}}{|\vec{Px}|}, x \in \Gamma_{1P} \right\}$

$$\begin{aligned} \int_{\omega_{1P}} |u(x)|^2 r(\omega) d\omega &\leq \int_{\omega_{1P}} \int_{r(\omega)}^{\infty} |\nabla u(s\omega)|^2 s^2 ds d\omega \\ &\leq \int_{\Omega} |\nabla u(x)|^2 dx. \end{aligned}$$

Denoting by  $dS$  the surface element of  $\Gamma_1$ , since  $dS = r(\omega)^2 \frac{r(\omega)}{\overrightarrow{Px} \cdot \mathbf{n}(x)} d\omega$  the condition

$$\sup_{x \in \Gamma_{1P}} \sigma(x) \frac{|\overrightarrow{Px}|^2}{\overrightarrow{Px} \cdot \mathbf{n}(x)} \leq C$$

implies

$$\begin{aligned} \int_{\Gamma_{1P}} \sigma(x) |u(x)|^2 dS &\leq C \int_{\omega_{1P}} r(\omega) |u(x)|^2 d\omega \\ &\leq C \int_{\Omega} |\nabla u(x)|^2 dx. \end{aligned}$$

Suppose that  $P_j, j=1, 2, \dots, n_1$  satisfy

$$\bigcup_{j=1}^{n_1} \Gamma_{1P_j} \supset \Gamma_1 \quad \text{and} \quad \sum_j \frac{\sigma(x) |\overrightarrow{P_j x}|^2}{\overrightarrow{P_j x} \cdot \mathbf{n}(x)} \leq (1-\varepsilon)(n_1+n_1)^{-1}.$$

Then it holds that

$$\begin{aligned} \int_{\Gamma_1} \sigma(x)^+ |u(x)|^2 dS &\leq \sum_{j=1}^{n_1} \int_{\Gamma_{1P_j}} \sigma(x)^+ |u(x)|^2 dS \\ &\leq \frac{n_1(1-\varepsilon)}{(n_1+n_2)} \int_{\Omega} |\nabla u(x)|^2 dx, \end{aligned}$$

where  $\sigma(x)^+ = \max(\sigma(x), 0)$ . By the same way we have

$$\int_{\Gamma_2} \sigma^+(x) |u(x)|^2 dS \leq \frac{n_2(1-\varepsilon)}{(n_1+n_2)} \int_{\Omega} |\nabla u(x)|^2 dx.$$

Then it follows that

$$\begin{aligned} \int_{\Gamma} \sigma(x) |u(x)|^2 dS &\leq \int_{\Gamma} \sigma(x)^+ |u(x)|^2 dS \\ &\leq (1-\varepsilon) \int_{\Omega} |\nabla u(x)|^2 dx. \end{aligned}$$

Thus (2.11) is derived. Therefore Lemma is proved.

### 3. Properties of broken rays

For  $x \in \Gamma$ ,  $\mathbf{n}(x)$  denotes the unit outer normal of  $\Gamma$  at  $x$ , and

$$\Sigma_x^+ = \{\xi; |\xi|=1, n(x) \cdot \xi \geq 0\}.$$

We denote by  $\mathcal{X}(x, \xi)$  the broken ray according to the law of the geometrical optics starting at  $x \in \Gamma$  in the direction  $\xi \in \Sigma_x^+$ , by  $X_1(x, \xi)$ ,  $X_0(x, \xi)$ , ... the points of reflection of the broken ray and by  $\Xi_j(x, \xi)$  the direction of the ray reflected at  $X_j(x, \xi)$ . More precisely, if

$$\{x + l\xi; l > 0\} \cap \Gamma = \phi$$

we set  $L_0(x, \xi) = \{x + l\xi; l \geq 0\}$ . If  $\{x + l\xi; l > 0\} \cap \Gamma \neq \phi$  we set

$$\begin{aligned} l_0(x, \xi) &= \inf \{l; l > 0, x + l\xi \in \Gamma\} \\ L_0(x, \xi) &= \{x + l\xi; 0 \leq l \leq l_0(x, \xi)\} \\ X_1(x, \xi) &= x + l_0(x, \xi)\xi \\ \Xi_1(x, \xi) &= \xi - 2(n(X_1(x, \xi)) \cdot \xi)n(X_1(x, \xi)). \end{aligned}$$

When  $\{X_1 + l\Xi_1; l > 0\} \cap \Gamma = \phi$ ,  $L_1(x, \xi) = \{X_1 + l\Xi_1; l \geq 0\}$ . Otherwise we set

$$\begin{aligned} l_1(x, \xi) &= \inf \{l; l > 0, X_1 + l\Xi_1 \in \Gamma\} \\ L_1(x, \xi) &= \{X_1 + l\Xi_1; 0 \leq l \leq l_1\} \\ X_2(x, \xi) &= X_1 + l_1\Xi_1 \\ \Xi_2(x, \xi) &= \Xi_1 - 2(n(X_2) \cdot \Xi_1)n(X_2). \end{aligned}$$

Thus we define successively  $l_j(x, \xi)$ ,  $X_j(x, \xi)$ ,  $\Xi_j(x, \xi)$ ,  $L_j(x, \xi)$  until  $\{X_j + l\Xi_j; l > 0\} \cap \Gamma = \phi$ . If there exists  $j_0$  such that for  $j \leq j_0$ ,  $l_j(x, \xi)$ ,  $X_j(x, \xi)$ ,  $\Xi_j(x, \xi)$  are defined and  $\{X_{j_0} + l\Xi_{j_0}; l > 0\} \cap \Gamma = \phi$ , then we define

$$\begin{aligned} \mathcal{X}(x, \xi) &= \bigcup_{j=0}^{j_0} L_j(x, \xi) \\ {}^*\mathcal{X}(x, \xi) &= j_0. \end{aligned}$$

Otherwise

$$\begin{aligned} \mathcal{X}(x, \xi) &= \bigcup_{j=0}^{\infty} L_j(x, \xi) \\ {}^*\mathcal{X}(x, \xi) &= \infty. \end{aligned}$$

We set for  $(x, t) \in \Gamma \times \mathbf{R}$ ,  $\xi \in \Sigma_x^+$

$$\begin{aligned} \mathcal{L}_j(x, t, \xi) &= \{(X_j(x, \xi) + l\Xi_j(x, \xi), t + \bar{l}_j(x, \xi) + l); 0 \leq l \leq l_j(x, \xi)\} \\ \bar{\mathcal{L}}_j(x, t, \xi) &= \{(X_j(x, \xi) + l\Xi_j(x, \xi), t + \bar{l}_j(x, \xi) + l); 0 \leq l\} \\ \mathcal{L}(x, t, \xi) &= \bigcup_j \mathcal{L}_j(x, t, \xi), \end{aligned}$$

where  $\bar{l}_j(x, \xi) = \sum_{p=0}^{j-1} l_p(x, \xi)$ .

The following two lemmas are trivial.

**Lemma 3.1.** For  $\mathcal{O}_1, \mathcal{O}_2$  there exists  $\delta_1 > 0$  with the following property: If  $x \in \Gamma_1, \xi \in \Sigma_x^+$  and

$$0 \leq -n(X_1(x, \xi)) \cdot \xi \leq \delta_1,$$

the reflected ray does not pass the  $d_0/2$  neighborhood of  $\mathcal{O}_1$ , i.e.,  $L_1(x, \xi)$  is a half line and

$$L_1(x, \xi) \cap \{y; \text{dis}(y, \mathcal{O}_1) \leq d_0/2\} = \phi.$$

**Lemma 3.2.** For each  $x \in \Gamma, \xi \in \Sigma_x^+$  verifying  ${}^*\mathcal{X}(x, \xi) \geq J$  and

$$n(X_j(x, \xi)) \cdot \Xi_j(x, \xi) > 0, \quad j = 1, 2, \dots, J,$$

there exists  $\varepsilon > 0$  such that if  $y \in \Gamma, \eta \in \Sigma_y^+$  and

$$|x - y| + |\xi - \eta| < \varepsilon,$$

it holds that

$${}^*\mathcal{X}(y, \eta) \geq J$$

and

$$X_j(y, \eta) \rightarrow X_j(x, \xi)$$

$$\Xi_j(y, \eta) \rightarrow \Xi_j(x, \xi)$$

for  $j=1, 2, \dots, J$  when  $(y, \eta) \rightarrow (x, \xi)$ .

**Corollary.** Let  $L$  be the line passing  $a_1$  and  $a_2$ , where  $a_j, j=1, 2$  are those in Theorem 2.2. Denote by  $S_j(\delta)$  the connected component containing  $a_j$  of

$$\Gamma_j \cap \{x; \text{dis}(x, L) \leq \delta\}, \quad \delta > 0.$$

For any  $\varepsilon_0 > 0$ , if we choose  $\delta_2 > 0$  sufficiently small, every ray starting at  $x \in S_1(\delta_2)$  in the direction  $\xi \in \Sigma_x^+$  and fitting  $S_2(\delta_2)$  (starting at  $x \in S_2(\delta_2)$  and hitting  $S_1(\delta_2)$ ) satisfies

$$-\xi \cdot n(X_1(x, \xi)) > 1 - \varepsilon_0.$$

Proof. Note that for  $\xi_0 = \frac{a_2 - a_1}{|a_2 - a_1|} \in \Sigma_{a_1}^+$

$$-\xi_0 \cdot n(X_1(a_1, \xi_0)) = 1.$$

Then from Lemma 3.2 there exists  $\varepsilon > 0$  such that

$$|y - a_1| + |\eta - \xi_0| < \varepsilon, \quad y \in \Gamma_1, \eta \in \Sigma_y^+$$

implies that

$$|\eta \cdot n(X_1(y, \eta)) - \xi_0 \cdot n(X_1(a_1, \xi_0))| < \varepsilon_0.$$

For any  $y \in S_1(\delta_2)$ ,  $\eta \in \sum_y^+$  such that  $X_1(y, \eta) \in S_2(\delta_2)$  we have  $|y - a_1| \leq 2\delta_2$  and  $|X_1(y, \eta) - a_2| \leq 2\delta_2$ . Then

$$\eta - \xi_0 = \frac{X_1(y, \eta) - y}{|X_1(y, \eta) - y|} - \frac{a_2 - a_1}{|a_2 - a_1|}$$

must be small if  $\delta_2$  is chosen so small. Then for any  $\varepsilon > 0$  there exists  $\delta_2 > 0$  such that  $y \in S_1$ ,  $\eta \in \sum_y^+$  and  $X_1(y, \eta) \in S_2$  imply

$$|y - a_1| + |\eta - \xi_0| < \varepsilon.$$

Using the above remark we have

$$-\eta \cdot n(X_1(y, \eta)) \geq -\xi_0 \cdot n(X_1(a_1, \xi_0)) - \varepsilon_0 = 1 - \varepsilon_0.$$

Q.E.D.

**Lemma 3.3.** For each  $\delta_2 > 0$  of Corollary of the previous lemma there exists a positive integer  $K$  such that  $x \in \Gamma - S(\delta_2)$ ,  $\xi \in \sum_x^+$  and  $\mathcal{X}(x, \xi) \cap S(\delta_2) = \emptyset$  imply

$$(3.1) \quad {}^*\mathcal{X}(x, \xi) \leq K,$$

where  $S(\delta_2) = S_1(\delta_2) \cup S_2(\delta_2)$ .

Proof. Let  $\delta_1 > 0$  be that of Lemma 3.1. Suppose that  $L = \{(0, 0, l); l \in \mathbf{R}\}$ . For  $x = (x_1, x_2, x_3)$  denote by  $x'$  the point  $(x_1, x_2, 0)$ . The strict convexity assures that

$$(3.2) \quad n(x) \cdot x' \geq c > 0, \quad \forall x \in \Gamma_l - S_l$$

if  $n(x) \cdot ((-1)^{l-1}(a_2 - a_1)) \geq 0$ , where  $c > 0$  depends on  $\delta_2$ . Suppose that

$${}^*\mathcal{X}(x, \xi) \geq J.$$

Then by Lemma 3.1 we have

$$(3.3) \quad -n(X_j) \cdot \Xi_{j-1} \geq \delta_1 \quad \text{for } j = 1, 2, \dots, J-1.$$

Note that, if  $x \in \Gamma_l - S_l$ ,

$$(3.4) \quad n(X_j) \cdot ((-1)^{j+l-1}(a_2 - a_1)) \geq 0, \quad l = 1, 2, j = 1, 2, \dots.$$

Let  $x(s) = (x_1(s), x_2(s), x_3(s))$  be the representation of  $\mathcal{X}(x, \xi)$  by the length of the broken ray from  $x$  to  $x(s)$ . Then for  $s \in \{\bar{l}_j; j = 0, 1, 2, \dots\}$

$$\frac{d}{ds} |x(s)'|^2 = 2 \left( x_1(s) \cdot \frac{d}{ds} x_1(s) + x_2(s) \cdot \frac{d}{ds} x_2(s) \right)$$

and if  $x(s) \in L_j$

$$= 2(X_j + (s - \bar{l}_j)\Xi_j)' \cdot \Xi_j \geq 2X_j' \cdot \Xi_j.$$

And we have

$$\begin{aligned} X'_j \cdot \Xi_j &= X'_j \cdot (\Xi_{j-1} - 2(n(X_j) \cdot \Xi_{j-1})n(X_j)) \\ &= X'_j \cdot \Xi_{j-1} + 2(-n(X_j) \cdot \Xi_{j-1})(X'_j \cdot n(X_j)). \end{aligned}$$

From (3.2) and (3.3) it follows that

$$(3.5) \quad X'_j \cdot \Xi_j \geq X'_j \cdot \Xi_{j-1} + 2c\delta_1.$$

Since  $X'_1 \cdot \xi \geq -|X_1| \geq -\max\{|x|; x \in \Gamma\} (= -A)$ , we have

$$X'_j \cdot \Xi_j \geq -A + 2jc\delta_1.$$

Then for  $j \geq j_0 = \left\lceil \frac{A}{2c\delta_1} \right\rceil + 2$ , we have  $X'_j \cdot \Xi_j \geq 2c\delta_1$ , which implies

$$\frac{d}{ds} |x(s)'|^2 \geq 4c\delta_1, \quad s \geq \bar{l}_{j_0}.$$

Therefore for  $s \geq \bar{l}_{j_0}$ , integrating the above inequality from  $\bar{l}_{j_0}$  to  $S$ , we have

$$|x(s)'|^2 - |x(\bar{l}_{j_0})'|^2 \geq 4(s - \bar{l}_{j_0})c\delta_1.$$

Then if  $x(s) \in \Gamma$  we have  $4(s - \bar{l}_{j_0})c\delta_1 \leq A^2$ , then

$$s \leq \bar{l}_{j_0} + \frac{A^2}{4c\delta_1} = s_0,$$

from which it follows that for any  $X_j = x(\bar{l}_{j-1})$

$$\bar{l}_{j-1} = \sum_{h=0}^{j-1} l_h \leq s_0.$$

Since  $l_j \geq |a_2 - a_1|$  for all  $j$ ,  ${}^*\mathcal{X}(x, \xi) \leq s_0/|a_2 - a_1| = K$ . Thus the lemma is proved. Q.E.D.

**Corollary.** *If we choose  $\delta_3$  as  $\delta_3 > \delta_2$  and sufficiently close to  $\delta_2$ , it holds that for any  $x \in S(\delta_3)$ ,  $\xi \in \sum_x^+$  such that  $X_1(x, \xi) \in S(\delta_3) - S(\delta_2)$*

$${}^*\mathcal{X}(x, \xi) \leq K + 1,$$

where  $K$  is the one in Lemma 3.3.

**Proof.** By the process of the proof of Lemma 3.3 we see that for  $x, \xi$  such that  $x' \cdot \xi \geq 0$  it must hold the estimate

$${}^*\mathcal{X}(x, \xi) \leq K.$$

Therefore, if we can show that

$$(3.6) \quad X'_1 \cdot \Xi_1 \geq 0,$$

we have

$$^*\mathcal{X}(x, \xi) \leq K+1.$$

Then we will show (3.6). Suppose that  $x \in S_1(\delta_3)$ ,  $X_1(x, \xi) \in S_2(\delta_3) - S_2(\delta_2)$  and  $x' \cdot \xi < 0$ . This means that

$$(3.7) \quad x_2^1 + x_2^2 \leq \delta_3^2$$

$$(3.8) \quad \delta_2^2 \leq (x_1 + l_0 \xi_1)^2 + (x_2 + l_0 \xi_2)^2 \leq \delta_3^2$$

$$(3.9) \quad x_1 \xi_1 + x_2 \xi_2 < 0.$$

Since

$$(x_1 + l_0 \xi_1)^2 + (x_2 + l_0 \xi_2)^2 = x_1^2 + x_2^2 + l_0 x' \cdot \xi + l_0 X_1' \cdot \xi,$$

(3.7) and (3.8) imply

$$\delta_2^2 \leq \delta_3^2 + l_0 x' \cdot \xi + l_0 X_1' \cdot \xi.$$

By taking account of (3.9) we have

$$(\delta_2^2 - \delta_3^2)/l_0 \leq X_1' \cdot \xi.$$

Since  $X_1 \notin S_2(\delta_2)$  we can use (3.5) and obtain

$$X_1' \cdot \Xi_1 \geq (\delta_2^2 - \delta_3^2)/l_0 + 2c\delta_1.$$

Then, if we choose  $\delta_3$  sufficiently close to  $\delta_2$ , we have (3.6). Thus our assertion is proved. Q.E.D.

#### 4. The curvatures of the wave front of reflected rays

For  $\varphi(x)$  such that  $|\nabla \varphi| = 1$ , we will call the surface  $\{y; \varphi(y) = \varphi(x)\}$  the wave front of  $\varphi$  passing  $x$  and denote it by  $\mathcal{G}_\varphi(x)$ . Let  $\mathcal{U}$  be a neighborhood in  $\mathbf{R}^3$  of  $s_0 \in \Gamma$  and  $\varphi^+$ ,  $\varphi^-$  be functions defined in  $\mathcal{U}$  satisfying  $|\nabla \varphi^\pm| = 1$ . Suppose that

$$(4.1) \quad \varphi^+(x) = \varphi^-(x) \quad \forall x \in \Gamma \cap \mathcal{U}$$

$$(4.2) \quad \frac{\partial \varphi^+}{\partial n} > 0, \quad \frac{\partial \varphi^-}{\partial n} < 0 \quad \text{on } \Gamma \cap \mathcal{U}.$$

We will consider the relation between the principal curvatures of the wave fronts  $\mathcal{G}_{\varphi^+}(s_0)$  and  $\mathcal{G}_{\varphi^-}(s_0)$ . Hereafter the principal curvatures of  $\mathcal{G}_\varphi(x)$  signify those with respect to  $-\nabla \varphi(x)$ . Suppose that

$$n(s_0) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

and  $\Gamma$  is represented in  $\mathcal{U}$  as  $\{y(\eta); \eta \in U\}$ ,  $U \subset \mathbf{R}^2$ ,  $y(0)=s_0$

$$(4.3) \quad \frac{\partial y}{\partial \eta_1}(0) = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad \frac{\partial y}{\partial \eta_2}(0) = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

Let  $\{\omega(\sigma); \sigma \in \tilde{U}\}$  be a representation of the wave front  $\mathcal{C}^- = \mathcal{C}_{\varphi^-}(s_0)$  such that  $\omega(0)=s_0$  and

$$(4.4) \quad \frac{\partial \omega}{\partial \sigma_i}(0) \cdot \frac{\partial \omega}{\partial \sigma_j}(0) = \delta_{ij}, \quad i, j = 1, 2$$

$$(4.5) \quad \frac{\partial i}{\partial \sigma_j}(0) = \kappa_j \frac{\partial \omega}{\partial \sigma_j}(0), \quad j = 1, 2$$

where  $i(\sigma) = (\nabla \varphi^-)(\omega(\sigma))$  and  $\kappa_j$ ,  $j=1, 2$  are the principal curvatures of  $\mathcal{C}^-$  at  $s_0$ . Then  $\mathcal{C}^+ = \mathcal{C}_{\varphi^+}(s_0)$  is represented as

$$\tau(\sigma) = y(\eta(\sigma)) - l(\sigma)r(\sigma)$$

where

$$r(\sigma) = i(\sigma) - 2\{i(\sigma) \cdot n(y(\eta(\sigma)))\}n(y(\eta(\sigma))),$$

and  $\eta(\sigma) = (\eta_1(\sigma), \eta_2(\sigma)) \in U$ ,  $l(\sigma) \in \mathbf{R}$  are determined by

$$(4.6) \quad y(\eta) = \omega(\sigma) + l(\sigma)i(\sigma) \quad \text{for } \sigma \in \tilde{U}.$$

Evidently  $r(\sigma) = (\nabla \varphi^+)(\tau(\sigma))$ . When we denote

$$(4.7) \quad -i(0) = \begin{bmatrix} \sin \nu \\ 0 \\ \cos \nu \end{bmatrix}$$

we can write

$$(4.8) \quad \frac{\partial \omega}{\partial \sigma_1}(0) = \begin{bmatrix} -\cos \theta_0 \cdot \cos \nu \\ \sin \theta_0 \\ \cos \theta_0 \cdot \sin \nu \end{bmatrix}, \quad \frac{\partial \omega}{\partial \sigma_2}(0) = \begin{bmatrix} \sin \theta_0 \cdot \cos \nu \\ \cos \theta_0 \\ -\sin \theta_0 \cdot \sin \nu \end{bmatrix}$$

for some  $\theta_0$ . Differentiate (4.6) by  $\sigma_j$  and we have

$$\frac{\partial y}{\partial \eta_1} \frac{\partial \eta_1}{\partial \sigma_j} + \frac{\partial y}{\partial \eta_2} \frac{\partial \eta_2}{\partial \sigma_j} = \frac{\partial \omega}{\partial \sigma_j} + \frac{\partial l}{\partial \sigma_j} i + l \frac{\partial i}{\partial \sigma_j}.$$

Substituting (4.3), (4.5), (4.7), (4.8) and  $l(0)=0$  we obtain

$$(4.9) \quad \frac{\partial l}{\partial \sigma_1}(0) = \tan \nu \cdot \cos \theta_0, \quad \frac{\partial l}{\partial \sigma_2}(0) = -\tan \nu \cdot \sin \theta_0$$

$$(4.10) \quad \begin{bmatrix} \frac{\partial \eta_1}{\partial \sigma_1} \frac{\partial \eta_1}{\partial \sigma_2} \\ \frac{\partial \eta_2}{\partial \sigma_1} \frac{\partial \eta_2}{\partial \sigma_2} \end{bmatrix}_{\sigma=0} = T\Theta,$$

where

$$T = \begin{bmatrix} (\cos \nu)^{-1} & 0 \\ 0 & 1 \end{bmatrix}, \quad \Theta = \begin{bmatrix} -\cos \theta_0 & \sin \theta_0 \\ \sin \theta_0 & \cos \theta_0 \end{bmatrix}.$$

Using (4.9) we have

$$(4.11) \quad \begin{bmatrix} \frac{\partial \tau}{\partial \sigma_1} \frac{\partial \tau}{\partial \sigma_2} \end{bmatrix}_{\sigma=0} = [Y_1 \quad Y_2]\Theta$$

where

$$Y_1 = \begin{bmatrix} \cos \nu \\ 0 \\ \sin \nu \end{bmatrix}, \quad Y_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

Denote by  $K_1, K_2$  the principal curvatures with respect to  $-n(s_0)$  of  $\Gamma$  at  $s_0$  and by  $\psi$  the angle from  $\frac{\partial y}{\partial \eta_1}(0)$  to the principal direction corresponding to  $K_1$ . Then it holds that

$$\left[ \frac{\partial n(y(\eta))}{\partial \eta_1} \frac{\partial n(y(\eta))}{\partial \eta_2} \right]_{\eta=0} = \left[ \frac{\partial y}{\partial \eta_1} \frac{\partial y}{\partial \eta_2} \right]_{\eta=0} {}^t\Psi K \Psi$$

where

$$\Psi = \begin{bmatrix} \cos \psi & -\sin \psi \\ \sin \psi & \cos \psi \end{bmatrix}, \quad K = \begin{bmatrix} K_1 & 0 \\ 0 & K_2 \end{bmatrix}.$$

Therefore we have

$$(4.12) \quad \begin{aligned} & \left[ \frac{\partial n(y(\eta(\sigma)))}{\partial \sigma_1} \frac{\partial n(y(\eta(\sigma)))}{\partial \sigma_2} \right]_{\sigma=0} \\ &= \left[ \frac{\partial n}{\partial \eta_1} \frac{\partial n}{\partial \eta_2} \right]_{\eta=0} \left[ \frac{\partial \eta_i}{\partial \sigma_j} \right]_{\substack{i,j=1,2 \\ \sigma=0}} = \left[ \frac{\partial y}{\partial \eta_1} \frac{\partial y}{\partial \eta_2} \right]_{\eta=0} {}^t\Psi K \Psi T \Theta. \end{aligned}$$

Note that

$$\frac{\partial r}{\partial \sigma_j} = \frac{\partial i}{\partial \sigma_j} - 2 \left( \frac{\partial i}{\partial \sigma_j} \cdot n \right) n - 2 \left( i \cdot \frac{\partial n}{\partial \sigma_j} \right) n - 2 (i \cdot n) \frac{\partial n}{\partial \sigma_j}.$$

Since at  $\sigma=0$

$$\frac{\partial i}{\partial \sigma_1} - 2 \left( \frac{\partial i}{\partial \sigma_1} \cdot n \right) n = \kappa_1 \begin{bmatrix} -\cos \theta_0 \cos \nu \\ \sin \theta_0 \\ -\cos \theta_0 \sin \nu \end{bmatrix} = \kappa_1 [Y_1 \quad Y_2] \begin{bmatrix} -\cos \theta_0 \\ \sin \theta_0 \end{bmatrix}$$

$$-2\left(i \cdot \frac{\partial n}{\partial \sigma_1}\right)n - 2(i \cdot n) \frac{\partial n}{\partial \sigma_1} = 2 \begin{bmatrix} \cos \nu \\ 0 \\ \sin \nu \end{bmatrix} \left(\frac{\partial n}{\partial \sigma_1}\right)_1 + 2 \cos \nu \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \left(\frac{\partial n}{\partial \sigma_1}\right)_2,$$

where  $\left(\frac{\partial n}{\partial \sigma_1}\right)_j$  denotes the  $j$ -th component of  $\frac{\partial n}{\partial \sigma_1}$ , we have

$$\frac{\partial r}{\partial \sigma_1}(0) = [Y_1 \quad Y_2](\kappa_1 I + 2 \cos \nu T^t \Psi K \Psi T) \begin{bmatrix} -\cos \theta_0 \\ \sin \theta_0 \end{bmatrix}.$$

By the same way

$$\frac{\partial r}{\partial \sigma_2}(0) = [Y_1 \quad Y_2](\kappa_2 I + 2 \cos \nu T^t \Psi K \Psi T) \begin{bmatrix} \sin \theta_0 \\ \cos \theta_0 \end{bmatrix}.$$

Namely

$$\left[ \frac{\partial r}{\partial \sigma_1} \frac{\partial r}{\partial \sigma_2} \right]_{\sigma=0} = [Y_1 \quad Y_2](\tilde{K} + 2 \cos \nu T^t \Psi K \Psi T) \Theta$$

where

$$\tilde{K} = \begin{bmatrix} \kappa_1 & 0 \\ 0 & \kappa_2 \end{bmatrix}.$$

Taking account of (4.11) we see that the principal curvatures of  $\mathcal{C}^+$  at  $s_0$  are the eigenvalues of the matrix

$$\tilde{K} + 2 \cos \nu T^t \Psi K \Psi T.$$

Thus we have

**Lemma 4.1.** *The principal curvatures  $\kappa_1^+$ ,  $\kappa_2^+$  of  $\mathcal{C}^+$  at  $s_0$  satisfy the inequalities*

$$(4.13) \quad \min_{j=1,2} \kappa_j^- + 2 \min_{j=1,2} K_j \leq \kappa_j^+ \leq \max_{j=1,2} \kappa_j^- + \frac{2}{\cos \nu} \max_{j=1,2} K_j$$

where  $\kappa_j^-$ ,  $K_j$ ,  $j=1, 2$  are the principal curvatures of  $\mathcal{C}^-$  and  $\Gamma$  at  $s_0$  respectively and  $\cos \nu = -n(s_0) \cdot \nabla \varphi^-(s_0)$ .

Next we will show

**Proposition 4.2.** *Let  $\mathcal{U}$  be a neighborhood of  $x_0 \in \Gamma_1$  (or  $\Gamma_2$ ) and  $\varphi \in C^\infty(\mathcal{U})$  and  $|\nabla \varphi| = 1$  in  $\mathcal{U}$ . Suppose that*

(i) *for all  $x \in \Gamma_1 \cap \mathcal{U}$  the principal curvatures of  $\mathcal{C}_\varphi(x)$  is positive.*

(ii)  ${}^* \mathcal{X}(x, \nabla \varphi(x)) \geq p+1, \quad \forall x \in \Gamma_1 \cup \mathcal{U}.$

*Then there exist  $\varphi_j(x)$ ,  $j=0, 1, 2, \dots, p$  such that*

$$(a) \quad \varphi_j \in C^\infty(\omega_j), \quad |\nabla \varphi_j| = 1 \quad \text{in } \omega_j$$

$$\text{where } \omega_j = \bigcup_{y \in \Gamma_1 \cap \mathcal{U}} L_j(y, \nabla \varphi(y)).$$

$$(b) \quad \varphi_0 = \varphi \quad \text{in } \omega_0 \cap \mathcal{U}$$

$$(c) \quad \varphi_j(x) = \varphi_{j-1}(x), \quad \frac{\partial \varphi_j}{\partial n}(x) = -\frac{\partial \varphi_{j-1}}{\partial n}(x), \quad \forall x \in \bigcup_{y \in \Gamma_1 \cap \mathcal{U}} X_j(y, \nabla \varphi(y)).$$

Moreover it holds that

(d) the principal curvatures  $\kappa_{jl}(y)$ ,  $l=1, 2$  of  $\mathcal{C}_{\varphi_j}(X_j(y, \nabla \varphi(y)))$  verify

$$(4.14) \quad 2 \min_{\substack{h=1,2 \\ x \in \Gamma}} K_h(x) \leq \kappa_{jl}(y) \leq 1/d_0 + 2/\delta_1 \max_{\substack{h=1,2 \\ x \in \Gamma}} K_h(x)$$

where  $d_0, \delta_1$  are the constants in Theorem 2.2 and Lemma 3.1 respectively.

Proof. In the proof we write  $L_j(y, \nabla \varphi(y))$  and  $X_j(y, \nabla \varphi(y))$  as  $L_j(y)$  and  $X_j(y)$  in brief. Note that  $\varphi(x)$  can be extended as  $C^\infty$ -function verifying  $|\nabla \varphi|=1$  in  $\{y+l\nabla \varphi(y); l \geq 0, y \in \Gamma_1 \cap \mathcal{U}\}$  because (i) holds. Suppose that there exist  $\varphi_0, \varphi_1, \dots, \varphi_j$ ,  $j \leq p-1$ , verifying (a), (b), (c) and (d). Taking account of (ii) it holds that

$$-n(X_{j+1}(y)) \cdot \Xi_j(y) \geq \delta_1,$$

from which it follows that

$$-\frac{\partial \varphi_j}{\partial n}(X_{j+1}(y)) \geq \delta_1.$$

Then there exists  $\psi(x)$  defined in a neighborhood of  $\bigcup_y X_{j+1}(y)$  verifying  $|\nabla \psi|=1$  and

$$\psi = \varphi_j, \quad -\frac{\partial \psi}{\partial n} = \frac{\partial \varphi_j}{\partial n} \quad \text{on } \bigcup_y X_{j+1}(y).$$

Since the principal curvatures of  $\mathcal{C}_{\varphi_j}(X_{j+1}(y))$  at  $X_{j+1}(y)$  are given by

$$\frac{\kappa_{jl}(y)}{1+L_j(y)\kappa_{jl}(y)}, \quad l=1, 2,$$

we see from Lemma 4.1 that the principal curvatures  $\kappa_{j+1l}(y)$  of  $\mathcal{C}_\psi(X_{j+1}(y))$  at  $X_{j+1}(y)$  satisfy

$$2 \min_{h,y} K_h(y) \leq \kappa_{j+1l}(y) \leq 1/l_j(y) + 2/\delta_1 \max_{h,y} K_h(y).$$

Taking account of  $l_j(y) \geq d_0$ ,  $\kappa_{j+1l}(y)$  satisfies (4.14). Now since  $\kappa_{j+1l}(y) > 0$ ,  $\psi(x)$  can be extended to  $C^\infty$ -function in  $\{X_{j+1}(y)+l\nabla \psi(X_{j+1}(y)); y \in \Gamma_1 \cap \mathcal{U}, l \geq 0\}$  verifying  $|\nabla \psi|=1$ . Setting this extended  $\psi(x)$  as  $\varphi_{j+1}$ , we see that  $\varphi_{j+1}(x)$  satisfies the required properties. Thus Proposition is proved.

### 5. Estimates of derivatives of the phase functions

Let the principal curvatures of  $\bar{\theta}_0 = \bar{\theta}_\varphi(x_0)$  be  $\kappa_1 \geq \kappa_2 > 0$  and let  $\{\omega(\sigma); \sigma \in U\}$  be a representation of  $\bar{\theta}_\varphi(x_0)$  near  $x_0$  such that

$$(5.1) \quad \begin{cases} \omega(0) = x_0 \\ \frac{\partial \omega}{\partial \sigma_i}(0) \cdot \frac{\partial \omega}{\partial \sigma_i}(0) = \delta_{ij} \\ \frac{\partial \nu}{\partial \sigma_j}(0) = \kappa_j \frac{\partial \omega}{\partial \sigma_j}(0), \quad j = 1, 2, \end{cases}$$

where  $\nu(\sigma) = (\nabla \varphi)(\omega(\sigma))$ . Take  $l > 0$  and set

$$x_1 = x_0 + l \nabla \varphi(x_0).$$

Let  $\{\tilde{\omega}(\eta); \eta \in \tilde{U}\}$  be a representation of  $\bar{\theta}_1 = \bar{\theta}_\varphi(x_1)$  near  $x_1$  satisfying

$$(5.2) \quad \begin{cases} \tilde{\omega}(0) = x_1 \\ \frac{\partial \tilde{\omega}}{\partial \eta_j}(0) = \frac{\partial \omega}{\partial \sigma_j}(0), \quad j = 1, 2. \end{cases}$$

Note that we can choose  $\omega(\sigma)$  and  $\tilde{\omega}(\eta)$  satisfying (5.1) and (5.2) in such a way

$$(5.3) \quad \begin{aligned} \sum_{|\gamma| \leq m} |(D_\sigma^\gamma \omega)(0)| &\leq C_m \sum_{|\gamma| \leq m} |(D_x^\gamma \varphi)(x_0)| \\ \sum_{|\gamma| \leq m} |(D_\eta^\gamma \tilde{\omega})(0)| &\leq C_m \sum_{|\gamma| \leq m} |(D_x^\gamma \varphi)(x_1)| \end{aligned}$$

where  $C_m$  is a constant independent of  $x_0, x_1$  and  $l$ .

On the other hand, since  $\varphi(x + l \nabla \varphi(x)) = \varphi(x) + l$  we have another representation of  $\bar{\theta}_1$

$$\bar{\theta}_1 = \{\omega(\sigma) + l\nu(\sigma); \sigma \in U\}.$$

Therefore for  $\eta$  sufficiently near 0 there corresponds  $\sigma \in U$  uniquely by the relation

$$(5.4) \quad \tilde{\omega}(\eta) = \omega(\sigma) + l\nu(\sigma).$$

Denote this correspondance as  $\sigma(\eta) = (\sigma_1(\eta_1, \eta_2), \sigma_2(\eta_1, \eta_2))$ . Now from (5.1) and (5.2) we have

$$(5.5) \quad \frac{\partial \sigma_i}{\partial \eta_j}(0) = (1 + l\kappa_j)^{-1} \delta_{ij}, \quad i, j = 1, 2.$$

Let us set

$$S^2 = \{(\xi_1, \xi_2, \xi_3); \xi_1^2 + \xi_2^2 + \xi_3^2 \leq 1\}.$$

For  $a=(a_1, a_2, a_3) \in S^2$  we set

$$X_a = a_1 \frac{\partial}{\partial x_1} + a_2 \frac{\partial}{\partial x_2} + a_3 \frac{\partial}{\partial x_3}.$$

DEFINITION 5.1. We define  $|\nabla \varphi|_m(x)$  for  $m=1, 2, \dots$  by

$$|\nabla \varphi|_m(x) = \max_{j \leq m} \max_{a(r) \in S^2} |(X_{a^{(1)}} X_{a^{(2)}} \cdots X_{a^{(j)}} \nabla \varphi)(x)|.$$

We may assume without generality that

$$\frac{\partial \omega}{\partial \sigma_1}(0) = (1, 0, 0), \quad \frac{\partial \omega}{\partial \sigma_2}(0) = (0, 1, 0), \quad \nu(0) = (0, 0, 1).$$

The correspondance from  $(\sigma, \tau)$  near  $(0, 0)$  to  $x$  near  $x_0$  given by

$$x = \omega(\sigma) + \tau \nu(\sigma)$$

verifies a relation

$$(5.6) \quad \frac{\partial \sigma_j}{\partial x_k}(x_0) = \delta_{jk}, \quad j, k = 1, 2, 3,$$

where we denote  $\tau$  as  $\sigma_3$ . Then we have

$$(X_a \nabla \varphi)(x_0) = \sum_{j=1}^2 a_j \frac{\partial \nu}{\partial \sigma_j}(0)$$

because  $\frac{\partial \nu(\sigma)}{\partial \tau} = 0$ . By the same way we have

$$(5.7) \quad (X_a \nabla \varphi)(x_1) = \sum_{j=1}^2 a_j \frac{\partial \mu}{\partial \eta_j}(0)$$

where  $\mu(\eta) = (\nabla \varphi)(\tilde{\omega}(\eta))$ . Since  $\mu(\eta) = \nu(\sigma(\eta))$  it follows from (5.5) and (5.7) that

$$(X_a) \nabla \varphi(x_1) = \sum_{j=1}^2 a_j (1 + l \kappa_j)^{-1} \frac{\partial \nu}{\partial \sigma_j}(0).$$

Taking account of (5.1) we have

$$\begin{aligned} |(X_a \nabla \varphi)(x_0)| &= \left( \sum_{j=1}^2 |a_j \kappa_j|^2 \right)^{1/2} \\ |(X_a \nabla \varphi)(x_1)| &= \left( \sum_{j=1}^2 |a_j \kappa_j (1 + l \kappa_j)^{-1}|^2 \right)^{1/2}. \end{aligned}$$

Thus we have

**Lemma 5.1.** For  $l > 0$  it holds that

$$(5.8) \quad |\nabla \varphi|_1(x_0 + l \nabla \varphi(x_0)) \leq \frac{1}{1 + l \kappa_2} |\nabla \varphi|_1(x_0).$$

Differentiating (5.4) by  $\eta_k$  and  $\eta_h$  we have

$$\begin{aligned} & \sum_{j=1}^2 \left( \frac{\partial \omega}{\partial \sigma_j} + l \frac{\partial \nu}{\partial \sigma_j} \right) \frac{\partial^2 \sigma_j}{\partial \eta_k \partial \eta_h} + \sum_{i,j=1}^2 \left( \frac{\partial^2 \omega}{\partial \sigma_i \partial \sigma_j} + l \frac{\partial^2 \nu}{\partial \sigma_i \partial \sigma_j} \right) \frac{\partial \sigma_i}{\partial \eta_k} \frac{\partial \sigma_j}{\partial \eta_h} \\ &= \frac{\partial^2 \tilde{\omega}}{\partial \eta_k \partial \eta_h}. \end{aligned}$$

Substitute  $\eta=0$  and take the scalar product with  $\frac{\partial \omega}{\partial \sigma_j}(0)$ . Then it follows that

$$\begin{aligned} & (1+l\kappa_j) \frac{\partial^2 \sigma_j}{\partial \eta_k \partial \eta_h} + l \frac{\partial \omega}{\partial \sigma_j} \frac{\partial^2 \nu}{\partial \sigma_k \partial \sigma_h} (1+l\kappa_k)^{-1} (1+l\kappa_h)^{-1} \\ &+ \frac{\partial \omega}{\partial \sigma_j} \frac{\partial^2 \omega}{\partial \sigma_k \partial \sigma_h} (1+l\kappa_k)^{-1} (1+l\kappa_h)^{-1} = \frac{\partial \omega}{\partial \sigma_j} \cdot \frac{\partial^2 \tilde{\omega}}{\partial \eta_k \partial \eta_h}. \end{aligned}$$

Taking account of (5.3) we have

$$(5.9) \quad \begin{cases} \frac{\partial^2 \sigma_j}{\partial \eta_k \partial \eta_h}(0) = -(1+l\kappa_j)^{-1} (1+l\kappa_k)^{-1} (1+l\kappa_h)^{-1} l \frac{\partial \omega}{\partial \sigma_j}(0) \frac{\partial^2 \nu}{\partial \sigma_k \partial \sigma_h}(0) + I_{j,k,h} \\ |I_{j,k,h}| \leq C_2 \{ |\nabla \varphi|_1(x_0) + |\nabla \varphi|_1(x_1) \} \end{cases}$$

where  $C_1$  is independent of  $x_2$ ,  $x_0$  and  $\varphi$ . Using (5.5) we have

$$\left. \frac{\partial^2(\nabla \varphi)(\tilde{\omega}(\eta))}{\partial \eta_k \partial \eta_h} \right|_{\eta=0} = \frac{\partial^2 \nu}{\partial \sigma_k \partial \sigma_h}(0) (1+l\kappa_k)^{-1} (1+l\kappa_h)^{-1} + \sum_{j=1}^2 \frac{\partial \nu}{\partial \sigma_j} \frac{\partial^2 \sigma_j}{\partial \eta_k \partial \eta_h}$$

Take the scalar product with  $\frac{\partial \omega}{\partial \sigma_j}(0)$  and substitute (5.9). Then we have

$$\begin{aligned} (5.10) \quad & \frac{\partial \omega}{\partial \sigma_j}(0) \cdot \left. \frac{\partial^2(\nabla \varphi)(\tilde{\omega}(\eta))}{\partial \eta_k \partial \eta_h} \right|_{\eta=0} \\ &= \frac{\partial \omega}{\partial \sigma_j}(0) \cdot \frac{\partial^2 \nu}{\partial \sigma_k \partial \sigma_h}(0) (1+l\kappa_k)^{-1} (1+l\kappa_h)^{-1} (1+l\kappa_j)^{-1} + \kappa_j I_{j,k,h}. \end{aligned}$$

Since  $\frac{\partial \nu(\sigma)}{\partial \sigma_j} \cdot \nu(\sigma) = 0$  we have

$$\frac{\partial^2 \nu}{\partial \sigma_j \partial \sigma_k} \cdot \nu = \left| \frac{\partial \nu}{\partial \sigma_j} \frac{\partial \nu}{\partial \sigma_k} \right| \leq C |\nabla \varphi|_1^2(x_0).$$

For  $a=(a_1, a_2, a_3)$ ,  $b=(b_1, b_2, b_3) \in S^2$

$$(5.11) \quad (X_b X_a \nabla \varphi)(x) = \sum_{i=1}^3 \sum_{j=1}^3 \sum_{k,h=1}^3 b_j a_i \left( \frac{\partial^2 \nu}{\partial \sigma_k \partial \sigma_h} \frac{\partial \sigma_k}{\partial x_j} \frac{\partial \sigma_h}{\partial x_i} + \frac{\partial \nu}{\partial \sigma_k} \frac{\partial^2 \sigma_k}{\partial x_j \partial x_i} \right).$$

Then using  $\frac{\partial \nu}{\partial \sigma_3} = 0$  we have

$$(5.12) \quad \begin{cases} (X_a X_b \nabla \varphi)(x_0) = \sum_{i,j=1}^2 a_i b_j \frac{\partial^2 \nu}{\partial \sigma_i \partial \sigma_j}(0) + II_0 \\ |II_0| \leq C |\nabla \varphi|_1(x_0). \end{cases}$$

By the same method we have

$$(5.13) \quad \begin{aligned} (X_a X_b \nabla \varphi)(x_1) &= \sum_{i,j=1}^2 a_i b_j \frac{\partial^2 (\nabla \varphi)(\tilde{\omega}(\eta))}{\partial \eta_i \partial \eta_j} \Big|_{\eta=0} + II_1 \\ |II_1| &\leq C \nabla |\varphi|_1(x_1). \\ &\left| \sum_{k,h=1}^2 a_k b_h \frac{\partial^2 \mu}{\partial \eta_k \partial \eta_h}(0) \right|^2 \\ &= \sum_{j=1}^2 \left| \sum_{k,h=1}^2 a_k b_h \left( \frac{\partial \omega}{\partial \sigma_j}(0) \cdot \frac{\partial^2 \mu}{\partial \eta_k \partial \eta_h}(0) \right) \right|^2 + II_1 \end{aligned}$$

using (5.10)

$$\begin{aligned} &= \sum_{j=1}^2 (1+l\kappa_j)^{-2} \left| \sum_{k,h=1}^2 \frac{a_k b_h}{(1+l\kappa_k)(1+l\kappa_h)} \left( \frac{\partial \omega}{\partial \sigma_j} \cdot \frac{\partial^2 \nu}{\partial \sigma_k \partial \sigma_h}(0) \right) \right|^2 + \tilde{II}_1 \\ &\leq (1+l\kappa_2)^{-2} |X_{\tilde{a}} X_{\tilde{b}} \nabla \varphi(x_0)|^2 + \tilde{II}_0 \end{aligned}$$

where

$$|\tilde{II}_j| \leq C |\nabla \varphi|_1(x_j), \quad j = 0, 1$$

and

$$\tilde{a} = \left( \frac{a_1}{1+l\kappa_1}, \frac{a_2}{1+l\kappa_2}, 0 \right), \quad \tilde{b} = \left( \frac{b_1}{1+l\kappa_1}, \frac{b_2}{1+l\kappa_2}, 0 \right).$$

Combining (5.11), (5.12) and (5.13) it holds that

$$\begin{aligned} &\max_{a,b \in S^2} |(X_a X_b \nabla \varphi)(x_1)| \\ &\leq \max_{a,b \in S^2} (1+l\kappa_2)^{-1} |(X_{\tilde{a}} X_{\tilde{b}} \nabla \varphi)(x_0)| + C \sum_{j=0}^1 |\nabla \varphi|_1(x_j) \\ &\leq (1+l\kappa_2)^{-3} \max_{a,b \in S^2} |(X_a X_b \nabla \varphi)(x_0)| + C \sum_{j=0}^2 |\nabla \varphi|_1(x_j). \end{aligned}$$

Thus, with the aid of (5.8), we have

**Lemma 5.2.** *For  $l > 0$  it holds that*

$$(5.14) \quad |\varphi|_2(x_0 + l \nabla \varphi(x_0)) \leq (1+l\kappa_2)^{-3} |\nabla \varphi|_2(x_0) + C_2 |\nabla \varphi|_1(x_0),$$

where  $C_2$  is a constant independent of  $x_0$  and  $\varphi$ .

REMARK. By the same reasoning we can show for  $m=3, 4, \dots$

$$(5.15) \quad |\nabla \varphi|_m(x_0 + l \nabla \varphi(x_0)) \leq (1+l\kappa_2)^{-(2m-1)} |\nabla \varphi|_m(x_0) + C_m |\nabla \varphi|_{m-2}(x_0).$$

Next consider the estimate of reflected rays. Let  $\varphi^\pm$  satisfy (4.1) and (4.2), and  $\omega(\sigma)$  and  $\tau(\sigma)$  be the same ones in § 4. We also suppose that  $\omega(\sigma)$  verifies (5.3). Since  $|(D_\sigma^\gamma i)(0)| \leq C_\gamma |\varphi^-|_{|\gamma|}(s_0)$  we have from (4.6)

$$|(D_\sigma^\gamma l)(0)| \leq C_\gamma |\varphi^-|_{|\gamma|-1}(s_0).$$

Using this estimate we have

$$|(D_\sigma^\gamma r)(0) - (D_\sigma^\gamma i)(0) + 2((D_\sigma^\gamma i)(0) \cdot n(s_0))n(s_0)| \leq C_\gamma |\nabla \varphi^-|_{|\gamma|-1}(s_0).$$

Evidently

$$\begin{aligned} \left| (X_a \nabla \varphi^-)(s_0) - \sum_{j=1}^2 a_j \frac{\partial i}{\partial \sigma_j}(0) \right| &\leq C |\nabla \varphi^-|_0(s_0) \\ \left| (X_a X_b \nabla \varphi^-)(s_0) - \sum_{k,j=1}^2 a_j b_k \frac{\partial^2 i}{\partial \sigma_j \partial \sigma_k}(0) \right| &\leq C |\nabla \varphi^-|_1(s_0). \end{aligned}$$

Concerning derivatives of  $\varphi^+$ , denoting

$$X_a = \xi_1 \frac{\partial}{\partial y_1} + \xi_2 \frac{\partial}{\partial y_2} + \xi_3 \frac{\partial}{\partial y_3}$$

where  $\xi_1 \frac{\partial \tau}{\partial \sigma_1}(0) + \xi_2 \frac{\partial \tau}{\partial \sigma_2}(0) + \xi_3 r(0) = a$ , we have

$$\begin{aligned} \left| (X_a \nabla \varphi^+)(s_0) - \sum_{j=1}^2 \xi_j \frac{\partial r}{\partial \sigma_j}(0) \right| &\leq C |\nabla \varphi^+|_0(s_0) \\ \left| (X_a X_b \nabla \varphi^+)(s_0) - \sum_{k,j=1}^2 \xi_j \gamma_k \frac{\partial^2 r}{\partial \sigma_j \partial \sigma_k}(0) \right| &\leq C |\nabla \varphi^+|_1(s_0). \end{aligned}$$

Taking account of  $|a| = |\xi|$  and that fact

$$|D_\sigma^\gamma i(0)| = |D_\sigma^\gamma i(0) - 2(D_\sigma^\gamma i(0) \cdot n(s_0))n(s_0)|$$

we have immediately

**Lemma 5.3.**

$$(5.16) \quad \begin{aligned} |\nabla \varphi^+|_1(s_0) &\leq |\nabla \varphi^-|_1(s_0) + C_1 |\nabla \varphi^+|_0(s_0) \\ |\nabla \varphi^+|_2(s_0) &\leq |\nabla \varphi^-|_2(s_0) + C_2 |\nabla \varphi^-|_1(s_0) \end{aligned}$$

REMARK. By the same reasoning we can show for  $m=2, 3$ ,

$$(5.17) \quad |\nabla \varphi^+|_m(s_0) \leq |\nabla \varphi^-|_m(s_0) + C_m |\nabla \varphi^-|_{m-1}(s_0).$$

Using the above lemmas let us show

**Proposition 5.4.** *Suppose that  $\varphi(x)$  satisfies the assumptions of Proposition 4.2. Let  $\varphi_j$ ,  $j=0, 1, 2, \dots, p$  be functions verifying (a)~(d) of Proposition 4.2. Then an estimate*

$$(5.18) \quad |\nabla \varphi_j|_m(x) \leq C_m |\nabla \varphi|_m(y)$$

holds for  $x \in L_j(y, \nabla \varphi(y))$ ,  $j=1, 2, \dots, p$ , where  $C_m$  is a constant independent of  $\varphi$  and  $p$ .

Proof. Note that (5.18) holds for  $m=0$  because  $|\nabla \varphi_j|=1$ ,  $\forall j$ . Suppose that (5.18) holds for  $m=1, 2, \dots, h$ . By Proposition 4.2 the principal curvatures  $\kappa_{ji}(y)$  of  $\mathcal{G}_{\varphi_j}(X_j(y))$  at  $X_j(y)$  satisfy

$$\kappa_{ji}(y) \geq K_0 = 2 \min_{\substack{j=1,2 \\ x \in \Gamma}} K_j(x).$$

Then by Lemma 5.2 and its remark we see that for  $j=1, 2, \dots, p$

$$|\nabla \varphi_j|_{h+1}(X_{j+1}(y)) \leq (1 + d_0 K_0)^{-2h-1} |\nabla \varphi_j|_{h+1}(X_j(y)) + C_h |\nabla \varphi_j|_h(X_j(y)),$$

and

$$|\nabla \varphi_0|_{h+1}(X_1(y)) \leq |\varphi|_{h+1}(y).$$

On the other hand Lemma 5.3 and its remark assure that

$$|\nabla \varphi_{j+1}|_{h+1}(X_{j+1}(y)) \leq |\nabla \varphi_j|_{h+1}(X_{j+1}(y)) + C_h |\nabla \varphi_j|_h(X_{j+1}(y))$$

holds for  $j=1, 2, \dots, p$ . Using the assumption on the estimate of  $|\varphi_j|_h$  we have an estimate

$$\begin{aligned} |\nabla \varphi_{j+1}|_{h+1}(X_{j+1}(y)) &\leq (1 + lK_0)^{-j(2h+1)} |\nabla \varphi|_{h+1}(y) \\ &\quad + 2C_h (1 + (1 + lK_0)^{-(2m+1)} + \dots + (1 + lK_0)^{-j(2m+1)}) |\nabla \varphi|_h(y). \end{aligned}$$

Thus our assurance is proved.

## 6. On the transport equation

First we will arrange some properties of asymptotic solution for  $\square w=0$  in the free space.

Let  $\varphi \in C^\infty(D)$ ,  $D \subset \mathbf{R}^3$  such that  $|\nabla \varphi|=1$ . Let us set

$$(6.1) \quad w(x, t; k) = e^{ik(\varphi(x)-t)} v(x, t; k)$$

$$(6.2) \quad v(x, t; k) = \sum_{j=0}^{\infty} v_j(x, t; k) k^{-j}.$$

Apply  $\square$  to  $w$  of (6.1) and we have

$$\begin{aligned} (6.3) \quad \square w &= -e^{ik(\varphi(x)-t)} \left\{ (ik)^2 ((\nabla \varphi)^2 - 1) v \right. \\ &\quad \left. + ik \left( 2 \frac{\partial v}{\partial t} + 2 \nabla \varphi \cdot \nabla v + \Delta \varphi \cdot v \right) - \square v \right\}. \end{aligned}$$

Substituting (6.2) into the right hand side of (6.3) and setting all the coefficients of  $k^{-j}$  equal to zero we obtain so called transport equations, that is,

$$(6.4)_0 \quad 2 \frac{\partial v_0}{\partial t} + 2 \nabla \varphi \cdot \nabla v_0 + \Delta \varphi \cdot v_0 = 0,$$

$$(6.4)_j \quad 2 \frac{\partial v_j}{\partial t} + 2 \nabla \varphi \cdot \nabla v_j + \Delta \varphi \cdot v_j + i \square v_{j-1} = 0, \quad \text{for } j \geq 1.$$

In an attempt to obtain successively the estimate of  $v_j$ , we consider the solution of the equation

$$(6.5) \quad 2 \frac{\partial u}{\partial t} + 2 \nabla \varphi \cdot \nabla u + \Delta \varphi \cdot u = h(x, t).$$

When  $\{x + l \nabla \varphi(x); l \in [0, l_0]\} \subset D$  the solution of (6.5) is represented as

$$(6.6) \quad u(x + l \xi(x), t + l) = \left[ \frac{G(x, l)}{G(x, 0)} \right]^{1/2} u(x, t) \\ + \int_0^l \left[ \frac{G(x, l)}{G(x, s)} \right]^{1/2} h(x + s \xi(x), t + s) ds$$

where  $\xi(x) = \nabla \varphi(x)$  and  $G(x, l)$  denotes the Gaussian curvature of  $\mathcal{C}_\varphi(x + l \xi(x))$  at  $x + l \xi(x)$ . This formula is due to Luneberg [7], Keller, Lewis and Seckler [4]. Let us denote the principal curvature of  $\mathcal{C}_\varphi(x)$  by  $\kappa_1(x)$  and  $\kappa_2(x)$ . Then those of  $\mathcal{C}_\varphi(x + l \xi(x))$  are  $\kappa_j(1 + l \kappa_j)^{-1}$ ,  $j = 1, 2$ . Thus we have

$$(6.7) \quad \frac{G(x, l)}{G(x, s)} = \frac{(1 + s \kappa_1)(1 + s \kappa_2)}{(1 + l \kappa_1)(1 + l \kappa_2)} \\ = \frac{1 + 2sH(x) + s^2G(x)}{1 + 2lH(x) + l^2G(x)},$$

where  $H(x)$  and  $G(x)$  denote the mean and Gaussian curvature of  $\mathcal{C}_\varphi(x)$  at  $x$ .

In the same way as § 5, we introduce the following norms. Set  $S^3 = \{(\xi_1, \xi_2, \xi_3, \xi_4); \sum_{j=1}^4 \xi_j^2 \leq 1\}$  and for  $a = (a_1, a_2, a_3, a_4) \in S^3$

$$X_a = \sum_{j=1}^3 a_j \frac{\partial}{\partial x_j} + a_4 \frac{\partial}{\partial t}.$$

DEFINITION. For  $w \in C^\infty(\mathcal{U} \times \mathbf{R}^1)$ ,  $\mathcal{U} \subset \mathbf{R}^3$  we set

$$|w|_m(x, t) = \max_{p \leq m} \max_{a^{(j)} \in S^3} |(X_{a^{(1)}} X_{a^{(2)}} \cdots X_{a^{(p)}} w)(x, t)|.$$

When  $x \in \mathcal{U} \cap \Gamma$  we also define another norm by

$$|w|'_m(x, t) = \max_{p \leq m} \max_{a^{(j)} \in S^{3'}} |(X_{a^{(2)}} X_{a^{(3)}} \cdots X_{a^{(p)}} w)(x, t)|,$$

where  $S^3 = S^3 \cap T_{(x,t)}(\Gamma \times \mathbf{R})$ , where  $T_{(x,t)}(\Gamma \times \mathbf{R})$  denotes the tangent space of  $\Gamma \times \mathbf{R}$  at  $(x, t)$ . And we set also for a subset  $\omega \subset \mathcal{U} \times \mathbf{R}$

$$|w|_m(\omega) = \max_{(x,t) \in \omega} |w|_m(x, t)$$

and for  $\omega' \subset \Gamma \times \mathbf{R}$

$$|w|'_m(\omega') = \max_{(x,t) \in \omega'} |w|'_m(x, t).$$

**Lemma 6.1.** *Let  $m$  be a non negative integer. For the solution (6.6) it holds that an estimate*

$$(6.8) \quad \begin{aligned} & |w|_m(x+l\xi(x), t+l) \\ & \leq (1+l\kappa_2(x))^{-1} |w|_m(x, t) + \int_0^l |h|_m(x+s\xi(x), t+s) ds \\ & \quad + C_m |\nabla \varphi|_m(x) \left\{ |w|_{m-1}(x, t) + |w|_{m-1}(x+l\xi(x), t+l) \right. \\ & \quad \left. + |h|_{m-1}(x+l\xi(x), t+l) + \int_0^l |h|_{m-1}(x+s\xi(x), t+s) ds \right\}, \end{aligned}$$

where  $C_m$  is a positive constant independent of  $w$  and  $\varphi$ . Especially, for  $h=0$  and  $m=0$  we have

$$(6.9) \quad |w(x+l\xi(x), t+l)| \leq (1+l\kappa_2(x))^{-1} |w(x)|.$$

**Proof.** Note that (6.9) follows immediately from (6.6) and (6.7). Let  $\{\omega(\sigma); \sigma \in U\}$  be a representation of  $\mathcal{C}_\varphi(x_0)$  which verifies  $\omega(0)=x_0$  and the condition (4.4), (4.5) and (5.3), where  $i(\sigma)=(\nabla \varphi)(\omega(\sigma))$ . Suppose that  $\frac{\partial \omega}{\partial \sigma_j}(0), j=1, 2$  and  $\xi(x_0)$  coincide with  $x_1, x_2$  and  $x_3$  axis respectively. Consider a mapping  $\mathbf{R}^4 \ni (\sigma_1, \sigma_2, l, \tau) \rightarrow (x_1, x_2, x_3, t) \in \mathbf{R}^4$  defined by

$$\begin{cases} x = x(\sigma, l) = \omega(\sigma) + li(\sigma) \\ t = \tau + l. \end{cases}$$

Then the Jacobian matrix of the mapping at  $\sigma=0$  is

$$(6.10) \quad \begin{pmatrix} 1+l\kappa_1 & 0 & 0 & 0 \\ 0 & 1+l\kappa_2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

Moreover, the inverse mapping satisfies

$$(6.11) \quad |D_{x,t}^\gamma \sigma_j| \leq C_\gamma |\nabla \varphi|_{|\gamma|}, \quad j = 1, 2, 3, 4$$

where we denote  $l=\sigma_3, \tau=\sigma_4$ . Let us set for some  $l>0$

$$x_1 = x_0 + l\xi(x_0), \quad t_1 = t + l.$$

First we show (6.8) for  $x=x_0$  in the case  $h \equiv 0$ . Suppose that for  $a^{(j)} = (a_i^{(j)})_{i=1,2,3,4} \in S^3, j=1, 2, \dots, p \leq m$  it holds that

$$(6.12) \quad |w|_m(x_1, t_1) = |(X_{a^{(1)}} X_{a^{(2)}} \cdots X_{a^{(p)}} w)(x_1, t_1)|.$$

On the other hand, taking account of (6.10) and (6.11) we have

$$(6.13) \quad \left| \left( \left( \prod_{j=1}^p X_{a^{(j)}} \right) w \right)(x_1, t_1) - \left[ \left( \prod_{j=1}^p \left( \sum_{i=1}^p a_i^{(j)} \frac{1}{1+l\kappa_i(x_0)} \frac{\partial}{\partial \sigma_i} + a_3^{(j)} \left( \frac{\partial}{\partial l} - \frac{\partial}{\partial \tau} \right) + a_4^{(j)} \frac{\partial}{\partial \tau} \right) \right) w(\omega(\sigma) + li(\sigma), \tau + l) \right]_{\sigma=0}^{\tau=t} \right| \\ \leq C_p |\nabla \varphi|_p(x_1) |w|_{p-1}(x_1, t_1).$$

Note that it follows from (6.7) that for  $0 \leq s \leq l$ .

$$\left| D_{\sigma, s}^\gamma \frac{G(\omega(\sigma), l)}{G(\omega(\sigma), s)} \right| \leq C_\gamma |\nabla \varphi|_{|\gamma|}(\omega(\sigma)).$$

Then using (6.6)

$$(6.14) \quad \left| [Y_1 Y_2 \cdots Y_p w(\omega(\sigma) + li(\sigma), \tau + l)]_{\sigma=0}^{\tau=t} - \left( \frac{G(x_0, l)}{G(x_0, 0)} \right)^{1/2} \left[ \prod_{j=1}^p \left( \sum_{k=1}^2 a_k^{(j)} \frac{1}{1+l\kappa_k(x_0)} \frac{\partial}{\partial \sigma_k} + (a_4^{(j)} - a_3^{(j)}) \frac{\partial}{\partial \tau} \right) \right] w(\omega(\sigma), \tau) \right]_{\sigma=0}^{\tau=t} \right| \\ \leq C_p |\nabla \varphi|_p |w|_{p-1}(x_j, t),$$

where

$$Y_j = \sum_{k=1}^2 a_k^{(j)} \frac{1}{1+l\kappa_k(x_0)} \frac{\partial}{\partial \sigma_k} + a_3^{(j)} \left( \frac{\partial}{\partial l} - \frac{\partial}{\partial \tau} \right) + a_4^{(j)} \frac{\partial}{\partial \tau}.$$

By the same consideration we have

$$\left| \left( \left( \prod_{j=1}^p X_{b^{(j)}} \right) w \right)(x_0, t) - \left[ \left( \prod_{j=1}^p \left( \sum_{k=1}^2 b_k^{(j)} \frac{\partial}{\partial \sigma_k} + (b_4^{(j)} - b_3^{(j)}) \frac{\partial}{\partial \tau} \right) \right) w(\omega(\sigma), \tau) \right]_{\sigma=0}^{\tau=t} \right| \\ \leq C_p |\nabla \varphi|_{p-1}(x_0) |w|_{p-1}(x_0, t).$$

Therefore, setting

$$b^{(j)} = \left( a_1^{(j)} \frac{1}{1+l\kappa_1(x_0)}, a_2^{(j)} \frac{1}{1+l\kappa_2(x_0)}, a_3^{(j)}, a_4^{(j)} \right)$$

we have with the aid of (6.13) and (6.14)

$$\left| \left( \left( \sum_{j=1}^p X_{a^{(j)}} \right) w \right)(x_1, t_1) - \left[ \frac{G(x_0, l)}{G(x_0, 0)} \right]^{1/2} \left( \prod_{j=1}^p X_{b^{(j)}} \right) w(x_0, t) \right| \\ \leq C_p |\nabla \varphi|_p(x_0) \{ |w|_{p-1}(x_0, t) + |w|_{p-1}(x_1, t_1) \}.$$

From (6.12) we have

$$|w|_m(x_1, t_1) \leq \left[ \frac{G(x_0, l)}{G(x_0, 0)} \right]^{1/2} |((\prod_{j=1}^p X_{\delta(j)})w)(x_0, t)| \\ + C_p |\nabla \varphi|_p(x_0) \{ |w|_{p-1}(x_0, t) + |w|_{p-1}(x_1, t_1) \}.$$

Since  $\left( \frac{G(x_0, l)}{G(x_0, 0)} \right)^{1/2} \leq \frac{1}{1+l\kappa_2(x_0)}$  and  $|w|_m(x_0, t) \geq |((\prod_{j=1}^p X_{\delta(j)})w)(x_0, t)|$  we have (6.8).

When  $h \neq 0$ , setting for each  $0 \leq s \leq l$

$$h_s(x + l\xi(x), t + l) = \left[ \frac{G(x, l)}{G(x, s)} \right]^{1/2} h(x + s\xi(x), t + s)$$

we have from the above result for all  $0 \leq s \leq l$

$$|h_s|_m(x_1, t_1) \leq \left( \frac{G(x_0, l)}{G(x_0, s)} \right)^{1/2} |h_s|_m(x_0 + s\xi(x_0), t + s) \\ + C_m |\nabla \varphi|_{m-1}(x_0) \{ |h_s|_{m-1}(x_1, t_1) + |h_s|_{m-1}(x_0 + s\xi(x_0), t + s) \}.$$

Using this estimate we have immediately (6.8) from (6.6).

**Lemma 6.2.** Let  $\varphi^+, \varphi^-$  be functions verifying  $|\varphi^\pm| = 1$ , (4.1) and (4.2). Suppose that  $h^+, h^- \in C^\infty(\mathcal{U} \times \mathbf{R})$  and  $w^+, I^-$  satisfy

$$(6.15) \quad 2 \frac{\partial w^\pm}{\partial t} + 2 \nabla \varphi^\pm \cdot \nabla w^\pm + \Delta \varphi^\pm \cdot w^\pm = h^\pm \quad \text{in } \mathcal{U} \times \mathbf{R}$$

$$(6.16) \quad w^+ = aw^- + f(x, t) \quad \text{on } (\Gamma \cap \mathcal{U}) \times \mathbf{R},$$

where  $|a| = 1$ . Then it holds that for all  $(x, t) \in (\Gamma \cap \mathcal{U}) \times \mathbf{R}$

$$|w^+|_m(x, t) \leq |w^-|_m(x, t) + |f|'_m(x, t) + C_m |\nabla \varphi^-|_m(x) \{ |w^+|_{m-1}(x, t) \\ + |w^-|_{m-1}(x, t) + |h^+|_m(x, t) + |h^-|_{m-1}(x, t) \}$$

where  $C_m$  is a constant independent of  $w^\pm, h^\pm$  and  $\varphi_\pm$ .

Since we can prove this lemma as almost the same way as Lemma 5.3, we omit the proof.

## 7. Construction of asymptotic solutions

Let  $s_0 \in \Gamma_1$  and  $\mathcal{U}$  be a small neighborhood of  $s_0$  in  $\mathbf{R}^3$ . Suppose that  $\theta(x, \eta, \beta), \rho(x, \eta, \beta)$  are defined for  $x \in \mathcal{U}, \eta \in \Sigma = \{(\eta_1, \eta_2); \eta_1^2 + \eta_2^2 = 1, \beta \in [-\beta_0, \beta_0]\}$  ( $\beta_0 > 0$ ) and they satisfy

$$(7.1) \quad \begin{cases} (\nabla \theta)^2 + \rho(\nabla \rho)^2 \equiv 1 & (\text{mod } \beta^\infty) \text{ in } \Omega^{(1)} \cap \mathcal{U} \\ \nabla \theta \cdot \nabla \rho \equiv 0 & (\text{mod } \beta^\infty) \text{ in } \Omega^{(1)} \cap \mathcal{U} \\ \rho = -\beta & \text{on } \Gamma_1 \cap \mathcal{U} \end{cases}$$

$$(7.2) \quad (\nabla\theta)(s_0, \eta, 0) = \eta_1 \frac{\partial}{\partial\sigma_1} + \eta_2 \frac{\partial}{\partial\sigma_2},$$

where  $\Omega^{(j)} = \mathbf{R}^3 - \mathcal{O}_j$ ,  $j = 1, 2$  and  $\{y(\sigma); \sigma \in U\}$  be a representation of  $\Gamma_1$  near  $s_0$ .

**Lemma 7.1.** *Let  $u$  be an oscillatory data given by*

$$(7.3) \quad \begin{aligned} u(x, t; \eta, \beta, k) &= e^{ik(\theta(x, \eta, \beta) - t)} f(x, t; k) \\ f(x, t; k) &\in C_0^\infty((\Gamma_1 \cap \mathcal{U}) \times \mathbf{R}). \end{aligned}$$

*For any positive integer  $N$ , there exists an asymptotic solution  $w^{(N)}(x, t; \eta, \beta, k)$  with the following properties:*

$$(7.4) \quad \text{supp } w^{(N)} \subset \bigcup_{(x, t) \in \text{supp } f} \{x + l\nabla(\theta + 2/3 \rho^{3/2})(x), t + l; l \geq 0\}$$

$$(7.5) \quad |w^{(N)}|_m(\Omega_R^{(1)} \times \mathbf{R}) \leq C_{N, m, R} k^m \left( \sum_{j=1}^N |f|'_{20j+m}(\Gamma_1 \times \mathbf{R}) k^{-j} \right)$$

$$(7.6) \quad \text{In } \{x; \text{dis}(x, \mathcal{O}_1) > d_0/2\} \quad w^{(N)} \text{ is of the form}$$

$$\begin{aligned} w^{(N)} &= e^{ik(\varphi - t)} v, \quad v = \sum_j v_j k^{-j} \\ |v_j|_m(\Omega_R^{(1)} \times \mathbf{R}) &\leq C_{j, m, R} |f|'_{20j}(\Gamma_1 \times \mathbf{R}) \end{aligned}$$

*and the principal curvatures of  $\mathcal{O}_\varphi(x)$  are positive.*

$$(7.7) \quad |\square w^{(N)}|_m(\Omega_R^{(1)} \times \mathbf{R}) \leq C_{N, m, R} k^{-N+m} |f|_{N'+m}(\Gamma_1 \times \mathbf{R})$$

$$(7.8) \quad |Bw^{(N)} - u|'_m(\Gamma_1 \times \mathbf{R}) \leq C_{N, m} k^{-N+m} |f|'_{N'+m}(\Gamma_1 \times \mathbf{R})$$

where  $\Omega_R^{(1)} = \Omega^{(1)} \cap \{x; |x| \leq R\}$  and  $N' = 20N$ .

Since this lemma may be proved by the same process treated in §4 and 6 of [2] we omit the proof.

**Lemma 7.2.** *Let  $\mathcal{U}$  be a neighborhood of  $\mathcal{O}_1$  and let  $y(x, t; k)$  be a function defined in  $\mathcal{U} \times \mathbf{R}$  of the form*

$$\begin{aligned} y(x, t; k) &= e^{ik(\varphi - t)} v(x, t; k) \\ \text{supp } y &\subset \mathcal{U} \times [0, \infty), \quad |\nabla\varphi(x)| = 1. \end{aligned}$$

*Suppose that for some  $\delta_4 > 0$  it holds that*

$$(7.9) \quad |\nabla\varphi(x) \cdot n(x)| \leq 2\delta_4, \quad \forall x \in \text{Proj}(\text{supp } y \cap \Gamma_1 \times \mathbf{R}),$$

*where Proj denotes the projection to the  $x$ -space. Then we can construct for any  $N$  positive integer a function  $z^{(N)}(x, t; k)$  defined in  $\Omega^{(1)} \times \mathbf{R}$  with the following properties:*

$$(7.10) \quad |z^{(N)}|_m(\Omega_R^{(1)} \times \mathbf{R}) \leq C_{N,m} k^{m+1} |v|'_m(\Gamma_1 \times \mathbf{R})$$

$$(7.11) \quad \text{supp } z^{(N)} \subset \bigcup_{(x,t) \in \text{supp } v} \bigcup_{\xi \in \Sigma_x^+} \mathcal{L}_0(x, t; \xi)$$

$$(7.12) \quad |\square z^{(N)}|_m(\Omega_R^{(1)} \times \mathbf{R}) \leq C_{N,m,R} k^{-N+m} |v|'_{N'+m}(\Gamma_1 \times \mathbf{R})$$

$$(7.13) \quad |B(y + z^{(N)})|'_m(\Gamma_1 \times \mathbf{R}) \leq C_{N,m} k^{-N+m} |v|'_{N'+m}(\Gamma_1 \times \mathbf{R}).$$

Proof. We may suppose that  $\text{supp } y \cap (\Gamma_1 \times \mathbf{R})$  is sufficiently small. Then, by using  $\mathcal{V}_j$  of § 4 of [3] we have

$$y(x, t; k) = \sum_{j=1}^4 \mathcal{V}_j y \quad \text{on } \Gamma_1 \times \mathbf{R}.$$

The definition  $\mathcal{V}_j$  gives

$$\begin{aligned} (\mathcal{V}_j y)(s(\sigma), t) &= \omega(s(\sigma), t) \int_{\mathbf{R}^1} d\tau \int_{-1}^{\infty} d\alpha \int_{\Sigma} d\xi' \int_{I_\sigma} d\sigma' \int dt' \\ &\quad \exp \{i\tau(-t+t' + (1+\alpha)\langle \sigma - \sigma', \xi' \rangle)\} \chi_j(1+\alpha^2) k^2 (1+\alpha) \\ &\quad \tilde{\omega}(s(\sigma'), t') e^{ik} v(s(\sigma'), t'; k). \end{aligned}$$

Taking out the integration with respect to  $t'$ , if  $|\tau - k| \geq \varepsilon k$  we have an estimate for any  $M$

$$\left| \int e^{i(\tau-k)t'} v(s(\sigma'), t'; k) dt' \right| \leq C_{\varepsilon, M} k^{-M} |v|'_M(\Gamma_1 \times \mathbf{R}).$$

For verifying  $|\tau - k| \leq \varepsilon k$ , since  $\varphi$  verifies (7.9) it holds that for  $j \neq 2$

$$|\text{grad}_{\sigma'}(\varphi(s(\sigma'))) - (1+\alpha)\langle \sigma', \xi' \rangle| \geq \alpha_0/2$$

when  $\chi_j(1+\alpha) \neq 0$  if we choose  $\varepsilon$  and  $\delta_4$  sufficiently small. Therefore we have for  $j \neq 2$

$$\begin{aligned} &\left| \int i^{\tau\{\varphi(s(\sigma')) - (1+\alpha)\langle \sigma', \xi' \rangle\}} \chi_j(1+\alpha^2) v(s(\sigma'))(t'; k) d\sigma' \right| \\ &\leq C_M |\tau|^{-M} |v|'_M(\Gamma_1 \times \mathbf{R}). \end{aligned}$$

These estimates show that when (7.9) holds for small  $\delta_4 > 0$

$$|\tilde{\mathcal{V}}_2 y - y|'_M(\Gamma_1 \times \mathbf{R}) \leq C_{N,m} k^{-N+m} |v|'_N(\Gamma_1 \times \mathbf{R}),$$

where  $\tilde{\mathcal{V}}_2$  denotes an operator replacing  $\int_{-\infty}^{\infty} d\tau$  in the definition of  $\mathcal{V}$  by  $\int_{|\tau-k| \leq \varepsilon k} d\tau$ . By using the change of variables of page 87 of [3] it follows that

$$\begin{aligned} (\tilde{\mathcal{V}}_2 y)(s(\sigma), t) &= \omega(s(\sigma), t) \int_{|\tau-k| \leq \varepsilon k} d\tau \int_{\Sigma} d\eta \int_{|\beta| \leq \beta_0} d\beta \int_{I_\sigma} d\sigma' \int dt' \\ &\quad \exp \{i\tau(\theta(s(\sigma), \eta, \beta) - \theta(s(\sigma'), \eta, \beta)) + t - t'\} \end{aligned}$$

$$\begin{aligned} & \cdot \chi_2(1+\alpha)^2 k^2(1+\alpha) \frac{D(\xi', \alpha)}{D(\eta, \beta)} \tilde{\omega}(s(\sigma'), t') e^{ik(\varphi(s(\sigma'))-t')} \\ & \cdot v(s(\sigma'), t'; k). \end{aligned}$$

By applying Lemma 7.1 to an oscillatory boundary data

$$u(x, t; \beta, \eta, k) = e^{ik(\theta(x, \eta, \beta)-t)} \omega(x, t) \chi_2(1+\alpha) k^2(1+\alpha) \frac{D(\xi', \alpha)}{D(\eta, \beta)}$$

we obtain  $w^{(N)}(x, t; \eta, \beta, k, \sigma')$  verifying (7.4)~(7.8). Let us set

$$\begin{aligned} z^{(N)}(x, t; k) = & \int_{|\tau-k| \leq \varepsilon k} d\tau \int_{\Sigma} d\eta \int_{|\beta| \leq \beta_0} d\beta \int_{I_\sigma} d\sigma' \int dt' \\ & w^{(N)}(x, t; \eta, \beta, k, \sigma') e^{i\tau(\theta(s(\sigma'), \eta, \beta)-t')} e^{ik(\varphi(s(\sigma'))-t')} v(s(\sigma'), t'; k). \end{aligned}$$

Then we see immediately that  $z^{(N)}$  is the required solution.

**DEFINITION.** Let  $\mathcal{U}$  be a neighborhood of  $s_0 \in \Gamma_1$  and let  $\psi(x)$  be a real valued function on  $\Gamma_1 \cap \mathcal{U}$ . We say that  $\psi(x)$  satisfies Condition C in  $\mathcal{U}$  if there exists a real valued function  $\varphi(x)$  defined in  $\mathcal{U}$  verifying

$$\begin{cases} \varphi(x) = \psi(x) & \text{in } \Gamma_1 \cap \mathcal{U} \\ |\nabla \varphi| = 1 & \text{in } \mathcal{U} \\ \frac{\partial \varphi}{\partial n}(x) \geq \delta_4 & \text{on } \Gamma_1 \cap \mathcal{U} \end{cases}$$

the principal curvatures of  $\mathcal{G}_\varphi(x)$  are positive for all  $x \in \Gamma_1 \cap \mathcal{U}$ .

**Lemma 7.3.** Let  $u$  be an oscillatory data on  $\Gamma_1 \times \mathbf{R}$  in the form

$$(7.14) \quad u(x, t; k) = e^{ik(\psi(x)-t)} f(x, t; k)$$

where  $\psi$  satisfies the condition C in  $\mathcal{U}$  and  $f(x, t; k) \in C_0^\infty((\Gamma_1 \cap \mathcal{U}) \times \mathbf{R})$ . Then we have  $v_j(x, t; k) \in C^\infty(\Omega^{(1)} \times \mathbf{R})$ ,  $j=0, 1, 2, \dots$  with the following properties:

$$(7.15) \quad \text{supp } v_j \subset \bigcup_{(x,t) \in \text{supp } f} \overline{\mathcal{L}_0}(x, t; \nabla \varphi(x))$$

$$(7.16) \quad |v_j|_m(\Omega_R^{(1)} \times \mathbf{R}) \leq C_{j,m,R} |\nabla \varphi|_{2m+j}(\Omega^{(1)}) |f|'_{m+2j}(\Gamma_1 \times \mathbf{R})$$

For an  $N$  positive integer, if we set

$$(7.17) \quad w^{(N)}(x, t; k) = e^{ik(\varphi(x)-t)} \sum_{j=0}^N v_j(x, t; k) k^{-j}$$

it holds that

$$(7.18) \quad \square w^{(N)} = e^{ik(\varphi-t)} k^{-N} \square v_N \quad \text{in } \Omega^{(1)} \times \mathbf{R}$$

$$(7.19) \quad Bw^{(N)} - u = e^{ik(\psi-t)} k^{-N} Bv_N \quad \text{on } \Gamma_1 \times \mathbf{R}.$$

REMARK. Combining (7.16) and (7.18) we have

$$(7.20) \quad |\square w^{(N)}|_m(\Omega_R^{(1)} \times \mathbf{R}) \leq C_{N,m,R} k^{-N+m} |f|'_{m+2N+2}(\Gamma_1 \times \mathbf{R})$$

and from (7.16) and (7.19) we have

$$(7.21) \quad |Bw^{(N)} - u|_m(\Gamma_1 \times \mathbf{R}) \leq C_{N,m,R} k^{-N+m} |f|'_{m+2N+2}(\Gamma_1 \times \mathbf{R}).$$

When  $B=B_1$  from method of the construction we have  $Bv_N=0$ .

Proof. If we look for any asymptotic solution in a form (6.1) from the consideration in the beginning of § 6, we require that  $v_j$  is the solution of (6.4)<sub>j</sub> in  $\Omega^{(1)}$ . In the case of  $B=B_1$  we set a boundary condition

$$v_0 = f, \quad v_j = 0 \quad \text{for } j \geq 1 \quad \text{on } \Gamma_1 \times \mathbf{R}.$$

In the case of  $B=B_2$ , since for  $w^{(N)}$  of (7.17)

$$Bw^{(N)} = e^{ik(\varphi-t)} \sum_{j=1}^N \left( ik \frac{\partial \varphi}{\partial n} v_j + \frac{\partial v_j}{\partial n} + \sigma v_j \right) k^{-j}$$

we set the following boundary condition for  $v_j$  on  $\Gamma_1 \times \mathbf{R}$

$$v_0 = (ik)^{-1} f$$

and for  $j \geq 1$

$$v_j = i \left( \frac{\partial v_{j-1}}{\partial n} + \sigma v_{j-1} \right) \left( \frac{\partial \varphi}{\partial n} \right)^{-1}$$

With the aid of Lemma 6.1 we obtain successively the estimate (7.16), and using (6.6) we have also successively (7.15). On the other hand, since  $v_j$  satisfies (6.4)<sub>j</sub> we have (7.18) from (6.3). And (7.19) follows from the boundary condition which  $v_j$  satisfy. Q.E.D.

**Lemma 7.4.** Let  $w^{(N)}(x, t; k)$  be a function constructed in the previous lemma. Suppose that

$$(7.22) \quad -n(X_1(x, \nabla \varphi(x))) \cdot \nabla \varphi(x) \geq \delta_4 \quad \forall x \in \text{Proj}(\text{supp } f).$$

Then there exists a function  $\tilde{w}^{(N)}(x, t; k)$  defined in  $\Omega^{(2)} \times \mathbf{R}$  in the form

$$(7.23) \quad \tilde{w}^{(N)} = e^{ik(\tilde{\varphi}-t)} \sum_{j=0}^N \tilde{v}_j(x, t; k) k^{-j}$$

satisfying

$$(7.24) \quad \text{supp } \tilde{v}_j \subset \bigcup_{(x,t) \in \text{supp } f} \mathcal{L}_1(x, t; \nabla \varphi(x))$$

$$(7.25) \quad |\tilde{v}_j|_m(\Omega_R^{(2)} \times \mathbf{R}) \leq C_{j,m,R} |f|'_{m+2j}(\Gamma_1 \times \mathbf{R})$$

$$(7.26) \quad \square \tilde{w}^{(N)} = e^{ik(\tilde{\varphi}-t)} k^{-N} \square \tilde{v}_N$$

$$(7.27) \quad B(w^{(N)} + \tilde{w}^{(N)}) = e^{ik(\varphi-t)} k^{-N} (Bv_N + B\tilde{v}_N).$$

Proof. Taking account of Proposition 4.2 there exists  $\tilde{\varphi}$  defined in  $\omega_1 = \bigcup_{\text{Proj } x \in \text{supp } f} L_1(x, \nabla \varphi(x))$ , which verifies

$$\begin{cases} |\nabla \tilde{\varphi}| = 1 & \text{in } \omega_1 \\ \tilde{\varphi} = \varphi & \text{on } \Gamma_2 \\ \frac{\partial \tilde{\varphi}}{\partial n} = -\frac{\partial \varphi}{\partial n} & \text{on } \Gamma_2. \end{cases}$$

Let  $\tilde{v}_j$  be a solution of the equation (6.4)<sub>j</sub> replaced  $\varphi$  by  $\tilde{\varphi}$ . We set a boundary condition for  $\tilde{v}_j$  on  $\Gamma_2$ .

$$(7.28) \quad \tilde{v}_j = \begin{cases} -v_j & \text{when } B = B_1 \\ v_j + i \left( \frac{\partial \tilde{\varphi}}{\partial n} \right)^{-1} (Bv_{j-1} + B\tilde{v}_{j-1}) & \text{when } B = B_2. \end{cases}$$

By using Lemma 6.2 we have

$$\begin{aligned} |\tilde{v}_j|_m(\Gamma_2 \times \mathbf{R}) &\leq |v_j|_m(\Gamma_2 \times \mathbf{R}) + |v_{j-1}|'_m(\Gamma_2 \times \mathbf{R}) \\ &\quad + C_m |\nabla \varphi|_m \{ |v_{j-1}|_m(\Gamma_2 \times \mathbf{R}) + |v_{j-1}|_{m-1}(\Gamma_2 \times \mathbf{R}) \\ &\quad + |\square \tilde{v}_{j-1}|_{m-1}(\Gamma_2 \times \mathbf{R}) \}. \end{aligned}$$

And from Lemma 6.1 we have

$$|\tilde{v}_j|_m(\Omega_R^{(2)} \times \mathbf{R}) \leq \tilde{C}_{m,R} |\nabla \tilde{\varphi}|_m \{ |\tilde{v}|_m(\Gamma_2 \times \mathbf{R}) + |\square \tilde{v}_{j-1}|_m(\Omega_R^{(2)} \times \mathbf{R}) \}.$$

Then the estimate (7.25) can be derived inductively from the above two estimates. Taking account of

$$\bar{\mathcal{L}}_0(X_1(x), \Xi_1(x)) = \bar{\mathcal{L}}_1(x, \nabla \varphi(x)),$$

we have immediately (7.24). (7.26) and (7.27) follows from the equations and boundary conditions which  $v_j, j=0, 1, 2$ , satisfy.

REMARK. Applying (7.25) to (7.26) we have

$$(7.29) \quad |\square \tilde{w}^{(N)}|_m(\Omega_R^{(2)} \times \mathbf{R}) \leq C_{N,m,R} k^{-N+m} |f|'_{m+2N+2}(\Gamma_1 \times \mathbf{R}).$$

Similary we have

$$(7.30) \quad |B(w^{(N)} + \tilde{w}^{(N)})|_m(\Gamma_2 \times \mathbf{R}) \leq C_{N,m} k^{-N+m} |f|'_{m+2N+2}(\Gamma_1 \times \mathbf{R}).$$

**Proposition 7.5.** *Let  $J$  be a positive integer and let  $u(x, t; k)$  an oscillatory data on  $\Gamma_1 \times \mathbf{R}$  of the form (7.3) or of the form (7.24) with  $\psi$  satisfying the condition C. Suppose that*

$$(7.31) \quad {}^{\sharp} \mathcal{X}(x, \nabla \varphi(x)) \leq J \quad \text{for all } x \in \text{Proj supp } f,$$

where for  $u$  of the form (7.3)  $\varphi(x)$  means  $\theta + 2/3 \rho^{3/2}$ . Then for any  $N$  positive

integer there exists a function  $w^{(N)}(x, t; k) \in C^\infty(\bar{\Omega} \times R)$  such that

$$(7.32) \quad \text{supp } w^{(N)} \subset \bigcup_{(x,t) \in \text{supp } f} \mathcal{L}(x, t; \nabla \varphi(x))$$

$$(7.33) \quad |w^{(N)}|_m(\Omega_R \times R) \leq C_{J,N,m,R} |\nabla \varphi|_{m+N'} k^{m+1} \sum_{j=0}^{N'} k^{-j} |f|_{m+2j+40}(\Gamma_1 \times R)$$

$$(7.34) \quad |\square^{(N)}|_m(\Omega_R \times R) \leq C_{J,N,m,R} |\nabla \varphi|_{m+N'} k^{-N+n} |f|_{N'+m}(\Gamma_1 \times R)$$

$$(7.35) \quad |B^{(N)} - u|_m(\Gamma \times R) \leq C_{J,N,m,R} |\nabla \varphi|_{m+N'} k^{-N+m} |f|_{N'+m}(\Gamma_1 \times R)$$

where  $N' = 20N + 40$  and for  $u$  of the form (7.3) we set  $\theta$  in the place of  $\varphi$  of the above estimates.

Proof. First we consider the case of  $u$  of the form (7.24). Let  $(s_0, t_0) \in \text{supp } f$  and let us set

$$*\mathcal{X}(s_0, \nabla \varphi(s_0)) = K_0$$

and  $X_j^0 = X_j(s_0, \nabla \varphi(s_0))$ ,  $\Xi_j^0 = \Xi_j(s_0, \nabla \varphi(s_0))$ . Taking account of Lemma 3.1, for fixed  $\varepsilon > 0$  there are three cases:

Case 1.  $n(X_j^0) \cdot \Xi_j^0 \geq \delta_1$ ,  $j = 1, 2, \dots, K_0$  and if  $X_{K_0}^0 \in \Gamma_1(\Gamma_2)$   $L_{K_0}(s_0, \nabla \varphi(s_0))$  does not pass the  $\varepsilon$ -neighborhood of  $\mathcal{O}_2(\mathcal{O}_1)$ .

Case 2.  $n(X_j^0) \cdot \Xi_j^0 \geq \delta_1$ ,  $j = 1, 2, \dots, K_0$  and if  $X_{K_0}^0 \in \Gamma_1(\Gamma_2)$   $L_{K_0}(s_0, \nabla \varphi(s_0))$  passes the  $\varepsilon$ -neighborhood of  $\mathcal{O}_2(\mathcal{O}_1)$ .

Case 3.  $n(X_j^0) \cdot \Xi_j^0 \geq \delta_1$ ,  $j = 1, 2, \dots, K_0 - 1$  and

$$0 \leq n(X_{K_0}^0) \cdot \Xi_{K_0}^0 \leq \delta_4.$$

Case 1. Taking account of Lemmas 3.1 and 3.2 we can take  $\mathcal{U}_0$  a neighborhood of  $s_0$  such that

$$*\mathcal{X}(x, \nabla \varphi(x)) = K_0 \quad \text{and} \quad n(X_j) \cdot \Xi_j \geq \frac{1}{2} \delta_1, \quad j = 1, 2, \dots, K$$

hold for all  $x \in \mathcal{U}_0 \cap \Gamma_1$  and  $L_{K_0}(x, \nabla \varphi(x))$  does not pass the  $\varepsilon/2$  neighborhood of  $\mathcal{O}_2$ . Suppose that  $\text{supp } f \subset \mathcal{U}_0 \cap \Gamma_1$ . Applying Lemma 7.3 we have

$$w_0^{(N)} = e^{ik(\varphi-t)} \sum_{j=0}^N v_{0j}(x, t; k) k^{-j}$$

satisfying (7.15)~(7.19). Applying Lemma 7.4 to  $w_0^{(N)}$  we have  $w_1^{(N)}$  with the properties (7.23)~(7.27). Next apply to  $w_1^{(N)}$  we have  $w_2^{(N)}$ . Repeating this process, we obtain  $w_j^{(N)}$ ,  $j = 0, 1, 2, \dots, K_0$  such that for  $j \geq 1$

$$\begin{aligned}
|w_j^{(N)}|_m(\Omega_R \times \mathbf{R}) &\leq C_{j,m,N,R} \sum_{l=0}^{2N} |f|'_{m+2l} k^{-l+m} \\
\text{supp } w_j^{(N)} &\subset \bigcup_{(x,t) \in \text{supp } f} \mathcal{L}_j(x, t; \nabla \varphi(x)) \\
|\square w_j^{(N)}|_m(\Omega_R \times \mathbf{R}) &\leq C_{j,m,N,R} k^{-N+m} |f|'_{2N+2m}(\Gamma_1 \times \mathbf{R}) \\
|B(w_j^{(N)} + w_{j+1}^{(N)})|_m(\Gamma_{(j)} \times \mathbf{R}) &\leq C_{j,m,N,R} k^{-N+m} |f|'_{2N+2+m}(\Gamma_1 \times \mathbf{R}),
\end{aligned}$$

where we set

$$\Gamma_{(j)} = \{X_j(x, \nabla \varphi(x)); x \in \mathcal{U}_0\}.$$

Taking account of

$$\text{supp } w_{K_0}^{(N)} \cap (\Gamma_\epsilon \times \mathbf{R}) = \phi, \quad \epsilon = (1 + (-1)^{K_0+1})/2 + 1$$

we see immediately

$$w^{(N)} = \sum_{j=0}^{K_0} w_j^{(N)}|_{\omega_j \times \mathbf{R}}$$

is the desired one.

For the case 2, we construct  $w_j^{(N)}$ ,  $j=0, 1, \dots, K_0$  by the same way as case 1. And for  $w_{K_0}^{(N)}|_{\Gamma_\epsilon \times \mathbf{R}}$  using Lemma 7.2 we have  $w_{K_0+1}^{(N)}$  such that

$$|B(w_{K_0}^{(N)} + w_{K_0+1}^{(N)})|_m(\Gamma_\epsilon \times \mathbf{R}) \leq C_{K_0,m,N} k^{-N+m} \sum |f|'_{N'+m}(\Gamma_1 \times \mathbf{R}).$$

Then

$$w^{(N)} = \sum_{j=0}^{K_0} w_j^{(N)}|_{\omega_j \times \mathbf{R}}$$

satisfies (7.32) (7.35).

For the case 3, we may show by the same process.

For  $u$  of the form (7.3), first construct  $w_0^{(N)}$  according to the process of Lemma 7.1. Then concerning

$$w_0^{(N)}|_{\Gamma_{(1)} \times \mathbf{R}} = e^{ik(\psi-t)} \sum_{j=0}^N v_{0j} k^{-j},$$

$\psi = \varphi|_{\Gamma_{(1)}}$  satisfies the condition C. Then we may apply the consideration for  $u$  of the form (7.24). On the other hand from (7.6)

$$|v_{0j}|_m(\Gamma_2 \times \mathbf{R}) \leq C_{j,m} |f|'_{20j}(\Gamma_1 \times \mathbf{R}).$$

Combining this estimate and the results for  $u$  of the form (7.24) (7.32)~(7.35) follow immediately.

## 8. Decay of asymptotic solutions

In this section we will prove the following

**Proposition 8.1.** *Let  $u$  be an oscillatory data of the form (7.3) or (7.14). Suppose that*

$$(8.1) \quad \text{supp } u \subset \Gamma_1 \times (T, T+1).$$

Then for any  $N$  positive integer there exists a function  $z^{(N)} \in C^\infty(\bar{\Omega} \times \mathbf{R})$  with the following properties:

$$(8.2) \quad \text{supp } z^{(N)} \subset \bar{\Omega} \times (T, \infty)$$

$$(8.3) \quad |z^{(N)}|_m(\Omega_R, t) \leq C_{N,m,R,\varepsilon} e^{-(c_0 - \varepsilon)(t-T)} \times k \sum_{j=0}^N k^{-j} |f|'_{m+2j+40}(\Gamma_1 \times \mathbf{R})$$

$$(8.4) \quad |\square z^{(N)}|_m(\Omega_R, t) \leq C_{N,m,R,\varepsilon} e^{-(c_0 - \varepsilon)(t-T)} k^{-N+m} |f|'_{m+2N+40}(\Gamma_1 \times \mathbf{R})$$

$$(8.5) \quad |Bz^{(N)} - u|_m(\Gamma, t) \leq C_{N,m,\varepsilon} e^{-(c_0 - \varepsilon)(t-T)} k^{-N+m} |f|'_{m+2N+40}(\Gamma_1 \times \mathbf{R}),$$

where  $c_0$  is the constant in Theorem 2.2,  $\varepsilon$  is an arbitrary positive constant and  $C_{N,m,R,\varepsilon}$  depends on  $N, m, R, \varepsilon$  and  $\{|\psi|_m; m=1, 2, \dots\}$ .

First we note a simple lemma without proof.

**Lemma 8.1.** Let  $\alpha$  be a positive constant  $< 1$ . Suppose that  $a_j, j=0, 1, 2, \dots$  be a sequence of positive numbers verifying for some  $A > 0$

$$a_j \leq \alpha a_{j-1} + A\alpha^{j-1}, \quad j = 1, 2, \dots$$

Then for any  $\varepsilon > 0$  we have an estimate

$$a_j \leq (a_0 + AC_\varepsilon)(\alpha\varepsilon^j)^j, \quad j = 0, 1, \dots$$

where  $C_\varepsilon$  depends only on  $\varepsilon$ .

Let us fix  $\varepsilon_0 > 0$  and we will show (8.2)~(8.5) for this  $\varepsilon_0$ . Let us choose  $\delta > 0$  so that we have

$$\left| \frac{K(y)}{l(1+lK(y))} - \frac{K(a_j)}{l_0(1+l_0K(a_j))} \right| < \delta, \quad \forall y \in S_j(\delta), j = 1, 2,$$

where  $l = \max_{\substack{x \in S_1(\delta) \\ y \in S_2(\delta)}} |x - y|$ ,  $K(y) = \min(\kappa_j(y))$ ,  $\kappa_j(y), j=1, 2$  are the principal cur-

vatures of  $\Gamma$  at  $y$ . Fix  $\delta_2$  and  $\delta_3$  so that  $\delta > \delta_3 > \delta_2$  and Corollary of Lemma 3.3 holds. Hereafter  $K$  denotes the one of Lemma 3.3 determined by  $\delta_2$ . Let us set  $S = S(\delta_2)$  and  $\tilde{S} = S(\delta_3)$ . Let  $v_{ij}(x)$ ,  $i, j=1, 2$  be functions defined on  $\Gamma_i$  such that

$$v_{i1} = \begin{cases} 1 & x \in S_i(\delta_2) \\ 0 & x \notin S_i(\delta_3) \end{cases}$$

and  $v_{i1} + v_{i2} = 1$  on  $\Gamma_i$ ,  $i=1, 2$ .

First we construct  $z^{(N)}$  for  $u$  of the form (7.14) verifying

$$\text{supp } f \subset S_1(\delta_2) \times (T, T+1).$$

Let us set for  $j=1, 2, 3, \dots$

$$\begin{aligned}\Gamma_{(j)} &= \tilde{S} \cap \bigcup_{x \in S_1(\delta_2)} X_j(x, \nabla \varphi(x)) \\ \tilde{\omega}_j &= \{X_j(x, \nabla \varphi(x)) + l\Xi_j(x, \nabla \varphi(x)); l \geq 0 \\ &\quad x \in S_1(\delta_2) \text{ such that } X_j(x, \nabla \varphi(x)) \in \tilde{S}\}.\end{aligned}$$

Then using Proposition 4.2 we have  $\varphi_j(x)$  defined in  $\tilde{\omega}_j$  such that

$$(8.6) \quad |\nabla \varphi_j| = 1 \quad \text{in } \tilde{\omega}_j$$

and

$$(8.7) \quad \begin{cases} \varphi_j = \varphi_{j-1} & \text{on } \Gamma_{(j)} \\ \frac{\partial \varphi_j}{\partial n} = -\frac{\partial \varphi_{j-1}}{\partial n} > 0 & \text{on } \Gamma_{(j)}. \end{cases}$$

Applying Proposition 5.4 we have

$$(8.8) \quad |\nabla \varphi_j|_m(\tilde{\omega}_j) \leq C_m |\nabla \varphi|_m(\tilde{\omega}_0) \leq \tilde{C}_m |\psi|'_{m+1}(\Gamma_1).$$

Let

$$w_0^{(N)} = e^{ik(\varphi-t)} \sum_{j=0}^N v_{0j}(x, t; k) k^{-j}$$

be the function constructed for  $u$  according to the process of Lemma 7.3. Set

$$u_{1p} = v_{2p} B w_0^{(N)}|_{\Gamma_2 \times R}, \quad p = 1, 2.$$

In the case of the Dirichlet boundary condition

$$u_{1p} = e^{ik(\varphi-t)} \sum_{j=0}^N v_{2p} v_{0j} k^{-j}.$$

In the case of the third kind boundary condition

$$u_{1p} = e^{ik(\varphi-t)} \sum_{j=0}^N v_{2p} \left( i \frac{\partial \varphi}{\partial n} v_{0j} + B v_{0j-1} \right) k^{-j+1}.$$

For  $u_{11}$  we can apply Lemma 7.3 and we have

$$w^{(N)}(x, t; k) = e^{ik(\varphi_1-t)} \sum_{j=0}^N v_{1j}(x, t; k) k^{-j}.$$

Set

$$u_{2p}(x, t; k) = v_{1p}(x) B w_1^{(N)}|_{\Gamma_1 \times R}, \quad p = 1, 2,$$

and construct for  $u_{21}$

$$w_2^{(N)}(x, t; k) = e^{ik(\varphi_2-t)} \sum_{j=0}^N v_{2j}(x, t; k) k^{-j}$$

following the process of Lemma 7.3. Repeating this process construct for  $q=0, 1, 2, \dots$ .

$$w_q^{(N)}(x, t; k) = e^{ik(\varphi_q(x)-t)} \sum_{j=0}^N v_{qj}(x, t; k) k^{-j}.$$

**Lemma 8.2.** *Let us set*

$$a = \max_{y \in S} \frac{1}{1+2LK(y)}, \quad l = \min_{\substack{x \in S_1(\delta_2) \\ y \in S_2(\delta_2)}} |x-y|.$$

*Then it holds that for  $q=0, 1, 2, \dots$*

$$(8.9) \quad |v_{qj}|_m(\Omega_R \times \mathbf{R}) \leq C_{j,R,m,\epsilon} (ae^\epsilon)^q |f|'_{m+2j}(\Gamma_1 \times \mathbf{R}).$$

*Proof.* First we show (8.9) for  $j=0$  and  $m=0$ . Note that  $v_{q0}$  satisfy

$$2 \frac{\partial v_{q0}}{\partial t} + 2 \nabla \varphi_q \cdot \nabla v_{q0} + \nabla \varphi_q \cdot v_{q0} = 0 \quad \text{in } \tilde{\omega}_q$$

and for  $q \geq 1$  they verify the boundary condition on  $\Gamma_{(q)}$

$$v_{q0} = \begin{cases} v_{\epsilon 1} v_{q-1,0} & \text{if } B = B_1 \\ -v_{\epsilon 1} v_{q-1,0} & \text{if } B = B_2, \end{cases}$$

where  $\epsilon = ((-1)^q + 1)/2$ . Consider the case of  $B=B_1$ . Then

$$v_{00} = f \quad \text{on } \Gamma_1.$$

Then applying (6.9) we have

$$|v_{00}|_0(\Gamma_2 \times \mathbf{R}) \leq |f|'_0(\Gamma_1 \times \mathbf{R}).$$

Note that from Lemma 4.1 we see the principal curvatures of  $\mathcal{C}_{\varphi_j}(X_j) \geq 2K(X_j)$ . From the definition we have for  $x \in \Gamma_2$

$$v_{10}(x, t) = v_{21}(x) v_{00}(x, t).$$

Then

$$|v_{10}|_0(\Gamma_2 \times \mathbf{R}) \leq |f|'_0(\Gamma_1 \times \mathbf{R}).$$

Applying (6.9) for  $v_{20}$  we have

$$|v_{20}|_0(\Gamma_1 \times \mathbf{R}) \leq a |v_{10}|_0(\Gamma_2 \times \mathbf{R}) \leq a |f|'_0(\Gamma_1 \times \mathbf{R}).$$

Repeating this process we have for all  $q$

$$|v_{q0}|_0(\Gamma_\epsilon \times \mathbf{R}) \leq a^{q-1} |f|'_0(\Gamma_1 \times \mathbf{R}).$$

Next, suppose that (8.9) holds for  $m$  when  $j=0$ . Note that

$$|v_{00}|_{m+1}(\Gamma_1 \times \mathbf{R}) \leq C |f|'_{m+1}(\Gamma_1 \times \mathbf{R}).$$

Then applying Lemma 6.1

$$|v_{21}v_{00}|_{m+1}(\Gamma_2 \times \mathbf{R}) \leq C |f|'_{m+1}(\Gamma_1 \times \mathbf{R}),$$

from which it follows that

$$|v_{21}v_{00}|_{m+1}(\Gamma_2 \times \mathbf{R}) \leq C \{ |f|'_{m+1}(\Gamma_1 \times \mathbf{R}) + |v_{00}|_m(\Gamma_1 \times \mathbf{R}) \}.$$

Since from the definition

$$v_{q0} = v_{\epsilon(q)1} v_{q-1,0} \quad \text{on } \Gamma_{(q)}$$

we have by applying Lemma 6.2

$$\begin{aligned} |v_{q0}|_{m+1}(\Gamma_{(q)} \times \mathbf{R}) &\leq |v_{q-1,0}|_{m+1}(\Gamma_{(q)} \times \mathbf{R}) \\ &\quad + C \{ |v_{q0}|_m(\Gamma_{(q)} \times \mathbf{R}) + |v_{q-1,0}|_m(\Gamma_{(q)} \times \mathbf{R}) \}. \end{aligned}$$

And from Lemma 6.1

$$\begin{aligned} |v_{q0}|_{m+1}(\Gamma_{(q+1)} \times \mathbf{R}) &\leq a |v_{q0}|_{m+1}(\Gamma_{(q)} \times \mathbf{R}) \\ &\quad + C \{ |v_{q0}|_m(\Gamma_{(q+1)} \times \mathbf{R}) + |v_{q0}|_m(\Gamma_{(q)} \times \mathbf{R}) \}. \end{aligned}$$

Combining these estimates and using the assumption

$$|v_{q0}|_{m+1}(\Gamma_{(q+1)} \times \mathbf{R}) \leq a |v_{q-1,0}|_{m+1}(\Gamma_{(q)} \times \mathbf{R}) + 4C \cdot C_{\epsilon}(ae^{\epsilon})^{q-1}.$$

Applying Lemma 8.1 we have

$$|v_{q0}|_{m+1}(\Gamma_{(q+1)} \times \mathbf{R}) \leq C_{m,\epsilon}(ae^{\epsilon})^q \quad \text{for all } q.$$

Taking account of (6.8) the estimate (8.9) for  $j=0$  follows from the above estimate.

Next we will show (8.9) for  $j \geq 1$ . Note that  $v_{qj}$ ,  $q=0, 1, 2, \dots$  satisfy

$$2 \frac{\partial v_{qj}}{\partial t} + 2 \nabla \varphi_q \cdot \nabla v_{qj} + \Delta \varphi_q \cdot v_{qj} = i \square v_{qj-1} \quad \text{in } \tilde{\omega}_q$$

and on  $\Gamma_{(q)}$

$$v_{qj} = \begin{cases} v_{\epsilon 1} v_{gj-1} & \text{if } B = B_1 \\ -v_{\epsilon 1} v_{qj-1} + (Bv_{qj-1} - v_{\epsilon 1} Bv_{q-1,j-1}) \left( i \frac{\partial \varphi_q}{\partial n} \right) & \text{if } B = B_2. \end{cases}$$

Suppose that (8.9) holds for  $j$  and  $m=0, 1, \dots$ . From the assumption

$$|\square v_{qj}|_m(\Omega_R^{(\epsilon)} \times \mathbf{R}) \leq C_{j,m,R}(ae^{\epsilon})^q |f|'_{m+2j+2}(\Gamma_1 \times \mathbf{R})$$

and

$$|Bv_{qj} - Bv_{q-1,j}|_m(\Gamma_{(q)} \times \mathbf{R}) \leq C_{j,m}(ae^{\epsilon})^q |f|'_{m+2j+1}(\Gamma_1 \times \mathbf{R}).$$

Then we have from Lemma 6.1

$$|v_{q,j+1}|_m(\Gamma_{(q+1)} \times \mathbf{R}) \leq a |v_{q,j+1}|_m(\Gamma_{(q)} \times \mathbf{R}) \\ + C(ae^\varepsilon)^q |f|'_{m+2j+2}(\Gamma_1 \times \mathbf{R}).$$

And applying Lemma 6.2 to  $v_{q+1,j+1}$  and  $v_{q,j+1}$  on  $\Gamma_{(q+1)}$  we have

$$|v_{q+1,j+1}|_m(\Gamma_{(q+1)} \times \mathbf{R}) \leq |v_{q,j+1}|_m(\Gamma_{(q+1)} \times \mathbf{R}) + C(ae^\varepsilon)^q |f|'_{m+2j+2}(\Gamma_1 \times \mathbf{R}).$$

Then we have (8.9) for  $j+1$  by combining the above two estimates and by using Lemma 8.1. By the induction we have (8.9) for all  $j$  and  $m$ . Q.E.D.

Since

$$w_q^{(N)} = e^{ik(\varphi_q - t)} \sum_{i=0}^N v_{q,i} k^{-i} \quad \text{and} \\ \square w_q^{(N)} = e^{ik(\varphi_q - t)} k^{-N} \square v_{qN},$$

we have

$$(8.10) \quad |w_q^{(N)}|_m(\Omega_R^{(\varepsilon)} \times \mathbf{R}) \leq C_{N,m,R,\varepsilon} (ae^\varepsilon)^q \sum_{j=0}^N |f|'_{m+2j+2}(\Gamma_1 \times \mathbf{R}) k^{-j+m}$$

$$(8.11) \quad |\square w_q^{(N)}|_m(\Omega_R^{(\varepsilon)} \times \mathbf{R}) \leq C_{N,m,R,\varepsilon} (ae^\varepsilon)^q |f|'_{m+2N+2}(\Gamma_1 \times \mathbf{R}) k^{-N+m}.$$

Taking account of

$$\text{supp } v_{qj} \subset \bigcup_{(x,t) \in \text{supp } f} \mathcal{L}_q(x, t; \nabla \varphi(x))$$

we have

$$\text{supp } w_q^{(N)} \cap (\Omega_R \times \mathbf{R}) \subset \Omega_R \times [ql_0, (q+1)l_1 + R],$$

therefore we can rewrite (8.10) and (8.11) as

$$(8.12) \quad |w_q^{(N)}|_m(\Omega_R^{(\varepsilon)} \times \mathbf{R}) \leq C_{N,m,R,\varepsilon} e^{-(c_0 - \varepsilon)t} \sum_{j=0}^N |f|'_{m+2j+2}(\Gamma_1 \times \mathbf{R}) k^{-j+m}$$

$$(8.13) \quad |\square w_q^{(N)}|_m(\Omega_R^{(\varepsilon)} \times \mathbf{R}) \leq C_{N,m,R,\varepsilon} e^{-(c_0 - \varepsilon)t} k^{-N+m} |f|'_{m+2N+2}(\Gamma_1 \times \mathbf{R}).$$

Concerning the estimate on the boundary, we see that from the process of the construction of  $w_q^{(N)}$

$$(8.14) \quad Bw_0^{(N)}|_{\Gamma_1 \times \mathbf{R}} = f + e^{ik(\varphi - t)} k^{-N} Bv_{0N}$$

and for  $q \geq 1$

$$(8.15) \quad [Bw_{q+1}^{(N)} + v_{\varepsilon 1} Bw_q^{(N)}]|_{\Gamma \times \mathbf{R}} = e^{ik(\varphi_q - t)} k^{-N} (Bv_{q+1,N} + v_{\varepsilon 1} Bv_{qN}).$$

Remark that for  $q \geq 1$  it holds that

$$^* \mathcal{X}(x, \nabla \varphi_q(x)) \leq K, \quad \forall x \in \text{Proj supp } u_{q2}.$$

Indeed,  $x \in \text{Proj supp } u_{q2}$  is represented as

$$x = X_1(\tilde{x}, \nabla \varphi_{q-1}(\tilde{x})), \quad \tilde{x} \in \text{Proj supp } u_{q-1,1}$$

and  $x \notin S$ . Then Lemma 3.3 and its corollary assure our assertion.

Now we can apply Proposition 7.5 to  $u_{q2}$ ,  $q \geq 1$  taking  $J=K$ . Then we have  $y_q^{(N)}(x, t; k)$  such that

$$(8.16) \quad \begin{aligned} |y_q^{(N)} + u_{q2}|'_m(\Gamma \times \mathbf{R}) &\leq C k^{-N+m} |f_{q2}|'_{m+2N+40}(\Gamma_\epsilon \times \mathbf{R}) \\ |y_q^{(N)}|_m(\Omega_R \times \mathbf{R}) &\leq C k^{m+1} \sum_{j=0}^N |f_{q2}|'_{m+2j+4}(\Gamma_\epsilon \times \mathbf{R}) k^{-j} \\ |\square y_q^{(N)}|_m(\Omega_R \times \mathbf{R}) &\leq C k^{-N+m} |f_{q2}|'_{m+2N+40}(\Gamma_\epsilon \times \mathbf{R}) \\ \text{supp } y_q^{(N)} &\subset \bigcup_{(x,t) \in \text{supp } f_{q2}} \mathcal{L}(x, t; \nabla \varphi_{q+1}(x)), \end{aligned}$$

where we set  $u_{q2} = \iota^{ik(\varphi_q - t)} f_{q2}$ . Then we have from the definition

$$f_{q2} = \begin{cases} v_{\epsilon 2} \sum_{j=1}^N v_{qj} k^{-j} & \text{if } B = B_1 \\ v_{\epsilon 2} \sum_{j=0}^N \left( ik \frac{\partial \varphi_q}{\partial n} v_{qj} + B v_{q,j-1} \right) k^{-j-1} & \text{if } B = B_2. \end{cases}$$

Using (8.9) it follows that

$$(8.17) \quad |f_{q2}|'_m(\Gamma_\epsilon \times \mathbf{R}) \leq C(ae^\varepsilon)^q \sum_{j=0}^N |f|'_{m+2j}(\Gamma_1 \times \mathbf{R}) k^{-j}.$$

Taking account of the support of  $u_{q2}$  and (8.16) we have

$$\text{supp } y_q^{(N)} \cap (\Omega_R \times \mathbf{R}) \subset \Omega_R \times [l_0 q, l_1(q+K)+R].$$

Then it follows that

$$(8.18) \quad |y_q^{(N)} + u_{q2}|'_m(\Gamma \times \mathbf{R}) \leq e^{-(c_0 - \varepsilon)t} k^{-N+m} \sum_{j=0}^N |f|'_{m+2N+40+2j}(\Gamma_1 \times \mathbf{R}) k^{-j}$$

$$(8.19) \quad |y_q^{(N)}|_m(\Omega_R \times \mathbf{R}) \leq C k^{m+1} e^{-(c_0 - \varepsilon)t} \sum_{j=0}^{2N} |f|'_{m+40+2j}(\Gamma_1 \times \mathbf{R}) k^{-j}$$

$$(8.20) \quad |\square y_q^{(N)}|_m(\Omega_R \times \mathbf{R}) \leq C e^{-(c_0 - \varepsilon)t} k^{-N+m} \sum_{j=0}^N |f|'_{m+40+2(N+j)} k^{-j}$$

Then we see immediately that

$$z^{(N)}(x, t; k) = \sum_{q=0}^{\infty} w_q^{(N)}(x, t; k) + y_q^{(N)}(x, t; k)$$

is well defined and it verifies the properties of Proposition because for each  $t$  only  $w_q^{(N)}$  such that  $tl_0 \geq q \geq t - R/l_1 - K$  do not vanish.

Now we will remove the assumption

$$\text{supp } f \subset S_1(\delta_2) \times (T, T+1).$$

Let  $s_0 \in \text{Proj supp } f$ . If

$$*\mathcal{X}(s_0, \nabla \varphi(s_0)) \leq K,$$

Proposition 7.5 proves Proposition 8.1 if  $\text{supp } f$  contained in a small neigh-

borhood of  $s_0$ . If  $\# \mathcal{X}(s_0, \nabla \varphi(s_0)) \geq K$ , Lemma 3.3 assures for some  $j \leq K$

$$X_j(s_0, \nabla \varphi(s_0)) \in S.$$

Then we may suppose that

$$X_j(s, \nabla \varphi(s)) \in S, \quad \forall s \in \text{Proj supp } f.$$

Using the process of Proposition 7.5 we construct  $w_0^{(N)}, w_1^{(N)}, \dots, w_j^{(N)}$ , then we have

$$\begin{aligned} Bw_j^{(N)}|_{\Gamma_{\epsilon(j)} \times \mathbf{R}} &= e^{ik(\varphi_j - t)} f_j \\ \text{supp } f_j &\subset S_{\epsilon(j)}(\delta_2) \times \mathbf{R} \end{aligned}$$

and the estimate

$$|f_j|'_m \leq C \sum_{l=1}^N |f|'_{2l+40+m}(\Gamma_1 \times \mathbf{R}) k^{-l}$$

holds. Thus we may apply the process of the construction of  $z^{(N)}$  to  $e^{ik(\varphi_j - t)} f_j$  and we have the desired function.

**Corollary.** *For an oscillatory boundary data of Proposition there exists  $z^{(N)}(x, t; k)$  satisfying*

$$\square z^{(N)} = 0 \quad \text{in } \Omega \times \mathbf{R}$$

and (8.2), (8.3) and (8.5).

Proof. Let  $z^{(N)}$  be a function verifying (8.2)~(8.5). Take a function  $h(x, t; k)$  such that

$$\begin{aligned} h(x, t; k) &= -\square z^{(N)} \quad \text{in } \Omega \times \mathbf{R} \\ |h|_m(\mathcal{O}, t) &\leq C_{N, m, R, \epsilon} e^{-(\epsilon_0 - \epsilon)(t - T)} k^{-N+m} |f|'_{2N+m+40}(\Gamma_1 \times \mathbf{R}). \end{aligned}$$

Let  $g(x, t; k)$  be the solution of

$$\begin{cases} \square g = h & \text{in } \mathbf{R}^3 \times \mathbf{R} \\ g(x, t; k) = 0 & \text{for } t < T. \end{cases}$$

And we have the estimate

$$|g|_m(x, t) \leq \int_{t-2R}^t |h|_m(|x| \leq R, s) ds.$$

Then we see immediately that

$$z^{(N)} = z^{(N)} + g$$

satisfies the required properties.

### 9. Proof of Theorem 2.2

Let  $s_0 \in \Gamma_0$  and let  $\mathcal{U}$  be a small neighborhood in  $\mathbf{R}^3$  of  $s_0$  such that for all

$$(9.1) \quad h(x, t) \in C_0^\infty((\Gamma_1 \cap \mathcal{U}) \times \mathbf{R})$$

the representation

$$h(x, t) = \sum_{j=1}^4 (\mathcal{W}_j h)(x, t)$$

of § 4 of [3] is applicable. We will define  $\mathcal{W}_j$  an operator from  $C_0^\infty((\Gamma_1 \cap \mathcal{U}) \times \mathbf{R})$  into  $C^\infty(\bar{\Omega} \times \mathbf{R})$ . We set

$$(\mathcal{W}_1^{(N)} h)(x, t) = \int dk \int_{-1}^{\infty} d\alpha \int_{\Sigma} d\xi' \\ \mathfrak{z}^{(N)}(x, t; \alpha, \xi', k) \chi_j(1 + \alpha)^2 k^2 (1 + \alpha) \hat{h}(\xi, k),$$

where

$$\hat{h}(\xi, k) = \int d\sigma' \int dt' e^{i(\xi \cdot \sigma' - t' k)} h(s(\sigma'), t'),$$

and  $\mathfrak{z}^{(N)}(x, t; \alpha, \xi', k)$  is the function of Corollary of Proposition 8.1 constructed for an oscillatory data

$$u(s(\sigma), t; k) = e^{ik((1+\alpha)\langle \sigma, \xi' \rangle - t)} \omega(s(\sigma), t).$$

Indeed,  $\psi(s(\sigma)) = (1 + \alpha)\langle \sigma, \xi' \rangle$  satisfies the condition  $C$  for  $\alpha \leq -\alpha_0$ . And we set

$$(\mathcal{W}_2^{(N)} h)(x, t) = \int dk \int_{\Sigma} d\eta \int_{|\beta| \leq \beta_0} d\beta \int d\sigma' \int dt' \mathfrak{z}^{(N)}(x, t; \beta, \eta, k) \\ \cdot k^2 e^{ik(\theta(s(\sigma'), \eta, \beta) - t')} \tilde{\omega}(s(\sigma'), t') h(s(\sigma'), t')$$

where  $\mathfrak{z}^{(N)}$  is the function of Corollary of Proposition 8.1 constructed for an oscillatory boundary data

$$u(s(\sigma), t') = e^{ik(\theta(s(\sigma), \eta, \beta) - t')} \omega(s(\sigma), t) \chi_2(1 + \alpha(\sigma, \sigma', \eta, \beta))^2 \\ \cdot (1 + \alpha(\sigma, \sigma', \eta, \beta)) \frac{D(\xi', \alpha)}{D(\eta, \beta)}(\sigma, \sigma', \eta, \beta).$$

And we set  $\mathcal{W}_j + \mathcal{Z}_j$ ,  $j=3, 4$  of § 4 of [3] as  $\mathcal{W}_j^{(N)}$ ,  $j=3, 4$  respectively. Remark that  $\mathcal{W}_j + \mathcal{Z}_j$  of [3] satisfies

$$\square(\mathcal{W}_j + \mathcal{Z}_j) = 0 \quad \text{in } \Omega^{(1)}$$

and for any  $N'$

$$|(\mathcal{W}_j + \mathcal{Z}_j)h|_m(\Gamma_2 \times \mathbf{R}) \leq C_{m, N'} |k|^{-N'} |h|_0(\Gamma_1 \times \mathbf{R}) \\ (\mathcal{W}_j + \mathcal{Z}_j)h = 0 \quad \text{for } |x| \geq t - 2l_1.$$

Then we have for  $j=1, 3$

$$|\mathcal{W}_j^{(N)}h|_m(\Omega_R, t) \leq C_{m,R,\varepsilon} e^{-(c_0-\varepsilon)(t-1)} \\ \cdot \int dk \int d\alpha \int d\xi' \left( \sum_{j=0}^{N'} k^{-j} A_{m+2j+40} \right) |k|^{m+1} \chi_j(1+\alpha)^2 k^2 (1+\alpha) |\hat{h}(\xi, k)|,$$

where  $A_j$  depends on the derivatives of  $\omega(x, t)$ , and

$$|\mathcal{W}_2^{(N)}h|_m(\Omega_R, t) \leq C_{m,R,\varepsilon} e^{-(c_0-\varepsilon)(t-1)} \\ \cdot \int dk \int d\eta \int d\beta \left( \sum_{j=0}^{N'} |k|^{-j} A_{m+2j+40} \right) k^{m+1} \chi_2(1+\alpha)^2 \\ \left( \int |\tilde{h}(s(\sigma'), k)| d\sigma' \right),$$

where  $\tilde{h}(s(\sigma), k) = \int e^{-ikt'} h(s(\sigma), t') dt'$ . Simillary it holds that

$$|\mathcal{W}_4 h|_m(\Omega_R, t) \leq C_{m,R,\varepsilon} e^{-(c_0-\varepsilon)(t-1)} \\ \cdot \int dk \int d\alpha \int d\xi' |\xi|^m \chi_4(1+\alpha)^2 k^2 (1+\alpha) \hat{h}(\xi, k).$$

By the same way we have, for  $j=1, 3$

$$|\mathcal{W}_j^{(N)}h - \mathcal{V}_j h|'_m(\Gamma, t) \leq C_{N,m} e^{-(c_0-\varepsilon)(t-1)} \int dk \int d\alpha \int d\xi' \\ \cdot A_{m+2N+40} |k|^{-N+m+1} \chi_j(1+\alpha)^2 k^2 (1+\alpha) |\hat{h}(\xi, k)|, \\ |\mathcal{W}_2^{(N)}h - \mathcal{V}_2 h|_m(\Gamma, t) \leq C_{N,m} e^{-(c_0-\varepsilon)(t-1)} \int dk \int d\beta \int d\eta \\ \cdot A_{m+2N+40} k^{m+1} \chi_2(1+\alpha)^2 \left( \int |\tilde{h}(s(\sigma'), k)| d\sigma' \right),$$

and

$$|\mathcal{W}_4^{(N)}h - \mathcal{V}_4 h|'_m(\Gamma, t) \leq C_{N,m,\varepsilon} e^{-(c_0-\varepsilon)(t-1)} \int dk \int d\alpha \int d\xi' \\ \cdot |k|^{-N} A_{m+2N+40} |\xi|^m \chi_4(1+\alpha)^2 (1+\alpha) k^2 |\hat{h}(\xi, k)|.$$

Set

$$\mathcal{W}^{(N)}h = \sum_{j=1}^4 \mathcal{W}_j^{(N)}h.$$

Then we have

$$(9.2) \quad \mathcal{W}^{(N)}h = 0 \quad \text{for } t < 0$$

$$(9.3) \quad |\mathcal{W}^{(N)}h|_m(\Omega_R, t) \leq C_{N,m,R,\varepsilon} e^{-(c_0-\varepsilon)(t-1)} \\ \cdot \int \{ (1+|k|)^{m+3} \int |\tilde{h}(s(\sigma), k)| d\sigma + \int |\xi|^m |\hat{h}(\xi, k)| d\xi \} dk$$

$$(9.4) \quad \square \mathcal{W}^{(N)}h = 0 \quad \text{in } \Omega \times \mathbf{R}$$

$$(9.5) \quad |D_t(B\mathcal{W}^{(N)}h-h)|'_m(\Gamma, t) \leq C_{N,m,R,\varepsilon} e^{-(c_0-\varepsilon)(t-1)} \|h\|_{-N+m+2, L^2(\Gamma \times \mathbf{R})}.$$

For  $p = ik + \mu$ ,  $\mu > -c_0$  define  $\mathcal{Z}^{(N)}(p)h$  by

$$(\mathcal{Z}^{(N)}(p)h)(x) = \int_{-\infty}^{\infty} e^{-pt} (\mathcal{W}^{(N)}h)(x, t) dt.$$

From (9.4) it follows that

$$(9.6) \quad (p^2 - \Delta) \mathcal{Z}^{(N)}(p)h = 0 \quad \text{in } \Omega.$$

Taking account of (9.2) and (9.3) we have for  $\operatorname{Re} p > 0$

$$\mathcal{Z}^{(N)}(p)h \in \bigcap_{m \geq 0} H^m(\Omega)$$

and

$$(9.7) \quad \|\mathcal{Z}^{(N)}(p)h\|_{m, L^2(\Omega)} \leq C_m (\operatorname{Re} p)^{-1} \|h\|_{m, L^2(\Gamma \times \mathbf{R})}$$

From (9.3) we have for  $\operatorname{Re} p \geq -(c_0 - \varepsilon)$

$$(9.8) \quad \|\mathcal{Z}^{(N)}(p)h\|_m(\Omega_R) \leq C_{N,m,R} \int_{-\infty}^{\infty} \{|k|^{m+3} \|\tilde{h}(\cdot, k)\|_{L^2(\Gamma)} + \|\tilde{h}(\cdot, k)\|_{m+3, L^2(\Gamma)}\} dk,$$

and also from (9.5) we have for  $\operatorname{Re} p \geq -(c_0 - \varepsilon)$

$$(9.9) \quad |p \{B(\mathcal{Z}^{(N)}(p)h - \tilde{h}(\cdot, p))\}|'_m(\Gamma) \leq C_{N,m,\varepsilon} \|h\|_{-N+m+1, L^2(\Gamma \times \mathbf{R})}$$

We see immediately that by using the partition of the unity we can define  $\mathcal{W}^{(N)}$  for all  $h \in C_0^\infty(\Gamma \times (0, 1))$  and (9.8) and (9.9) hold.

Define  $U_q^{(N)}(p)g$  an operator from  $C^\infty(\Gamma)$  into  $C^\infty(\bar{\Omega})$  for  $p$  and  $q(t)$  such that  $\operatorname{Re} p > -c_0$ ,  $\hat{q}(p) \neq 0$  by

$$(9.10) \quad \tilde{U}_q^{(N)}(p)g = \frac{1}{\hat{q}(p)} (\mathcal{Z}^{(N)}(p)h),$$

where

$$h(x, t) = g(x)q(t), \quad g \in C^\infty(\Gamma), \quad q \in C_0^\infty(0, 1).$$

Let  $m(t)$  be a function of  $C_0^\infty(0, 1)$  verifying

$$\int m(t) dt = 1.$$

Set

$$q_k(t) = e^{-ikt} m(t).$$

Since  $\hat{q}_k(ik' + \mu) = \hat{m}(\mu + i(k' - k))$  there exist  $a_0 > 0$  and  $C_0 > 0$  such that for all  $|k' - k| \leq a_0$ ,  $1 \geq \mu \geq -c_0$

$$(9.11) \quad |\hat{q}_k(ik' + \mu)| \geq C_0.$$

From the definition we have

$$(9.12) \quad \int |k'^m q_k(ik' + \mu)| dk' \leq C_m |k|^m, \quad \forall k \in \mathbf{R}$$

where  $C_m$  depends only on  $m(t)$ . Then by using (9.11) and (9.12) we have from (9.8) for  $1 \geq \operatorname{Re} p \geq -(c_0 - \varepsilon)$ ,  $|k' - k| \leq a_0$

$$(9.13) \quad |\tilde{U}_{q_k}^{(N)}(\mu + ik')g|_m(\Omega_R) \leq C_{N,m,R,\varepsilon} \{ |k|^{m+3} \|g\|_{L^2(\Gamma)} + \|g\|_{m+3, L^2(\Gamma)} \}$$

and from (9.9)

$$(9.14) \quad |\mu + ik'| |B\tilde{U}_{q_k}^{(N)}(\mu + ik')g - g|_m(\Gamma) \leq C_{N,m,\varepsilon} \|g\|_{L^2(\Gamma)}.$$

Then, when  $C_{N,m,\varepsilon}/|k| \leq 1/2$ , for  $p = ik' + \mu$  such that  $1 \geq \mu \geq (c_0 - \varepsilon)$  and  $|k' - k| \leq a_0$

$$A_k^{(N)}(p) = I + \sum_{j=1}^{\infty} (I - B\tilde{U}_{q_k}^{(N)}(p))^j$$

is well defined as a mapping from  $C^m(\Gamma)$  into  $C^m(\Gamma)$  and for all  $g \in C^m(\Gamma)$  we have

$$(9.15) \quad B\tilde{U}_{q_k}^{(N)}A_k^{(N)}(p)g = g.$$

Define  $U_k^{(N)}(p)g = \tilde{U}_{q_k}^{(N)}(p)A_k^{(N)}(p)g$ . Then it holds that

$$(9.16) \quad \begin{cases} (p^2 - \Delta)U_k^{(N)}(p)g = 0 & \text{in } \Omega \\ BU_k^{(N)}(p)g = g & \text{on } \Gamma, \end{cases}$$

$$(9.17) \quad |U_k^{(N)}(p)g|_m(\Omega_R) \leq 2C_{N,m,R,\varepsilon} \{ |k|^{m+3} \|g\|_{L^2(\Gamma)} + \|g\|_{m+3, L^2(\Gamma)} \}.$$

For  $1 \geq \operatorname{Re} p > 0$  we have from (9.7)

$$(9.18) \quad \|U_k^{(N)}(p)g\|_{m, L^2(\Omega)} \leq C_m (\operatorname{Re} p)^{-1} \{ |p|^m \|g\|_{L^2(\Gamma)} + \|g\|_{m, L^2(\Gamma)} \}.$$

For each  $k$  verifying  $C_{N,m,\varepsilon}/|k| < 1/2$ ,  $U_k^{(N)}(p)$  is analytic in  $1 \geq \mu \geq -(c_0 - \varepsilon)$ ,  $|k' - k| \leq a_0$ . Since in  $\operatorname{Re} p > 0$  the solution of

$$\begin{cases} (p^2 - \Delta)u(x) = 0 & \text{in } \Omega \\ Bu = g & \text{on } \Gamma \end{cases}$$

is unique in  $L^2(\Omega)$ , we see that  $U_k^{(N)}(p)$  is independent of  $k$  and  $N$  in  $\operatorname{Re} p > 0$ . Thus  $U_k^{(N)}(p)g$  is the analytic continuation of  $U(p)g$ . Then we have

**Proposition 9.1.** *For any  $\varepsilon > 0$ , there exists  $k_\varepsilon > 0$  such that for any  $g \in C^m(\Gamma)$ ,  $m \geq 3$ ,  $U(p)$  can be prolonged analytically into a region*

$$\{\mu + ik; \mu \geq -(c_0 - \varepsilon), |k| \geq k_\varepsilon\}$$

and the estimate

$$|U(p)g|_m(\Omega_R) \leq C_{m,R,\varepsilon} \{ \|k\|^{m+3} \|g\|_{L^2(\Gamma)} + \|g\|_{m+3, L^2(\Gamma)} \}$$

holds.

Proof. First set  $m=0$ . Then we can choose  $k_\varepsilon$  as the proposition holds. By using the regularity theorem for  $\Delta$  we get the estimate for all  $m \geq 1$  if  $p$  belongs to this region.

## Appendix

Let  $\mathcal{O}_1$  be a convex and bounded obstacle with smooth boundary  $\Gamma_1$  such that

$$\begin{aligned} \Gamma_1 &\supset \{(x, y, z); x = 0, y^2 + z^2 \leq 1\} \\ \mathcal{O}_1 &\subset \{(x, y, z); x \leq 0\} \end{aligned}$$

and  $\mathcal{O}_2$  be a convex and bounded obstacle with smooth boundary  $\Gamma_2$  such that

$$\begin{aligned} \Gamma_2 &\supset \{(x, y, z); x = 1, y^2 + z^2 \leq 1\} \\ \mathcal{O}_2 &\subset \{(x, y, z); x \geq 1\}. \end{aligned}$$

For these  $\mathcal{O}_j$  we consider the problem with the Dirichlet boundary condition. Set

$$p_{m,R}(t) = \sup_{\{u_0, u_1\} \in C_0^\infty(\mathcal{Q}_R)} E(u, R, t)^{1/2} (\|u_0\|_{m+1, L^2(\Omega)} + \|u_1\|_{m, L^2(\Omega)})^{-1}.$$

We like to show

**Theorem A.** *Let  $R \geq 2$ . For every positive integer  $m$  it holds that*

$$\overline{\lim}_{t \rightarrow \infty} p_{m,R}(t) t^{m(1+\varepsilon)} > 0$$

for any positive number  $\varepsilon$ .

To prove this theorem we consider the behavior of an asymptotic solution  $U^{(N)}(x, y, z, t; k)$  of the problem

$$(A.1) \quad \begin{cases} \square U = 0 & \text{in } \Omega \times (0, \infty) \\ BU = 0 & \text{on } \Gamma \times (0, \infty). \end{cases}$$

First we construct an asymptotic solution of the problem

$$(A.2) \quad \square w = 0 \quad \text{in } \mathbf{R}^3 \times (0, \infty)$$

in the form

$$(A.3) \quad w = e^{ik(x-t)} \sum_{j=0}^{\infty} v_j(x, y, z, t) k^{-j}.$$

Then  $v_j$  must satisfy

$$(A.4)_0 \quad 2 \frac{\partial v_0}{\partial t} + 2 \frac{\partial v_0}{\partial x} = 0 \quad \text{in } \mathbf{R}^3 \times (0, \infty)$$

and for  $j \geq 1$

$$(A.4)_j \quad 2 \frac{\partial v_j}{\partial t} + 2 \frac{\partial v_j}{\partial x} = -i \square v_{j-1} \quad \text{in } \mathbf{R}^3 \times (0, \infty).$$

**Lemma A.1.** *Let  $f(x, y, z) \in C_0^\infty(\mathbf{R}^3)$ . If we require*

$$(A.5)_0 \quad v_0(x, y, z, 0) = f(x, y, z)$$

and for  $j \geq 1$

$$(A.5)_j \quad v_j(x, y, z, 0) = 0$$

$v_j, j=0, 1, 2, \dots$  the solutions of (A.4)<sub>j</sub> are determined successively and they are represented as

$$(A.6)_j \quad v_j(x, y, z, t) = \sum_{l=0}^j t^l (P_{jl} f)(x-t, y, z)$$

where  $P_{jl}$  are differential operators with constant coefficients of order  $\leq 2j$ .

**Proof.** Let  $v_0$  satisfy (A.4)<sub>0</sub> and (A.5)<sub>0</sub>. Then for any  $(\xi, y, z) \in \mathbf{R}^3$  and  $s \geq 0$

$$\frac{\partial}{\partial s} v_0(\xi+s, y, z, s) = \left( \frac{\partial v_0}{\partial t} + \frac{\partial v_0}{\partial x} \right)(\xi+s, y, z, s) = 0.$$

Then we have for all  $s \geq 0$

$$v_0(\xi+s, y, z, s) = v_0(\xi, y, z, 0) = f(\xi, y, z).$$

Then taking  $\xi = x-t, s=t$  it follows that

$$v_0(x, y, z, t) = f(x-t, y, z).$$

This is nothing but (A.6) for  $j=0$ . Suppose that (A.6) holds for  $j=h$ . Then we have

$$\begin{aligned} \square v_h(x, y, z, t) &= \sum_{l=0}^h t^l \left( \left( -\frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} \right) P_{hl} f \right)(x-t, y, z) \\ &\quad + \sum_{l=0}^h \left\{ l(l-1) t^{l-2} (P_{pl} f)(x-t, y, z) - 2l t^{l-1} \left( \frac{\partial}{\partial x} P_{hl} f \right)(x-t, y, z) \right\} \\ &= \sum_{l=0}^h t^l (\tilde{P}_{h+1,l} f)(x-t, y, z), \end{aligned}$$

where  $\tilde{P}_{h+1,l}$  is a differential operator of order  $\leq 2(h+1)$  with constant coefficients. Now from (A.4)<sub>h+1</sub> and (A.5)<sub>h+1</sub> it follows that

$$\begin{cases} 2 \frac{\partial}{\partial s} v_{h+1}(\xi+s, y, z, s) = \sum_{l=0}^h s^l (\tilde{P}_{h+1,l} f)(\xi, y, z) \\ v_{h+1}(\xi, y, z, 0) = 0, \end{cases}$$

from which we have

$$v_{h+1}(\xi+s, y, z, s) = \sum_{l=0}^h s^{l+1} (2(l+1))^{-1} (\tilde{P}_{h+1,l} f)(\xi, y, z).$$

Setting  $s=t$ ,  $\xi=x-t$  and  $P_{h+1,l}=(2(l+1))^{-1}\tilde{P}_{h+1,l}$  we see that (A.6) holds for  $j=h+1$ . Q.E.D.

Suppose that

$$(A.7) \quad \text{supp } f \subset \{(x, y, z); y^2 + z^2 \leq 1/16, 1/4 \leq x \leq 3/4\}$$

$$(A.8) \quad \int |f(x, y, z)|^2 dx dy dz = 1.$$

Let us define  $w^{(N)}$  by

$$w^{(N)}(x, y, z, t; k) = e^{ik(x-t)} \sum_{j=0}^N v_j(x, y, z, t) k^{-j}.$$

Then by using (A.6) we have from (A.7)

$$(A.9) \quad \text{supp } w^{(N)} \subset \{(x, y, z, t); y^2 + z^2 \leq 1/16, 1/4 \leq x-t \leq 3/4\}.$$

We have immediately

$$\begin{aligned} (A.10) \quad \square w^{(N)} &= e^{ik(x-t)} \square v_N \\ &= e^{ik(x-t)} k^{-N} \sum_{l=0}^N t^l (\tilde{P}_{N+1,l} f)(x-t, y, z). \end{aligned}$$

Set

$$\omega = \{(x, y, z); 0 \leq x \leq 1, y^2 + z^2 \leq 1\}.$$

Then for any  $(x, y, z, t) \in \omega \times \mathbf{R}$  (A.9) implies that the number of positive integer  $p$  such that

$$w^{(N)}(2p-x, y, z, t) - w^{(N)}(2p+x, y, z, t) \neq 0$$

is at most one. Now we can define a function  $\tilde{U}^{(N)}$  in  $\omega \times \mathbf{R}$  by

$$\begin{aligned} &\tilde{U}^{(N)}(x, y, z, t; k) \\ &= w^{(N)}(x, y, z, t; k) - \sum_{p=1}^{\infty} \{w^{(N)}(2p-x, y, z, t) - w^{(N)}(2p+x, y, z, t; k)\}. \end{aligned}$$

We see immediately

$$U^{(N)} = 0 \quad \text{on } x = 0 \text{ and } x = 1$$

$$\square U^{(N)} = g^{(N)}(x, y, z, t; k) - \sum_{p=1}^{\infty} \{g^{(N)}(2p-x, y, z, t) - g^{(N)}(2p+x, y, z, t)\}$$

where  $g^{(N)} = \square w^{(N)}$ . Since we have

$$\int_{\omega} |D^{\gamma} g^{(N)}(2p-x, y, z, t; k)|^2 dx dy dz \leq C_{N, \gamma} t^{N+|\gamma|} k^{-N+|\gamma|}$$

it holds that

$$(A.11) \quad \int_{\omega} |D^{\gamma} \square \tilde{U}^{(N)}(x, y, z, t; k)|^2 dx dy dz \leq C_{N, \gamma} t^{N+|\gamma|} k^{-N+|\gamma|}.$$

Let us set

$$U^{(N)}(x, y, z, t; k) = \begin{cases} \tilde{U}^{(N)}(x, y, z, t; k) & \text{if } (x, y, z) \in \omega \\ 0 & \text{if } (x, y, z) \in \Omega - \omega. \end{cases}$$

Evidently we see that  $U^{(N)} \in C^{\infty}(\Omega \times (0, \infty))$  and

$$\square U^{(N)} = \begin{cases} \square \tilde{U}^{(N)} & \text{in } \omega \times (0, \infty) \\ 0 & \text{in } (\Omega - \omega) \times (0, \infty). \end{cases}$$

Let  $V^{(N)}$  be the solution of

$$\begin{cases} \square V^{(N)} = -\square U^{(N)} & \text{in } \Omega \times (0, \infty) \\ V^{(N)} = 0 & \text{on } \Gamma \times (0, \infty) \\ V^{(N)}|_{t=0} = V_t^{(N)}|_{t=0} = 0. \end{cases}$$

Then we have from (A.11)

$$(A.12) \quad \|V^{(N)}(\cdot, t)\|_{l, L^2(\Omega)} \leq C_{N, l} t^{N+l+2} k^{-N-l} \quad t \geq 0.$$

Setting  $W^{(N)} = U^{(N)} + V^{(N)}$  we have

$$\begin{cases} \square W^{(N)} = 0 & \text{in } \Omega \times (0, \infty) \\ W^{(N)} = 0 & \text{on } \Gamma \times (0, \infty). \end{cases}$$

Taking account of the properties of  $U^{(N)}$  and  $V^{(N)}$  we have

$$\begin{aligned} W^{(N)}(x, y, z, 0; k) &= e^{ikx} f(x, y, z) \\ W_t^{(N)}(x, y, z, 0; k) &= -(ikf(x, y, z) + \frac{\partial f}{\partial x}(x, y, z))e^{ikx}. \end{aligned}$$

Then it follows that

$$(A.13) \quad \|W^{(N)}(\cdot, 0)\|_{m+1, L^2(\Omega)} + \|W_t^{(N)}(\cdot, 0)\|_{m, L^2(\Omega)} \leq C_m k^{m+1}.$$

From the form of  $U^{(N)}$  we have

$$U^{(N)}(x, y, z, 2n; k) \\ = e^{ik(x-2n)}(f(x, y, z) - \sum_{j=1}^N \sum_{l=0}^j (2n^l)(F_{jl}f)(x, y, z)k^{-j}).$$

Then we have

$$(A.14) \quad E(U^{(N)}, R, 2n)^{1/2} \geq k(1 - \sum_{j=1}^N C_{N,j} n^j k^{-j}).$$

Therefore it follows from (A.12) and (A.14) that

$$E(W^{(N)}, R, 2n)^{1/2} \geq E(U^{(N)}, R, 2n)^{1/2} - E(V^{(N)}, \infty, 2n)^{1/2} \\ \geq k(1 - \sum_{j=1}^N C_{N,j} n^j k^{-j}) - C_N(2n)^{N+1} k^{-N+1}.$$

Then we have

$$p_{m,R}(2n) \geq \max_{k,N} \{k(1 - \sum_{j=1}^N C_{N,j} n^j k^{-j}) - C_N(2n)^{N+1} k^{-N+1}\} (C_m k^{m+1})^{-1}.$$

Suppose that  $\varepsilon > 0$  is given. Then take  $N > (1 + \varepsilon)/\varepsilon$  and  $k = n^{1+\varepsilon}$ . Then we have for large  $n$

$$p_{m,R}(2n) \geq (2C_m)^{-1} (2n)^{-m(1+\varepsilon)}$$

from which we have

$$\varlimsup_{t \rightarrow \infty} p_{m,R}(t) t^{m(1+\varepsilon)} \geq \varlimsup_{n \rightarrow \infty} p_{m,R}(2n) (2n)^{m(1+\varepsilon)} \geq (2C_m)^{-1}.$$

Thus Theorem A is proved.

## References

- [1] F. Asakura: *On the Green's function for  $\Delta - \lambda^2$  with the boundary condition of the third kind in the exterior domain of a bounded obstacle*, J. Math. Kyoto Univ. **18** (1978), 615–625.
- [2] M. Ikawa: *Problèmes mixtes pour l'équation des ondes*, II, Publ. Res. Inst. Math. Sci. Kyoto Univ. **13** (1977), 61–106.
- [3] ———: *Mixed problems for the wave equation*, III, *Exponential decay of solutions*, Publ. Res. Inst. Math. Sci. Kyoto Univ. **14** (1978), 71–110.
- [4] ———: *Mixed problems for the wave equation*, IV, *The existence and the exponential decay of solutions*, J. Math. Kyoto Univ. **19** (1979), 375–411.
- [5] J.B. Keller, R.M. Lewis and B.D. Seckler: *Asymptotic solution of some diffraction problems*, Comm. Pure Appl. Math. **9** (1956), 207–265.
- [6] P.D. Lax and R.S. Phillips: *Scattering theory*, Academic Press, New York, 1967.
- [7] P.D. Lax, C.S. Morawetz and R.S. Phillips: *Exponential decay of solutions of the wave equation in the exterior of a star-shaped obstacle*, Comm. Pure Appl. Math. **16** (1963), 447–486.
- [8] R.K. Luneberg: *The mathematical theory of optics*, Brown Univ., 1944.

- [9] S. Mizohata: *Theory of partial differential equations*, Iwanami, Tokyo, 1965 (Japanese).
- [10] C.S. Morawetz: *Decay for solutions of the exterior problems for the wave equation*, Comm. Pure Appl. Math. **28** (1975), 229–264.
- [11] C.S. Morawetz, J.V. Ralston and W.A. Strauss: *Decay of solutions of the wave equation outside nontrapping obstacles*, Comm. Pure Appl. Math. **30** (1977), 447–508.
- [12] J. Ralston: *Solutions of the wave equation with localized energy*, Comm. Pure Appl. Math. **22** (1969), 807–823.
- [13] H.F. Walker: *Some remarks on the local energy decay of solutions of the initial-boundary value problem for the wave equation in unbounded domains*, J. Differential Equations **23** (1977), 459–471.

Department of Mathematics  
Osaka University  
Toyonaka, Osaka 560  
Japan

