

Title	Metal Ion-dependent Effects of Clioquinol on the Fibril Growth of an Amyloid eta Peptide
Author(s)	Raman, Bakthisaran; Ban, Tadato; Yamaguchi, Kei- ichi et al.
Citation	Journal of Biological Chemistry. 2005, 280(16), p. 16157-16162
Version Type	VoR
URL	https://hdl.handle.net/11094/71294
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Metal Ion-dependent Effects of Clioquinol on the Fibril Growth of an Amyloid β Peptide*

Received for publication, January 10, 2005, and in revised form, February 16, 2005 Published, JBC Papers in Press, February 16, 2005, DOI 10.1074/jbc.M500309200

Bakthisaran Ramanद, Tadato Ban‡, Kei-ichi Yamaguchi‡, Miyo Sakai‡, Tomoji Kawai||, Hironobu Naiki**§§, and Yuji Goto‡ ‡‡

From the ‡Institute for Protein Research, Osaka University, and CREST, Japan Science and Technology Agency, Yamadaoka 3-2, Suita, Osaka 565-0871, Japan, the §Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad 500 007, Andhra Pradesh, India, the ∥Institute for Scientific and Industrial Research, Osaka University, Osaka 567-0047, Japan, the **Faculty of Medical Science, University of Fukui, and §§CREST, Japan Science and Technology Agency, Matsuoka, Fukui 910-1193, Japan

Although metal ions such as Cu²⁺, Zn²⁺, and Fe³⁺ are implicated to play a key role in Alzheimer disease, their role is rather complex, and comprehensive understanding is not yet obtained. We show that Cu²⁺ and Zn²⁺ but not Fe^{3+} renders the amyloid β peptide, $A\beta_{1-40}$, nonfibrillogenic in nature. However, preformed fibrils of $A\beta_{1-40}$ were stable when treated with these metal ions. Consequently, fibril growth of $A\beta_{1-40}$ could be switched on/off by switching the molecule between its apo- and holo-forms. Clioquinol, a potential drug for Alzheimer disease, induced resumption of the Cu²⁺-suppressed but not the Zn²⁺-suppressed fibril growth of A β_{1-40} . The observed synergistic effect of clioquinol and Zn² suggests that Zn²⁺-clioquinol complex effectively retards fibril growth. Thus, clioquinol has dual effects; although it disaggregates the metal ion-induced aggregates of $A\beta_{1-40}$ through metal chelation, it further retards the fibril growth along with Zn²⁺. These results indicate the mechanism of metal ions in suppressing Aβ amyloid formation, as well as providing information toward the use of metal ion chelators, particularly clioquinol, as potential drugs for Alzheimer disease.

Alzheimer disease $(AD)^1$ is a neurodegenerative disease characterized by cerebral deposits of extracellular amyloid plaques, intracellular tangles, and intravascular or extravascular deposits (1, 2). The deposits are comprised of a mixture of 39-43-amino-acid polypeptides generally designated as amyloid- β (A β) peptide (3, 4). A β peptides are secreted by proteolytic processing of several membrane proteins, which are alternative splicing products of the gene coding the amyloid precursor protein (APP) (5–7). Accumulation of the A β peptides as insoluble amyloid deposits occurs in AD due to yet unclear mechanism(s).

The role of metal ions, notably Cu^{2+} , Zn^{2+} , and Fe^{3+} , in AD

and metal ion chelators as therapeutic agents has been the subject of interest over recent years (8–10). Supplementing Cu^{2+} in drinking water leads to development of the disease in rabbit model (11). High concentrations of Cu^{2+} (0.4 mM) and Zn^{2+} (1 mM) have been reported in the senile plaques of the diseased brain, extracellular Cu^{2+} concentration reaching to 15 μ M (12). In contrast, Cu^{2+} has been reported to decrease A β deposits in APP23 transgenic mice (13). Moreover, overexpression of human A β peptides in transgenic mice leads to a decrease in brain Cu^{2+} (14). APP knock-out mice show an increase in Cu^{2+} (15). Thus, the role of metal ions is rather complicated; it is not clear whether the accumulation of metal ions is the cause or the consequence of AD or a biological defense response to decrease progression of the disease.

defense response to decrease progression of the disease. The $A\beta$ peptides bind Cu^{2+} and Zn^{2+} with high affinity (16-21). Although histidine residues, especially His-13, are involved in binding Cu²⁺ and Zn²⁺ (20-22), Cu²⁺ and Zn²⁺ differ in their coordination to His-13 (22). Cu^{2+} and Zn^{2+} trigger aggregation and precipitation of A β peptides (19, 23– 28), which are distinct from the amyloid fibrils, especially in the β -sheet content (27, 28). Toxicity of granular aggregates is a target of recent extensive studies (29, 30). In this respect, the use of "metal-protein attenuation compounds" is proposed as a potential strategy for treating AD (31, 32). The antibiotic, clioquinol (CQ), which is also known to be a transition metal ion chelator (33, 34), has been found to dissolve deposits of the AD brain tissue (8) and to yield promising results in an animal model for AD in decreasing the number of deposits (8, 36). CQ has been the focus of attention as a potential drug (8). However, its effect on the amyloid fibril formation, especially in conjunction with Cu²⁺ or Zn²⁺, has not been understood so far. Therefore, it is important to obtain comprehensive understanding of the interactions with metal ions and their impact on the aggregation and amyloid fibril formation of $A\beta$ peptides.

In the present study, we investigated the role of some metal ions on the aggregation and fibrillogenic propensities of $A\beta_{1-40}$. The results show that Cu^{2+} and Zn^{2+} , but not Fe^{3+} , render $A\beta_{1-40}$ less fibrillogenic or non-fibrillogenic under conditions in which no significant metal ion-induced aggregation occurs. We used this system to examine the effects of metal ion chelators, EDTA and CQ, on the fibril growth of $A\beta_{1-40}$. The results revealed the dual effects of CQ, explaining the underlying mechanism of clioquinol as a potential drug for Alzheimer disease.

EXPERIMENTAL PROCEDURES

Materials—Human $A\beta_{1-40}$ was purchased from the Peptide Institute, Inc. (Osaka, Japan). CuCl₂ dihydrate, anhydrous ZnCl₂, and anhydrous FeCl₃ were purchased in their purest forms from Aldrich. CQ (5-chloro-7-iodo-8-hydroxyquinoline, commonly known as clioquinol)

^{*} This work was supported in part by Grants-in-aid for Priority Areas (15076206, 15032228) and scientific Research (B) (13480219) from the Japanese Ministry of Education, Culture, Sports, Science and Technology. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

[¶] Supported by a Post Doctoral Fellowship from the Japan Society for Promotion of Science.

^{‡‡} To whom correspondence should be addressed. Tel.: 81-6-6879-8614; Fax: 81-6-6879-8616; E-mail: ygoto@protein.osaka-u.ac.jp.

¹ The abbreviations used are: AD, Alzheimer disease; A β , amyloid β ; APP, amyloid precursor protein; CQ, clioquinol; ThT, thioflavin T; TIRFM, total internal reflection fluorescence microscopy.

was procured from Calbiochem. "Protein assay Coomassie Brilliant Blue solution" was purchased from Nacalai Tesque Inc. (Kyoto, Japan).

Effects of CQ and Buffer Conditions on Metal Ion-induced Aggregation of $A\beta_{I-40}$ —50 μ M $A\beta_{I-40}$ in the indicated buffers in the presence or absence of indicated substances was incubated at 37 °C for 15 min. The final concentrations of Cu²⁺ or Zn²⁺ in the samples were 50 μ M. Stock CQ solution was prepared in methanol, and it was added to a final concentration of 100 μ M. The samples were centrifuged at 13,000 × g for 15 min. Peptide concentration in the supernatant was determined by the Bradford method using the commercial protein assay Coomassie Brilliant Blue solution and was represented as a percentage of recovery relative to the control without metal ion and CQ. We did not measure the trace contamination of the metal ions in our buffer system or from the peptide preparations. Therefore, the mentioned concentrations of the metal ions represent the added concentrations of the metal ions. Unless otherwise mentioned, the final concentration of methanol in our experiment due to the addition of CQ did not exceed 2%.

Atomic Force Microscopy—Samples of 50 μ M A β_{1-40} in 20 mM HEPES buffer, pH 7.4, containing 100 mM NaCl and 50 μ M metal ions were incubated for 5 min. An aliquot of the sample was placed on a mica surface and incubated for 1 min. Excess sample was drained and washed quickly with water to remove salts. The mica surface was air-dried. Atomic force microscopy imaging was performed using a dynamic force microscope, SPI3700-SPA300, Seiko Instruments. The scanning tip was a silicon microcantilever (SI-DF20, Seiko Instruments, spring constant = 13 newton/m, resonance frequency = 135 kHz). The scan rate was 0.5 Hz.

Amyloid Fibril Growth of $A\beta_{1-40}$ —Samples of 50 μ M (~0.22 mg/ml) A β_{1-40} in 50 mM sodium phosphate buffer, pH 7.4, containing 100 mM NaCl (referred to as buffer A henceforth) in the absence and presence of required concentrations of metal ions were incubated at 37 °C with 5 μ g/ml sonicated fibrils (fibril seed). Aliquots (5 μ l) of the samples were withdrawn at different time points and added to 1 ml of 5 μ M thioflavin T in 50 mM glycine-NaOH buffer, pH 8.5 (ThT solution). Fluorescence intensity at 485 nm, which is proportional to the extent of fibril-bound ThT (37) and thus the amyloid fibril growth, was measured with an excitation wavelength of 445 nm using a Hitachi F-4500 fluorescence spectrophotometer. To test the effect of EDTA and CQ, the required concentrations of the chelators were added at the indicated time points.

Total Internal Reflection Fluorescence Microscopy (TIRFM)—ThT was added to the samples in buffer A to a final concentration of 5 μ M, and the sample was placed on a glass slide. The TIRFM system to observe ThT-bound amyloid fibrils was developed based on an inverted microscope (IX70; Olympus, Tokyo, Japan) as described earlier (38, 39).

Circular Dichroism—Samples of 0.22 mg/ml $A\beta_{1-40}$ peptide alone in buffer A and the samples of the peptide and fibril seeds (5 µg/ml) incubated in the absence or presence of required concentrations of metal ions at 37 °C for 80 min were prepared. These samples were diluted 1:1 (v/v) with buffer A, and the far UV CD spectra were recorded using a Jasco-600 spectropolarimeter at 37 °C with thermo-stated cell holder and a 0.1-cm path length quartz cuvette. The data shown are the mean residue ellipticity.

Sedimentation Velocity Measurements—Sedimentation velocity measurements were performed using an Optima XL-I analytical ultracentrifuge (Beckman Coulter) with An-60 rotor and two-channel charcoal-filled Epon cells. Samples (0.3 ml) of 30 μ M A β_{1-40} in buffer A in the absence and in the presence of 50 μ M metal ions or 100 μ M CQ were incubated at 37 °C for 15 min followed by centrifugation at the same temperature at 224,000 × g. The protein boundary was scanned at 12-min intervals for its absorbance at 225 nm. The boundary curves at every 24-min intervals are shown. The sedimentation data were analyzed using the software Origin 4.1 (OriginLab).

RESULTS

Effects of CQ and Buffer Conditions on Metal Ion-induced Aggregation of $A\beta_{1-40}$ —In HEPES buffer containing 100 mM NaCl (pH 6.6 or 7.4), both 50 μ M Zn²⁺ and 50 μ M Cu²⁺ largely decreased the percentage of recovery of $A\beta_{1-40}$ (Fig. 1, *A* and *B*). The Cu²⁺-induced decrease in the percentage of recovery was more significant at pH 6.6 than at pH 7.4, indicating the pH dependence of the Cu²⁺ effects. Although 100 μ M CQ did not itself exhibit notable effect on the aggregation of $A\beta_{1-40}$, it suppressed the metal ion-induced aggregation of the peptide, increasing its recovery (Fig. 1, *A* and *B*). Thus, in HEPES buffer, Cu²⁺ and Zn²⁺ induce aggregation of $A\beta_{1-40}$, in agreement with the earlier studies (19, 23–28), which can be sup-



FIG. 1. Effect of buffer conditions and CQ on Cu²⁺- or Zn²⁺induced aggregation of $A\beta_{1-40}$. Buffers used were 20 mM HEPES buffer, pH 6.6, containing 100 mM NaCl (A), 20 mM HEPES buffer, pH 7.4, containing 100 mM NaCl (B), and 50 mM sodium phosphate buffer, pH 7.4, containing 100 mM NaCl (buffer A) (C). The final concentrations of Cu²⁺, Zn²⁺, and CQ were 50, 50, and 100 μ M, respectively. D, atomic force microscopy images (area corresponds to 2 μ m²) of amyloid fibrils of $A\beta_{1-40}$ (*image 1*) and the aggregates of $A\beta_{1-40}$ formed in 20 mM HEPES buffer, pH 7.4, containing 100 mM NaCl and either 50 μ M Zn²⁺ (*image 2*) or 50 μ M Cu²⁺ (*image 3*).

pressed by CQ. The atomic force microscopy images (Fig. 1D) indicated that metal ion-induced aggregates are amorphous in nature, supporting earlier reports (27, 28).

Importantly, metal ion-induced aggregation of $A\beta_{1-40}$ depended on buffer conditions. In buffer A, we did not observe metal ion-induced loss of $A\beta_{1-40}$ (Fig. 1*C*). Furthermore, sedimentation velocity measurements showed that the metal ion-treated or untreated $A\beta_{1-40}$ do not exhibit significant change in their sedimentation property (Fig. 2). The apparent sedimentation coefficients of $A\beta_{1-40}$ in buffer A alone and in the presence of 50 μ M Zn²⁺ and 50 μ M Cu²⁺ were calculated to be 0.96, 0.86, and 1.05 *s*, respectively. Thus, our result shows that the metal ions do not promote significant aggregation of $A\beta_{1-40}$ in buffer A.

One possible reason for the observation made in buffer A could be that the formation of the phosphate complexes of these metal ions prevents their interaction with the $A\beta$ peptide. However, our following results on the effects of Cu^{2+} and Zn^{2+} on the fibril growth and β -sheet formation of the peptide indicate that the peptide interacts with these metal ions in buffer A. Moreover, Cu^{2+} binding to the peptide in buffer A is further supported by the fluorescence quenching studies presented later on in this study. Thus, taking together these evidences, we presume that the observed lack of metal ion-induced aggregation in buffer A is not due to the lack of metal ion binding. Rather, it represents the conditions in which the metal ionbound peptide exhibits less or no aggregation. In this context, it is important to note that two different laboratories have earlier made contradictory observations on the susceptibility of A β peptide toward the Zn²⁺-induced aggregation (23, 25). Huang et al. (26) clarified that the susceptibility of the $A\beta$ peptide to the Zn²⁺-induced aggregation is sensitive to "complex factors in buffer milieu that impact upon the peptide's



FIG. 2. Effects of metal ions on sedimentation property of $A\beta_{1-40}$. *A*-*C* represent scans of boundary movement of 30 μ M $A\beta_{1-40}$ in buffer A alone and in the presence of 50 μ M Zn^{2+} and 50 μ M Cu^{2+} , respectively, in an analytical ultracentrifugation experiment. *D*, analysis of the sedimentation pattern of $A\beta_{1-40}$. The In(*r*/*rm*) versus $\omega^2 t$ plot for the $A\beta_{1-40}$ in buffer A alone (\bigcirc) and in the presence of 50 μ M Zn^{2+} (\blacktriangledown) and 50 μ M Cu^{2+} (\bigstar) is shown.

conformation and polymerization state." Therefore, interestingly, our observation that metal ion-induced aggregation is less pronounced in buffer A is useful to understand the relative fibrillogenic propensities of the metal-free (apo) and metal-bound (holo) $A\beta_{1-40}.$

Effect of Cu^{2+} , Zn^{2+} , and Fe^{3+} on the Fibril Growth of $A\beta_{I-40}$ —The relative fibrillogenic propensities of the apo- and holo-forms of $A\beta$ peptides have not been understood so far. Since Cu^{2+} or Zn^{2+} does not induce detectable aggregation in buffer A, this system is useful to address the issue. We have investigated the effects of Cu^{2+} , Zn^{2+} , and Fe^{3+} on the amyloid fibril growth of $A\beta_{I-40}$ in buffer A (pH 7.4). Even at substoichiometric concentrations, Cu^{2+} retarded the fibril growth (Fig. 3A). At a molar ratio less than 0.4 of the metal ion to $A\beta_{I-40}$, Cu^{2+} increased the time lag of the onset of the fibril growth (despite providing the fibril seed for nucleation). At the molar ratio more than or equal to 0.4, fibril growth was not observed significantly during the observed time period. However, the fibril growth occurred slowly to the extent of the

control (without the metal ion) by 2 days of incubation at the molar ratio less than 0.6 (data not shown). At the molar ratio of 1.0, Cu^{2+} prevented the fibril growth completely; there was no fibril growth even after 5 days (data not shown). Zn^{2+} also exhibited a similar effect in preventing the amyloid fibril growth of $A\beta_{1-40}$ (Fig. 3B). On the other hand, Fe^{3+} did not prevent the fibril growth of $A\beta_{1-40}$ at concentrations comparable with those of Cu^{2+} or Zn^{2+} (Fig. 3B) as well as at its molar ratio as high as 4.0 (data not shown).

We examined by TIRFM the samples of ThT-bound fibril seed, the $A\beta$ amyloid fibrils, and the samples of the $A\beta$ peptide and the seed incubated along with Cu²⁺ or Fe³⁺ (Fig. 3C). Fibril growth of $A\beta_{1-40}$ occurred to the order of a few micrometers in buffer A alone (Fig. 3C, *image 2*) and even in the presence of 0.2 mM Fe³⁺ (Fig. 3C, *image 3*). On the other hand, we could see only fibril seeds in Cu²⁺-treated (Fig. 3C, *image 4*) or Zn²⁺-treated (data not shown) samples. Thus, Cu²⁺ or Zn²⁺ but not Fe³⁺ prevents the fibril growth of $A\beta_{1-40}$.

 Cu^{2+} and Zn^{2+} Prevent the β -Sheet Formation—Amyloid fibrils exhibit characteristic well ordered cross- β -sheet structure (40, 41). $A\beta_{1-40}$ freshly dissolved in buffer A exhibited random coil conformation as revealed in its far UV-CD spectrum (Fig. 4, *curve 1*). Incubation of the peptide in the presence of fibril seeds for 80 min generated a characteristic far-UV CD spectrum for β -sheet structure with a minimum around 218 nm (Fig. 4, *curve 2*). Such an association-induced generation of β -sheet structure also occurred in the presence of Fe³⁺ (Fig. 4, *curve 3*). However, induction of β -sheet structure did not occur in the presence of Cu²⁺ or Zn²⁺ (Fig. 4, *curves 4* and 5). Thus, Cu²⁺ or Zn²⁺ but not Fe³⁺ prevents the generation of β -sheet structure, consistent with the effects on fibril growth monitored by ThT fluorescence (Fig. 3).

Copper Binding to $A\beta_{I-40}$ in Buffer A—Prevention of the fibril growth and the association-induced β -sheet formation shows that the metal ions, Cu^{2+} and Zn^{2+} , interact with the $A\beta$ peptide in buffer A, where phosphate complexation of the metal ions is also favored. We further investigated the Cu^{2+} binding by fluorescence spectroscopy.

The sequence of $A\beta_{1-40}$ does not contain any tryptophan residue but a single tyrosine residue at position 10 in addition to two phenylalanine residues at positions 19 and 20. The intrinsic fluorescence of the peptide thus predominantly represents the fluorescence of the sole tyrosine residue upon exciting at 280 nm. The intrinsic fluorescence of $A\beta_{1-28}$ is known to be quenched upon binding to Cu^{2+} (20). We have investigated the effect of the metal ions on the intrinsic fluorescence of $A\beta_{1-40}$ peptide. The fluorescence intensity of the peptide progressively decreased upon the addition of the paramagnetic ion, Cu²⁺ (Fig. 5A). This observed quenching of the fluorescence is not due to the general collisional quenching of the fluorescence by the paramagnetic ion, $\mathrm{Cu}^{2+},$ as the fluorescence of N-acetyl tyrosine amide was not significantly quenched under the concentration range of the metal ion used (Fig. 5B). Thus, the specific quenching of the intrinsic fluorescence of $A\beta_{1-40}$ observed in buffer A is due to the specific binding of Cu²⁺, leading to proximal interactions between the quencher (Cu^{2+}) and the fluorophore (the tyrosine residue of the peptide). Using the fluorescence quenching data (Fig. 5C), the dissociation constant (K_d) of copper binding to $A\beta_{1-40}$ was calculated to be $\sim 8 \times 10^6$ M. The addition of the diamagnetic ion, Zn^{2+} , however, did not lead to quenching of the fluorescence of $A\beta_{1-40}$ (Fig. 5C). Thus, the fluorescence experiment shows that Cu^{2+} binds to the A β peptide in buffer A, under the condition in which the metal ion-induced prevention of fibril growth but not the metal ion-induced aggregation was observed. We presume that Zn^{2+} also interacts with or binds to $A\beta_{1-40}$ in buffer A

FIG. 3. Effect of metal ions on the amyloid fibril growth of $A\beta_{1-40}$. A, the amyloid fibril growth of A β_{1-40} (50 μ M) in buffer A in the absence (\bigcirc) and in the presence of 5 (\bullet), 10 (\triangle), 15 (\blacktriangle), and 20 μ M (\Box) Cu²⁺. *B*, the percentage of prevention of the fibril growth as a function of the molar ratio of metal ion to $A\beta_{1-40}$. Metal ions used are Cu^{2+} (O), Zn^{2+} (\bullet), and Fe^{3+} (\triangle). The percentage of prevention of the fibril growth was calculated using the formula, $[1 - (F_{m90} - F_{m0})/(F_{90} - F_0)] \times 100$, where F_0 and F_{m0} are the ThT fluorescence intensities at zero time in the absence and in the presence of the metal ions and F_{90} and F_{m90} are the fluorescence intensities after a 90-min incubation in the absence and in the presence of the metal ions, respectively. C, TIRFM images of fibril seeds (image 1); fibrils formed in buffer A alone (image 2); and the $A\beta_{1-40}$ samples incubated along with fibril seeds and 0.2 mM Fe^{3+} (*image 3*) and 30 μ M Cu²⁺ (*image 4*). The yellow bar in the images represents 5 μ m.





FIG. 4. Effect of metal ions on the fibril growth-associated generation of β -sheet. Far-UV CD spectra of $A\beta_{1-40}$ peptide (*curve 1*) and the sample of the $A\beta$ peptide (50 μ M) incubated at 37 °C with 5 μ g/ml fibril seed for 80 min in the absence (*curve 2*) and presence of 200 μ M Fe³⁺ (*curve 3*), 50 μ M Zn²⁺ (*curve 4*), or 30 μ M Cu²⁺ (*curve 5*) are shown.

since ${\rm Zn}^{2+}$ behaved similarly to ${\rm Cu}^{2+}$ on the fibril growth of ${\rm A}\beta_{1-40}~({\rm Fig.~}3B)$

Switching On / Off the Fibril Growth—One of the reasons for our observation that fibril growth is prevented by the metal ions is that the metal ions destabilize the fibrils. However, incubation of the fibrils of the A β peptide with the metal ions at 37 °C did not result in significant dissociation of the fibrils as judged by ThT binding (data not shown), thus ruling out the possibility of fibril instability as one of the reasons for the observed metal-induced prevention of the fibril growth. Since Cu^{2+} -treated fibrils are stable but Cu^{2+} prevents the fibril growth, it was possible to switch on/off the fibril growth by switching $A\beta_{1-40}$ to its fibrillogenic apo- and non-fibrillogenic holo-forms (Fig. 6). The results show that one of the key factors determining the $A\beta$ fibril growth is the environmental concentrations of metal ions.

Metal Ion-dependent Effects of CQ on the Fibril Growth—The antibiotic CQ, which can bind both Cu^{2+} and Zn^{2+} with the same stoichiometry but with different coordination geometry (33), has been proposed to serve as a potential drug for AD (8, 31, 32). It would be pertinent to study the effects of CQ on the A β amyloid fibril growth as well as on the metal ion-induced suppression of the fibril growth.

 \overline{CQ} alone did not affect the fibril growth of $A\beta_{1-40}$ significantly (Fig. 7A). The addition of CQ or EDTA to the sample of



FIG. 5. Cu^{2^+} binding to $A\beta_{1-40}$ in Buffer A monitored by quenching of the intrinsic fluorescence of the peptide. A, intrinsic fluorescence spectra of 10 μ M $A\beta_{1-40}$ (from top to bottom) in the absence and in the presence of 4, 8, 12, 18, 32 μ M Cu^{2^+} , respectively. B, the fluorescence spectra of N-acetyl tyrosine amide in the absence (*curve 1*) and in the presence (*curve 2*) of 50 μ M Cu^{2^+} . C, the relative change in the fluorescence intensity at 304 nm of $A\beta_{1-40}$ as a function of increasing concentrations of Cu^{2^+} (\bigcirc) and Zn^{2^+} (\triangle). The samples were excited at 280 nm.

Cu²⁺-treated A β_{1-40} led to resumption of the Cu²⁺-suppresed fibril growth (Fig. 7*B*). Interestingly, the addition of EDTA, but not CQ, led to resumption of the fibril growth suppressed by Zn²⁺ (Fig. 7*C*). Since 8-hydroxyquinoline derivatives are well known transition metal ion chelators (33, 34) and CQ forms complexes with both Zn²⁺ and Cu²⁺ with a similar stoichiometry of 2:1 (33), it is unlikely that the lack of resumption of the Zn²⁺-suppressed fibril growth is because CQ does not chelate



FIG. 6. Switching on/off the fibril growth of $A\beta_{1-40}$ by Cu^{2+} and EDTA monitored by ThT fluorescence. The *left vertical dashed line* represents the addition of 30 μ M Cu²⁺, and the *right vertical line* represents the addition of 50 μ M EDTA. Buffer A was used.



FIG. 7. Effects of metal chelators on the metal ion-induced suppression of the fibril growth of $A\beta_{1-40}$. A, the effect of CQ on the amyloid fibril growth of $A\beta_{1-40}$. The fibril growth in the absence (\bigcirc) and presence (\bigcirc) of 160 μ M CQ. B, the effect of the metal chelators, 100 μ M EDTA (\bigcirc) and 160 μ M CQ (\bigcirc), on the 50 μ M Cu²⁺-suppressed fibril growth. C, the effect of 100 μ M EDTA (\bigcirc) and 160 μ M CQ (\bigcirc) on the 50 μ M CQ (\bigcirc) on the 50 μ M CQ (\bigcirc) on the 50 μ M CR²⁺-suppressed fibril growth. The arrows indicate the addition of chelators. The final concentration of methanol due to the addition of CQ in this experiment was ~4% (this did not affect the fibril growth significantly).

Zn²⁺. Since complexes of 8-hydroxyquinoline derivatives with Zn²⁺ are relatively less stable when compared with those with Cu²⁺ (33, 34), a second possibility could be that the possible lesser stability of the Zn²⁺-CQ complex may lead to shuttling of Zn²⁺ between CQ and the A β peptide, which may transiently increase the population of holo-form of A β_{1-40} that is less competent for fibril growth. The third possibility is that the Zn²⁺-CQ complex exhibits novel retardation effect on fibril growth. We, therefore, attempted to verify these possibilities as follows.

At 10 μ M Zn²⁺, the fibril growth of the peptide was prevented only partially (Fig. 8A). Since CQ itself does not exhibit significant effect on the fibril growth, the addition of CQ to the sample should not have any effect in the presence of Zn²⁺ if the first possibility of the lack of metal chelation by CQ is valid. If the second possibility is the primary reason, then in the presence of Zn²⁺ and CQ, the fibril growth profile would be expected to be in between that of the control and that of the sample in the presence of Zn²⁺ alone. Surprisingly, the addition of CQ led to further retardation of the fibril growth, indicating a synergistic effect of CQ and Zn²⁺ on the fibril growth



FIG. 8. Distinct effects of CQ complexation of \mathbb{Zn}^{2+} and \mathbb{Cu}^{2+} on the fibril growth of $A\beta_{1-40}$. *A*, synergistic effect of cliquinol and \mathbb{Zn}^{2+} in retarding the fibril growth of $A\beta_{1-40}$. *B*, effect of CQ and \mathbb{Cu}^{2+} on the $A\beta_{1-40}$ fibril growth. The conditions of amyloid fibril growth of $A\beta_{1-40}$ were in buffer A alone (\bigcirc) , in the presence of 20 μ M CQ($\textcircled{\bullet}$), 10 μ M metal ions (\triangle) , or a mixture of 20 μ M CQ and 10 μ M respective metal ion (\bigstar).



FIG. 9. Analysis of the sedimentation pattern of $A\beta_{1-40}$ in the presence of CQ complexes of metal ions. The $\ln(r/rm)$ versus $\omega^2 t$ plot for the $A\beta_{1-40}$ in buffer A alone (\bigcirc) and in the presence of 50 μ M Zn^{2+} and 100 μ M CQ (\blacktriangle) and 50 μ M Cu^{2+} and 100 μ M CQ (\bigstar). The apparent sedimentation coefficients of the peptide in buffer alone, in the presence of Zn^{2+} -CQ and Cu^{2+} -CQ, were calculated to be 0.9, 0.88, and 1.2 *s*, respectively.

of $A\beta_{1-40}$ (Fig. 8A). On the other hand, we did not observe such synergistic effect of CQ in the presence of Cu²⁺; CQ completely eliminates the suppressive effect of Cu²⁺ on fibril growth (Fig. 8B).

From the observed synergistic effect of CQ and Zn^{2+} , it appears that the third possibility is responsible for the observed phenomenon of distinct effects of CQ on the Cu²⁺- and Zn²⁺-suppressed fibril growth; Zn²⁺-CQ complex, but not Cu²⁺-CQ complex, retards the fibril growth. Sedimentation velocity experiments of the A β peptide in the absence and presence of metal ions and CQ revealed no significant changes (Fig. 9), suggesting that metal ion-CQ complexes do not promote other non-fibrillar aggregates not detected by ThT fluorescence. Thus, our results demonstrate that the effect of CQ on the fibril growth process is metal ion-dependent.

DISCUSSION

There has been an increasing interest in the role of transition metal ions in AD (8–10) as $A\beta$ peptides exhibit high affinity binding to Cu^{2+} and Zn^{2+} (16–22). Our study provides two important findings toward the understanding of the role of metal ions and metal ion chelators in AD.

Firstly, our study demonstrates that the soluble holo-forms (Cu²⁺- or Zn²⁺-bound) of $A\beta_{1-40}$ are less fibrillogenic or non-fibrillogenic in nature, which was not understood earlier. Al-

though Fe³⁺ is also found at high concentration in AD brain (12), it does not exhibit a significant effect on the fibrillogenic propensity of $A\beta_{1-40}$. It appears that high affinity binding of Cu^{2+} and Zn^{2+} may actually be beneficial as it does not favor fibril formation. On the other hand, high concentrations of metal ion under certain buffer conditions, as shown in Fig. 1, lead to aggregation and precipitation of the peptide, which may be toxic (29, 30).

The second important finding is that CQ exhibits unique metal ion-dependent effects on the fibril growth of the $A\beta_{1-40}$. Chelation of the metal ions such as Cu^{2+} and Zn^{2+} by EDTA leads to conditions favorable for amyloid fibril growth. Animal experiments with CQ have provided encouraging results as a potential drug (8, 31, 32, 36). Our in vitro study shows that the addition of CQ leads to resumption of Cu²⁺-suppressed but not Zn^{2+} -suppressed amyloid fibril growth of $A\beta_{1-40}$. CQ and Zn^{2+} together exhibit synergistic retardation of the fibril growth, suggesting that retardation of fibril growth by the Zn²⁺-CQ complex is responsible for the observed distinct effects of CQ on the Cu²⁺- and Zn²⁺-suppressed fibril growth. However, we cannot completely rule out the possibility of shuttling of Zn²⁺ between CQ and $A\beta_{1-40}$ due to the possible lesser stability of Zn^{2+} complexes of 8-hydroxyquinoline derivatives (33, 34), leading to the transient formation of the less fibrillogenic holoform. Both of these mechanisms should prove advantageous for the therapeutic use of CQ in AD. Monomeric $A\beta$ is not toxic to the cells (29, 30, 35). Toxicity has been reported to be associated with β -sheet generation and fibril formation (35). On the other hand, granular aggregates, formed before fibril formation, were found be more toxic than the fibrils, although the toxicity depends on the size of the aggregates (29, 30). In this respect, CQ can disaggregate the metal-induced aggregates of $A\beta_{1-40}$ under certain buffer conditions (Fig. 1) and, along with Zn²⁺, it can retard fibril growth (Figs. 5 and 6).

Although our results suggest that the Zn²⁺-complex, but not the Cu²⁺-complex of CQ, retards the fibril growth of A β_{1-40} , how the complexes distinctly affect fibril propagation is not clear. We consider that the difference in the observed coordination geometry of the complexes of Cu^{2+} and Zn^{2+} (33) may have some role to play. The planar nature of the complex of Cu²⁺ may lead to stacking of the aromatic ring, promoting self-association of the complex. Such stacking is not favorable due to the fifth coordination involving a water molecule as well as due to puckered configuration of the Zn²⁺ complex of CQ (33), which can increase the availability of the complex molecule to interact or interfere in the amyloid fibril growth of the A β peptide, probably through hydrophobic interactions.

We conclude that Cu^{2+} and Zn^{2+} , but not Fe^{3+} , significantly prevent the amyloid fibril growth of $A\beta_{1-40}$. Our study demonstrates for the first time that the soluble holo-form of the peptide is non-fibrillogenic in nature and that the fibril growth of the peptide can be switched on/off by switching the molecule between its apo- and holo-forms. CQ exhibits distinct effects on the Cu²⁺- and Zn²⁺-induced suppression of the fibril growth. Particularly, complexation of Zn²⁺ with CQ may have dual beneficial effects in not only disaggregating the metal ioninduced aggregates but also in retarding the fibril growth of $A\beta_{1-40}$. Our study should prove useful in understanding the role of the transition metal ions in amyloid fibril propagation of $A\beta$ peptides in AD and provide information concerning the use of metal chelators, particularly CQ, as potential drugs for AD.

REFERENCES

- 1. Hardy, J., and Selkoe, D. J. (2002) Science 297, 353-356
- 2. Selkoe, D. J. (1997) Science 275, 630-631
- 3. Glenner, G. G., and Wong, C. W. (1984) Biochem. Biophys. Res. Commun. 120, 885-890
- Masters, C. L., Simms, G., Weinman, N. A., Multhaup, G., Mc Donald, B. L., and Beyreuther, K. (1985) Proc. Natl. Acad. Sci. U. S. A. 82, 4245–4249
 Shoji, M., Golde, T., Ghiso, J., Cheung, T., Estus, S., Shaffer, L., Cai, X.,
- McKay, D., Tintner, R., Frangione, B., and Younkin, S. (1992) Science 258, 126 - 129
- 6. Busciglio, J., Gabuzda, D., Matsudaira, P., and Yankner, B. (1993) Proc. Natl. Acad. Sci. U. S. A. 90, 2090-2096
- 7. Seubert, P., Vigo-Pelfrey, C., Esch, F., Lee, M., Dovey, H., Davis, D., Sinha, S., Schlossmacher, M., Whaley, J., Swindlehurst, C., McCormack, R., Wolfert, R., Selkoe, D., Lieberberg, I., and Schenk, D. (1992) Nature 359, 325–327 8. Bush, A. I. (2003) Trends Neurosci. 26, 207-214
- 9. Gnjec, A., Fonte, J. A., Atwood, C., and Martins, R. N. (2002) Front Biosci. 7,
- d1016-d1023
- 10. Doraiswamy, P. M., and Finefrock, A. E. (2004) Lancet Neurol. 3, 431-434 11. Sparks, D. L., and Schreurs, B. G. (2003) Proc. Natl. Acad. Sci. U. S. A. 100,
- 11065-11069 12. Lovell, M. A., Robertson, J. D., Teesdale, W. J., Campbell, J. L., and Markesbery, W. R. (1998) J. Neurol. Sci. 158, 47-52
- 13. Bayer, T. A., Schafer, S., Simons, A., Kemmling, A., Kamer, T., Tepest, R., Eckert, A., Schussel, K., Eikenberg, O., Sturchler-Pierrat, C., Abramowski, D., Staufenbiel, M., and Multhaup, G. (2003) Proc. Natl. Acad. Sci. U. S. A. 100, 14187-14192
- 14. Maynard, C. J., Cappai, R., Volitakis, I., Cherny, R. A., White, A. R., Beyreuther, K., Masters, C. L., Bush, A. I., and Li, Q. X. (2002) J. Biol. Chem. 277, 44670-44676
- 15. White, A. R., Reyes, R., Mercer, J. F., Camakaris, J., Zheng, H., Bush, A. I., Multhaup, G., Bevreuther, K., Masters, C. L., and Cappai, R. (1999) Brain Res. 842, 439-444
- 16. Atwood, C. S., Scarpa, R. C., Huang, X., Moir, R. D., Jones, W. D., Fairlie, D. P., Tanzi, R. E., and Bush, A. I. (2000) J. Neurochem. 75, 1219-1233
- 17. Dong, J., Atwood, C. S., Anderson, V. E., Siedlak, S. L., Smith, M. A., Perry, G., and Carey, P. R. (2003) Biochemistry 42, 2768-2773
- Curtain, C. C., Ali, F., Volitakis, I., Cherney, R. A., Norton, R. S., Beyreuther, K., Barrow, C. J., Masters, C. L., Bush, A. I., and Barnham, K. J. (2001) J. Biol. Chem. 276, 20466–20473
- 19. Bush, A. I., Pettingell, W. H., Paradis, M.-d., Jr., and Tanzi, R. E. (1994) J. Biol. Chem. 269, 12152-12158
- 20. Syme, C. D., Nadal, R. C., Rigby, S. E. J., and Viles, J. H. (2004) J. Biol. Chem. 279, 18169-18177
- 21. Liu, S. T., Howlett, G., and Barrow, C. J. (1999) Biochemistry 38, 9373-9378 22. Miura, T., Suzuki, K., Kohata, N., and Takeuchi, H. (2000) Biochemistry 39,
- 7024-7031
- 23. Bush, A. I., Pettingell, W. H., Multhaup, G., Paradis, M., Vonsattel, J. P., Gusella, J. F., Beyreuther, K., Masters, C. L., and Tanzi, R. E. (1994) Science 265, 1464-1467
- 24. Atwood, C. S. Moir, R. D., Huang, X., Scarpa, R. C., Bacarra, N. M., Romano, D. M., Hartshorn, M. A., Tanzi, R. E., and Bush, A. I. (1998) J. Biol. Chem. **273,** 12817–12826
- 25. Esler, W. P., Stimson, E. R., Jennings, J. M., Ghilardi, J. R., Mantyh, P. W., and Maggio, J. E. (1996) J. Neurochem. 66, 723-732
- 26. Huang, X., Atwood, C. S., Moir, R. D., Hartshorn, M. A., Vonsattel, J. P., Tanzi, R. E., and Bush, A. I. (1997) J. Biol. Chem. 272, 26464-26470
- 27. Yoshiike, Y., Tanemura, K., Murayama, O., Akagi, T., Murayama, M., Sato, S., Tanaka, N., and Takashima, A. (2001) J. Biol. Chem. 276, Sun, X., 32293-32299
- 28. Klug, G. M., Losic, D., Subasinghe, S. S., Aguilar, M. I., Martin, L. L., and Small, D. H. (2003) Eur. J. Biochem. 270, 4282-4293
- 29. Kayed, R., Head, E., Thompson, J. L., McIntire, T. M., Milton, S. C., Cotman, C. W., and Glabe, C. G. (2003) Science 300, 486-489
- 30. Hoshi, M., Sato, M., Matsumoto, S., Noguchi, A., Yasutake, K., Yoshida, N., and Sato, K. (2003) Proc. Natl. Acad. Sci. U. S. A. 100, 6370-6375
- 31. Cuajungco, M. P., Faget, K. Y., Huang, X., Tanzi, R. E., and Bush, A. I. (2000) Ann. N. Y. Acad. Sci. 920, 292-304
- 32. Ritchie, C. W., Bush, A. I., Mackinnon, A., Macfarlane, S., Mastwyk, M., MacGregor, L., Kiers, L., Cherny, R., Li, Q. X., Tammer, A., Carrington, D., Mavros, C., Volitakis, I., Xilinas, M., Ames, D., Davis, S., Beyreuther, K., Tanzi, R. E., and Masters, C. L. (2003) Arch. Neurol. 60, 1685–1691
 33. Di Vaira, M., Bazzicalupi, C., Orioli, P., Messori, L., Bruni, B., and Zatta, P.
 - (2004) Inorg. Chem. 43, 3795-3797
- 34. Lane, T. J., Sam, C. S. C. A., and Kandathil, A. J. (1960) J. Am. Chem. Soc. 82, 4462 - 4465
- 35. Lorenzo, A., and Yankner, B. A. (1994) Proc. Natl. Acad. Sci. U. S. A. 91, 12243-12247
- 36. Cherny, R. A., Atwood, C. S., Xilinas, M. E., Gray, D. N., Jones, W. D., McLean, C. A., Barnham, K. J., Volitakis, I., Fraser, F. W., Kim, Y., Huang, X., Goldstein, L. E., Moir, R. D., Lim, J. T., Beyreuther, K., Zheng, H., Tanzi, R. E., Masters, C. L., and Bush, A. I. (2001) Neuron 30, 665-676
- 37. Naiki, H., Higuchi, K., Hosokawa, M., and Takeda, T. (1989) Anal. Biochem. 177, 244-249
- 38. Ban, T., Hamada, D., Hasegawa, K., Naiki, H., and Goto, Y. (2003) J. Biol. Chem. 278, 16462–16465
- 39. Ban, T., Hoshino, M., Takahashi, S., Hamada, D., Hasegawa, K., Naiki, H., and Goto, Y. (2004) J. Mol. Biol. 344, 757-767
- 40. Wetzel, R. (2002) Structure 10, 1031-1036
- 41. Sunde, M., and Blake, C. C. (1998) Q. Rev. Biophys. 31, 1-39

Acknowledgments-B. Raman thanks Dr. T. Ramakrishna and Dr. Ch. Mohan Rao, Centre for Cellular and Molecular Biology, India for useful discussions.

Metal Ion-dependent Effects of Clioquinol on the Fibril Growth of an Amyloid β Peptide

Bakthisaran Raman, Tadato Ban, Kei-ichi Yamaguchi, Miyo Sakai, Tomoji Kawai, Hironobu Naiki and Yuji Goto

J. Biol. Chem. 2005, 280:16157-16162. doi: 10.1074/jbc.M500309200 originally published online February 16, 2005

Access the most updated version of this article at doi: 10.1074/jbc.M500309200

Alerts:

- When this article is cited
- When a correction for this article is posted

Click here to choose from all of JBC's e-mail alerts

This article cites 41 references, 18 of which can be accessed free at http://www.jbc.org/content/280/16/16157.full.html#ref-list-1